# **The Drinched Book**

# UNIVERSAL LIBRARY OU\_174179 AWARININ AWARINA

## A DICTIONARY

OF

# APPLIED CHEMISTRY

VOL. II.

# A DICTIONARY

OF

# APPLIED CHEMISTRY

BY

SIR EDWARD THORPE, CB, LLD, FRS

INTELLES FLORESSON OF CENERAL CHEMISTRY AND DESCRIPTION OF THE CHEMISTE LABORATORIES OF THE IMPERIAL COLLECT OF SCHENE AND LECTRODORY, SOUTH REINSTREAM, LONDON, LATE TRIVELPAL OF THE GOVERNMENT LABORATORY, AND A LAST LIFESPEED OF THE CHEMICAL SOCIETY AND OF THE SOCIETY OF CHEMICAL INDUSTRY

ASSISTED BY EMINENT CONTRIBUTORS

VOL. II.

REVISED AND ENLARGED EDITION

WITH ILLUSTRATIONS

LONGMANS, GREEN, AND CO.
9 PATERNOSTER ROW, LONDON
FOURTH AVENUE & 30111 STREET, NEW YORK
BOMBAY, CALCUTTA, AND MADRAS

1921

All rights reserved

### **ABBREVIATIONS**

OF THE TITLES OF JOURNALS AND BOOKS.

```
Amer Chem. J.
                       American Chemical Journal
Amer. J Pharm
                       American Journal of Pharmacy
Amer. J. Sci
                       American Journal of Science
Anal Fis. Quim.
                       Anales de la Sociedad Española Fisica y Quimica.
Analyst . . .
                       The Analyst.
Annalen
                       Annalen der Chemie (Justus Liebig).
Ann. Chim anal. .
                       Annales de Chimie analytique appliquée à l'Industrie, à l'Agriculture.
                           à la Pharmacie et à la Biologie
Ann. Clam. .
                       Annales de Chimie.
Ann. Falsif. .
                       Annales des Falsifications
Ann Inst. Pasteur.
                       Annales de l'Institut Pasteur.
Ann. Physik.
                       Annalen der Physik
Ann. Physique .
                       Annales de Physique.
Annalı Chim. Appl
                       Annali di Chimica Applicata
Apoth. Zert. . . . . Arch. Pharm. . .
                       Apotheker-Zeitung.
                       Archiv der Pharmazie.
Attı R. Accad.
  Lincer .
                       Atti della Reale Accademia dei Lincei.
Bentl. a. Trim. .
                       Bentley and Trimen Medicinal Plants
                       Berichte der Deutschen chemischen Gesellschaft.
Ber Deut. pharm
                       Berichte der Deutschen pharmazeutischen Gesellschaft.
  Ges
Bicd. Zentr. .
                      Biedermann's Zentralblatt für Agrikulturchemie und rationellen
                           Landwirtschafts-Betrieb
Bro-Chem. J
                       The Bio-Chemical Journal
Brochem Zertsch
                       Biochemische Zeitschrift
Brewers J
                       Brewei's Journal.
Bull Imp Inst
Bull. Soc chim
                       Bulletin of the Imperial Institute
                       Bulletin de la Société chimique de France.
Chem. Ind .
                      Chemische Industrie
Chem. News
                      Chemical News
Chem Soc Proc
                      Journal of the Chemical Society of London.
                                                                     Proceedings.
Chem Soc. Trans
Chem Weekblad
                      Journal of the Chemical Society of London
                                                                     Transactions
                      Chemisch Weekblad
Chem Zert. . . . Chem Zenti . .
                      Chemiker Zeitung
                      Chemisches Zentralblatt.
Compt. 1 end.
                      Comptes rendus hebdomadaires des Seances de l'Académie des
                           Sciences.
                      Dingler's polytechnisches Journal.
Dingl poly. J. .
Farber-Zeit .
                      Farber-Zeitung
Flück. a. Hanb.
                      Fluckiger and Hanbury Pharmacographia.
Frdl
                      Friedlander's Fortschritte der Teerfarbenfabrikation.
Gazz chim. ital.
                      Gazzetta chimica italiana.
Helv. Chim. Acta .
                      Helvetica Chimica Acta.
                      Neues Jahrbuch fur Mineralogie, Geologie und Palaeontologie.
Jahrb. Min..
J. Amer Chem. Soc.
                      Journal of the American Chemical Society.
J. Bd. Agric.
                      Journal of the Board of Aguculture.
J. Franklin Inst.
J Ind Eng. Chem.
J. Inst Brewing.
                      Journal of the Franklin Institute.
                      Journal of Industrial and Engineering Chemistry.
                      Journal of the Institute of Brewing.
J Pharm. Chim. .
J. Phys. Chem. .
                      Journal de Pharmacie et de Chimie.
                      Journal of Physical Chemistry.
J. pr. Chem.
J. Russ. Phys. Chem
                      Journal fur praktische Chemie.
  Soc. .
                      Journal of the Physical and Chemical Society of Russia.
J. Soc. Chem. Ind.
                      Journal of the Society of Chemical Industry.
J. Soc. Dyers.
                      Journal of the Society of Dyers and Colourists.
J. Tokyo Čhem Soc
                      Journal of the Tokyo Chemical Society.
J. Washington Acad.
  Sci
                      Journal of the Washington Academy of Sciences.
Kollord Zertsch.
                      Kolloid-Zeitschrift.
Met & Chem. Eng
                      Metallurgical and Chemical Engineering
Min. Mag. .
                      Mineralogical Magazine and Journal of the Mineralogical Society.
Monatshefte fur Chemie und verwandte Theile anderer Wissen-
Monatsh. . .
                          schaften.
Pharm. J.
                      Pharmace 'cal Journal.
Pharm, Zert.
                      Pharmaceutische Zeitung.
```

### vi ABBREVIATIONS OF THE TITLES OF JOURNALS AND BOOKS.

Phil. Mag	
Phil. Trans	Philosophical Transactions of the Royal Society.
Phot. J. $\cdot$	Photographic Journal.
Proc. Roy. Soc	Proceedings of the Royal Society.
Proc. Roy. Soc. Edin.	Proceedings of the Royal Society of Edinburgh,
Rec. trav. chim	Receuil des travaux chimiques des Pays-Bas et de la Belgique.
Trans Faraday Soc.	Transactions of the Faraday Society.
Zertsch. anal. Chem.	Zeitschrift für analytische Chemie.
Zertsch.angew.Chem	Zeitschrift für angewandte Chemie.
Zertsch. anorg. Chem	Zeitschrift für anorganische Chemie.
Zertsch. Nahr.	·
Genussm	Zeitschrift fur Untersuchung der Nahrungs- und Genussmittel.
Zertsch. offentl	, , , , , , , , , , , , , , , , , , ,
Chem	Zeitschrift fur offentliche Chemie.
Zeitsch. physikal.	
Chem	Zeitschrift für physikalische Chemie, Stochiometrie und Verwandt- schaftslehre.
Zertsch physiol.	
Chem	Hoppe-Seyler's Zeitschrift für physiologische Chemie.

### LIST OF CONTRIBUTORS

### TO VOLUME II.

- Dr. E. F. ARMSTRONG, FIC, F.RS. [CARBOHYDRATES; DEXTRINS, DIASTASE; EMULSIN.]
- HARRY BAKER, Esq., F.I.C. (The Castner Kellner Company, Weston Point). [Chlorine.]
- Professor C. O. BANNISTER, The University of Liverpool [Copper, Metallurgy of.]
- Dr. GEORGE BARGER, M.A., F.R.S., Professor of Chemisty in relation to Medicine in the University of Edinburgh. [Calumba; Cascara Sagrada; Cascarilla Bark, Cheiroline; Choline and Related Substances; Chrysarobin, Colchicum; Colocynth; Conine and the Alkaloids of Hemlock, Curara; Cusparia Bark; Dandelion Root; Digitalis, Dulcamara; Elaterium; Ergot of Rye; Eriodiamine, Erythrophleine; Eserine; Euonymus.]
- G. S. BLAKE, Esq, ARSM, B.Sc, M.IMM. [CHROMIUM.]
- BERTRAM BLOUNT, Esq., F.I.C. [CEMENT.]
- Professor W A BONE, D.Sc, F.R.S, Imperial College of Science and Technology, South Kensington, and ERNEST BURY, Esq, M.Sc, O B.E., The Skinningrove Iron Co, Ltd. [Coke Manufacture and the Recovery of By-Products]
- Professor W. A. BONE, D.Sc., F.R.S., and Professor R. V. Wheeler, D.Sc. [Gaseous Explosion.]
- Professor J S. S BRAME, FIC. [PROTECTION OF METALS FROM CORROSION.]
- JOSEPH BURTON, Esq. (Pilkington Tile and Pottery Co., Clifton Junction, Manchester).
  [Clay]
- Dr. J. C. CAIN, F.I.C, Editor of the Journal of the Chemical Society, London. [Disazo- and Tetrazo- Colouring Matters]
- Sir DUGALD CLERK, K B.E , LL D., F.R.S. [EVAPORATION.]
- Dr. JULIUS B COHEN, FR.S., Professor of Organic Chemistry in the University of Leeds. [Diphenylamine, Diphenylmethane.]
- W. H. COLEMAN, Esq. [CARBOLIC ACID, CREOSOTE.]
- JAMES CONNAH, Esq., B.A., B.Sc., FI.C., Deputy Principal, The Government Laboratory, London [Chartreuse, Chocolate; Cocoa, Coffee, Cordials and Liqueurs, Denaturants.]
- C. F. CROSS, Esq., B.Sc, F.I.C., F.R.S. [CELLULOSE]
- Sir JAMES J. DOBBIE, M.A., D Sc, LL D, F.R S, late Principal of the Government Laboratory. [Corydaline and the Corydalis Alkaloiis.]
- Dr. THOMAS EWAN, M Sc. (Cassel Cyanide Co., Ltd., Glasgow). [Cyanides ]
- Dr. ARTHUR J EWINS (Messrs. May & Baker, London). [CHLOROFORM; CYTISINE; DAMASCENINE.]
- SAMUEL FIELD, Esq, Northampton Polytechnic Institute, Clerkenwell, London. [Electrodeposition and Electroplating.]
- Dr. J. A. NEWTON FRIEND, F.I.C [CORROSION OF METALS.]
- W. M. GARDNER, Esq., M.Sc., Dyeing Department, Technical College, Bradford. [Dyeing; Dye-stuffs, Identification of, on Fabrics.]
- JOHN GOLDING, Esq., D.S.O., F.I.C., University College, Reading. [Cheese.]
- Dr. J. N. GOLDSMITH, M.Sc., F.I.C. [CELLULOID.]
- Dr. ERNEST GOULDING, F.I.C., Imperial Institute, South Kensington. [Cotton.]
- SIR ALFRED D. HALL, K C.B, F R S., Board of Agriculture. [Cereals; Crops]
- Professor G. G. HENDERSON, M.A., D.Sc., LL.D., F.R.S., The University, Glasgow. [Catalysis in Industrial Chemistry.]
- T. H. P HERIOT, Esq , F.I.C., Ronal Technical College, Glasgow. [Decolourising Carbons]
- J. W. HINCHLEY, Esq., A R.S M, Professor of Chemical Engineering, Imperial College of Science and Technology South Kinsington. [CRYSTALLISATION.]

- GEORGE T. HOLLOWAY, Esq., A.R C.S., F.I.C. [COLUMBIUM.]
- BERNARD F. HOWARD, Esq., and OLIVER CHICK, Esq. (Messrs. Howard & Sons, Ilford). [CINCHONA ALKALOIDS.]
- Dr. HARRY INGLE. [CORK.]
- HERBERT INGLE, Eq., B.Sc., FI.C., late Chief Chemist to the Transvaal Agricultural Department. [Carrot; Cassava; Castor Seeds; Celery; Cherry; Chestnut, Chick Pea; Chicory; Clover; Cranberry; Cucumber; Date; Egg Fruit; Elderberry]
- HAROLD KING, Esq., M.Sc. [CAMPHOR.]
- VIVIAN B. LEWES, Esq , F.I.C. [CORROSION AND FOULING OF STEEL AND IRON SHIPS AND THEIR PREVENTION.]
- Dr. JULIUS LEWKOWITSCH, M.A, FI.C. (Revised by C. A. Mitchell, Esq., M.A., F.I.C.) [Cameline Oil; Candles, Candle-nut Oil, Castor Oil, Chaulmoogra Oil Group, Chinese Vegetable Tallow; Chocolate Fats, Coco-nut Oil Group; Cod Liver Oil; Cotton Seed Oil; Curcas Oil, Dégras or Sod Oil; Dika Fat Group; Driers, Edible Oils and Fats.]
- WILLIAM D. McCREATH, Esq (The Quantock Vale Cider Co., Ltd., North Petherton, Bridgwater). [Cider.]
- CHARLES A. MITCHELL, Esq., M.A , F.I.C. [CANDELILLA WAX; COPYING OR INDELIBLE INK PENCILS.]
- PROFESSOR GILBERT T. MORGAN, O B.E., D.Sc, F.R.S, Professor of Chemistry in the University of Birmingham. [Cerium; Cerium Metals and Earths; Diazo-Compounds; Dysprosium; Erbium, Europium.]
- L. MYDDELTON NASH, Esq , F.I.C. [CASEIN.]
- Professor A. G. PERKIN, F.R.S., F.I.C., Department of Color r Chemistry in the University of Leeds. [Camwood; Catechu, Catellagic Acid; Cedrela Toona, Chamomile Flowers, Chay Root; Chestnut Extract, Chica Red and Carajura, Chromone; Clover Flowers; Coprosma; Coto Bark; Cotton Flowers; Cotton Seed, Colouring Matter of, Courmaranone Group, Cyanellagic Acid; Daphnetin, Datisca Cannabina; Delphinium Consolida, Diphenyl-dimethylolid, Divi-Divi, Drosera Whittakeri; Dyer's Broom; Ellagic Acid, Ellagitannic Acid, Eriococcus Coriaceus; Erythrin; Evodia Meliafolia]
- Professor A. G. PERKIN, F.R.S., and Dr. ARTHUR E. EVEREST. [Cochineal, Evernic Acid and Everninic Acid.]
- Professor J. C. PHILIP, O.B E., D Sc , F.R.S., Imperial College of Science and Technology. [Chemical Affinity; Colloids; Cryoscopy, Diffusion.]
- Dr. SAMUEL RIDEAL, FI.C., and Dr. ERIC K RIDEAL, M.B.E., M.A., F.I.C. [DISINFECTANTS.]
- Dr. F. W. RUDLER, I.SO. [ENAMELS.]
- Dr. SYDNEY SMITH, B Sc , F.I.C. (Wellcome Chemical Works, Dartford). [Cocaine and the Coca Alkaloids]
- L. J. SPENCER, Esq., M.A, F.G.S, Mineral Department, British Museum. [Carbonite, Carclazite, Carnallite, Carnotite, Cassiterite, Catapleite, Celestite; Cerargyrite, Cerite, Cerussite, Chalcanthite; Chalcedony, Chalk, Chalmersite, Chalveite; China-Stone; Chiolite, Chloanthite, Chronite, Chrysoberyl, Chrysocolla; Chucchite; Cinnabar; Cinnamon Stone; Clayite; Cobaltite; Colemanite; Collophanite; Columbite; Copper-glance; Copper Pyrites; Cordylite; Crudum; Covellite; Cristobalite; Crocidolite; Crocoite, Crookesite; Cryolite; Cryolithionite, Cuprite, Cylindrite, Danalite; Datolite, Descloizite, Diabase; Diallage; Diamond; Diaspore, Diatomite; Dietzeite, Diopside, Dioptase; Diorite; Dolerite; Dolomite; Domeykite; Dysanalyte; Dyscrasite; Emboyite, Emery; Enstatite; Epidorite; Epidote, Epsomite; Erythrite; Eucairite; Euclase; Eucryptite; Euclalyte; Euxenite.]
- Dr. J. SUDBOROUGH, F.I.C., Professor of Chemistry in the Indian Institute of Science. [Esterification.]
- Professor JOCELYN F. THORPE, C B.E., F.R.S., Imperial College of Science and Technology, South Kensington. [Diphenyl.]
- Professor EDWIN R. WATSON, M.A., D.Sc., Dacca College, East Bengal. [COLOUR AND CHEMICAL CONSTITUTION]
- Dr. M. A. WHITELEY, O.B.E., F.I.C., A.R.C.S., Assistant Professor of Organic Chemistry, Imperial College of Science and Technology, South Kensington. [Carnine; Carnitine; Carnosine; Creatine; Creatine; Cystine; Dialuric Acid; Dilituric Acid]
- Dr. SYDNEY YOUNG, Sc.D., F.R.S, F.I.C., Professor of Chemistry, Trinity College, Dublin [Distillation.]

### DICTIONARY

OF

### APPLIED CHEMISTRY.

CALAFATITE v. ALUNITE.

CALCULI v. URINE.

CALCUSOL. Trade name for a preparation of acid potassium carbonate and piperidine-p-

sulphamine benzoate

CALIATURWOOD. This wood, the botanical origin of which appears to be unknown, is very similar to though somewhat darker in colour than sanderswood It was imported from the East Indies and is stated to have been chiefly employed on the Continent According to Franchimont and Sicherer it contains santalin but in larger amount than sanderswood.

CALICHE. The term by which the impure native sodium nitrate or Chile saltpetre is known throughout South America. Also applied to a limestone hardpan found in Arizona and else-

where (v Sodium)

CALLISTEPPIN v ANTHOCYANINS. CALLUTANNIC ACID v TANNINS. CALMATAMBIN v. GLUCOSIDES.

CALMONAL. Calcium methylbromide. CALOMEL. Mercurous chloride (v. MERCURY).

CALOMELOL. A colloidal preparation of

CALOPHYLLUM INOPHYLLUM (Linn.). A tree (ord Gutisferæ), Alexandrian laurel, dilo of Fiji, tamarınd of E. Polynesia, and sultana champa (Hind.) of India), the seeds of which yield 60 p c. of a fragrant green oil, fluid at ordinary temperatures, solidifying when cooled below 10°. The tree bears fragrant white flowers. The oil, the woondel or bitter oil of Indian commerce, used in medicine and for lamp oil, is of a greenish tint, and a very little imparts its colour to a whole cask of coco-nut oil (Pharm. J. [3] 8, 363).

CALOTYPE v. PHOTOGRAPHY.

Vol. II.—Т.

transversely and dried. It was introduced into Europe in the 17th century as an antidote for poison, and found a place in the London Pharmacopæia of 1788. It possesses mild bitter tonic properties, for which purpose it is now chiefly employed in medicine.

The drug contains 30-35 p c. of starch, some mucilage, traces of a volatile oil, but no tannin; the root gives 4-7 p.c., the rhizome 12-17 p.c of ash. There are further present two bitter

principles and three alkaloids

Columbia  $C_{28}H_{32}O_{9}$  (according to Ulrich, Annalen, 1907, 351, 363; but cp Feist, lc.), the best-known bitter principle, occurs to the extent of 08 pc., and is extracted by hot ether and recrystallised from the same solvent. It forms colourless prisms, mp 182°, hardly soluble in hot or cold water, little in cold alcohol or ether, readily in hot ether, chloroform, and alcohol. Columbin adds directly 8 bromine atoms. A crystalline diacetyl derivative melts at 218°. Bolling 5 pc potassium hydroxide yields columbic acid C<sub>28</sub>H<sub>32</sub>O<sub>10</sub>, crystals, m.p 220° (Frey, Annalen, 1907, 351, 372), bolling 15 pc. hydrochloric acid gives an amorphous yellow substance and yellowish green fluorescence. A second bitter principle, m.p. 246°, almost insoluble in most solvents, was isolated by Feist (Arch. Pharm 1907, 245, 586).

The alkaloids were at first mistaken for berberine, to which they are closely related (Gordin, Arch. Pharm. 1902, 240, 146; Gunzel, ibid. 1906, 244, 257; Gadamer, ibid. 1902, 240, 450; and particularly Feist, ibid. 1907, 245, 586). All three yield yellow quaternary salts The aqueous solution of an alcoholic extract of the drug is extracted with ether to remove columbin (see above), and concentrated potassium iodide then precipitates the iodides, which CALUMBA. Calumbæ radæ, B.P.; Calumba,
U.S.P. Colombo, Columbo. (Racene de Colombo,
Fr.; Kalumba or Columbowurzel, Ger.)
The root of the Jateorrhiza Columba (Miers),
a • herbaceous climbing plant, inhabiting the
forests of Eastern Africa (Fluck. a. Hanb. 23;
Bentl. a. Trim. 13). The fleshy root is sliced

2 CALUMBA

groups and one hydroxyl group; it is a methyl ether of the second, which contains three methoxy and two hydroxyl groups, for methylcolumbamine and dimethyljateorrhizine are Palmatine iodide contains four methoxyl groups, but no hydroxyl, and hence remains undissolved on treating the mixed iodides with sodium hydroxide. Its exact relationship to the other two is unknown Reduction of the corresponding nitrates leads in all three cases to colourless tetrahydio derivatives. Methylcolumbamine is oxidised by permanganate to corydaldine (an oxydation product of corydaline, qv), a trimethoxy-o-phthalic acid (gallo-earboxylic acid?), mp 200°, and a nitrogenous acid, mp 200°-202° The free bases corresponding to the above salts are unknown, so that it is doubtful whether the alkaloids are ammonium hydroxides or the corresponding anhydrides

For the identification of columbin and the alkaloids under the microscope, see Tunmann, Pharm Zentr H. 1914, 55, 775 G B

CALYCANTHINE C<sub>11</sub>H<sub>14</sub>N<sub>2</sub>, crystals, mp 243°-244° (dry), occurs with isocalycanthine, prisms, mp 235°-236° (dry), in the seeds of Calycanthus glaucus (Wild), 2 p.e. One mitrogen is secondary, the other tertiary. Toxic (Gordin, J. Amer. Chem Soc 1905, 27, 144, 1418; 1909, 31, 1305; 1911, 33, 1626)

CAMELINE OIL, Dodder oil, or German

CAMELINE OIL, Dodder oil, or German Sesamé oil, is prepared from the seeds of Myagrum sativum (Crz.), Camelina sativa (Linn), ('gold of pleasure') Several centuries ago the seed was grown all over Germany as a crop, but at present it is only cultivated on a small scale in some part of South Germany, Belgium, Holland, Hungary, the Balkan States, and South Russia. The seed is occasionally expressed in Hull, but, as a rule, the press cakes arrive in this country from Odessa, and are used (in Liverpool) in the manufacture of 'compound cakes' The average composition of cameline seed is stated as follows.—

Oil			-294 рс
Water			75,
Proteins			25 9 ,,
Carbohyd	rates		173,
Crude fib	e		105,
$\mathbf{A}\mathbf{s}\mathbf{h}$			92,

The oil has a golden-yellow colour and a pungent taste and smell. On exposure to the air it dries slowly. It is a semi-drying oil, and therefore yields, on boiling with litharge or manganese, a slowly drying varnish. The low saponification value of the oil points to the presence of crucic acid

The 'cold-drawn' oil is used for edible purposes in Russia. The oil expressed in this country, as also the oil expressed by heat in Hungary and the south of Russia, is used for soap making

It has the following analytical characters sp gr 0 9224-0 9260 at 15°; solidification point, -15°-19°; saponification value, 185-188; iodine value, 133-136; and mp. of fatty acids, 18°-20°; molecular equivalent of fatty acids, 296 A sample examined by Grimme (Chem Rev. Fett Ind 1912, 19, 102) contained 1 16 pc of unsaponifiable matter According to van Ketel and Antusch (Zettsch.

angew Chem 1896, 581), linseed is liable to be contaminated with cameline seed, and so may yield an oil with a low iodine value. J. L

CAMPHOID. A preparation of one part of guncotton dissolved in 20 parts each of absolute alcohol and camphor.

alcohol and camphor.

CAMPEACHY WOOD v. LOGWOOD
CAMPHANIC ACID v. LACTONES.
CAMPHENE, CAMPHINE v. TERPENES.
CAMPHOL v. CAMPHORS.

CAMPHOPHORONE v. KETONES.

CAMPHOR. The term camphor was applied exclusively, in the sixth century, by Arabian and Chinese writers, to Borneo or Malayan Camphor (Borneol), but in the sixteenth century it included Laurel or Common Camphor, for Garcia d'Orta mentions the latter as well known in Europe in 1563, and points out what indeed is true at the present day, that the much more costly Borneo or Sumatra ('amphor is entirely consumed in the East (cf Fluck a Hanb 510) The similarity in properties between laurel camphor and the crystalline separations from the volatile oils of plants, led, in the eighteenth century to an extension of the term camphor so as to include these solids. An attempt by Berzelius (Lehrbuch VI 580) to simplify the nomenclature, by calling all solid portions of volatile oils, 'stearoptenes,' and the liquid portions 'clæoptens' has only resulted in the ' continued use, almost synonymously, of the words stearontene and camphor. The discovery that laurel camphor contains oxygen has still further amplified the meaning of the word, which in its most comprehensive sense now includes most of the oxygenated products, both liquid and solid, of volatile oils.

Camphors are volatile oxygenated hydrocarbons, usually possessing a peculiar characteristic odour. They are mostly crystalline solids insoluble in water, but soluble in alcohol Except in the case of a few which are produced artificially they are obtained from plants where they occur associated with hydrocarbons from which they are probably derived. Most plants when distilled in steam yield volatile oils composed of heavier oxidised compounds held in solution more or less by lighter hydrocarbons. The former often separate in crystals during the condensation of the oil or afterwards on standing.

or they may be obtained by fractional distillation. The more important camphors, using the term in its larger signification, may be considered, for convenience, in 4 groups, as isomerides or homologues of: (1) Peppermint camphor, menthol  $C_{10}H_{20}O$ ; (2) Borneo camphoi, borneol  $C_{10}H_{18}O$ ; (3) laurel camphor  $C_{10}H_{16}O$ ; and (4) thyme camphor, thymol  $C_{10}H_{16}O$  For the most part camphors behave like alcohols or phenols, but laurel camphor and menthone are ketones, cineol is an ether and alant camphor The majority of camphors belong to a lactone saturated or unsaturated cyclic systems, but in 1891 Semmler (Ber 24, 210) introduced the expression 'olefinic camphors' for certain substances which possess an open chain structure, are closely related to the cyclic camphors, and are responsible for the aroma of many essential oils. The chief members of this group are citronellol C<sub>10</sub>H<sub>20</sub>O, geraniol, nerol and citronellas C10H18O, and citral C10H16O. (See l also Perfumes.)

Peppermint camphor. Menthol C<sub>10</sub>H<sub>19</sub>OH. le-6-180propyl-1-cyclohexanol. The lævo 3-Me-6-isopropyl-1-cyclohexanol. modification of this camphor is the chief con stituent of peppermint oil, the product of distillation of the herbs Mentha piperita (Linn) and M. arvensis (Linn), varieties piperascens and glabrata. It is also a minor constituent of a few other volatile oils. A menthol of unrecorded rotatory sign predominates in the oil from Hyptis suaveolens (Linn) (Bacon, Phillipine J. Sc. 1909, 4, A, 130). The proportion of menthol n normal Japanese oil of peppermint varies between 69 and 91 pc, whilst English and American oils contain 48 to 68 pc A portion, about 10 p c., of the menthol is present in ester form combined with acetic and isovalerianic acids The camphor is extracted by freezing oils rich in menthol, whereby the crystals separate and are recovered either by pressure or by centrifugal treatment. The more important papers dealing with menthol in its bearing on oil of peppermint are: Beckett and Wright (Chem. Soc Trans. 1876, 29, 1), Atkinson and Yoshida (ibid. 1882, 41, 50), Moriya (ibid. 1881, 39, 77), Power and Kleber (Arch. Pharm. 1894, 232, 639), Power and Fluckiger (Pharm J 1880 [3] 11, 220), Andres and Andred (Ber 1892, 25, 609), Schindelmeiser (Apoth Zeit 1906, 21, 927), and the semi-annual reports of Schimmel and Co, and of Roure Bertrand Fils.

Menthol crystallises in prisms belonging to the hexagonal system, and in appearance resembles magnesium sulphate It has a strong peppermint odour and taste, and produces a sensation of warmth on the tongue which is replaced by a sensation of cold when air is replaced by a sensation of cold when all is inhaled M p 42°-43°, 43 5°-44 5° (Gildemeister and Hoffmann, Die Atherischen Ole, 1910, i. 405), b p 215 5° at 758 mm (P. a K.), sp gr. 0 88 at 20° (Long, J Amer Chem. Soc 1892, 14, 151),  $[\alpha]_{\rm D}^{20^{\circ}} -50$  7° in 10 p c alcoholic solution (Beckmann, J. pr Chem 1897, [2] 55, 15). Menthol is very slightly soluble in water, but soluble in the usual organic solvents and in concentrated hydrochloric acid. It rotates on the surface of water in a similar manner to laurel camphor Mixed with chloral, thymol, or laurel camphor, menthol reacts forming a liquid (Kyle, Amer J Pharm 1885, 429; Becker, bul. 1886, 283) When menthol is boiled with sulphuric acid diluted with half its volume of water it acquires an indigo-blue or ultramarine colour, the acid becoming brown (Brit. Pharm

Menthol is a saturated secondary alcohol which under the influence of dehydrating agents, e.g. phosphorus pentoxide, zinc chloride, potassium bisulphate, passes into the hydrocarbon menthene, C<sub>10</sub>H<sub>18</sub>. A very large number of esters of menthol are known, of which the acetate C<sub>10</sub>H<sub>19</sub>OAc, b p 227°-228°, [a]<sub>D</sub>-79 26°, (cf. Kishner, J. Russ Phys. Chem Soc. 1895, 27, 480), and isovalerianate, C<sub>10</sub>H<sub>19</sub>OOC C<sub>4</sub>H<sub>9</sub>, occur in oil of peppermint (P. a. K.). For the identification of l-menthol by means of derivatives, the phenylcarbamate C<sub>10</sub>H<sub>19</sub>OCONHPh, m.p. 111°-112° (Leuckart, Ber 1887, 20, 115), the hydrogen phthalate C<sub>10</sub>H<sub>19</sub>OCC C<sub>7</sub>H<sub>5</sub>O<sub>3</sub>, m.p. 122° (Pickard and Littlebury, Chem. Soc. Trans. 1912, 101, 118), and benzoate C<sub>10</sub>H<sub>9</sub>OBz, m.p. 55°-56° (P. a. L.) are characteristic.

Two atoms of hydrogen are removed from 1-menthol by treatment with chromic acid mixture (Beckmann, Annalen, 1889, 250, 325) with formation of 1-menthone, a liquid ketone which occurs to the extent of about 20 pc. in oil of peppermint (B a W.), and with an isomeride, d-isomenthone, forms 50 pc of American pennyroyal oil obtained from Hedeoma pulegoides (Linn) (Barrowcliff, Chem. Soc. Trans. 1907, 91, 875). Menthol has been obtained synthetically from menthone and pulegone (Beckmann and Pleissner, Annalen, 1891, 262, 32, Haller and Martine, Compt. 1end 1905, 140, 1298), also from thymol by use of the Sabatier and Senderens method of reduction By this latter method the four possible menthols corresponding to menthone, namely, d- and l-menthol and dand l-neomenthol, have been prepared (Pickard and Littlebury, Chem Soc Trans. 1912, 101, A small proportion of d-neomenthol was isolated at the same time from Japanese oil of peppermint.

In its medicinal action menthol resembles phenol—Its chief uses are as a local anæsthetic for relief of superficial pains of various kinds, also dissolved in oil as a spray for influenza, hay fever, &c., and as a local application in

certain cutaneous diseases.

Camphors isomeric with menthol Citronellol  $C_{10}H_{20}O$ , a liquid isometide of menthol occurs as l-citronellol in rose oil, Rosa damascena (Mill) (Tiemann and Schmidt, Ber 1896, 29, 922), and as d-citronellol in Java citronella oil, Andropogon Nardus Java de Jong (Schimmel's Rept 1902, 114), and in the oil from Barosma pulchella (Linn) (ibid 1910, 117), whilst both forms occur in geramium oil, Pelargonium graveolens (Ait.) (Tiemann and Schmidt, Ber 1896, 29, 924).

Borneo and Allied Camphors. Malayan, Barus or Dryobalanops camphor; Borneol, camphol C<sub>10</sub>H<sub>1.7</sub>OH, 1, 7, 7,-trimethyl [1, 2, 2]-bicyclo 2-heptanol This camphor occurs in nature in the dextro- and levorotatory varieties, esterified chiefly in the lavorotatory modification Common borneol, which is dextrorotatory, is the product of Dryobalanops aromatica (Gartn), a majestic tree indigenous to Sumatia, Borneo, and Labuan. The camphor is deposited in fissures in the old wood, and to obtain it the The tree is cut down trees have to be sacrificed and the wood is examined by a laborious process of splitting The collected camphor averages 3 to 11 lbs from each tree Owing to the reck-less manner in which the trees have been destroyed without the planting of others, the forests of Sumatra now contain few that are worth working (Pharm. J. 1881, [3] 12, 83). The states of Sarawak and Brunei are much richer in trees, and at the present time produce borneol by similar methods (cf Hanb. and Fluck 516; Kremers, Pharm Review, 1905, 23, 7; Gildemeister and Hoffmann, Die atherischen Öle, 1916, iii 178) The crude camphor is best purified by sublimation (Pelouze, Annalen, 1841, 40, 326).

d-Borneol is also a constituent of the following volatile oils: Rosemary, Rosmarinus officinalis (Linn.) (Gildemeister and Stephan, Arch. Pharm. 1897, 235, 586); Siam Cardamom Amomum cardamomum (Linn.) (Schimmel's Rept. 1897, ii. 9), Nutmeg, Myristica fragrans (Houtt.) (Power and Salway, Chem. Soc. Trans.

1907, 91, 2037); Lavender, Lavandula vera (D.C.) (Bruylants, J. Pharm. Chim. 1879, [4] 30, 139; cf. Charabot, Bull. Soc. Chim. 1897, [3] 17, 380); Spike Lavender, Lavandula Spica (Linn.) (Bouchardat, Compt rend 1893, 117, 53 and 1094), and Virginian Snake-root, Aristolochia serpentaria (Linn.) (Spica, Gazz. chim. ital. 1887, 17, 314). In addition to these natural sources d-borneol may be obtained, by the action of reducing agents on laurel camphor when borneol, of the same sign of rotation as the camphor used, is obtained mixed with a small proportion of tsoborneol of opposite sign (Haller, Compt. rend. 1887, 105, 227; Beckmann, J. pr. Chem. 1897, [2] 55, 31), and in small proportion with l-borneol it is found among the products of the distillation of amber with potash (Berthelot and Buignet, Annalen, 1860, 115, 245)

Borneol is harder and less volatile than laurel camphor, and does not sublime in the bottles in which it is kept. It has a peppery camphoraceous odour and burning taste M p. 203°-204° (preparations containing isoborneol melt 206°-208°), b p. 212°, sp. gr. I 011 (Plowman, Pharm. J. 1874, [3] 4, 711). It crystallises in the hexagonal system (Traube, J. pr. Chem. 1894, [2] 49, 3); [a]<sub>D</sub> +37.7° (Beckman, Appeler, 1899, 250, 252; Heller, Competer, 253, 254; Heller, Competer, 254, 254; Heller, Competer, 254, 254; Heller, Competer, 255, 255; Heller, Competer, 255, 255; Heller, Competer, 255; Heller, 255; Hell mann, Annalen, 1889, 250, 353; Haller, Compt rend 1891, 112, 143; Pickard and Littlebury, Chem Soc. Trans. 1907, 91, 1973). Borneol is a saturated secondary alcohol, and on oxidation yields camphor,  $C_{10}H_{16}O$  of the same rotatory sign. Powerful exidising agents produce camphoric acid C<sub>10</sub>H<sub>18</sub>O<sub>4</sub>, and further degradation products (Laurent, Annalen, 1837, 22, 135, Pelouze, *ibid.* 1841, 40, 328; Kachler, *ibid.* 1878, 191, 143) Removal of a molecule of water by phosphoric anhydride gives rise to a solid terpene, camphene, C<sub>10</sub>H<sub>16</sub> (Wallach, *ibid*. 1885, 230, 239); for the mechanism of this change, see Meerwein, ibid. 1914, 405, 129) Concentrated hydrochloric acid (Berthelot, ibid 1859, 112, 366), or phosphorus pentachloride (Kachler, *ibid* 1879, 197, 89) reacts with borneol producing *bornyl chloride* C<sub>10</sub>H<sub>17</sub>Cl, a product identical with pinene hydrochloride or 'artificial camphor.' For the characterization of borneol the phenylcarbamate, C<sub>10</sub>H<sub>1</sub>,0CONHPh, mp 138°-139° (Bertram and Walbaum, J pr Chem 1894, [2] 49, 5), the acetate C<sub>10</sub>H<sub>17</sub>OAc, m p. 29° (B. and W. Arch Pharm. 1893, 231, 304), and the addition products with bromal, mp 105°-108° (Minguin, Compt rend 1893, 116, 889), and chloral, mp 55°-56° (Haller, ibid 1891, 112, 145), are employed

l-Borneol is a constituent of the following oils Valenan, Valeriana officinalis (Linn) (Bruylants, Ber. 1878, 11, 452, Ohiviero, Bull, Soc Chim 1895, [3] 13, 917), Canadian Snakeroot, Asarum canadense (Linn) (Power and Lees, Chem Soc Trans 1902, 81, 63), Arborvitæ, Thuja occidentalis (Linn) (Wallach, Annalen, 1907, 353, 213), Citronella, Andropogon nardus (Linn) (Schimmels, Rept. 1912, 1, 43), Pinus maritima (Mill) (Belloni, Chem Zentr 1906, 1, 1552), Yellow Pine, Pinus palustris (Mill.) (Schimmels, Rept. 1910, 1, 106), and Feverfew, Matricaria Parthenium (Linn.) (ibid. 1894, 1, 71). It is also contained in the fusel oil of the crude spirit obtained by the fermentation of the sugar of madder, and formerly known

as madder camphor (Jeanjean, Annalen, 1857, 101, 95), and in Ngai camphor, the name given to the solid part of the essential oil of Blumea balsamifera (D. C.) (Plowman, Pharm. J. 1874, [3] 4, 712; Fluckiger, ibid. 828; Schimmels, Rept 1910, 1, 149). From this latter source the l-borneol had a higher rotatory power  $[\alpha]_{\rm D}$  — 39° 25′, than is usually observed (Schimmels Rept 1895, 1, 74). On oxidation with intricacid l-borneol yields levorotatory camphor.

Numerous esters of borneol are known. Those of l-borneol, which occur in nature, are bornyl acetale, in oil of valerian (Bruylants), Siberian fir oil, Abies sibirica; Ledeb (Hirschsohn, Chem Zentr 1892, 11 793), hemlock spruce oil, Abies canadensis; Michx. (Hunkel, Pharm. Review, 1896, 14, 35), and coriander oil, Coriandrum sativum (Linn) (Walbaum and Muller, Wallach Festchrift, 1909, 654), bornyl formate, butyrate and isovalerianate in oil of valerian (Bruylants).

Inactive borneol, the racemic mixture of d- and l-borneol has been obtained by the distillation of the crude product of the action of sulphuric acid on turpentine (Armstrong and Tilden, Chem. Soc Trans 1879, 35, 752); an ester of i-borneol is also obtained by treating pinene with oxalic acid (Schindelmeiser, J Rus. Phys Chem Soc 1902, 34, 954).

Isoborneol, a stereoisomeride of borneol and an intermediate product in the manufacture of synthetic camphor, was first obtained by Bertram and Walbaum (J pr Chem 1894, [2] 49, 1) from camphene obtained by the action of sodium acetate and glacial acetic acid on pinene hydrochloride at 200°. Camphene is warmed with glacial acetic acid and a little, 50 pc, sulphuric acid, and the resulting acctate is decomposed by alcoholic potash. Isoborneol can now be prepared from pinene hydrochloride directly, by the action of boiling glacial acetic acid in the presence of a small quantity of a metallic salt such as anhydrous zinc chloride (Weizmann and Clayton Aniline Co, Eng Pat. 8266, 1906) It is also formed to a small extent in the reduction of laurel camphor (Beckmann, J pr Chem 1897, [2] 55, 35) It differs from borneol, physically, in its greater volatility, its higher m p 212° (in sealed tube), and its greater solubility in benzene and light petroleum, chemically in its greater stability towards oxidising agents, and in the ease with which it loses water forming camphene. Its bromal addition compound melts at 71°-72°. The optically active isoborneols melt at 214°, and have [α]<sub>n</sub>±34 3° (Pickard and Littlebury, Chem Soc. Trans. 1907, 91, 1979).

Camphors isomeric with borneol. Cineol, Cajeputol or Eucalyptol C<sub>10</sub>H<sub>10</sub>O, anhydro 1-Me 4 (a-Me a-oxyethyl) 1-oxyeyclohexane, is a constituent of a great number of volatile oils of Eucalyptus globulus (Labill.) (Jahne, Ber. 1884. 17, 2941), Wormseed, Artemisia maritima (Linn.) var. Stechmann (Wallach and Brass, Annalen) 1884, 225, 291, Hell and Sturcke, Ber. 1884, 17, 1970); Cajeput, Melaleuca Cajeputi (Roxb.) (Wallach and Gildemeister, ibid. 1888, 246, 276); M. viriaflora de Brogn. et Gris (Bertrand, Bull. 820. Chim. 1893, [3] 9, 432). Cincol melts at 1°-15° (Scnimmel & Co.); bp. 1/6°

(Wallach, Annalen, 1888, 245, 195) It has a camphor-like odour and is optically inactive. It is characterised by its addition compounds. with the halogens, with hydrobromic acid m.p. 56°, which also serves for the isolation of small quantities (Wallach, Annalen, 1888, 246, 280), with phosphoric acid (Scammell, D. R. P. 80118), with arsenic acid (US Pat. 705545, Thoms and Molle, Arch Pharm 1904, 242, 172), with iodole (tetraiodopyirol), m.p 112° (Bertram and Walbaum, Arch. Pharm. 1897, 235, 178), by means of which very small quantities can be detected (Hirschsohn, Chem Zentr. 1893, 1. 503), and with resorcinol (Baeyer and Villiger, Ber 1902, 35, 1209). The compounds with hydrobromic acid, phosphoric acid, arsenic acid, and resoremol are used for its quantitative estimation (cf Turner and Holmes, Amer J Pharm. 1915, 87, 101), the question as to their relative efficiency being still a matter of dispute. Dehydrating agents convert cineol into dipentene C<sub>10</sub>H<sub>16</sub> (Wallach and Brass, Hell and Sturcke). Oxidation by permanganate gives cincolic acid C<sub>10</sub>H<sub>16</sub>O<sub>5</sub>, m p 197 (Wallach and Gildemeister), but towards reducing agents it is very stable The oxygen atom in cincol is present as an oxide or ether linking. The aitificial production of cincol has been effected by the action of dilute acids on terpineol or terpin hydrite (Wallach, Annalen, 1887, 239, 18).

Cincol is used in medicine as a local antiseptic diessing, and for catarrhal conditions of

the respiratory mucous membranes.

The volatile oils of the following contain other isomerides of borneol; Citionella, Andropogon Nardus (Linn.) (Semmler, Ber 1893, 26, 2254) and Eucalyptus cutrodora (Hook) (Schimmels, Rept 1907, 2, 35) contain d-Citronellal, whilst 1-Citronellal is contained in 'Java lemon oil' (ibid 1903, 1, 21). Linaloe, Ocotea caudata (Mez.) (Barbier, Compt 1end 1895, 121, 168), Lavender, Lavandula vera (D.C.) (Bertram and Walbaum, J. pr. Chem. 1892, [2] 45, 590); Bergamot, Citrus Aurantium (Linn) (Bertram and Walbaum, ibul 602) contain l-linalol, whilst d-linalol ('corundrol') is contained in oil of coriander, Coriandrum sativum (Linn) (Barbier, Compt. rend. 1893, 116, 1460). Geranium, Pelargonium graveolens (Ait.) (Bertram and Gildemeister, J. pr Chem. 1894, [2] 49, 191), Citronella, Andropogon Nardus (Linn) (Schimmels, Rept. 1893, 2, 12); Lemon grass, Cymbopogon flexuosus (Stapf) (Tie-mann and Semmler Ber 1893, 26, 2721), contain geraniol ('rhodinol') Neroli oil from Citrus Aurantium (Linn), subspec. amara (Linn) (Hesse and Zeitschel, J. pr. Chem. 1902, [2] 66, 481) contains nerol. d-a-Terpineol, m p. 37°-38°, is a constituent of oil of sweet orange, Citrus Aurantium (var) dulce (Linn.) (Stephan, J. pr. Chem. 1900, [2] 62, 523) of neroli and petit-grain oils; 1-a-terpineol of oil of camphor, Cinnamomum Camphora (Nees et Eberm.) (Schimmels, Rept. 1903, 2, 39); 1-Terpineol, m p. 35°, of caleput oil, Melaleuca cajeput (Roxb) (Voiry, Bull. Soc. Chim. 1888, [2] 50, 108)

Common and allied camphors. Laurel Camphor; Dertro Camphor. (Camphre, Fr; Campher, Ger.) C<sub>10</sub>H<sub>10</sub>O 1.7.7-Trimethyl [1-2]-bicyclo 2-heptanone. Common camphor is contained in all pa ts of the camphor la rel, Ci immomum

Camphora (Nees et Eberm), a tree inhabiting Japan, Formosa and Southern China It occurs also as a constituent of the volatile oils of Spike, Lavandula spica (D.C.) (Lallemand, Annalen, 1860, 114, 198); Rosemary, Rosmarinus officinalis (Linn) (ibid. 197); Sage, Salvia officinalis (Linn) (Mur, Chem Soc. Trans. 1880, 37, 678); Siam Cardamom, Amomum Cardamomim (Linn) (Schimmels, Rept. 1897, 2, 9); American Wormseed, Chenopodium ambrosoides (Linn.) var anthelminthuum (Gray) (ibid. 1908, 1, 112), and Sassafras, Sassafras officinalis (Nees) (Power and Kleber, Pharm Review, 1896, 14, 101)

Although camphor is at the present time still manufactured artificially, most of the camphor of commerce is obtained from the camphor laurel, the chief industry being confined to Japan. Formosa and China The fined to Japan, Formosa and China experimental plantations which were undertaken for the production of natural camphor m Florida, California, Jamaica, East Africa, Malay States, Algeria and Ceylon have at least in one case, that of Florida, begun to furnish camphor on the commercial scale (Oil, Paint and Drug Rep., March 29, 1915) In Formosa the camphor industry before the China-Japanese war was nominally in the hands of the Chinese, but owing to the savage state of the interior of the island, and the unscrupulous methods of the dealers, it had declined to small and unimportant proportions Afterwards Japan assumed the control of the industry, and it was made a government monopoly in 1899, and this was extended to Japanese camphor in Since the establishment of the monopoly the old crude methods of obtaining the camphor (Fluck. and Hanb 513) have been replaced by more modern ones. The camphor wood in the form of chips is steamed in roughly built furnaces, and the camphor which is driven off is condensed in cooled wooden vessels, and freed from most of the oil by draining, crude of 'grade B' camphor being thus obtained. For the production of refined or 'grade A' camphor, the crude material is heated in large non retorts, through which a current of air is passed the first 48 hours only sufficient heat is applied to drive off the water and oil. The retoits are then connected with a condensing chamber, the roof of which is cooled by running water, and are heated to a high temperature in order to volatilise the camphor. It is then formed into blocks in wooden moulds and pressed, first by steam power and finally by a very high hydraulic pressure (Collins, J. Soc. Chem. Ind. 1904, 75) The yield of camphor is greater in winter than in summer, while the reverse is true of the oil. In summer a charge of 120 kilos, of wood gives 2.4 kilos, or 2 p c. of camphor, and 18 litres of oil; in winter, 3 kilos., or 2.4 p.c. of camphor and 0 5-0.7 litre of oil.

Crude camphor appears in commerce as a coarse wet powder of a greyish-white colour Chinese crude camphor is packed in round wooden pots of about 150 lbs net content, Japanese in lead-lined wooden boxes containing about 130 lbs. net. The principal impurities of crude camphor are water, camphor oil, iron, sand, wood, &c. Light oil of camphor has a sp gr. of 0.895-0.920; b.p. 170°-180°, and contains pinene, camphene, dipentene, phellandrene and

fenchene. Heavy oil of cumphor, having a sp.gr. of 0 960-0 970, and b p. 240°-300°, contains cadinene, safrol, eugenol, cincol, fenchone, terpincol, &c. Besides these compounds, both of the oils contain a large amount of camphor in solution Before refining crude camphor, the oil and water must be got rid of, as the oil gives the refined camphor a fatty touch, and the water renders it opaque. The elimination of water and oil is often effected by washing in a centrifugal machine with small quantities of water; the liquid diaming off contains both oil and water. The separation is also done by hydraulic pressure, or by crystallisation from boiling benzene or light Resublimation in Europe is done principally in glass retorts, yielding a round concave cake weighing 6-8 lbs, having an opening in the centre In America the camphor is sublimed from flat iron pans with iron covers, yielding plates of about 18 inches square and 1 inch thick, weighing from 8 to 10 lbs. The process consists in mixing the camphor with a little lime, charcoal or iron filings, according to the quality, and charging each pan with about 12 lbs of material, the subliming pans are heated slowly for 2 hours, and then fired rapidly up to 190° this temperature being kept for from 12 to 16 After cooling, the refined product is removed from the cover by slightly heating, and is then cut into the desired shape (Drobegg, J. Soc Chem Ind. 1907, 381)

Purified camphor comes on to the market in three forms, 'bells,' 'blocks' and 'flowers.' Block camphor consists of colourless translucent masses traversed by numerous cracks. It can be broken but is too tough to admit of being powdered by trituration. This may, however, be readily effected if a little alcohol be added, which escapes again during the operation. Camphor finds very extensive commercial application in the preparation of celluloid, in medicine where it is used externally as a mild rubefacient and to destroy parasites, internally, as a carminative and intestinal disinfectant, and to a minor extent in the manufacture of explosives Sp gr 0 986-0 996; mp. 178.75° corr. (Lenz. Arch. Pharm. 1911, 249, 289); b p 204° It sublimes to some extent at ordinary temperatures and condenses in hexagonal crystals (cf. Folger, Y.B. Pharm 1886, 232);  $[\alpha]_{\rm D}$  44·22° in 20 p.c. alcoholic solution (Beckmann, Annalen, 1889, 250, 353). Refraction coefficient (Kanonnikof, J. pr. Chem 1885, [2] 31, 348) Camphor in common with chloral hydrate and some other substances rotates in a peculiar manner on the surface of water (lowering of surface tension by solution), (Tomlinson, Phil Mag. 1873, [4] 46, 376, Pharm. J. 1874, [3] 4, 654, 672; Chem. News, 1885, 52, 50). It is very sparingly soluble in water (1:700), readily in 90 pc. alcohol (1·1), also in chloroform and ether. Mixed with chloral hydrate, it forms an oily liquid  $C_{10}H_{16}O$ ,  $CCl_{3}CHO,H_{2}O$ , which has been employed in CCI<sub>3</sub>CHO,H<sub>2</sub>O, which has been employed in medicine as an anodyne pigment (Brown, Pharm. J. 1873, [3] 4, 729; Saunders, *ibid.* 1876, [3] 7, 89; Cazeneuve and Imbert, Bull Soc. Chim 1880, [2] 34, 209; Zeidler, J. 1878, 645; Albright, Amer. J. Pharm. 1886, 282) Similar combinations take place between camphor and chloral alcoholate (Zeidler), phenol camphor and chloral alcoholate (Zeidler), phenol (Léger, Bull Soc Chim 1890, [3] 4, 725); menthol (Kyle, Amer J. Pharm. 1885, 429), and thymol (Symes, Pharm. J 1879, [3] 9, 598).

thymol (Symes, Pharm. J 1879, [3] 9, 598).

Camphor is a saturated hydroaromatic ketone, but does not combine with sodium bisulphite For identification purposes use is made of the oxime  $\rm C_{10}H_{16}NOH,~m.p.~118^\circ-119^\circ$  (Auwers, Ber. 1889, 22, 605), the semicarbazone (Auwers, Ber. 1889, 22, 605), the semicarouzone  $C_{10}H_{16}=N$  NHCONH<sub>2</sub>, m.p. 236°-238° (Themann, Ber 1895, 28, 2191), the oxymethylene compound  $C_{10}H_{14}O=CHOH$ , mp 80°-81°, and the benzylidene compound  $C_{10}H_{14}O=CHC_6H_5$ , mp 96° Of its substitution products, one, a-bromocamphor  $C_{10}H_{15}OBr$  has been employed in medicine as a sedative in nervous excitement, and is an article of trade It is best prepared by treating powdered camphor with biomine at 100°, pouring the liquefied product into water and recrystallising from spirit (Armstrong and Matthews, Chem News, 1878, 37, 4) It has a weak camphoraceous odour and taste, mp 76°; bp 274° It is insoluble in water, but soluble in the common organic solvents, in fixed oils, and in sulphuric acid By the action of reducing agents camphor is converted into a mixture of borneol and isoborneol, the former predominating (Beckmann, J pr Chem. 1897, [2] 55, 35, Haller, Compt rend 1887, 105, 227). On oxidation with nitric acid (Kachler, Annalen, 1878, 191, 143) or hot alkaline permanganate (Grossei, Ber. 1881, 14, 2507), camphor yields camphoric acid C<sub>8</sub>H<sub>14</sub>(COOH)<sub>2</sub>.

Synthetic Camphor, Inactive Camphor Camphor can be manufactured from turpentine oil, and has from time to time appeared on the market. It is termed 'synthetic' camphor to distinguish it from the so-called 'artificial camphor,'which consists of pinene hydrochloride, and is unsuitable for the chief purposes to which camphor is applied. Only when the prices of the natural product are abnormally high has synthetic camphor been able to compete with it During the war, in Germany, owing to the failure of supplies of d-camphor, the Schering interests and probably other firms (Houseman, Amer. J. Pharm. 1915, 87, 58) produced synthetic camphor in quantity, whilst the American Camphor Corporation contend that they can supply the requirements of the celluloid trade with synthetic camphor (Oil, Paint and Drug Rep, March 29, 1915) Natural camphor and synthetic camphor only differ in their action on polarised light, the latter being either mactive or showing only a slight activity; in their physiological action they are identical (Joachimoglu, Ber deut Pharm Ges 1916, 26, 427). By working at low temperaturer, however, Darmois (Compt rend 1910, 150, 925) has obtained synthetic camphor with the rotati n of natural camphor, French oil of turpentine yielding l-camphor, whilst turpentine from the Aleppo pine, *Pinus halepenis* (Mill) yielded d-camphor The various routes taken by patented processes for the manufacture of synthetic camphor are shown by Fig. 1.

The first commercial process consisted essentially in converting oil of turpentine into pmene hydrochloride, which compound by elimination of hydrogen chloride yields camphene. Camphene is then converted into isoborneol, which, like natural borneol, yields camphor on oxidation. For the preparation of pmene

hydrochloude, oil of turpentine, which contains 70-90 p.c. of punene, depending on its origin, is treated with dry hydrogen chloride, whereby a solid hydrochloride is obtained together with a




Fig 1.

liquid hydrochloride which is not suitable for the manufacture of camphene. This liquid hydrochloride contains a considerable proportion of the solid hydrochloride in solution, which may be recovered by oxidising the only product with dilute nitric acid, or by extraction with petroleum after treatment with sulphune acid at 40°, which polymerises the oil (Eng. Pat. 7319, 1908) The preparation of camphene by the removal of hydrogen chloride from pinene hydrochloride (Wallach, Annalen, 1887, 239, 6, Bruhl, Ber 1892, 25, 147; Reychler, Bull Soc Chim 1896, [3] 15, 371) was at first attended with difficulty. the yield was small and the product was liable During the last two to contain chloride decades the following methods have been patented for the production of camphene in farge quantities, many of which are claimed to give almost theoretical yields and a product free from chloride. Heating the hydrochloride with bases of the aliphatic series such as methylamine, or with piperazine or piperidine (US Pat 707270, 1902), or with tertiary bases of the pyridine and quinoline group (Eng. Pat. 5032, 1906); by heating with alcoholic, aqueous, or gaseous ammoma (U.S. Pat. 707271, 1902), or ammonia in presence of a phenol (D. R' P 264246, 1912), by heating with aqueous sodium hydroxide or ammonia in presence of a soap (US Pat 725890, 1903); by heating with lead acetate in glacial acetic acid solution (Fi Pat 349896, 1904); by heating with glacial phosphoric acid in glacial acetic acid solution (D R P 207888, 1908), by heating with a metallic salt of a higher fatty acid (Eng. Pat. 19960, 1906; 1948, 1907), or with a mixture of metallic salts of the fatty acids (Eng. Pat. 2488, 1913); by heating with a metal and an oxidising agent, such as zinc and barium peroxide in molecular proportions (Fr. Pat 361333, 1905); by heating with salts of phenols or naphthols either anhydrous or in presence of water (Eng Pat 16429, 1906; 21180, 1906; 5674, 1908) by herting with lime or other similar oxide (Fr Pat 375897, 1907; Eng Pat. 25453, 1907), in the presence of phenol at 180° (Eng. Pat. 22810, 1906), in the presence of polyvalent alcohols or carbohydrates (Eng. Pat 10784, 1907), by heating with alkali, and alkali or alkaline earth salt of a sulphonic acid of the benzene or naphthalene series (Eng. Pat. 10783, 1907); by heating with alkali or alkaline earth salts of organic sulphamides or with phenolsulphonic acids (D. R. P. 197346, 197805, 1907); by heating with alkali or alkaline earth hydroxides in presence of compounds which combine with water such as alkalı or alkalıne earth oxides (Fr Pat. 413282, 1909); by heating with zinc oxide in presence of alkalı nydroxide (Eng. Pat. 11029, 1911); by

heating with  $\alpha$  or  $\beta$  -naphthylamine (D. R. P. 206386, 1907), or with aniline (D. R. P. 205850, 1907); by heating with a meta- or pyro-borate, silicate, phosphate or arsenate in aqueous solution (D. R. P. 205295, 1908); by heating with phenol, cresol or naphthol and the carbonate of an alkalı or alkalıne earth metal (Eng Pat. 13902, 1909); by heating under reflux until the hydrochloric acid has gone (US Pat. 923967, 1909); by the Grignard reaction (Hesse, Ber 1906, 39, 1127) According to Eng. Pat. 10227, 1910, pinene vapour when passed over a catalyst at 200°–300° is converted into camphene

Beitram and Walbaum (J pr. Chem 1894, [2] 49, 1) made the discovery that camphene is converted into isoborneol by warming with a mixture of acetic acid and a little sulphuric acid, and decomposing the resulting isobornyl acetate by alcoholic potash. This method is employed on the manufacturing scale variant of the method is described in Eng. Pat 19961, 1906, where camphene is heated with oxalic acid in a suitable solvent with a condensing agent such as sulphune or phosphoric acids with formation of normal and acid esters of isoborneol. Camphene has also been directly converted into isoborneol by boiling in acctone with half its weight of 20 pc H<sub>2</sub>SO<sub>4</sub> for 10 hours, or in an autoclave at 100° for a shorter time (Fr. Pat The isoborneol purified by 385341, 1907) crystallisation from light petroleum or benzene is then oxidised to camphor. For this purpose various oxidising agents are employed, many of which have been patented Amongst these are . potassium permanganate in glacial acetic acid (Ber. 1900, 33, 3430); aqueous solution of permanganate employed with a benzene solution of isoborneol (F1. Pat 341513, 1904), potassium permanganate in alkaline solution (D. R. P. Anm. 38311, 1904), chlorine, either gaseous or in solution (Eng. Pat. 28035, 1904, U.S. Pat. 802792, 1905), hypochlorite, either alone (US Pat 864162, 1907), or in presence of copper and ferric chlorides (Eng. Pat. 20050, 1908); chromic acid or chromates (Eng. Pat. 26779, 1904, 14550, 1907); in acetone solution (Eng. Pat. 21946, 1907); or in presence of an indifferent solvent (Eng. Pat. 5513, 1908; 24779, 1912); nitric acid in which the presence of nitrous acid is essential (Eng. Pat. 9857, 1907); nitric acid in presence of vanadic acid and oxygen (Eng. Pat. 18047, 1908); nitric acid in presence of a halogen (U.S. Pat. 1313661, 1919), oxygen (U.S. Pat. 801485, 1905); oxygon, oxides of heavy metals or peroxides (Fr. Pat. 385352, 1907); basic oxidising substances, e.g. CuO, HgO (Eng Pat. 3750, 1908), metals or oxides which can liberate hydrogen from isoborneol (Eng Pat 26708, 1908); electrolytic methods (U.S. Pat. 875062, 1907; Fr. Pat. 387539, 1907); ozone (US Pat. 801483, 1905), nitrous anhydride (U.S. Pat. 802793, 1905); metallic copper at 300° in absence of air (Eng. Pat. 17573, 1906). According to US Pat. 790601, 1905, and Eng. Pat 21171, 1906, 180bornyl acetate and other isobornyl esters can be directly oxidised to camphor without previous saponification.

Camphene can also be directly oxidised to camphor (Armstrong and Tilden, Ber 1879, 12, 1756), and the following processes have been patented: Treatment with ozone at 80° (Eng Pat. 21294, 1890); treatment with hot air or

oxygen (Eng. Pat 3555, 1896), oxidation by potassium permanganate in acetone solution (addition to Fr Pat 349896, 1904); treatment of camphene at 180° with an oxidising agent, such as chromic acid, converts it into oxonide C<sub>10</sub>H<sub>16</sub>O<sub>2</sub>, which loses oxygen on treatment with water, and is converted into a lactone, camphenolide, which yields camphor on heating in presence of water (Fr. Pat 361333, 1905), oxidation with a per-salt of potassium and sulphuric acid (Fr. Pat 389092, 1908)

Another method of obtaining synthetic camphor depends on the formation of bornyl and isobornyl esters by the direct action of various acids on pinene or turpentine oil following processes have been patented: Treatment of anhydrous turpentine at 120°-130° with anhydrous oxalic acid (US Pat 698761, 1902, cf. Pond, J Soc Chem Ind 1907, 385), by heating pinene with anhydrous oxalic acid in presence of anhydrous chlorides, eg. AlCl<sub>8</sub> with the addition of a diluent (Eng. Pat. 14408, 1908); by heating turpentine with an alcohol and an acid anhydride, e g B<sub>2</sub>O<sub>3</sub> at 120°-140° (Eng Pat. 11248, 1908); by heating pinene with an aromatic polysubstituted acid (Eng Pat 18049, 1908); terpenes or oils containing terpenes heated at 160° with propionic or butyric acid (Fr Pat. 408106, 1909) Similar methods, based on the direct formation of bornyl or isobornyl esters from pinene hydrochloride, have been patented. by heating with an alkaline formate or oxalate in alcohol and water for 10 hours to 120° (Eng. Pat 5549, 1904); by heating with metallic formates and formic acid (Eng Pat 14379, 1907), solid pinene hydrochloride is heated with an alkaline earth oxide in presence of excess of a lower member of the fatty acids (Fr Pat. 393504, 1908); turpentine treated with dry hydrochlone acid gas is boiled with an alkali, alkaline earth or metallic butyrate or propionate in presence of excess of the corresponding fatty acid (Fr Pat 408065, 1909), by heating pinene hydrochloride in acetic acid solution with lead acetate (Fr Pat 349896 1904); by heating in acctic acid solution with an acetate and fused CaCl2 or KHSO4 (Eng. Pat. 1936, 1908); by heating with acetic acid in presence of zinc chloride (Eng. Pat. 8266, 1906), or some other zinc salt (Eng. Pat. 10798, 1900), and with addition of an alkali or alkaline earth salt of a formate, acetate or butyrate (Eng Pat. 18280, 1906); by heating with the acetate of an alkaline earth for 20 hours at 180° (Fr Pat 375590, 1907), by heating with a fatty acid in presence of metallic zinc (Eng. Pat. 16605, 1907), or by heating with organic acids in presence of zinc (Eng. Pat. 14752, 1908); by heating with fatty acids and zinc chloride in presence of cobaltous chloride or other suitable inorganic salt (Eng. Pat. 22129, 1907); solid pinene hydrochloride is shaken with a salt of acetic or stearic acid or phenol with an excess of either acetic, stearic acid or phenol in presence of a dehydrant (Fr Pat 382790, 1906)

The direct conversion of pinene hydro-chloride into camphor is claimed in Eng Pat 27317, 1913, by heating with nitric acid or its salts in presence of a catalyst, whilst two processes for the conversion of turpentine into camphor have been patented. They are: turpeutine is treated with a halogen at a low

temperature,  $-25^{\circ}$ , until the increase in weight is 25-50 p.c., and the product, still at a low temperature, is mixed with one-eighth to onefifth its weight of NaOH, when it is converted into camphor (Fr Pat. 392182, 1907); turpentine in a neutral solvent is treated with nascent oxygen (Fr Pat 406005, 1908)

(For analyses and points of difference between natural and commercial synthetic camphor, see Crane and Joyce (J. Soc. Chem Ind. 1907, 386), Lohmann (Ber. deut Pharm. Ges. 1909, 19, 222), and Lenz (Aich. Pharm. 1911, 249, 289).)

L $\alpha vo$ camphor; Matricaria camphor This camphor occurs naturally in the volatile oils of Feverfew, Matricana Parthenium (Linn) (Chautard, J. pr. Chem 1848, 45, 45; Compt rend 1853, 37, 166); Tansy, Tanacctum vulgure (Linn.) (Schimmels, Rep 1895, ii. 35), and Artemisia Nana (Pursh) (Whittelsey, Chem Zentr. 1909, 11 2160). It is also a constituent of Ngai camphor, the solid portion of the oil of Blumea balsamifera (D.C.) (Schimmels Rep. 1910, 1 149). It can be prepared by oxidation of camphene obtained from lavorotatory pinene (Riban, Bull, Soc Chim. 1900, 24, 19, Darmois, Compt. rend 1910, 150, 925), and by oxidation of l-borneol with nitric acid (Pope and Harvey, Chem Soc Trans 1901, 79, 76). It differs from common camphor only in its behaviour towards polarised light.

Inactive camphor is a constituent of the oil of Chrysanthemum sinense (var.) japonicum (Keimatsu, J. pharm Soc. Japan, 1909,

236, 1)

The following are the more important isomerides of camphor which occur naturally in volatile oils:  $Pulegone \ C_{10}H_{16}O$ , is only found in oils of the Labiate, chiefly in European Pennyroyal, Mentha pulegium (Linn) (Beckmann and Pleissner, Annalen, 1891, 262, 1); Fenchone in its dextro modification is a constituent of oil of Fennel, Foeniculum vulgare (G) (Wallach, Annalen, 1891, 263, 129), whilst the laro variety is found in oil of Thuja, Thuja occidentalis (Linn) (Wallach, Annalen, 1892, 272, 99), Thujone occurs in three forms, as d-a-thujone in Thuja oil (Paolini and Divizia, Atti. R. Accad dei Lincei, 1912, 21, [1] 570), as 1-a-thujone in the same oil (Wallach, Annalen, 1893, 272, 99); and as d-\$\textit{d-\textit{\textit{d-b-thu}}}\textit{one in oil of Tansy, Tanacetum vulgare} (Linn) (Semmler, Bei. 1892, 25, 3343), and oil of Wormwood Artemisia Absinthium (Junn) (ibid); Citral is a frequent constituent of volatile oils and a major ingredient of the oil from Backhousia citriodora (F v M) (Bertram, Schimmels, Rep. 1888, ii. 17), and of Lemon-gr ss oil, Cymbopogon flexuosus (Stapf.) (ibid.); Sabinol is the most important constituent of oil of Sav.n, Juniperus Sabina (Linn.) (Fromm, Ber. 1898, 31, 2025).

The more important homologues of common camphor belong to the sesquiterpene alcohols (see Terrenes) Owing to their capacity for crystallising from volatile oils some of them have long been known

Matico camphor C<sub>15</sub>H<sub>26</sub>O, was formerly a constituent of oil of matico, Piper angustifolium (Ruiz et Pavon), but appears to be absent from present day oils. M.p.  $94^{\circ}$ ,  $[a]_{D}$ -28  $7^{\circ}$  (in chloroform). Boiling with 25 p.c. sulphuric acid converts it into a blue sesquiterpene

C<sub>15</sub>H<sub>24</sub> (Kugler, Ber 1883, 16, 2841, Thoms, Arbeit Pharm. Inst. Berlin, 1905, 2, 125).

Cedar camphor, Cypress camphor, Cedrol  $C_{1b}H_{2d}O_{,}$  is a frequent constituent of the volatile oil of  $Juniperus\ virginiana\ (Linn\ ),\ m\ p\ 86°-87°, <math>[a]_{D}\ 9.5\ (in\ chloroform).$  It yields cedrene  $C_{1b}H_{2d}$ when treated with phosphoric anhydride (Walter, Annalen, 1841, 39, 247, cf also Semmler and Mayer, Ber. 1912, 45, 1384)

Cubebs camphor C<sub>15</sub>H<sub>26</sub>O, is a constituent of the volatile oil of Piper Cubeba (Linn), when it has been exposed to the air for some time M.p. 68°-70°, bp 148°, with loss of water, a change which very readily takes place under various conditions of treatment forming a sesquiterpene (Schmidt, Ber 1877, 10, 188. Schaer and Wyss, Arch Pharm 1875, 206, 316)

Ledum camphor C<sub>15</sub>H<sub>26</sub>O, obtained from the volatile oil of Marsh Tea, Ledum palustre (Linn ). Needles, m p  $104^{\circ}$ - $105^{\circ}$ , b p.  $282^{\circ}$ - $283^{\circ}$  The sesquiterpene, ledene C<sub>15</sub>H<sub>24</sub>, is readily produced by the action of dehydrating agents (Rizza, Chem. Zenti 1887, 1257; Hjelt, Ber 1895, 28, 3087; Lomidse, Chem. Zeit Repert, 1903, 27, 284)

Patchouli camphor, patchouli alcohol  $C_{15}H_{26}O,\ crystallises$  out from patchouli oil, the product of *Pogostemon Patchouli* (Pell), m p 56°, [a]<sub>p</sub>-977° (in chloroform) It readily loses the elements of water forming patchoulene  $C_{15}H_{24}$  (Montgolfier, Compt rend 1877, 84, 88, Wallach, Annalen, 1894, 279, 394, Schimmels, Rep 1905, 1 62)

Eudesmol C<sub>15</sub>H<sub>26</sub>O, is a constituent of the oils of various species of Eucalyptus, but especially of E macrorhyncha (F v M) (Baker and Smith, J. Linn Soc. N S W. 1899, 33, 86) M p 84°, [a], 31 3° (in chloroform) (Semmler and Tobias; Ber 1913, 46, 2026, Semmler and Risse, ibid 2303).

Thyme and allied camphors. Thymol  $C_{10}H_{14}O$ ,

Isopropyl m-cresol
Thymol is a constituent of the volatile oils of Ajowan, Carum cophicum (Benth) (Haines, Chem Soc Trans 1856, 8, 289; Stenhouse, Annalen, 1855, 93, 269, ibid 1856, 98, 309, Muller, Ber 1869, 2, 130), Thyme, Thymus vulgaris (Linn) (Doveri, Annalen, 1848, 64, 374; Lallemand, *ibid* 1857, 102, 119); Wild Thyme, T. Serpyllum (Linn) (Jahns, Ber 1882, 15, 819, Ocimum viride (Willd) (Goulding and Pelly, Proc Chem Soc 1908, 24, 63); Horsemint, Monarda punctata (Linn) (Arppe, Annalen, 1846, 58, 41), Satureia Thymbra (Linn) (Schrmels, Rep 1889, n 55), Origanum hirtum, (Link.) (Holmes and Umney, Perf. and Essent. Dir Rec., 1913, 4, 73) On the manufacturing scale it is prepared from Ajowan and Horsemint Oils (cf. Holmes, Perf and Essent, Oil Rec, 1916, 7, 311; also U S Dept. Agric. Bull. 372, 1916), by extraction with caustic soda solution, when the thymol forms a water soluble sodium salt, and is thus separated from the hydrocarbons alkaline solution is acidified with hydrochloric acid, which precipitates the crude camphor. may then be purified by crystallisation from glacial acetic acid or alcohol. Another plan is to submit the oil to fractional distillation and to freeze out the thymol from the heavier portions of the distillate. A possible source for syn-

thetic thymol is cymene (U.S. Pat. 1306512,

Thymol has the odour of thyme and a warm pungent taste; m.p. 50·5°-51 5°; b p. 232° at 752 mm. It is sparingly soluble in water (1:1200) and glycerine (1:1000), but readily in the usual organic solvents and alkaline solutions. It forms a liquid with menthol (Kyle, Amer J. Pharm. 1885, 429); camphor (Symes, Pharm. J. 1879, [3] 9, 598), berberne (Lloyd, New Remedies, 1881, 195), but not with chloral hydrate (Symes, cf. Mazzaia, Gazz chim ital. 1883, 13, 272) If a solution containing thymol be warmed with half its volume of glacial acetic acid, together with more than its volume of sulphuric acid, a deep red-violet colour is developed. This reaction is obtained even in very dilute solutions For the further identification of thymol use is made of the phenyl carbamate  $\rm C_{10}H_{13}OCONHPh$ , mp 107° (McKenzie and Semmler, Ber 1906, 39, 1158), and nitrosothymol C<sub>10</sub>II<sub>13</sub>O(NO), m p 160°-162° (Schuff, Ber 1875, 8, 1500) The methyl ether of thymol C10H13OMe occurs naturally in the essential oil of sea-samphire, Crithmum maritinum (Linn) (Delépine and Belsunce, Bull Soc Chim. 1918 [ıv ] 23, 24)

For estimation of thymol in essential oils, Kremers and Schremer (Chem Zentr. 1897, n 147, cf Messinger and Voitmann, Ber. 1890, 23, 2753) recommend the following process known weight of the sample is mixed with an equal volume of light petroleum in a stoppered burette, and agitated with 5 p c aqueous caustic soda till no further diminution of the oily layer takes place, and a drop of the oil dissolved in 20 drops of chloroform, on gently warming with a small piece of sodium hydroxide, does not turn red The decrease in volume is observed or the alkaline liquor can be titrated with rodine, but it appears the results obtained by titration are about 3 pc lower (For iodometric estimation, see also Messinger (J pr Chem 1900,

[2], 61, 247)

(For the estimation of the purity of thymol, see Seidell (Amer Chem J 1912, 47, 508), Ehon (J Amer Chem Soc 1917, 39, 1513), Redman, Weith and Brock (J Ind Eng Chem

1913, 5, 831))

Thymol is extensively used in medicine on account of its strong antiseptic properties. In its general behaviour it resembles carbolic acid It has also been used internally as a vermicide Of its derivatives a great number have been tried in medicine, but only three have been used to any extent: Thymol volude (Messinger and Vortmann, Ber 1890, 23, 2754; Carswell, Chem. News, 1893, 68, 181, DRP. 49739, 1889) obtained by condensation of two molecules and the introduction of two atoms of iodine is used as a substitute for iodoform; thymoform  $[C_{10}H_{13}O]_2CH_2$  (D R P. 99610, 1898) used as an antiseptic dusting powder; thymol carbonate (C<sub>10</sub>H<sub>13</sub>O)<sub>2</sub>CO (Richter, J. pr Chem 1883, [2] 27, 505) is used as an anthelminthic

Camphors isomeric or homologous with thyme camphor. Carvone, formerly carvol C<sub>10</sub>H<sub>14</sub>O, b.p 230°, is a liquid isomeride contained in the following volatile oils Caraway seed, Carum carvi (Linn.) (Wallach, Annalen, 1893, 277, 107), and Dill-fruit Anethum graveolens (Linn) (Gladstone, Chem. Soc. Trans. 1872, 25, 1), both

containing 50 to 60 p.c. d-carvone, whilst 1-carvone is contained in the volatile oils of Spearmint, Mentha viridis (Linn) (Gladstone, ibid. Fluckiger, Ber. 1876, 9, 473); Kuromoji, Lindera sericea (Bl.) (Kwasnik, Arch. Pharm 1892, 230, 265). 1-Carvone has been found in Ginger-grass oil, Cymbopogon Martini (Stapf) (var) Sofia (Walbaum and Huthig, Schimmels, Rep. 1905, i. 34). For the isolation of carvone, its power of combining with sulphuretted hydrogen to form a crystalline hydrosulphide (C<sub>10</sub>H<sub>14</sub>O)<sub>2</sub>H<sub>2</sub>S, mp. 210°-211°, is employed (Fluckiger, Ber, 1875, 9, 468, Wallach, Annalen, 1899, 305, 224), but a simpler process is to form the water soluble compound with neutral sodium bisulphite (cf Sadtler, J Soc Chem. Ind 1904. 23, 303) On reduction carvone yields dihydrocarveol C<sub>10</sub>H<sub>1</sub>,O, a constituent of Caraway Oil (Schimmels, Rep. 1905, i 50)

Carvacrol, cymophenol C<sub>10</sub>H<sub>14</sub>O, Isopropyl o-cresol is a thick oil, solidifying at 0°, b p 236°-237°, closely related to thymol It is formed by distilling carvone in presence of sulphune acid, phosphoric acid, or caustic potash, and occurs mainly in various thyme, origanum and monarda oils. It may also be obtained by chemical means from Spruce Turpentine (US) Pat. 1265800, 1918, cf also J Ind Eng Chem 1918, 10, 982) For its identification the phenylcarbamate C<sub>10</sub>H<sub>13</sub>OCO-NHPh, mp. 140° (Gildemeister, Arch. Pharm. 1895, 233, 188) and its nitroso-derivative C<sub>10</sub>H<sub>13</sub>O(NO), m.p. 153° (Klages, Ber. 1899, 32, 1518), are especially suitable

Irone  $C_{1,3}H_{20}O$  is a constituent of oil of orris, the product of distillation of Iris germanica florentina (Linn), I pallida (Lam), and I (Linn) (Tiemann and Kruger, Ber 1893, 26, In the diluted state it has an odour of Bp 111°-112° at 2 mm, 144° at 2675) violets 16 mm. It has been synthesized from acetone and A4-cyclocitral (Merling and Welde, Annalen, 1909, 366, 119), see Perfumes.

Santalol C15H24O is a liquid camphor obtained from the volatile oil of sandalwood, Santalum album (Linn) The commercial product known under various trade names is prepared by steam distillation of the finely divided sandalwood The bulk of the world's supply comes from Mysore, where it was being distilled locally (Perf and Essent Oil Rec 1917, 8, 359) The physiological action of the oil is attributed to its alcoholic constituent, santalol, which is present to the extent of about 90 pc., and has been separated into a-santalol, a thick colourless liquid with a faint odour of sandalwood, bp 301°-302°, [a]<sub>D</sub>+1° 13′, and  $\beta$ -santalol, b p. 158° at 5 mm.,  $[\alpha]_D$  -41°47′ (von Soden, Arch Pharm 1900, 238, 353, Schimmels, Rep 1910, n. 107).

The oxygen content of the foregoing camphors is one atom. The following contain two atoms of oxygen: Diosphenol, Buchu camphor C<sub>10</sub>H<sub>16</sub>O<sub>2</sub>, a crystalline separation from oil of Buchu leaves, *Barosma betulina* (Bartl), *B. crenulata* (Hook), B. serratifolia (Willd.) (Fluckiger, Pharm. J. 1880, [3] 11, 174, 219), is closely related to thymol and carvacrol, into which it passes quantitatively by the action of concentrated hydrochloric acid at 150°-180° M p. 83°-84°, b.p. 109°-110° at 10 mm. It is characterised by its reduction of ammoniacal silver solution and Fehling's solution (Semmler and McKenzie, Ber. 1906, 39, 1158; Konda-koff and Bachtscheef, J. pr. Chem. 1901, [2] 63, 61)

Inula or Alant camphor, alantolactone  $C_{15}H_{20}O_2$  is the chief constituent of oil of Dissert. Rostock, 1895). Mp. 76°, bp. 275°, with partial decomposition (Bredt. and Posth, Annalen, 1895, 285, 349). It dissolves in alkalis with formation of the corresponding hydroxy acid, which melts at 94°. The lactone has been used in medicine, under the name 'helenin,' as an antiseptic for internal use. (See also Sprinz, Arch. Pharm 1901, 239, 201)

H. K. CAMPHORAL. Trade name for the camphoric acid ester of santalol.

CAMPHOSAN. A solution of camphoric acid methyl ester in santalol.

CAMWOOD. Camwood or 'cambe wood,' stated to be derived from a variety of Baphia nutida (cf banwood), is very similar in general properties to the other 'insoluble red' dyewoods It is, however, more expensive, yields deeper shades on dyeing, and its colouring matter is said to be more soluble than that present in the other woods

It has been examined by O'Neill and Perkin (Chem Soc. Trans. 1918, 113, 126), who employed for this purpose similar methods to those found

serviceable with sanderswood (lc)

Iso-santalin, the main colouring matter, forms a chocolate-coloured powder, which on grinding becomes redder in appearance, and is readily soluble in boiling methylated spirit When heated, it shows no sign of melting, darkens at 280°, and is fully decomposed at 290°-300°, being then a carbonaceous powder. It possesses the formula  $C_{22}H_{16}O_6(OCH_3)_2$ , gives with alcoholic potassium acetate the salt C<sub>96</sub>H,<sub>3</sub>O<sub>32</sub>K or C<sub>72</sub>H<sub>65</sub>O<sub>24</sub>K, shows evidence of the formation of soluble oxonium salts, and in general properties closely resembles its isomer santalin. The colour reactions and also the dyeing properties of the two compounds indicate, however, that they are distinct substances .-

Santalın Iso-santalin Alcoholic hydrobromic acid Crimson Reddish-violet Dilute sodium Dull red. Dull violet. hydroxide. ferric Alcoholic chloride Violet Bluish-violet

For dyeing the colouring matter in alcoholic solution was added to the water in the dye-beth and the dyeings were carried out (a) employing wool alone, (b) employing wool alone and subsequently saddening with bichrome, (c) employing wool mordanted with bichrome and cream of tartar, and (d) employing wool mordanted with bichrome and sulphuric acid.

Acetyl-180-santalin  $\mathrm{C_{24}H_{18}O_8(C_2H_3O)_4}$ , consists of a deep salmon-coloured powder, and does not possess a definite melting-point, being gradually decomposed without fusion between 250°–280°. A molecular weight determination employing naphthalene gave the high figure 2344, a result very similar to that given in these circumstances by acetyl-santalin itself, and which may possibly be due to the production of a colloidal solution

Deoxy-iso-santalin C<sub>24</sub>H<sub>18</sub>O<sub>2</sub>(OCH<sub>3</sub>)<sub>2</sub>, corresponding to the deoxysantalin of sanderswood, is a scarlet amorphous powder which has not yet been obtained in a definitely crystalline condition. When heated, it did not show a distinct melting-point, but decomposed at 160°–165° with evolution of gas. A solution of this substance in absolute alcohol gives no immediate precipitate with potassium acetate, as happens in the case of 130-santalin, and only when excess of the reagent is employed is a gelatinous deposit formed.

A comparison of the colour reactions and dyeing properties of deoxy-iso-santalin (a) and deoxy-santalin (b) is given in the following table—

(a)**(b)** Alcoholic solution\* . Orange Orango-brown Alcoholic hydrobiomic acid . Bught crimson Scarlet. Dilute sodium hydroxide . Crimson-scarlet Scarlet Alcoholic ferric . Violet. chloride Maroon.

The dyeing experiments were carried out by the same methods as those described above.

Deoxy-iso-santalin	(a) Pale red- violet	$^{(b)}_{ m Red}$ puce.
Deoxy-180- santalın	(c) Dull red- violet	(d) Pale dull red-violet.
Deoxy-santalın	$\stackrel{(a)}{\mathrm{Red}}$	(b) Dull bluish- red
Deoxy-santalın	. Dull crimson	(d) Red.

 $Acetyl-deoxy\text{-}iso\text{-}santalin \quad C_{24}H_{20}O_7(C_2H_3O)_4, \\ \text{when heated, fused and decomposed at }170^\circ\text{-}175^\circ\text{-}15$  It consists of an almost colourless powder differing considerably in appearance from acetyl-iso-santalin. A molecular weight determination, employing naphthalene as solvent, gave the figure 1324, which is approximately half that found in the same circumstances for acetyl-iso-santalin.}

By exhaustion with alcohol the sample of camwood employed by these authors gave 16 p.c. Camwood does not appear to contain either ptero-carpin of homopteio-carpin. A. G. P

CANADA BALSAM v BALSAMS, and OLEO-RESINS.

CANADA PITCH or HEMLOCK SPRUCE RESIN v RESINS.

CANADINE. A member of the isoquinoline group of alkaloids, occurring together with hydrastine and berbeijne in *Hydrastis Canadensis Sec* under Hydrastine

CANADIUM. A term formerly applied to a supposed new element, now used to denote an alloy of 1 part palladium, 2 parts of platinum, and 6 parts of mckel, used as a substitute for platinum. It has similar resisting powers, and melts at about 1570° Its cost is about half that of platinum (Heath, Met. and Chem. Eng. 1917, 17, 666)

CANADOL. Petioleum ether or ligioin That portion of refined petroleum which boils at about 60° and has sp gr of 0 65 to 0 70, consists mainly of noimal hexane (v Paraffin, and Petroleum).

### CANANGA OIL v OILS, ESSENTIAL.

**CANARIN.** A yellow colouring matter obtained by the action of bromine or a mixture of potassium chlorate and hydrochloric acid or ammonium persulphate upon potassium or ammonium throcyanate, it probably consists of  $C_8H_6ON_8S_7$  (Goldberg, J pr Chem 1901, ii, 63, 465, 64, 439, cf H Schmid, Dingl poly J 251, 41) (For details of mode of preparation, v. ibid. 253, 130, also J. Soc. Chem. Ind. 3, 476). Cloth dyed with canaim resists light and soap. Neither concentrated acids not alkalis destroy it or dissolve it out of the fibre. Bleaching powder has no action upon it. Acts as a moddant for basic aniline dyes.

CANDELILLA WAX. Candelilla occurs as a deposit on the surface of the candelilla plant (probably Euphorbia antisyphilitica), which grows wild in Mexico and Texas ally the wax was separated by boiling the parts of the plant with water, but of recent years a method of extraction with chloroform has been introduced, some of the works being capable of dealing with 5 tons a day From 2 5 to 6 p c of crude wax is obtained, and is purified by straining and melting in water containing 0.5 to 1 p c of sulphuric acid. The crude grey product contains up to 65 pc of water, 3 to 15 pc of dirt, and 0 5 to 1 5 p c of mineral matter purer red or brown variety is probably obtained by the extraction method (Berg, Chem Zeit 1914, 38, 1162)

Several methods of extraction have been patented In Sharp's process (US Pat. 1018589, 1912) the wax plant is boiled with petroleum spirit in the presence of fuller's earth, the extract clarified, and the solvent evaporated In Watson's patent the plants are heated to 149°, and subjected to the action of a downward current of steam (US Pats 1042992, 1912, and 1058648, 1913). A special form of extractor devised by Thatcher (US Pat 1056378, 1913) comprises a digesting compartment, reservoir for the solvent, and vaporisers to evaporate the solvent

The crude greenish-grey wax melts at 67°-68°, and has sp gr. at 15°, 0 9825-0 9850; acid value 12.4-14.39; sapomification value,

46'1-64'9; nodine value, 16'6-36'8; unsapomfiable matter, 77-91'2 p.c., largely consisting of hydrocarbons. The constituents identified include hentriacontane, dotriocontane (m.p. 71°), myricyl alcohol, two cholesterols, an oxylactone (m.p. 88°), and resmous constituents giving the Liebermann-Storch reaction (Fraps and Rather, J Ind Eng. Chem. 1910, 2, 454, Hare and Bjerregaard, ibid. 1910, 2, 203, Sanders, Chem. Soc. Proc. 1911, 250; Meyer and Soyka, Monatsh. Chem. 1913, 34, 1159; Berg. Lc.)

A difficulty attending the use of the pure blown or red wax, is that it can only be bleached after the addition of a considerable quantity of paraffin wax, and is therefore only suitable for the manufacture of inferior kinds of varnish, for insulating telephone wires, and especially for boot polishes, for which its low m p is an advantage. For other purposes, such as candlemaking, it is inferior to carnauba wax (m p 83-91°) (\* WALES)

CANDLES. Candles represent some of the most ancient and most useful forms of illuminants. The excellence of a candle depends on the nature of the wick and of the combustible matter, and on the manner and extent in which these are apportioned. The prototype of the candle was undoubtedly the torch, which must be regarded as a huge wick with the minimum amount of combustible matter, this proportion being gradually altered until the ratio used at present, viz. about 1 part wick to 50 parts of combustible matter, is reached.

The following table shows the stages in the development of candles.

Torch Pine branches or slips (Lat. tæda), saturated, naturally or artificially, with resinous or fatty matter

Link Rope-strands steeped in iosin, tar, or pitch; in olden times, doubtless, in asphalt or bitumen.

Flambeau. A core of hemp, soaked in rosin and coated with crude beeswax. Later, the outer coat was made of bleached wax

Dips. Wicks of rush-pith (rush-lights), clude flax or cotton, and subsequently of twisted cotton yarn, coated with beeswax or tallow by repeated dippings.

Mould's (Introduced about the 15th century by the Sieur de Brez.) Hard tallow, spermaceti, stearine, or parafiin wax, cast round the wick in a mould.

In addition to these varieties may be mentioned rolled, poured, and drawn candles

Dips, the cheapest, as well as the oldest form of candle, were made until comparatively lately of tallow or household grease. Beeswax candles were usually poured, drawn, or rolled. The great advantage of the 'dip' lies in the ease with which it can be made. The thrifty house-keeper was wont to set aside the superfluous kitchen fat for melting day, when part went to make soap, whilst the harder portions were formed into candles. The fat was thrown into boiling water and boiled three or four times with a little salt to 'render' it. The dirt and impurities being removed by subsidence, the clean liquid fat was run through a strainer into the candle pan. Some skill and experience were required to regulate the temperature of the melted fat. If too hot, the liquid fat would run

off the wicks; if too cold, the material would congeal too quickly and adhere to the wicks irregularly and in lumps. In households these would be rushes, divested of their piel with the exception of a thin strip which was left to give strength to the pith. These were usually tied in bunches of four, so that one wick could be held between each pair of fingers, and thus be immersed in the liquid fat. A short time was allowed for each dipping to cool the last coat Four dippings mostly sufficed, when the finished candles were hung up in an airy loft to harden and whiten

In dips as manufactured on a large scale, the wicks are of twisted cotton, and are strung on rods, each rod holding eight. The workman, holding an end of the rod in either hand, first immerses the wicks in very hot tallow, in order that their fibres may be thoroughly saturated, and then, as each rod is ready, it is slid on to a cross frame to cool; the wicks are then redipped in tallow at a temperature little above its solidifying point, till the desired weight is obtained. As a rule four 'dispures' are required


obtained As a rule four 'dippings' are required.

Many devices exist for reducing labour and ensuring uniformity of result in manufacturing Of these contrivances, the 'Edinburgh wheel' is the best known It consists of a long pole, pivoted vertically, and having mortises cut about its centre through which pass long wooden bars, each pivoted at its centre on an These bars carry frames at their exiron pin tremities, each of which, in its turn, holds about eight wick-rods. The post revolves, and as each bar with its frame passes over the melting pan, it is pulled down, so as to immerse the wicks. The opposite frame restores equilibrium, and the rotating and dipping are con-tinued until the proper weight is obtained, which is signalled by some continuance on the post, itself the object of much inventive in-The most perfect apparatus of this kind was patented by Price's Candle Company and worked at the Inventions Exhibition in London in 1885.

Rolled, poured, and drawn candles. Rolled candles are now almost obsolete. Formerly, the larger sizes of church candles were made by 'rolling' a cake of warm wax, kneaded until it was plastic, round a wick, and imparting the necessary smoothness and uniformity by means of a rolling-pin. This process has now been entirely superseded by that of 'pouring.' 'Drawing' is resorted to only in the case of 'cables' of small sizes, and tapers.

In pouring, an apparatus similar to that used in dipping is employed, but instead of the wicks being dipped, the melted beeswax is poured over them while the frame is kept in rotation (Fig. 1). In the larger sizes, the wicks have to be reversed from time to time to ensure a uniform thickness. The workman ascertains the correct dimensions by measuring with a piece of tape. After every two or three pourings, according to the weather, the hoops are hung in a current of air to cool. When the candles are of the requisite thickness, they are severed from the frame and placed on a smooth marble slab The operator then rolls them to and fro under a board, on which he leans with his full weight. By this process, which requires much skill and practice, the irregularities are smoothed away,

and the candles, in the hands of a skilful workman, assume the evenness of outline of moulded candles, with a peculiar softness, indescribable, though unmistakable when once identified.



his 1

much skilled labour, must render beeswax candles very expensive. They are, indeed, the most costly form of illuminant. Their use is confined to churches and, to a small extent, to carriage lamps, in which formerly no other candle could be employed, by reason of the strong spring, although now several efficient substitutes exist and are in use.

To understand the favour which beeswax continued to enjoy long after the introduction of stearine and paraffin wax candles, which are not only much cheaper, but give a better light, we must take several circumstances into account The first and, doubtless, the most cogent fact lies in the sharp contrast which formerly obtained between beesway and tallow, a contrast which finds mention in many a tale of social life. even as far back as the times of Haroun Alras-The candle at one period was absolutely the only means of illumination available, and formed a very considerable item in the household expenses As only the rich could afford this luxury, the beeswax candle became intimately associated with the other signs of wealth, as introduction of spermaceti, in the middle of the 18th century, greatly affected the use of beeswax, and this was quickly followed by the adoption of gas in houses. Within the last decades, the electric light has replaced both beeswax and spermaceti candles in houses where for centuries no other light was known

Tc adapt the beeswax as it comes from the hive to the requirements of the chandler, but Lttie is needed beyond the ordinary cleansing; processes. The clean beeswax is run from the 'copper' over a rapidly rotating drum into cold water, by which means it becomes divided into fine shavings. These are exposed to the influences of sun and air for several weeks, with occasional re-meltings and pourings, after which the now cream-white wax is ready for use.

Drawn candles are made by drawing a considerable length of wick through a pan of beeswax (Fig. 2), maintained at a regulated temperature by (fire or) steam. The wick is wound from one drum on to another. As it leaves the pan, it traverses a plate perforated with holes in-

creasing from one sixteenth of an inch to half an inch in diameter, whereby the superfluous wax is stripped off, the coated wick as it emerges being of the diameter of the hole. When the wick is all wound off, the drums are reversed, the end of the coated cotton passed through the




Fig. 2.

hole next in size, and the operation continued till the requisite thickness is obtained. This is seldom over half an inch, as the risk of cracking increases considerably with the diameter of the cable. In fact, this process is resorted to mainly for manufacturing 'spills' or lighting-wicks and the little tapers used for decorating Christmastrees, but here also stearine in the former and paraffin wax in the latter have gradually displaced beeswax.

Mould Candles -The chief variety of candles now made are mould candles. In order to ensure the proper burning of a candle, great attention must be bestowed upon the preparation of the wick, which is to the candle what the burner is to the gas. Till about 1820 the only wicks used were made of twisted cotton yarn, still to be seen in tallow dips No means for consuming the wick being employed, snuffers had to be used to remove the charred and glowing end Several expedients were devised to bring the wick automatically into contact with the air, and thus cause it to be consumed as the candle burnt The best of these consisted in twisting the wick round a rod, and while in a state of tension, coating one side with rize or other stiffening. This imparted a tendency to turn outwards and curl over, which to a great extent obviated the use of snuffers. Palmer, in 1852, invented a simple means for achieving the same result more thoroughly by introducing a fine thread coated with metallic bismuth into the substance of the wick, which thread he called the 'doctor' When ignited, the easily fusible metal formed a globule on the end of the cotton, which by its weight bent the wick out of the flame into the air, when the oxygen could combine with the incandescent carbon, whilst the bismuth was volatilised. So successful was this device, that Palmer's 'metallic wick' candles achieved great popularity, especially for burning in candle lamps, for which large candles, some-times weighing two pounds, were employed. The same device was again patented in Germany, by Varenkamp, as late as 1899

But the introduction of 'braided' or 'planted' wicks by Cambacères (in 1825) rendered such precautions unnecessary, the flat form of the wick, as now made, imparting a sufficient tendency to curve and bend outside the flame, where the cotton can burn completely. Now all except tallow dip candles are provided with these cores.

Before the wick can be used by the candle maker, it has to be 'pickled,' i.e. soaked in a

solution of certain chemicals, which vary with cated apparatus, consists of a cluster of pewter the nature and purpose of the cotton. The moulds, introduced in 1724 by Freitag, held preparation of these solutions forms one of the chief items in the manufacture of a good candle, and to carry it out thoroughly demands considerable practical experience; for the wick itself varies much in its composition, especially as regards the percentage of mineral matter The plait, too, will vary in tightness, however carefully woven Then again, a paraffin candle requires a thin wick to check a too-rapid supply of the liquid fuel. A beeswax or sperm candle, on the other hand, the melted material of which is less mobile, or has less tendency to smoke, needs a thicker wick. The nature of the combustible, its melting-point, viscosity, and burning powers; the compounding of a suitable material by mixing candle materials from several sources; the selection of a wick that in number of threads, structure, and substance is adapted to the special material chosen; the subsequent treatment of this with such salts as will correct or enhance its tendencies, -these are some of the points which must be considered before proceeding with the more mechanical operations involved in candle making

The wicks, which arrive at the factory from the cotton spinner in hanks, are placed two or three days before being wanted in the appropriate pickling solution. The object of 'pickor 'smoking,' two great defects to which the candle is liable. The 'pickling' retards the too rapid combustion of the cotton, and vitrifies it slightly, so that when the wick protrudes from the flame any ash can drop off readily In order to ensure the ready lighting of a candle, it has been proposed (Haase, Fr Pat 342527) to impregnate the protruding end of the wick with a solution of celluloid in acetone nitre, sal ammoniac, potassium chloride or chlorate, and ammonium phosphate, are the principal salts employed for pickling; though almost every manufacturer has his own recipe. The proportion of the salt to water is usually about 2 oz to 1 quart The wicks are steeped in the solution for about 24 hours, after which they are placed on a perforated shelf, to allow the bulk of the liquid to drain off, and then transferred to a centrifugal machine rotating at a speed of nearly 1000 revolutions a minute, where the residual water is expelled without the slightest torsion or other disturbance in the structure of the threads, which would be almost inevitable in any other process. The hanks, still slightly damp, are placed in a cupboard heated by steam, and finally hung up in an airy room until required. The amount of salt remaining in the fibre must, of course, be very small, but it is quite sufficient to influence materially the burning of the candle The wick is next transferred to the spools, from which it passes to the moulding machine. This task is performed by boys, who wind the wick off the This task is hank previously stretched on a circular rotating frame, on to 'spools' or bobbins on spindles (Fig 3) As they wind, they let the wick run through their fingers, to detect any knots or other irregularities, which might interfere with the burning of the candle. The spools are now ready for transport to the machine, invented by Cahouet in 1850. The old 'hand-frame,' the forerunner and prototype of the present complimoulds, introduced in 1724 by Freitag, held




Fig. 3


little loop of cotton, through which a wire is passed

The butt ends of the moulds open into a trough about 1 inch deep. Into this the hot material is poured until it is quite filled, and the whole frame is then placed in a tank of water, the temperature of which varies with the quality of candle, as will be explained presently When quite cold, the candles are removed from the pipes by simply inverting the frame, the cake of superfluous candle material having been previously scraped off, and the wires and pegs withdrawn As the pipes are made slightly conical and as the substance shrinks in cooling, there is seldom any difficulty in emptying the frame; if one or two of the candles should refuse to leave the moulds, a slight tap with a mallet speedily loosens them This method is obviously slow and cumbrous, and were it not that some buyers will have hand-made goods, also that a few odd sizes for which it would not be worth while to keep special machines can only be made in hand-frames, these frames would have been abandoned long since hand-frame can be filled and emptied once an hour, whilst a modern machine will do thice the amount of work in the same time

Fig 4 represents a candle machine of modern manufacture, for making candles with oidinary, i.e. not self-fitting, ends. It embodies the conbined inventions and improvements made successively by Sampson (Eng. Pat 1796); Binns (Eng. Pat. 2488, 1801); Morgan (Eng. Pat. 6610, 1834); Tuck (Eng. Pat. 7409, 1837); Palmer (Eng. Pat. 12077, 1848); Cowper (Eng. Pat. 1988, 1856); Humiston (Eng. Pat., in the name of Newton, 2641, 1857); Stainthorp (Eng. Pat, in the name of Pitman, 2556, 1857); Stainthorp (Eng. Pat. 740, 1860) The machine consists mainly of a metal tank, G, in which the pipes are fixed with their butts opening in a trough and their tips emerging below. The tips are not of one piece with the moulds, but can slide up and down, fitting water-tight when drawn home. The motion is communicated by pistons, D, to which the tips are soldered, and which are perforated longitudinally.

The wicks wound on the above-mentioned spools, which rotate on a series of spindles at

the base of the machine, pass up these perforations D. The tank G, which surrounds the



pipes, is preferably connected with hot and cold water cisterns, or with cold water and steam. The moulds are made of pewter (an alloy of tin

and lead), other materials, such as tin, brass, glass, &c, not having been found suitable in practice Porcelain moulds have been patented for paraffin candles (Ger Pat 195702), but it is very unlikely that they will be adopted in candle works, as they are too costly; moreover properly cleaned pewter moulds impart the same lustre to a candle that the porcelain mould gives. To begin the operation of casting, the wicks are drawn through the pistons, tips, and pipes, and temporarily secured in the troughs. These, and the moulds opening into them, are now filled roughly, merely to afford a startingpoint for the wick, and emptied as soon as the material is cold. To effect this, the trough is scraped clear with a 'spud' (a blunt kinfe of the width of the trough), after which the handle E is slowly turned. This imparts, by means of rack-and-pinion gear, a vertical motion to the pistons, which mount up the pipes, carrying the dummy candles with them, and unwinding the wick from the spools As the candles emerge from the pipes, they are received in a clamp frame c, which is held open by a spring handle When the tips are clear of the trough, the clamp is closed, gripping the candles firmly handle is now turned the leverse way, causing the pistons to descend in the moulds till the tips close the ends securely. Thus the wicks




Fig. 5.-Multiple moulding machine.

are left in the centre of the pipes, held by the most recent type of this machine is shown in dummies above and the spools below. The Fig. 5. The next step depends on the nature of

the material used. If stearine candles are to be made, the tanks are charged with tepid water, the temperature of which the operator determines by his hand in preference to other methods. The reason for this precaution depends upon the highly crystalline structure of 'stearic' acid ('stearme'), which renders it hable to crack if chilled too suddenly, or to crystallise if cooled too slowly, whereby the appearance if not the structure of the candle would be impaired Therefore the melted stearine is stirred, with the aid of simple machinery, while in the melting pans, till it assumes the consistency and appearance of gruel, in which state it is transferred to the 'jacks,' cans having a flat, wide spout. and thence to the moulds. When these are full, they are allowed to rest about 25 minutes until they are sufficiently hard to permit of their being withdrawn without fracture. The troughs are not scraped in this case, as the tops are hard enough if the candles are raised 2 inches. As soon as the pipes are filled again and the material set sufficiently to hold the wick without extraneous acid, the upper row of candles, often while still warm, is removed into trays where they cool and bleach. This tendency of stearine to crack was a great obstacle to the success of these candles when they were first introduced. Among the various devices employed to overcome the defect was that of stirring a small proportion of arsenic into the melted steamne This certainly broke the grain, but it was of course diffused into the air on burning the candles When this generally known, the prejudice against stearine candles was such as to affect injuriously the success of the new substitute for tallow, and to this day stearine candles enjoy but a comparatively limited popularity in this country

In a patent process for preparing the fatty acids for stearine candles (Reuter Process Co, Eng Pats 9393 and 9394, 1915), the glycerides are fractionally saponified by boiling with sulphonic acids, such as Twitchell's reagent (Eng Pat. 4741, 1908). It is claimed that candles made from the resulting fatty acids are not decolourised by an exposure of several

days to the sun and air.

Paraffin Candles - At present, stearine candles are replaced in this country by candles made of a mixture of stearine and paraffin wax, except in the case of hollow candles (see below) In France, however, the stearine candle, being protected by customs duties on paraffin wax, holds its own against paraffin candles; and in that country, as also in Italy, stearine candles are manufactured in considerable quantities In Germany and Austro-Hungary, the stearine candle is gradually being ousted by the paraffin candle When, in 1854, James Young obtained from shale oil a pure white paraffin, which was subsequently made into candles under a patent of J K. Field, the immediate popularity of the novel illuminant left little doubt as to what would be the material of the candle of the future. Even the most enthusiastic optimist of that day would have hesitated before accepting the prophecy that paraffin candles would be sold at half the price of tallow dips, as is the case to-day.

The main difficulty experienced in the early stages of the manufacture of paraffin candles arose from want of precaution in regulating the

temperature while casting. The same machine can be employed for either paraffin or stearing candles, but when paraffin wax is to be used. the water in the tanks is raised to a temperature of nearly 200° F., by passing steam through the pipe. Also the paraffin wax itself is heated to about 170°-180° F., and poured in a highly mobile state. When the pipes are filled, the hot water is discharged from the tanks and cold water immediately introduced, thus suddenly chilling the fluid material. If the exact point be missed, failure will result. Thus, if the pipes are colder than the inflowing paraffin wax, it will congeal as it touches the metal, assuming a dull. striated aspect. On the other hand, if the moulds are allowed to cool slowly, the candles will shrink on to, instead of from, the walls, and will be injured in removal from the moulds, if indeed they can be withdrawn at all in the solid When the operation is properly carried out, and the pipes are not worn by usage, and further, if the paraffin is highly refined and of a high melting point, the candles leave the moulds with a lustre which hardly suffers by ordinary handling. This lustre, combined with the translucency, snowy whiteness, and ready adaptability to ornament in shape and colour, gives paraffin wax the pre-eminence over all other materials for candle making It has, however, its disadvantages. The readiness with which it liquefies by heat, gives to even the best paraffin candles a tendency to 'gutter' Moreover, the paraffin wax has the peculiar property, which is independent of the melting-point, of becoming plastic after exposure to a moderate degree of heat. The time required to develop this tendency varies, it is true, with the hardness or fusibility of the paraffin wax, which ranges from 105° F to 140° F; but even ceresin, of a melting-point of 142°-143° F, is affected by pro-The inferior varieties of paraffin tracted heat candles, from the ease with which their substance melts, are very prone to smoke, the wick is apt to be too abundantly supplied with combustible, and is consequently chilled below the point at which complete combustion can take place

By judiciously mixing stearin and paraffin wax, the candle maker is able to obviate many of the defects of the substances, whilst retaining their advantages Such candles as 'petrostearine,' 'palmitine,' and other popular varieties and especially the paraffin candles of the Saxo-Thuringian paraffin-wax industry, are made on

this principle.

In the United States the method of preparing such composite candles is to melt the paraffin wax by means of steam, to add a small proportion of stearine, and to clarify the naxture by the addition of oxalic acid, which precipitates calcium salts. The clear upper layer of wax is then drawn off into a steam-jacketed pan, where it is heated, and impurities are detected and separated, after which it is transferred by means of a 'swimmer' to a 'jack' or bucket, which delivers it in a regulated stream to the moulding machine (Campbell, J. Inst. Petrol. Techn. 1916, 2, 274)

In Laursen-Schjodt's process (Fr. Pat. 466027, 1913), oleine or stearme is dissolved in turpentine or other solvent, the solution treated with the calculated amount of alkali for sapomification, and the resulting mass incorporated with paraffin wax.

The use of tilmethylamine in the manufacture of composite candles has been patented by Herzmann (D. .R P 275344, 1911), the trimethylamine being treated first at the ordinary temperature, and then under pressure at 150° with chlorohydroxy fatty acids or sulphuric esters of hydroxy fatty acids such as Turkey-red oil. The resulting product can be saponified

In Burke's patent (Eng. Pat. 14975, 1914) about 0.25 p.c. of a heavy metal salt of an organic acid (e.g. lead oleate or palmitate) is added to

the parafhn wax

FIG. 6

The transparency of stearine and paraffin candles may be reduced by adding from 1 to 6 p.c of 'denatured' ethyl or methyl alcohol,

the amount depending upon the degree of opacity desired (Moretti, Fr. Pat

431490, 1911)

One of the greatest improvements in candle making is the self-fitting end, by which simple device a candle will fit any normally constitueted sconce without being sciaped or using paper or any other expedient to enlarge or diminish the butt. The self-fitting end patented by J. L. Field (Eng. Pat 3032, 1861), is now in universal use

Fig. 6 gives an example of the conical butt, whilst Fig. 7 represents the machine patented by E. Cowles for its manufacture. The difficulty to be overcome lies in the butt projecting beyond the stem of the candle, which renders the ordinary method of withdrawal impracticable. To meet this, the butts are cast




Fig. 7.

in a separate frame, which can be fitted to or removed from the pipes at will. When the machine is to be filled, the butt frame is lowered by the chain till the butts rest upon Vol. II.—T.

and fit tightly into their respective pipes. The wicks are then drawn through the butt moulds with a wire and clamped in a centric position. after which the filling is proceeded with in the usual manner. When cold the wicks are east loose, the tops scraped off, and the upper frame hoisted off the butts, as shown in the illustration, the operation being concluded as before. It will be noticed that this apparatus precludes running a continuous wick, as is done in the making of candles having plain ends; and in a large factory, a serious loss is incurred in the course of the year by the meh or two of wick snipped off at each operation. A very ingenious machine for saving this waste is due to E Cowles, who invented a split bed-plate holding the butts When ready for withdrawal, the butt




Fig. 8


moulds are raised and opened out by a simple lever motion, and the candles can thus be passed through without further trouble (Figs. 8 and 9)

Numerous improvements have been made in the construction of candle machines by the makers in this country, in France, the United States, and Germany, but all these improvements affecting the principle All these machines work discontinuously. A continuous candle machine, for a description of which the reader must be referred to Engineering, 1901, Feb. 1st, has been constructed by Fournier et Cie, Marseille. The machine is very complicated, and would seem only to commend itself to works where a very large output can be relied upon, and where, at the same time, the works suffer from a shortage of cooling water.

An improvement, which has found more favour on the Continent than in this country, consists in perforating the stem of the candle

0

longitudinally so as to afford an internal escape for any combustible which would otherwise have




guttered down the outside The up-draught of air which takes place in the tubes has a tendency to ensure more perfect combustion. By reason of the hardness required in their manufacture, these candles must be made with stearine.


A more recent process to prevent guttering is that devised by Powell and Bush (Eng. Pat 110579, 1916), according to which the candles are varnished with a colour varnish or with paraffin wax stiffened with a combustible powder such as cellulose

('andles made for ordinary purposes are ready for packing immediately after leaving the Better-class candles are subcandle machine jected to polishing and finishing processes in special machines, which are much more used on the Continent than in this country

Much art has been expended on the ornamentation of candles Before the introduction of aniline dyes, candles were coloured with such pigments as vermilion, ultramarine, and verdigus, these, besides causing a dull, heavy appearance, rumed the wick by choking it with meom-bustible matter. In beeswax candles, this evil



by Field, was made by turning the plain candle in a lathe of complicated structure, which could fashion over thirty different sizes and patterns of candles, e.g. cabled, spiral,



The demand for cabled candles is now so large that they are made in moulds, the pistons while ascending being caused to rotate, so as to screw the candles out of their spiral moulds

Candles burning with coloured flames have been patented by Scheurer (D R P 216338, 1907), the candle material being impregnated with nitrates or other salts readily decomposing when heated, and metallic salts or wires capable of imparting a colour to the flame

Night lights are short thick cylinders of a fat or wax, furnished with small wicks, their object being to give a constant and lasting rather than a bright light They are usually cast direct in paper cases, and are recommended to be placed on water Price's Candle Co have adopted the plan of casting the cylinders in a machine similar to a candle frame, and employing a material so hard (pressed coco-nut stearine) as to retain its torm in hot weather without external support These lights are burnt in glass cups, thus shedding their full light and rendering the precaution of water unnecessary Another very popular form of this illuminant is Clark's 'pyramid' night light, which, as its name indicates, is coniform. In this, the material employed is pressed tallow of great purity, and the wick is a rush-pith with two lateral strips of the peel left adhering These turn outwards in burning, giving a neat and shapely flame. The base of the cone is a disc of plaster of Pans, which holds the wick upright till the whole of the fat is consumed; a result achieved less surely in the other kinds, where the wick is of 'inkle' (flax), held by a little strip of tin, into which the base of the fibre is fixed

Candles intended for heating purposes, and so burning with a hot smokeless flame, and giving only a feeble light, are prepared from esters of amino or imino acids melting between 50° and 130°, and decomposing between 200° and 300° Ammonium nitrate or similar substance is also added (Scheuble, D. R. P. 234340, 1909)

The use of viscose filaments as a material for candle wicks has been patented by Wilson (Eng Pat 5170, 1913).

The strides made in gas and electric illumination have not materially affected the prosperity of candle manufacture. On the contrary, every year sees a considerable increase in the number of candles made This is especially due to the increasing consumption of candles in mines (Transvaal, India, Australia, &c). An inquiry made by the writer, in 1903, as to the consumption of candle material in this country, led to an estimated quantity of 45,600 tons, which may be distributed over the different materials mentioned above, as follows tallow (for dips), 1000 tons; stearine, 2190 tons, paraffin wax, 42,200 tons, ceresin, 100 tons. France produces about 30,000 tons of candles, chiefly stearine candles; Germany, about 11,000 tons, chiefly paraffin candles; Holland, about 6000 tons, chiefly stearine candles; Belgium, about 4000 tons, chiefly stearine candles. Austro-Hungary,



Italy, and Russia produce much smaller quantities than the last-named countries.

Candles were exported from Germany to the value of £56,000 in 1912, principally to Russia, Mexico, Biazil and the Mediterranean countries. In 1912 the exports from Austria-Hungary, chiefly to the Balkan States, Russia, Turkey, and Egypt, were valued at £14,000; whilst the total exports from the United Kingdom (1913) were of the value of £435,000 (Board of Trade, Bull. No 89)

CANDLE-NUT OIL is obtained from the fruits of the candleberry tree, Aleurites moluccana (Willd), belonging to the family of spurgeworts (Euphorbiacca) The tree attains the height of 30 or 40 feet, is a native of the Moluceas and South Pacific Islands, and is widely distributed over the tropics of the eastern hemisphere, it also occurs in the Cameroons In India it is known as 'Indian akhrout.' The fruits resemble walnuts in their appearance, but the shell of the candle-nuts is far thicker. The name 'candle-nut' is derived from the practice of the South Sea Islanders to dry the kernels, thread them on reeds, forming a candle, several of these candles being wrapped in a leaf of pandanus (screw-pine) so as to make a torch

The average composition of the kernels is as follows

JW5					Per cent.
Oıl					. 610
Water					. 50
Proteins					. 230
Ash					. 34
Carbohyo	irates	and	crude	fibre	76
_					
					100 0

The cold-drawn oil is limpid, colourless, or yellowish, has a pleasant odour and bland taste, but cannot be used for edible purposes on account of its purging properties

The oil extracted from a typical sample of the nuts had the following characters sp gr at 155°, 0 9276; refractive index (25°), 1 4970, acid value, 08, saponification value, 1882, and rodine value, 1516 (Tests for Candle-nut American Soc for Testing Materials, 1914)

The oil is obtainable in enormous quantities, but hitherto it has not been used to such an extent as its drying properties would warrant

The fruit of A titloba (Forst) is two-celled, fleshy, ohve-coloured, and about two and a half inches in diameter. Each cell contains one nut, resembling in appearance and taste a walnut, the shell of which is very hard The seeds yield about 50 pc of an oil known as 'kekuna oil' Seeds from Hong Kong, where the tree is cultivated as one of the best shade trees, were found to contain 60 p c. of an edible oil very similar to that obtained from A. moluccana; whilst Mauritius seeds yielded 68 l p c. of oil with the following characters. sp gr. at 15°, 0 927, saponification value, 1937; and iodine value, 151. The nuts are worth about £28 per ton in Europe (Bull Imp. Inst 1912, 10, 44). CANELLA BARK. The dued

The dued bark of Canella alba (Murr), obtained from Florida and the West Indies in large buff-coloured quills or pieces of about an inch broad. Has a bitter pungent acrid taste and odour of a mixture of cloves and cinnamon, due to a volatile oil con-

The bark contains a bitter principle, cloves canellin, together with resin, mannite, and starch. Used in medicine as an aromatic bitter and tonic, and in rheumatism and gout

CANELLE or BISMARCK BROWN v. Azo-COLOURING MATTERS.

CANFIELDITE v. ARGYRODITE

CANNABIN or INDIAN HEMP RESIN v.

CANNABINENE v. RESINS.

CANNABINOL. The active constituent of Indian hemp resin or 'hashish' (v Resins).

CANNABIS INDICA v. RESINS. CANNONITE v EXPLOSIVES

CANTHARIDES. Spanish flies (Lytta vesicatoria) Dried coleopterous insects used in medieme, in the form of plasters, tineture, limments, &c, on account of their vesicating properties, due to a peculiar acrid principle, cantharidin.

Employed also as a stimulant diuretic, as an aphrodisiac, and as an emmenagogue Obtained principally from Hungary, Russia, and the South of France

(Gadamer, Arch Pharm 1914, 252, 609, 1917, 255, 277, 1917, 255, 315) is obtained by treating the powdered insects with chloroform, ether, or ethyl acetate, acidined with sulphure acid, evaporating the extract and treating the residue with carbon disulphide or light petroleum, to remove fat, or by mixing the canthandes with water and magnesia, drying, treating with dilute sulphuric acid, and extracting with ether. The yield varies from 03 to 103 pc of the insects taken Crystallises in trimetric plates, melts at 218°, but begins to sublime at 85° It is tasteless and modorous, blisters the skin strongly, especially when in solution or mixed with fat Yields the alkaline solution or mixed with fat salts of cantharie acid C10H11O5, by heating with aqueous solution of potash or soda For other derivatives of canthandin, see Rudolph, Arch. Pharm 1916, 254, 423. Gadamer has shown that the isocantharidin and isocantharidic acid of Andeilini and Ghira are in reality acetyl hydrato cantharic anhydride and acetyl hydrato

canthanc acid respectively
CANTON'S PHOSPHORUS v. CALCIUM CAOUTCHOUC v RUBBER; also TERPENIS CAOUTCHOUC OIL. An oil which has proved efficient in preventing rust - It is spread with a piece of flannel in a very thin layer over the metallic surface which is to be preserved, and allowed to dry. To remove it, the article taining eugenic acid which is also found in oil of has simply to be treated with caoutchouc oil again, and washed after 12 to 24 hours (J. Soc. Chem Ind 1, 315)

CAP COMPOSITIONS v EXPLOSIVES. CAPE BERRY WAX & WAXES CAPRIC ACID or DECOIC ACID v DECATOIC ACLD

CAPROIC OF HEXOIC ACIDS C'5H11 COOH

Seven isomerides are known

1 n-Caproic acid CH3(CH2)4COOH occurs in the fats, in cheese, and, possibly, mixed with certain of its isomerides, in cocoa-nut oil, in St John's bread, in the fruit of Heracleum sphondylium and in the flowers of Satyrium hircinum, and in the fusel-oil from the fermented molasses of mangel wurzel best prepared by fractionating crude fermentation butyric acid (Lieben, Rossi, Annakn, 159,
75, 165, 118), b p 204 5°. sp gi 0 9446 at 0°
2 Isobatyl acetic acid, or γ-methylicaleric acid

### (CH,),CHCH,CH,COOH

is present in butter, may be obtained by the action of hydroide acid upon hydroxyisocaproic acid (Mick & Annalen, 180, 57). Fittig Ruhlmann, ibid 226, 347), or by the distillation of isobutylmalome acid (Konig, Monatsh 15, 20 , Bentley and Perkin, Chem. Soc. Trans  $1898,\,487)$  , b p  $200^\circ\text{--}201^\circ$  , sp gi. 0.925 at 0°

3 Diethylaectic acid, pseudocaproic acid, or a-ethylbutynic acid (C<sub>2</sub>H<sub>3</sub>)<sub>2</sub>CH COOH, may be obtained by the action of sodium upon acetic ester, and treatment of the product by othyl iodide (Frankland and Duppa, Annalen, 138, 221), by boiling diethylcarbinol eyanide with potassium cyanide (Saytzeff, ibid 193, 349), by reduction of dichlorethyl acetic ester by sodium amalgam (Markownikow, Ber 1873, 1175), by heating sodium ethylate and sodium acetate in a stream of carbon monoxide (Geuther and Frohlich, Annalen, 202, 308), by heating diethylmalonic acid at 170°-180° (Conrad, ibid 204, 141), by heating α-diethyl-β-hydroxy-butyric acid (Schnapp, ibid 201, 70), by heating diethyleyanacctic acid with concentrated hydrochloric acid at 160° (Hesse, ibid 18, 749), bp  $190^{\circ}$  (756.5 mm), sp gr 0.9196 at  $20^{\circ}/0^{\circ}$ 

4 Methylpropylactic acid or a-methylcalcic acid CH<sub>3</sub> CH<sub>2</sub> CH<sub>2</sub> CH<sub>2</sub> CH<sub>3</sub> CO<sub>2</sub>H is obtained by the oxidation of its aldehyde, and by the reduction of methylacitylic acid (Licben and Zeisel, Monatsh 4, 37, 63), by heating a-methyl valerolactone or isosacchain with hydrodic acid and red phosphorus (Liebermann and Scheibler, Ber 1883, 1823, Kiliani, *ibid* 185, 632), by heating  $\alpha$ -methylpropyl- $\beta$ -hydroxybutyric acid (Jones, Annalen, 226, 292), and by the distillation of turpentine, bp 193°-194°, sp gi 0 9279 at 18°/0°, 1 part water dissolves 0 5693 part at 17°

5 Methylisopropyl acetic acid, isocaprone

acid, or ab-dimethylbutyric acid

### $(CH_3)_2CH$ $CH(CH_3)CO_2H$

may be formed from methylasopropylearbinol cyanide (Markownikow, Zeitsch Chem 1860, 1 205), by the oxidation of the corresponding hexylalcohol from Roman camomile oil (Kobig, Annalen, 195, 102), or by melting camphoric and with caustic alkali (Crossley and Perkin,

Chem. Soc Trans 1898, 16), bp 189°-191°.
6 Dimethylethylacetic acid or aa-dimethylpromonic acid (CH<sub>3</sub>)<sub>2</sub>C(C<sub>2</sub>H<sub>5</sub>)CO<sub>2</sub>H is formed by

the action of hydrochloric acid upon dimethylethylearbinol cyanide (Wischnegradsky, Annalen, 174, 56), and by the oxidation of the methylethyl ketone prepared from pinacolm (Lawimowitsch, *ibid* 185, 126) A colour-less liquid, bp 187°, mp — 14°

7 (a) B-thylbutyne acid of active caproic acid CH<sub>3</sub> CH(C<sub>2</sub>H<sub>3</sub>)CH<sub>2</sub>COOH is obtained by the oxidation of the active hexylalcohol obtained from Roman camomile oil (Romburgh, Rec trav chim 5, 221, Kobig, Annalen, 195, 102), b p 196°-198° (770 mm); sp gr 0 930 at 15°

(b) Inactive \(\beta\)-ethylbutyric acid may formed by heating secondary butylmalonic acid (Romburgh, Rec trav chim 6, 153) or from secondary butyliodide, diethylmalonic ester and sodium ethoxide (Kuhlisch, Monatsh. 14, 561, Bentley, Chem Soc Trans 1895, 267).

CAPRYL ALCOHOL V OCTYL ALCOHOL

CAPRYLIC ACID V OCTOIC ACIDS

CAPSAICIN V RESINS

CAPSICUM (Bird pepper, Cayenne pepper) The dried tipe fruit of Capsicum minimum (Roxb), which was probably a native of tropical America, but is now known only in cultivation, the chief supply coming from Africa and India

The fruit is 1-2 cm long and about onethird as broad, somewhat flattened oblongovoid in shape, and is covered with a crimson or scarlet, semi-transparent, wrinkled corraccous epicarp It is two-celled and contains about 20 yellow, flat, reinform seeds, 3.5 mm long and 2.5 mm broad. It is prepared for use by gunding the whole fruit

Cayenne pepper is sometimes adulterated with flour, corn-meal, or even sawdust Entire capsicum is also frequently substituted by larger Adulteration of the powder may be detected by the estimation of oleo-resin, which should not be less than 5 pc , and of the ash, which should not exceed 7 pc , and by microscopical examination (v Bitto, Chem Soc Absti 1893, D, n 546)

The active principle of capsicum is capsaicin (capsiein, capsicol)  $C_1, H_2, O_3N = (C_1, H_2, O_3N,$ Lapworth and Royle), which is present to the extent of 0.05-0.07 p  $\epsilon$  (Morbitz, Chem. Zentr 1897, n 593) It is obtained from the ethercal extract of capsicum, and crystallises from light petroleum in colourless plates, m p 63 5°, easily soluble in ether, alcohol, chloroform, &c., but almost insoluble in hot water and insoluble in cold water. It is not volatile in steam, and has the properties of a weak, phenol-like acid The benzoyl derivative forms silky needles, m p 74° (Micko, Chem Zenti 1899, i 293)

The other constituents are a volatile, nonacrid alkaloid, red colouring matter, wax, resin, fixed oil, oleic, palmitic, and steams acids, and a minute amount of a volatile oil having the odour of paisley (Pabst Aich Phaim 230, 108)

Capsicum is largely employed as a condiment In medicine, it is used, externally as a powerful subefacient, and internally in the treatment of atonic dyspepsia, chronic alcoholism, and delirium tremens

CAPSICUM RESIN v Resins

CAPTOL. Trade name for a condensation product of chloral and tannin; used in dermatology

CAPUT MORTUUM. A term employed by

CARAMEL 21

the alchemists to denote the residue remaining tion occurs in the retort or alembic after the volatile matter of the former, acquires an orange-yellow colour, has passed away

CARAGHEEN MOSS. Irish pearl moss (Chondrus crispus) (v Carrageen, Alge)

CARAJURÚ, CARCURU, or CRAJURU Brazilian dyestuff of a red colour, said to be identical with chica-red, used by certain Indian tribes to stain their skins, and which is obtained from the leaves of the Bignonia chica (Humb et Bonpland). It is a light, mealy, odourless, tasteless powder, which acquires a coppery lustre on tuturation Insoluble in water, but soluble in alcohol, ether, and alkalis (Viiey, J. Pharm 1844, 151)

CARAMEL. The brown substance produced by the action of heat on sugar glucose or invert sugar, used for colouring liquors, beer, sweet-

meats and gravies

Preparation —On the small scale common sugar is heated, with constant stirring, in a metal pan capable of containing about ten times the quantity taken, till the whole mass melts, turns brown, and at last suddenly froths up, when the heat must be immediately stopped. Not more than 15 pc of the weight of the sugar should be driven off, as the product is hable to become insoluble if further heated

For preparation of caramel colours, Asrymusty recommends the following processes (long kept secret) --

Colour for rum

Potassium or sodium hydroxide . 3 kilos Water Or potassium or sodium carbonate Water

Dissolve with heat in large iron pan, and add Glucose 120 kilos Or syrup of glucose 130

Boil over the fire till irritating vapours begin to appear, then decrease the heat and stir continuously till the required shade is produced, then stop the process by adding 30 to 40 litres of water in a fine jet

### Colour for beer or vinegar

Ammonium carbonate	6	kilos
Water	6	,,
Glucose	120	

Process exactly as above

In the modern method of manufacture the process is conducted in a closed vessel, generally with the addition of ammonia or one of its salts (cf Salamon and Goldie, J Soc Chem Ind. 1900, 19, 301) The colouring power of the caramel seems to depend upon the amount of amn.onia used in its preparation In McCowan's process glucose is heated with from 5 to 10 p c of its weight of liquor ammonia (sp gr 0 8803 in a closed digester to 100° for above 24 hours, and the mixture concentrated

According to Magalhaes (Compt 1end 1896, 123, 896), caramel prepared from cane sugar differs considerably from caramel made from glucose, in that it behaves with certain reagents exactly like a coal-tar colour Thus, when treated with basic lead acetate and then shaken with amyl alcohol, the former caramel communicates an orange-yellow colour to the solver,, whilst with glucose criamel, no coloura- According to Gélis its composition is C<sub>36</sub>H<sub>4</sub>,O<sub>24</sub>

Ether, when shaken with a solution whilst with the latter no colour is extracted Mordanted wool is dyed orange by the former, yellow by the latter Genuine wine, to which no caramel has been added, does not give these reactions (cf D'Aquiar and Da Silva, Compt rend 1897, 124, 408)

reddish-brown. Properties - Amorphous, brittle mass, porous and deliquescent; completely soluble in water, partially soluble in alcohol, of a more or less bitter taste and non-

fermentable Melts at 134°-136°

Little is known of the chemistry of caramel Gelis (Ann Chim Phys 1858 [iii ] 52, 352; 1862, [m] 65, 496) separated it into three amorphous substances, caramelan, caramelen, and caramelin, differing in solubility and colouring power, but not of fixed composition without definite melting-points, or crystalline derivatives and readily forming colloidal solutions formation from sucrose has been represented by the following equations

(i) 
$$6C_{12}H_{22}O_{11} = 12H_2O_{-2} = 6C_{12}H_{13}O_{9}$$
 (caramelan)

$$\begin{array}{ll} \text{(i)} & 6C_{12}H_{22}O_{11} - 12H_2O \\ & = 6C_{12}H_{11}O_{9} \text{ (caramelan)} \\ \text{(ii)} & 6C_{12}H_{22}O_{11} - 18H_2O \\ & = 2C_{36}H_{18}O_{21} \text{ (caramelan)} \\ \text{(iii)} & 6C_{12}H_{22}O_{11} - 27H_2O \\ & = 3C_{24}H_{26}O_{13} \text{ (caramelin)} \\ \end{array}$$

Stolle, by heating caramelan with dilute (3 p c) hydrochloric acid, obtained a hexose, which gave an osazone crystallising in yellow needles, melting at 179°

Cryoscopic observations by Sabanéoff and Antushevitch (J Russ Chem Soc 25, 23) would seem to indicate that the molecular weight of caramelan is much higher than the above formula indicates They point out that carbon dioxide, in addition to water, is given off in its formation. They allege that it is not a carbohydrate, but has the composition

and they represent its production from sugar by the equation

$$11C_{12}H_{22}O_{11} = C_{125}H_{183}O_{50} + 7CO_2 + 27H_2O$$

Gills found on heating sucrose to a temperature of 180°-190° that a loss of weight of 12 p c gave mostly caramelan, a loss of 15 p.c. Caramelen, with a loss of 22 pc caramelin, was the main product. On treating the caramel with 84 pc alcohol the caramelan was extracted, caramelen and caramelin being insoluble in alcohol of this strength. The caramelen could be removed by cold water, when the caramelin could be dissolved out by 60 p.c. alcohol or hot water

Caramelan, the main constituent of ordinary caramel, is a brown brittle solid of bitter taste, deliquescent and very soluble in water Softens at 108° The aqueous solution gives no precipitate with metallic salts, but reduces silver nitrate and Barreswil's (Fehling's) solution An alcoholic solution of lead acetate forms a precipitate C<sub>12</sub>H<sub>16</sub>O<sub>8</sub>,PbO an ammoniacal solution of lead acetate gives C<sub>12</sub>H<sub>16</sub>O<sub>8</sub>,2PbO (Gélis)

Caramelen is of a darker brown than caramelan, and is not deliquescent. It reduces Barreswil's (Fehling's) solution and forms compounds with lead similar to those of caramelan

22 CARAMEL

Caramelin is of a still darker brown. appears to exist in three modifications of which A is soluble in cold water, B in boiling water, and C is insoluble. The modification A passes into B on evaporating its aqueous solution. The soluble forms reduce Baireswil's solution and give precipitates with metallic salts All varie-

ties have the composition  $C_{96}H_{102}O_{51}$  (Gélis)

The molecular weight of 'caramel' was found by Gladstone and Tribe to be 1700, corresponding with Gélis' formula for caramelin.

Stolle (Zeitsch Ruben-Zuckei Industrie, 1899, 49, 800; 1901, 51, 836, 1903, 53, 1149) found on heating sucrose that no carbon dioxide or other volatile products were formed below 180°, and that caramelan is the sole product when this temperature is not exceeded and the loss is not greater than 12 pc, and that its formula 15 C<sub>12</sub>H<sub>18</sub>O<sub>9</sub>

The action of heat on sucrose has been further studied by Cunningham and Dorée with special reference to the production, properties and composition of caramelan (Chem Soc Trans 1917, 111, 589) They found on heating sucrose to temperature 170°-180° that, in addition to water, pungent acid vapours, furfuraldehyde and carbon dioxide were evolved, but that if the loss of weight did not exceed 12 p.c., caramelan free from sucrose and higher caramel products was the sole product When quite div it melts at 136°, is readily soluble in water, 84 p.c. alcohol, pyridine, methyl alcohol, or hot glacial acetic acid, insoluble in ether and benzene A concentrated aqueous solution acidified with hydrochloric acid gives a red precipitate with resorcinol, soluble in alcohol or alkalis Phloroglucinol gives a similar precipitate of a deeper colour It forms a tetra-acctate, yellow powder, insoluble in water or ether, but soluble in benzene, hot alcohol, or glacial acetic acid, mp 107°, a tetra-benzoate, a pale buffcoloured powder, mp 105°-108°, soluble in acetone, chloroform, alcohol or benzene, insoluble in water, ether, or light petroleum, a tetranitrate, a yellow powder which inflames violently on warming, readily soluble in ether, alcohol or benzene, insoluble in water

It appears to be a tetra-atomic alcohol, of the molecular formula C<sub>12</sub>H<sub>18</sub>O<sub>9</sub>, or, more probably, C21H36O1, Compounds formed with phenylhydrazine and semicarbazide indicate the existence of one CO- or CHO-grouping per C21 unit Concentrated nonoxidising acids convert it into caramelin C<sub>24</sub>H<sub>26</sub>O<sub>13</sub>; weaker solutions hydrolyse and dehydrate, producing dextrose,

methylfurfuraldehyde and humic acid

### Detection and estimation of caramel.

Detection of caramel in wines, &c -10 c c of the liquid to be tested is mixed with 30 to 50 c c of paraldehyde, and enough alcohol added to make the liquids mix, if caramel is present, it forms a duty-brown precipitate sticking to the bottom of the vessel, the solution is poured off, the precipitate washed with alcohol, dissolved in a little hot water, and the solution evaporated to 1 cc, when different quantities of caramel may be compared by comparison of colours to confirm, this solution is mixed with one of 2 parts hydrochloride of phenylhydrazine in 20 of water, with excess of acetate of soda Caramel yields an amorphous brown precipitate,

To detect caramel in vinegar 50 c c, are mixed with an excess of calcium carbonate and evaporated to dryness. The mass is then extracted with 20 cc of other; the othereal solution is filtered into a test-tube containing 10 c c of resorginol reagent (1 gram of resorginol in 100 e.e. of hydrochloric acid of sp.gr. 1 125). Should caramel be present in the vinegar, a red zone is obtained at the junction of the two liquids (Ronnel, Ann Falsif 1912, 5, 517) reaction is due to the presence of hydroxy-methylfurfural, which also yields a violet colour with  $\beta$ -naphthol in sulphuric acid, and an orange-yellow colouration with phenol in sul-phuric acid. The liquid to be tested, wine, vinegai, brandy, &c, is neutralised if acid, concentrated to a small bulk, extracted with ethor, the ethereal extract treated with a few drops of a 5 pc sol of phenol in ether, and The residue is mixed allowed to evaporate with about 5 cc sulphuric acid, when the characteristic colour appears (Schenk, Apoth Zeit 1914, 29, 202, J Soc Chem Ind 1914,

Amthor's method, which is the only one capable of distinguishing between caramel and the colouring matters of colour malts, is as follows: 10 cc. of the beer or other liquid is treated with 30 50 c c of paraldehyde, according to the intensity of colour, and absolute alcohol is added until the liquids mix If the mixture is left for 24 hours a brownish precipitate forms in presence of caramel. This precipitate is washed with absolute alcohol, dissolved in hot water, filtered, and the solution concentrated to I ce in vacuó over sulphume acid. From the intensity of its colour the quantity of caramel can be estimated. Or the filtered solution of the paraldehyde precipitate may be poured in a solution of phenyl-hydrazine acetate, when if caramel be present a precipitate slowly forms, more quickly on warming / Heuss (Zeitsch ges Biauw 1918, 41, 108 124, Abti J Inst Brewing, 1918, 312)

Rough test for caramel in beer -Add a quantity of tannic acid, and shake thoroughly If malt only is present, the colour is precipi-Caramel colour remains unchanged

Contrary to the statement of Pasquero and Cappa the presence of furfural is not a sufficient proof of caramel unless the caramel has been added after fermentation

Fradiss (Chem Zentr 1899, 1 1160) determines the amount of caramel in sugar products by treating the dry substance with methyl alcohol of 95°, evaporating on the water-bath, drying the residue in vacub, or in a current of dry air, and again digesting it with methyl alcohol for 2 hours. The caramel is precipi ated from the solution by amyl alcohol, the precipitate redissolved, reprecipitated once or twice more, and finally dried at 90° and weighed caramel may also be determined by titiating with Barreswil's solution, but if dextrose is present, it is necessary to titrate the original solution before and after treating with excess of lead acetate Then if  $\Lambda - c$  c of copper solution required, a=c c of original solution required for 5 e e of copper solution, the volume of the copper solution, corresponding with the caramel, is 5 (a-A) Allowance must also be made for the amount of lead acetate solution ( vol ) used The dextrose may also be destroyed by boiling with calcium carbonate and then precipitating with carbon dioxide Traces of caramel are estimated by treating 200 c c of the solution with excess of ammoniacal lead acetate. The precipitate, after washing with water free from carbonic acid, is suspended in water and decomposed with hydrogen sulphide. The filtrate is again treated with lead acetate, and the solution, from which the hydrogen sulphide has been expelled by evaporating to 10 c c, is treated with copper solution.

Stolle (Chem Zentr 1899, ii 1099) bases a method of determining caramel by means of the spectroscope on the fact that aqueous solutions of that substance partially absorb the rays of the blue end of the spectrum. The results are

said to be accurate to within 0.05 p.c.

Methods for estimating the amount of caramel on the surface of coffee bernes roasted with sugar, have been suggested by Neubauer, Konig, Stutzer, and Hilger These methods have been tested by Fresenius and Grunhut (Zeitsch anal Chem 1897, 36, 225), who give the preference to the method of Hilger

Stutzer's method, which consists in shaking 10 grams of the unground coffee for 5 minutes with 250 c e of cold water, making up to 500 c e ofecanting, filtering, and determining the solid contents (dired at 100°), and the ash, was recommended by the Conference of Bavarian Technical Chemists Neubauer's and Konig's processes, which depend on the extraction of the betties with hot water, give too high results

In Hilger's process, 10 grams of whole coffee are digested three times for half an hour each time, with 100 c c of a mixture of equal vols of water and 90 p c alcohol at ordinary temperatures. Each portion of the extract is decanted, and the united extracts are made up to 500 c c, filtered, and residue and ash determined. It, from the numbers thus obtained, there is deducted a constant 11 07 of dry residue, or 0.83 of ash-free residue per 100 grams of dry coffee, the remainder represents the amount of caramel present.

Upsher Smith (Amer J Pharm 1911, 83, 411) suggests employing the coloured solution prepared by adding Nessler's reagent to a solution of an ammonium salt as a coloumetric test. As an arbitrary value for the colour intensity of caramel, a '100 p e' caramel is regarded as that which has such a colouration that 10 c e of its 0.1 p e solution when treated with 2 c e of the Nessler reagent and diluted to 50 c e matches the colour obtained by mixing 10 c e of a solution containing 0.0417 gram of ammonium oxalate in a litre, 38 c e of water and 2 c e of Nessler solution

CARANNA v. OLEO-RESINS

CARAPA BARK. The bark of Carapa Guanensis, has a bitter taste, and is used as a febrifuge—Said to contain an alkiloid, carapine (Petroz and Robinet, J. Pharm. 7, 293, 349)

CARAPA OIL. 'Carap Oil' or 'Crab Oil' (Fr Hude Carape), is obtained from the seeds of the 'carap,' 'crappo' or 'crab-wood' tree, Carapa Guianensis (Aubl), a native of the West Indies and Central America

The oil is prepared in Trinidad for local consumption by very primitive methods. Accord-

ing to de Verteuil (Trinidad, London, 1858, 272), the seeds are gathered in June and July, boiled for 6 hours, then laid in heaps for 8 or 10 days, during which time they undergo a sort of fermentation, they are then broken, and the pulp they contain carefully taken out and kneaded into lumps of thick paste, each about 15 lbs This paste is laid on boards slightly incurvated and inclined, and placed in a sheltered place, when the oil oozes through the mass, and runs into a vessel placed for its reception paste is carefully remoulded every morning and evening, so as to favour the disengagement of the oil After 12 days, boiling water is poured on the mass, and a fresh quantity of oil of inferior quality is thereby obtained It is a clear yellow oil with a faint vegetable odour, and an intensely bitter taste which precludes its use as an chible oil. It is used locally as an embiocation against rheumatism and for the destruction of ticks which attack domestic animals (de Verteuil The Agricultural Record, Trinidad, August, 1899, 17) It has considerable reputation as a finiment for dressing wounds on horses and cattle Its excessive bitterness repels all insects, and mixed with tai is used for preserving wood. It is well suited for the manufacture of soap, and it is in this direction that its future possibilities lie

The oil from C grandifloid, from Uganda, has been examined by Lewkowitsch (Analyst, 1908, 33, 184). Good kernels contained 30.26 p.e. of oil. The colour of the cold-pressed oil was almost white, with a tinge of pale yellow At the ordinary temperature, it solidities. The hot-pressed oil is much darker in colour, and remains solid at the ordinary temperature

Lewkowitsch (Analyst, 1909, 34, 10) has also examined the oil from *C proceia* (D.C.), the seeds being obtained from Sicra Leone The sound keinels yielded by extraction with ether 57.26 p.c. of oil, whilst the yield of oil by cold and hot extraction at a pressure of 150 atmospheres was 46.7 p.c.

The following analytical results are due to C. H. Wright (Bull Dept Agric Trimdad, ix 1910).—

•		1	11
Specific gravity at 40°/4	10°C	0.9149	)
Specific gravity at 15 5°	7/15 5°C	0 9249	0.9211
Acid value .		27.5	19 4
Saponification value		197 2	196 I
Todine value		67.7	58.5
Reichert Meissl value		38	3 6
Unsaponifiable matter			1.16

CARAWAY FRUIT. Carm fractus, B P. The dued tipe fruit of Carum Carri (Linn)

CARAWAY OIL v OILS, ESSENTIAL CARBAMIC ACID CH<sub>1</sub>NO<sub>2</sub> or NH<sub>2</sub>COOH Ammoforme acid An acid not known in the free state Ammonium carbanate is, however, met with in freshly sublimed commercial ammonium carbanate, and may be prepared by mixing gaseous ammonia and carbon dioxide Obtained also by the action of alkaline solution of potasium permanganate on albumen, leucine, tyrosine, and glycocoll

CARBAZOLE C<sub>1,2</sub>H<sub>9</sub>N or C<sub>0</sub>H<sub>4</sub> NH A white erystalline substance found amongst the products of the distillation of coal tar, and also met with in crude anthracene Carbazole

24 CARBAZOLE

compounds are most easily recovered from crude anthracene by converting the former into the easily soluble nitrosocarbazole (Wirth, Eng. Pat 14462; J Soc Chem Ind 1901, 464) May be obtained by passing vapour of aniline or diphenylamine through a red-hot tube, by boiling immodiphenyl sulphide HN  $\subset C_6H_4$   $\subset S_6H_4$ with reduced copper; by passing the vapour of orthoaminodiphenyl over lime heated to a dull redness (Blank, Ber 1891, 306), or by acting on orthoammodiphenylamine with nitrous acid and heating the phenylazimidobenzene so formed to 360° (Graebe and Ullmann Annalen, 291, 16) Forms white lamina, mp 238°, bp 3515° (con) Carbazole dissolves in concentrated sulphune acid to a yellow solution which changes Addition of formaldeto red-violet on heating hyde gives a blue solution changing to a greenishblue on addition of large quantities (Gabietti, Chem Zentr 1907, n 98) 1 mol carbazole, 2 mols, sale ylaldehyde and concentrated sulphuric acid, heated to 100°-110°, give an intense blue colour (Carrara, Gazz chim ital 1899, н [535)

By melting together carbazole and oxalic acid, and extracting with hot water and benzene, dissolving the residue in warm alcohol, filtering and evaporating, carbazole blue is obtained The alcoholic solution of carbazole blue, when treated with glacial acetic acid and zine-dust, is decolourised. Filtered into caustic soda solution, the leuco- base separates out as a white flocculent precipitate. On oxidation with any of the usual reagents, the blue colouring matter is again formed (Bamberger and Muller, Ber 1887, 1903, J Soc Chem Ind 1887, 660) zole ydlow is prepared from the tetrazo-derivative of diaminocarbazole. It is easily soluble in water, and dyes unmordanted cotton in a neutral bath a fast golden yellow and wool in a neutral or acid-bath a yellow, fast to milling (Bad Anil and Sod Fab Eng Pat 14478 and 14479, J Soc Chem Ind 1889, 771) (v Azo- colouring MATTERS)

Nitrocarbazole is obtained by treating nitrosocarbazole with 10 times its weight of benzene and its own weight of intric acid, spgr 1 36 (With, Eng Pat 2899, D R P 128853, J Soc Chem Ind 1901, 890) Dimitrocarbazoles, v Bad Anil u Sod Fab (D R P 46438, Frdl ii 447), With (D R P 128853, Frdl vi 58), Trinitrocarbazole (Akt Ges f anilinfab D R P 295817) These nitro-compounds may be reduced to the amino- compounds by means of alkalı sulphides (Wirth, D. R. P. 139568, Chem. Zentr. 1903, 1. 746). By Bvheating carbazole with concentrated sulphuric acid, a mixture of disulphonic and trisulphonic acids is obtained. Nitration and subsequent reduction of the former yields aminocarbazoledisulphonic acid, which, when diazotised and coupled with  $\beta$ -naphthol, yields a dyestuff dyeing red from an acid-bath. If the nitic acid is added direct to the products of sulphonation, on reduction a second amino- acid is obtained which, when diazotised and coupled with β-naphthol, gives a red dyestuff, and with a-naphthylamine a violet (Schultz and Hauenstein, J pr Chem 1907, 336) The sulphonic acids, preferably the disulphonic, condense with tetralkyldiamino diphenylearbinol to yield

valuable green dyest iffs (Cassella & Co. Eng. Pat 20709; J. Soc. Chem. Ind. 1909, 697). By allowing nitrosophenol, its homologues or derivatives to act on carbazole in the presence of cone sulphuric acid, compounds are obtained which yield blue sulphide dyestuffs with alkali polysulphides (Haas, Eng. Pat. 2918; J. Soc. Chem. Ind. 1909, 517) For N-acyl derivatives of carbazole, see Copisarow. Chem. Soc. Trans. 1918, 816

A method has been patented (Ch. Fab. Act. Gesell in Hamburg, D. R. P. 81237, Ber. 1895, iv 692) for obtaining potassium cyanide or feirocyanide, by heating the potassium derivative of carbazole in the absence or presence of an

CARBITE. An explosive consisting of nitroglycerine (25), potassium nitrate (34), flour (38-5), barium nitrate (1), powdered bark (1), and sodium carbonate (0.5 part)

CARBODYNAMITE V EXPLOSIVES CARBOHYDRATES. The term is applied to polyhydroxy-aldehydes and ketones, and to the substances which give these when hydrolysed by heating with mineral acids Dextrose. ordinary sugar or sucrose, starch, and cellulose are carbohydrates universally distributed in plants Dextrose and glycogen enter into the composition of animal tissues. The simple carbohydrates, monosacchandes, have the empirical composition CH2O, the most important being those containing 5 or 6 atoms of earbon The complex carbohydrates, polysacchandes, are built up from two or more simple carbohydrates with the elimination of water, they have the composition  $(C_8\Pi_{10}O_5)_n\Pi_2O$  (Kiliani, Chem Zenti 1908, 32, 366) Compounds in which nequals 1, 2, 3, or 4 are distinguished by names having the suffix -ose They may be conveniently grouped as follows -

1 Monosaccharides

(a) Pentoses  $C_5H_{10}O_5$ , eg arabinose, xylose (b) Hexoses  $C_6H_{12}O_6$ , aldoses, eg dextrose, mannose, galactose, ketoses, eg lavulose, sorbose

2 Disacchandes n=2  $C_{12}H_{22}O_{11}$ (a) Non-reducing, e.g. sucrose, trehalose

(b) Reducing,  $e \stackrel{g}{g}$  maltose, lactose, &c 3 Trisaccharides n=3  $C_{13}H_{32}O_{16}$ tianose, raffinose, melezitose

4 Tetrasacchandes n=4 C<sub>24</sub>H<sub>12</sub>O<sub>21</sub> Stachyose

5 Polysaccharides

(a) Glycogen, mulin, dextrins, n unknown

(b) Starch, cellulose, n very large

Related to the carbohydrates are those substances which yield simple carbohydrates, together with other substances when hydrolysed Such are glucosides, gums, pectins, &c (q c)

General characters -The members of classes 1, 2, 3, 4, and 5a are soluble in water They have a more or less pronounced sweet taste The monosaccharides are colourless crystalline compounds, characterised by the power of reducing alkaline solutions of the heavy metals, for example, copper They form characteristic compounds with phenylhydrazine

The di- and poly-sacchandes are converted into simple carbohydrates when hydrolysed This change may be effected by means of mineral acids, or, as a rule, more quickly by enzymes The complicated polysacchandes are usually amorphous, they tend to form colloidal solutions, and are chemically relatively mert. In virtue of the hydroxyl- groups which they contain, they react with acetyl chloride, benzoyl

chloride, nitric acid, &c, forming esters

Carbohydrate solutions are generally optically The direction and amount of the active rotation vary with the substance, and also with the strength of the solution, the nature of the medium, the temperature, &c The optical rotatory power of carbohydrates is made use of in their analyses (r Saccharimetry), and as a guide to their structure

Interesting generalisations have been drawn by Hudson affecting the molecular rotations of  $\alpha$  and  $\beta$  forms of sugars now generally accepted that the difference between the molecular rotations of the  $\alpha$  and  $\beta$  forms of mutarotatory sugars is a constant The compounds selected show considerable diversity of type and discordant results have from time to time been explained away by showing that the compound in question is derived from other than the  $\alpha$  and  $\beta$  forms of the sugar (of Hudson, J Amer Chem Soc 1917, 39, 1013) The maximum values for the specific iotations are now arrived at by the indirect method based on solubility measurements (Lowry, Chem Soc Trans 1904, 85, 1551, Hudson, J Amer Chem Soc 1904, 26, 1065) The asymmetric system attached to the  $\alpha$ -carbon atom excits a preponderating influence on the activity of these compounds and determines the sign of the rotation

Hudson's generalisation have enabled the optical rotatory power of carbohydrate derivatives to be used as an important guide to their structure

Constitutional formula of Dextrose -In addition to those carbohydrates which occur naturally, a number of others have been prepared artificially by Emil Fischer, and it is largely to his work that the present complete knowledge of the group is due (Fischer's original work has been republished, Untersuchungen über Kohlenhydrate, Berlin, 1909, J. Springer)

Dextrose and its isomerides C6H12O6, conveniently termed glucoses, have 5 oxygen atoms present as 'hydroxyl,' since they form esters such as dextrose pentacetate

#### C<sub>6</sub>H<sub>7</sub>O(O CO CH<sub>3</sub>)<sub>5</sub>

The sixth oxygen behaves as an aldehyde in dextrose and galactose, as a ketone in Levulose and sorbose. The two classes of compounds are accordingly referred to as aldoses and ketoses Reduction converts the glucoses into alcohols  ${\rm C_6H_{14}O_6},$  dextrose yielding soi bitol and mannitol, and galactose forming dulcitol Secondary hex liodide, a derivative of normal hexane, is formed by the action of hydrogen iodide, hence the glucoses must possess all their six carbon atoms united in normal chain

Dextrose on oxidation yields, in the first place, gluconic acid C5H6(OH)5 CO2H which retains all six carbons and, like dextrose, forms a pentacetyl ester Hurther oxidation converts it into saccharic acid  $C_0H_{10}O_3$ . Galactose behaves similarly, yielding the isomeric galactonic, and mucic acids. Taking further into consideration the reducing power of dextrose, the constitutional formula may be written

 $\mathrm{CH}^{3}(\mathfrak{O}\Pi)$  СП(ОН) СП(ОП) СП(ОП) СП(ОН) СПО

This formula does not fully express the behaviour of dextrose, which lacks many of the well-attested properties of the aldehydes; for example, it is not oxidised in the air, and does not show their various colour reactions Tollens, in 1883, therefore, proposed the lactonic or  $\gamma$ -oxide formula (i), which may also be written (II).

This formula (1 or 11) is in entire agreement with the reactions of dextrose, and has been generally adopted It is supposed that in presence of alkalis and other reagents the  $\gamma$ -oxide ring opens, forming the aldehyde

The discovery of a third methyl glucoside (Fischer, Ber 1914, 47, 1980) and of additional isomerides of other glucose derivatives (Hudson, J Amer Chem Soc 1915, 37, 1589, 1591, 1916, 38, 1223) has involved the recognition of a different internal linking in the sugar molecule The oxygen atom may connect other pairs of atoms as well as the 1st and 4th The new form of sugar termed  $\gamma$ -glucose (dextrose) obtained from this glucoside has not been isolated in the free state, but its tetramethyl derivative is known (Irvine, Chem. Soc. This is far more Trans 1915, 107, 524) reactive than the corresponding derivatives of α or β-dextrose it condenses with alcohol with extreme ease and instantly reduces alkaline permanganate solution, giving rise to a new form of tetramethyl gluconic acid

The work of Tanret (Compt rend 120, 1060) has shown that dextrose exists in more than In addition to the hydrated and anhydrous modifications, he isolated two other anhydrous modifications These differed particularly in optical rotatory power Taniet described as a dextrose having  $[a]_D + 110^\circ$  when first dissolved, and falling in aqueous solution, a  $\beta$ -dextrose having  $[\alpha]_D + 19^\circ$ , and increasing in solution, and a dextrose having  $[a]_D + 52.5^{\circ}$ , and remaining constant in solution modification was called  $\gamma$ -dextrose by Tanret It is not the same, of course, as the  $\gamma$ -glucose mentioned above lt had long been known that the optical rotatory power of fieshly dissolved dextrose diminished to about one-half its initial value in solution, and the term mutarotation or bi-rotation was applied to this phenomenon

Dubrumfaut ascribed the change to purely physical causes Fischer considered that the aldehyde dextrose underwent hydration to an alcohol C<sub>6</sub>H<sub>44</sub>O<sub>7</sub> of lower rotatory power view of Taniet's discoveries, it is now considered that the change in rotatory power is due to the mutual interconversion of the  $\alpha$ - and  $\beta$ -dextioses, until equilibrium is attained Tanret's y-dextrose represents such an equilibrium mixture. It is supposed that both  $\alpha$  and  $\beta$ -dextrose have a closed y-oxide ring structure, and it is still a matter of controversy whether some proportion of open-chain aldehyde is present in solution in the equilibrium mixture (For the full discussion of this question, see E. F. Armstrong's The Simple Carbohydrates, Longmans, 1919)

To understand fully the relation of  $\alpha$ - to 8dextrose, it is necessary to refer to the van't Hoff-Le Bel conception of space isomerism Chemical compounds which rotate the plane of polarised light contain an asymmetric carbon atom, that is, one in which the four affinities are satisfied by four distinct radicals earbon atom is considered as lying in the centre of a tetrahedron, and each of the four different radicals with which it is combined as being at the four solid angles thereof. Two modifications of the compound are possible, which are related to one another like an object and its reflected image

In the Tollens formula for dextrose, the carbon atom (C) has the four different groups, H, OH, the oxygen of the ring, and the complex  ${
m C_5H_{10}O_4}$  attached to it—It is asymmetric, and purplicating acid  ${
m C_4H_9}$  (HMcCO.H—This betwo isomerides are possible, which are representation establishes Levulose as a ketohexose. It sented on a plane surface by writing H and OH on one or other side of the carbon atom, as the following formulæ show

Corresponding to these two isomenic forms, dextrose gives use to a double senes of denvatives

The compound represented by the formula си<sub>2</sub>(он) си(он) си(он) си(он) си(он) сио contains four asymmetric carbon (marked \*), and should, therefore, be capable of existing in 16 stereoisometic forms, 8 of which would be mirror images of the other 8, and of equal but opposite rotatory power

The carbohydrate group has afforded a unique opportunity of testing this hypothesis, and although only three of the isomerides occur naturally, no less than 12 have been prepared artificially by Fischer, who has, in addition, made them in such ways that then structure is established

The known aldohexoses are the dextro- and lævo- isomerides of mannose, dextrose, idose, gulose, galactose, and talose Two further isomerides, namely, d-allose and d altrose, were obtained by Levene and Jacobs in 1910 (Ber 43, 3141), starting from d-ribose This is a component of many nucleosides, and so a considerable quantity had been accumulated for the purpose of applying the cyanohydrin process It will suffice to give the constitutional formula as established by Fischer of the natural members -

Mannose and galactose, like dextrose, exist in  $\alpha$ - and  $\beta$ - forms, and it is more correct to write their formulæ in the form adopted for  $\alpha$ and B-dextrose

As is to be expected, the isomeric glucoses differ but little in their chemical behaviour

Formula of Lavidose —Lævulose, when oxidised, forms glycollic and trihydroxyglutaric acid, or other acids with less than 6 carbon atoms It forms an additive compound with hydrogencyanide, which yields lævulose carboxylic acid C<sub>7</sub>H<sub>14</sub>O<sub>8</sub> on hydrolysis, and this, when boiled with hydriodic acid, is converted into methyl-. has the formula

Fructose shows a pronounced tendency to exist in reactive forms other than the normal  $\gamma$ -oxide type—probably many of its compounds, including sucrose, are derived from the ethylene oxide type

The ethylene oxide form of methyl fructoside readily condenses with acctone and rapidly reduces permanganate

Formulae of the Pentoses and Tetroses —Eight aldopentoses are possible, of which seven have been artificially prepared, viz d- and l-arabinose and xylose, d-lyxose, and d- and l-ribose. The two most common naturally occurring pentoses have the constitutional formula-

Like dextrose, they occur in two forms, and are more correctly written with a y-oxide A pentose believed to be d-arabinose structure has been isolated from the glucoside barbaloin

(Léger). Levene (Ber 1909, 42, 3247; 1910, 43, 3147) obtained d-ribose from nucleic acid

Three out of the four possible aldotetroses are known, viz. d- and l-crythrose and l-threose, but they have not been found in nature

Reaction with Phenylhydrazine—The behaviour of the carbohydrates with phenylhydrazine is particularly characteristic, and it has further an historical interest, as phenylhydrazine, in the hands of Fischer, served as one of the chief aids in the elucidation of the relationships of the group.

Dextrose reacts with the reagent in acctic acid solution in two stages. In the cold a phenylhydrazone is formed

$${\rm C_6H_{12}O_2}{+}{\rm C_6H_5} {\rm NH~NH_2 \atop C_5H_{12}O_5~CH}$$
 . N NH  ${\rm C_6H_5}{+}{\rm H_2O}$ 

This is a colourless substance, soluble in water, and is obtained in two modifications, one or the other predominating according to the method of preparation. Similar compounds, with asymmetrically disubstatuted hydrazine—for example, benzylphenylhydrazine—are often less soluble and more characteristic. Mannose is remarkable in forming an almost insoluble phenylhydrazone. The hydrazones of lactose and maltose are very soluble.

At the temperature of boiling water, dextrose reacts with excess of phenylhydrazine to form the phenylosazone

Dextose (1 molecule) is heated with 3-4 molecules of phenylhydrazine, which should be almost colourless, and free from oxidation products, and an equal volume of 50 p.c. acetic acid in a flask immersed in boiling water for an hour or more. Usually the osazone begins to separate in 20 minutes or less. In the case of the disaccharides, no separation of osazone takes place from the hot solution.

The yellow osazone is filtered and carefully washed, it may be crystallised from most organic solvents, but the most suitable appears to be a dilute solution of pyridine. Dextrose, mannose and Levulose form the same phenylosazone, as they differ only in the relative positions of groups which are destroyed in the formation of the osazone.

Too much dependence must not be placed on the melting-points of the osazones in identifying unknown sugars in solution, as these depend very largely on the rate of heating, and the method of purification adopted. Disubstituted hydrazines do not directly form osazones with dextrose, though these may be prepared from the osones.

Diphenylmethanedimethyldihydrazine

(J. von Biaun, Bei 1917, 50, 42, Votoček, Ber 1917, 50, 35) appears likely to be of use in determining the configuration of aldoses Hydrazones are only produced when at least two of the CH(OH) groups following the aldehyde group have the same spatial airangements Thus hydrazones are formed with airabinose, lyxose, rhamnose, ribose, rhodeose, fucose, galactose, mannose, but not with xylose, isorhodeose and glucose

To recover the original sugar from the

phenylhydrazone, it is boiled in water with a slight excess of benzaldehyde. The phenylhydrazine is removed as benzaldehyde phenylhydrazone, the excess of aldehyde extracted by ether, and the sugar solution concentrated in warno. With the disubstituted hydrazones better results are obtained on using formaldehyde.

The phenylhydrazine residues may be removed from the osazones by heating with fuming hydrochloric acid. Compounds termed osones are formed.

### CH<sub>2</sub>(OH) [CH (OH)]<sub>3</sub> CO CHO

These are colourless syrups, they act as strong reducing agents, combine directly with phenylhydrazine, and are non-ferimentable. The slightly soluble osazones of the disaccharides, and of the pentoses, are converted into osones on boiling with benzaldehyde.

When reduced by means of sodium amalgam, the osones are converted into ketoses, and not into the aldoses from which they were formed. Thus it is possible to pass from dextrose to levulose—a transformation of great importance—the stages being

dextrose→ phenylosazone→ osone→ lævulose

Another important transformation is that from dectrose to mannose, which is based on the fact that gluconic acid (q|r|), when heated with quinoline, is converted into a mixture of gluconic and mannonic acids. The stages in this case are

$$\begin{array}{c} \text{oxidation} & \text{reduction} \\ \text{Dextrose} \rightarrow \left\{ \begin{array}{c} \text{gluconic} \\ \text{acid} \end{array} \right\} \rightarrow \left\{ \begin{array}{c} \text{mannonic} \\ \text{acid} \end{array} \right\} \rightarrow \text{mannose} \end{array}$$

By the aid of these two reactions, Fischer was able to pass from the synthetical  $\alpha$ -across to d-glucose, d-fructose, and d-mannose, and so effect the complete synthesis of the naturally occurring hexoses

The scheme on next page shows the successive operations

Cyanohydrin synthesis.—The aldoses combine directly with hydrogen cyanide, forming nitriles, which, when hydrolysed, give rise to acids containing one carbon atom more than the original carbohydrate. The lactones of these acids, when reduced with sodium amalgam, yield the corresponding aldoses with one carbon atom more than the original carbohydrate:

$$\begin{array}{l} C_5 H_{11} O_5 \text{ CHO} + \text{HCN} = C_5 H_{11} O_5 \text{ CH(OH) CN} \\ C_5 H_{11} O_5 \text{ CH(OH) CO}_2 H \longrightarrow C_5 H_{11} O_5 \text{ CH(OH) CHO} \end{array}$$

In this manner arabinose is converted into dextrose, dextrose into glucoheptose. Fischer has continued the process as far as the aldononose.

Actually two isomerides are formed at each stage, and arabinose gives both dextrose and mannose

Degradation—The converse change from dextrose to arabinose has been effected by several different methods

Wohl (Ber 1893, 26, 730, &c) heated dextrose oxime with concentrated sodium hydroxide, so converting it into glucome acid nitrile, from which, on further heating hydrogen cyanide was eliminated and d-arabinose formed

$$C_5H_{11}O_5 CHO \Rightarrow C_5H_{11}O_5 CH NOH \Rightarrow C_5H_{10}O_5$$



In practice the method adopted is to heat the oxime with acetic anhydride and a grain of zinc chloride; a vigorous reaction ensues, and the pentacetate of glucome acid mitrile is formed, from which hydrogen cyanide is eliminated by treatment with ammoniacal silver oxide

Ruff (Ber 1898, 31, 1573) oxidises the calcium salt of gluconic acid, with hydrogen peroxide in presence of a ferrous salt (Fenton's method). The carboxyl-group is eliminated, and arabinose formed

Neuberg (Biochem Zeitsch 1908, 7, 527) effects the climination of the elements of formic acid from glucome and similar acids by electrolysis

Weerman (Chem Soc Abstr 1915, 1 387) obtains gluconamide by saturating gluconolactone in alcoholic solution with ammonia On subsequent treatment with hypochlorous and the amide is decomposed according to the scheme

R CH(OH) CO NH<sub>2</sub>
$$\rightarrow$$
R CH(OH) N C O $\rightarrow$ R CHO

According to Livine the reactions proceed with the utmost smoothness and constitute a great improvement on the methods litherto in use

The reaction has been applied to glucose, galactose, l-mannose and l-arabinose

Action of Acids —Dilute acids are without action on the aldoses, but when submitted to the continued action of acids, hexoses yield levulic acid (acetopropionic acid)

and formic acid, together with brown or black humus substances The first product from the 2-ketohexoses is 4-hydroxymethylfurfuraldehyde

and it is to the condensation products of this aldehyde that the colour reactions of the ketoses with phenols are due—Blanksma and Ekenstein have shown that this aldehyde is also produced on heating mannose or dextrose with oxalic acid, and Fenton obtained 4-halogenmethylfurfurol from Levulose, and more slowly from the aldoses, on treatment with dry halogen hydride—The aldoses apparently undergo a 1-, 2-enolisation and conversion into 2-ketohexose.

Action of Alkalis —Although absolutely stable in pure aqueous solution, dextrose, nannose,

and fructose pass over into one another in presence of traces of alkalis in the cold. The optical rotation of a dextrose solution containing alkali, for example, slowly falls to zero. This change was observed by Lobry de Bruyn and Alberda van Ekenstein (Rec. trav. Chim. 1895, 14, 150, 204). It is due to the conversion of the sugar originally present into enol.

 $CH_2(OH)[CH(OH)]_3 C(OH) CH(OH)$ 

which is reconverted into all three hexoses

Guandine (Morrell and Bellars, Chem. Soc. Trans 1907, 91, 1010-1033) brings about similar changes.

The action of alkalis on the carbohydrates is exceedingly complicated, and leads to the formation of a great variety of substances, in addition to humus compounds. The concentration of the alkali is of great importance, a larger number of products being formed when the relatively sparingly soluble lime and bartya (Kihani) are used, than when concentrated sodium hydroxide is employed (Nef, Annalen, 1910, 376, 1, 1914, 403, 204) The latter paper contains a summary of many years' work, and is the main source of information on this complex subject In contact with dilute alkalis protound changes ensue in a hexose, an equilibrium being finally established between the various enolic forms The number of possible forms is diminished as the concentration of the alkali 18 reduced The unsaturated molecules formed undergo fission at the double bond

The decomposition takes place quickly on heating, slower in the cold, with the production of acetol, acetone, lactic, acetoe and formic acids, also glucinic and saccharumic acids, and other substances. With calcium hydroxide saccharim or calcium saccharimate is produced

In addition to these substances, intermediate products are formed, more closely related to the monosaccharides, and from which these are reconstructed by the synthetic action of the alkali. It is established that hexoses break down into a molecule each of diose and aldotetrose, or into 2 molecules of glycerine aldehyde, that is, the 2:3 and 3 1-dienois decompose. The 1:2-dienois never break down into aldopentose and form aldehyde. Conversely, hexoses are never built up from pentoses.

Pentoses form 1.2-dienols, which, in the main, give addoted oses and hydroxymethylene on decomposition, they also form 2.3-dienols, breaking down to diose and glycerine aldehyde

Dextroses exclusively form 2 3-dienols, breaking down to 2 molecules of diose, they never give 1:2-dienols yielding hydroxymethylene and glycerine aldehyde. The strongly dissociated zine hydroxide ammonia acts on the monosaccharides in the cold, producing methyl-

glyoxalıne | CMe NH GH•N (Wındaus, Beı 1905,

38, 1166, 1906, 39, 3886, 1197, 40, 799)

Saccharinic acids. According to Net (Annalen, 1910, 376, 1) there are 24 isomeric saccharinic acids, with 6 carbon atoms, namely a-meta-saccharinic acids ( $a\gamma\delta\epsilon$ -tetrahydroxy-hexoic acids).

CO<sub>2</sub>H CH(OH)·CH<sub>2</sub> CH(OH) CH(OH) CH<sub>2</sub>OH derived from the 16 aldohexoses, 4 to sacchar-

mic acids (a-hydroxymethyl  $\alpha\gamma\delta$ -trihydroxyvalerie acids).

 ${\rm CO_2H~C(OH)(CH_2~OH)~CH_2~CH(OH)~CH_2(OH)}$  derived from the 8  $\beta$ -ketohexoses , 8 saccharing acids ( $\alpha$ -methyltetrahydroxyvalene acids).

CO<sub>2</sub>H CMc(OH) CH(OH) CH(OH) CH<sub>2</sub>(OH) derived from the 8  $\gamma$ -ketohexoses; and 1 parasaccharinic acids (a,  $\omega$ -hydroxyethyltinhydroxybutyiic acid).

CH<sub>2</sub>(OH) CH<sub>2</sub>·C(OH)(CO<sub>2</sub>H) CH(OH) CH<sub>2</sub>(OH) The lactones of these acids are known as saccharins. They give use to characteristic alkaloid salts and phenylhydrazides, and possess optical activity. The saccharins are obtained by the action of lime on dextrose, maltose, or lactose.

Synthesis of Monosaccharides —The condensation of formaldehyde under the influence of bases to carbohydrates has been the subject of repeated study. Butlerow, using milk of lime, obtained an uncrystallisable syrup with a sweet taste, which behaved as a sugar, but was nonfermentable and optically mactive Ho termed the product methylemian Locw improved the technique of the method by saturating a 4 p c. solution of formaldehyde with slaked lime in the cold, and allowing it to stand The syrupy product was termed 'formose,' and supposed by Loew to be homogeneous, but subsequently Fischer showed that on treating it with phenylhydrazine, a mixture of osazones was formed, one of which was identical with a-acrosazone Net has given reasons for believing formose (prepared by means of lead oxide) to be a mixture in approximately equal parts of isometic hexoses and pentoses its composition obviously depends on the alkali employed and its concentration

Fischer and Tafel effected the condensation of acrolem dibromide by means of baryta, and showed the syrupy product to consist of two sugars, distinguished as a- and B-acrose. Subsequently, crude glycerose was made the starting-point of the synthesis. It is a mixture of glycerine aldehyde and dihyroxy acctone, both of which substances have, in after-years, been separately used as the initial material. Fenton has used glycollic aldehyde in the same manner

A product of synthesis by all these methods is a acrose, which is partly fermentable by yeast, and this Fischer isolated as phenylosazone, and showed to be identical with mactive dl-lævulose. From this by the series of reactions already given, he obtained the natural d-glucose, d-mainose, and d-fluctose (lævulose), as well as their optical antipodes and a number of new isomerides. The application of the methods of degradation, elsewhere described, enabled pentoses and tetroses to be synthetically prepared from these hexoses, whilst the cyanolydim reaction made it possible to obtain monosaccharides with 7, 8, and 9 carbon atoms.

The full explanation of the synthesis of carbohydrates in the green leaves of growing plants has still to be given. It is generally supposed that formaldehyde, the first product of assimilation, undergoes polymerisation to carbohydrates, a theory advanced by Bacyer in 1870 (Ber. 3, 63). Various compounds, such as glycolhe and glyceric aldehydes and dihydroxy

actione, have been suggested as intermediate stages in the polymerisation, but it has never been possible to detect any of these in the plant. Apparently the synthesis is asymmetric, hexoses of the dextro-series, and their polysaccharides being the only products of assimilation. It is supposed that the formaldehyde elements are superposed under the influence of the asymmetry of the protoplasm molecule, according to a definite plan, and are split off when six are minted.

Synthesis of Disaccharides—Though often attempted, the synthesis of disaccharides, by purely chemical means, has only been partially successful. Fischet and Armstrong have prepared a syrupy compound, probably identical with the natural sugar melibiose, by the interaction of acctochlorogalactose with sodium glucosate.

Fischer and Delbruck found that when actiobromoglucose is shaken in dry ethered solution with silver carbonate and traces of water, the octacetyl derivative of a disaccharide is obtained. This is hydrolysed by cold barium hydroxide to a non-reducing disaccharide of the type of trehalose. The octacetyl-derivative is a mixture of isomerides, and yields isomeric disaccharides.

disacchandes.

In the plant the di-acchandes are formed from hexose by the agency of enzymes, which exercise synthetic as well as analytic functions. It is undecided whether the same enzyme can cause both synthesis and hydrolysis, or whether special enzymes alone act as synthetic agents. It is proved, however, that an enzyme effects the synthesis of a carbohydrate isometre with that which it normally hydrolyses, the two compounds being apparently related in the same manner as the  $\alpha$ - and  $\beta$ -methyl glucosides (v- Croft Hill, Chem. Soc. Trans. 1898, 73, 634, 1903, 83, 587, E. F. Armstrong, Prot. Roy. Soc. 1905, 76, B, 592, and monograph on Carbohydrates, 1919)

It has been established by Bourquelot that the equilibrium between a sugar and a glucoside when in contact with the appropriate enzyme can be displaced in the direction of the glucoside when the solvent contains excess of the hydroxy component of the glucoside. A large number of glucosides have been synthesised by Bourquelot from sugars dissolved in alcohols containing varying amounts of water in every case the glucoside synthesised being that normally hydrolysed. This reaction, however, is not on all fours with that taking place in a concentrated solution of sugar in the absence of an alcohol, and so far all attempts to synthesise maltose or sucrose in this manner have been definitely without result.

In the following pages the occurrence, preparation, properties, and derivatives of the natural, mono-, di-, tii-, and tetra-sacchandes, also glycogen and mulin, are dealt with Sucrose, the dextrins, starch, and cellulose, are described in separate articles.

**Dextrose.** Dextroglucose, grape sugar, starch sugar  $C_6H_{12}O_6$  and  $C_6H_{12}O_2$ ,  $H_2O$ .

Occurrence—Reducing sugars are very widely distributed in nature, but dextrose is almost always accompanied by lævulose, the two hexoses being probably derived from sucrose. It is thus found in honey, in the juice of most

fruits, and generally in vegetable tissues in small quantities. In the animal kingdom it is found in blood and in diabetic urine.

Formation — Dextrose is formed on hydrolysing most of the known polysaccharides and glucosides, either with acids or enzymes, it is similarly formed from glycogen. Thus cellulose, statch, dextrins, the disaccharides maltose, lactose, and sucrose, all yield dextrose.

Preparation —Dextrose is usually prepared from starch or sucrose—It can be obtained from any of the substances mentioned, but they are less suited for the preparation in quantity

Crude commercial dextrose is purified by dissolving in 8-10 pc of hot water, and adding dry boiling methyl alcohol, until a cloud is produced. This solution may be decolourised by shaking with animal charcoal, and filtering. On standing, anhydrous dextrose in a pure state separates abundantly. To prepare dextrose from sucrose, the latter must first be inverted. This is best effected by treatment at a temperature not exceeding 50°, with a small quantity of invertase or of pressed yeast. After a few hours the solution is evaporated in vacu6 to a syrup, which solidities on standing. The mass is washed with cold ethyl alcohol (sp.gr. 0-830), and the residue dissolved in more alcohol. Dextrose crystallises on cooling.

Sulphure acid may be used for inversion, it must be neutralised with baryta before evaporation. Soxhlet advises the use of alcoholic hydrochloric acid at 50° for 2 hours. Dextrose crystallises in a few days without neutralisation.

Properties - Dextrose crystallises from strong (93-95 pc) ethyl or methyl alcohol in the anhydrous state,  $({}^{\circ}_{3}\Pi_{12}O_{6}, \text{ and from concentrated aqueous solutions at }30^{\circ}-35^{\circ}$ , with special treatment, crystals of this composition may also be obtained. The crystals formed in aqueous solutions at ordinary temperatures contain a molecule of water of crystallisation.  $C_6H_{12}O_6, H_2O_6$ The anhydride forms white crystalline crusts or powder and, when it separates from the solvent slowly, fairly well-defined glassy prisms, the hydrate crystallises in cauliflower-like masses, but with care and slow erystallisation well-defined crystals can be obtained Both bodies dissolve easily in water, the anhydride with evolution of heat. The hydrate dissolves freely in strong ethyl or methyl alcohol, especially on the application of heat, but the anhydride is practically insoluble in those liquids

a-Dextrose exists in contact with boiling saturated alcoholic solutions it forms hard brittle needles. From aqueous solutions monohydrate crystallises in six sided leaflets. As purified by crystallisation from dilute methylaleohol a mixture of the different forms is obtained. A homogenous product is obtained by allowing the mixture to soak several days or weeks at a constant temperature with the solvent until the whole of the  $\beta$ -sugar present has been converted into the  $\alpha$ -isomeride Hudson and Dale (J. Amer Chem Soc 1917, 39, 320) obtain the  $\alpha$ -form by dissolving two parts of sugar in one part of water and adding 4 parts of glacial acetic acid, crystallisation being allowed to take place at the ordinary temperature

β-Dextrose is forme by heating dextrose at

105°-110° for some hours or by crystallisation from water above 98° From a solution in boiling pyridine a monopyridine salt separates which on exposure rapidly loses pyridine An even better method of prepara-(Behrend) tion is said to be the following (Hudson and Dale) 10 parts of dextrose are dissolved in 1 part of water on a water bath, and 12 parts of glacial acetic acid heated to 100° are added. The whole is well mixed and removed from the water bath when crystallisation at once commences After four such crystallisations pure  $\beta$ -dextrose is obtained

 $\gamma$ -Dectrose (the equilibrium mixture) is obtained on precipitating with alcohol a concentrated aqueous solution which has stood for some time

The initial optical rotatory powers of the two forms and that of the equilibrium mixture are as follows in aqueous solution -

$$a-[a]_{D}+113 \ 4^{\circ}$$
  
 $\beta-[a]_{D}+19 \ 0^{\circ}$   
 $\gamma-[a]_{D}+52 \ 2^{\circ}$ 

(Hudson and Dale, J. Amer. Chem. Soc. 1917, 39, 320)

An aqueous solution of dextrose containing 10 grams anhydrous dextrose in 100 cc of solution at 155°, has a spgr 10385 Its solutions are optically active, ie possess the power of rotating the plane of polarisation of a ray of polarised light. The rotation is righthanded Hesse (Annalen, 176, 106) and Tollens (Ber. 17, 2234) have carefully determined this constant, and the latter's results are expressed by the formula

 $[a]_D = 52.5^{\circ} + 0.018796P + 0.00051683P^2$ for the dry sugar, and

 $[\alpha]_D = 47.73^{\circ} + 0.015534P + 0.0003883P^2$ 

for the hydrate  $C_6H_{12}O_6, H_2O$ , in which P =the percentage of anhydride and hydrate in solution In dealing with solutions containing 10 p e diy sugar, the factors  $[a]_D = 52.7^{\circ}$  and [a] = 58 3° may be used The optical activity of dextrose solution is not materially affected by temperature. The activity is nearly twice as great in freshly prepared solution, diminution taking place slowly in cold, but rapidly in boiling solutions, until the constant value is arrived at It is only crystallised dextrose that exhibits this: property of bi-rotation, as it is called, dextrose fused and allowed to solidify does not show it (Hesse, Annalen, 176, 113) The solution of devtrose in strong alcohol retains the power of bi-rotation.

When dextrose is heated to 170°, the elements of a molecule of water are eliminated, dextrosan (glycosan)  $C_6H_{10}O_5$  being produced. If the temperature is increased, that body is caramelised and chaired Dextrosan is again converted into dextrose by digestion with water or dilute acids It is a hygroscopic, dextro-rotatory, non-fermentable substance If the temperature is raised to 200°, and maintained at that point without being increased until these products calciumdextrosates are described containing cease to be evolved, the brownish-black residue varying proportions of dextrose, lime, and (caramelised mass) becomes gradually solid, water. Dextrose is not precipitated with lead

This substance is soluble in water, completely if the temperature was not too high, and constitutes the colouring matter used in winc, beer, cookery, &c Should the heat be further continued so that the body is destructively distilled, carbon mon- and di-oxide, and methane are evolved, aldehyde, furfuraldehyde, acetone, propionic, acetic, and formic acids, furane and furfurol distil over, and a black carbonaceous V CARAMEL mass is left

The action of alkalis and of acids on dextrose has been discussed elsewhere. When reduced by sodium amalgam, the main product (35-10 pe) is sorbitol, some quantity of mannitol is also formed in alkaline, but not in neutral or acid solution. It is derived from the mannose produced by the action of alkali on dextrose

Oxygen has no action on dextrose in neutral or acid solutions, but alkaline solutions absorb the gas Ozone oxidises it to formic acid, saccharic acid, &c., being produced. Strong nitric acid converts dextrose into saccharic acid, and finally into exalie, carbonic, and formic acids Treated with chlorine or bromine in the presence of water, the resulting product yields gluconic acid when submitted to the action of silver oxide

Dextrose in alkaline solutions reduces the oxides of gold, platmum, silver, mercury, bismuth, lead, copper, non, manganese, &c , in the case of gold, platmum, silver, and mercury, the reduction is complete, the metal being formed, in the other cases lower oxides are produced Ferrocyanide of potassium, indigo, pieric acid, litmus, and such substances are also reduced by dextrose in alkaline solution. The products from the dextrose are not the same in all cases, but they consist chiefly of carbonic, formie, oxalie, and glycollic acids, other products being also formed

Nearly all the true yeasts (Saccharomyces membrana factors is an exception), a great number of torulæ, moulds, and other organisms ferment dextrose, yielding carbon dioxide and alcohol as the chief products—Glycerol, succinic acid, and other bodies (fuscl oil) are formed at the same time, the proportions of these bodies varying materially with the organisms and the conditions under which the growth takes place The work of Ehrlich suggests that most of these by-products are derived from amino-acids, produced from the yeast and not from dextrose (v. Fermentation)

Pressed yeast-juice, zymase, which contains no living protoplasm, also ferments dextrose

In the presence of growing bacteria of various species, dextrose yields many products, one species converts it into lactic acid, another into butyric acid, another into cellulose, the so-called slimy fermentation, and so on Other oxidise dextrose (v Fermentation) Other organisms

Dextrose combines with sodium chloride, bromide, and iodide, forming well-defined crystalline compounds, these decompose in aqueous solution. Compounds with bases are indefinite in character, and unstable Sodium and potassium dextrosate (6H11MO6 are for a little time, gas, water, and volatile acids obtained on adding sodium or potassium ethoxide are given off, and if the heating is continued to an alcoholic solution of dextrose. Several acetate, but basic lead salts are obtained in presence of ammonia.

The methylglucosides. These are of importance, as types both of glucosides generally, and of the two isomeric series of dextrose derivatives. They are regarded as stereoisomeric y-oxides, and have the following structural formula.—

 $\alpha\text{-Mothylglucoside}$  crystallises in long needles, in p. 165°  $[\alpha]_D + 157^\circ$ ,  $\beta\text{-methylglucoside}$  forms rectangular prisms, in p. 104°  $[\alpha]_D$  –33°

They are hydrolysed by warm mineral acids to methyl alcohol and dextrose, the \$\beta\$-isomeride being attacked 18 times as rapidly as the \$\alpha\$-isomeride. Each glucoside requires its own particular enzyme to hydrolyse it \$\alpha\$-Methylglucoside is hydrolysed by maltase alone, \$\beta\$-methylglucoside only by emulsin. They are used as test materials in studying enzyme action.

The isomerides are both formed when dextrose is heated with methyl alcohol containing

hydrogen chloride.

To obtain a-methylglucoside, 1 part of dextrose is heated with 4 parts methyl alcohol containing 0.25 p.c. hydrogen chloride, for 50 hours in an autoclave. The solution deposits crystals of the glucoside on evaporation, and the mother liquors, after further heating, give a second and third crystallisation, as the  $\beta$ -isomeride present is converted into the  $\alpha$ - by the acid. A yield of 80 p.c. or more is obtained

To obtain B-methylglucoside, solid anhydrous glucose is dissolved in methyl alcohol, nearly saturated with dry hydrogen chloride, at room temperature As soon as etherification is complete, as witnessed by the absence of reducing power, the solution is neutralised and evaporated. and a mixture of  $\alpha$ - and  $\beta$ -isomerides obtained The a- is partly in about equal quantities separated by crystallisation from strong alcohol The residue is dissolved in water, and fermented by a pure culture yeast The a-methylglucoside is hydrolysed, and the dextrose fermented, the  $\beta$ -methylglucoside remains untouched, and is recovered and punfied

In addition to the crystalline a- and  $\beta$ -methylglucosides obtained in the reaction between a-glucose and methyl alcohol containing 1 pc HCl, a considerable amount of syrup is formed which after distillation in a vacuum is an oil,  $C_7H_{14}O_6$ , bp  $200^\circ-215^\circ/0.2$  mm,  $[a]_D-4^\circ$ . It is named  $\gamma$ -methylglucoside to differentiate it from the a- and  $\beta$ -forms (E

Fischer, Ber. 1914, 47, 1980)

It is stable to alkalis, Fehling's solution or hot water, scarcely attacked by emulsin or by yeast extract at 37° C., but is hydrolysed to

glucose and methyl alcohol by dilute acetic acid at 100°, or by dilute hydrochlone acid at 100° or 18°

The other carbohydrates form analogous methylglucosides, and similar compounds are obtained with ethyl alcohol (For other artificial glucosides, v Glucosides)

The methyl derivatives of the sugars have proved of great importance in the clucidation of their structure. Purdie and Irvine's method of methylation is employed, consisting in dissolving the sugar in methyl iodide and a little acetone and boiling the mixture with silver oxide. By protecting some of the hydroxyl groups from methylation those remaining may be methylated and the configuration thus established.

A very large number of dextrose compounds have been prepared, in some of which the aldehydic hydroxyl is concerned, whilst others, eg the acetates, involve all five hydroxyls. As already stated, in most cases two isomerides derived from a- and  $\beta$ -dextrose have been prepared. Few of these compounds are really characteristic of dextrose, and it must suffice

in the space at disposal to indicate the properties of some of them in tabular form.

01 00120 01 1110111		•	
Compound	Appearance	Мр	[a]D
Pentanitrate	amoi phous		1 98°
a-Pentacetate .	colourless	112°-113°	$+100^{\circ}$
	ncedics		
<b>β-Pentacetate</b>	granular	134°	$+3^{\circ}$
p 2 circuloctato	crystals		
a-Acetochlorodex-	long needles	63°-64	
trose	tong meeting	00 0.	
B-Acetochlorodex-	tall to umn.	73°-71°	+165°
		10 -11	T 103
trose	gates	79°-80°	
	pusms	79*80*	
trose			1.4000
β-Acetobromodex-	lustrous needles	88°89°	198°
trose			
a-Acetonitrodex-	needles	92°	115°
tiose			
B-Acetonitrodex-	lustrous thombic	150°-151°	$+149^{\circ}$
trose	prisms		
Pentabenzoate .	needles .	$179^{\circ}$	
Monomethyl-a-		161°	+ 108°
dextrose		101	1 100
Monomethyl-β-		135	$+24^{\circ}$
		100	7 44
dextrose		85° -87°	+82°
αβ-Dimethyl-a-		09 -01	T 04
dextrose		1000 1100	1.40
ав-Dimethyl-в-		108°-110°	4 0°
dextrose			. =
Trimethyldex-	syrup		$+79^{\circ}$
troses	syrup .		8°
Tetramethyl-a-		81°	$+101^{\circ}$
dextrose			
Tetramethyl-β-		81°	$+73^{\circ}$
dextrose			
Pentamethyldex-	stellate crystals	42°-43°	−7°
trose			
Dextroscethyl-	colourless matter	1 127°-128°	
mercaptal	needles		
a-chloralose .	bunches of	186°	
a-cinoratoso .	needles	200	
B-chloralose	fatty plates .	230°	
Dextrose diacetone	noodles	107°-109°	
Dextrose ureide	rhombic plates	207°	
Dextrose dreide	and needles	201	
G 1 1		175°	
Semicarbazone	needles		
Thiosemicarbazone		204°	
Anilide .	needles .	147°	
Oxime	needles	137°	
a-Phenylhyda-	needles	159-160°	70°
zone			
β-Phenylhyda-	needles .	140-141°	5°
zone			
p-Bromophenyl-		164-166	
hydrazone			
Methylphenylhy-	colourless plates	130°	
drazone	District D Parton		
a-Amylphenylhy-	light brown	128°	
w Amjilandham.	TIBLE OLUMI		

needles

light yellow needles 155°

drazone

a-Allylphenylhydrazone

Compound	Appearance	Mр
a-Benzylphenylhy- drazone	light yellow needles	165°
Diphenylhydia- zone	colourless oblique prisms	161°
β-Naphthylhydia- zone	brown needles.	95°
	aggregates of vellow needles	206°
p-Bromophenyl- osazone	yellow needles	222°
Nitrophenylosa- zone	reddish needles	25 <b>7°</b>
Dextrose-o-diami- nobenzene	colourless lustrous needles	_
Didextrose-o-dia- minobenzene	slender colour- less needles	
Dextrose-γ-diami- nobenzoic acid	lustrous silver plates	$243^{\circ}$

Since acetobromoglucose is the starting-point for many other glucose derivatives its preparation is given in detail 200 grams powdered crystalline anhydrous glucose are heated on the water-bath with 100 grams of alcoholic sodium acetate and 1 litre of acetic anyldride for 2\{\} hours. The product is poured into 4 litres of ice water, and the crude \(\beta\)-pentacetylglucose is recrystallised from 96 p.c. alcohol after freeing it from acetic anhydride by water (yield 74 p.c.).

150 grams of the penta-acetate are treated with the commercial solution of hydrobromic acid in glacial acetic acid (300 grams) for 2 hours at ordinary temperatures, 600 c c chloroform added and the whole poured into 2 libres of ice water. The chloroform layer is separated, and the rest shaken with 150 c c chloroform; the united chloroform extracts are washed with 1 libre of water, dried over calcium chloride, concentrated in a vacuum, and the residue gradually treated with light petroleum. The resulting crystals are rapidly recrystallised from 75 c c amyl alcohol, washed with light petroleum and stored in a vacuum desiccator over soda-lime. (See E. Fischer, Ber. 1916, 49, 584.)

The reduction of accobromoglucose with zinc-dust and a ctu acid yields the acetyl derivative of glucal,

$$CH_2(OH)$$
  $CH$   $CH_2$   $CH(OH)$   $C=CH(OH)$ 

a soluble viscid syrup, b p 170°-180°/0 2 mm Glucal can be further reduced by hydrogen and palladium to hydroglucal,

which melts at 86°–87°, tastes slightly sweet, and has  $\lceil a \rceil_D + 16$ °.

d-Glucosumine (Chitosamine) or aminodextrose has the formula

It is obtained by boiling the shells of lobsters, particularly the claws, with concentrated hydrochloric acid, when the hydrochloride results. The shells of lobsters consist of carbonate of lime and chitin. Chitin is a constituent of various invertebrate skeletal structures, and of the cell walls of fungi; it is considered by Irvine to contain acetylamino-dextrose and amino-dextrose residues in the proportion of 3 to 1  $(C_{30}H_{50}O_{19}N_4)n$ . It is prepared from the hydrochloride by decomposing if with Vol., II—T.

diethylamine (Breuer, Ber 1898, 31, 2193) or sodium methoxide (Lobiy de Bruyn, did 1898, 31, 2476) Fischer and Leuchs (Ber 1903, 36, 24) effected the synthesis by the combination of d-arabinose and ammonium eyanide to d-glucosamine acid, the lactone of which was reduced to d-glucosamine. Glucosamine forms a colourless crystalline powder, or long needles; mp. 115°  $[a]_{\rm p}+48^{\circ}$  The aqueous solution reacts alkaline, but is stable. It is not fermentable. The pentacetate exists in two forms, needles, mp. 183.5° and 133° respectively; the former is optically inactive, the latter has  $[a]_{\rm p}+86.5^{\circ}$ . The oxime forms prisms, mp. 127°. Two modifications of the hydrochloride are known glistening monoclinic crystals  $[a]_{\rm p}^{23}+100^{\circ}$ , falling to  $+72.5^{\circ}$  on standing, and hexagonal needles  $[a]_{\rm p}^{23}$  constant at 72.5°.

It cannot be converted into dextrose directly, though it gives dextrose phenylosazone when heated with phenylhydrazine. Irvine and Hynd (Chem. Soc. Trans. 1912, 101, 1128) have converted it into aminomethyl glucoside, and this by the action of baryta into methylglucoside and ultimately to d-glucose. When methylglucosamine is treated with sodium nitrite a mannose derivative is obtained almost quantitatively. Probably a Walden inversion takes place (see Chem. Soc. Trans. 1914, 105, 698). By the action of nitrous acid, a compound, C<sub>6</sub>H<sub>10</sub>O<sub>5</sub>, is formed, which was at one time regarded as a carbohydrate, and termed chitose, but was shown by Fischer and Andreae (Ber. 36, 2587) to be a hydrated furfurane derivative

Chitose forms an oxime and phenylhydrazone and is oxidised by bromine to chitome acid (flucosamine was formerly regarded as a derivative of chitose, and termed chitosamine

An isomeride of glucosamine is reglucosamine CH<sub>2</sub>(OH) [CH (OH)]<sub>3</sub> CO CH<sub>2</sub> NH<sub>3</sub>, which Fischer obtained by reducing phenylglucosazone with zinc-dust, and which is converted by the action of introis acid into Levulose

The glucosumines were formerly believed to be immo compounds, but Irvine, Thomson and Garrett (Chem Soc Trans 1913, 103, 238) suggest that they are amino-glucosides. Levene (J Biol. Chem 1916, 24, 59) shows that the nitrogen content of glucosumine, galactosimine, xylosimine, and lyxosimine as given by (i) Kjeldahl's method, and (ii) by van Slyke's nitrous acid method indicates the presence of a primary amino group.

Chondrosamine (Levene, J. Biol Chem. 1916, 26, 143) gives galactosazone, and is therefore probably a lyxohexosamine

By condensing bromotriacetylglucosamine with hydroxy compounds in the presence of a base such as morphine, acctylated amino glucosides are obtained. The compounds formed are of two types: stable N-cyclic compounds and normal glucosides, such as a-aminohelicin and a-aminosalicin (Irvine and Hynd, Chem. Soc. Trans 1913, 103, 41).

A new glucosamine derivative, lycoperdin  $C_{13}H_{24}O_{9}N_{2}$ , has been isolated from the fungus

Lycoperdum gemmatum (Kotake and Sera,

Zeitsch physiol Chem 1913, 88, 56)

Manufacture of commercial dextrose-se. destro-glucose, glucose, saccharum, saccharine, &c -These products, varying much in character and composition, are all prepared by the action of acids on starch or starchy substances, such as rice, maize, &c., and when these latter are the raw material, it has been proposed to employ the combined action of diastase and acids When starch is acted upon by dilute acids, it is dissolved, dextrins and maltose being produced These bodies, by the further action of the acid, yield dextrose; this itself does not resist the action of acids, so that during the process of conversion, a portion of it is attacked, with the production of substances imperfectly studied (v ISOMALTOSE)

The phases of the action of dilute acids on starch are easily followed. After heating has been continued for a short time, a portion of the solution cooled ceases to give a blue colouration with iodine, this indicates absence of starch, in the place of this a reddish-brown colouration is produced, this indicates the presence of a-dextrin After a time this disappears, and at length the solution ceases to give a precipitate, when to a portion of it twice its bulk of alcohol (sp.gr 0 \$20) is added, or when to a portion from which the sulphuric acid has been separated by baryta-This estabammoniacal lead acetate is added lishes the absence, not of maltose, but only of For syrup the boiling is stopped as soon as the jodine test gives a port-wine colour; for solid dextrose it is continued till alcohol The processes involve causes no precipitate in the manufacture may be conveniently arranged as follows -

- (a) The conversion
- (b) The neutralisation
- (c) The filtration(d) The decolourisation.
- ( ) The concentration
- (f) The purification.
- (a) The conversion—In this country the materials chiefly employed in the manufacture of commercial dextrose are sago, maize, and rice starch, finely ground rice itself, as well as granulated maize and rice being at times used Potato starch is also used in Germany In America green-maize starch is the chief material, hence the term 'corn sugar' The acid usually employed in Europe is sulphuric; hydrochloric acid is generally used in America, and nitric acid as well as a mixture of nitric and sulphuric acids have been recommended The use of sulphurous and orthophosphoric acids has also been The proportions of material, water, advocated and acid generally employed in the case of sulphuric acid are 100 · 250 · 5, and when green starch, ie. undried purified starch, is used, 14 parts are taken instead of 1 part dry starch. When pressure converters are used, the acid can be reduced to 0.5 p c.

Formerly open converters made of wood, lined with lead and provided with steam coils of lead and stirring gear, were used, but these are now abandoned nearly everywhere for closed These are made of cast iron, copper, or gun metal, and are worked at a prossure of 30 lbs of steam.

The operations involved in the manufacture. whatever converter is used, are as follows: Half the sulphuric acid is mixed with half the water in a separate vat constructed as the converter, with or without coil, and at such a temperature that the heat of the dilute acid will be less than the gelatinising point of the starch of the material employed. This vat is furnished with stirring gear, and placed under a hopper, in which is stored the starch or starchy substance When the dilute acid is at the right temperature, the material is allowed to flow into it, the stirring gear being kept in motion; a milk of starch in dilute acid is thus obtained Meanwhile the remaining half of the acid and water are introduced into the converter, and heated to boiling, then the starch milk is run in at such a rate that the temperature does not fall far below boiling, and the introduction should be effected as rapidly as possible

The time required for conversion depends on the rate of introduction of the starch milk, and on the pressure employed, &c. For corn syrup an average time from the commencement of the starch introduction to the discharge is a little over an hour, for solid sugar about 14 hours is required When converted, the liquid

is blown into the neutralising vats

(b) The neutralisation -Finely-ground chalk is used to neutralise the acid employed in the conversion The chalk is converted into a milk with water, and gradually introduced into the converted liquid About 12 lbs of powdered chalk are taken for every 10 lbs sulphuric acid employed: this is more than the theoretical quantity, but it is found to answer in practice. owing, no doubt, to the insolubility of a portion The neutralised liquid is kept of the chalk stirred for some time, and is then allowed to rest It is found best to allow it to cool and settle gradually, and then draw off the clear portion, but where vat room is a consideration, the neutralised liquid is run immediately to the filter presses (r FILTRATION)

(c) (d) The clear filtered solution is passed through bone-char filters of the same construction as those described under Sugar acid is sometimes used in bleaching the product.

(e) Concentration -This operation is effected in vacuum pans (1 Sugar), and is considered complete when the syrup contains about 82 p c solid matter, ie reaches a sp gr of 150 then run into moulds, usually cone-shaped vessels, in which, on standing and cooling, it solidifies after a short time. This is hastened by the introduction of some previously solidified The solidified mass extracted from the mould by a short immersion in hot water is broken into irregular lumps, in which form it appears in the market. The colour of the commercial parcels varies from pure white to darkbrown, this depends on the character of the material employed (the purity and variety of the starch), on the mode of conversion, and the extent to which the decolourisation was carried The composition is also very variable, but the constituents in all are the same, viz. dextrose, maltose, dextrin, unfermentable organic bodies, and ash Very few samples contain dextrin, some few maltose, and the proportion of unfermentable bodies varies very much, being sometimes as high as 15 p.c. The syrups,

which are used in confectionery, contain a large percentage of dextrin and maltose

In the United States the conversion is carried out by stirring the green corn starch, made in the same factory, with a large body of water, and subjecting this mixture to a high heat under pressure, the process being a hydrolytic one. A small amount of pure hydrochloric acid is added to facilitate the catalytic action. The conversion is performed in large vessels 6 feet in diameter, and about 20 feet high, made of hammered copper 1 inch in thickness.

After the required conversion, the traces of acid are neutralised with soda ash, the mixture being left very slightly acid to litmus, and excess of alkali being very carefully avoided, as the least trace colours the liquid. In the case of glucoso syrup, conversion is complete in less than 10 minutes; conversion into anhydrous dextrose takes considerably longer The converted starch, now an almost white liquor, is run from the neutralisers through filter presses, to remove the dissolved gluten, &c, which is precipitated during neutralising, and then through an evaporator, which reduces the gravity to 30° B It is then run successively over three bone-char filters, each 10 feet in diameter and 24 feet high, and containing 70,000 lbs of bone-char each The liquors receive a much greater amount of bone-char treatment than is the custom in sugar refineries. passing, in all, over 72 feet. The remaining traces of acid are completely neutralised, and the liquid has a brilliant crystal-white appear-The bone-char is revivited after each The liquor is concentrated in vacuum pans to the required density 42°-45° B, and crystallises on cooling (A full account of the American industry of corn products is given by Wagner (J. Soc. Chem. Ind `1909, 28, 343) )

The average composition of these products, according to the older analyses, is as follows —

	Corn syrup	70 sugai	89 sugar	An- hydrous sugar
Water . Dextrose . Dextrin . Ash	P 6	P c	P c	P 6
	19 0	19 7	11 2 1	4 0
	38 5	70 2	79 91	94 6
	42 0	9 3	18 0	0 7
	0 5	0 8	0 9	0 7

The ash consists principally of chlorides Wesener and Teller (J Ind Eng Chem 1916, 8, 1009) give the following analysis as representing the modern product—

and Dextria (or dextrins) not fermented by yeast, but made fermentable by pancreatin, takadiastase, &c.

From which it appears that the proportion of maltose present generally exceeds that of dextrose. In the older analyses all the reducing sugar was regarded as dextrose

(f) Purification.—Behr 'Eng. Pat. 9797,

1884; 1767, 1885) manufactures pure commercial crystalline anhydrous dextrose by adding crystals of dextrose to a solution concentrated to 85 p.c., and allowing the mixture to stand at about 35°. After a short time an abundant crop of crystals separates, and by treatment in a centrifugal machine, the uncrystallised syrup is removed (v. Sugar)

In the older process of Soxhlot, the concentrated syrup was dissolved in about an equal weight of boiling methyl alcohol, and allowed to stand. The crystals were drained, and the loaves washed with methyl alcohol, which was recovered from the loaves and mother liquous by distillation in a vacuum.

Dextrose from Cellulose —Cellulose is disintegrated by boiling with acids only with great difficulty, and at high temperatures, with the formation of dextrose, and the process cannot compete industrially with the preparation from starch. Numerous attempts have been made to obtain dextrose from wood cellulose, with the industrial objective of the manufacture of alcohol.

Simonsen (Zeitsch angew Chem 1898, 195, 219, 962, 1007) found the most favourable conditions for the conversion of 40 grams of cellulose to be a digestion for 2 hours at 6–8 atmospheres, with 1000 c c. of 0.5 p.c. sulphung acid 45 p.c. of reducing sugar was obtained, but, according to Korner (ibid 1908, 2353), this is not all termentable dextrose.

Ekstrom (Fr. Pat. 380358, July 29, 1907) treats sawdust with 95 p c sulphune acid at the ordinary temperature for 20 minutes, dilutes to 1 p c free acid, and heats from 1 to 5 hours under a piessure of 3–8 atmospheres , 55–57 p c, conversion is claimed

Ewen and Tomlinson (US Pat. 938308, Oct. 26, 1909) heat sawdust with a minimum of 5 p e acid to a temperature between 135° and 162°. The use of 2–6 p c. hydrofluorie acid has also been suggested (Orlowski, Fr. Pat. 405187, July 17, 1909)  $d\text{-Mannose} \ C_6 H_{12} O_6 \ \text{occurs in nature in the}$ 

form of anhydride-like condensation products, known as mannosans It is best prepared from the vegetable ivory nut (sternnuss) which is used in Beilin and elsewhere for the manufacture of buttons and collar studs. The waste turnings are hydrolysed to mannose by boiling for 6 hours with 2 parts of hydrochloric acid It was originally obtained by Fischer and Hirschberger by oxidising mannitol with nitric acid. Mannose is characterised by forming an insoluble yellow phenylhydrazone, mp. 186°-188°, and use is made of this compound in isolating it from solution The hydrazone is subsequently decomposed with benzaldehyde, or better, formaldehyde, and the resulting solution concentrated in vacuô till the sugar crystallises It may even be isolated in the solid form without the intermediate formation of the phenyl hydrazone (Hudson and Sawyer, J Amer Chem Soc. 1917, 39, 470)

Mannose separates in colourless rhombic crystals  $[a:b:c=0\ 319:1\cdot 0\ 826]$  of sweet taste, mp. 132°; spgr. 1·539 100 cc of the saturated aqueous solution at 17·5° contain 24 8 grams, in absolute alcohol it is almost insoluble. It shows muta-rotation, the equilibrium value being  $[a]+14\cdot5°$ . The a-form has

 $[a]_{\rm p}+34.0^{\circ}$ , and the  $\beta$ -form  $[a]_{\rm p}-17.0^{\circ}$  (Hudson | (III) m p. 98°,  $[a]_{\rm p}-42^{\circ}$  (in chloroform); and Dale, J. Amer. Chem. Soc. 1917, 39, 320)

It is very similar to dextrose in all its properties, yielding the same phenylosazone fermented by all yeasts which attack dextrose and levulose Bromine oxidises it to mannonic acid, sodium amalgam reduces it to mannitol. a-Methyl mannoside had m p 190°,  $[\alpha]_{\rm p}^{20} + 80^{\circ}$ The oxime crystallises in colourless needles, mp 167°-180° Like the phenylhydrazone, the substituted hydrazones are all sparingly soluble

Caroubmose, a carbohydrate isolated from the seeds of the carob bean, is identical with

mannose.

d-Galactose  $C_2H_{12}O_6$  Milk sugar, when hydrolysed by means of acids or enzymes, yields dextrose and galactose in equal proportions On a commercial scale, sulphuric acid is used to effect hydrolysis under pressure at 105°, after which the solution is neutralised and concentrated until an amount which contained 500 grams of milk sugar weighs 640 grams syrup is set aside with a few crystals of galactose, and, after some days, the crystalline mass is washed with 80 p.c. alcohol, and the galactose purified by recrystallisation from aqueous alcohol Even the very best commercial galactose contains minute quantities of dextrose; to remove this it must be treated with a pure culture of some yeast, such as S Ludwign, which ferments dextrose, but not galactose

Galactose is a constituent of raffinose and stachyose, and is widely distributed in the polymene form, galactan, particularly in gums and seaweeds It occurs in a few glucosides It is a colourless crystalline powder,  $+1\rm{H}_2\rm{O}$ , m p  $118^\circ$ - $120^\circ$  (from water), anhydrous m p  $165^\circ$ (from alcohol), less soluble than dextrose, and has the same general properties as dextrose The most characteristic behaviour is the formation of mucic acid on oxidation with nitue acid.

Galactose exhibits muta-rotation; the aisomeride has  $[\mathbf{a}]_{\mathrm{D}}^{20} + 144^{\circ}$ , the  $\beta$ -modification +52°. The equilibrium value is  $[a]_{D}^{20}$  +80 5°. The influence of concentration and temperature on this value is expressed by the equation

 $[\alpha]_{\rm D}\!=\!83\!\cdot\!883\!+\!0.0785p\!-\!0.209t$ 

where p and t denote concentration and temperature respectively

Galactose derivatives are prepared similarly to the corresponding dextrose compounds The a-pentanitrate (Will and Lenze, Bei. 31, 68) forms needles, mp 115°-116°; the  $\beta$ -isomeride has mp. 72°-73°; the  $\beta$ -pentacetate has mp 142°, amethylgalactoside has mp  $[a]_{D}^{110}-112^{\circ}$ ,  $[a]_{D}^{20}+179.3^{\circ}$ , and crystallises in transparent doubly refractive needles; the  $\beta\text{-isomeride}$  forms needles, m p  $173^\circ\text{--}176^\circ\text{,}$  $[a]_{D}^{20}+2.6^{\circ}$ ; the phenylhydrazono has mp 158°, the phenylosazone when quickly heated melts at 193° -194°.

Galactose penta-acetate has been isolated in

(1) m p.  $142^{\circ}$ ,  $[a]_{p} + 25^{\circ}$  (in chloroform),  $+8^{\circ}$  (in

benzene); (ji) m p. 95.5°,  $[\alpha]_D + 107^\circ$  (in chloroform);

(iv) m.p.  $87^{\circ}$ ,  $[a]_{D} + 61^{\circ}$ .

(Hudson and Parker, J Amer Chem. Soc 1915, 37, 1589, 1591; Hudson and Johnson, *ibid*. 1916, 38, 1223).

Some yeasts ferment galactose, but not all those which ferment dextrose and Levulose is apparently possible to grow a yeast in presence of galactose and dextrose, until it acquires the power of fermenting galactose. Galactose is identical with the sugar cerebose, isolated from the brain by Thudichum, where it is a constituent of the cereborsides, phrenosin and kerasin

Lævulose or d-Fructose ('<sub>6</sub>H<sub>12</sub>O<sub>6</sub> is a keto-hexose or ketose sugar. It is found together with dextrose in honey and in many fruits, and, combined with dextrose, it occurs in sucrose, raffinose, stachyose, &c Inulin, found in dahlia tubers is completely converted into fructose on hydrolysis. It has not so far been produced by the hydrolysis of any glucoside

It is termed d-fructose, in spite of the lavorotation, to indicate that it belongs to the d-mannitol series

After prolonged heating at 165°-170°, cane sugar becomes converted into a mixture of levulosan (C<sub>6</sub>II<sub>10</sub>O<sub>5</sub>) and dextrose, without change of weight. On fermenting the aqueous solution of the mass with yeast, the dextrose is decomposed, while the levulosan remains unchanged, and may afterwards be converted into Levulose by treatment with dilute acids

Preparation —It is best prepared from mulin 100 grams of mulin with I pc ash are heated with 250 c.c. water and 0.5 gram hydrochloric acid for half an hour in a boiling water-bath, neutralised with 15 grams sodium carbonate evaporated on the water-bath at 60°, and finally over sulphurie acid in vacuô to a thick syrup This is extracted with absolute alcohol, put aside for 24 hours, the solution poured off clear, and inoculated. Complete crystallisation of the lievulose is obtained in 3 days, and it is quite pure after one more crystallisation

Inulin may also be hydrolysed with 0.5

vol p c. oxalic acid.

(For the preparation from Jerusalem artichokes, see U.S. Pat 789422, Feb. 18, 1908.)

To obtain lævulose from sucrose, the latter is first inverted by means of mineral acids or invertase, and the acid removed by neutralisation with baryta or silver oxide. The invert sugar resulting contains both dextrose and lavulose, which may be separated as follows · 6 grams of finely powdered slaked lime is added to every 100 cc. of a neutral ree-cold 10 pc solution of invert sugar, and the solution, which at first sight is merely milky, is agitated, the vessel being preferably immersed in ice-cold water until it acquires a creamy consistency. The mass, which consists of the difficultly soluble calcium levulosate CaO C<sub>6</sub>H<sub>12</sub>O<sub>6</sub> H<sub>2</sub>O, tegether with the easily soluble calcium dextrosate, is filtered through linen, and strongly pressed to remove the greater part of the liquid, and is suspended in water and decomposed by addition of oxalic acid or by passage of carbonic acid gas. The filtered solution is evaporated in vacuo over sulphuric acid, with production of anhydrous lavulose. Calcium lavulosate dissolves in 1.27

parts of water at 15°, and is almost insoluble at  $0^{\circ}$ .

Levulose, when pure, crystallises from alcohol, and even from aqueous solution, in needles, also in crusts of transparent prisms, which are not hygroscopic It has mp 95°, D<sup>17</sup>5 1 669 It sublimes unchanged but very slowly at 100°/1-2 mm (Ghosh, J. C. S. Abstr. 1917, 112, (1) 80). It is very soluble in water, almost insoluble in cold absolute alcohol. It is much sweeter than dextrose, and somewhat sweeter than sucrose It exhibits muta-rotation, and, no doubt, exists in solution as an equilibrated mixture of stereoisomeric y-oxide forms, The initial rotation  $[\alpha]_{\rm p}^{20} = -104^{\circ}$  falls to  $\left|\alpha\right|_{D}^{20}$  -92° (Parkus and Tollens, Annalen, 257. The specific rotatory power is very considerably influenced by changes of temperature and concentration, which cause an alteration in the proportions of the isomerides in the equili-The rotatory power usually brium mixture taken is  $\left[\alpha\right]_{D}^{15}$  =93.8° It falls 0.6385° for each degree centigrade of increase in temperature, and at 873° is equal but opposite to that of dextrose. This property is taken advantage of m determining its proportion in presence of other sugars, the rotation being observed at two different temperatures

According to Soxhlet, the relative reducing powers of levulose and dextrose on Fehling's solution, in 1 p.c. solution, are 475–513, but Allihn finds that if the solutions be boiled for 30 minutes (instead of for 2 minutes, as in Soxhlet's experiments), their reducing powers are identical. On Knapp's mercurial solution their action is the same, but on Sachsse's mercurial solution their action are 100 (levulose)

and 148 6 respectively

Oxidation converts levulose into acids with fewer carbon atoms, glycollic and trihydroxybutyric. On reduction with sodium amalgam, mainited and sorbited are formed. Acids attack it far more readily than dextrose with the production of levulic and formic acids, alkalis somewhat less readily. Heating concentrated solutions with oxalic acid forms 1-hydroxymethylfurfurol, whilst it interacts with hydrogen bromide in othercal solution, forming 4-bromomethylfur

CH . C(CH<sub>2</sub>Br) O (Fenton and Gostling, CH : C(CHO)

Chem Soc Trans 1899, 75, 423, 1901, 79, 301)
It is fermented by all yeasts which attack destrose.

By the action of methyl alcohol and hydrogen chloride (0.5 pc) at  $35^\circ$ , a syrup is obtained which probably represents a mixture of methyl fructosides. It is partially hydrolysed by yeast extract

Fructose diacetone  $C_{12}H_{20}O_{6}$ , prepared in a similar manner from acetone, forms needles, mp 119°-120°  $[\alpha]_{\rm D}$ -161°. An isomeride has mp 97°  $[\alpha]_{\rm D}$ -33 7°.

The osazones of lavulose are identical with those of dextrose. Lavulose is more easily oxidised by disubstituted phenylhydrazines, and forms a characteristic osazone with methylphenylhydrazine, long yellow needles, mp. 158°-160°.

To prepare this, 4 grams of methylphenylhydrazine are added to 1 8 grams levulose, in 10 c.c. water, and enough alcohol to make a clear solution further 2 c.c. of acetic acid are added, and the mixture heated 5-10 minutes at most on the water-bath. It is then covered and allowed to stand. Within 2 hours red crystals are obtained, which may be recrystallised from 10 p.c. alcohol.

Fructose penta-acetate melts, when pure, at  $108^\circ-100^\circ$ , has  $\lfloor \alpha \rfloor_D = 121^\circ$ , and crystallises in billiant colourless needles, 1 cm. in length (Hudson and Brauns, J Amer Chem. Soc

1915, 37, 1283)

Sorbose (Sorbinose)  $C_6H_{12}O_6$  is the ketohexose obtained from the juice of the mountainash berries, which has been exposed to the air and allowed to ferment. The juice contains sorbitol, this is oxidised by bacterium xylinum to sorbose

The juices are evaporated to 1.05 sp gr, allowed to stand some days till the sugar is fermented, and the clear liquid poured into flat dishes and moculated with an active culture of B-xylinum (Bertrand, Ann Chim Phys 1904, [viii ] 3, 181). This is left at 30° till the reduction is a maximum, clarified with lead acetate, excess of lead precipitated as sulphate, and the neutral filtrate concentrated in view When pure sorbitol is available, 100 grams are dissolved in 500 c.c. water, some nutrient solution for the bacterium added, and the fermentation carned out as above

Sorbose forms thombic crystals of a sweet taste, in p 151°, it reduces Fehling's solution, and generally behaves as Levulose, but yields sorbitol on reduction. It is not fermentable by yeasts. It is Levo-rotatory, having  $[\alpha]_D - 42^\circ$ 

Sorbose phenylosazone is similar to the dextrose compound, but has m p 164°.

The properties are in agreement with the structural formula

### HO H OH CH<sub>2</sub>OH CO C · C · C · CH<sub>2</sub>OH H OH H

Methyl sorbinoside crystallises in transparent thick plates, in p 120°-122° (Fischer, Ber. 28, 1159)

#### HEPTOSES.

d-Mannoketoheptose Occurs in the Avocado pear (Persat gratissima) (La Forge, J Biol. Chem 1917, 28, 511) It has in p 152°, [a]<sub>p</sub>+29°. It is not fermented by yeast, and does not show mutarotation. It gives a colour reaction with oremol and hydrochlone acid like other heptoses. The p-bromophenylhydrazone melts at 179°, the phenilosazone is identical with that from d-mannoaldoheptose.

On reduction the sugar yields d-persital and

d-mannoheptitol

Since the p-bromophenylhydrazone of d- $\alpha$ -mannoaldoheptose melts at  $207^{\circ}$ - $208^{\circ}$ , and that of d- $\beta$ -mannoaldoheptose is an oil, the configuration of the sugar must be :—

 $\begin{array}{cccc} \textbf{Perseulose} \ (l.c.) \ \text{is shown to be } l\text{-}galaheptose: --- \\ & \text{H} & \text{OH} & \text{OH} & \text{H} \\ & \text{CH}_2(\text{OH}) \ \text{CO} \ \text{C} & \text{C} & \text{C} & \text{C} & \text{CH}_2(\text{OH}) \\ & \text{OH} & \text{H} & \text{H} & \text{OH} \\ \end{array}$ 

Sedoheptose Occurs in Sedum spectabile (La Forge and Hudson, J. Biol Chem 1917, 30, 61) as a non-fermentable, reducing syrup

Phenylosazone, mp 197°; p-bromophenyl-hydrazone, mp 227°-228°; the osone is a syrup, its o-phenylenediamine derivative melts at 163°-165°

Bromine water does not oxidise the sugar, but it reduces with sodium amalgam to two alcohols  $C_7H_{18}O_7$ —

a-Sedoheptitol, m p  $151^{\circ}-152^{\circ}$ ,  $[\alpha]_{D}+225^{\circ}$ .

 $\beta\text{-Sedoheptitol}, \quad \text{m p} \quad 127^\circ\text{--}128^\circ, \quad \text{optically mactive}$ 

The two alcohols are separable by their benzylidene derivatives

These reactions indicate that the sugar is a ketose

Anhydrosedoheptose  $C_7H_{12}O_6$  is formed when sedoheptose is heated with dilute acids. It forms long white prisms, melting at 155°,  $\lfloor a \rfloor_D - 146$ °. It shows no mutatotation, does not reduce Fehling's solution, and, when boiled with dilute acid, gives an equilibrium mixture which contains 20 p.c. of sedoheptose

These are widely distributed in plants, where they are present in the skeletal structure, and do not, like the hexoses, serve as food products Pentoses also appear to be a constituent of many of the tissues of the animal body, and are sometimes found in urine. Both the natural pentoses, *l*-arabinose and *l*-xylose, belong to the lavo- senes Dextrose (dextro-glucose), when degraded by the methods of Ruff or Wohl, gives rise to the isomeric d-arabinose. However, the natural pentoses are closely related to the natural hexoses For example, the space arrangement of the groups attached to the upper four carbon atoms is the same in d-glucose as it is in l-xylose

A similar relationship exists between galactose and arabinose, which occur together in many gums, whilst some polysaccharides yield both xylose and glucose on hydrolysis. The pentoses do not occur as such, but in the form of condensation products of high molecular weight, termed 'pentosans' (araban or xylan). These are comparable with the hexose condensation products, starch and cellulose, as a rule, the pentosans contain both C<sub>5</sub> and C<sub>6</sub> carbohydrates.

The mechanism of the formation of pentoses

in the plant is not yet clear. Some authorities consider they are derived by oxidation from the hexoses, others advocate the view that they are duectly formed in the cell, like the hexoses, but independently of these Such a transformation as that from dextrose to l-xylose has not been effected in the laboratory the chemical degradation of dextrose by oxidation leads to d-arabinose. The pentoses show all the general chemical properties of the hexoses Biochemically. they are different, being not fermentable by yeasts, whilst their glucosidic derivatives (e q. the methyl arabinosides and xylosides) are not hydrolysed by any of the better-known plant enzymes. They are attacked by a number of bacteria Characteristic of the pentoses is the formation of furfurol on distillation with hydrochloric acid They, therefore, show colour reactions with original and phloroglucinol. Equal parts of concentrated hydrochloric acid and water are taken, together with the pentose and a little oremol or phloroglucinol, and the mixture warmed in the water-bath. With oromol the solution becomes red, then violet, and finally blue or blue-green, with the separation of a precipitate which dissolves in amyl alcohol, giving a blue-green solution, this shows an absorption band between C and D. With phloroglucinol the solution becomes cherry-red, and a precipitate is formed, giving a red solution in amyl alcohol, with an absorption band between D and E

l-Arabinose  $\mathrm{C_5ll_{10}O_5}$  is obtained by hydrolysis of most gums, particularly gum arabic and cherry gum

Preparation -A solution of gum arabic, or allied gum (dextro-rotatory, and yielding little mucie acid by the action of nitric acid, by preference), containing 30-40 grams of the gum, and 2-3 c c sulphune acid in 100 c c, is boiled for 7-12 minutes, cooled, the acid neutralised with calcium or battum hydroxide or carbonate and alcohol, sp gr 0 83 added as long as a precipitate is produced. This precipitate soon settles, leaving the alcoholic solution clear. On distilling off the alcohol and evaporating to a syrup, arabinose crystallises out. The substance is obtained in well-defined thombic prisms, with monoclinic termination, by recrystallising from water. With some gums the digestion of the acid solution can be continued for 2 hours with increased yield of arabinose, and without impairing the purity of the product. The crystals are anhydrous, mp. 160°

Arabinose has  $[a]_D+104^\circ$  to  $+105^\circ$ , the activity increasing with the strength of the solution, and decreasing as the temperature at which it is observed is raised

It exhibits muta-rotation,  $\alpha$ -arabinose having the initial rotatory power  $[\alpha]_D + 175^\circ$ , decreasing to  $+104^\circ$   $\beta$ -Arabinose has not been isolated. Accordingly, like dextrose, it has a  $\gamma$ -oxidic structure.

The heat of combustion is 558 cals

Arabinose reduces Fehling's solution, the amounts of copper oxide produced by equivalent weights of arabinose and dextrose being as 111:100

Bromine oxidises it to l-arabonic acid  $C_5H_{10}O_6$ , which crystallises as the lactone  $C_5H_8O_5$ , m p. 95°-98°,  $[a]_D$ -73 9°. Nitric acid

forms oxalic acid as the final product, but in dilute solution, firstly arabonic acid, and secondly *l*-trihydroxyglutaric acid, in p 127°, are formed. On reduction, amongst other products, *l*-arabitol, in p 102°, is formed, which is optically inactive, but becomes law-rotatory in presence of borax

Arabinose forms an additive compound with hydrogen cyanide, which is hydrolysed to arabinose carboxylic acid. This is, in reality, a mixture of two acids, of which the lactones on reduction give rise to l-glucose and l-mannose.

The diphenylhydrazone, mp 218°, is almost insoluble The other hydrazones have the following melting-points: phenyl, 151°; p-biomophenyl, 150°, α-methylphenyl, 161°, α-othylphenyl, 153°; α-benzoylphenyl, 170°, β-naphthyl, 141° Arabinose phenylosazone, mp 160°, is prepared similarly to the dextrose compound. It is slightly more soluble, and, when boiled with benzaldehyde, arabinosone is formed. The p-biomophenylosazone has mp 196°-200°.

Arabinose forms crystallisable compounds with the aromatic diamines and diamino-acids

l-Xylose is obtained by the hydrolysis of wood gum (xylan) extracted from wood by dilute soda solution

According to Bertrand (Bull Soc chim [3] 5, 554), it may be obtained from wheat or oat straw by first extracting the straw with tepid water, and then boiling it for several hours with dilute (1–2 p c) sulphuric acid. The liquoi, after removal of the sulphuric acid by baryta, is concentrated on the water-bath and treated with alcohol; this extracts the xylose, which forms a syrup on evaporation of the alcohol, solidifying on the addition of a crystal of xylose. The yield is about 2 p c from wheat straw, and 4 p c from oat straw

Another convenient source are oat hulls, which are cleaned by soaking in dilute ammonia, washed and hydrolysed with sulphume acid Hudson and Harding (J Amer Chem Soc 1917, 39, 1038) obtained a yield of 8-12 p c of xylose by direct acid hydrolysis of cottonseed hulls A more suitable source is maize cobs (*idem* 1918, 40, 1601) Xylose crystallises in orthorhombic prisms, in p. 144°–145° or 150°–154° It exhibits muta-rotation, a xylose having  $[a]_p$ +79°, falling to +19°, which is the rotation of the equilibrated mixture The heat of combustion is 561 cals Bromine oxidises it to xylonic acid, a syrup which is at first levo-rotatory, but reaches  $[\alpha]_D + 17.5^\circ$ , as it is transformed into the lactone in solution It forms a very characteristic insoluble compound C5H9O6CdBr, H2O with cadmium bromide Nitrie acid in the cold forms an optically mactive trihydroxyglutaric acid

On reduction, mactive xylitol, a syrup, is formed

a-Methylxyloside has mp 90°-92°,  $[a]_{\rm D}+152$ , the  $\beta$ -isomeride has mp 156°-157°,  $[a]_{\rm D}-65$ 9° With hydrogen cyanide two xylose carboxylic acids are formed, which yield l-gulose and l-idose on reduction.

The phenylhydrazone has m p.  $116^{\circ}$ ; the  $\beta$ -naphthylhydrazone, m p  $70^{\circ}$ , and the phenylosazone, m p  $166^{\circ}$ 

The methyl pentoses are pentose carbohydrates, in which one of the hydrogen atoms in

the primary alcohol group is replaced by methyl They appear to be widely distributed, but their brochemical significance is not yet understood. They show the same reactions as the pentoses, and, like these, are not ferimented by yeasts. Characteristic is the formation of methylfurfuraldehyde on distillation with acids

Rhamnose (Isodulettol) C<sub>6</sub>H<sub>12</sub>O<sub>5</sub>,H<sub>2</sub>O, on account of its empirical formula, and the ease with which it crystallised, was formerly classed with dulettol and mannitol, hence the original name of isodulettol. Its chemical behaviour shows it to be a methyl pentose. It appears to be widely distributed in nature in the form of glucosides, the best known of which are xanthorhamnin, the colouring matter of Persian berries and querettiin. Other glucosides containing rhamnose are baptism, fisetin, frangulin, fustin, datiscin, glycyphyllin, hesperidin, natingin, tutin, sophorm (v. Glucosides).

It is prepared by extracting Persian berries with hot alcohol, distilling off the alcohol, and heating the xanthorhammin on the water-bath for 2 hours, with 6 pc sulphune acid. The precipitated rhamnetin is filtered, the filtrate neutralised with barium carbonate, and, after removal of the barium sulphate, evaporated to a syrup. This is diluted hot with absolute alcohol, and the solution allowed to stand, when it deposits hard lustrous crystals of rhamnose About 15–20 grams are obtained per kilo of berries.

It is advantageously prepared from the residues of the manufacture of quereitron extract, when these are available

Rhamnose has the constitutional formula

Until recently the positions of the groups attached to the carbon atom next to the  $\mathrm{CH_{1}}$ -group were uncertain, but Fischer and Zach (Ber 1912, 45, 3761) have obtained d-isothamnose by a series of reactions from triacetyl dibromoglucose, in which compound the two halogen atoms are attached to the terminal carbon atoms of the sugar chain.

The hydrate softens at 89°; m p 92°-93° When anhydrous, it crystallises from acctone in needles, m p 122°-126° (Fischer), 108° (Tarret) Like dextrose, it exists in  $\alpha$ - and  $\beta$ - modifications, which are in equilibrium in solution. The  $\alpha$ -torm has  $\lfloor \alpha \rfloor_{\rm p} -17^{\circ}$ ; the  $\beta$ - form  $\lfloor \alpha \rfloor_{\rm p} +31.5^{\circ}$ , and the equilibrated mixture has  $\lfloor \alpha \rfloor_{\rm p} +9^{\circ}$ . It sublimes slowly and unchanged at 105°/1-2 mm

Bromine oxidises it to rhamnonic acid, with intric acid it yields a mixture of formic, oxalic, and trihydroxyglutaric acids C<sub>5</sub>H<sub>4</sub>O<sub>7</sub> On reduction, the corresponding alcohol rhamnitol is formed

The phenylhydrazone has m p 159°; methylphenylhydrazone, m p 124°, ethylphenylhydrazone, m p. 124°; allylphenylhydrazone, m p. 135°, amylphenylhydrazone, m p. 99°, diphenylhydrazone, m p 134°, benzylphenylhydrazone, m.p 121°,  $\beta$ -naphthylhydrazone, m p 170°. Rhamnose phenylosazone crystallises in slender yellow needles, slightly soluble in bolling water; m p. 190°–192°.

Methylrhamnoside has m.p.  $108^{\circ}-109^{\circ}$ ,  $[\alpha]_{D}-62^{\circ}$ ; acctone rhamnoside, m.p.  $90^{\circ}-91^{\circ}$ .

Other derivatives are rhamnose ethylmercaptal, m p 135°-137°; rhamnose oxime, m p 127°-128°; rhamnose anilide, m p 118°.

Fucose C<sub>6</sub>H<sub>12</sub>O<sub>5</sub>

is obtained by hydrolysing the polymer fucosan, which is a component of the cell-wall of many marine algae. It is also present in Fucus vesiculosus, in laminatia and Carragheen moss. The seaweed is washed with water and dilute hydrochloric acid, and boiled for 12 hours with 3 p c sulphuric acid. The acid is neutralised with baryta, the filtrate concentrated to a syrup, taken up with alcohol, concentrated again, and the sugar isolated as hydrazone Reduction with sodium amalgam gives d-fucitol (Votoček, 1913, 46, 3653), mp 153°–154°, which combines with ihodeitol to form i-fucitol.

Fucose crystallises slowly in minute needles, in p 130°-140° . It exhibits muta-rotation [ $\alpha_D$ ], being initially  $-112^\circ$ , falling to  $-74.4^\circ$  to  $-77^\circ$ .

The phenylhydrazone has mp 170°–172°, the phenylmethylhydrazone, mp 177°, phenylbenzylhydrazone, mp 172°–173°; diphenylhydrazone, mp 198°; p-bromophenylhydrazone, mp 181°–184°. Fuconic acid lactone has mp 106°–107°,  $\lfloor a \rfloor_p + 73$ ° to + 78 3°.

Epifucose. Fucome acid lactone, heated with pyridine at 145°-150°, is converted to epifuconic acid lactone (Votoček and Cervený, Ber 1915, 48, 658), which on reduction with sodium amalgam gives epifucose

$$\begin{array}{cccc} \mathrm{CH_3} \; \mathrm{CH(OH)} & \begin{array}{ccccc} \mathrm{H} & \mathrm{H} & \mathrm{OH} \\ \mathrm{C} & \mathrm{C} & \mathrm{C} & \mathrm{CHO} \\ \mathrm{OH} \; \mathrm{OH} \; \mathrm{H} \end{array}$$

a syrup,  $\begin{bmatrix} a \end{bmatrix}_0 - 9^\circ$ , whose phenylhydrazone and p-bromophenylhydrazone are identical with those of fucose

Rhodeose is the optical antipode of fucose (Muther and Tollens, Ber. 1904, 37, 306). It has  $\lfloor a \rfloor_p + 73^\circ$  to +75 2°, and forms derivatives having the same melting-point as those from fucose just described. Rhodeome and lactone has mp 105 5°,  $\lfloor a \rfloor_p - 76$  3° to -69 4°. Rhodeose is obtained from the glucoside convolvulin

The configuration of thodcose is shown by the following reactions (Hudson, J. Amei. Chem Soc. 1911, 33, 405, Votoček, Ber. 1910, 43, 476).—

(a) Oxidation of rhodeonic acid lactone gives l-trioxyglutanic acid

(b) Reduction of rhodeose gives rhodeitol, not attacked by sorbose bacterium

(c) The rhodeohexonic acid produced by applying the cyanohydrin reaction to ihodeose yields, on oxidation, saccharic acid

Hence thodeose is

Epirhodeose (Votoček, Ber 1911, 44, 362) Rhodeonic acid, heated with pyridine at 150°-160°, is epimerised to epithodeonic acid, the lactone of which on reduction gives epithodeose:

$$\mathrm{CH^3\,CH(OH)} \cdot \overset{H}{\overset{\circ}{\mathrm{C}}} \overset{H}{\overset{\circ}{\mathrm{C}}} \overset{H}{\overset{\circ}{\mathrm{C}}} \overset{H}{\overset{\circ}{\mathrm{C}}} \overset{H}{\overset{\circ}{\mathrm{C}}}$$

The sugar is a syrup which gives the same osazone as rhodeose, it oxidises to trioxyriboglutaric acid, m p  $~184^{\circ}\text{--}185^{\circ}, \left[\alpha\right|_{D} + 12^{\circ}.$ 

Isorhodeose (Votoček, Ber. 1910, 43, 476; 1911, 44, 819, 3287)

This occurs combined with rhodeose, rhamnose, and other compounds in the glucoside convolvulin, it is the optical antipode of Fischer's isonhamnose. The configuration is shown by the following reactions.—

(a) Oxidation with intric acid produces

i-trioxyglutaric acid

(b) Oxidation with bromine water gives isorhodeonic acid, which reacts —

(1) With pyridine at 150° to give antirhamnonic acid, which reduces to anti-hamnose. HCN converts this sugar to anti-hamnohexomic acid, which may be further oxidised, giving macic acid

(ii) With H<sub>2</sub>O<sub>2</sub> in presence of iron to give a methyltetiose, which oxidises to l-taitaric acid

Isorhodcosc is therefore

$$\mathrm{CH^3\,CH(OH)} \, \cdot \, \overset{\mathrm{OH}}{\overset{\mathrm{CH}}{\overset{\mathrm{CH}}{\overset{\mathrm{OH}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{CHO}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{\mathrm{Cho}}{\overset{Cho}}}{\overset{Cho}}{\overset{Cho}}{\overset{Cho}}{\overset{Cho}}{\overset{Cho}}}{\overset{Cho}}{\overset{Cho}}{\overset{Cho}}}{\overset{Cho}}{\overset{Cho}}{\overset{Cho}}}{\overset{Cho}}}{\overset{Cho}}}{\overset{Cho}}}}}{\overset{Cho}}}}}}}}}}}}}}}}}}}}}}$$

it has  $[\alpha]_p+31.5^\circ$ , and the melting-points of the phenylosazone and p-biomophenylosazone are respectively  $186^\circ-187^\circ$  and  $222^\circ$ 

#### Tetroses

By reduction of dioxyvalerolactone, a mixture of a methyltetrose and the corresponding alcohol were produced. The methyltetrose is a syrup, giving an osazone, in p. 140°-142°, and a phenylbenzylhydrazone, in p. 99°-100°. The methyltetritol is also a syrup, its tetrabenzoate melting at 136°-137° (Gilmour, Chem Soc Trans 1914, 105, 73)

Rhodeotetrose. Rhodeose is degraded by Wohl's method to a rhodeotetrose, the p-bromophenylhydrazone of which melts at 143°-144° (Votoček, Ber. 1917, 50, 35)

Chinovose  $C_8H_{12}O_5$  is only known as a constituent of the glucoside chinovin, which, on hydrolysis with alcoholic hydrochloric acid, yields ethylchinovoside  $C_8H_{11}O_4$   $OC_2H_5$ , originally termed chinovite,  $[\alpha]_D + 78^\circ$ .

Chmovite, when hydrolysed with dilute sulphuric acid, yields chinovose, which is a syrup, showing all the reactions of the carbohydrates. Phenylchmovosazone forms yellow needles, in p. 193°–194°.

Aplose  $C_5^\dagger H_{10}O_5$  is an altogether remarkable sugar, obtained together with dextrose on hydrolysing the glucoside apin present in the seeds and leaves of parsley. It contains a branched chain of carbon atoms, having the

constitutional formula

$$\begin{array}{c} \mathrm{CH_{2}(OH)} \\ \mathrm{CH_{2}(OH)} \end{array} \hspace{-0.5cm} \hspace{-0.5c$$

(Vongerichten, Annalen, 1901, 318, 121, 1902, 321, 71, Bei 1996, 39, 235).

It is a pale yellow syrup, optically mactive,

non-fermentable, does not yield furfurol, and yields isovaleric acid when reduced by hydrogen iodide and phosphorus. Bromine oxidises it to apionic acid.

The phenylhydrazone has mp. 135°; phenylosazone, m p 156°, and the bromophenylosazone,

mp 209°-212°

Digitoxose C<sub>6</sub>H<sub>12</sub>O<sub>4</sub> is obtained on hydrolysing digitoxin, one of the glucosides of digitalis crystallises in plisms or plates, in p 102°, and is dextro-rotatory It has the composition (Kiliani)  $CH_3 CH(OH) [CH(OH)]_2 CH_2 CHO$ 

Digitalose C7H14O5 is similarly obtained from

digitalin.

#### THE CARBOHYDRATE ACIDS

are those containing the same number of carbon atoms as the aldose carbohydrates from which they are derived. They are either mono- or dibasic. Ketoses yield acids with fewer carbon atoms than the original carbohydrate on oxidation.

The monobasic acids are formed on oxidising the aldehyde group to acid, generally by means of bromine or dilute nitric acid. More energetic oxidation converts also the primary alcohol

(-CH<sub>2</sub>OH) group into acid (CO<sub>2</sub>H)

Glucuronic acid, which is of importance physiologically, is a type of monobasic acid, in which the aldehyde group is intact, and the primary alcohol group oxidised to acid monobasic acids are transformed in aqueous solution, particularly on evaporation on the water-bath, into lactones which crystallise well. The dibasic acids similarly sometimes form double lactones

The most important property of gluconic and isometic acids is their behaviour on heating with pyridine or quinoline, when they are transformed into isomerides, rearrangements of the groups attached to the a-carbon atom taking place. Thus, when glucome acid is heated with quinoline or pyridine at 130°-150°, it is partially converted into mannonic acid. This behaviour has been of the utmost importance in effecting

the synthesis of the carbohydrates.

Glueonic acid  $C_6H_{12}O_7$  always contains a proportion of lactone. It is slightly Levorotatory at the instant of solution, but becomes dextro-rotatory as the transformation to lactone takes place, the final figure for a mixture in equilibrium of acid and lactone being about +19°. The lactone crystallises in slender needles, in p 130°-135° -1t has  $[\alpha]$  +68° when  $+19^{\circ}$ . first dissolved, falling to +19° for the solution in equilibrium with the acid. The calcium salt is sparingly soluble and characteristic, likewise the phenylhydrazide C<sub>6</sub>H<sub>12</sub>O<sub>6</sub> N<sub>2</sub>H<sub>2</sub>Ph, which forms colourless prisms, m p 200°

Mannonic acid. The lactone crystallises in

needles, m p.  $149^{\circ}$  –  $153^{\circ}$ ,  $[\alpha]_{\rm p}$  +  $53.8^{\circ}$ . The phenyl

hydrazide has m p 214°-216°.

Galactonic acid is obtained by oxidising galactose or hydrolysed lactose with bromine In the latter case, 100 grams lactose are boiled 4 hours with 100 grams 5 pc. sulphuric acid, the acid precipitated as barium sulphate, and the filtrate evaporated to 300 c c, cooled to 35°, and 200 grams of bromme added with constant shaking. After standing for a short time, the excess of bromine is removed by a current of an, hydrog n bromide by lead carbonate and silver oxide, and the solution is sturated with cadmium carbonate, and concentrated till the cadmium galactonate crystallises. The mother liquors yield a second crop, about 50 grams in all being obtained. This salt is purified and decomposed

with sulphure acid

Galactonic acid crystallises in needles, which, on heating to 100', lose water, forming the lactone The lactone crystallises with difficulty, it has mp 90°-92°, and forms a crystalline hydrate, mp 65°. The acid has  $\lfloor a \rfloor_D - 10.5$ °, which becomes -57° on heating. The lactone has  $[a]_D$  70°, and the crystalline hydrate  $[a]_D$  -58°. The cadmium salt is characteristic, the phenylhydrazide has mp 200°-205°. On heating with pyridine at 150°, galactonic acid is partially converted into d-talonic acid.

Lactobionic and maltobionic acids, obtained by the action of bromine on the corresponding disacchandes, are colourless syrups or amorphous

compounds

Glucuronic acid (glycuronic acid) C<sub>6</sub>H<sub>10</sub>O<sub>7</sub> is frequently found in urine, combined with a variety of substances, forming compounds of a glucosidic nature. It was synthesised by Fischer (Ber. 24, 521), by reduction of saccharic glucosidic nature acid lactone, and has the formula

## CO, II [CH OH], CHO

It is conveniently prepared from euvanthic acid, which is readily hydrolysed by dilute acids to glucuronie acid and euxanthon

$$C_{10}H_{16}O_{10}+H_{2}O-C_{6}H_{10}O_{7}+C_{13}H_{8}O_{4}$$

Euxanthic acid, known in commerce as Indian yellow or puri, is a pigment made almost exclusively in Bengal, and is obtained from the urine of cows which have been fed upon mango leaves (v. Indian Yellow)

According to Thierfelder (Zeitsch physiol Chem, 11, 388), I part of euxanthic acid is heated with 200 parts of water in an autoclave at 120°, the euxanthone separated by filtration, and the solution concentrated Owing to the introduction of aniline dyes, piuri has become very rare

and expensive

Neuberg and Lachmann (Brochem Zeitsch 1910, 24, 416) propose to obtain glucuronic acid by feeding menthol to rabbits. Menthol has the advantage that it is not poisonous, the paried glucurome acid is fairly easily isolated and readily hydrolysed 2 grams menthol are dissolved in 1 c c alcohol and shaken with 20 ce lukewarm water, this emulsion is administered to rabbits, and the urine collected, rendered acid, and extracted with other. Excess of ammonia is added to the ethereal solution, and the ether distilled off. The residue of insoluble ammonium menthylglucuronate is dissolved in water, precipitated by lead acetate, the precipitate decomposed with hydrogen sulphide, and the filtrate evaporated when pure menthylglucurome acid is obtained is decomposed by boiling with dilute sulphuric acid to give glucuronic acid

Glucuronic acid is a syrup, and is transformed into the crystalline lactone on evaporation. This is very soluble in water, it has m p. 175°, [a]<sub>D</sub>+19°. It reduces Fehling's solution, yields furfurol on distillation with mineral acids, showing positive reactions with oremol and

phloroglucinol, and is not fermentable

A number of substances when introduced

into the animal body are excreted in the urine as Levo-rotatory-paned glucuronic acid derivatives. These are isolated by precipitation with basic lead acetate, of which an excess should be avoided.

Saceharic acid 
$$C_6H_{10}O_2$$
  
 $OH H OH OH$   
 $CO_2H-C-C-C-C-CO_2H$   
 $H OH H H$ 

is obtained on oxidising dextrose and all carbohydrates which contain dextrose, also from sorbitol, glucome acid and glucurome acid. It is prepared from starch or dextrose (50 grams) by heating with nitric acid (350 grams) (sp gr 1 15) on the water-bath, and evaporating to a syrup whilst stirring continually. A little water is added, and the solution again evaporated The residue is dissolved in 150 grams of water, neutralised with a concentrated solution of potassium carbonate, 25 c c acetic acid (50 p c) added, and the solution evaporated to 80 c c On standing, the monopotassium salt separates, and is recrystallised from a little hot water the mother liquois, Neuberg (Biochem. Zeitsch 1910, 28, 355) has recently found 10 pc of an acid identical or isomeric with glucuronic acid

Saccharic acid is a syrup,  $[a]_D + 8^\circ$ , increasing to  $+22.5^\circ$ . The lactone  $C_6H_4O_7$  has m p  $130^\circ$ – $131^\circ$ ,  $[a]_D +37.9^\circ$ , decreasing to  $2.25^\circ$  for the equilibrium mixture with the acid. Characteristic is the monopotassium salt  $C_6H_9O_8K$ , formed on the addition of acetic acid in excess to a solution of the normal saccharate. This crystallises in small orthorhombic needles, sparingly soluble in water (1.12 parts per 100°). The diphenylhydiazide has m p 210°.

The continued action of intric acid forms tartaric and racemic acids, and, finally, ovalic acid. On dry distillation it decomposes with the evolution of carbon dioxide, yielding pyromucic and isopyromucic acids.

Mucie acid  $C_6 H_{10} O_8$ 

is obtained on oxidising galactose, lactose, laffinose, dulcitol, and all carbohydrates which yield galactose when hydrolysed.

100 grams of lactose are heated on the waterbath with 1200 c c, of nitic and sp gr 1 15 till the volume is reduced to 200 c c, this residue is taken up with an equal volume of water and left to crystallise, the yield is about 38 grams

Mucie acid is a microcrystalline powder, m p 213°-214°, it dissolves in 300 parts of water at 14°, or 60 parts at 100°. It is optically mactive, and cannot be resolved into optically active isomerides

It forms normal and acid salts with metals, and normal salts with alkaloids. The monophenylhydrazide has mp 190°-195°, the diphenylhydrazide, prepared by heating with phenylhydrazine acetate on the water-bath, has m.p 240°

### THE CARBORYDRATE ALCOHOLS

are widely distributed in the vegetable kingdom, where they occur, as a rule, uncombined with other substances. As a class they are readily soluble in water, crystallise well, and have a small

notatory power. On cautious oxidation they give in turn a reducing sugar, monobasic acid, and dibasic acid. They are obtained by reduction of the corresponding aldose or ketose carbohydrates with sodium amalgam, but, as a rule, owing to transformations in the alkaline solution, a mixture of alcohols is produced. They are not fermentable, and do not reduce Fehling's solution. Only those which occur naturally will be described here, though the alcohols corresponding to the synthetical sugars are known.

Erythritol C4H10O4

$$\begin{array}{c} \text{OH OH} \\ \text{CH}_2(\text{OH}) - \text{C} - \text{C} - \text{CH}_2(\text{OH}) \\ \text{H} \quad \text{H} \end{array}$$

occurs in a number of lichens, particularly the varieties of  $Roccella\ timetoria$ , where it is present in the form of the ether crythrin  $C_{20}H_{22}O_{10}$ , which is a diorsellmate of crythritol. This is saponified by boiling with baryta or lime

 $\mathbf{C_{20}H_{22}O_{10}+2H_{2}O=C_{4}H_{10}O_{4}+2C_{7}H_{8}O_{2}+2CO_{2}}$ 

Erythrin is heated for about 2 hours at 150° in a closed iron vessel, with a quantity of milk of lime somewhat less than sufficient to decompose it, the exclusion of air prevents the formation of a large quantity of resinous matter The liquid filtered from calcium carbonate, on evaporation at a gentle heat, deposits orcinol in fine crystals, and the mother liquors on further evaporation yield a crystalline mass of orcinol and crythritol, from which the former is extracted by ether (*i* De Luynes, Compt rend. 56, 803) Stenhouse (Chem Soc. Trans [2] 5, 222) dissolves the erythrin in a slight excess of milk of lime, boils for half an hour in a vessel with a long condensing tube to exclude air, precipitates the lime in the resulting solution with carbon dioxide, or (on a large scale) by exact neutralisation with dilute sulphune acid, and evaporates the filtered solution to dryness On digesting the residue for 20-30 minutes with a mixture of benzene, toluene, &c (bp. 110°-150°), in a metallic vessel provided with a condenser, an undissolved portion is left free from ordinol, which, on evaporation and filtering from resin, &c, deposits crystals of erythritol. These may be purified by washing with cold alcohol, pressing, and recrystallising once or twice from hot water

Erythitol crystallises in large, colourless, transparent quadratic prisms, having an adamantine lustre Sp gr 1 542-1 449. It melts at 126° (Liebermann, Ber 17, 873), and may be cooled far below its melting-point without solidifying

It is optically mactive even in presence of boax, and has no action on litmus, but the addition of borax renders it acid. Its taste is sweet, but less so than that of oremol. It is very readily soluble in water, its heat of solution being at 18°, 5·2 cals, and at 9°, 5·12 cals (Colson, Compt. rend. 104, 113), sparingly in cold alcohol and insoluble in other. It does not lose water below 100°, and does not piccipitate neutral or basic lead acetate, copper salts, or animonium lead nitrate (Stenhouse).

Erythitol is not altered by ammonia, by bromine, or by chloride of lime. Heated to 260° with solid potassium hydroxide, it yields oxalic and acetic acids with evolution of hydrogen. With fuming hydriodic acid, it gives secondary butyl iodide C<sub>4</sub>H<sub>8</sub> III It is attacled by atmospheric oxygen ir presence of platinum

black, forming trihydroxybutyric and oxalic acids (De Luynes, Compt. rend. 56, 803, Sell, ibid. 61, 741).

With dilute nitile acid, oxalic acid, taitaric acid, and a large quantity of hydrocyanic acid are produced; and with chromic acid and potassium permanganate, or with chromic mixture, formic acid, carbon dioxide, and oxalic acid are obtained (Przybytek, Bull Soc chim [2] 35, 108)

E Fischer and J. Tafel (Ber. 20, 1088), by the oxidation of crythritol with dilute nitric acid, procured a reducing sugar, erythrose, which formed a crystalline phenylosazone C16H18N4O2, m p 166°-167°, sparingly soluble in water, more readily in other and benzene. The tetranitrate C4 II6(NO3)4 forms large shining laminæ, m p 61°

When eightritol is dissolved in concentrated sulphuric acid, and the mixture heated to 60°-70°, the compound  $C_8H_{11}O_2(SO_1H)$ , is produced, which forms salts with inetals (Hesse,

Annalen, 117, 329)

Erythritol dissolves easily in chlorosulphonic acid, forming eightritol tetrasulphune acid C4H6(SO4H)4, which crystallises in snow-white masses consisting of small prisms - It is decomposed by hot water with regeneration of erythritol. The metallic salts are sparingly soluble

By heating crythritol for about 100 hours 12-15 parts concentrated hydrochloric acid, De Luynes (Ann Chim Phys [4] 2385, J 1864, 497) obtained erythritol dichlorohydrin C<sub>4</sub>H<sub>6</sub>(OH)<sub>2</sub>Cl<sub>2</sub> By raising the temperature to 120°-130°, the result is more quickly attained, or the compound may be produced by heating eightritol to 120°-130° in a current of hydrochloric acid (Przybytek, Ber 14, 2072) It melts at 124°-125° (m p 145°, De Luynes)

Eigthritol tetrachloride C4H6Cl4, produced by the action of phosphorus pentachloride, melts at 73°, and crystallises in prisms (Henninger, Bull Soc chim 34, 194, Bell, Ber 12, 1271) Dinitro-

dichlorhydiin  $C_4\Pi_6(Cl_2)(NO_2)_2O_2$ , has m.p. 60°. Erythritol dibromhydiin  $C_4\Pi_6(Br_2)(OH)_2$ is formed by heating with a saturated solution of hydrobiomic acid for 3 hours at 100° in a sealed tube. It is insoluble in water, and melts at 130° (Champion, Compt 1cnd 73, 114) Dinitiodibromhydiin  $C_4H_4(Bi_2)(NO_2)O_2$  is a flaky precipitate, mp  $75^{\circ}$  It does not detonate under the hammer (Champion)

The tetraformate forms long slender needles, m p  $150^{\circ}$ , the tetracetate has m p  $85^{\circ}$ , and

the tetrabenzoate, m p 186°-187°. (Characteristic is the dibenzylideneacetal  $_4^{\rm H}_6{\rm O}_4({
m C}_7{
m H}_6)_2$ , obtained on shaking with 2 parts of benzaldehyde and 3 parts of 50 p c sulphuric acid (Fischer, Bei 27, 1524) which forms slender needles sparingly soluble in hot water

It is not fermentable by yeasts Bacterium xylinum transforms it into a reducing keto-tetrose (Bertrand, Compt. rend. 126, 762), and it is attacked by bacteria in cow-dung and hay water, succinic acid in the one case and volatile acids in the other being the main products of change (Fitz, Bei 11 1890, 12, 474)

The anhydride αδ-oxy-βγ-butanediol is obtained by heating erythritol for some hours with acids, it is a liquid, b p 154°-155° at 18 mm

The second anhydride  $C_4H_6O_2$  was prepared by Przybytek (Ber. 17, 1091) by the action of potash on erythritol dichlorhydrin. It is a colourless mobile volatile liquid, b.p. 138°, sp.gr. 1.1322 It combines with water, acids, and ammonia, and forms compounds with hydroand aniline Its constitutional CH<sub>2</sub> · CH CH · CH<sub>2</sub> cyanic acid and aniline formula is

When heated with anhydrous barium oxide for 3 days at 200°-220°, the anhydride yields furfurane, when heated with phosphoius pentasulphide it gives thiophene.

Adonitol C<sub>5</sub>H<sub>12</sub>O<sub>5</sub>

$$\begin{array}{c} \text{OH OH OH} \\ \text{CH}_2\text{(OH)} - \text{C} - \text{C} - \text{C} - \text{CH}_2\text{(OH)} \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$$

is the only naturally occurring pentose alcohol, it corresponds to ribose, and is obtained on reducing this It is found in Adonis vernalis (Podwyssotzki, Arch Pharm 1889, 141) It separates from water in transparent prismatic crystals mp 102° It has a sweet taste, and is When shaken with benzaldeoptically mactive hyde and 50 pc sulphune acid, the dibenzylidene acctal is formed quantitatively, it crystal-lises in slender needles, mp 164°-165°. The The corresponding diformylideneacetal has m p 145° It is oxidised to a mixture of pentoses (Fischer, Ber 1893, 26, 633, 27, 2491, Meick, Aich Pharm 1893, 231, 129).

Mannitol or d-mannite C<sub>6</sub>H<sub>8</sub>(OH)<sub>6</sub>

is the alcohol corresponding to mannose was first found in the dried exudation from the manna ash, Fravinus ornus, by Proust, in 1806, and is widely distributed in plants. It is contained in the sap of the laich, in monkshood, in the leaves of Syringa vulgaris, in celery, sugar cane, in fungi, where it exceeds dextrose in quantity or even replaces it Thus Lactarius pallidus, when dried, contains from 10 to 15 p c (Bourquelot, Compt rend 108, 568), and Agaricus integer is also a good source Kylin (Zeitsch physiol Chem 1913, 83, 171) has found it in several Fucus and Laminaria species It also occurs in the sap of French beans on keeping It is prepared from manna by extraction with boiling alcohol Mannitol is obtained on reducing mannose, Levulose, or dextrose with sodium amalgam. In many cases it seems to be a fermentation product derived from trehalose (Busolt, Chem Soc Abstr 1913, 803) Its formation may be avoided by preserving plant extracts under sterilised conditions

It crystallises in colourless thombic prisms, тр 166° 100 parts of water dissolve 15 parts at 16°, 100 parts of alcohol dissolve 0 07 part at 14°. It is optically mactive, but on the addition of boiax it becomes strongly dextro-rotatory, a mixture containing 10 p c mannitol and 8 p c borax having  $[a]_D + 225^\circ$ . This mixture With caustic soda and other is strongly acid alkalis it becomes levo-rotatory

Mannitol has a slightly sweet taste, is not fermentable by yeast, sorbose bacteria oxidise it to Levulose, and it is attacked by many other It does not reduce Fehling's solution, or races interact with phenylhydiazine

Mannitol is easily oxidised An in presence of platinum black forms mannosaccharic acid, and a mixture of mannose and Levulose, which is also formed with hydrogen peroxide and an iron salt. Bromine gives mainly Levulose. Nitric acid transforms it into mannosaccharic acid and finally into glycollic acid and oxalic acids.

Derivatives—On heating at 280°, mannitol is converted by loss of water into mannitan  $C_6H_{12}O_5$ , which is formed at 100° in presence of strong acids. The hexanitrate, also known as intromannitol  $C_6H_*(NO_3)_6$ , is prepared by the action of a mixtune of sulphuric and nitric acids; it forms colourless accular crystals, in p 112°-113°,  $[\alpha]_0+40$ °, which explode violently on being struck or when suddenly heated

The dichlorohydrin has m p  $174^{\circ}$ ,  $[\alpha]_{\rm D} - 3.75^{\circ}$ , and its tetranitiate forms slender needles, m p  $145^{\circ}$ . The hexacetate (Franchimont, Ber 12, 2059) separates from alcohol in orthorhombic crystals, m p  $119^{\circ}-120^{\circ}$ ,  $[\alpha]_{\rm D} + 18^{\circ}$  The hexacetate

benzoate has m p 149°

The tribenzylidene acetal crystallises in slender needles, insoluble in water, in p. 218°–222° (Fischer, Ber. 28, 1975). Triacetone manntol (Fischer, thid. 28, 1167) forms colourless prisms, in p. 68°–70°,  $[\alpha]_p+12.5$ °. It tastes bitter. Mannitol monoacetone, treated with benzoyl chloride and quinoline at 15° for five days, yields tetrabenzoylmanntol monoacetone, in p. 122°–123°, which hydrolyses partially to tetrabenzoylmannitol, in p. 122°–123°,  $[\alpha]_p+8$ ° (E. Fischer, Ber. 1915, 48, 266)

For the conversion of mannitol into methyl-

a-pyren

sec Windaus and Tomich (Chem. Zentr. 1918, n 109).

The anhydride mannitan exists in amorphous and crystalline modifications, the latter having m p 137°,  $\left[a\right]_{\rm D}-24^{\circ}$  A second anhydride is mannide  ${\rm C_6H_{10}O_4}$ , which exists in several forms One obtained by reducing the dichlorohydrin crystallises in slender needles or large prisms, m p 119°,  $\left[a\right]_{\rm D}+94^{\circ}$ ; iso-mannide produced by the dry distillation of mannitol in vacuó, forms prisms, m p 87°,  $\left[a\right]_{\rm D}+914^{\circ}$ 

$$\begin{array}{cccc} \textit{d-Sorbitol} \text{ or sorbit } \mathrm{C_6H_8(OH)_6} \\ & \text{OH H} & \text{OH OH} \\ \mathrm{CH_2OH-C-C-C-C-CH_2OH} \\ & \text{H} & \text{OH H} & \text{H} \end{array}$$

is the alcohol corresponding to dextrose, from which it is obtained on reduction with sodium amalgam in faintly alkaline solution. Sorbitol was discovered in mountain-ash berries by Boussingault (Compt rend 1872, 74, 939), and occurs in the fruits of a large number of plants belonging to the Rosaccæ. It was found as a crystalline network on the dried heads of the fungus Boletus bovinus (von Lippmann, Ber. 1912, 45, 3431)

It is prepared from the expressed juice of mountain ash berries, which is fermented to remove carbohydrate, defecated with basic lead ace ate, and concentrated to a syrup in vacuo.

This syrup is taken up with alcohol, and usually gives crystals of sorbitol. In other cases an equal weight of 50 pc. sulphuric acid, and a little less benzaldehyde is added, and the mixture agitated. After 24 hours it sets to a crystalline mass of dibenzylidene sorbitol. This is separated, washed, and decomposed by boiling with dilute sulphuric acid.

The benzaldehyde is removed by steam distillation, sulphuric acid by baryta, and the benzoic acid extracted by ether. The liquid is concentrated to a syrup, and inoculated with a crystal of sorbitol (Vincent and Delachanal, Compt. 1916, 147). This method permits of the separation of sorbitol in small quantities

from plants

Sorbitol crystallises with 1 mol  $H_2O$  in slender needles, mp 55°. The anhydride has mp 110°. It is very soluble in water, sparingly so m alcohol. It has  $[\alpha]_D - 17.3$ °, which becomes +1.4° in presence of borax. It is faintly sweet, non-fermentable, but is oxidised by B-xylinum to sorbose  $(q, \nu)$ .

The hexacetate has m p 99°. The dibenzylidene acetal as obtained above is a mixture of two isomerides, one of which has m p. 163°-164°

Triacetone sorbitol has m p 45°

Styracitol, a naturally occurring substance isolated by Asahina (Ber 1912, 45, 2363) from Styrax abassia, is isomeric with the anhydrosorbitol obtained by reduction and hydrolysis of anhydromethylglucoside (Fischer and Zachs Ber 1912, 45, 456, 2068). This suggests that such anhydrides of glucose and of glucoside occur in nature.

**Dulcitol.** Duloin, Dulcose, Melampyrite, Melampyrin

is the alcohol corresponding to galactose

Occurrence —Found m a manna or clude preparation from a bulbous root of unknown origin from Madagascar (Laurent, Annalen, 76, 358, Jacquelain, ibid. 80, 345, in Scrophularia nodosa and Rhinanthus Crista galli (Eichler, Ueb. d Melampyrin, Moscow, 1885), in Melampyrim nemorosum (Hunefeld, J. pr. Chem. 4, 233, Gilmer, Annalen, 123, 372), in Euonymus europæa (Kubel, P. Pharm Chim. 85, 372), and in dulcite manna (Hecht, Annalen, 165, 148)

Preparation—Dulcitol is prepared from the Madagascar material by extracting with boiling water and allowing the filtrate to crystallise. To prepare it from the plants above-mentioned, they are gathered at a suitable time and dired A decoction is made of them, rendered alkalino with milk of lime, boiled, filtered, concentrated, and acidified with hydrochloric acid. Dulcitol crystallises out on cooling. Dulcite manna is extracted with boiling alcohol, dulcitol crystallises from the solution filtered hot. The manna yields 72 p.c. dulcitol (Hecht).

Dulcitol may also be prepared by the reduction of galactose with sodium amalgam, the soda produced being from time to time neutralised with sulphuric acid. It is obtained mixed

with mannitol on reducing lactose.

Properties.—Dulcitol, both of natural and synthetical origin, crystallises in colourless, highly lustrous monoclinic prisms with octahedral summits The crystals are gritty between the teeth, and taste very faintly sweet, m p. 118° One part dulcitol dissolves in 34.41 parts water at 165°; it is but slightly soluble in alcohol, even at a boiling temperature. It is optically mactive, even in presence of borax, the denivatives are also mactive. Its solutions are not coloured brown by potash, nor do they reduce alkaline copper solutions. Nitrie acid oxidises it to mucic acid, oxalic and racemic acids are also produced (Carlet, Compt rend. 51, 137. 53, 343) Yeasts do not ferment it, but other organisms, such as exist in cheese, convert it, in presence of calcium carbonate, into alcohol, lactic, and butyric acids A saturated solution containing dulcitol and borax in the proportion of 2 mols dulcitol to 1 mol borax is neutral, but if the proportion of borax is less, the solution is strongly acid, when the proportion of borax is greater the solution is alkaline (dulcitol being neutral and borax slightly alkaline) A solution of a mixture of dulcitol and sodium paratungstate has an acid reaction after boiling (Klein, Compt. rend, 99, 144)

Dulcitol sublimes partially on careful heating, at 200° it gives off water, and is converted into dulcitan; at 275° it decomposes

Sodium, potassium, calcium, barium and

copper compounds exist

Neither lead acctate nor subacctate precipitates dulcitol, but a white precipitate is thrown down with an ammoniacal solution of acctate. This is an unstable lead compound

Dulcitol heated with hydriodic acid in an atmosphere of carbon dioxide yields hexyliodide,

water, and iodine.

The hexantrate is obtained by the action of fuming intrie and sulphuric acids on dulcitol it crystallises from alcohol in beautiful colourless flexible needles, which melt between 68° and 72°, and give off intrie acid vapours continually, being thereby converted into the tetranitrate. This latter is soluble in alcohol, from which it crystallises in translucent prisms, mp. 130°–140°, and decomposes at 145°.

Dulcitol forms with hydrochloric acid an unstable hydrochloride  $C_0H_{14}O_0$ - $HC_13H_2O$  Similar hydrobromides and hydroiodides exist, they are rather more stable (Bouchardat, Ann.

Chim Phys [4] 27, 145)

Dulcitol yields chloro- and bromo- hydrins  $C_6H_{12}O_4Cl_2$  crystallises in tables insoluble in water,  $C_6H_{12}O_4Br_2$  and  $C_6H_{12}O_4BrCl$  are of much the same nature. The corresponding dulcitan bodies  $C_6H_{11}O_4Cl$ , &c, exist. When these bodies are dissolved in fuming nitric or sulphuric acids, the corresponding tetranitrates are formed, as  $C_6H_8(l_2(NO_3)_4)$ , &c

With the acids of the fatty and aromatic groups, dulcitol forms a number of neutral compound ethers. Some of these are ethers of

dulcitol and some of dulcitan

The hexacetate forms hard crystalline plates, mp 171°; the hexabenzoate has mp. 147°, both are insoluble in water and ether, sparingly soluble in cold alcohol

The dibenzylidene acetal forms needles, mp 215°-220°. Dulcitol diacetone crystallises in prisms of bitter taste, mp. 98°.

Dulcitol diacetone, m p 98°, may be fractionally crystallised into . Dulcitol  $\alpha$ -diacetone, m p.  $145^{\circ}$ – $146^{\circ}$ ; and dulcitol  $\beta$ -diacetone, m p.  $113^{\circ}$ – $114^{\circ}$ 

The α- form, when treated with benzoyl chloride and quinoline at 15°, gives monobenzoylduleitol α-diacetone, an oil which, when shaken in chloroform solution with 5N/HCl, gives monobenzoylduleitol, in p 155°-156°.

If the benzoylation is conducted at 100°, the first product is dibenzoylduleitol a-diacetone, mp 185°-186°, which with 5 pc HCl in cold acetic acid gives dibenzoylduleitol, mp 210° (see

E Fischer, Ber 1915, 48, 266)

Perseitol  $C_7\Pi_{16}O_7$  was discovered in the fruits of Laurus persea, and occurs in Persea quatissima. It is identical with d-mannoheptitol, synthesised by Fischer (Ber 23, 930, 2226) by reducing the lactone of d-mannoheptonic acid. It crystallises in colourless slender needles, in p. 188°; dissolves slightly in cold water (6 parts per 100), readily in hot water, and is insoluble in absolute alcohol. It has  $[a]_D = 1.3^\circ$ , but becomes slightly dextro-rotatory in presence of borax.

Dilute nitile acid in the cold converts it into mannoheptose, the sorbose bacterium oxidises

it to a ketcheptose

The heptanitrate forms needles, m p 138°;

the heptaacetate has m p 119°.

The optical isomeride of persented has not been found naturally, but is prepared artificially from *l*-mannoheptose.

**Volemitol**  $C_7\dot{H}_{16}O_7$  was obtained by Bourquelot by extracting dired *Lactarius volemus* with alcohol. It crystallises in slightly sweet slender needles, in p. 151°–153°,  $[\alpha]_D+1$  92°, which is not altered on the addition of borax. The constitution is not yet determined

An octitol  ${}^{\prime}_{s}H_{1s}O_{8}$  was obtained by Vincent and Meunier (Compt' rend 127, 760) from the mother liquous of the sorbitol preparation from the fruit of some of the Rosacea. The syrup had  $[a]_{i}=3.42^{\circ}$ , the compound with benzaldehyde crystallised in slender needles, in p. 230°; an isomeride formed at the same time having in p. 140°.

#### THE DISACCHARIDES.

The disaccharides consist of two monosaccharides, united through an oxygen atom. One residue is fixed through the oxygen of the aldehyde group in just the same manner as is dextrose in the glucosides. In the second residue, the aldehyde or ketonic group may be still present, in which case the junction takes place through one of the alcohol groups, or it may have disappeared, being concerned in the junction. The disaccharides are accordingly of In the first type, in which the aldetwo types hyde group is present, they reduce cupric salts, and behave just as dextrose does. In the second type, all these properties are absent Most of the best-known disaccharides are derived from hexoses.

In establishing the structure of a disaccharide, three points demand attention: (1) the nature of the constituent hexoses; (2) whether it represents an a- or  $\beta$ -glucoside; (3) which hydroxyl- group is concerned in the attachment of the two hoxose residues. As a rule, they form no really characteristic derivatives, so that

their isolation and identification is far from easy. Those of Type I. contain, like dextrose, an aldehyde group or its equivalent, and since they exhibit muta-rotation and exist in two modifications, there is no doubt that, like dextrose, they possess a closed ring structure rather than a free aldehyde group. In solution, therefore, they exist as an equilibrated mixture of interconvertible isomerides. This is in agreement with their crystallising properties and in marked contrast to the carbohydrates of Type II, equivorose and trehalose, of which only one form exists, and which are characterised by crystallising readily.

Accordingly, both halves of the molecule possess a  $\gamma$ -oxidic structure, but only one section retains the aldehydic grouping potentially functional

The disaccharides are selectively hydrolysed by enzymes, and are attacked with very different degrees of readiness by mineral acids

Important conclusions as to their structure are based on this behaviour, for the full discussion of which reference should be made to The Simple Carbohydrates, by E. F. Armstrong, ch. v. and vi. (Longmans, 1919)

The following table gives a list of the di, tiland tetra-saccharides and their products of hydrolysis —

### DISACCHARIDES

Carbohydrate Components

Tune I Ald hade group motentially

Type V Aldehyde group, potentially functional

Maltose Dextrose-a-glucoside Dextrose-β-glucoside Isomaltose Gentiobiose Dextrose-\(\beta\)-glucoside Cellose Dextrose-\(\beta\)-glucoside Lactose Dextrose-\(\beta\)-galactoside Isolactose Dextrose-galactoside Melibiose Dextrose-galactoside Turanose Dextrose and Levulose Vicianose Dextrose and arabinose Galactobioses Galactose-galactosides

Type II. No reducing properties.

Sucrose Dextrose and Levulose Trehalose Dextrose (2 mols )
Glocoxylose Dextrose and xylose

TRISACCHARIDES.

Type I

Mannotriose Dextrose and galactose (2 mols) Rhamninose Dextrose and rhamnose (2 mols.)

Type II

Raffinose Galactose and dextrose and lævulose

Gentianose Dextrose (2 mols) and lævulose Melezitose Dextrose (2 mols) and lævulose

TETRASACCHARIDES

Type II.

Stachyose Lævulose and dextrose and galactose (2 mols)

Sucrose, Saccharose or cane sugar. The preparation and properties of this are dealt with under sugar, accordingly only the more theoretical aspects of its properties will be considered here In contrast to most of the sugars it crystallises

exceedingly well, this is almost certainly due to the fact that one substance only is present in solution, and not a mixture of isomerides lacks both aldehydic and ketonic properties, being without action on Fehling's solution Considerable difficulty has been experienced in ascribing a constitutional formula to sucrose which is entirely satisfactory. It is probably a derivative of α-glucose, but since it is not attacked by maltase, it is not a simple  $\alpha$ -glucoside. The methyl fructosides of the butylene oxide type are not attacked by invertase, and are far more stable to acids than sucrose. These facts are far more in agreement with the assumption that the fructose in the disaccharide has an ethylene oxide structure. and this is confirmed by the hydrolysis of octamethyl sucrose to tetramethyl glucose and tetramethyl fructose The former was of the stable butylene oxide type, the latter of the reactive ethylene oxide type

Sucrose may be formulated accordingly as

On hydrolysis of sucrose the following series of changes take place, the final products consisting of the equilibrium mixtures A and C together, probably, with a small proportion of B (cf. Hawoith and Law, Chem. Soc. Trans. 1916, 109, 1314)

Sucrose  $\rightarrow \alpha$ -glucose  $\rightarrow \alpha$ -fructose  $\rightleftharpoons \alpha$ -fructose

 $\begin{array}{c|cccc} & & & & & & & & & \\ & & & & & & & & \\ & \beta \text{-glucose} & & \beta \text{-fructose} & & \beta \text{-fructose} \\ & A & Butylene- & B & Ethylene- & C & Butylene- \\ & \text{oxide forms} & \text{oxide forms} & \text{oxide forms} \end{array}$ 

Sucrose contains eight hydroxyl groups, as evidenced by the formation of oetacetyl-, octanitro-, and octamethyl- derivatives. It forms saccharates  $C_{12}H_{21}O_{11}Na$  with sodium or potassium hydroxides, and more complex saccharates with lime, strontia and baryta

Lactose or milk sugar  $C_{12}H_{22}O_{11}$  was first separated from whey by Fabrice Bartoletti in 1619, and termed nitrum seri lactis. It is found in the milk of all mammals. Human milk contains from 5 to 8 p.c. of milk sugar, cow's and ewe's milk, from 4½ to 5 p.c.; mare's and ass's milk, about 6 to 7 p.c. It has not been found in plants

Lactose is hydrolysed by mineral acids to dextrose and galactose; it is a  $\beta$ -galactoside, that is, the active aldehyde group is in the dextrose residue. It is uncertain whether the primary alcohol group of the dextrose molecule takes part in the union with the galactose residue. Accordingly, lactose has one of the following formula:—

 $C_6H_{11}O_6$ — $O \cdot CH_2 \cdot CH(OH) \cdot CH[CH \cdot OH]_2 \cdot CH(OH)$ (1)

 $C_6H_{11}O_5$ —O·CH[CH $_2$ ·OH] CH[CH OH] $_2$  CHOH)

The following details of the manufacture of lactose from whey are given by Aufsberg (Chem Zeit. 1910, 34, 885). Cheese makers usually send their supply of whey to a central factory, where the lactose is separated. Before the whey is concentrated, it is heated and treated with

from 1 to 3 p.c of acid (whey vinegai), in order to coagulate any albuminous substances present. The whey is then concentrated at a temperature of from 60° to 70° under reduced pressure, until it contains about 60 pc. of total solids, the concentrated liquid is run into rectangular vessels, each holding about 700 lities and surrounded by a cold-water jacket, and after the lapse of 10 hours the thick liquid is stirred, this operation being repeated at intervals the end of 24 hours, the pasty mass is submitted to centuringal action, the crude, moist lactose amounting to about 3.85 p.e. of the weight of the original whey, a further quantity of crude lactose, amounting to about 0 5 p c, is obtained by evaporating the mother liquor and allowing it to crystallise The crude lactose is now refined by dissolving it in water so as to form a solution having a density of 13° to 15° B, adding to the solution a quantity of bone charcoal and about 0 2 p c of acetic acid, heating the mixture to 90°, then adding a small quantity of magnesium sulphate, and heating the whole to boiling The flocculent precipitate is removed by filtration, and the filtrate is evaporated by heating under reduced pressure until it has a density of 35° B. Crystallisation is then allowed to take place, and the crystals are separated The lactose thus obtained is again subjected to a similar refining treatment, the final yield of crystalline lactose being dried in a centrifugal machine supplied with a current of hot air The dry product, amounting to about 25 pc of the weight of the original whey, is then ground and sifted See also MILK SUGAR

Milk sugar is deposited from its aqueous solutions in hemihedral rhombic or monoclinic crystals. The crystals are white, semi-transparent, hard, and grate between the teeth, their sp gr is 1.53–1.54. They contain 1 molecule of water ( $\rm C_{12}H_{22}O_{11},H_{2}O$ ), which they retain at 100° over  $\rm H_{2}SO_{4}$ , but give up completely at 130°, leaving a colourless melted mass of anhydrous milk sugar, which solidifies in the crystalline state on cooling, and is strongly hygroscopic.

If a solution of milk sugar is evaporated to dryness on the water-bath, an anhydrous milk sugar remains behind, which is not hygroscopic

Hydrated milk sugar is slightly hygroscopic, soluble in 5 or 6 parts of cold or  $2\frac{1}{2}$  of boiling water, from the latter solution the crystals only separate out slowly on cooling. An aqueous solution saturated at  $10^{\circ}$  has a density of 1055, and contains 1455 p.c. of hydrated milk sugar; when lett to evaporate it begins to deposit crystals as soon as the density =1063, and the solution contains 2164 p.c. of sugar. It is insoluble in alcohol or ether.

According to Hudson (J. Amer. Chem Soc. 1908, 30, 1767), lactose hydrate crystalises from aqueous solution at the ordinary temperatures, whilst above 95° the  $\beta$ -anhydrous form separates Schmoeger, and later Hudson, have described three forms of lactose. The  $\alpha$ -hydrate  $[\alpha]_{\rm D}+86^{\circ}$  and the  $\beta$ -anhydride  $[\alpha]_{\rm H}+35^{\circ}$  both change in solution into the equilibrium mixture or  $\gamma$ -isomeride of constant rotatory power  $[\alpha]_{\rm D}+55^{\circ}$ . Lactose was the first carbohydrate of which more than one modification was observed, and the relauonship of these isomerides has been the

subject of much discussion (see v Lippmann, Chemie der Zuckeiaiten). It is to be explained in the same way as the muta-rotation of dextrose.

Lactose loses the water of crystallisation at  $130^{\circ}$  when heated, turns brown at  $160^{\circ}$ , begins to decompose at  $175^{\circ}$ , forming lactocaramel  $C_{12}H_{20}O_{10}$ , and melts at  $203^{\circ}$ 

On oxidation with biomine water, lacto-biomic acid,  $C_{12}H_{22}O_{12}$  is formed, and this is converted into galactose and gluconic acid, when hydrolysed by boiling with mineral acids, proving that lactose is a galactoside. Nitric acid first causes hydrolysis, and then forms mucie acid from the galactose, and saccharic acid from the dextrose. Take dextrose, lactose gives rise to two series of derivatives, corresponding to the  $\alpha$ - and  $\beta$ - forms of the sugar already mentioned. It contains 8 hydroxyl- groups, and therefore forms octacetates on heating with acetic anhydride and a dehydrating agent. These are colourless, crystalline substances, and well characterised. Other lower acetates are less definitely characterised.

Lactose octacetate may be converted into acetochlorolactose, and this into methyl lactoside, which crystallises in needles, in p. 171°. The octacetate, obtained by heating with acetic anhydride, crystallises in rectangular plates or needles, in p. 86° (Herzfeld), 98° (Kremann, Schmoeger), 106° (Fischer and Armstrong), according to the method of purification. It is obviously a mixture. Acetochlorolactose forms prisms, in p. 57°-59° (Fischer and Armstrong, Bei. 35, 841), [a]<sup>20</sup>+76·2°. The isomeride

forms prisms, m.p.  $118^{\circ}-120^{\circ}$ ,  $[a]_{D}^{20}+73.5^{\circ}$  (see also Bodart, Monatsh. 23, 1, Dittmar, ibid 23, 865)

On heating with phenylhydrazine, acetic acid and sodium chloride in a boiling water-bath for 1½ hours, phenyllactosazone  $C_{24}H_{32}O_{9}N_{4}$  is formed. This does not separate until the solution is cold, and is thus readily distinguished from phenylglucosazone. The yellow needles melt at 200°, and dissolve in 80 parts of boiling water. With very dilute sulphuir acid they form an anhydride  $C_{24}H_{30}O_{8}N_{4}$ , which melts at 223°–224°, and is almost insoluble in water (Fischer, Ber. 20, 830). The phenylhydrazine residues are eliminated by boiling with benzaldehyde, lactosone being formed:

# C<sub>6</sub>H<sub>11</sub>O<sub>5</sub> O CH<sub>2</sub> [CH OH]<sub>3</sub> CO CHO

This is obtained as a colourless syrup on concentration of the aqueous solution in a vacuum, and is similar in properties to glucosone

Lactose forms characteristic hydrazones with some of the substituted phenylhydrazines. The amylphenylhydrazone has mp 123°, allylphenylhydrazone, mp. 132°, benzovlphenylhydrazone, mp. 128°,  $\beta$ -naphthylhydrazone, mp. 203°. Lactal  $C_{20}H_{12}O_{9}$ , which is analogous to glucal from glucose, has mp 184°–186°,  $[\alpha]_{\rm D}+27^{\circ}$ .

Alkalis decompose lactose, giving a variety of products. Strong alkalis produce ultimately lactic acid; on fusion with potassium hydroxide, oxalic acid is formed. The prolonged action of lime results in the formation of saccharins  $(q\ v)$ 

Hydrolysis —Lactose is hydrolysed with difficulty by cold mineral acids, and resists even

boiling with weak organic acids. Towards sulphune acid it is 1240 times as stable as sucrose, and slightly more stable than maltose. Stronger acids hydrolyse it to dextrose and galactose, the optical rotatory power of the solution changing from  $+52^{\circ}$  to  $+68^{\circ}$ .

Lactose is also hydrolysed by an enzyme lactase (q v.) which is present in Kephir grains, in almonds, and in some yeasts. It is not fermented by yeasts, with the exception of a few torula (S fragilis, &c; see Armstrong, Proc Roy Soc 1905, 76 B, 600) which contain lactase It is more susceptible to bacterial attack, and readily undergoes lactic and butying acid fermentations

The synthesis of lactose has not been effected, but an isometide named provisionally iso-lactose, was obtained by Fischer and Armstrong by the synthetical action of Kephir lactase on a concentrated mixture of equal parts of galactose and dextrose. It was isolated as phenylisolactosazone A galactosidoglucose was prepared by the same workers from acetochlorogalactose, and shown to be probably identical with melibiose

Maltose C<sub>12</sub>H<sub>22</sub>O<sub>11</sub> was perhaps obtained from starch by De Saussure in 1819 · it was prepared by Vairy, who failed to recognise it as a distinct substance. Dubrunfaut, in 1847, further examined it and gave the name maltose, but regarded it as a trisaccharide O'Sullivan showed, in 1872, that it is a disaccharide, and gave a tuller description of its characters

Maltose is primarily a decomposition product of starch, and its occurrence in nature is mainly under conditions where it has been derived from Thus it is found in the leaves of plants, in germinating seeds, and in malt, &c abnormal conditions, it is found in the urine It is a constituent of commercial glucose. Its occurrence in the form of a glucoside is doubtful, amygdalin does not contain maltose, and it is

improbable that lotusin does

Starch is converted by the enzyme diastase first into dextims and then into maltose Diastase also acts on glycogen, forming maltose amongst other products. It is also obtained when starch is hydrolysed by acids workers have denied the presence of maltose amongst the products of acid hydrolysis, but Fernbach and Schoen (Bull Soc chim 1912, 11, 303) have completely identified it They regard the processes of hydrolysis of starch by diastase and by acids as essentially identical in the first

Preparation —Maltose is always prepared by the action of diastase, usually malt extract, on starch paste; its purification from the dextrins present, which are soluble in alcohol, is often a matter of difficulty. Baker and Day (But Ass Report, Dublin, 1908, 671) recommend the use of ungerminated barley diastase, which converts starch into maltose, about 60 p.c., and a-amylodextrin: this can easily be separated by one precipitation with alcohol Precipitated barley diastase is added to a 3 pc starch paste, and hydrolysis conducted at 50° for 5 or 6 hours, and subsequently in the cold. The solution is evaporated to a thin syrup and poured into 95 p.c. cold alcohol, so that the resulting strength is about 80 p.c. The a-amylodextrin is precipidistilled off the maltose syrup, seeded with a few crystals, and a little alcohol added few hours it sets to a solid magma of pure maltose. To purify it further the sugar is triturated with strong alcohol, filtered, dissolved in a little water, and poured into boiling 95 p c alcohol, again filtered and crystallised from the filtrate

When malt diatase is used, the procedure is similar, but the entire removal of admixed dextrins requires repeated purification. Maltose of this degree of purity separates readily

from strong aqueous solution

Properties.—Maltose crystallises from water in plates which contain C12H22O11,H2O, from alcohol (sp. gr. 0.810) it crystallises in crusts, and at times in cauliflower-like or waity aggregations which contain  $C_{12}H_{22}O_{11}$ . The hydrate casily parts with its water at  $100^\circ$  in a current of dry air. The anhydride is but slightly soluble in strong ethyl alcohol, it is more soluble in methyl alcohol The hydrate dissolves more freely in both solvents. Both bodies are very soluble in water, but less so than dextrose The solution is optically active, the activity being less when the solution is freshly prepared than after standing

The specific rotatory power for dry maltose is  $[a]_i = 154^{\circ} - 155^{\circ}$ ,  $[a]_D = 139^{\circ} - 140^{\circ}$  at 15.5° in solutions containing 10 grams per 100 c.c. and under (O'Sullivan, Chem Soc Trans. 35, 771, and 45, 5) Meissel (J. pr. Chem [2] 25, 114)

gives the formula

$$[a]_{D} = 140.375^{\circ} - 0.01837P - 0.095T$$

to represent the specific rotatory power of dry maltose at 0°, in which P=percentage of maltose in solution, and T the temperature of observation Brown, Morris, and Millar give +137 93° The value depends on the concentration as well as on the temperature The initial rotation is  $[a]_D = +119^{\circ}$ Brown and Moriis have shown that maltose is first formed in this low rotatory modification

Maltose reduces Fehling's solution, 100 parts being equivalent to 625 parts of dextrose (O'Sullivan), or 619 parts (Brown and Heron, Chem Soc Trans 1879, 35, 618) It is inactive towards cupric acetate (Barfoed's reagent). The density of a solution containing 10 grams maltose in 100 c c at 15 5° is 1 0393.

Maltose is hydrolysed on heating with mineral acids to two molecules of dextrose. Sulphuric acid at 80° attacks it 1.27 times as readily as lactose. The weaker organic acids are almost without effect except after prolonged

Bromine oxidises maltose to maltobioric acid (Fischer and Meyer, Ber. 22, 194) This has the formula C<sub>12</sub>H<sub>22</sub>O<sub>12</sub>, and is hydrolysed by mineral acids to dextrose and gluconic acid. The behaviour towards maltase classes maltose as dextrose-a-glucoside, and the following constitutional formula is provisionally applied .-

The compounds of Maltose.—The octonitrate tated and removed by filtration. The alcohol is (Will and Lenz', Ber 31, 68) crystallises in lustrous needles, m p.  $163^{\circ}-164^{\circ}$ ,  $[a]_{D}^{20}+128^{\circ}$ The octacetate exists in two isomeric forms. Prepared by heating maltose with acetic anhydride and sodium acetate, it has m p. 159°-160°,  $[a]_{\rm D}+63^{\circ}$  in chloroform The a-isomeride has m p. 125°,  $[\alpha]_p+122^\circ$  (Hudson and Johnson, J Amer. Chem. Soc 1915, 37, 1276)

 $\alpha$ - and  $\beta$ -heptacetylchloro- and -biomomaltose have been described (Fischer and Armstrong, Ber. 34, 2895, 35, 840; Foerg Monatsh.

23, 44),

a-Methyl maltoside (Foerg, ibid 23, 44) crystallises in slender plates, in p 125°-127°. β-Methyl maltoside (Fischer and Aimstiong, Ber 13, 34, 2895; Konig and Knorr, wid. 34, 4343) forms concentrically grouped needles, m p. 93°-95°,  $[a]_D^{20}+70^\circ$ .  $\beta$ - Phenyl maltoside, m p. 96°, and menthyl maltoside (Fischer, Ber 1910, 43, 2521), have also been described

The hydrazones are not characteristic.

The phenylosazone C24H32O9N4 is prepared by heating maltose with excess of phenylhydrazine and acetic acid for 11 hours on the waterbath. It separates after the solution has cooled (difference from dextrosephenylosazone) cording to Fischer (Ber. 17, 579; 20, 821) it crystallises in slender bright yellow needles and not in aggregates, having m p 206°, Ost (Chem Zeit 19, 1503) gives 202°–208°.

J. L. Baker states that the osazone prepared from his pure maltose had mp 195°, and consisted entirely of characteristic broad prisms, resembling ribbons in the microscopic field The stellate aggregates of crystals present in the osazone, from maltose containing traces of dextrm, were absent Maltosazone does not form an anhydride

Maltose-p-bromophenylosazone has m p. 198°. The p-mitrophenylosazone crystallises in ied needles, m p 261° (Hyde, Ber 32, 1815).

Maltose-p-diammobenzoie acid

$$C_6H_3(CO_2H) \underbrace{\begin{array}{c} NH \\ NH \\ \end{array}} C_{12}H_{20}O_{10}$$

is formed in small quantities on prolonged boiling of concentrated solutions of the components. It crystallises anhydrous in plates, in p 235°, or hydrated with 1 mol,  $\rm H_2O$  in needles. The metallic compounds of maltose are amorphous and ill-characterised. It does not form double salts with the alkali chlorides.

Fermentation —Maltose is fermented only after hydrolysis to dearrose. This hydrolysis is effected by maltase, which is present in most yeasts but absent in S. Ludwigu, S. Marxianus, S. eviquus, S membranæfactens, some forms of S. anomalus, W. saturnus, and many others. Accordingly, these yeasts do not ferment maltose. Maltose is also fermented by most moulds, and readily undergoes lactic and butyric acid fermentation

Synthesis -- Maltose is perhaps formed to-gether with isomaltose by the action of concentrated hydrochloric acid on dextrose. It is also formed by the action of emulsin on concentrated solutions of dextrose (E F Armstrong, Proc Roy. Soc 1905, 76 B, 592) In each case, however, only amorphous or syrupy products could be obtained.

Isomaltose (Revertose, Gallisin) was obtained and repeatedly crystallised from 95 p.c. alconol Vol II -T

by Fischer (Ber. 23, 3024, 3687) by the action of 400 grams of cold concentrated hydrochloric acid on dextrose (100 grams) at 10°-15°. It is a viscid syrup,  $[a]_D + 70^\circ$ , and characterised by being non-fermentable and forming a crystalline phenylosazone, mp from 140° to 155° (Fischer),

according to the purity.

Scheibler and Mittelmeier (Bei 23, 3075) claimed that the gallism of Schmitt and Cobenzl (Ber 17. 1000, 2456) consists mainly isomaltose.

Gallisin is found in commercial glucose may be obtained as a white amorphous deliquescent powder, by treating the unfermentable syrup, after evaporation, with absolute alcohol, and finally with ether alcohol. It has a slightly sweet taste; is insoluble in ether, slightly soluble in absolute ethyl alcohol, more easily soluble in methyl alcohol and acetic acid. It reduces silver nitrate, and Fehling's solution, the cupric reducing power of 10 98 grams-that of 5 giams of glucose. It gives no precipitates with solution of lead acetate, mercuin chloride, ferric chloride, or barium chloride. On boiling with dilute acids it is converted into dextrose

Subsequently, a large number of products obtained in the hydrolysis of starch or of glycogen have been stated to be identical with isomaltose, but, failing any characteristic derivative, definite proof of its presence in such cases is lacking (v Lippmann's Chemie der Zuckeiarten) It is possible that gallisin and most of these products represent dextrins of low molecular weights or admixtures of these with maltose

By the action of maltase on a concentrated solution of dextrose, Croft Hill (Chem Soc. Trans 1903, 83, 578) has obtained a disaccharide which he terms Revertose Possibly this is

identical with isomaltose

By a similar process, E F Armstrong (Proc. Roy Soc 1905, 76, B, 592) has obtained an asomaltose which, like that obtained by the action of acids, is hydrolysed by emulsin isoMaltose is not attacked by maltase, and the behaviour towards enzymes suggests that it is dextrose-\(\beta\)-glucoside

According to Gatterbauer (Zeitsch. Nahr. Genussm 1911, 22, 265), the difficultly fermentable part of starch-syrup, formerly termed gallisin, consists of a new carbohydrate glucosin and small amounts of maltose, dextrins are

not present

Glucosin is isomeric but not identical with maltose, it is a yellowish-white powder, hydrolysed with difficulty by beer-yeast, more readily by yeast maltase or emulsin, to glucose

Dilute mineral acids give glucose, with concentrated mineral acids the reaction is reversible The phenylosazone melts at 97°-100°, the p-nitrophenylosazone at 240°

Cellose (Cellobrose) is prepared by hydrolysis of the octacetate formed by the action of acetic

anhydride on filter paper

To prepare the octacetate C<sub>12</sub>H<sub>14</sub>O<sub>11</sub>(OC<sub>2</sub>H<sub>3</sub>)<sub>8</sub>, 7 5 grams of finely divided filter paper are shaken with 20 c c acetic anhydride, the mixture cooled to 70°, and at this temperature a mixture of 7 c c acetic anhydride and 4 c.c. concentrated sulphuric acid is added. The temperature rises to 110°, the product is poured into 500 c c water, and the amorphous acetate collected, washed,

in colourless needles, m p 228°

Five grams are moistened with alcohol and hydrolysed with 15 pc alcoholic potassium hydroxide. A granular powder separates is dissolved in a little water, excess of alkali neutralised with acetic acid, and the solution evaporated to a syrup, which, after treatment with absolute alcohol and inoculation with a crystal, ultimately yields cellose as a colourless microcrystalline powder It tastes faintly sweet, dissolves in 8 parts of cold and 15 parts of hot water, and reduces Fehling's solution It has  $[\alpha]_D^{20}$  about  $+25^\circ$  when freshly dissolved, increasing to  $+35\,0^\circ$ . It is not fermentable, acids hydrolyse it to 2 molecules of dextrose, as also does the emulsin of almonds

The phenylosazone forms long yellow needles, m p.  $208^{\circ}-210^{\circ}$ ,  $[\alpha]_{D}-17.5^{\circ}$ , soluble in

135 parts of boiling water

Hudson and Johnson (J Amer Chem Soc 1915, 37, 1276) consider that cellose is composed of two glucose residues united similarly to the glucose and galactose residues in lactose. They quote an a-octacetate, mp 229°, [a], -|-42°, and a \$-octacetate prepared with acetic anhydride and sodium acetate, mp 202', [a] -15° On reduction of ocetobromocellose with zinc and acetic acid and hydrolysis of the acetyl derivative cellobial is obtained, in p 175°-176°, [a]<sub>D</sub>+1° Emulsin hydrolyses this to glucose and hydroglucal (Fischer and Fodor, Ber 1914, 47, 2057)

Hudson and Dale ( $l\epsilon$ ) give the specific iotation in water of the a- and B- forms of cellose as +72 0° and + 16 0° respectively

Gentiobiose is obtained by the action of invertase or of very dilute acids on the tri-saccharide gentianose. It may be prepared from old gentian roots in which this hydrolysis has already taken place by extraction with water and removal of the Levulose by fermentation with top-fermentation yeast separates from alcohol in anhydrous crystals, mp 190°-195°. It exhibits muta-rotation, the initial value for  $[\mathfrak{a}]_{D}^{20}$  being  $+9.6^{\circ}$  It is not fermented by yeast Emulsin, derived either from almonds or Aspergillus niger, hydrolyses it to dextrose The phenylosazone has mp

The  $\beta$ -octacetate, m p 195°,  $[\alpha]_D$ -6°, crystallises well, and may be used to isolate gentiobiose from gentian roots or other plant products The phenylosazone crystallises in stellate needles, in p  $160^{\circ}$ - $170^{\circ}$ , [a]<sub>D</sub>- $76^{\circ}$ . An isomeride obtained on heating this with acetic anhydride and zinc chloride has m.p.  $188^{\circ}-189^{\circ}$ ,  $[\alpha]_{D}+52^{\circ}$ .

Melibiose was first obtained, together with lævulose, by the partial hydrolysis of raffinose (Scheibler and Mittelmeier, Ber 22, 1678, 23, 1438); this may be effected by dilute acids of certain bottom-fermentation years. Bau gives the following methods of preparation

20 grams of raffinose in 250 cc water are sterilised and fermented for 1 day at 30° with a pure culture of a top-fermentation yeast The filtrate is sterilised and fermented during several days with a further portion of the yeast. It is

or from ethylacetate It is ultimately obtained then concentrated, poured into hot alcohol, and, after cooling, precipitated with ether syrup, after further treatment with alcohol and inoculation, crystallises with difficulty after

Alternatively, 20 pc solutions of raffinose are hydrolysed by boiling with 2 pc acetic acid, and concentrated to a syrup in a porcelain vessel. This is rubbed with 2 vols of 95 p c alcohol, the alcohol decanted, and ether added till a cloud appears After 2 days, this solution is decanted from any precipitate, inoculated with a few crystals, and set aside in closed vessels

Melibiose hydrate C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>,2H<sub>2</sub>O forms doubly refractive monoclinic plates:

[a b  $c=1.19227.20124, \beta=7716^{\circ}$ ]

m p 84°-85° The hydrate has  $\left[\alpha\right]_{D}^{20} + 129.5^{\circ}$ as stable value, and shows muta-rotation, the ınıtıal value, 5 minutes after solution, being The anhydride has  $\left[\alpha\right]_{D}^{20}+143^{\circ}$ very soluble in water, sparingly so in ethyl alcohol Strong acids hydrolyse it to dextrose and galactose it is slightly more resistant than lactose It is very sensible to traces of alkali It is reduced by sodium amalgam to melibiositol, a syrup which is hydrolysed to galactose and mannitol. It is attacked by bottom- and not by top-fermentation yeasts, and thus affords a means of distinguishing between these two varieties of yeast. The former variety contains the enzyme melibiase. It is slowly hydrolysed by emulsin

The following derivatives are known phenylhydrazone, light-yellow needles, m p 145°, allylphenylhydrazone, m.p. 197°; 8-naphthylhydrazone, m.p. 135°, phenylosazone, yellow needles from toluene, m.p. 178°–179°. This dissolves in 110 parts of boiling water, and interacts with benzaldehyde to form melibiosone The p-bromophenylosazone has mp 181°-182° The octacetate by  $\left[\alpha\right]_{D}+102^{\circ}$ 

Melibiose was the first natural disaccharide to be obtained synthetically (Fischer and Armstrong, Ber 35, 3144) It was prepared by the interaction of acetochlorogalactose with dex-

Trehalose (mycose, trehalabiose) was originally discovered in ergot, and has since been proved to be very widely distributed in fungi (Bourquelot) It apparently replaces sucrose in those plants which contain no chlorophyll and do not manufacture starch. Trehalose was found in 142 out of 212 species investigated, but only in certain parts of the fungus and at certain stages of the vegetative period. The quantity is a maximum just before the formation of spores When fungi are picked, trehalose is rapidly converted into mannitol, and after a few The trehala hours it has entirely disappeared manna from Syria and Persia is stated to contain 20 pc of trehalose. To prepare trehalose, suitable fungi (particularly Agaricus and Boletus species) are boiled with much strong alcohol immediately after picking. It was found as an incrustation on flowers of Carex brunescens (flowering rush) after a sudden sharp July frost (von Lippmann, Ber. 1912, 45, 3431).

Trehalose  $C_{12}H_{23}O_{11}$ ,  $2H_2O$  crystallises in lustrous glass-like rhombic prisms

[ $a \cdot b : c = 0.6814 \cdot 1.04171$ ;  $\beta = 1111^{\circ} 31'$ ] m p. 97°. It becomes anhydrous at 130°. It has  $[a]_{\rm b}^{20} + 178^{\circ}$  for the hydrate and  $+197^{\circ}$  for the anhydride, without any muta-rotation Acids hydrolyse it to 2 molecules of dextrose, but according to Winterstein, only with considerable difficulty, and it contrasts markedly in this respect with sucrose

It does not reduce Fehling's solution or form a phenylosazone, and has the formula

It is not hydrolysed by invertase and most enzymes, but is affected by a special enzyme, named trehalase, present in most fungi, moulds, and some yeasts, and conveniently prepared from Aspergillus inger. Accordingly, trehalose is fermented by those yeasts which contain trehalase. It forms few derivatives the octanitrate has m.p.  $124^{\circ}$ ,  $|\alpha|_{\rm p}^{18}+173.8^{\circ}$ ; the octacetate has m.p.  $97^{\circ}$ ,  $|\alpha|_{\rm p}^{1}+162^{\circ}$ , the tetrabenzoate, m.p.  $81^{\circ}-83^{\circ}$ 

Turanose  ${\rm C}_{12}{\rm H}_{22}{\rm O}_{11}$  is obtained from the trisaccharide melezitose, together with dextrose on hydrolysis with 20 p c acetic and solution. The dextrose is destroyed by yeast, and the turanose isolated from an alcoholic extract of the residue in transparent non-crystalline rounded grains of the composition.

They melt at  $60^{\circ}$ - $65^{\circ}$ , lose their alcohol at  $100^{\circ}$  and yield the pure sugar  $[\alpha]_{\rm p}$ + 7.18° without muta-rotation (Taniet, Compt. rend. 1906, 142, 1424)

Turanose has a reducing power of 60 (dextrose—100), and is hydrolysed by mineral acids to an equal molecular mixture of dextrose and levulose, and not to dextrose only, as originally stated by Alekin (Ann Chim Phys [6] 18, 532) Invertase, maltase, emulsin, and diastase are without action. The phenylosazone has mp. 215°-220° (Fischer, Ber. 27, 2486). It thus differs from sucrose in containing a free aldehydic group.

# OTHER LESS-KNOWN DISACCHARIDES.

Glucoapiose  $C_{11}H_{20}O_{10}$  is contained in the glucoside apin. It has not yet been isolated, as hydrolysis of apin by dilute acids yields only apiose and gluco-apigenin, and enzymes are without action

Galacto-arabinose was obtained by Ruff and Ollendorf (Ber 32, 352, 33, 1806) by oxidation of lactobionic acid from lactose—It is a dextrototatory syrut, the phenylosazone has m. p. 238°

10tatory syrup—the phenylosazone has mp 238° Glucoxylose. By hydrolysis of the glucoside dihenzoylglucoxylose, occurring in Daviesta latifolia, the carbohydrate glucoxylose  $C_{11}H_{20}O_{10}$ , is produced as an amorphous horny mass, very soluble in water (Power and Salway, Chem Soc Trans 1914, 105, 767, 1062)—It has  $\left[\alpha\right]_{\rm D}$ -36°, does not reduce Fehling's solution, and does not

give an osazone  $\,$  Acid hydrolysis yields glucose and xylose. Its constitution is therefore given as —

Strophantobiose  $C_{12}H_{22}O_{10}$  is obtained in the form of the methyl-derivative from the glucoside strophantin (Fiest, Ber. 31, 535, 33, 2063, 2069, 2091). It forms colourless crystals, in p. 207°, does not reduce Fehling, and is not fermentable. Acids hydrolyse it to mannose, thamnose, and methyl alcohol.

Vicianose  $C_{11}H_{20}O_{10}$  is prepared by Bertrand and Weisweiller (Compt. 1end. 1910, 150, 180) by hydrolysing the glucoside in the seeds of Vicia angustifolia by the enzyme occurring in them. It crystallises in aggregates of small needles, in purpose about 210°, has  $\lfloor a_{\rm D}^{20} \rfloor + 39.7$ °, the initial value being much higher. It reduces Fehling, and is converted into dextrose and arabinose on hydrolysis

Diglucose from amygdalin When amygdalin is hydrolysed by the digestive junce of Helix pomatia, the first product is a biose obtained as an amorphous powder. This is nonferimentable, and does not reduce Fehling solution. It probably has a constitution resembling that of trehalose (Giaja, Compt. rend 1910, 150, 793–796)

### TRISACCHARIDES

Mannotriose  $C_{18}H_{32}O_{16}$  is obtained by the action of either invertase or acetic acid on mannotetrose (stachyose). It is a colourless, faintly sweet crystalline substance, in p. 150°  $[\alpha]_{\rm b}+167^{\circ}$ . It reduces Fehling's solution Bromine oxidises it to mannotrionic acid, which is hydrolysed on warming with acids to 2 molsid-galactose and 1 mol gluconic acid. It is slowly hydrolysed by emulsin and by some yeasts, and is then ferimented, but whether completely so is not known.

The phenylosazone has mp 192°-194°, according to Neuberg, and 122° as given by Tamet Bierry (Compt rend 1911, 152, 465) confirms mp 122°-124°

Rhamniñose  $C_{18}H_{34}O_{14}$  is only obtained from the glucoside xanthorhammin present in the Persian berry (Rhamnus infectoria). In addition to the glucoside, the fruits contain an enzyme, termed rhamnunase, which hydrolyses it to the trisaccharide and rhamnetin. Rhamninose forms colourless crystals of faintly sweet taste, in p. 135°–140°,  $\lceil \alpha \rceil_D - 41^\circ$ , it reduces Fehling's solution. Acids hydrolyse it to galactose and 2 molecules of rhamnose. Sodium amalgam reduces it to the alcohol rhamnintol  $\lceil \alpha \rceil_D - 57^\circ$ , which is hydrolysed by acids to dulettol and 2 molecules of rhamnose. Bromine oxides it to rhamninome acid  $C_{18}H_{32}O_{15}$ , which is obtained, mixed with its lactone, as an amorphous substance; in p. 125°,  $\lceil \alpha \rceil_D - 94$  3°. It is hydrolysed to d-galactonic acid and rhamnose.

This behaviour points to the two rhamnose residues being united in the trisaccharide, but this disaccharide has not been isolated. Rhamninose

is not fermentable, and enzymes are without The octacetate has m p. 95°, [a]<sub>p</sub>-31°. action

Gentianose C<sub>18</sub>H<sub>32</sub>O<sub>16</sub> is found in the roots of a number of gentians, from which it is prepared by extracting the fresh roots with 95 p c alcohol It crystallises in colourless plates, m.p.  $209^{\circ}$ – $210^{\circ}$ , tastes faintly sweet, and has  $[a]_{\rm D}+31\cdot2^{\circ}$  to  $+33\cdot4^{\circ}$ .

It does not reduce Fehling's solution, but is hydrolysed by acids to lævulose and 2 molecules of dextrose, the final product having  $[a]_D-20$  2°. The enzyme invertase hydrolyses it to levulose and the disacchande gentiobiose, that of Aspergillus niger converts it into dextrose and sucrose.

Raffinose (also termed melitose, gossypose, melitriose) C<sub>18</sub>H<sub>12</sub>O<sub>16</sub> was first discovered by Johnston in 1843, in the manna from Eucalyptus mannifera, found in Tasmania. It is often found in considerable amount in the sugar beet, where it was discovered by Loiseau, Ritthausen found it in cotton-seed cake, it is also present in wheat and in oats To prepare raffinose from molasses, strontium oxide is added until a molecule is present for every molecule of sucrose After some hours, the strontium saccharate formed is separated by filtration and the liquid heated with excess of strontia, whereby the raftmose distrontionate is precipitated. The raffinose distrontionate is precipitated strontia compound is decomposed by carbon dioxide, the solution concentrated, and the process repeated to remove most of the sucrose The syrup finally obtained is dissolved in hot alcohol, the product which separates on cooling is kept under alcohol in a closed vessel, when raffinose slowly crystallises

According to Hudson and Harding (J Amer Chem Soc 1914, 36, 2110), cotton-seed meal is extracted with water, and, after purification, the solution is evaporated and barium hydrate added. The barium raffinate is powdered and washed with methyl alcohol The barium compound is suspended in water and decomposed with the exact amount of 50 pc phosphoric acid. The filtrate is completely freed from barium by sulphuric acid, and concentrated in a vacuum, seeded with inflinose hydrate and left to crystallise The yield is 4-5 pc of the weight of cotton-seed meal employed

Raffinose forms small needles, soluble in 6 parts of water at 16°, it is less soluble in ethyl alcohol than sucrose It is difficult to recognise when it crystallises with sucrose The hydrate melts at 80°, the anhydrous forms at 118°-119° The optical rotatory power [a]<sub>D</sub> is +104° without muta-rotation It has no reducing action, and behaves exactly as sucrose Dilute mineral acids hydrolyse it to levulose and melibose strong acids yield a molecule each of lævulose, dextrose, and galactose Invertase converts it into lævulose and melibiose, emulsin hydrolyses it to sucrose and galactose Accordingly, the formula is established as.

$$\begin{array}{c} C_6H_{11}O_5 - O - C_6H_{10}O_4 - O - C_6H_{11}O_5 \\ \text{Lee vulose} & \underline{Dextrose} & \underline{\textbf{Melbiose}} \end{array}$$

The undeca-acetyl derivative has mp 100°,  $[\alpha]_D + 92^\circ$ 

closely related sugars may be effected by means of selected enzymes For details of the method, see Davis (J Soc Chem Ind 1915, 35, 201)

Melezitose (melicitose) C<sub>18</sub>H<sub>32</sub>O<sub>16</sub>'2H<sub>2</sub>O, is

found in Briancon manna, which is obtained as an exudation from the young twigs of the larch (Abies tarix; Fi Mélèze) The manna from the Douglas fir consists mainly of melezitose (Hudson and Sherwood, J. Amer. Chem. Soc 1918, 40, 1456) It was investigated by Berthelot (Ann (him Phys [3] 55, 282) and by Alekhine (ibid [6] 18, 532). It is found also in turanibin or terenjabine, an exudation from Alhagi maurorum, growing in Afghanistan, Persia, and in the deserts of the East, and used in Northern India as a laxative (Villiers, ibid. [5] 12, 43) This substance is a brown, semisolid mass, of a slightly sweet taste, due mainly to the presence of dextrose. To obtain melezitose from this product it is treated with four times its weight of lukewaim water, and the liquid passed through a sieve to separate husks, leaves, &c, and after standing to allow of the deposition of earthy matter, it is evaporated to half its bulk. On standing for a few days, the greater part of the melezitose crystallises out The crystals are redissolved in twice their weight of hot water, the solution heated on the waterbath, diluted with its own volume of alcohol, heated to boiling to separate carthy and coagulable matter, and filtered whilst hot On cooling, melezitose separates out and may be obtained pure by two or three recrystallisations It forms large rhombohedral prisms

 $C_{18}H_{32}O_{16}, 2H_{2}O$ 

which effloresce in an and become opaque, they lose the whole of their water at 110°. Dehydrated melezitose has mp 148°-150°, 100 pts of water dissolve of anhydrous melezitose 26.8 pts at  $17.5^{\circ}$ , 34.4 pts at  $25^{\circ}$ , 75.6 pts at  $100^{\circ}$ , sparingly soluble in alcohol, insoluble in ether. The rotatory power is  $[a]_D + 88.5^{\circ}$ . Melezitose yields no crystalline compounds with alkaline chlorides. On boiling with dilute sulphuric acid it yields turanose and dextrose, the iotation falling to +63°. It has no action on an alkaline solution of copper, and is unchanged by diastase or yeast - Nitric acid transforms it into oxalic acid without the intermediate formation of mucic acid. It does not interact The undeca-acetate with phenyl hydrazine  $C_{18}\Pi_{21}O_{16}(OC_2H_3)_{11}$  crystallises in shining monoclinic prisms, m p  $170^{\circ}$ ,  $[a]_{D}^{20} + 110 4^{\circ}$ .

Stachyose (Mannotetrose, Lupcose) C24H42O21 was discovered by Schulze and Planta in the tubers of Stachys tubifera, and thought to be a trisaccharide. Tanret obtained it from ash manna, and termed it mannotetrose It has also been found in the twigs of white jasmine and in the subterranean parts of Lamium album. It constitutes from 60-75 p c. of the dry matter of the stachys tubers The tetrahydrate crystal lises in lustrous hard doubly refractive colourless plates (a b c=1.0512.1.0.4213,  $\gamma=90^{\circ} 46'$ ). It tastes quite sweet and dissolves at 13° in 0.75 part of water. The anhydride has m.p.  $167^{\circ}$ – $170^{\circ}$ ,  $[a]_{D}+148^{\circ}$ , without muta-rotation; the hydrate has [a]<sub>D</sub>+133°.

Stachyose is not acted on by Fehling's solu-The analysis of rafinose in admixture with tion or by alkalis. Acetic acid hydrolyses it

Sulphuric acid to lævulose and mannotriose. converts it into 4 monosaccharides, viz.: 1 molecule lavulose, 1 molecule dextrose, 2 molecules galactose. By the action of nitric acid, about 38 pc of mucic acid is formed Invertase hydrolyses it to levulose and mannotriose, which latter is slowly attacked by the emulsin of almonds or of Aspergillus niger

According to recent work of Schulze, stachyose is probably identical with the carbohydrate lupeose, present in Lupinus luteus and angustifolius, as lupeose has  $[a]_{D}+148^{\circ}$ , and gives the

same products on hydrolysis

Neuberg (Brochem Zeitsch 1910, 24, 173) prepares it by pressing the juice from the tubers, precipitating impurities with mercuric acetate, removal of the metal by hydrogen sulphide, neutralisation with ammonia, and concentration to a syrup, which is precipitated with alcohol The precipitate is dissolved in water, impurities precipitated with phosphotungstic acid, and the filtrate concentrated Barium hydroxide and alcohol are added alternatively After 24 hours, the barrum salt is collected, washed with alcohol, and decomposed by carbon dioxide. Reducing sugars are removed by this treatment and the stachyose readily crystallises

Polyamyloses have been obtained from potato starch paste by the action of Bacillus Macerans (Schardinger, Chem Soc Absti 1912, i 181, Pringsheim and Langhans, Ber 1912, 45, 2533) Tetra-amylose  $(C_6H_{10}O_5)_4$  or a-dextrin crystallises in colourless hexagonal plates,  $[\alpha]_D + 128^\circ$ , hexa-amylose or  $\beta$ -dextrin forms rhombic crystals,  $\lceil \alpha \rceil_D + 136^\circ$ On acetylation hydrolysis of the dextrins takes place, and from the acetyl derivatives formed diamylose and triamylose were obtained These crystallise in needles, and do not reduce Fehling's solution Rice starch gives similar definite polyamyloses on degradation (Pringsheim, Ber 1914, 47, 2565)

## Polysaccharides.

Glycogen  $(C_6H_{10}O_5)n$ , the reserve carbohydrate of the animal organism in which it appears to take the place of statch, is a regular constituent of all developing cells It is present in the liver up to 10 p c and in muscle

Preparation -Liver is finely minced and thrown into boiling water acidified with acetic The coagulated proteins are filtered off, and the remaining proteins precipitated from the filtrate with trichloracetic acid or potassium mercuric iodide (Brucke's reagent) Glycogen is precipitated from the filtrate with alcohol

Fresh finely minced liver is extracted for 2 hours with boiling 60 p c potassium hydroxide An equal volume of 95 pc alcohol is added to the clear filtrate, containing 15 pc KOH, and the precipitated glycogen washed with a mixture of 15 pc KOH and 2 vols of 95 pc alcohol.

It is purified in either case by solution and reprecipitation with alcohol, and obtained as an amorphous snow-white powder [a]<sub>p</sub>+191°, soluble in cold water, giving an opalescent solution. It is not fermentable, does not reduce Fehling's solution, and is not acted on by alkali, even by concentrated solutions at the boiling-With iodine it gives a brown-red colouration. Acids hydrolyse it ultimately to dextrose, but dextrins and maltose are formed as intermediate products Diastase derived from either plants or animals also converts it into dextrins and maltose (see Tebb, Journ. of Physiol 1898, 22, 423)

Glycogen may be obtained from yeast free from nitrogen, yeast-gum, and ash (Harden and Young, Chem Soc Trans 1912, 101, 1928) The crude preparation from yeast is freed from nitrogen and ash by repeated solution in water and reprecipitating with alcohol, the yeast-gum is removed by dissolving the crude glycogen in water and saturating with ammonium sulphate The last traces of ash are removed several times by several reprecipitations from water by alcohol

To estimate glycogen, the tissue is minced and boiled 2-3 hours with 30 pc potassium hydroxide The solution is diluted, filtered, and precipitated with alcohol The glycogen is again dissolved in dilute alkali and precipitated with alcohol Finally it is dissolved in water and estimated by hydrolysis with acids into dextrose (1 part = 0 927 part glycogen) or with the polarimeter

A 5 pc solution of sterilised glycogen treated with a potato culture of Bacillus macerans at 37° C, gave two crystalline cyclic polysaccharides (dextrin-\$\beta\$- or \$\beta\$-hexa-amylose (25 p c) and dextrin-a- or tetra-amylose (75 p c)) See Pringsheim and Lichtenstein, Ber 1916, 49,

Inulin. Inulin is widely distributed as a reserve material in plants having tubers or tuberous or fleshy roots, particularly in the antumn in dahlia tubers, Jerusilem artichoke chicory, garlie, onions, and other plants and bulbs

To prepare it, a mash is made of the ripe tubers, a little calcium carbonate is added, and the mixture extracted with boiling water filtrate is frozen and the sediment which separates dissolved in hot water and again frozen. Tanret advises precipitation with barium hydroxide, decomposition of this precipitate with carbon dioxide and subsequent precipitation with 95 p c. alcohol

Inulin, precipitated by alcohol, is a compact powder, resembling starch colourless molecular composition is uncertain, as the ordinary methods do not yield trustworthy results Kiliani gives  $(C_6H_{10}O_5)_6, H_2O$ , Brown and Morris,  $(C_6H_{10}O_5)_{12}, 2H_2O$ , Dull,  $(C_6H_{10}O_5)_{18}, H_2O$ , Tantet,  $(C_6H_{10}O_5)_{10}, 5H_2O$  It melts at 178°, having  $[a]_D-38^\circ$  to  $-40^\circ$ , sp gr In cold water it is sparingly soluble, but very soluble in hot water, and readily forms supersaturated solutions. It is precipitated by alcohol, when the percentage of alcohol in the liquid reaches 65.

The barium salt C<sub>36</sub>H<sub>62</sub>O<sub>31</sub>·3BaO (Tanret, Compt rend 116, 415) is insoluble

Inulin is hydrolysed by boiling the aqueous solution under pressure, by dilute acids, and also by the enzyme mulase, discovered by Reynolds Green, which is usually present in the same plant when in the germinating condition, but mert in the resting stage. In every case, levulose is the sole product of hydrolysis. Yeasts are without action

A number of products are described as

Levulins and Levulans, which probably represent polymerides of Levulose of lower molecular weight than inulin

Manna is the dried saccharine exudation from the stem of the flowering ash Frazinus ornus and F. rotundifolia, small trees cultivated for the purpose in Sicily A vertical series of oblique incisions is made in the bark when the trees are about 10 years old, the nuce slowly exudes and either dries on the stem (flake manna) or in wet seasons drops from the trunk, where it is caught upon tiles of cactus leaves, yielding an inferior variety. Flake manna is obtained in commerce in brittle stalactitic pieces, about 4-6 inches long and I inch wide It is yellowishwhite in colour, has a slight agreeable odour and a sweet taste It is used as a gentle laxative for infants and children

It consists principally of mannitol (70-80 p.c.) other constituents are mannotetrose (12-16 p.c.), mannotriose (6-16 p.c.) which is formed from mannotetrose, and small quantities of hexoses

The term is not restricted to the ash, and a number of saccharine exudations of varying origin and composition are classed as 'mannas' Amongst these are the Persian and Turkestan manna, derived from Alhagi mauroium of camelorum, the tamarisk manna, also of Persian origin, from the Tamaric gallica, oak manna, procured from Quercus valonia and Quercus persicus in Kuidistan, Australian manna, Eucalyptus viminalis, Briançon manna, from the larch, Larre europæa in Southern France, Astragalus manna, from various species of astragalus in Persia

Authorities -E von Lippmann, Die Chemie der Zuckerarten, 3rd ed 1904, L Maquenne, Les Sucres et leurs principaux dérivés, Paus, 1900; E Frankland Armstrong, The Simple Carbohydrates, London, 1919, Fischer, Untersuchungen über Kohlenhydrate, 1884-1908, Berlin, 1909 EFA

CARBOLIC ACID Phenol C. H. OH
Occurrence and Formation — Carbolic acid occurs in small quantities in castoreum (Wohler, Annalen, 67, 360), in urine (Stadeler, ibid 77, 18), and is a constant product of the putiefaction of albumen (Baumann, Ber 10, 685) formed when organic substances are strongly heated, and is contained in the tars obtained by the dry distillation of bones, wood (Duclos, Annalen, 109, 163), shale (brown coal), and coal (Runge, Pogg Ann 31, 69, 32, 308; Laurent, Ann Chim Phys [3] 3, 195)

The tar produced in gas manufacture was until the late war the chief source of carbolic Owing to the demand for carbolic acid as a disinfectant and for the manufacture of dyes and explosives, it has become usual, in tar distillation, to collect a special fraction, the 'carbolic oil' containing as much carbolic acid as possible before collecting the 'creosote oil,' which contains the other phenolic constituents of coal tar The 'carbolic oil' begins to come over when the specific gravity of the distillate attains that of water According to Watson Smith (Chem Soc Trans 1886, 21), ordinary Lancashire coal tar yields about 5 p.c by volume of crude phenols, containing 65 p c. by volume of carbolic acid crystallisable at ordinary temperatures

('oke ovens are now constructed so that the tar produced during the coking of coal can be collected, the following data, having reference to the percentage of carbolic acid in such tais, are given by Lunge (Coal Tar and Ammonia, ed. 1909), which has been largely drawn upon in the compilation and revision of this article. he says, The tar from Hussener's modification of the Carvès oven gave 1 37 p c of pure carbolic acid, the tar from the Simon-Carvès oven gave 0.05 p e. of crude carbolic acid, and similar ovens working with another kind of coal and at higher temperature gave 0 305 pe of crude acid The tar collected from blast furnaces, working with coal (raw) and also that obtained from Mond and Duft producers, contains a high percentage of phenohe constituents, but the quantity of carbolic acid present is very small?

Carbolic acid can be obtained by fusing potassium benzenesulphonate with caustic potash, the yield is proportional to the temperature and quantity of alkali employed, and is at best (96 23 p c of that theoretically possible), when I mol prop of the former is fused with 6 mol props of the latter at 252° Caustic soda cannot be employed instead of potash, since only a poor yield is obtained by its use (Degener, 7 pi Chem [2] 17, 394

Carbolic acid can also be obtained by heating the diazo-compounds of benzene with water

Preparation --- Carbolic acid is almost exclusively obtained from coal tar, and is worked up either from the special fraction known as carbolic oil' or from the oils intermediate in specific gravity (sp gr = 1.000-1.020) between the light and heavy oils, that is, distilling between 170° and 230°. The acid is present also in the crossote oils, distilling between 240° and 270°.

The preparation of carbolic acid from coal tar is effected by washing the oils obtained on distilling the tar with a solution of caustic soda, sp gr 1075-1100, when the tar acids are dissolved as sodium compounds and form a heavy solution upon which the now neutral oils float. It is not advisable to use a stronger solution of caustic soda, as in that case a considerable quantity of naphthalene would also be dissolved. If it is desired to extract carbolic acid only, without any of the higher homologues, the oils are often washed fractionally For example, a quantity of caustic soda solution more than enough to extract all the carbolic acid but insufficient to take up all the cresylic acid, is employed, and the phenate of soda solution drawn off is used to wash another portion of oil when the cresylate of soda is decomposed and the cresylic acid is replaced by carbolic acid, and a solution consisting almost entirely of carbolate of soda is obtained

Another method, suggested by Davis, is to employ, for the first wash, only one-third of the quantity of caustic soda required to extract all the tar acids when carbolic acid only is extracted. In many works, however, the whole of the tar acids contained in the light oil, carbolic oil, and creosote oil fractions are extracted, and after separating the tar acids in one of the ways about to be described, the mixture of carbolic and cresylic acids obtained is separated by fractional distillation. The phenate or carbolate of soda, drawn off from the oil is, in many c ses, first purified by blowing steam through it. This

carries away naphthalene, neutral oils, and pyridine bases, and the blowing with steam is continued until the water condensing from the escaping steam ceases to have a milky appearance. The purified phenate of soda is then treated with an acid to decompose it and set free the tar acids which float on the surface and are either skimmed off or are separated in an apparatus similar to a Florentine receiver.

Formerly the decomposition was almost always effected by sulphuric or hydrochloric acid, in which case the soda was lost as sulphate or chloride, which it did not pay to recover. The more common practice to-day, however, is to use carbon dioxide to effect the decomposition. If this is used under pressure, complete decomposition is said to be effected, but more generally the process is completed by the addition of a small quantity of sulphuric acid, this is the more necessary, as both carbolic acid and carbolate of soda are mutually soluble in each other.

The separated solution of sodium carbonate, which contains about 1 pe of carbolic acid, is recausticised with lime and is used over again to wash another quantity of oil A twofold saving is thus effected, in that the loss of the soda is avoided and also the carbolic acid which remained dissolved in the sodium sulphate of chloride solution, and was run to waste, is now recovered, masmuch as it goes forward with the regenerated caustic soda and is regained in the next washing operation. The clude carbolic thus obtained contains, besides phenol (carbolic acid), about 14 to 15 p.c. of water and variable quantities of eresplic acid. It occurs in commerce in three qualities, known as 75 s, 60's, and 50's crude carbolic acid, and its value is determined by 'Lowe's' test This is performed as follows 100 e e of the crude tar acid is slowly distilled in an 8-oz glass retort, and the receiver is changed when 10 e.e. of oil have come over, these 10 ce are accompanied by a varying amount of water, and by reading the quantity of water obtained, the percentage of water is ascertained The distillation is continued until 621 c.c. have collected in the graduated tube which is used as a receiver, the distillation is The 621 cc are then cooled, and then stopped a minute crystal of pure phenol is dropped into the tube, and the cooling continued until the contents begin to crystallise. The temperature is noted when the crystallisation appears to spread through the tube, and this is the measure of the quality of the carbolic acid

The preparation of pure carbohe acid from the crude article is always commenced by a distillation in wrought-iron stills, formerly without a column, but at the present day fractionating columns are almost universally employed. The water comes over first and then the carbohe acid, which is collected apart, so long as it crystallises on cooling. Some uncrystallisable acid comes over next, and the distillation is then stopped and the residue is mostly pitch.

In some cases, a little strong sulphuric acid and potassium dichromate is added to the contents of the still. The condensing worm must be made of zinc, silver, or stoneware. The distillate is allowed to cool and crystallise, and the crystals are drained from the liquid portion,

either on a drainer or in a centrifugal machine. The liquid portion is either sold as liquid carbolic acid or is worked up again with the next batch. The drained crystals are treated with a small quantity of concentrated sulphuric acid and potassium dichromate and redistilled, when pure' crystallised carbolic acid of commerce is obtained. In some cases the separation of the last traces of cresylic acid is effected by adding a small quantity of water, and the mixture is cooled, when a hydrate of carbolic acid crystallises out; this is separated from the liquid portion, which contains the cresylic acid and the crystals are redistilled Water comes over first, and then the boiling-point rises and pure carbolic acid distils over For medicinal purposes, a final distillation in glass retorts is performed

Properties —Pure carbolic acid crystallises in long, colourless needles, has a characteristic odour, melts at 42.5° or 43° and boils at 178.5° (Choay, Compt rend 118, 1211), Lunge (Coal Tar and Ammonia, 1909 ed 210), gives m.p. 42.2° and b.p. 184° under 760 mm. pressure. Ordinary pure carbolic acid of commerce, which contains traces of cresols, melts at a somewhat lower temperature and boils at 182° to 184°. The specific gravity of carbolic acid at 18° is 1.065 (Laurent); its expansion for temperature to at 760 mm. pressure is stated by H. Kopp as follows —

 $abla t = 1 + 0.0006744t + 0.000001721t^2 \\
-0.00000000050408t^3$ 

The presence of small quantities of impunities such as water, naphthalene, or pyridine, materially lowers the melting-point. The red colour of ordinary commercial acid seems to be induced by the presence of minute traces of metals, particularly lead (Meyke, Ber 16, 2513, of however, Ebell, abid 17, Referate 69, Hager, abid. 18, Referate 111, Yvon, Pharm. J Trans 1881, 1051, Richardson, J Soc Chem. Ind 12, 415, Bach, Mon Ser (A), 8, 508, Kraemer and Spilker, Ber 1890, 648) (Carbolic acid is a deliquescent substance, and in damp air forms a hydrate ('6H5 OH, H2O, which melts at 17 2° (Allen, Analyst, 3, 319). At the same time, however, it is not very readily soluble in water, 100 parts of water dissolving 4 83 parts at 11°; 5 36 parts at 35°, 6 19 parts at  $45^{\circ}$ , 7 33 parts at  $58^{\circ}$ , 11 83 parts at  $77^{\circ}$ , and at 84° both liquids mix in all proportions, whilst conversely, 100 parts of phenol dissolve 23 3 parts of water at 9°, 26 75 parts at 32°; 31 99 parts at 53°; and 40 72 parts at 71° (Alexejeff, Ber. 10, 410) According to Hamberg (*ibid* 4, 751), however, it dissolves in 15 parts of water at 16°-17°. It is soluble in all proportions in alcohol, ether, benzene, glacial acetic acid, glycerol, olive oil, chloroform, carbon disulphide, &c. The aqueous solution becomes colouted violet by ferric chloride and is without action on litmus According to Kramers, a mixture of benzene, toluene, xylene, naphthalene, anthracene, and phenanthrene, is obtained when the vapour of carbolic acid is passed through a red-hot tube (Annalen, 189, 129) Two isomeric mononitrophenols are obtained by the action of dilute nitric acid (sp gr 1.34) (Fritzsche, *ibid.* 110, 150); and at a lower temperature, the para- and at higher temperature

the ortho-derivative appear to be chiefly formed (Goldstein, Ber 11, 1943) By the further action of nitric acid, dinitrophenol (Kolbe, 4, 147, 67), and trinitrophenol (pierie acid) (Schmitt

and Glutz, ibid. 2, 52), result

Chlorine (Laurent, Annalen, 23, 60, 43, 209, Faust and Muller, ibid 173, 303, Benedikt, Monatsh 4, 233, Benedikt and Schmidt, ibid. 4, 606) and bromine (Korner, Annalen, 137, 200, 205, 208, 209, 210, Hubner and Brenken, Ber 6, 171) convert carbolic acid at the ordinary temperatures into mono-, di-, and trisubstitution derivatives, and at higher temperatures or in the presence of halogen carriers, such as antimony pentachloride, &c, into tetra- and penta- derivatives The nitro-, chloro-, and bromo- derivatives of carbolic acid are fairly strong acids, and the acid character increases with the number of nitro- groups or chlorine or bromine atoms introduced into the molecule Iodine yields mono- and di- derivatives in the presence of mercury (Hlasiwetz and Weselsky, Ber 2, 524) or rodic acid (Korner, Annalen, 137, 213) Sulphuric acid converts carbolic acid into two isomeric monosulphonic acids (Kekulé, Zeitsch f Chem [2] 3, 107), of which the orthoacid is the chief product at the ordinary and the para- acid at higher temperatures When heated with ammonium zinc chloride at 280°-300°. carbolic acid yields aniline, together with diphenylamine and phenyl ether (Merz and Weith, Ber 13, 1299) When fused with caustic potash, salicylic acid, metahydroxy-benzoic acid and two isomeric diphenols are formed (Baith and Schreder, ibid 11, 1332), whilst fusion with caustic soda results in the formation of pyrocatechol (catechin), resorcinol (resorcin), and phloroglucol (phloroglucin) (*ibid* 12, 417). The behaviour of sodium and potassium phenates (carbolates), when heated in a current of carbon dioxide, has been studied by Kolbe (J. pr. Chem. [2] 10, 89, of also Schmitt, J. pr. Chem. [2] 31, 410, D R P 29939, June 24, 1884), with sodium phenate, the reaction commences below 100°, is most rapid and complete about 180°. and continues up to the temperature of the decomposition of sodium salicylate with the formation of that compound as sole product, but with potassium phenate, salicylate is formed only at temperatures below 150°, whilst at higher temperatures the isomeric potassium parahydroxy-benzoate constitutes almost the entire product, in these reactions, one half of the phenol is recovered. When carbolic acid is heated with phthalic anhydride and a dehydrating agent, such as zinc chloride or sulphuric acid, it yields phenolphthalein (Baeyer, Ber 9, 1230, Annalen, 202, 68)

If the acid is heated with oxelic acid and sulphuric acid, it is converted into coralline (Zulkowsky, Annalen, 194, 119, 202, 184), from which pure aurine can be extracted by suitable methods (Dale and Schoilemmer, ibid. 196, 77), whilst benzaurine results from heating carbolic acid with benzotrichloride (Doebner-Ber 12, 1462)

Reactions -(1) In aqueous solution, carbolic acid gives a violet colour with ferric chloride; this, however, is neither developed in very dilute solutions (Sarauw, Ber 15, 46) nor in the presence of alcohol (Hesse, Annalen, 182, 161); limit of sensitiveness, 1 in 2000 (Polacci, Ber

7, 360) (2) The aqueous solution, on treatment with 1 vol. of ammonia and then with a few drops of an aqueous solution of bleaching powder (1 part in 20 parts of water), gives either at once or on standing a blue colouration; limit of sensitiveness, 1 in 4000 (Salkowski, Zeitsch. anal Chem 11, 316). (3) Bromine water, added to an aqueous solution of carbolic acid, gives either at once or on standing a yellowishwhite flocculent precipitate of tribromophenol bromide (Landolt, Ber. 4, 770); limit of sensitiveness, 1 in 43,700 (ibid ), 1 in 80,000 (Lunge) (1) A solution of carbolic acid assumes a deepred colour when boiled with one-third to one-half its bulk of Millon's reagent (a 10-15 p c. solution of mercurous nitrate containing nitrous acid), limit of sensitiveness, reaction distinct with 1 in 60,000, and still appreciable with 1 in 200,000 Salicylic acid gives a similar reaction (Plugge, Zeitsch anal Chem 11, 173, Almen, J. 1878, 1079)

Quantitative Estimation —Carbolic acid occurs in commerce in different qualities, as ciude acid, as liquid acid, and as crystallised acid of varying degrees of purity The percentage of phenols in crude carbolic acid can be approximately determined by shaking it with twice its volume of 10 pc soda ley added gradually, the separation of the oily and aqueous layers becomes more accurate if a known quantity of petroleum spirit equal in volume to that of the layer of indifferent and resmous substances, is added, and the amount added afterwards deducted. phenols are estimated in a portion of the aqueous layer by treatment with hydrochloric acid and subsequent addition of salt to ensure a complete separation. Such a correction is, however, hardly called for, since the phenols will dissolve about as much water as the water will dissolve phenols (Beckurts, Arch Pharm [3] 24, 572). Beckurts (*ibid* [3] 24, 580) also states that Koppeschaar's method (*v infra*) gives satisfactory results for testing liquid carbolic acid when the specimen is free from cresols thelaz (Bull Soc Chim 42, 574) tests measured samples of crude or liquid carbolic acid in tubes with water, with sulphune acid (1 to 1), and with soda solution (1 part soda ley (sp. gr. 138) to 9 parts of water), and finally fractionates a fourth sample

The exact quantitative estimation of carbolic acid in the pure product is always made by precipitating it with biomine. The composition of the precipitate was believed to be tribromophenol (Landolt) until it was ascertained by Benedikt that under certain conditions it was partially composed of tribromphenol bromide (cf Weinreb and Bondi, Monatsh 6, 506). Koppeschaar's method (Zeitsch anal Chem 15, 233), the one usually employed, requires the following solutions, (a) solution of sodium thiosulphate equivalent to a solution of iodine containing 5 pc. of rodine, (b) solution of starch, (c) bromine water (titiated with pure carbolic acid), or preferably a solution of bromine in caustic soda prepared by adding an excess of bromine to the soda and subsequently removing the excess either by boiling (Allen, J. Soc. Chem. Ind 1884, 64) or by evaporating the solution to dryness and again dissolving in water; (d) solution of potassium iodide containing 125 grams in the litre The process consists in treating 25 c.c.

CARBON 57

of the aqueous solution of carbolic acid (4 grams in the litre) with 100 c c, either of the solution of bromate and bromide with 5 c c, of concentrated hydrochloric acid to liberate the bromine, or of bromine water in a stoppered bottle, which is then well shaken, allowed to remain for 15 minutes, treated with 10 c c of potassium iodide solution, again shaken, and titrated with thio-sulphate. The reactions involved are the following —

(Weinreb and Bondi,  $l\,\epsilon$ ), so that, as Beckurts points out (lc infra), the assumption made by Koppeschaar in the preceding method, that 6 atom, props of bromine enter into combination with I mol prop of phenol, is in practice correct Beckurts (Arch Pharm [3] 24, 562) gives the following as the best mode of estimating carbolic acid. The solutions employed are (a) 1120 normal potassium bromide (5 939) grams in the litre): (b) centinormal potassium bromate (1 666 grams in the litre), (c) potassium iodide solution containing 125 grams in the litie, (d) decinormal sodium thiosulphate (24 8 grams of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>,5H<sub>2</sub>O in the litre) titration, 25 to 30 c c of the phenol solution (1 m 1000) are treated with 50 c.c. each of the potassium bromate and bromide solutions, shaken with 5 cc of concentrated sulphuic acid, allowed to remain 15 minutes, treated with 10 cc of the potassium iodide solution, and finally titrated with thiosulphate to estimate the amount of rodine liberated. The calculation is simple: from the 50 cc each of bromate and bromide solution, 0 2397 gram of bromide is liberated by the sulphuric acid, and this is capable of converting 0 0469 gram of carbolic acid into tubromophenol, I ce of the decinormal thiosulphate is equivalent to 0 008 gram of bromine, a quantity capable of converting 0 00156 gram carbolic acid into tribromomultiplying now 0.00156 by the number of cc of thiosulphate used, and subtracting the product from 0.0469, gives the weight in grains of phenol in the quantity of solution originally taken This process is accurate whenever solutions of pure carbolic acid, or mixtures from which the pure acid can be separated, are to be tested, but it fails in all cases (eq crude carbolic acid) when other phenols, particularly cresols, are also present Methods for the estimation of carbolic acid have also been proposed by Waller (Chem. News, 43, 152), Degener (J. pr. Chem. [2] 17, 390), and Chandelon (J. Soc. Chem. Ind. 1882, 203): but these are maccurate (Beckurts) Other methods have been proposed by Moeik (Chem Zenti Ind 1917, 36, 842; idem 1918, 265 T); and A biological method Petrie (idem 1919, 132 T) for the determination of phenol has been proposed by Blyth and Goodban (J. Soc Chem. Ind. 1907, 632), and an iodometric method by Skirrow (ibid 1908, 58); and Wake and Inglis have investigated the iodine values of the phenols (*ibid* 1908, 315) A method has been devised by Allen for determining phenols in carbolic soap (Analyst, 1886, 103)

Uses.—Carbolic acid has powerful antiseptic properties and is used on the large scale as a disinfectant. Large quantities of carbolic acid are also employed in the manufacture of salicylic acid, whilst in the colour industry, a considerable demand for the acid exists, masmuch as it is the source of pieric acid and coralline, and it is used in the preparation of some azo- colours has, of late years, been very largely used for the 'Lyddpreparation of modern high explosives ite; and 'melinite,' consisting very largely of piene acid and picrates, although trinitrotoluene appears now to be replacing pieric acid—Phenol forms condensation products with formaldehyde which are bodies of resinous appearance and properties, and which are proposed as substitutes for natural gums and shellac

Compounds with Metals —Potassium phenate  $C_6H_5$  OK, formed by dissolving potassium in earbolic acid (Hartmann, J. pr Chem [2] 16, 36), or by heating carbolic acid with caustic potash (Baumann, Ber. 10, 686), crystallises in slender white needles, readily soluble in water, alcohol, and ether Sodium phenate can be prepared by similar methods and may be used in the manufacture of saheyhe acid and as a disinfectant Compounds with barium (Laurent) and lead (Calvert, Zeitsch Chem 1865, 531) have also been described. W. H. C.

CARBOLINEUM. A preparation of crude carbolic acid Used as an antiseptic and preservative

CARBON. Sym. C At wt 12 Carbon bols at 3600° (Violle, Compt rend 1892, 115, 1273, 1895, 120, 868) Carbon occurs in nature in the free state, and very abundantly in combination, notably in the form of carbonates, and as an essential constituent of organic bodies. In the free state it is a solid, without taste or smell, exhibiting great diversity in the physical characteristics of its three allotropic forms—diamond, graphite, and charcoal

The diamond, up to the time of Bergman, was supposed to be a kind of rock-crystal, although Newton regarded it as probably an unctuous substance coagulated. It was shown to be combustible by the members of the Academy del Cimento in 1694, and Lavoisier proved that the sole product of its combustion was earbon dioxide He had previously ascertained that this gas was a compound of oxygen and an element to which he gave the name of carbone, contained in coal and wood Graphite, as its synonyms, blacklead and plumbago (from the Italian grafio prombino-the writing lead), imply, was long considered to be a kind of lead, or as related to antimony, Scheele regarded it as a compound of iron and carbon, Kastner Carbon proved that it was essentially carbon combines directly with hydrogen at 1100°-1200°, forming hydrocarbons (Bone and Jerdan, Chem. Soc Proc 1901, 162, Bone and Coward, *ibid* Trans 1908, 1975; 1910, 1219, Proc 1908, 222, Pring and Hutton, Trans 1906, 1591, Pring, ibid. 1910, 498) At high temperatures, such as the electric furnace, it is dissolved by the alkaline earth carbides and by rhodium, iridium, palladium, and platinum (Kahn, Compt rend. 1906, 143, 49; 1907, 144, 197, Moissan, ibid. 122, 1479). At the temperature of the electric aic, carbon reduces alumina (Moissan, ibid. 1894, 119, 935).


58 CARBON.

Manufacture of Wood Charcoal —The earliest | meiler' (Figs. 2 and 3) Both are distributed in plan of coaling wood, as the manufacture of much the same way.

charcoal was termed and is still called, is carried on as follows. A piece of ground is levelled at some convenent spot in the forest, which is termed the 'hearth' or 'earth'. In the centre of this a thick pole or bundle of brushwood is placed, around which the wood is arranged, some of the pieces being laid horizontally and others set up at an inclination, or the wood may be placed.

altogether at any steep angle, sloping outwards from the centre to form a flattened cone, which, when complete, is usually called a heap, the object, whichever way the wood is placed, is to obtain a free circulation of air under the heap to communicate with the chimney in the centre, which is formed by then withdrawing the centre pole or bundle of brushwood The large wood should, if convenient, be at the bottom of the heap, and the outside packed as close as possible, the heap is then covered with small brushwood, and afterwards with turt, or the material most impervious to air which can be conveniently obtained. A fire is lighted in the centre chimney, and by leaving openings in the outside covering at the bottom of the heap, the fire soon extends, and can be guided to any part by making temporary openings to admit the air. When the heap is sufficiently fired, all the openings are closed, and lastly the chimney itself. The fire will always extend most rapidly on the side facing or towards the wind, and great care must be taken to watch and check this, by keeping the covering on that side in good order charcoal burner must always be careful to spread the fire as evenly as possible through the heap, and after it is coaled to stop it down carefully, he can always accelerate the process in any part of the heap, if well built, by opening the outside to admit air freely, but if he finds this does not act, from any fault in setting the wood, he had better open a hole with a bar at the place required, and light a fire in the hole, this will soon communicate with the main fire in the heap As soon as the smoke and white flame cease to escape at the vents, the whole heap must be closed from the air as carefully as possible until the charcoal is quite cooled, and is ready to draw. The fire must never burn too fast, the slower the process, if the fire is steady and regular, the better the yield of charcoal Hard close-grained woods take a longer time to coal than soft open-grained woods, and should be placed in the heap accordingly These technical instructions, handed down in the forests for ages as secrets from father to son amongst the 'coalliers' in every country in Europe, are the results of long practical experience, and strictly accord with the true principles on which the process is based

To carbonise wood under a movable covering, the plan of meller, or heaps, is employed in Germany. The wood is arranged either in horizontal layers or in nearly vertical ones with a slight slope, so as to form conical rounded heaps of different sizes. The former are called 'lying meller' (Fig. 1); the latter, 'standing



In districts where the wood can be transported by means of rivers or mountain slides, a dry flat space must be pitched upon (screened from storms and floods), which may be walled round, having a slight declivity made in the ground towards the centre (v Fig 4). Into this space the tarry acid will partially fall, and may be conducted outwards through a covered gutter beneath into a covered tank. The mouth of the tank must be shut during the coking with an

Fig. 4 represents a walled meder station a, the station, b, the gutter, c, the tank, which is covered with the slab d, e, a slab which serves to keep the gutter clear of coals. The cover of the heaps is formed of earth, sand,

iron or stone slab, luted with clay A square

iron plate is placed over the inner orifice of the


gutter to prevent it being choked with coal ashes



Fig 4

ashes, or such other matter as may be most readily found in the woods. They should be kindled in the centre. From 6 days to 4 weeks may be required for charing a heap, according to its size, haid wood requiring more time, and the slower the process the better and greater is the product, generally speaking.

Chairing of wood in mounds (haufe or liegende worke) (Figs 5 and 6) differs from that in the meder, because the wood in the haufe is successively charted, and the charcoal




Figs 5 and 6

is raked out little by little. The product is said to be greater in this way, and also better. Uncleft billets, 6 or 8 feet long, being laid over each other, are covered with ashes, and then


CARBON 59

carbonised The station is sometimes horizontal and sometimes made to slope. The length may be 24 feet, the breadth 8 feet, and the wood is laid crosswise. Piles are set perpendicularly to support a roof made of boughs and leaves covered with ashes. Pipes are occasionally laid within the upper parts of the mounds, which serve to catch and carry off some of the liquid.

Fig. 7 is a vertical section, and Fig 8 a half bird's-eye view and half cross-section at the




height of the pit bottom, of Chabeaussière's kiln for making wood charcoal a is the oven, b, vertical air-pipes, c, c, horizontal flues for admitting air to the kilns, d, d, small pits which communicate by short horizontal pipes, c, e, with the vertical ones, f, the sole of the kiln, a circle of buckwork upon which the cover or hood h reposes, i, a pipe which leads to the cistem k;



l, the pipe destined for earrying off the gaseous matter; m, m, holes in the iron cover of hd

The distribution of the wood is like that in the horizontal meiler or heaps; it is kindled in the central vertical canal with buining fuel, and the lid is covered with a few inches of earth At the beginning of the operation all the draught flues are left open, but they are progressively closed as occasion requires. In 8 kilns of this kind 500 decasters of oak wood are carbonised, from which 15,000 hectolitres of charcoal are obtained, equal to 64,000 lbs. French, being about 25 p.c., besides tar and 3000 velts of wood vinegar of from 2° to 3° Baumé.

At Crouy-upon-the-Ourcq, near Meaux, there is a well-constructed kiln for making turf-chai-It resembles most nearly a tar kiln. In Fig 9, a is the cylindrical coking place whose surrounding walls are heated by the flame which passes through the intermediate space b. The place itself is divided by partitions of fire tiles into three stages, through the apertures in which the flames of the fire, c, c, rise, and heat the exterior of the coking apartment In order to confine the heat, there is in the enclosing walls of the outer kiln a cylindrical hollow space, d, where the air is kept stagnant Through the apertures left in the upper end at e, the turf is introduced; they are then shut with an iron plate, f, which is covered with ashes or sand The fireplace opens above this aperture, and its outlet is provided with a movable iron cover,  $q_i$ in which there is a small hole for the issue of the gases. The sole of the kiln consists of a cast-iron slab, h, which may be raised by means of a hook, i, upon it. This is drawn back after the carbonisation is completed, whereby the charcoal falls from the coking space into a




subjacent vault. The volatile products are carried off by the pipe k, and led into the condensing cistern, the gases escaping to the fire-place, where they are burned. The iron slab is protected from the corrosion of the acid vapours by a layer of coal ashes.


Charcoal obtained by the action of a rapid fire in close vessels is not so solid and so good a fuel as that which is made in the ancient way by the slow calcination of pyramidal piles covered with earth. According to Juon (Stahl and Eisen, 1904, 24, 1230), this is due merely to the difference in temperature of the two processes, the higher temperature used in the heapmade carbon increasing the carbon content One of the most economical ovens for making wood charcoal is that invented by M Foucauld, To construct which he calls a shroud or abri one of these, 30 feet in diameter at the base, 10 feet at its summit, and from 8 to 9 feet high, he forms, with wood 2 inches square, a frame 12 feet long, 3 feet broad at one end, and 1 foot at the other end.

The Figs. 10 and 11 will explain the

construction. The uprights, A, B, and C, D, of this frame are furnished with three pairs of wooden handles, a, a, a, by means of which they can be joined together, by passing through two contiguous handles a wooden fork, the frame being previously provided with props, as shown in Fig. 11, and covered with loam mixed with grass A flat cover of 10 feet diameter, made of planks well joined, and secured by four crossbars, is mounted with two trap doors, M, N (Fig. 12), for giving egress to the smoke at the



commencement of the operation, a triangular hole, P, cut out in the cover, receives the end of the conduit, Q, R, s (Figs. 12 and 13), of wood formed of three deals destined to convey the gases and condensed liquids into the casks F, G, H



Lastly, a door, r, which may be opened and shut at pleasure, permits the operator to inspect the state of the fire—The charcoal calcined by this abri has been found of superior quality

When it is wished to change the place where the abri is erected, and to transport it to a store of new-felled timber, the frame is taken down, after beating off the clay which covers it, the joints are then cut by a saw, as well as the ends of the fork which fixed the frames to one another. This process is economical in use and simple and cheap in construction, since all the pieces of the apparatus are easily moved about, and may be readily mounted in the forests. For obtaining a compact charcoal for the use of artisans, this mixed process of Foucauld is said to be preferable to either the close iron cylinder or the pile.

The fierce process for the production of charcoal, in which the wood is heated in brick kilns shaped like a bee-hive, of 32 feet diameter and 16 feet high in the centre, by means of the combustion of the permanent gases produced during a previous process, is said to produce a far higher percentage of charcoal and to involve a smaller loss of by-products (Dudley, J. Anal and Appl. Chem. May, 1891; Armour, Bowers, J. Soc. Chem. Ind. 1892, 152)

The charcoal made at Ontano from the local wood is of excellent quality for blast-furnace use. The wood is here heated in half-iron cylinder retorts, set honzontally and about 9 feet long and 4 feet high. The retorts are charged once a day, and are heated underneath by the tar produced mixed with the waste from the charcoal. The distillate is condensed by

straight condensers, one of which is attached to each retort (Goodwin, J. Soc Chem. Ind. 1902, 743)

In other processes wood is carbonised by superheated steam (Elfetrom, Proc Inst Civil Eng 1906, 164, 60, Zwilliger, J Soc. Chem Ind 1891, 129).

According to Rubands (J. Soc Chem. Ind. 1902, 1466), a hygienic and antiseptic wood charcoal is obtained by heating a mixture of aromatic plants, such as jumper, thyme, broom, &c., in an oven until the cresols begin to come off. The mass is then thrown into sheet-iron boxes, sealed to prevent combustion, and allowed to cool, the empyreumatic matters thus being reabsorbed

A large number of improvements in the apparatus employed for obtaining wood charcoal have been proposed (Jones, Eng Pat 1485, 1894, J Soc Chem Ind 1895, 24; Wise Waworth, Eng Pat 10522, 1897, ibid 1899, 128, Schenk, Eng Pat 23085, ibid. 1900, 321, Burey, ibid 1902, 909; Mourlot, ibid 1902, 1448; Blacher, Chem Ind 23, 508)

Chaicoal is also prepared from sawdust, wood, chips, and peat. The raw material is fed and pressed into and through heated retoits in a continuous stream, and comes out at the other end of the furnace as hard chaicoal. The retorts are provided with pipes for drawing off the by-products (Heidonstam, J. Soc. Chem. Ind. 1898, 445). Many other methods of converting peat into charcoal have been described (Rose, ibid. 1896, 643, Eng. Pat. 17568, 1895, Calmont, ibid. 1892, 257, Eng. Pat. 3978, Cohrab. ibid. 1893, 437, 1897, 1006, Eng. Pat. 3619, 1892, Jurgensen and Bauschlicker, Chem. Zeit. 1901, 25, 635, Stauber, J. Soc. Chem. Ind. 1906, 685; Jurgensen, ibid. 1907, 685)

For blast-furnace work, peat charcoal and that obtained from birchwood is the best

According to Maubér (*ibid* 1890, 881), charcoal is also produced from the by-products in the manufacture of spirits from grains and cereals Charcoal suitable for use as filters, respirators, and as wick for candles, can be prepared by placing vegetable fibres, such as cotton, cotton fabric or waste threads into drums, which are placed one by one into an inclined furnace, down which they roll to the lower end, which is provided with an exit door When no more gases are given off, the drums are withdrawn and the charcoal washed with water (Cooper, J. Soc Chem Ind 1902, 421)

For making gunpowder charcoal the lighter woods, such as the willow, dogwood, and alder, answer best, and in their carbonisation care should be taken to let the vapours freely escape, especially towards the end of the operation, for when they are reabsorbed, they greatly impair the combustibility of the charcoal (Taylor and Challon, ibid 1890, 106

The charcoal of some woods contains silica, and is therefore used for polishing metals. Being a bad conductor of heat, charcoal is employed sometimes in powder to encase small furnaces and steam-pipes. It is not affected by water, and hence the extremities of stakes driven into moist ground are not liable to decomposition. In like manner casks when charred inside preserve water much better than common

casks, because they furnish no soluble matter for termentation or for food to animalcules.

For making crayons of charcoal the willow is the best wood that can be employed, as the softness is uniform in all its parts The durability of charcoal may be seen in several of our old churchyards, where the letters made with lampblack are still perfect, though the white lead with which the body of the stones was painted is entirely destroyed. This property of carbon is shown, however, in a more striking manner by the writings that were found in the ruins of Herculaneum, which have retained their original blackness for two thousand years. The ancients wrote with mk made from ground charcoal.

If it be required to purify any carbonaceous matter to render it fitter for delicate pigments, this may be done by first calcining it in a close vessel, and then lixiviating it in water slightly

acidulated by nitric acid.

The incorruptibility of charcoal was well known to the ancients, and they availed themselves of this property upon all important occasions.

Some years ago a quantity of oak stakes were found in the bed of the Thames, in the very spot where Tacitus says that the Britons fixed a vast number of such stakes to prevent the passage of Julius Cæsai and his army These stakes were chaired to a considerable depth, had retained their form completely, and were firm at the heart

Desmond found that wood charcoal, when submitted to a high temperature out of contact with air, evolved a considerable quantity of gas consisting of 9:14 p.c.  $CO_2$ , 18 08 p.c. CO, 49 11 p.c.  $H_2$ , 16:04 p.c.  $CII_4$ , 0-26 p.c.  $O_2$ , and 737 pc. N<sub>2</sub>, its antiseptic properties being superior to carbon monoxide (Compt. rend 1894, 119, 733).

Most of the houses in Venice stand upon piles of wood, which have all been previously charied for their preservation In this country estates were formerly marked out by charred stakes driven to a considerable depth into the ground These are occasionally found, and usually the charred portions are quite perfect, although every other part is decayed (r Wood, DESTRUCTIVE DISTILLATION OF).

Wood charcoal possesses in a remarkable degree the power of absorbing gases in its poies, and in many cases of determining their combi-This power increases with its density As ordinary charcoal contains atmospheric air in its pores, it must for this purpose be prepared by heating it to redness in a close vessel, and cooling over mercury. The following table shows the absorptive power of Saussure boxwood charcoal of sp gr 157 for different gases.—

Ammonia gas .	. {	0 vols.
Hydrochloric acid gas	. 8	5 ,,
Sulphur dioxide	(	5,,
Hydrosulphuric acid	5	5 ,,
Nitrous exide	4	.0 ,,
Carbon dioxide .	3	5 ,,
Ethylene .	3	5 ,,
Carbon monoxide .		9.42 vols.
Oxygen		9.25 ,,
Nitrogen		6.5 4 .,
Hydrogen .		1.25 ,,

cocoa-nut has still greater absorptive properties, absorbing, according to Hunter,

Ammonia	. 171 7 vols
Carbon dioxide	677,
Carbon monoxide	21 2 ,,
Oxygen .	179,

In consequence of this absorption of gases in the pores of charcoal, their chemical activity is greatly increased When a piece of charcoal which has absorbed a considerable quantity of sulphuretted hydrogen is introduced into oxygen, a violent reaction immediately takes place, water and sulphur dioxide being set free. If air be employed instead of oxygen, sulphur is deposited.

This property is utilised in the removal of feetid and bad-smelling gases from rooms, and in the sweetening of bad-smelling liquids and of clothes For this purpose charcoal of average porosity is found to be the most efficient, and it should be in moderate-sized pieces, its absorbent power being greater in this state than when

in fine powder or in very large pieces

Potassium and sodium alloy absorbs chaicoal, and the suspension so obtained has a variety of uses, and constitutes an intermediate product in a variety of processes such as the production of potassium cyanide and so forth (Weintraub, J Soc Chem Ind 1909, 791)

When charcoal is boiled in a solution of platinum tetrachloride, it becomes impregnated with platinum, and is known as platinised charcoal. This substance possesses in a greatly increased degree the power of inducing chemical combination

Charcoal containing 2 p c of platinum causes oxygen and hydrogen to unite completely in about a quarter of an hour, the rapidity increasing with the percentage of platinum, charcoal containing a larger quantity of platinum acting like platinum sponge Platinised charcoal may be applied to the preparation of air-filters and respirators, and also as a mild caustic (Stenhouse, Chem Soc Trans 8, 105).

Charcoal is readily oxidised by dilute aqueous solutions of calcium hypochlorite, and potassium chlorate, especially in presence of osmium tetroxide, forming carbon dioxide, carbon monoxide, mellogen, and mellitic acid.

A further property of charcoal is its power of depriving most coloured liquids of then colouring matters, by absorbing them within its The colouring of red wine, cochineal, madder, or indigo, can thus be removed from solutions

This property was discovered in 1790 by Lowitz, and is made use of on the large scale in several industrial operations, particularly in the process of refining sugar. It is most strikingly possessed by bone-black or animal charcoal Charcoal of great decolourising power can be obtained by the dry distillation of birchwood or other substances containing carbon mixed with calcium or magnesium chloride, without access of air. The charcoal so obtained is washed with water, then with hydrochloric acid, and then again with water (Osticjho, J Soc Chem Ind 1900, 1099, 1902, 58) Chaicoal of higher decolourising power is

obtained from blood, horns, hoofs, clippings of hides, glue, &c, in contact with pearl-ash. Charcoal obtained from the shell of the good decolourising charcoal is also obtained by

carbonising vegetable matters mixed with chalk, calcined flints, or other earthy substances.

According to Halse (J. Soc Chem. Ind. 1903, 504), charcoal, especially applicable for use in decolourising and purifying saccharine and other liquids, and which can be used in acid solutions, is obtained by heating wood or a similar material with sulphuric acid to a temperature not exceeding 200°. The charted mass is then washed and dried, and, when spent, it can be revivified by treatment with a smaller amount of sulphuric acid.

Charcoal also precipitates or absorbs certain substances from solution, notably iodine, lime, and its salts (being used for the purification of highly calcarcous waters), lead salts and most metallic subsalts, besides many organic substances, as the bitter principles of hop, gentian, and aloc, tannin, alkaloids and resins, from their alcoholic solution (de Connick, Compt. rend. 130, 1627. Laval, Pharm. J. 1900, 65, 213., Malmegac, J. Pharm. Chim. 1900, 12, 5; Davis, Chem. Soc. Trans. 1907, 1666).

Crude alcohol is occasionally allowed to remain for some time in contact with charcoal

Carbon is used for the reduction of metals from their ores, and in the manufacture of glass for the reduction of alkaline sulphates.

Porous carbon can also be used with advantage in electrolytic cells (Lob. Zeit. f. Elect. Chem 1896, 3, 185) When used as an anode in strong hot sulphuric and other acids, which give off oxygen at the anode, carbon is dissolved, and this property can be used for depositing carbon on suitable cathodes (Coehn, J. Soc. Chem. Ind 1897, 1445)

(For the treatment of carbon articles before use, compare Thomson Houston, *ibid* 1905, 1230, Eng Pat 28062, 1904; Acheson U.S. Pat 74918, 1904)

Gas-carbon (clance coal) is a very dense form of carbon, deposited in the upper part of the retorts in the manufacture of coal gas and in blast furnaces. It often exhibits the lustre and sonority of a metal, is very hard, and a good conductor of heat and electricity. It is used to form the negative element in a Bunsen battery.

Materials used for manufacture of carbon electrodes are coke, hard and soft pitch, coal tar, and petroleum oil, in some cases, to obtain electrodes of better mechanical wearing power, a small portion of the coke is replaced by soot. The proportions of the various substances depend on the kind of electrode required Roush (J. Ind. Eng. Chem. 1909, 1, 286, Lake, J. Soc. Chem. Ind. 1900, 1001, Cooper, Shrewsbury and Marshall, ibid. 1896, 25, Starts, ibid. 1906, 544, U.S. Pat. 819606, 1906).

According to Nieweth (J Soc Chem Ind 1894, 1200), carbons of less resistance and greater flummating power when used in arc lamps, can be obtained by mixing the carbon with aluminous earths, magnesia, or other metalliferous minerals before moulding, after which process it is heated by means of an electric current. The carbon, when being heated, is supported in a holding sleeve of non-conducting material to prevent warping. Castner (J. Soc. Chem. Ind. 1894, 1067; Acheson, ibid. 1894, 407) adds 10 p.c. carbide of silicon to the carbon used for arc lamps; Rivers (ibid. 1906, 483; Kuffrath, ibid. 1907, 1903; Smith, J. Cennard, J. Soc. 300) adds yttrium salts. Langville (ibid. 1892, ibid. 1902, 1084).

935) describes a process for obtaining carbon for electrical purposes from paper pulp (Eng. Pat. 13847, 1892).

Carbon filaments are also prepared for use in meandescent lamps and electric lamps (Thomson Houston, J Soc. Chem. Ind 1905, 62; Pring and Fielding, Chem Soc Trans 1909, 1497, Chatelier and Wolagdine, Compt rend. 1909, 148, 1715, Howell, Electrician, 1905, 55, 588; Levis, J Soc Chem Ind 1905, 721, 882, Eng Pat 6959, 1904)

Graphte is formed in the manufacture of carborundum  $(q\ v)$ , and its production on a commercial scale has been established by Acheson (US Pat 568523). It is used to form lamp carbons, crucibles, electrodes, &c For the lower qualities of graphite articles anthracite may be employed, for the finer kinds, and for the manufacture of lubricating graphite, petroleum coke mixed with a catalytic oxide is used. For a discussion of the properties, mode of formation, and structure of graphitic carbon, v Kohlschutter, Zeitsch anorg. Chem. 1919, 105, 35

Lampblack is prepared on the large scale by burning fat, oil, resin, tar, &c, with an imperfect supply of air, either in a brick furnace or in cast-iron chambers.

The resulting dense black smoke is conducted into large chambers, where, on account of the extremely sluggish draught, it is allowed to roll about until the particles gradually coalesce into masses which after some time tall on the floor as soot. The condensation of the particles may be greatly facilitated by sending a current of electricity through the atmosphere of smoke, or else it can be agitated by mechanical means (Irvine, J. Soc. Chem. Ind. 1890, 1110)

For an account of the history and development of the lampblack and carbon-black industry, see Cabot, 8th Int. Cong Appl Chem 1912, Ve, 12-13

When commercial lampblack is strongly heated, a large amount of empyreumatic matter is given off, which condenses into a brown greasy mass consisting of chrysene, pyrene, capnomat, &c This causes ordinary lampblack to cohere when pressed. After calcination it loses this property.

Cressote or other suitable carbonaceous fucl is burnt in a furnace with an adjustable an supply, and the products of combustion are sent through a cooler, where by-products are condensed in centrifugal separators, the carbon deposited being collected in underlying receivers (Adam, J. Soc. Chem. Ind. 1907, 701, Eng. Pat. 13301, 1900)

An arrangement of lamps for the preparation of an especially fine quality of lampblack from waste fat and mineral oils is described in Payen-Paul, 64, 65.

Numerous improvements in the apparatus for the production of lampblack have been described (Wegelin, J. Soc. Chem. Ind. 1898, 747, 1899, 925, 1900, 56; 1902 1022; 1903, 1190, 1908, 81, 579, Fr. Pat. 380983, 1907; Irvine, J. Soc. Chem. Ind. 1890, 1110; Sanders, ibid. 1900, 753; Lembdner, ibid. 1903, 1238; U.S. Pat. 741726, 1903; Smith, J. Soc. Chem. Ind. 1909, 430; Meiser, ibid. 1908, 1212; D. R. P. 203711; Lennard, J. Soc. Chem. Ind. 1899, 1009: Ogilvy, ibid. 1902, 1084).

Carbon has also been obtained direct from smoke (Ward, bud. 1897, 902) livine (bud 1894, 532) obtains lampblack by spraying tars or oils rich in carbon over incandescent coke, mixing the gases so produced with air and steam, and burning them with a limited supply of air.

Lampblack, carbon black, also sometimes spoken of as gasblack, hydrocarbon black, jetblack, and so forth, is used in the production of printer's ink, paints, immeral-black, stove polish, shoe leather, rubber goods, fertilisers, colouring cement, mortal pulp, and artificial stone, harness oil, stenciling, &c. It varies in composition and properties with the material and method used in its production. When produced from the natural gas in American oil wells, the carbon-black differs from ordinary lampblack, by mixing with water on being shaken with it, ordinary lampblack will not do this. It is also quite different in appearance from lampblack obtained from oils, having a fine velvet-like gloss

(See description of figure and apparatus for manufacture, Cabot, J. Soc. Chem. Ind. 1894, 130. Cf. also Purtle and Rowlands, ibid. 1908, 234; U.S. Pat. 877596, 1908, Prosch, J. Soc. Chem. Ind. 1903, 640, Buller, Min. and Eng. World, Oct. 28, 1911.)

A good lampblack, said to be superior to the best American blacks, is also obtained by sparking a mixture of hydrocarbons under pressure and with or without admixture of carbon disormon-oxide (Koppers, J. Soc. Chem. Ind. 1906, 13; Machtolf, ibid. 894, 1909, 482, 1050, D. R. P. 207520, 1907, Gollwert, J. Soc. Chem. Ind. 1909, 1050). According to Frank (ibid. 1900, 545, 1905, 29, 1044), the carbon, at the moment of liberation, is readily taken up by metals, it can thus be made to pass into silver, imparting to the latter the appearance of so-called oxidised silver.

According to Peters, lampblack suitable for use in the preparation of paints, can also be obtained by heating the waste carbons from aic lamps to 1200°-1400°, and then allowing the mass to cool slowly (see also Acceptane black)

Colloidal carbon. By the careful addition of sugar to concentrated sulphune acid, filtering and dialysing the liquid, a clear black stable solution of colloidal carbon may be obtained (Sabbatam, Kolloid-Zeitsch 1914, 14, 29).

Carbon monoxide, carbonous of carbonuc orde; forma anhydrade (O, sp gr 0 96716 (Raylengh), 0 96702 (Leduc) (arr = 1) V D 14 (-186°) (Wroblewski, Compt rend 98, 982) S H p 0 2346 S H v 0 16844 (Wredemann, Pogg Ann 157, 1) C E 0 003667 (Regnault) S (6°) 0 0287, (9°) 0 0269; (18 5°), 0 02315 (Bunsen) S alcohol (2°) 0 20356; (13°) 0 20416, (16°) 0 20566, (24°) 0 20452 (Bunsen),  $\mu_{\rm c} = 1~000301$ ;  $\mu_{\rm g} = 1~00035$ ;  $\mu_{\rm u} = 1~00030$  (Croullebois, Ann Chim Phys [4] 20, 136) Does not exactly obey Boyle's law, PV/P<sub>1</sub>V<sub>1</sub> = 1 00293 (Regnault) Its critical temperature is  $-138~7\pm0$  1, and the corresponding pressure  $34~6\pm1$  atmospheres, critical density 0 3110 It boils at  $-190^\circ$ 

Discovered by Lassonne, and independently by Priestley. Composition first established by Clement and Desormes.

Carbon monoxide is present in coal- and water-gas, from both of which it is sometimes

obtained industrially (Williams, J. Soc Chem. Ind 1891, 270, Claus, ibid 1890, 170)

Produced by the oxidation of charcoal at high temperatures (if Johnson and McIntosh, Trans Roy Soe Canada, 1913, 7, m 161) By the reduction of carbon dioxide by means of hydrogen, carbon, metals, &c, at a red heat, or over carbon made incandescent electrically and coated with a bad conducting material (Engels, J Soc Chem Ind 1901, 350) On the rate of reduction of carbon dioxide by carbon, see Rhead and Wheeler (Chem. Soc. Trans. 1912, 101, 831, which contains a number of references to previous work on the subject) In the reduction of metallic oxides by charcoal or coke By the dry distillation of many organic compounds, and together with hydrogen and carbon dioxide when steam is passed over red hot charcoal. By passing electric sparks through charcoal By passing electric sparks through carbon dioxide (Buff and Hofmann, Annalen, 113, 140) or by heating carbon dioxide to  $1300^{\circ}$ (Deville, Compt rend 59, 873) By heating dehydrated oxalie acid, or by heating dry alkaline formates and acetates with formic acid (Loim, ibid 82, 750). By heating an oxalate or formate or these acids with concentrated sulphuric acid

Carbon monoxide is also produced in considerable quantities from various explosives, such as mellinite, gun cotton, &c. It is evolved by an alkaline pyrogallol solution during the absorption of oxygen, unless a large excess of potash is employed (Clowes, Chem. Soc. Proc. 1895, 200)

Prepared by the decomposition of (i) oxalic or formic acid, (2) potassium ferrocyanide, or (3) potassium cyanide, by concentrated sulphuric acid (Boudouard, Compt. rend. 1899, 128, 98, 307, 822, Wade and Panting, Chem. Soc. Trans. 1898, 257)

$$\begin{array}{l} \text{(1)} \ \ (^{1}_{2}\text{H}_{2}\text{O}_{4} = (^{1}\text{O} + (^{1}\text{O}_{2} + \text{H}_{2}\text{O} \\ \text{(2)} \ \ (^{1}\text{H}_{2}\text{O}_{2} = (^{1}\text{O} + \text{H}_{2}\text{O} \\ \text{(3)} \ \ \text{K}_{4}\text{Fe}((^{1}\text{N})_{6} + 6\text{H}_{2}\text{SO}_{4} + 6\text{H}_{2}\text{O} \\ -6(^{1}\text{O} + 2\text{K}_{2}\text{SO}_{4} + 3(\text{NH}_{4})_{2}\text{SO}_{4} + \text{FeSO}_{4} \\ \end{array}$$

In the first case the gas is mixed with carbon dioxide, which must be removed by passing through potash or lime-water. In the last reaction it is free from carbon dioxide if the temperature does not exceed 200°, but above this point this gas and sulphur dioxide are produced by the reducing action of carbon monoxide upon sulphuric acid

$$H_2SO_1+CO-CO_2+SO_2+H_2O$$

It is hable also to contain a small quantity of hydrocyanic acid vapour

Dry calcium oxalate or barium oxalate is mixed with about  $\frac{1}{10}$  dry lime, and strongly heated, the gas passed through lime and dried  $\mathrm{CaC}_2\mathrm{O}_4 = \mathrm{CaC}_2\mathrm{I}_4 + \mathrm{CO}_2$ , any carbon droxide formed being absorbed by the lime Carbon monoxide is evolved in the manufacture of carbides, and is collected and stored. It is also produced in the decomposition of sugars

Carbon monoxide is manufactured practically pure by bringing a mixture of oxygen and carbon dioxide in contact with incandescent graphite, coke, anthracite, &c. As thus obtained, it may be used for the manufacture of special steels, the reduction of refractory oxides, the treatment of iron ores in the blast furnace, and for other

purposes (Loiseau, J Soc. Chem Ind 1908, 1112; Pat. 390673, 1908). Carbon monoxide passed over heated sodium or potassium hydroxide, soda lime, or calcium hydroxide, yields a formate and hydrogen, together with methane and ethylene. Carbon monoxide may be converted into hydrogen and hydrocarbons to the extent of 995 pc. by passage over hot lime several times, and it is suggested that in this way illuminating gas might be freed from this toxic constituent (Vignon, Bull Soc chim 1911, [iv.] 9, 18).

Properties - Colourless gas of metallic odour and taste, combustible, non-supporter of combustion. Absorbed by carbon, by several metals, and by a solution of cuprous chloride in hydrochloric acid or ammonia It acts as an energetic reducer It is a highly poisonous gas, producing giddiness, and ultimately asphyxia when inhaled 0 43 pc in air is fatal to man (Mosso, J Gas Lighting, 1902, 80, 1334), 0.2 pc. is also poisonous if breathed for any

When carbon monoxide is passed over dry silver oxide, the latter is reduced thus.  $CO + Ag_2O$ = Ag<sub>2</sub>+CO<sub>2</sub>. Ammoniacal silver oxide is also reduced, and the reaction may be employed for the detection of small quantities of carbon monoxide in air (Dejust, Compt. rend. 1905, 140, 1250; Berthelot, *ibid.* 1891, 112, 597; Habermann, Zeitsch. angew. Chem. 1892, 324) It reduces Barreswil's (Fehling's) solution, and in contact with metallic copper moistened with aqueous potassium hydroxide is rapidly oxidised when mixed with air The copper is converted into a peroxide, Cu<sub>2</sub>O<sub>3</sub> or CuO<sub>2</sub>, which is the active oxidising agent The rate of oxidation is increased when the copper surface is coated with iridium

0 02 pc carbon monoxide may be detected by the decolouration of a mixture of silver nitrate, and weak permanganate solution aciditied with nitric acid (Merwet, Compt. rend 1897,

According to Grehant (ibid 1891, 113, 289; 114, 309), 1 part of carbon monoxide may be detected in 10,000 parts of air, by aspirating the air through dog's blood, and testing the respiratory capacity of the blood before and after the experiment. When the air is passed under a pressure of 5 atmospheres, the test becomes still more delicate. It can also be detected by passing the gas through palladium chloride, when a metallic precipitate is formed and the colour indicates the quantity present 30000 to 300000 carbon monoxide can be detected in the atmo sphere by passing several litres of the suspected air through rodic acid, when the following reaction occurs .-

$$5CO + 2HIO_3 = 5CO_2 + H_2O + I_2$$

The iodine is absorbed in chloroform or carbon disulphide, when the relative tint indicates the quantity of carbon monoxide, or else the carbon dioxide is absorbed by standard potash solution and the carbon monoxide calculated (Gautier, Compt. rend. 1898, 126, 793, 871, 931, 1299; St Martin, Compt. rend. 14, 1036; Kinnicutt and Sandford, J Amer Chem. Soc 1900, 22, 14, Fillanger, Chem Zeit. 1903, 27, Rep. 126; Levy and Pécoul, Compt. rend. 1905, 140, 98; Fr. Pat. 349714, 1904; 142162, 1906; Morgan

and McWhorter, J Amer. Chem. Soc. 1906, 29, 1589; Goutal, Ann. Chim. anal. 1910, 15; Graham and Winmill, Chem. Soc. Trans. 1914, 105, 1996; Sinnatt and Cramer, Analyst, 1914, 163; Frobose, Zeitsch. anal. Chem 1915, 54, 1; Graham. J Soc Chem. Ind. 1919, 10 T.).

Carbon monoxide is oxidised quantitatively to carbon dioxide by passing it over yellow mercuric oxide heated to 100° (Moser and Schmid, Zeitsch anal Chem. 1914, 53, 217); Clowes recommends the flame cap test as an accurate and rapid method for detecting carbon monoxide (Brit Assoc Report, Chem Zeit. 1896), whilst to detect the presence of carbon monoxide in mines, Haldane and also Herman (J Soc Chem Ind. 1896, 854) recommend the use of mice, which are very rapidly affected by small quantities of carbon monoxide. For a method of determining small quantities of carbon monoxide in hydrogen as used for ammonia synthesis, see E K Rideal and Taylor, Analyst, 1919, 89

According to Leblane (Ann Chim. Phys. [3] 5, 223), it is chiefly the presence of this gas that causes the poisonous action of air in which charcoal has been burnt. The gases from a charcoal fire may contain up to 18 p c of carbon monoxide, those from a petrol engine 7.32 p c (in these gases the quantity of carbon monoxide exceeds that of the carbon dioxide), whilst the combustion products from gas-heaters contain from 1 part in 15,000 to 03 pc, according to the supply of air to the burner. A small proportion of carbon monoxide in a dusty atmosphere constitutes a dangerously explosive mixture. It combines with the red colouring matter of the blood-forming carboxyh@moglobin, and may be recognised by its absorption spectrum, which is almost identical with that of oxygenated blood, and is characterised by two bands between D and E. On the addition of ammonium sulphide, these disappear in the case of oxygenated blood, and the spectrum shows one band midway between D and E, but remains unchanged if carbon monoxide is present (Vogel, Ber. 11, 235; Hoppe-Seyler, Zeitsch anal Chem 3, 439, also Preyer, J 1867, 802; Nawrocki, J 16, 640, Gréhaut, Lindel Chem 1, 1868, 1864, July 1868, 1864, 1864, July 1868, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 1864, 18 Compt rend 87, 193; St Martin, ibid 1892, 1 4, 1006) According to Nicloux (ibid. 1898, 126, 1526, 1595), carbon monoxide is a normal constituent of dog's blood, but this has been denied by Buckmaster and Gardner (Proc. Roy. Soc. B, 1909, 81, 515). Carbon monoxide burns ordinarily with a blue flame, which, by previous heating, becomes red, generating carbon dioxide. The temperature of its flame in air is about 1400° (Valerius, J. 1874, 58). According to Wieland (Ber. 1912, 45, 679), formic acid is an intermediate product in the combustion of carbon monoxide, and can be detected by allowing the flame to play upon ice. When dry it is not changed by the electric current nor by ignited platinum wire, but when standing over water it is decomposed by a glowing platinum spiral (Buff and Hofmann, Chem. Soc. Trans. 1860, 12, 273). When not absolutely dry, it may be exploded with oxygen by the electric spark or by platinum wire heated to 300° or by spongy platnum at ordinary temperatures. 2 vols. CO unite with 'vol. O, forming 2 vols. CO<sub>2</sub>

Dixon (Phil. Trans 1884, 617) has shown that no action takes place when a spark is passed into a mixture of perfectly dry carbon monoxide and oxygen, but that a mere trace of moisture renders the mixture explosive The combination takes place very slowly in presence of small quantities of steam, and increases in rapidity with the quantity of steam present. Probable reactions:

$$\substack{ 2\text{CO} + 2\text{H}_2\text{O} - 2\text{CO}_2 + 2\text{H}_2 \\ 2\text{H}_2 + \text{O}_2 = 2\text{H}_2\text{O} }$$

Hence the steam acts as a carrier of oxygen to carbon monoxde (v Dixon, Chem. Soc Trans.

Small quantities of other gases than steam were tried. if the gas contained hydrogen, explosion occurred, if the gas contained no hydrogen, no explosion occurred. When a mixture of carbon monoxide and steam is heated to about 600°, a portion of the carbon monoxide is oxidised. If the carbon dioxide is removed as it is formed, the whole may be oxidised Meyer's experiments (Ber 19, 1099) seem to prove that a dry mixture of carbon monoxide and oxygen can be exploded if a very strong spark is used, and a sufficiently high temperature then obtained, and if the gases are under considerable pressure When sparks from an induction coil are passed through a mixture of carbon monoxide and steam, carbon dioxide, a little formic acid, and sometimes carbon, are formed (Dixon, Chem. Soc. Trans 49, 94) When to a mixture of dry carbon monoxide and hydrogen, oxygen insufficient for complete combustion is added, and the mixture exploded by the spark, carbon dioxide and steam are formed in a ratio dependent upon the shape of the vessel and the pressure up to a certain limit, called the 'critical pressure.' Above this pressure the ratio of the formation of carbon dioxide and water is independent of the shape of the vessel. The critical pressure becomes lower the larger the quantity of oxygen The ratio CO+H<sub>2</sub>O.CO<sub>2</sub>+H<sub>2</sub> remains constant so long as the vol. of the hydrogen is more than twice that of the oxygen, provided no steam condenses, and the pressure is above When the vol of hydrogen the critical pressure is less than twice the vol of oxygen, the above The presence of an mert ratio diminishes gas increases the formation of carbon dioxide and diminishes that of water, hence it lowers the value of the ratio  ${\rm CO} + {\rm H_2O} \cdot {\rm CO}_2 + {\rm H}_2$ . This ratio is called the 'coefficient of affinity' of the reaction (Dixon, Phil Trans. 1884, 617; Chem. Soc Trans. 49, 94). According to Brodie, when carbon monoxide and hydrogen are submitted to the action of the silent discharge, marsh gas is formed by synthesis, when pure and dry carbon monoxide is circulated through the induction tube, it is decomposed, carbon dioxide being formed together with other 'oxy-carbons' of the formulæ C<sub>4</sub>O<sub>3</sub> and C<sub>5</sub>O<sub>4</sub> (Brodie, Proc Roy Soc 21, 245).

Carbon monoxide is converted into carbon dioxide under the influence of nascent oxygen from chromic acid (Ludwig, Annalen, 142, 47), but not by ozone, either in diffused daylight or in direct sunshine (Remsen and Southwark, Amer. J Sci [3] 11, 136) It is likewise oxidised by palladium charged with hydrogen in presence of oxygen and water (Traube, Ber. 15, 2325, 2854; 16, 123; Remsen and Keiser, ibid. 17, Vol. II.-T.

83; Wieland, Ber. 1912, 45, 679); by mixing with oxygen and passing over platinum black; by nitric oxide (Hasenbach, J, pr. Chem [2] 4, 1), by heating with metallic oxides and with 'activated' many oxysalts Adsorbed on platinum it is rapidly converted into carbon dioxide in contact with oxygen.

When passed over finely divided metals, such as nickel, cobalt, or iron, the quantity of carbon dioxide formed depends on the temperature and the metal (Zimmermann, Ber. 1903, 36, 1231, Sabatier and Senderens, Bull Soc chim 1903, 29, 294, Berthelot, Compt rend 1891, 112, Van Breakedensen and Horst, Rec trav. 594 chim 19, 27, Smits and Wolff, Zeitsch. physikal. Chem 1903, 45, 199, Charpy, Compt rend. 1903, 137, 120; Boudouard, *ibid* 1899, 128, In some cases a volatile carbonyl compound is formed of the type M(CO)4 (Lange and Quincke, Chem Soc Proc 1890, 112, 1891, 117, Wartha, Chem Zeit 1891, 15, 915, Garnier, Compt rend 1891, 113, 189, Gautz, Bull Soc chim 1892, 13, 278, Bertholet, 1892, Accord-431, 434, Sabatier and Senderens, lc.) ing to Gautz (Compt rend 1892, 114, 115), at 400° carbon monoxide reacts with manganese and iron, thus. CO+Mn=MnO+C. Carbon monoxide passes through cast and wrought iron when heated to redness, and, according to Sieverts and Krumbhaai (Ber. 1910, 43, 893), it dissolves in nickel and cobalt, but not in copper

Minute quantities of carbon monoxide reduce gold chloride, and also gold, silver, and mercury oxides (Fay and Secker, J. Amer Chem Soc. 1903, 25, 641) Brunck (Zeitsch angew Chem. 1912, 25, 2479) has proposed to employ its reducing action on palladium chloride (sodiumpalladous chloride) as a method of estimating

small quantities of carbon monoxide

When an electric spark is sent through a mixture of ammonia and carbon monovide, ammonium cyanate is chiefly formed (Northall Laurie, Chem Soc Trans 1905, 433)

Pure carbon monoxide forms a colourless transparent liquid under 200-300 atmospheres at-139°, and solidifies to a snowy mass in vacuo at  $-211^{\circ}$  (Olszewski, Compt rend. 99, 706; 100, 350, Wroblewski and Olszewski, Ann Chim

Phys [6] 1, 112)

Carbon monoxide combines with potassium at about 80° to form an explosive compound, KCO (Brodie, Chem Soc Trans 1860, 12, 269). It is rapidly absorbed by a solution of cuprous chloride in ammonia or hydrochloric acid, by which means it may be directly estimated in a gaseous mixture (cf Thomas, Chem News, 37, 6). According to Manchot and Brandt (Annalen, 1909, 370, 286), it forms the compound CuCl CO,2H<sub>2</sub>O, with copper chloride, and the 2H<sub>2</sub>O can be replaced by 2NH<sub>3</sub> It is also absorbed by moist silver oxide. It unites directly with chlorine, forming phosgene gas (Schutzenberger, Zeitsch f Chem [2] 4, 321) It is absorbed by heated potassium hydroxide at about 200°, forming potassium formate (Berthelot, Ann. Chim. Phys [3] 61, 463) Frohlich and Geuther (Annalen, 202, 317) recommend passing carbon monoxide over soda-lime for this purpose. combines with sulphur to form carbon oxysulphide; with platinum tetrachloride to form C<sub>3</sub>O<sub>3</sub>PtCl<sub>4</sub> and C<sub>2</sub>O<sub>2</sub>PtCl<sub>2</sub> (Schützenberger, Ann. Chim. Phys. [4] 21, 430). It is rapidly absorbed

by anhydrous hydrocyanic acid cooled by a freezing mixture, two layers being formed, but the gas is evolved on removing the tube from the mixture (Bottinger, Ber. 10, 1122). It acts on metallic alcoholates with formation of acids by synthesis (Geuther, Annalen, 202, 288; Schroeder. ibid 221, 34)

Carbon monoxide has been used in electrolytic cells by Haber and Moser (Zeitsch Elektro-

chem. 1905, 11 593).

Carbon dioxide, Carbonic anhydride, Car bonic acid. Gaseous.—Sp.gr. 1 52909 (Rayleigh); 1 52894 (Leduc) V.D. 22 42 at 800°, 21 2 at 1180° (Meyer and Goldschmidt, Ber. 1882, 15, 1165).

Specific heat (const. vol.), 0.33 (equal vol. air=1); 0 2169 (equal wt air=1) (Regnault, Compt rend 36, 676, etc., Wiedemann, Pogg Ann 157, 24). Ratio of specific heat at const press to specific heat at constant volume =129-1:305 (Amagat and Rontgen, Compt rend 71, 336; 77, 1325) ('oefficient of expansion=0 0037 (Regnault, Magnus, and Joly) Coefficient of ex- $PV/P_1V_1 = 1\ 00722$  (Regnault, ibid 20, 975); at 200° carbon dioxide obeys Boyle's Law (Amagat, ibid 68, 1170, 73, 183)

 $u_{\rm C} = 1\,000395$ ,  $u_{\rm E} = 1\,000356$ ,  $u_{\rm G} = 1\,000496$ (Croullebois, Ann. Chim Phys [4] 20, 136, Chapman and Rivière, Compt rend. 103, 37). Heat of formation (const. press.)  $(C+O_2)$ =96 96; (CO+O)=67.96 heat of formation (const vol)  $(C + O_2) = 96.96$ ; (CO + O) = 67.67;  $\begin{array}{l} \text{(CHO}_2 + \text{Aq}) = 102.84 \; ; \; \; \text{(CO} + \text{O} + \text{Aq}) = 73.84 \; ; \\ \text{(CO}_2 + \text{Aq}) = 5.88 \; ; \; \text{(CO}_2 \text{Aq} \; + \; n \text{NaOHAq}) \\ = 11.016 \; (n=1) \; , \; \; 20.184 \; (n=2) \; ; \; \; 20.592 \; (n=4) \\ \end{array}$ 

Solubility of CO, in water (Bunsen, Annalen, 93, 1) -

at 0°-1 7967	at $7^{\circ} = 13339$	at $14^{\circ} = 10321$
$1^{\circ} = 17207$	$8^{\circ} = 1.2809$	$15^{\circ} = 1\ 0020$
$2^{\circ} = 16481$	$9^{\circ} = 1\ 2311$	$16^{\circ} = 0.9753$
$3^{\circ} = 1.5787$	$10^{\circ} - 1\ 1847$	$17^{\circ} = 0.9519$
$4^{\circ} = 15126$	$11^{\circ} = 1 \ 1416$	$18^{\circ} - 0.9318$
$5^{\circ} = 14497$	$12^{\circ} = 1\ 1018$	$19^{\circ} = 0.9150$
$6^{\circ} = 13901$	$13^{\circ} = 10653$	$20^{\circ} = 0.9014$

whence absorption coefficient

```
=1.7967-0.07761t+0.0016424t^2
```

For influence of non-electrolytes on the solubility of carbon dioxide in water, see Usher (Chem Šoc Trans 1910, 97, 66)

Solubility of CO2 in alcohol -

```
at 3.2^{\circ} = 4.0442
                                  at 14 3°=3 3257
   68^{\circ} = 37374
                                     18^{\circ} = 3.0391
                                     22.6^{\circ} = 2.8277
  10.4^{\circ} = 3.4875
```

whence absorption coefficient

$$=4.32955-0.09395t+0.00124t^2$$

For the solubility of carbon dioxide in beer, see Langer and Schultze (Zeitsch für des ges. Brauwesen, 1879, 2, 369); Emslander and Freundlich (Zeitsch physikal. Chem 1904, 49, 317); Findlay and Spen (Chem Soc Trans 1911, 1313).

Liquid — Sp.gr. 1.057 at  $-34^{\circ}$ ; 0.966 at  $-11^{\circ}$ ; 0.84 at  $+11^{\circ}$ ; 0.726 at +22 2° (Cailletet and Mathias, Compt. rend. 102, 1202). efficient of expansion very large, 120 vols. at -20° become 150 vols. at +30° (Thilorier, Ann. Chim. Phys 60, ii 427).

Critical temperature 30.9° (Andrews, Phil.

Trans. 1869, 575), 31 35° with a corresponding Trans. 1869, 575), 31 35° with a corresponding pressure of 72·0 atmos (Amagat) 31 9° and a pressure of 77 atmos. (Dewar). Vap press. in atmos (Regnault) 17·1 at  $-25^\circ$ , 30 9 at  $-5^\circ$ ; 35 4 at 0°; 40 5 at  $+5^\circ$ , 52 2 at 15°; 66 at 25. Bp  $-78\cdot2^\circ$  at atmos press (Regnault). Solid—Sp gr slightly under 1 2 (Landolt, Ber 1884, 17, 309) Vap press in atmos. (Faraday) 5 33 at  $-57^\circ$ , 2 2 at  $-70.5^\circ$ ; 1 14 at  $-99\cdot4^\circ$ . When solid CO<sub>2</sub> is exposed to the air, the temperature remains constant at  $-78^\circ$  to

the temperature remains constant at -78° to -79° (Villard and Jarry; Pouillet, Reg-

nault)

Carbon dioxide was known to Paracelsus and Van Helmont, and was carefully studied by Its true composition was first demonstrated by Lavoisier in 1775 It was liquefied by Davy and solidified by Thilorier It is formed by the combustion of carbon in oxygen or air (Dixon, Chem Soc Proc 1899, 118, Nauwaan, Zeitsch, angew Chem 1896, 200) According to Potter (Proc Roy Soc. 1908, 80, B, 239), amorphous carbon, such as charcoal, lampblack, coal, &c., when exposed to air, is slowly oxidised by the agency of bacteria, to carbon dioxide, the amount of which increases with rise in temperature The temperature at which carbon will form carbon dioxide and monoxide, when heated in oxygen, depends largely on whether the temperature has been reached by heating or cooling (Manville, Compt rend 1906, 142, 1190) The presence of moisture is necessary for the combustion of carbon in oxygen (Brereton Baker, Chem Soc. Trans 1885, 349, Rhead and Wheeler, Chem. Soc Trans. 1912, 101, 846; *ibid* 1913, 103, 1210) It is a constant product of ordinary processes of combustion. It is also formed by the respiration of animals, in various processes of fermentation, and by the decay of animal and vegetable substances It is evolved from fissures in the ground in volcanic districts, and from the craters of active volcanoes, and exists in solution in natural waters, some of which contain it in such quantities as to effervesce is found also in mines, quarries, wells, and caverns, particularly in limestone districts is a constant constituent of the atmosphere, which contains on the average about 0 034 p.c. The air in streets often contains as much as 0 05 to 009 pc In crowded rooms it may reach 03 pc. It is further produced by the decomposition of carbonates either by the action of heat or of stronger acids, and is frequently formed when organic bodies are subjected to high temperatures In the reduction of many metallic oxides by carbon by burning carbon monoxide; by heating together carbon monoxide and steam; and by the action of steam on calcium carbonate at red heat, by heating a mixture of potassium dichromate and sodium carbonate.

Best prepared by acting on chalk, marble, or magnesite with dilute hydrochloric acid solution. The gas may be collected over water or by displacement of air On the large scale it is obtained by heating chalk or limestone to redness in iron or earthen vessels (in lime burning) or by burning charcoal. It is manufactured also as a byproduct in the combustion of fuel. Also pure on large scale by heating carbonates with steam (J. Soc. Chem. Ind. 3, 568).

According to Thom and Pryor (ibid. 1909,

1087), pure carbon dioxide can be manufactured from limestone by the following process. The limestone is heated in closed retorts around a central combustion chamber, beneath which is a furnace, the whole apparatus being enclosed in a casting which excludes an from the retorts Superheated steam is admitted to the retorts at their lower ends, and this decomposes the limestone and carries the carbon dioxide through outlets at the top of each retort, leading to a collecting gas main running round the top of the The retorts have closely fitting mouthpieces, at which they are charged and discharged respectively. By a system of air and waste-gas flues, perfect combustion and the fullest use of heat from the waste gases is ensured (Eng. Pats. 20102, 24332, 1908).

Carbon dioxide is obtained commercially from natural waters from the Saratoga springs in New York, and also in South Germany North Germany the gas is obtained by the combustion of coke, which is generally considered to be the cheapest and best commercial method of producing carbon dioxide Coke is burnt in a generator, A (Fig. 14), the products pass into a combustion chamber, B, into which hot air is

it leaves, almost saturated, by the overflow siphon and accumulates in the tank L. The pump J then drives it through the tubular vessels M and o (where it is heated to 60° and 90° respectively) into the heater c, where its temperature is raised to 100°. The 'poor ley' circulates in the opposite direction back to the tank K. The carbon dioxide evolved from c passes into a cooler, then to a gasholder (Schwatolla, Zeitsch. angew Chem 1900, 1284)

Other methods of obtaining carbon dioxide from coke are described by Candia and Merlini (J Soe Chem Ind 1908, 847), Leshe (ibid. 1905, 240), Gravillon (ibid 1900, 164, Eng Pat.

8377, 1904), and many others


pend upon its absorption

by potassium or sodium

carbonate solution, from

which it is then boiled

Numerous patents also exist for obtaining carbon dioxide from furnace gases and from lime kilns, most of which de-



admitted to complete the combustion, thus oxidising all the carbon monoxide and hydrogen sulphide. The hot gases now pass through tubes contained in the vessel c, thus heating and decomposing the bicarbonate solution The gas now passes contained in this vessel through one or more wash towers, D, containing limestone over which water trickles, and is freed here from dust and sulphur dioxide E 18 a blower or fan, which sucks the gas through the furnace and drives it through the absorbing apparatus. F is a rectangular vessel filled with potash ley, and provided with a number of semicircular baffle plates, so airanged that the gas accumulates under each successively, thus remaining exposed for a considerable time under the pressure of the column of fluid to a large surface of liquid The absorption is further helped by an agitator, which throws the liquid into the upper portion of the vessel the gas passes into the tower c, where the absorption is practically completed, and then leaves the apparatus. The circulation of the potash lev starts from the tank K, whence it is pumped by H through the Korting's jets into the upper part of G, which is thus filled with fine spray. The siphon at the bottom of the tower delivers the partially saturated ley into F, which

compressed, after which they are cooled in a heat interchanger, from which they are led to a drying vessel, containing calcium chloride, into a refugerator, in which the carbon dioxide is condensed to snow, the other gases, still above their critical temperature, are passed back through the interchanger, whilst the carbon dioxide is liquefied and drawn off into tubes and cylinders (Eng. Pat. 11902, 1906)

A method of obtaining steribsed carbon

dioxide, with disinfecting properties, is described by Bouchaud-Praceiq (J Soc. Chem. Ind 1908, 994)

Many attempts have been made to collect the earbon dioxide given off during fermentation processes, and some of these have been successful from a commercial and industrial

point of view

Fig. 14

Mixtures of a carbonate or bicarbonate with a salt or double salt, which is readily hydrolysed in aqueous solutions, have been prepared for the generation of carbon dioxide (Menter and Sedhtzky, J. Soc Chem Ind 1908, 1151; Eng. Pat 14863, 1908).

Ageron and Remy (J. Soc. Chem. Ind. 1906, 847) have prepared a compound by heating a mixture of magnesium sulphate and sodium sulphate or bisulphate with sulphuric acid,

which they use as a substitute for acids in liberating carbon dioxide from carbonates.

Carbon dioxide is a colourless, inodorous gas. neither combustible nor a supporter of combustion. Strongly heated potassium or sodium and brightly burning magnesium burn in carbon dioxide

When quite dry it has no action upon litmus, but if moisture be present the litmus is coloured wine-red. The colour disappears on exposure to air, owing to escape of the gas. Passed into lime-water, carbon dioxide renders it turbid, owing to the formation of calcium carbonate, but if the gas be in excess, the neutral carbonate is converted into an acid carbonate, and the liquid becomes clear.

Carbon dioxide is rapidly absorbed by a solu-

tion of caustic potash

Between -85° and +54° absolutely dry carbon dioxide does not react with absolutely dry alkalı hydroxides. At or above 54°, the dry substances do react, and at 85° a trace of moisture at once starts the reaction, which is completed very rapidly (Moissan, Compt rend. 1903, 136, 723) It is also absorbed by sulphuric acid [100a = 926], where a = Bunsen's absorption coefficient (Bohr, Chem Zentr 1910, n. 1414).

Its solution in water has a sp gr. 1 0018 It has a slightly acid taste, colours litmus winered, and partially neutralises alkalis and dissolves carbonates of barium, strontium, calcium, and

magnesium, &c.

The solution corrodes iron Krohake (J. Soc Chem Ind 1900, 520) found that water containing about 40 mgrms. of CO<sub>2</sub> per litre reduced iron pipes in a short time from 26 mm to 7 mm diameter, owing to the formation of a brown crust, the pipe itself being corroded to a depth of 1-3 5 mm

Its solution in water probably contains carbonic acid  $\rm H_2CO_3$ . At ordinary temperatures, and under I atmosphere, the mass of carbon dioxide dissolved increases as the pressure. But at pressures of 2, 3, or more atmospheres, the mass of carbon dioxide dissolved is less than that calculated by Dalton and Henry's law (v Khanikoff and Louguinine, Ann Chim Phys [4] 11, 412). By the action of carbon dioxide under pressure on water at low temperatures, the crystalline hydrates CO<sub>2</sub>,6H<sub>2</sub>O (Villard, Compt. rend. 1894, 119, 368), CO<sub>2</sub>,8H<sub>2</sub>O, and CO<sub>2</sub>,9H<sub>2</sub>O (Hempel and Seidel, Ber 1898, 31, 2997) have been obtained

Water which has been saturated with carbon dioxide under pressure gives it up as soon as the pressure is removed This property is made use of in the manufacture of effervescing drinks and

aërated waters (v AERATED WATERS)

Under the same pressure the volume of gas absorbed by water diminishes as the temperature rises, the whole of the gas being expelled at boiling heat. Hence carbonic acid water, hold ing an earthy carbonate in solution, deposits it when the water is boiled This is the cause of the furring of kettles, boilers, &c, in which spring or river waters have been boiled.

Potassium burns in carbon dioxide at a red heat with a red incandescence, depositing charcoal mixed with potassium carbonate Sodium decomposes it in a similar manner, as do phosphorus and boron in presence of an alkalı.

It is decomposed by the electric spark into

carbon monoxide and oxygen, if hydrogen, or mercury, or some other metal is present to combine with the oxygen, otherwise the carbon monoxide and oxygen recombine to form carbon dioxide (Collie, Chem. Soc. Trans. 1901, 1063).

Partly decomposed by electric sparks, a condition of equilibrium is attained when change of CO, into CO+O equals that of CO+O into CO2 (Dixon and Lowe, Chem Soc Trans 47, 571).

Carbon dioxide is decomposed by the silent electric discharge, the amount of decomposition varying with the degree of dryness of the gas, the intensity of the discharge, and the form of ozoniser employed, but in all cases a diminished pressure of the gas results in a greater decomposition (Holt, Chem Soc Proc 1908, 24, 271).

Ultra-violet light decomposes dry carbon dioxide at the ordinary temperature into carbon monoride and oxygen, similar results are obtained by the action of radium emanations (Herschfinkel, Compt rend 1909, 149, 395; Ramsbottom, Chadwick, and Chapman, Chem. Soc Proc 1906, 23)

It is partly changed to carbon monoxide and oxygen by heating to 1300° in a porcelain tube (Deville, Compt rend 56, 729, also Berthelot, ibid 68, 1035)

When heated at 350° with hydrogen over reduced nickel or cobalt, it is completely reduced to methane (Sabatier and Senderens, *ibid.* 1902,

With carbon at 500°-650°, it forms carbon monoxide, equilibrium being reached when the mixture of gases contains 61 p c unreduced  $CO_2$  (Boudouard, *ibid* 1899, 128, 824, 1524; 1905, 141, 252)

Mixed with hydrogen and heated to bright redness, or submitted to induction sparks, carbon monoxide and water are formed, if the water is removed, the whole is converted to carbon monoxide (Dixon, Chem. Soc. Trans. 49, 94, Gautier, Bull. Soc. chim. 1906, 35, 929). A mixture of carbon dioxide and hydrogen passed over red-hot pumice yields carbon and water (Dubiunfaut, Compt, rend. 74, 125).

Mixed with sulphur-vapour and passed through a red-hot tube it gives carbon monoxide, sulphur dioxide, and a little carbon oxysulphide (Berthelot, Bull. Soc chim. [2] 40, 362) Passed through a red-hot tube with sulphuretted hydrogen, it forms carbon monoxide, water, and sulphur (Kohler, ibid. 11, 205) Reduced to carbon monoxide by hydrogen, iron, and zinc or copper which has occluded hydrogen (Tissandier, Compt rend 74, 531; Schrotter, Wied Ann. 34, 27); by potassium cyanide (Eiloart, Chem. News, 54, 88), by protoxides of iron and tin (Wagner, Zeitsch anal Chem. 1879, 559); and partially by ferrous sulphate, and a little water in a closed tube (Horsford, Ber 6, 1390).

Carbon dioxide is reduced to carbon by heating with potassium, sodium, magnesium, or silicon (Schmole, J. Soc. Chem. Ind. 1895, 1060). Alkaline carbonates heated strongly with phosphorus or boron give carbon dioxide, which is reduced to carbon (Tennant, Crelles Annalen, [1793] 1, 158; Dragendorff, J. 1861, 111; Leeds, Bull. Soc. Chim. 12, 1834 and 2131); decomposes moist potassium iodide at high temperatures, yielding hydriodic acid (Papasogli, Gazz. chm. ital. 1881, 227); it also decomposes alkaline aluminates (Ditte, Compt.

rend. 1893, 116, 386); and, according to Moore, liberates nitrous acid from nitrites (J. Amer. Chem. Soc. 1904, 26, 959). Solution of carbon dioxide in water yields sodium formate with metallic sodium (Kolbe and Schmidt, Annalen, 119, 251) Carbon dioxide is assimilated by the chlorophyll of plants under the influence of sunlight, carbon being abstracted and oxygen set free. Carbon dioxide diffuses through caoutchouc (Kobbe, Chem. Zot. 1890, 14, 1142)

The specific heat of carbon dioxide is greater at high than at ordinary temperatures (Berthelot and Vieille, Compt. rend. 98, 770 and 852). According to Hoppe-Seyler (Zertsch. physikal Chem. 10, 201, and 10, 401), the carbonic acid and marsh gas found in water-logged soils is due to a widespread process of fermentation of

cellulose

In any working- or living-room the carbon dioxide should not be more than 10 volumes per 10,000 of an.

To estimate the carbon droxide in air, a measured sample of the air is passed through a known quantity of barium hydroxide, coloured with phenolphthalein, and by noting the quantity of air needed to decolourise the hydroxide the carbon droxide can be calculated. The method is sometimes modified by using a limited volume of air, and titrating the hydroxide remaining unneutralised (Ballo, Ber. 1884, 1097; Wiener, Monatsh. 1894, 15, 429, Henriet, Compt. rend. 1896, 123, 125; Woodman, J. Amer. (hem. Soc. 1903, 25, 150., Jean, Compt. rend. 1902, 135, 746., J. Pharm. Chim. 1903, 27, 418., Pécoul and Levy, J. Soc. Chem. Ind. 1906, 655., Davies and McLellan, 454d. 1909, 232)

Carbon dioxide is used extensively in the arts for the manufacture of aerated waters, in sugar manufacture for separating lime from the juice of the sugar-cane, in bread making, and for raising

and clarifying beer.

It is also employed for protecting wines from moulds and acetifying organisms. It preserves their aroma and bouquet, and makes the wine fresher, sweeter, and stronger. Wines can also be revivified by the gas (Piaz, J Soc Chem Ind. 1894, 266, 1903, 644, Pini, Bull de l'Assoc. des Chim 1898, 15, 741) It also possesses antisoptic properties and retards the putrefaction of meat (Kolbe, J pr Chem [2] 28, 61) It has also been proposed to use it as a motive power where fuel is expensive (Herbert, Chem. Zentr. 1885, 543, 558, 572)

Mixed with acetylene (about 5-8 pc.), it decreases the smokiness of the flame and prevents the clogging of the burners (J. Gas Lighting,

1898, 72, 916).

Liquid carbon dioxide (Buchner, Zeitsch physikal Chem. 1906, 54, 665) can be obtained in large quantities by an apparatus described by Thilorier (Annalen, 30, 122) Natteier compressed carbon dioxide by a specially constructed air-pump (J pr Chem 35, 169), v also Goie, Phil Trans 1861, 63) It is prepared from bisulphates by the action of carbonates, the apparatus consisting essentially of a leaden vessel containing a solution of the bisulphate and having a stirring apparatus and gear fixed air-tight to it. By means of a tube and pump an equivalent quantity of calcium carbonate (obtained as a waste roduct in the manufacture of caustic soda) suspended in water in a second vessel is forced

in and the stirrer set in motion The liberated carbon dioxide is dried and passed into a gasometer from which it is afterwards condensed. An apparatus is also described which allows the evaporation and expansion of liquid carbon dioxide to take place round a tube containing a solution of calcium chloride. The latter is so cooled that it may be used for the manufacture The carbon dioxide thus used is passed over moist sodium carbonate, converting it into bicarbonate, which may be again used in the carbonic acid manufactory (Thomas, Zeitsch. angew. Chem 1900, 386) (For description and figure of apparatus for hquifying CO2 and for bottles in which it is stored and sold, see Sisson, J Soc. Chem Ind 1904, 242)

Liquid carbon dioxide may be bought in monor steel bottles containing 8 kilos. (about 4000 litres of the gas at ordinary temperature and pressure), costing about 1 shiper kilo. It furnishes the cheapest means for aerating waters, and is used for raising sunken ships, for driving torpedoes, for extinguishing fires, for cooling purposes, and, as 'Pictet's fluid,' for freezing machines, and by the firm of Krupp, in Essen, for the condensation of steel and other metals, and is further suggested as a motive power for traincars and balloons (J. Soc. Chem.

Ind 4, 610)

It is also used for making carbonates and other chemical products, for filtering and sterilising organic liquids (d'Arsonval, Compt. rend. 1891, 112, 667), and in Germany for the generation of pressure on storage casks of beer to the faucets above where the beer is drawn for consumption. In Engineering, Dec. 3, 1903, a description is given of its use in railway signalling in outlying stations, to supply the motive power for working the signals, so as to avoid the expense of compression machinery or of long lengths of connecting pipes.

Liquid carbon dioxide is colourless, very soluble in alcohol, ether, and volatile oils, but does not mix with water. When the pressure is suddenly relieved, part of the carbon dioxide immediately vaporises, producing sufficient cold to solidify the remainder. Landolt allows the liquid to evaporate freely into woollen bags, and compresses the solid carbon ploxide in coincal wooden moulds by wooden pistons (Ber 17, 309), of Purcell (J. Soc Chem Ind 1892, 936, Eng Pat 13684, 1891). A vessel for storing solid carbon dioxide is described by Heyl and Wultze (J. Soc. Chem Ind 1904, 1215, Eng Pat. 344957, 1904)

An apparatus for obtaining small and large quantities of solid carbon dioxide is described by Fechlu (J pr. Chem. 1903, 67, 423, J Soc. Chem. Ind. 1903, 626), also by Elsworthy (J. Soc. Chem. Ind. 1905, 1231). (For an apparatus for collecting solid carbon dioxide, v. Ducietet,

Compt rend 99, 235.)

Solid carbon dioxide is a white, flocculent, snow-like mass, melting at -567° at 5 atmospheres pressure, and may be left exposed to the air for some time without sensible evaporation. An air or spirit thermometer immersed in it sinks to -79°, it can, however, be placed on the hand without any acute sensation of cold. By mixing with ether its refrigerating power is greatly increased. This is denied by Villard and Jerry (l.c.) but the temperature sinks to

-85° when it is mixed with chloroform. The temperature can readily be reduced to  $-90^{\circ}$  by passing a current of air through the mixture. 5 mm pressure the temperature of solid carbon diovide soon falls to  $-125^{\circ}$ .

Magnesium powder burnt in a dish of solid CO<sub>2</sub> deposits carbon (Brunner, Ber 1905, 38,

1432).

#### COMPOUNDS OF CARBON

Carbon tetrabromide or Tetrabrommethane CBr4 White lustrous crystals of a characteristic pungent smell; m.p.  $91^{\circ}$ ; b.p.  $189.5^{\circ}$  with slight decomposition; b.p.  $101^{\circ}$  at 50 mm without decomposition Occurs in two enantiotropic forms (Rothmund, Zeitsch physikal. Chem 1899, 664). Insoluble in water, soluble Best obin alcohol, ether, and chloroform tained by action of bromine on carbon disulphide in presence of iodine (Bolas and Groves, Chem Soc Trans. 1870, 161, 1871, 773)

Carbon tribromide or Hexabromide, Tetrabrommethylene dibromide C2Br6 Small rect angular prisms obtained by action of bromine on ethylene dibromide, or by heating C2HBr5 with bromine and water at 170° Soluble in carbon disulphide; insoluble in alcohol and ether Decomposed at 200° into C<sub>2</sub>Br<sub>4</sub> and Br<sub>2</sub> (Reboul,

Annalen, 124, 271)

Carbon dibromide, Tetrabrommethylene C, Bi White crystals; mp 53°; formed by action of nascent hydrogen on C<sub>2</sub>Br<sub>6</sub> (Lowig, Annalen, 3, 292, Lennox, Chem Soc Trans 1862, 14, 209)

Carbon bromide, Dibromacetylidene C CBr2, b p 76°. Prepared from tetrabromoethane by removal of two molecules of hydrobromic acid Spontaneously inflammable in air and very explosive (Lemoult, Bull Soc chim 1905, 43, It contains a double bond (Lawrie, Amer Chem. J 1906, 36, 487)

Chlortribrommethane (CIBr, Colourless plates, mp 55°, bp 160° Prepared by heating chloroform with bromine at 250°, and retaining fraction distilling at 160° (Besson,

Compt rend 1892, 114, 223

Dichlordibrommethane CCl. Br. Fine needles, m p 22°, b p 135° Obtained from chloroform by action of bromine as above, retaining the fraction of the product boiling at about 135° (Besson, lc).

Trichlorbrommethane CCl<sub>3</sub>Br, mp b.p. 104°. From chloroform and bromine (Besson, lc) From bromine and trichloracetic acid

(J. H van't Hoff, Bet. 1877, 678)

Chlorotribromethylene CCIBr ('Br<sub>2</sub>; mp  $34^{\circ}$ , bp  $203^{\circ}$ – $205^{\circ}$  at 734 mm (Denzel,  $\psi_{ij}$ ) 1879, 2208)

aa-dichloro- $\beta\beta$ -dibromoethylene  $CCl_2 \cdot CBl_2$ , b p 194° (Bourgoin, Bull Soc chim 1875, 116, and Denzel, Annalen, 1879, 208).

 $\alpha\beta$ -dichloro- $\alpha\beta$ -dibromoethylene CCIBr CCIBr, b.p. 172° at 765 mm. (Swarts, Chem Zentr 1899,

i. 588).

Trichlorobromoethylene CCl<sub>2</sub> CClBr, mp -12° to -13°, bp 145°-148° (Besson, Compt rend. 1894, 119, 88)

Chloropentabromoethane C<sub>2</sub>ClBr<sub>5</sub>, m p. 170°

Denzel, Ber 1879, 2207)

#### aa-dichloro- $a\beta\beta\beta$ -tetrabromoethane CCl, Br CBr,

Colourless crystals; m.p 180° (Denzel, Ber 1879, 2207).

# $\alpha\beta$ -dichloro- $\alpha\alpha\beta\beta$ -tetrabromoethane CClBr. CClBr.

m p 191° (Swarts Chem. Zentr. 1899, [1] 588)  $\alpha\beta\beta$ -trichloro- $\alpha\alpha\beta$ -tribromoethane

CClBr, CCl, Br

m p. 178°-180° (Besson, Compt. rend 1894, 88).  $\alpha\beta\beta\beta$ -tetrachloro- $\alpha\alpha$ -dibromoethane

CClBr<sub>2</sub>·CCl<sub>3</sub>

Prisms from alcohol, smells like camphor. May be sublimed (Paternò, Gazz chim ital. I, 593, Bourgoin, Bull Soc. chim [2] 23, 4, Gossner, Chem Zentr 1903, n 1053)

 $aa\beta\beta$ -tetrachloro- $a\beta$ -dibromoethane

CCl<sub>2</sub>Br·CCl<sub>2</sub>Br

Tables from alcohol (Malaguti, Ann Chim Phys.

1846, 24, Gossner, Chem Zentr 1903, n. 1053).

Fluorotribromomethane CBr<sub>3</sub>F, a colourless, heavy, pleasant-smelling liquid, b p 107°/760 mm, obtained by heating carbon tetrabromide at 50°-60° with silver fluoride

Difluorodibromomethane CBr<sub>2</sub>F<sub>2</sub>, a mobile hquid, b p. 24  $5^{\circ}/760$  mm, mp below  $-80^{\circ}$ formed by the action of the preceding compound on silver fluoride (Rathsburg, Ber 1918, 51,

Carbon tetrachloride or Tetrachlormethane CCl<sub>4</sub> was first obtained by Regnault (Annalen, 1840, [33] 332), who prepared it by the action of chlorine on chlorform in sunshine

 $\mathrm{CHCl_3} + \mathrm{Cl_2} - \mathrm{HCl} + \mathrm{CCl_4}$ 

Dumas afterwards obtained it by the action of chlorine upon marsh gas (Annalen, 1840, [33] 187) It can also be prepared by passing carbon disulphide saturated with chlorine through a The mixture of carbon tetrared-hot tube chloride and sulphur chloride thus obtained is treated with potash or milk of lime and the tetrachloride distilled off (Kolbe, Annalen, 1843, 45, 41, 1845, 51, 146) The tetrachloride may contain carbon disulphide if that substance was in excess, or if the heat was insufficient may be removed by leaving the liquid for some time in contact with potash Geuther removes the carbon disulphide by converting it into potassium xanthate by dissolving the mixture in alcohol, adding alcoholic potash so long as the liquid becomes darker in colour, and heating gently, the unaltered tetrachloride is separated by water and purified by washing (Geuther, Annalen, 1858, 107, 212, and Selmitz-Dumont, Chem Zentr. 1897, n 265).

The complete chlorination of carbon disulphide and of many other carbon compounds can be more readily accomplished in the presence of a chlorine carrier Thus, in the case of carbon disulphide, bromine has been used (Crump, Chem News, 1886, 14, 154) and nodine, both alone (Morel, Compt. rend. 1877, 84, 1460) and in the presence of metallic iron (Serra, Chem. Xentr. 1899, 11. 1098) · antimony pentachloride reacts with carbon disulphide alone, without the addition of chlorine. The mixture becomes hot, and on cooling deposits crystals of antimony trichloride mixed with sulphur, whilst carbon tetrachloride remains in solution (Hofmann, Annalen, 1860, 115, 264)

A large number of other substances can be used as catalytic agents, e.g. molybdenum pentachloride (Aronheim, Ber. 1876, 1788), the chlorides of phosphorus and arsenic, and also a

series of chlorides of metals, especially those of the iron group. In the presence of these substances, sulphur chloride reacts readily with carbon and excess of chlorine, giving carbon tetrachloride A stream of chlorine is led over a red-hot mixture of charcoal and sulphur Carbon disulphide is first formed, then carbon tetrachloride and sulphur monochloride, the latter then breaks up into chlorine and sulphur, and the process begins over again The carbon tetrachloride prepared in this way is usually contaminated with sulphur chlorides and chlorethane compounds (Combes, Chem Zentr 1909, 1, 326, J Soc Chem Ind 1902, 272 and 1469) In the presence of manganese chloride the reaction can be carried out with precipitation of sulphur, which can be used to manufacture more carbon disulphide (Côte and Pierron, *ibid* 1903, 962. Febvre, *ibid* 1905, 1189) Other carbon compounds beside carbon disulphide can be completely chlorinated. Thus chloroform heated with iodine chloride at 165° gives carbon tetrachloride (Friedel and Silva, Bull Soc chim 1872, 537), the same reaction taking place, to a slight extent, in the presence of aluminium trichloride instead of iodine chloride (Monneyrat, ibid 1898, 262)

Perchlormethyl formate, when treated with aluminum chloride in the proportion of 50 parts to 1 of chloride, also yields carbon tetrachloride CClO<sub>2</sub> CCl<sub>3</sub>=CCl<sub>4</sub>+CO<sub>2</sub> (Hentschel,

J. pr Chem 1877, 306)

The technical preparation of carbon tetrachloride is generally carried out by the direct chlorination of carbon disulphide in the presence of powdered aluminium chloride, the quantity of catalyst necessary being only about 1 p c of the carbon disulphide used

$$CS_2+3CI_2 = CCI_4+S_2CI_2$$

The sulphur chloride reacts separately with more carbon disulphide in the presence of iron filings, and the crystalline sulphui which is formed is employed again in the manufacture of carbon disulphide (Urbain, Chem Zeit 1902, 1086; Breteau, J Pharm Chim 1908, 110, Urbain, J Soc Chem Ind 1902, 926, Eng Pat 13733)

Later modifications of the process consist in the further chlorination of the sulphur monochloride, in the presence of antimony chloride, before treatment with carbon disulphide (Haworth and Baker, J. Soc. Chem. Ind. 1906, 559), and the use of a solution of sulphur in sulphur monochloride, in a finely divided condition, as a chlorine carrier (Acker and Strue, ibid. 1908, 833). Instead of aluminum chloride, aluminum amalgam has been employed (Castner Co. ibid. 1905, 903). Under this patent, the resulting carbon tetrachloride is purified by treatment with a solution of alkaline sulphide containing free alkali, the carbon disulphide being removed as thiocarbonate (cf. Dow Chemical Co. U.S. Pat. 1204608; J. Soc. Chem. Ind. 1916, 35, 1271)

The Isco Chemical Co (US Pats. 1260621 and 1260622) treat carbon disulphide containing free sulphur with chlorine in presence of a catalyst to form sulphur dichloride and carbon tetrachloride, and an additional quantity of carbon disulphide is then added to convert the sulphur dichloride into sulphur monochloride,

with the formation of more carbon tetrachloride. The product is treated with sufficient tin to convert the sulphur monochloride into sulphur, with the formation of stannic chloride, and after separating the sulphur, water is added to hydrate the stannic chloride, and the carbon tetrachloride is recovered from the mixture by distillation.

A number of electric-heating processes have been suggested A mixture of sodium chloride and silica is heated in the lowest portion of a furnace lined with magnesia, and provided with three sets of electrodes, one above the other The middle electrodes heat a mass of coke, and the chlorine from the alkali chloride reacts with the incandescent carbon, forming carbon tetrachloride, which, after passing through the ares between the third set of electrodes, is distilled off The temperature does not exceed 2000° (Machalske and Lyon, J Soc Chem Ind 1903, 1298, Machalske, Chem Zentr. 1904, 1, 1069) The chlorine can also be supplied from an external source, the carbon only being heated by electricity (Maywald, J Soc Chem Ind 1907, 1253)

Another process consists in heating together hime, calcium chloride, and carbon in the absence of air in an electric furnace, in the proportions indicated by the equation

$$2\text{CaO} + 2\text{CaCl}_2 + 10\text{C} - 4\text{CaC}_2 + 2\text{COCl}_2$$

The carbonyl chloride is led over heated coke, animal charcoal, or pumice, where it is decomposed into carbon tetrachloride and carbon doxide by surface contact (Machalske and Darlington, J Soc Chem Ind 1906, 559)

A mixture of calcium chloride and coke, heated in the electric furnace in a stream of chlorine, also yields carbon tetrachloride (Matthews and Darlington, *ibid* 1906, 1232)

The halogenation of metallic or other carbides, such as acetylene, also yields carbon halides, so long as the temperature is kept below the dissociation point of the halide required (Blackmore, *ibid* 1908, 712)

Carbon tetrachloride is a transparent colourless liquid, with a pungent aromatic odour; mp — 22 7°, bp 76 74°, and spgr 1 63195, 0°/4° (Thorpe, Chem Soc Trans 1880, 201); 76 96° (corr) (Thorpe and Rodger); (Moles, Anal. Fis Quim 1912, 10, 30). Its specific heat increases regularly from 0 2010 at 0° to 0 2031 at 70° The specific heat of the vapour falls from 0 14 at 0° to 0 115 at 70° (Mills and MacRae, J Physical Chem 1911, 15, 54)

The vapour pressure of carbon tetrachloride has been measured by Schreinemakers (Zeitsch physikal Chem 1904, 48, 445). The compressibility has been measured by Richards and Stull (Amer. Chem J. 1904, 26, 408). In the solid state carbon tetrachloride is trimorphous, the three modifications melting at  $-28~6^{\circ}$ ,  $-23.77^{\circ}$ , and  $-21~2^{\circ}$  respectively, the second being the form usually obtained (Tammann, W. Ann. 1898, 480) The refractive index for the line F is 14726, and for the line D is 14658 (Gladstone, Chem Soc Trans 1886, 623).

Its dielectric constant has been measured by Turner (Zeitsch. physikal. Chem. 1900, 385), Drude (Zeitsch. physikal. Chem. 23, 267), and Veley (Phil. Mag. [6] 11, 73). Its solubility in water, which is extremely small,

has been measured by Rex (Zeitsch. physikal. Chem. 55, 355).

It is soluble in alcohol and ether, and dissolves a large number of organic substances, especially those of a fatty nature, whence its wide application in the arts. It is preferable to carbon disulphide on account of its non-inflammability. With sulphonated oils and resins it yields a gelatinous soap, which is miscible with water (Stockhausen, Chem. Zentr. 1906, ii. 731). It yields chlorine and lower chlorides of carbon on passing through a red-hot tube (Regnault, lc). When mixed with hydrogen and passed through a red-hot tube with pumice, it yields marsh gas and ethylene (Berthelot, Ann. Chim. Phys. 1858, 53, 69).

Phys 1858, 53, 69).

The same mixture passed over reduced nickel yields hexachlorethane (carbon truchloride) at 270°. With excess of hydrogen at the same temperature, tetrachlorethylene (carbon dichloride) is obtained (Sabatier and Mailhe, Compt.

rend 1904, 138, 407)

Mixed with sulphuretted hydrogen and passed through a red-hot tube, the vapour of carbon tetrachloride yields hydrochloric acid and carbon sulphochloride CSCl<sub>2</sub> (Kolbe, Annalen, 1843, 45, 41, and 1845, 54, 146)

Heated under pressure with oxygen and water at 250°, it forms carbonyl chloride and carbon dioxide (Goldschmidt, Ber 1881, [14] 928). In alcoholic solution, and treated with potassium amalgam, it yields chloroform, marsh gas, and methylene chloride (Regnault, *l.c.*) Phosphorus pentoxide at 200° acts in two ways in the first, carbonyl chloride and carbon dioxide are produced, in the second, in the presence of more phosphorus pentoxide, carbon dioxide alone is obtained.

$$\begin{array}{c} 2\text{CCl}_4 + \text{P}_2\text{O}_5 = \text{COCl}_2 + \text{CO}_2 + 2\text{POCl}_3 \\ 3\text{CCl}_4 + 2\text{P}_2\text{O}_5 = 3\text{CO}_2 + 4\text{POCl}_3 \end{array}$$

(Gustavson, Zeitsch, f. Chem. 1871, 615).

A similar reaction takes place at 100° with sulphur trioxide, carbonyl chloride being formed

$$CCl_4 + 2SQ_3 = S_2Cl_2Q_5 + COCl_2$$

(Erdmann, Ber. 1893, 26, 1993).

With the oxides of many metals, a similar production of carbonyl chloride, together with carbon droxide and the chloride of the metal, takes place (Demarcay, Compt. rend. 1887, 104, 111; Quantin, *ibid.* 1887, 104, 223, 1888, 106, 1074; Delafontaine and Linebarger, Amer Chem. J. 1896, 532).

With zine and dilute acid, carbon tetrachloride is converted into hydrochloric acid and chloroform (Geuther, \$lc\$). Alcoholic potash slowly converts it into potassium chloride and carbonate (Regnault, \$lc\$). By long-continued action at 100° in sealed tubes, alcoholic potash converts it partially into ethylene (Berthelot, Ann. Chim Phys 1859, 109, 118) With alcoholic potash and aniline it yields phenylsocyanide (cf. Chloroform). With phenylamine it yields diphenylaminobenzamide (Hofmann, Proc. Roy. Soc. 9, 284, and 10, 184, Wieth, Ber. 1877, 10, 358; Michler and Walder, Ber. 1881, 14. 2174), a reaction facilitated in presence of a coppermercury couple (Hartung, Chem. Soc. Trans. 1918, 113, 163). By the action of sulphur at 220°, a number of sulphochlorides of carbon disulphide (Klason, Ber. 1887, 20, 2376). It is

decomposed at 200° by molecular silver, being converted into hexachlorethane  $C_2Cl_6$  (Goldschmidt, Ber. 1881, 14, 927). By heating with aluminium bromide it is readily converted into carbon tetrabromide (Gustavson, Bull. Soc. chim. 1881, 556).

Heated in a sealed tube for 10 hours at 130° with hydrodic acid, carbon tetrachloride yields iodoform and hydrochloric acid (Walfisz, *ibid* 1893, 7, 256) Acetylene reacts with carbon tetrachloride, yielding free carbon in a very finely divided form, hydrochloric acid being formed at the same time (Sandman, Chem Soc Abstr. 1902, 82, 1–581) Carbon tetrachloride reacts with benzene in the presence of aluminium trichloride, giving, as chief product, diphenyldichloromethane, CCl<sub>2</sub>(C<sub>6</sub>H<sub>b</sub>)<sub>2</sub> (Boeseken, Chem Soc Abstr 94, 1, 189) Carbon tetrachloride absorbs ozone, giving a blue solution

In its physiological action, carbon tetrachloride strongly resembles chloroform.

Its popularity as a solvent in the arts has been somewhat diminished by the increasing manufacture of tetrachlorethane C<sub>2</sub>Cl<sub>4</sub>H<sub>2</sub>, as the latter is more indifferent toward metals, and is readily produced by the action of chlorine on a double compound of antimony pentachloride and acetylene (D. R. P. 154657, 1906).

Carbon trichloride, Carbon hexachloride, Tetrachlorethylene dichloride, Hexachlorethane, Perchlorethane C.Cl<sub>8</sub>, discovered by Faraday (Phil. Trans 1821, 47), also investigated by Regnault (Ann Chim Phys 1838, [2] 69, 166, and 1839, [2] 71, 371) Is obtained by the action of chlorine in sunshine upon various derivatives of ethyl and ethylene, on carbon dichloride, on ethylene chloride (Faraday, lc), and also (Liebig, Annalen, 1, 219) on ethyl chloride, first in the shade, afterwards in sunshine (Laurent, Ann. Chim Phys 1837, [2] 64, 328), on mono-, di-, or tri-chlorinated ethyl chloride (Regnault, lc), on ethyl sulphite (Ebelmen and Bouquet, ibid. 1846, [3] 17, 66), on ethyl oxide (Regnault); on hydrochloride of ethylamine (Geuther and Hofacker, Annalen, 1858, 108, 51) (in this process some chloride of nitrogen is produced), by passing the tetrachloride through a red-hot tube (Kolbe, Annalen, 1845, 54, 147), by distilling perchlorethylic oxide, and repeatedly treating the distillate with water (Malaguti, Ann Chim. Phys 1846, [3] 16, 14), by heating carbon tetrachloride with copper powder at 120° (Radziszewski, Ber. 1887, 17, 834) or with molecular silver at 200° (Goldschmidt, ibid. 1881, 14, 928), by heating acetyl chloride with excess of phosphorus pentachloride at 180° (Hubner and Muller, Zeitsch f Chem. 1870, 328); by heating propyl chloride or isobutyl chloride with iodine trichloride at 200° (Krafft and Merz, Ber. 1875, 8, 1298), by treating pentachlorethylchloroformic ester with aluminium chloride (Muller, Annalen, 1890, 258, 63).

Carbon trichloride crystallises from a mixture of alcohol and ether in right rhombic prisms (Gossner, Chem Zentr 1903, ii. 1052). Colour-less, transparent, and nearly tasteless, but having an aromatic odour resembling that of camphor, sp. gr. 20; mp. 184°-186° (Bolton, Zeitsch. Elektrochem. 1902, 8, 165; 1903, 9, 209), volatilising even at ordinary temperatures; b.p. 185° (Hahn, Ber 1878, II, 1735). Insoluble in water; soluble in alcohol and ether,

and in all oils. Converted, by repeated distillation, into the dichloride and free chlorine. Burns with a red light in the flame of a spirit lamp. Mixed with hydrogen and passed through a red-hot tube, it yields dichloride and hydrochloric acid (Geuther, l.c.) Metals heated in its vapour form metallic chlorides, chaicoal being deposited. Heated in sealed tubes with potassium hydroxide, it yields potassium oxalate and chloride

$$C_2Cl_6 + 8KHO = K_2C_2O_4 + 6KCl + 4H_2O$$

(Geuther, J. 1859, 277) The same products together with hydrogen and ethylene are formed when it is heated with alcoholic potash in sealed tubes (Berthelot, Annalen, 1859, 109, 118).

Reducing agents such as zinc and dilute sulphuric acid, or alcoholic potassium hydrogen sulphide (Regnault, lc), reduce it to tetrachlorethylene (carbon dichloride). The same reaction takes place when it is heated with silver at 280°. Sulphur trioxide acts upon it at 150° according to the equation

 $CCl_3 CCl_3 + 2SO_3 = CCl_3 COCl + S_2O_5Cl_2$ (Prudhomme, Annalen, 1870, 156, 342)

Carbon dichloride, Tetrachlorethylene C2(14 (Faraday, Phil Trans 1821, 47, Regnault, Ann Chim Phys 1839, [2] 70, [2] 71, 372), is formed as a liquid coloured with chlorine when the vapour of carbon trichloride is passed through a red-hot tube filled with fragments of glass. It is purified by passing it repeatedly through a red-hot tube, then shaking up with mercury and rectifying at a low temperature (Faraday) More easily obtained by adding carbon trichloride in small portions to an alcoholic solution of hydrogen potassium sulphide, as long as sulphuretted hydrogen is given off. The liquid is distilled and carbon dichloride separates from the distillate on dilution with water (Regnault, l.c); or by reduction of carbon trichloride with zinc and sulphuric acid Mobile liquid of sp gr 1 619 at 20° (Regnault), 1 612 at 10° (Geuther). b p 122° (Regnault), 116° (Geuther), 120 74° (corr.); sp gi 1 6551 0°/4° (Thorpe and Rodger), remains liquid at -18°, and does not conduct electricity Insoluble in water, acids, and alkalis, but dissolves in alcohol and other, and Decomposed at a red heat into the oils hexaclorobenzene C<sub>6</sub>Cl<sub>6</sub> and free chlorine When its vapour is passed over baryta at a red heat, it is decomposed with vivid ignition into barium chloride, carbon dioxide, and charcoal It absorbs bromine in the sunshine, forming carbon chlorol romide C<sub>2</sub>Cl<sub>4</sub>Br<sub>2</sub> It is converted by continued heating to 200° with potassium hydroxide into potassium oxalate and chloride, hydrogen being given off (Geuther, Annalen, 247) It absorbs dry chlorine in sunshine, forming the trichloride, but is converted into trichloracetic acid when exposed to an atmosphere of chlorine under water

$$\mathrm{C_2Cl_4} + 2\mathrm{H_2O} + \mathrm{Cl_2} = 3\mathrm{HCl} + \mathrm{C_2HCl_3O_2}$$

(Kolbe, Annalen, 1845, 54, 181)

as-Heptachloropropane  $C_5Cl_{10}$ , a crystalline substance, mp 32°, formed by the action of phosphorus pentachloride on pentachloroacetone; also obtained by the direct addition of chloroform to tetrachloroethylene under the influence of aluminium chloride (Boeseken and Prins).

Hexchlorobenzene C<sub>6</sub>Cl<sub>6</sub>, m p. 225°; discovered in 1821 by Julin and investigated by Phillips and Faraday (Phil. Trans 1821), and by Regnault (Ann. Chim. Phys. [2] 70, 144), who prepared it by passing the vapour of chloroform or tetrachlorethylene through a porcelam tube filled with fragments of porcelam and heated to redness Obtained from methylene chloride by ICl or ICl<sub>3</sub> (Holand, Annalen, 240, 234), the crystalline product is dissolved in ether, filtered, evaporated to dryness, and sublimed. Obtained by Hugo Muller as a final product of the action of antimony pentachloride or chlorine in presence of rodine on It is also obtained by sending a benzene. current of 5 5 amperes and 42 volts for ½ hour between carbon electrodes in an atmosphere of If a higher current and voltage are used, and it is continued for much longer, the trichloride C2Cl8 is formed Bromine yields similar compounds (Bolton,  $l \epsilon$ .). Forms white delicate needles with a silky lustre, tasteless, but with an odour resembling spermaceti. sublimes without fusion at 120°.

Insoluble in water, acids, and alkalis, but soluble in alcohol, ether, and hot oil of turpentine. Decomposed into chlorine and charcoal on passing through a red-hot porcelain tube filled with glass or rock crystal. Burns with a bluish colour in the flame Potassium burns on being strongly heated in its vapour Formerly regarded as carbon monochloride: molecular formula established by Bassett (Chem Soc. Trans. [2] 5, 443)

Carbon tetraiodide, Tetraiodomethane Cl<sub>4</sub>
Dark-red octahedra, sp gr 4 32, obtained by action of aluminium rodide on a mixture of carbon tetrachloride and disulphide (Gustavson, Ber 14, 1705) For other methods, see Spindler, Annalen, 231, 257, Moissan, Compt. rend 113, 19, Robineau and Rollin, Chem. Centr. 1894, 1 1140, Lantenois, Compt rend. 1913, 156, 1385.

Carbon nitride (v. Cyanogen, art. Cyanides) Carbon subnitride C<sub>4</sub>N<sub>2</sub> has been obtained by Moureu and Bongrand (Compt rend 1910, 150, 225) by removing two molecules of water from the diamide of acetylene dicarboxylic acid CONH<sub>2</sub>C CCONH<sub>2</sub>=N: CC:CC:N+2H<sub>4</sub>O. It forms fine white needles. mp 20 5°-21°, bp 76° at 753 mm, and its vapour has an odour and an irritating action resembling that of cyanogen. It is readily combustible, and takes fire spontaneously at 130°, burning with a puiple flame, also resembling that of cyanogen. It has an abnormally high molecular refraction and dispersion, and a density of 0 9703 at 25°/4° For its behaviour towards ammonia and amines, see Moureu and Bongrand, Compt. rend. 1914, 158, 1092.

Carbon oxychloride COCl<sub>2</sub>, Carbonyl dichloride, Chlorocarbonic acid, Phosgene gas. First obtained by J Davy by action of sunlight on mixture of carbon monoxide and chlorine. (For details of this mode of preparing it, r. Emmeling and Lengyel, Annalen Suppl 7, 101, and Wilm and Wischin, ibid. 147, 450.) By passing the gases over heated animal charcoal combination occurs in the dark. Also obtained by passing carbon monoxide into boiling antimony pentachloride (Hofmann, ibid. 70, 129; Butle row, Zeitsch. f. Chem. 1863, 484; Kraut,

Gm.-K. I 2, 380); by heating carbon tetrachloride with zinc oxide in closed tubes at 200° or by mutual action of carbon tetrachloride and carbon monoxide at 400°, or by heating a mixture of chloroform, potassium dichromate, and sulphuric acid (cf Plotnikow, J. Russ. Phys. Chem Soc 1916, 48, 457)

Carbon oxychloride is a colourless liquid, boiling at 82°, and of sp gr. 14320/4°. Soluble in acetic acid and benzene, decomposed by water  $COCl_2 + H_2O = CO_2 + 2HCl$  With alcohol forms chlorocarbonic ester Cl COOC 2H 5 bines with ammonia to form urea and ammonium chloride With aniline it forms diphenyl-Its presence in air may be detected by this reaction (Kling and Schmutz, Compt. rend.

1919, 168, 773)

Carbonyl cyanide CO(CN)2 is a yellow amorphous non-volatile solid, formed by subjecting a mixture of carbon monoxide and cyanogen to the ultra-violet rays from a mercury-quartz lamp It dissolves in alkalis, forming a yellow solution, and slowly hydrolyses on the addition of acid, giving carbon dioxide and hydrogen cyanide (D. Berthelot and Gaudechon, Compt. rend 1913, 156, 1766)

Carbon disulphide, Throcarbonic anhydride, Sulphocarbonic and CS, This compound was discovered by Lampadius, in 1796, by heating pyrites with charcoal Clement and Desormes, in 1802, in examining the action of sulphur on red-hot charcoal, obtained the same product

Its nature was established by Vauquelin.


Manufacture—Schrotter in Germany, and Peroncel in France, first prepared carbon disulphide on a manufacturing scale Schrotter employed a cylinder of Hessian clay in which the charcoal was heated, the sulphur was introduced through a hole near the bottom, to the upper end was fixed an exit tube for the vapour of the carbon disulphide formed The apparatus produced about 20 kilos of the disulphide in 12 hours

Deiss and Fisher employed cast-iron cylinders in which the vapour of sulphur passed over charcoal heated to redness Fireclay retorts glazed internally to prevent the escape of vapour through the pores were also used, each cylinder was about 5 feet high and nearly 20 inches diameter Four retorts were arranged in a single furnace in such a way that the flame of the fire could play entirely round Each retort was divided internally into two parts by means of a perforated shelf, the upper compartment, which was the larger of the two, served to contain the charcoal, which could be thrown in through a special opening in the lid without interrupting the working. A second opening in the top of the retort was provided with a tube about 2 mehes in diameter, which led directly into the lower compartment, and was used to throw in sulphur as required The vapour of carbon disulphide which was formed during the process was conducted through a delivery pipe connected with a third opening in the retort into the condensers four retorts, having been filled with charcoal and the lids closed, were heated to redness by a coal Pieces of sulphur wrapped in cylindrical paper packets were then dropped in the pipes leading to the lower partitions of the retorts Two such packets, each containing about 51 oz. of sulphur, were thrown in at intervals of

about 3 minutes, the opening at the end of each tube being closed at each successive addi-The charcoal was renewed every 7 hours, and was heated about an hour and three-quarters each time before the necessary temperature was attained for its combination The vapour of carbon disulphide with sulphui mixed with that of sulphur escaped through the delivery tubes, passing into the first row of condensers, where it was partially condensed, together with some sulphur, the undissolved sulphur was recovered and used over again. Thence the uncondensed vapours passed on into the second row of condensers, and so on till the The condensers, amounting to 18 or 20 m all, were connected together by pipes, the last of which was connected with the chimney of the works, or better, with vessels or tubes containing layers of pulverised lime to absorb the sulphuretted hydrogen which would otherwise create a nuisance in the neighbourhood of the factory

The condensers were made of sheet zinc, were cylindrical in shape and about 26 inches in diameter, bottomless, with slotted sides, and stood in a kind of shallow cistern containing water a little deeper than the openings, so as to form a water lute, permitting the passage of the condensed liquid while closing the exit against vapours. The lids of these condensers were furnished with rims forming a kind of saucer, containing water to assist in the condensation of Each hid had two openings in the the vapour form of tubulures, into which the pipes were fixed for the transmission of the vapours from condenser to condenser

Gérard employed a vessel of cast iron 63 feet in height, about 4 feet 8 inches in diameter, and sides 13 inch thick (Fig. 15). The retort was elliptical



in section, stood upon a brickwork arch, and was heated from below by a fire. Near the bottom was a pipe fitted with a valve, through which sulphur could be introduced into the cylinder. At the upper extremity was a wide pipe which could be closed with a lid, and served for the introduction of charcoal into the apparatus. Another pipe, cast in one piece with the upper one, sloped forward and passed into a receiver in which the sulphur which passed over during the

operation was condensed, and could be easily removed by opening a valve in a tubulus placed at a bend in the pipe; the more volatile carbon disulphide passed forward into the condensers. The condenser consisted of three vessels, cylindrical in form, arranged one above the other, and communicating by vertical pipes. The topmost condenser had an exit pipe for sulphuretted hydrogen, &c, and the condensed carbon disulphide could be drawn off into a receiving vessel beneath by opening a cock in a pipe proceeding from the lower condenser.

The three condensess were contained in a tank 5 feet high and 5 feet wide, filled with water,

which could be renewed at pleasure


The sulphur was added in portions of about 3 lbs at intervals of 3 minutes. The oven was heated during the night to volatilise the sulphur, and the residual charcoal was added to the new charge. This apparatus yielded in 24 hours about 570 lbs of carbon disulphide, consisting theoretically of 480 lbs of sulphur and 90 lbs of carbon, in practice 530 lbs of sulphur and 242 lbs of wood charcoal were used. Gerard found it advantageous to surround the generating

vessels with brickwork, which rendered them more durable.

Only retorts of a certain size could be employed, as, when they were too large, the heating was megular, causing waste of sulphur. When using unrefined sulphur, the retorts needed cleaning every fortnight, but with purified sulphur they could be kept working for two months

The present mode of manufacture is thus described by Ignatius Singer (J. Soc. Chem. Ind. 8, 1889, 83) (see Fig. 16).

A is a vertical east-non retoit, or, better still, made of earthenware, glazed inside, of elliptical shape, about 66 inches high, and 20 by 12 inches internal diameter. It rests on a support, B, made of firebricks, and is protected by a mantle of best firebricks, C, about 4 inches thick, leaving a space of from ½ to ¼ inch between it and the retoit. The metal of the retoit should not be less than 2 inches in thickness. D is the outer brickwork of the furnace lined on the miside with firebricks, D', E, E, fire-grates, E', ash pits, F, furnace doors, made of frames, a, into which a lire-proof slab, b, is inserted. The



F10 16.

furnace doors are suspended by cables running over pulleys, and are counterpoised, so that they can be raised or lowered At d is shown a small peep-hole, through which the temperature in the furnace may be watched. In the lid of the retort are cast two tubes, a and a', of about 5 inches internal diameter and not less than 1 inch in thickness, but it is advisable to have them made somewhat stronger, as the lid will then outlast several retorts Over the openings, G, a vent-pipe or flue, H, is suspended by a lever, so that it can be raised or lowered at will, one end of which passes through the roof fresh charges of charcoal are to be introduced into the generator A, which is done through the tubulure a', the lid closing the tube a is first removed, and the vent-pipe H quickly lowered, which serves the purpose of carrying off the noxious gases that would otherwise be injurious to the workmen G' may now be opened, and the carbon fed in without the slightest inconvenience

From the tubulure g' issues a pipe, I, inclined upwards, through which the disulphide of carbon vapours escape. Most of the free sulphur passing over will condense here and run back into the retort. But the partially cooled sulphur

falling on to the surface of the incandescent carbon would be immediately vapourised again, at the same time cooling down the top layer of the charcoal. To obviate this, Singer suggests that an opening should be made in the lower part of the pipe I, to which a hopper can be fitted, as shown at J, terminating in an earthernware pipe K, reaching nearly to the bottom of the retort. By this means the sulphur distilling over uncombined would be conducted back to the bottom of the generator, the pipes would be loss liable to be choked up, while the resulting product would be purer

From the lower end of the retort, close to the bottom, a pipe M branches off, a little upwards inclined, passing through the brickwork and terminating in a chamber N, with door n During work this pipe is carefully closed by a lid. It serves the purpose of raking the ashes out of the retort, which is done once a week. At m a flue is shown, carrying the noxious gases into the chimney. The ashes are allowed to cool here before their removal Adjoining this chamber is a hearth into which an iron vessel, O, is fitted, for melting the sulphur. It is heated by a flue from the furnace passing underneath it, provided with a damper

to regulate the heat. (In Fig. 16 this sulphur vessel is raised so as to make it visible in the drawing) A pipe, o', closed by a conical valve to which an iron rod, o, is attached, connects the sulphur pot with the arm M of the retort, by which means the sulphur can conveniently be

introduced into the latter

The sulphur flowing down the incline of the arm M arrives at the bottom of the retort, where it is volatilised: the vapours, passing upwards, combine with the carbon. The generated disulphide vapours escape through pipe I, and then pass down the vertical tube I' into a vessel P, where any excess of sulphur is deposited. The lid of this vessel rests in hydraulic seals, and is connected with the inlet and outlet pipes by a kind of telescopic tube made air-tight by water. When the vessel P is to be removed, the lid is raised, the vessel taken away, another similar one put in its place, and the hid lowered again. This should be done once a week only, and never while distillation is going on.

The carbon disulphide vapours, now freed from the greater part of uncombined sulphui, pass through a Liebig's condenser, L, about 30 feet long, into a receiver, s, partially filled The crosspiece T, connecting the with water Liebig's condenser with the receiver, is made so that it can be detached—being secured at v by flanges bolted together, and at n1 by water seal-for purposes of cleaning, &c. From here the disulphide is allowed to run through a siphon, p, direct to the storage tanks A better plan, however, is to keep the storage tanks for the crude product on a high level, which greatly facilitates the after-process of purification. this case the disulphide of carbon is allowed to flow into a montejus, v, as shown in drawing, A pipe q, reaching nearly to the bottom, is passed through the cover of the montejus to the high-level tank. A second pipe, r, connects the vessel u with a force-pump On air being pumped into v through r, the stop-cocks  $p^1$  and p<sup>2</sup> having been shut off, the disulphide is pressed up through pipe q into the tank. This is a far better method than using pumps, as in the latter it is difficult to prevent leakage, the ciude disulphide having a very corrosive action on most metals. For the same reason, the receivers, tanks, &c., for the crude substance should all be lined with sheet-lead, as wrought-iron vessels soon perish, while copper is even more energetically acted upon. Cast iron withstands corrosion much better, but is objectionable on account of its porosity This might be remedied, however, by giving the vessels repeated coatings of dilute silicate of soda, both inside and outside, the vessels being first slightly heated with

steam and the silicate applied while warm

Near to the top of the receiver s is a pipe, s, for carrying off the uncondensable vapours—chiefly sulphuretted hydrogen. The gases are conducted into a rectangular vessel, w, where they are made to circulate in zigzag over a series of shallow trays filled with a vegetable oil, to absorb any disulphide vapours which have escaped condensation. The oil can be made to trickle from a reservoir, x, as shown, through a goosenecked pipe, which, passing from tray to tray in an opposite course to that of the gases, finally trickles through x into a receiver Y. When a sufficient quantity has accumulated here, the

disulphide is distilled off and the oil returned to x. Before the gases enter this absorber they are 'washed' by making the end of the pipe s dip into the oil as shown at y whereby the apparatus is disconnected from contact with air. The gases are now passed through a second vessel, z, similar in construction to the one just described, only substituting lime or oxide of iron for the oil, to absorb the  $H_2S$ , and are then allowed to escape

into the open air through the pipe z.

Above the retort A, and between it and the chimney, set at right angles to each other, is a smaller retort, similar in construction to A, but only about one-third the capacity of the latter, and is heated by causing the products of combustion coming from the furnace to circulate round it before finally passing into the chimney (Only the discharge hole h', corresponding to the arm m of the retort A, is visible in the drawing.) As the generator a requires fresh charges of carbon every 8 hours, it is very economical to keep this small retort always filled with charcoal, which by the time it is required will be red hot, and no more time need be lost than is required for transferring it from the one into the other By this arrangement, part of what would otherwise be waste heat is utilised, and a great saving in time and fuel effected. A still better plan is to make the charcoal on the spot, E hours being more than sufficient to char the wood and to heat it up to bright redness. Spent dyewoods, tanner's refuse, or sawdust, are excellent for the purpose, and as these would not require crushing as when charcoal is bought in lumps—a saving in labour, possibly also in money, might be made, besides being more cleanly. By a small outlay, the vapours might be condensed, and pyroligneous acid obtained as

a by-product
The apparatus is worked as follows. After having allowed the brickwork to set and partially dry for a few days, the fires are started, at first gently, to prevent the brickwork from cracking, then gradually increasing the heat until the retort becomes a dull-red colour. The latter is now filled with small charcoal, and the smaller retort with chips, sawdust, &c., and the fires are now urged until the retort and the charcoal in it have become a 'cherry red'. This heat should never be exceeded, nor should it be allowed to fall below this, as in either case it would result

in a lesser yield of carbon disulphide When the proper temperature is attained, the two openings in the lid of the retort are closed, the hds being screwed down tightly, with some clay made into a thick paste as a lute. A better way would be to cast in the top of these tubulures grooves, and the lids with a rim loosely fitting into these grooves, forming what is called an hydraulic seal, only using lead in the place of water. The heat there is sufficient to melt lead, but not so great as to prevent its use. would form a much better lute, and could be opened and closed in much less time, which, in the case of disulphide of carbon manufacture, is a great desideratum. Having previously secured all the joints throughout the apparatus, sulphur is now run in by raising the rod o' in the sulphur pot. A bubbling will immediately be heard in the gas washer at y, caused by the escape of sulphuretted hydrogen, &c. In a few minutes, this bubbling ceases, and carbon disulphide


begins to distil over Distillation is now pro- which can be replenished through the hoppers n, ceeding, sulphur being charged in every 5 minutes, about  $1\frac{1}{2}$  to  $1\frac{3}{4}$  lbs. each time, for 7 hours, when the supply of sulphur is stopped, and one hour allowed to elapse before fresh charcoal is filled in. The lid is then removed from c, and the flue-pipe H lowered. c' may now be opened, and last of all the  $lid\ h'$  is removed from h of the carboniser, and the red-hot charcoal raked from the carboniser, by means of a funnel or hopper, into the retort A The retort is now closed, taking care that the opening communicating with the flue H is the last to be shut off, and work started again At the end of each week, neither coal nor sulphur is charged in for 8 hours, after which the whole apparatus is cleaned out in the following manner flue H having been lowered on to G as described above, the cover at the end of the delivery pipe 1 is removed, and wet bags or pieces of canvas are tightly rammed down the pipe I, and past its junction with the vertical pipe I', so as to isolate the condensers, &c, from the furnace ashes are now raked out through M, as described above, the retort refilled with fresh charcoal, and while this is getting heated up, the other portions of the apparatus are examined, and where sulphur is present it is removed.

With one such retort, from 4 to 5 cwts. of carbon disulphide can be made in a day, but it is much more economical, both in labour and fuel, to have several retorts-say four-in the same furnace


# MANUFACTURE OF CARBON DISULPHIDE BY THE ELECTRIC FURNACE.

Carbon disulphide is now made on the large scale, near Penn Yan, NY., America, by the Taylor Chemical Company, by means of an electric furnace, devised and patented by Mr. Edward R Taylor (US Pat 688364, Dec 10, 1901, 871971, Nov 26, 1907), who has kindly supplied the information upon which the following description of the process is based construction of the furnace is shown in Figs 17, 18, 19, 20, 21. Fig. 17 shows a section in elevation through the electrodes, and Fig. 18, shows a cross section in elevation at right angles, the figures being taken from the patent specifications In Fig 17 is shown the position of the electrodes d, D, near the bottom of the furnace shaft, which is represented as filled with charcoal; in Fig. 18, they are seen at d. At k (Fig. 17) are pipes through which are passed broken carbons, i erefuse from the factories of carbons for are lights, for the purpose of reinforcing the main carbons; these pass down, upon, and over the ends of the main carbons, and convey the current from the electrodes to the charcoal. The intermediate conductive material protects the electrodes proper from the intense heat and eroding action and thereby prolongs their life It also tends to steady the current, fluctuations being mainly due to variations in power and not to changes of resistance in the furnace itself. The charcoal for the reaction is fed into the furnace through the opening x (Fig. 17). Preferably four electrodes are used in this construction, supplied with two-phase alternating current. Provision is made for keeping the metallic portions of them comparatively cool by feeding cold sulphur on to them through the inlets shown as o (Fig. 17), and is needed. The furnace is 16 feet in diameter

the sulphur passing into the annular chambers




shown below the hoppers, seen in detail in Figs 19, 20, and 21. The interior of the furnace thus filled with sulphur, which is gradually melted and finds its way to the reaction zone. By this construction, the heat that would otherwise be radiated from the external walls of



the furnace is absorbed, and no outside lagging

and 41 feet high. It needs much less care and is easier to manage than one of the small non or clay retorts generally used


In working, charcoal is put into the charger above the bell and the opening scaled. The charcoal falls into the shaft of the furnace as often as there is room for it, the charger being replenished from time to time. The shaft should be kept filled with charcoal and the internal spaces filled with sulphur—thus there is no loss of heat by radiation and the heat



absorbed in melting the sulphur is returned to the furnace.

Electricity is supplied in such quantity that the electrodes are not submerged in the melted sulphur to such an extent as to restrict the passage of the current. The electricity is ordinarily passed from the one electrode to the other immediately opposite, but the direction can be changed at any time so as to pass to the adjacent electrode, thus burning out any piers or 'scaffolds' that may form in the corners and which would prevent the regular descent of the

charge. The sulphur within the working chamber z soon becomes melted, and its level rises more or less nearly to the top of the electrodes. As it approaches the heat zone, the sulphur is vapourised and rises through the charcoal which, when sufficiently heated, combines with it, forming the disulphide, the vapour of which ascends through the charcoal



in the shaft, and passes out through the pipe x to a condenser. The furnaces are run for about a year before renewing the electrodes or cleaning out. Aluminium bars, 6 inches wide and  $\S$  inch thick, connect the insulaced electrodes with two Stanley inductive type dynamos, each of 330 kilowatts capacity, run, as a rule, by water power. The average voltage is about 50. The yield is usually about 14,000 lbs. of the disulphide in 24 hours, but it is capable of being increased to 25,000 lbs.

An improved type of electric furnace, more


particularly adapted for a smaller installation, is seen in Figs. 22 and 23, reproduced from the

patent specifications.

Fig. 22 shows the furnace in sectional eleva-Fig 23 shows a sectional elevation with the plane of the section at right angles to that of Fig. 22, and with portions broken away at top to expose parts behind the plane of the section. Fig 24 is a top view on which the planes of Figs. 22 and 23 respectively are indicated by broken lines, A-B and C-D, and Fig 25 represents a horizontal section on the line m-F (Fig. 23). Like reference characters refer to like parts in all the figures There is an outer shell of iron or steel within which, at the bottom, is a flat hearth of floor 4 (Figs. 22, 23, and 25) of firebrick, which supports a circular wall of like material, within which is the working chamber a of the turnace, and an outer wall, 6, concentric with the wall 5, extending upward a short distance and terminating substantially on a level with the bolster, forming lower ends of a pair of diametrically opposite embrasures, 7 and 8.

The wall 5 of the working chamber a is continued upward to the top of the outer shell 1, 2, 3; a feed chamber b for sulphur being formed between the walls 5 and 6 at bottom and between the wall 5 of the working chamber and the outer shell above the top of the outer wall 6.

The shell 1, 2, 3 terminates at top in a horizontal annular top portion, 9 (Figs. 22, 23, 24),



which is provided at suitable intervals with hoppers b' and with openings 10 (Fig. 23) at their bottoms, communicating with said feed space b, each of the hoppers being provided with a suitable plug 11, having a handle, 11', extending above the top of the hopper, so that the hoppers may be filled above the plugs, and the plugs then lifted to drop the contents of the hoppers into the feed space b, and quickly replaced to prevent the escape of fumes and the admission of air. Within the central opening of the top portion 9, of the shell, a charcoal hopper, a' (Figs. 22, 23, 24) is supported: this is constructed with a conical top, 12, provided with a capped charging neck,

13, through which the charcoal is introduced into the hopper, and a conical bottom, 14, depending within the upper end of the working chamber and provided with a central opening closed by a bell, 15, the top of the hopper being further provided with the supporting link 16 of a weighted lever, 17, connected with the said bell 15, by a rod, 18, and rendering the hopper a' normally closed at bottom A discharge pipe, c (Figs. 23 and 24), extends from the working chamber a through the top portion 9 of the shell, in an oblique direction, as shown in Fig. 23.

Openings 19 and 20 (Figs 22, 23, and 25) for the sulphur, are formed in the working chamber




Fig. 25.

wall 5, immediately above the floor 4, and at one or more higher points, connecting the feed space b with the working chamber a, and pipe openings 21 and 22 extend through the outer wall 6 and the lower shell section 1 at different heights to allow the sulphur to enter the furnace proper. Metallic conductors, d and e (Figs. 22 and 24), connected with the source of electricity, are attached respectively to the protruding upper ends of metallic electrode stems f and g, which extend to the top of the outer wall 6, and rest upon the top of this wall, as in Fig 22 Each of the electrode stems d and e is made in three sections, numbered respectively, 1', 2', and 3' in Fig 22, the upper section 1' of each of the stems being hollow and cylindrical, and extending through a stuffing box, 23, mounted on the top of the shell, and preferably constructed with a water chamber, 21, for keeping the gland sufficiently cool The lower end of the upper section has flanges 25, on opposite sides, adapting it to be coupled to the partly closed upper end of a flat middle section 2', which is tubular, excepting its lower end, and this end is trough-shaped in common with the lowermost section 3', which is detachably bolted to the trough-shaped lower end of the middle section 2'.

A pair of manholes, 26 and 27 (Fig. 22), in the middle section 2' of the outer shell, afford access to the bolts connecting the middle and lowermost sections 2' and 3' of the electrode stems, and facilitate disconnecting the lowermost sections 3' when the latter need to be renewed

The electrodes proper, h and t, are of carbon, made m sections of different lengths, but all of one and the same size, and all rectangular m cross section (compare Figs 22 and 25). Such carbon sections project at right angles from the

lower ends of the stems f and g, and may be clamped within the trough-shaped stem sections 3'. A sufficient number of the longer carbon sections are united at bottom to form the main body of each electrode and above these the shorter sections are arranged in steps, as shown in Fig 22. The electrodes h and i project inwardly, as shown in Figs 22 and 25, and are supported beneath by the bolster-forming lower ends of the embrasures 7 and 8, through which the respective electrodes extend toward each other, into the working chamber a, as shown in Fig 22
The tubular or hollow-stem sections 1' and

2' provide for feeding the material downwards upon the electrodes. The upper end of each stem is provided with a conical plug, 29, and with a gallows or yoke, 30, between which and the top of the plug 29 a wooden wedge, 31, 1s

driven to render the closure air-tight

In charging the furnace, the lower part of the working chamber a is filled with broken sulphur to the plane of the electrodes h, and a sufficient quantity of charcoal is superimposed The lower part of the furnace is then tightly closed and sealed, and the working chamber a is thereafter filled with charcoal to any desired extent by way of the charcoal hopper a' by depressing the bell 15 by means of the lever 17 The hopper a' is then automatically closed at bottom by the re-elevation of the bell 15 Crushed sulphur is fed into the annular feed space b surrounding the working chamber wall from time to time as required by way of the hoppers b', which are filled with the sulphur, and may serve to measure the quantity introduced, and are emptied into the feed space b by partly withdrawing their plugs 11 and replacing them immediately to prevent the entrance of air The disulphide and the escape of fumes vapour, as it is formed, passes off through the pipe c into a suitable condenser

An upper manhole, 32, extending outward from the working chamber a through the shell sections 1 and 2, as shown in Figs. 22, 23, and 25, facilitates access to the interior of the furnace when it is cold for adjusting or renewing the electrodes h and  $\iota$ , and for arranging the charge at the beginning of an operation; and a subjacent manhole, 33 (Fig. 23), facilitates removing the ash from time to time as may be necessary. The electrode stems f and q, and their stuffing boxes 23, are insulated with mica and asbestos

paper

This furnace is designed to be 8 or 10 feet in diameter and 18 or 20 feet high, and will produce 5000 lbs. of carbon disulphide in 24 hours. Two

electrodes only are employed

Purification of Carbon Disulphide —Tho crude product, which contains considerable quantities of sulphur, may be purified by distillation. The distilling vessel is made of sheet zinc, and is fitted in a water-bath. The vapours of carbon disulphide are passed through a wide tube into a worm surrounded by cold water, and ending in a lap at the bottom; the liquid as it condenses is allowed to trickle into a vessel placed to receive it.

Deiss employed large boilers with flat bottoms, which were 10 feet in length,  $6\frac{1}{2}$  feet in diameter, and 31 feet high. They had domed covers, externally coated by badly conducting

material, so as to reduce to a minimum the amount of carbon disulphide returned to the boiler by condensation. The boiler was capable of receiving 5 tons of crude disulphide at a single charge, and had six delivery tubes which terminated in six vertically placed condensers. At the bottom of the boiler, there were two serpentme pipes, through one of which steam was passed until the crude carbon disulphide boils. For the purpose of distilling off the last portions steam was passed through the second serpentine pipe direct into the boiler, by which means carbon disulphide vapour and steam pass over together into the condensers, when the former accumulated in a layer below the surface of the water. The distillation of 5 tons lasted 3 or 4 days, and the products of the distillation at different stages were separately collected, and served for different purposes In the first portions, foul-smelling constituents predominate, such as sulphuretted hydrogen; the intermediate portions are the purest, while the last portions are contaminated with sulphur. Much of the sulphuretted hydrogen was removed by placing a small quantity of caustic soda in the boiler.

Bonière purified the crude material by running it into a still containing a strong solution of caustic soda and heated externally by steam; the vapour was then passed through several other vessels of a similar kind containing alkaline liquids, solutions of salts of iron, lead, or copper, from which it was distilled and condensed. Millon patented a process which consisted in mixing the disulphide with half its weight of milk of lime, and then carefully distilling.

Sidot first distils the crude product, and then agitates it with mercury till the shining surface of the metal is no longer blackened by Carbon disulphide, when pure, is not affected by contact with mercury for any length of time. Or it may be shaken with 0.5 p.c. of mercuric chloride, which removes a feetid compound of The clear liquid is then decanted, 0 02 of its weight of an inodorous fat is added, and the mixture distilled in a water-bath at a low

temperature (Cloez).

Another method of purification is to add to 100 parts of commercial product from 2 to 3 parts of dried copper sulphate, and to shake the The copper salt blackens and settles mixture down, with removal of the smell of sulphuretted hydrogen Absolute purity is obtained by again rectifying over dry copper sulphate. The latter can be rendered fit for further use by ignition, treating with sulphuric acid, and again igniting. The sulphide may be retained in a state of purity by allowing it to stand constantly over dry copper sulphate (J. Soc. Chem. Ind. 2, 246).

Singer has employed the following simple

method with very satisfactory results:—
A cylindrical vessel, about 30 inches in diameter and 6 feet high, is provided with a perforated coil of lead pipe at the bottom. Into this vessel the impure carbon disulphide is run to about one-third its height. Lime-water is then pumped into it by means of a force-pump through the perforated coil. The lime-water, being specifically lighter than the carbon disulphide, rises to the surface, and, while traversing the body of the disulphide in a finely divided spray, the lime combines with the sulphuretted

81

hydrogen, &c. This washing is continued until the lime-water, which leaves this vessel through an overflow pipe near to the top, is perfectly clear. The carbon disulphide is now run into a still, about 1 p.c. its weight of a cheap colourless oil added, and covered with a layer of about 1 inch of water, to which some sugar of lead may be added. The carbon disulphide is now distilled in a water-bath and condensed in the usual way

Carbon disulphide comes into the market in sheet-iron drums, the plates forming the top and bottom being bent inwards for their better protection from blows in moving about, and in the upper end there is an opening which can be closed by a screw stopper. Great care should be taken in the storing of carbon disulphide on account of its extreme volatility, the explosive nature of a mixture of the vapour with air, and the fact that its products of combustion are three irrespirable gases, viz. carbon dioxide, sulphurous oxide, and nitrogen. The store room ought to be isolated from other buildings, and should be well ventilated. The vessels in which the substance is kept ought to be tolerably large and placed where they are not likely to receive a blow or get thrown over When required for use, the carbon disulphide can be drawn off by means of a siphon.

Properties .- Carbon disulphide is a colourloss and extremely volatile liquid, the commercial product has a repugnant and feetid smell, but when purified it has a sweetish ethereal odour, and an acrid pungent taste is highly refractive, its power in this respect being 1 645; its sp.gr. is 1 29215, 0°/4° (Thorpe) Water dissolves about 1000 of its volume of the disulphide, and yields it up again unchanged on distilling, the solution in water possesses the odour of the disulphide, and has a slight burning The aqueous solution acts as an anti-Carbon disulphide dissolves sulphur, phosphorus, iodine, bromine, chlorine, camphor, caoutchouc, oils, and fats, and may be mixed in almost any proportions with alcohol, ether benzene, and the fixed and volatile oils Sulphur and phosphorus may be obtained in crystals by the spontaneous evaporation of their solutions When a rapid stream of in carbon disulphide air is passed through it, the vapour, as it rises, is condensed to cauliflower-like masses, which make their appearance on the surface of the liquid (Waitha, Bei 3, 80) When the whole of the liquid has disappeared, the resulting solid has a constant temperature of  $-12^{\circ}$ , so long as it remains unvolatilised. According to Wroblewski and Olzewski, it solidifies at -116°, and remains solid for a considerable time, emitting a peculiar aromatic odour

Carbon disulphide boils at 460° (Thorpe), 46 42° (corr) (Thorpe and Rodger), and the vapour ignites in air at 149°. When mixed vapour ignites in air at 149°. with three times its volume of oxygen, or an amount of air containing that volume of oxygen, it is very explosive. The flame of burning carbon disulplade is blue, and gives rise to sulphurous and carbonic acid gases: CS+302 =CO<sub>2</sub>+2SO<sub>2</sub>. When carbon disulphide is burnt in a lamp on the principle of the Bunsen burner, and special precautions taken to prevent explosions, it produces a flame with an actinic power superior to that of magnesium (Compt. rend. 79, M78).

Vol. II.—T.

A mixture of the vapour of carbon disulphide with nitric oxide burns with a blue flame, which is very rich in rays of high refrangibility, and on this account has been employed in photography.

Carbon disulphide is highly poisonous, inhalation of the vapour producing giddiness, vomiting, congestion, and finally coma. Even in small quantity only, it produces in time very serious effects on the nervous system. In the workmen employed in its manufacture, it causes weakness, depression, and loss of memory. Solution of ferrous carbonate in carbonic acid water has been found in some cases effectual Carbon disulphide vapour, when as a remedy passed through a tube heated to bright redness, is partly resolved into its elements, carbon depositing and sulphur passing off with the undecomposed compound

The following table represents its vapour tension at varying temperatures (Seyferth) .-

$45.5^{\circ}$			1 atn	osphere
57 7			1 1/2	,,
66.9			2 atn	ospheres
74.3			$2\frac{1}{2}$	,,
80.6			3	,,
86 1			$3\frac{1}{2}$	,,
90 9			4	,,
$95\ 5$			$4\frac{1}{2}$	,,
$99\ 0$			5	,,

(For table of vapour pressures between  $0^{\circ}$  and 50°, v. Ramsay and Young, Chem. Soc. Trans 47, 653, also Herwig, Pogg. Ann. 137, 69, 141, 83 , 147, 161 )

The vapour of carbon disulphide, when passed over metallic oxides at a red heat, yields sulphur dioxide and carbon, together with metallic sulphides, which are generally found crystallised and resemble those found in nature Carbon disulphide is one of the most powerful sulphurising agents known, and by means of it many sulphides may be produced which are not otherwise obtainable (Fiémy) It converts oxides into sulphides when heated with them in sealed tubes The vapour is strongly attacked by nitric acid, yielding sulphuric acid and nitrous vapours

A mixture of the vapour of carbon disulphide and sulphuretted hydrogen, when passed over red-hot copper, yields copper sulphide and maish as  $CS_2 + 2SH_2 + 4Cu = 4CuS + CH_4$ 

The vapour of carbon disulphide when passed over wood undergoing destructive distillation alters the character of the carbonaceous residue, producing a material having high conducting powers for heat and electricity, and extremely sonorous when struck

Carbon disulphide is a powerful disinfectant. Meat and other putrescible bodies have been kept in an atmosphere containing its vapour for

months without change.

Heated potassium burns in the vapour of carbon disulphide with formation of potassium sulphide and liberation of carbon

When brought in contact with a solution of an alkaline hydroxide, carbon disulphide is decomposed, a carbonate and a sulphocarbonate being formed:

 $6KHO + 3CS_2 = 2K_2CS_3 + 3H_2O + K_2CO_3$ When the vapour of carbon disulphide is

passed over heated calcium hydroxide, it is decomposed, carbon dioxide and sulphuretted hydrogen being evolved.

This reaction has been utilised in the removal

of carbon disulphide from coal gas.

Chlorme has but little action on carbon disulphide at the ordinary temperatures, but in the presence of iodine, the chlorides of antimony and molybdenum, &c., chlorine replaces the sulphur with conversion of the carbon disulphide into carbon tetrachloride, and, if the action is interrupted before it is complete, intermediate compounds or sulphochlorides are formed (J. Soc Chem. Ind 6, 729)

Carbon disulphide combines with triethylphosphine  $P(C_2H_5)_3$  to form a solid compound, crystallising in red crystals of the composition

P(C2H5)3CS2

Owing to its volatility it may be made to produce great cold by its own evaporation, and if this takes place very quickly under the airpump, a temperature of  $-60^{\circ}$  may be obtained

An aqueous or alcoholic solution of caustic potash, boiled with carbon disulphide, yields with lead salts a black precipitate of lead sulphide. This is a delicate test for the substance

Carbon disulphide may be detected in coal gas by passing the gas when completely freed from sulphuretted hydrogen over red-hot copper foil, when the copper acquires an indescent lustre, and its nitric acid solution diluted with water yields after some time a precipitate with barium chloride (A Vogel, Zeitsch anal Chem [2] 6, 253).

Carbon disulphide combines with metallic sulphides, forming sulphocarbonates

$$CS_2+Na_2S=Na_2CS_3$$

Sulphocarbonic acid is a yellow oily liquid, obtained by decomposing its potassium salt with hydrochloric acid

$$K_2CS_3+2HCl=H_2CS_3+2KCl$$

Potassium sulphocarbonate has been used for the destruction of phylloxera

Uses.—Carbon disulphide is applied to a large number of useful purposes, the chief of which is in the preparation of vulcanised caoutchouc, and as a solvent for the latter in the manufacture of waterproof goods, by the deposition of a thin layer of the dissolved caoutchouc on the fabric; for the extraction of fat from crude wool; fatty oils from oil seeds and pressed residues; and for the purification of paraffin It is used in the extraction of uncombined sulphur, and of bitumen from minerals A solution of guttapercha in carbon disulphide is employed in making seamless joints and for the preparation of caoutchouc cement, various colours, ammonium thiocyanate, &c. It is also used for filling prisms on account of its high dispersive power

Advantage is taken of its poisonous properties to expel weevils and other insects from stored grain without injuring the cereal. For this purpose, a small quantity of the carbon disulphide is sprinkled over the grain; both the larvæ and eggs are killed without injury to the grain (Doyère). It is also employed to destroy chicken lice and fleas, moths in furs, and in America to kill burrowing animals, such as gophers, ground squirrels, woodchucks, prairie dogs, rats, mice, moles, &c., and as a plant

insecticide. It is used in Brazil to expel the Sauba ant from the foundations of houses. It may also be used in the analyses of oil seeds as a solvent for the purpose of ascertaining the proportion of oil which the ground seeds contain.

As carbon disulphide dissolves iodine in large quantity, but does not appreciably dissolve in water, it is employed for determining the amount

of moisture in commercial iodine.

It may be used for dissolving quinine and other alkaloids; for extracting the aromatic principles from seeds and spices, and for the extraction of the scent of flowers. For these purposes, however, the purest quality of the

disulphide can alone be employed.

A solution of phosphorus in carbon disulphide has been employed in the electrotyping of very deheate objects, such as grasses, flowers, feathers, &c. These are dipped into the solution, when, by exposure to the air, the carbon disulphide evaporates and leaves a thin film of phosphorus on the surfaces; they are then dipped into a solution of silver nitrate, by which silver is precipitated in a thin film, upon which, by the electrotype process, any thickness of silver, gold, copper, &c, can be deposited Ιf a few drops of carbon disulphide are put into a solution of silver cyanide, from which the metal is being electro-deposited, the silver is deposited bright, whereas without the disulphide it would be dull

Deiss, who first employed carbon disulphide in the extraction of fatty oils from seeds, oilcakes, and other pressed residues, used the fol-

lowing airangement —

Carbon disulphide was introduced into a large cemented blickwork reservoir, with a manhole, which is generally kept closed, 22 feet long, 6 feet in diameter, and nearly 6 feet deep, and limed with lead up to the point to which it was filled with the disulphide and water, the latter being added as a thin layer to prevent the too rapid evaporation of the carbon disulphide. Immediately above this reservoir is a worm condenser, whose tail pipe dips into the liquid in the reservoir below; on one side of the condenser there is an extractor having a capacity of 4400 gallons, and capable of receiving a charge of 12 tons of oilcake This extractor is supplied by means of a pump with carbon disulphide from the reservoir below through a pipe leading up from the reservoir to the extractor apparatus has two perforated shelves, one nearly at the top and the other nearly at the bottom, between which the substance to be operated upon is placed. In the space between the lower perforated shelf and the bottom of the extractor is a coil of pipe through which steam can be passed. Just above the upper perforated shelf there are nine pipes on the same level for the purpose of carrying the vapours produced in the operation to the condenser. There are also operation to the condenser pipes just above this shelf which are in direct communication with a still on the opposite side of the condenser, to which disulphide saturated with oil is carried, the saturated disulphide rising to the surface in the operation owing to the difference in the sp gr of the carbon disulphide (1.29) and oil (about 0.900). The still in which this liquid is distilled is about 11 feet long, 5 feet wide, and 15 inches deep, holding when half filled about 400 gallons. The liquid

is heated by steam supplied through two pipes coiled several times round the bottom of the still, and the vapour of the volatilised carbon disulphide is led through nine pipes into the condenser, and thence into the reservoir below the condenser, and is ready to be used again in the extractor for macerating a fresh quantity of material.

The time taken to fill the extractor with carbon disulphide is 8 hours, the maceration of the material 4 hours, the emptying of the contents of the extractor into the reservon at the close of the operation 2 hours, and the steaming of the oil in the still in order to free it from carbon disulphide from 8 to 12 hours Working with an apparatus such as described, about  $2\frac{1}{2}$  tons of oil can be obtained in 30 hours from 25 tons of oilcake

Extraction of fat from wool -Morson and Jerome employ for this purpose a large cylindircal vessel with a close-fitting cover and double walls, between which hot water circulates The wool is placed on a perforated shelt near the bottom of the cylinder, whilst a perforated plate, a kind of piston, could be pressed upon the wool by an arrangement of sciews Carbon disulphide is pumped from a reservoir into the extractor, and as the liquid becomes saturated with fat it is led through a pipe into a still, heated by a coil of steam pipe at the bottom, the vapour being passed to a worm condenser and the liquid disulphide passed into the reservoir which is immediately beneath. By a second arrangement of steam pipes, steam can be passed directly into the still so as to remove the last portions of disulphide. A current of air, heated to 70°, is then forced through the wool by means of an an-pump As all the parts of the apparatus are in direct communication with each other, it is necessary to supply the diminished pressure on the reservoir When air is pumped out of it and sent through a tube several times bent and the last portion fitted with a hot-water jacket, the equilibrium of pressure should be maintained, and this is done by a supply of air from a gasholder in connection with the apparatus The last portions of carbon disulphide are thus swept out of the extractor and are carried through a pipe into a second worm condenser, and thence to the reservoir. At convenient points between the macerating vessel and the still, as well as at the ends of the condensing worms, there are windows in the pipes, which serve to form an idea of the state of the operation, and close to each of these windows is a cock from which a little of the liquid can be drawn and evaporated, in order to see if a residue of fat remains

The macerating vessel or extractor is charged with about 10 tons of wool, which is then reduced to about half its bulk by screwing down the perforated plate The carbon disulphide is pumped into the extractor, and, after filtering through the wool, ascends through the perforations and passes into a pipe leading to the condenser. This is continued until the liquid comes over colourless, and leaves no residue on evapo-When the extracting process is finished, the cock admitting carbon disulphide into the extractor is closed, and a partial vacuum is formed by drawing air out of the extractor by the air-pump, the greater part of the disulphide

the first few strokes. The water-jacket is then heated by water at 70°, and the cocks are so arranged as to allow the heated air to pass through the wool, and thence into the open air By this method a considerable quantity of fatty substance is obtained from sheep's wool which was formerly lost

The following materials are treated with carbon disulphide in order to obtain fatty

matters from them .-

1 The dark-coloured residues of stearm manufactories, which are products of the treatment with sulphuric acid. They yield from 18 to 20 p.c. of fatty acid, which was formerly almost entirely lost. The residues are mixed with sawdust to facilitate the filtration of the dissolved portion.

2 The dark-brown cart-grease from the axles of carts, carriages, &c This is first treated with

sulphuric acid, washed, and dried.

3. Tow and rags used in cleaning machinery. This treatment of rags, &c, has a threefold advantage, recovery of fat or fatty acids, purification of the rags so that they can be used again, and the prevention of spontaneous combustion, which these materials are hable to on exposure

4 The refuse of the preparation of beeswax, which on treatment with carbon disulphide yields a yellow wax useful for many purposes.

5 Sawdust that has been used for filtering

oils after purification with sulphuric acid.

6. The sediment produced by the treatment of various oils with sulphuric acid contains about 50 p c. of oil, which may be extracted by carbon disulphide after washing with boiling water, drying, and mixing it with sawdust

7 Bones from slaughter-houses and kitchens to be used in the preparation of bone-black yield to carbon disulphide from 10 to 12 p c

8. The expressed residues of oil seeds, such as rape, sesamé, flax, &c, when they cannot be properly employed as fodder In all cases it is necessary to break up the orleake into small pieces before treating it with carbon disulphide, so as to ensure thorough penetration of this substance. The residue left after extracting the fat is hardly suitable for fodder, but is valuable as a manure

9. The 'cracklings' or greaves obtained by

the melting of tallow.

10. The pressed cacao beans from which no further cacao butter can be obtained by pressing 11 The pressed residues obtained in the

preparation of olive oil.

Extraction of bitumen, &c , from minerals — Carbon disulphide can be employed in the extraction of bitumen from mineral bodies which contain so small a percentage as not to pay when obtained by the ordinary process of distillation. The apparatus used consists of a close reservoir for the carbon disulphide, over which is a cooling apparatus which serves as condenser. Below the reservoir are two filters provided near the bottom with perforated shelves. are introduced into the filters upon the shelf through doors near the bottom, which are then closed and made air-tight. Carbon disulphide is now admitted through a pipe at the bottom of each filter, and as the liquid rises it percolates adhering to the wool being carried with it at through the minerals, dissolving out the bitumen,

and when full flows out at the top through a pipe leading to a still, where the saturated liquid is distilled, the vapour of carbon disulphide ascending into the condenser, and the liquid carbon disulphide as it condenses trickles into the reservoir beneath. The bituminous matters, which remain in the still, are drawn off by a cock at the bottom, and the disulphide which remains absorbed by the minerals in the filters is expelled by a current of steam. This apparatus is practically continuous, very little loss of the disulphide taking place at each operation, and as much as 12 p c of bitumen can be obtained, whereas only 7 or 8 p c. could be had by distillation

Extraction of spices, &c.—Bonière has devised an apparatus for extracting the aromatic principles of various spices and other substances used for seasoning food, by means of carbon disulphide It is on the same principle as the apparatus already described, and consists of a series of wire-gauze sieves arranged one over the other in an air-tight case The carbon disulphide is made to pass through the sieves, which contain the substance to be operated upon, e.g. pepper, thus dissolving out the active principles, and as it reaches the top it flows through a pipe into a boiler The boiler contains salt, sugar, lactose, dextrin, saltpetre, or other material The liquid is distilled in the boiler, by admitting steam into a kind of jacket, when the disulphide distils over and is condensed in a worm condenser, while the salt or sugar remains behind, retaining all the aromatic or active principles of the spice.

Beigot has constructed an apparatus for rapidly determining the quantity of oil in seeds, &c, by means of carbon disulphide It consists of a glass vessel in the neck of which a cylindrical glass vessel is accurately fitted, and at the side in a tubulure is inserted a small air-pump 50 grams of the finely ground seed to be examined are placed in the cylindrical glass vessel, and are covered with a diaphragm upon which 50 more grams are placed and covered with a second diaphragm Carbon disulphide is now poured on until the seed is fully moistened, and after a few minutes a partial vacuum is made in the apparatus by means of the air-pump, when the pressure of the air drives out the disulphide and oil in solution into the vessel beneath. This is repeated until the carbon disulphide comes off colourless and leaves no oily stain on contact with filter About 400 or 500 grams of carbon dipaper sulphide are sufficient to exhaust the above amount of seed

The oily solution thus obtained is heated ver a water-bath, and as soon as the disulphide is expelled, the residue is allowed to cool and is weighed.

H. L. Greville has extracted with advantage the carbon disulphide absorbed in the purifiers of the gasworks The lime used for this purpose, after being discharged from the purifiers, is introduced into a boiler capable of receiving a charge of several hundredweights, fitted with a perforated false bottom, pressure gauge, and an arrangement for admitting steam. The yield of carbon disulphide from 100 tons of spent lime is about 11 tons In addition to the value of the crude product thus obtained, the lime in 1867; occurs in certain hepatic waters or

remaining is, with a small addition of fresh lime, fit for further use in the purification of coal gas. As is well known, oxide of iron, after continued use in the absorption of sulphuretted hydrogen from coal gas, contains about 50 pc of free sulphur. The crude carbon disulphide, obtained as described, can be used for the extraction of this sulphur, and the oxide which remains is revivified, and is again fit for purifying purposes The sulphur is recovered by distillation, and the carbon disulphide can be used over again (J. Soc. Chem. Ind. 2, 488).

The principal seats of manufacture of carbon disulphide in England are London, Manchester, certain towns in Yorkshire, and Ironbiidge, in France, Paris, Boideaux, and Marseilles. Germany, Austria, and Sicily have several manufac-

Carbon monosulphide (rCS) A reddish-brown powder obtained by exposing carbon disulphide to sunlight Insoluble in water, alcohol, turpentine, and benzene (Sidot, Compt rend. 69, 1303; 74, 180, 81, 32).

It is also formed when thiocarbonyl chloride reacts at ordinary temperature with nickel carbonyl  $x \text{CSCl}_2 + x \text{Ni}(\text{CO})_4 = x \text{NiCl}_2 + 4x \text{CO} + (\text{CS})_{x}$ The reaction also takes place at -20°, although more slowly The nickel chloride produced is extracted with water, and the brown residue is dried by heating at 100°, then at 150°-200° under reduced pressure.

Carbon monosulphide has sp gr. 1.6, but after compression into a solid block the sp gr is 183. It is a non-conductor of electricity. In concentrated sulphuric acid it forms a purplish-brown solution, the colour of which is destroyed on boiling, when carbon dioxide and sulphur di-oxide are evolved. When poured into water the purplish-brown solution yields the unchanged It also forms brown solutions monosulphide in aqueous or alcoholic ammonia, in ammonium sulphide, in potassium-hydroxide sulphide and hydrosulphide With nitric acid it gives a red solution. When heated in a vacuum at 360° or below, no change takes place, only a small quantity of hydrogen sulphide is evolved, but at a dull red heat it decomposes thus

$$2(CS)_x = xC + \tau CS$$

When carbon disulphide vapour at low pressure is subjected to the action of the silent electric discharge, sulphur is deposited, and gaseous carbon monosulphide is The gas can be collected in a tube evolved cooled with liquid air, when it condenses together with carbon disulphide If the temperature of the condensing tube is allowed to rise, the monosulphide polymerises with explosive violence to the brown solid carbon monosulphide  $(CS)_x$ . In presence of excess of carbon disulphide, the gaseous monosulphide is fairly stable under certain conditions even at high temperatures, but it is completely decomposed when passed through 10 cm of a red-hot tube packed with Roy. Soc. 1911, 85, 574.)

Carbon oxysulphide or Carboxyl sulphide or Carbonyl sulphide COS. Discovered by Than,

mineral springs Prepared (1) by passing a mixture of carbonic oxide and vapour of sulphur through a hot tube. CO+S=COS.—(2) By the action of sulphuric acid upon potassium or ammonium thiocyanate.

$$\begin{array}{l} \text{CNKS} + 2\text{SO}_4\text{H}_2 + \text{OH}_2 \\ = \text{COS} + \text{SO}_4\text{HK} + \text{SO}_4(\text{NH}_4) \end{array}$$

The gas is purified by passing it successively through strong caustic alkali, concentrated sulphuric acid, and finally through a mixture of triethylphosphine (1 part), pyridene (9 parts), and nitrobenzene (10 parts) It is then cooled to 10°-20°, and the last traces of impurities are removed by liquefying it, when the impurities remain uncondensed (Hempel, Zeitsch. angew Chem 1901, 14, 865).—(3) By heating together trioxide and carbon disulphide.  $CS_3 + 3SO_3 = COS + 4SO_3$  (Armstrong) —(4) By heating urea with carbon disulphide in a sealed tube to  $110^{\circ}$ :  $COH_4N_2 + CS_2 = COS + CNS NH_4$ . The oxysulphide is given off together with hydrogen sulphide on opening the tube, and the latter is absorbed by passing through lead accetate (Ladenburg)—It is also formed in the combustion of carbon disulphide with oxygen (Dixon and Russel, Chem. Soc Trans 1899, 610), and by the passage of the dark electric discharge through a mixture of carbon disulphide and carbon monoxide (Losanitsch and Jovitschitsch, Ber. 1897, 30, 135)

Properties.—It is a colourless gas, odourless when quite pure (Stock and Kuss) It acts strongly on the nervous system, a rapid overpowering action following the inhalation of small quantities It is readily inflammable, and forms with oxygen an explosive mixture (Russell, Chem Soc Tians 1900, 356) The gas is soluble in water, 1 cc dissolving 0.8 cc of the gas at 13 5° and 756 mm pressure The solution gradually decomposes, forming hydrogen sulphide and carbon dioxide A platinum wire heated to whiteness in the gas decomposes it into sulphur and carbonic oxide, the latter occupying the original volume of the gas It has a specific gravity of 2 1046, and may be easily poured from one vessel to another burnt in air, it produces carbon dioxide and sulphur dioxide With caustic alkalis, it yields a mixture of carbonate and sulphide

$$COS + 4KHO = CO_3K_2 + K_2S + 2H_2O$$

Carbon oxysulphide reacts with solutions of metallic salts, forming the metallic sulphide and carbon dioxide, and is readily oxidised to carbon dioxide and sulphure acid by bromine water or acid permanganate.

It is slowly decomposed by acid cuprous chloride, thus

$$COS + 2CuCl + H_0O = CO_0 + 2HCl + Cu_0S$$

(Berthelot, Ann. Chim. Phys. 1898, [7] 14, 205). With magnesium organic compounds, it can be employed in the synthesis of thio-acids and of s-trisubstituted carbinols (Wiegert, Ber. 1903,

36, 1007)

Its critical temperature is 105°, and its critical pressure is 60 atmospheres. It is liquefied at a pressure of 125 atmospheres and temperature 0° to a colourless, mobile, and highly refractive liquid, 1D—87° 124; m.p.—1382°, boiling at —50°2°, at 76° mm., which dissolves sulphur and mixes with alcohol and

ether, but not with water [Ilosvay, Bull Soc. chim. [2] 37, 294; Stock and Kuss, Ber 1917, 50, 159).

Carbonyl ketene; Malonic anhydride) C<sub>3</sub>O<sub>2</sub>. CARBON SUBOXIDE (Carbon carbonyl; pared by distilling ethyl malonate under reduced pressure into a vessel containing a large excess of phosphorus pentoxide and some glass wool, the vessel being kept at a temperature of 300° The resulting products are passed through a dry tube, and then into a vessel immersed in liquid air to condense the ethylene and carbon suboxide The ethylene volatilises from the mixture at the ordinary temperature, leaving a small quantity of colourless liquid. This is volatilised and collected in a tube kept at between 60° and 70° (Diels and Wolf, Ber 1906, 689). Diels and Meyerheim (191d. 1907, 355) showed that carbon suboxide may be obtained by similar methods from methyl, benzyl, and phenyl malonates and also from ethyl methanetucarboxylate; and that by heating malonic acid at 140°-150° there is formed acetic acid, carbon dioxide, and about 10-12 p c of carbon suboxide Staudinger and Bereza obtained carbon suboxide by treating malonyl chloride with silver, lead, or zinc oxides, the yield is only 10 pc (Ber. 1908, 4461) By treating an ethereal solution of dibrommalonyl chloride with zinc shavings, a method similar to that for preparing the ketenes. Staudinger and Klever obtained a 50-80 pc yield of carbon suboxide (Ber. 1908, 906) Stock and Stoltzenberg (Ber 1917, 50, 498) find that the poor yields of carbon suboxide usually obtained by the action of phosphoric oxide on malonic acid are largely due to the polymerisation of the suboxide under the catalytic influence of the dehydrating agent, and they describe an apparatus in which the suboxide is rapidly removed by carrying out the reaction in a vacuum and condensing the product by liquid air.

The pure oxide is a colourless, very mobile, refractive liquid with a pungent smell resembling acetaldehyde and mustard oil It is poisonous the bols at 7° under 761 mm, and its vapour density corresponds to the formula ( ${}_{3}O_{2}$ ; mp -107° (-111.3° Stock and Stoltzenberg), density of liquid 1.11. On passing the vapour through a constricted tube, a metallic mirror resembling arsenic is obtained. At low temperatures, carbon suboxide is quite stable, but between 0° and 15° a trace of impurity rapidly polymerises it to a red solid. This polymeride is very hygroscopic, and on addition of water, heat is developed, carbon dioxide evolved, and an eosin-red solution formed · by heating the red compound carbon dioxide, carbon monoxide and carbon suboxide are formed, and at higher temperatures, carbon monoxide and dioxide are produced, the residue being a mixture containing more carbon than is required for carbon suboxide. By keeping carbon suboxide for 2 days in a sealed tube at 15°, it completely changes into the polymeride, forming a dark-red solid mass, soluble in water to a dark-red solution; this change is instantaneous at 100°. Carbon suboxide burns with a bright-blue but smoky flame, forming carbon dioxide. Carbon suboxide does not combine with aldehydes, Schiff's bases or hydrogen cyanide (Diels and Lalin,

Ber. 1908, 3426), but in liquid sulphur dioxide at  $-40^{\circ}$  to  $-50^{\circ}$  forms a white crystalline compound with formic acid, and a colourless syrup with acetic acid; the constitution of these compounds is doubtful Water quickly reconverts it into malonic acid

Two formulæ have been proposed for carbon suboxide; Diels and Wolf represent it as the anhydride of malonic acid () C:C.C (), Michael (Ber. 1906, 1915, 1908, 425), as the lactone of  $\beta$ -hydroxypropiolic acid  $C \subset C \subset C$ .

The first formula seems to be correct, for the following reasons (v. Diels and Blumberg, Bei.

1908, 82, 1233) .-

(1) Bp 7° Propiolic acid boils at 144°, and if  $\beta$ -hydroxypropiolic acid had a boiling-point, it would boil higher still, and its lactone would certainly not boil at so low a temperature as 7°.

(2) The values for the molecular refraction and molecular dispersion are high:  $M_{\rm p}=16.6$ ,

 $\gamma - \alpha = 0.736$ , 0.739, 0.862 (three determinations). (3) Similarity of carbon suboxide to the

metallic carbonyls.

- (4) Similarity to the isocyanates and ketenes (5) Addition of four atoms of bromine to form dibrommalonyl bromide and the regeneration of carbon suboxide from this compound
  - (6) The following chemical reactions.
- O C C C O+2H2O=HOOC CH2 COOH(malonic acrt) O C C C O+2HCl=CloC CH2 COCl(malonylchloride) O C C C O+2NH3
- $\begin{array}{c} = H_{3} \text{N OC CH}_{2} \text{ CONH}_{2} \text{ (malonamide)} \\ \text{O C C O} + 2C_{6}H_{5}^{2}\text{NH}_{2} \\ = C_{6}H_{5}^{2}\text{HN OC CH}_{2} \text{ CONH } C_{6}H_{5} \text{ (malonanilide)} \end{array}$
- (7) The improbability of the existence of a  $\beta$ -lactone containing two carbon atoms triply linked together (v Ketones).

CARBONADO v DIAMOND.

CARBONITE, or natural coke A coke-like material formed by the baking action of intrusive igneous lock-masses on seams of bituminous coal Found in Ayrshire and in Chesterfield co, Virginia L. J S.

CARBONITE. An explosive consisting of nitroglycerine, 25-27 parts, barium nitrate and potassium nitrate, 30-36 parts, wood meal, 40-43 parts, sodium carbonate, 0.5 parts (vwood meal,

EXPLOSIVES)

CARBONYLS Metallic compounds of car-The existence of this class of bon monoxide compounds was first made known in 1890, when Mond, Langer, and Quincke discovered that carbon monoxide, passed over heated, finely divided nickel, yielded a colourless volatile liquid

Iron, cobalt, molybdenum, and ruthenium have since been found to yield carbonyls by

similar treatment.

The production of the carbonyl is facilitated by conducting the operation under pressure and, in most cases, at a fairly high temperature.

For the production of the iron and nickel compounds, the carbon monoxide is simply passed over the heated metal, but for the other compounds a special apparatus is used This consists of a nickel-steel tube, in the form of a test-tube, capable of withstanding a pressure of 500 atmospheres and a temperature of 450°. It is copper lined to prevent the action of carbon monoxide upon the iron. The carbon monoxide under pressure is led in through a pressure valve

and down a copper tube reaching almost to the bottom of the nickel-steel tube, where it comes in contact with the metal to be experimented upon, contained in a glass tube loosely fitted to the top of the apparatus The excess of the to the top of the apparatus carbon monoxide, and other vapours, escape through a pressure valve and pass through a tube immersed in a freezing mixture, in which any compound that is formed is collected (Chem. Soc. Trans. 1910, 97, 798).

The carbonyls have very similar chemical Upon heating, in some cases an properties intermediate compound is formed, but they all decompose ultimately, yielding the metal, which is deposited as a bright mirror, and carbon Dilute acids are practically without monoxide action upon them, but strong acids decompose them readily, giving the corresponding salt, with the evolution of carbon monoxide and hydro-The halogens also decompose them are decomposed by moist air, giving precipitates of the metallic hydroxides and, in some cases, what is probably a hydrated basic carbonate. They are more or less soluble in the common organic solvents, but all are insoluble in water

In physical properties, they vary considerably, thus, in volatility, they range from the non-volatile Fe<sub>2</sub>(CO)<sub>4</sub> of sp gr. 2 08 to the very volatile Ni(CO)<sub>4</sub> of sp gr. 1 32 They are highly coloured, except in the cases of molybdenum and nickel carbonyls, which are white and colourless respectively Some can exist in the three states of solid, liquid, and vapour, and others in only one or two of these phases

Owing to their diversity in composition, a general chemical formula cannot be assigned to them, but it is probable that they are similarly

constituted. A ring formula, e g Nic `CO:ĊO

does not explain all their properties, and the compound Fe<sub>2</sub>(CO)<sub>2</sub> cannot be brought into line with the other compounds which only contain one atom of the metal Presumably they have a molecular rather than an atomic constitution (Mond, Hertz and Cowap, Ic.). The various known carbonyls are described under the several metals.

CARBORUNDUM (Crystalline silicon car bide) SiC. This compound was discovered accidentally by Acheson, in 1891, and is now manufactured under his patent (Eng Pat 17911, 1892) by the Carborundum Co at Niagara Falls, USA, and to a smaller extent in Canada.

Preparation —Amorphous silicon carbide is formed by heating carbon and silica together at about 1200°, as a greenish powder (Schutzenberger, Compt rend 114, 1089), but carborundum is only formed at a temperature of 1950° (Tucker and Lampen, J Amer Chem Soc 1906, 28, 853) (1820° according to Gillett), which is attained by means of the electric arc thus prepared pure silicon carbide from the vapours of carbon and silicon (Compt. rend. 1893, 114, 1089)

The method of manufacture used at Niagara is as follows .—The furnaces are built of firebrick and are about 16 feet  $long \times 5$  feet  $high \times 5$  feet wide. The ends are solid, about 2 feet thick, and carry the terminals. Each terminal consists of 60 carbon rods, 30 minches long and

3 inches diameter, into the ends of which fit copper plugs which in turn fit into sockets in a square copper plate bolted to the outside of the wall, and connected with the leads Only the end walls and the bed are permanent, the sides are built up with the charge and taken down to remove the product.

In charging, the furnace is half filled with the materials, which must not touch the electrodes, and a cylindrical core 21 inches diameter, composed of pieces of coke 1-1 inch diameter, is built up between the electrodes Over this the charge is built up to a height of 8 feet. current passes through the coke and forms numerous arcs which heat the charge to a very high temperature The charge is composed of:

Coke .		34 2 parts
Sand .		54 2 ,,
$\mathbf{Sawdust}$		9.9 ,,
Salt .		17

The salt is added as a flux, the sawdust to increase the porosity of the charge and allow the escape of the carbon monoxide formed in the reaction ·

$$S_1O_2 + 2C = S_1 + 2CO$$
  
 $S_1 + C = S_1C$ 

During the run, about 6 tons of this gas are given off; it is allowed to burn at the top of the furnace.

The alternating current supplied by the Niagara Power Co at 2200 volts is transformed down to 165 volts, and a large water rheostat is used to regulate or interrupt the current, the usual means being too dangerous with the heavy currents used At the beginning of a run, the EMF. is 165 volts, but within 13 hours the resistance decreases and the EMF. is reduced to 125 volts, during this period, the current increases from 1700 amperes to 6000 amperes The conditions then remain steady during the whole run of 36 hours

A modification of the process consists in preheating the charge by combustion of gaseous or solid fuel (Tone and the Carborundum Co US Pat 908357, 1908) By moving the position of the arc relatively to the charge, the process may be made continuous (ibid US Pat 937119, 1909).

The charge per furnace is about 30,000 lbs. of which the core forms about 3 p c, the energy used is about 26,400 kilowatt-hours, and the product contains about 6700 lbs of carboiundum and 5000 lbs. of amorphous silicon carbide.

The carborundum is dug out in large masses and pounded with water in a mechanical crusher, and then digested with sulphuric acid (1:2) for 3 days at 100°, and finally washed with The finer portions, which are washed away, are collected separately and known as flours. The residue is dried in a kiln and graded through a system of screens Hand-washed powders are of various grades of fineness, according to the length of time they remain suspended in water (Kohn, J. Soc. Chem. Ind. 1897, 863).

Some analyses of these product are tabulated above :--

	I.	II.	III.	IV.	v.
Sı C	70 00 30 00	69·93 29 90	62·70 32·26	69·10 30·20	65 42 27 93
$\left. egin{array}{c} \mathrm{Al_2O_3} \ \mathrm{Fe_2O_3} \end{array}  ight\}$			0 93	0 46	5.09
CaO				0.15	0.38
MgO			0.11		0 21
mgo					

I. is the calculated composition for SiC; II is Moissan's pure product (Compt. rend. 1893, 114, 1089), III is commercial carborundum, IV is the same after purification by Muhlhauser's method-heating in oxygen for I hour and treating with hot sulphuric and hydrofluoric acids (Dingl poly J. 1893, 289, 164); V is amorphous silicon carbide

Weber has patented a method of production from kaolin and coke in the electric furnace. The product is treated with water, which decomposes the aluminium carbide, leaving the carborun-

dum (U.S. Pat 728528, 1903).

The production of a dense, compact variety, by the action of the vapours of carbon and silicon upon silicon carbide has been patented by the Carborundum Co. (US Pat. 913324) and by Bouvier (Fr. Pat 350369, 1904) It is very hard, resists acid, and is a good insu-

Its preparation from silicon monoxide and carbon has been described by Potter and Westing house (U.S. Pat 875673, 1907)

Articles may be formed in pure carbon and afterwards converted into carborundum by heating to a very high temperature in a bed of finely powdered carborundum or of sand and carbon (Bolling, Eng. Pat. 6693, 1905, D. R. P. 195533, Fr Pat 353017)

Properties —Carborundum crystallises in flattened hexagonal rhombohedia, having an adamantine lustre and a hardness of 9 5 (Muhlhauser, Zeitsch anal Chem 1893, 637). The colour of the commercial product is greenishgrey to yellow or blue; obtained pure from pure materials, the compound is colourless. Its sp gr is 3 23 at 15° (Fitzgerald, J Soc. Chem. Ind 1897, 246).

It is infusible and quite unaffected up to a temperature of 2220°, when it decomposes into silicon and graphitic carbon (Tucker and Lampen, J Amer. Chem Soc 1906, 28, 853). Acids have no action upon it, but it is attacked by fused alkalis with formation of a silicate and carbon.

The determination of silicon in carborundum is made by converting it to silicate as above, and precipitating as silica; carbon is estimated by oxidising the finely powdered substance with lead chromate (Matthews, J. Soc. Chem. Ind. 1895, 755, Muhlhauser, Zeitsch. anal. Chem. 1893, 637)

Uses —Carborundum is used chiefly instead of emery as an abrasive. Though it is more costly, it does the same work more effectively in one-third to one-fourth the time (Muhlhauser, lc.) The powder is used in glass cutting and grinding, and for polishing

It is made up into wheels, hones, &c., by mixing with moistened kaolin and felspar, moulding under hydraulic pressure, and firing the articles in a kiln for 7 days. Other binding materials, such as shellac, are used for special purposes, and papers, similar to emery paper, are made. The binding material may be dispensed with by moulding the carborundum with water and then heating the articles to 2500°F. In an oxidising flame (Imray, Eng. Pat. 9963, 1904, Tone, U.S. Pat. 772262)

Carborundum can be used instead of diamond for drill heads, if incorporated in a suitable metallic or ceramic matrix (Bouvier, Fr. Pat.

375338, 1907).

Silicon carbide is used instead of ferro-silicon as a source of silicon in steel making. About 0·1-0·4 p.c. is placed in the ladle and dissolves readily in the molten steel, ensuring solid castings (Kaufmann and Bouvier, Fr. Pat. 344906, 1904; Eng. Min J. 75, 481)

The amorphous silicon carbide, formerly a waste product, is used for making highly refractory firebricks and the retorts for zinc distilling (Chesneau, Ann Chim. anal appl. 1908, 13, 85). A refractory cement is made from carborundum 90-60 parts, fireclay 10-40 parts, lime 0-4 parts, and water-glass solution 47°Bć 20-50 parts, mixed, dried, and repowdered (Muller, Fr. Pat. 338914, 1903)

Neumann has shown that silicon carbide reduces silver, copper, nickel, and lead from the chlorides (Chem. Zeit 1900, 24, 1013)

By the action of silicon carbide on metallic oxides, many metallic silicides may be obtained. This reaction may be used for the preparation of special ternary or quaternary steels at a single operation (Baradac-Muller, Soc. Mg. Cie. de France, Proc. verb. 1998, 245)

o-CARBOXYPHENYLTHIO-G L Y C O L L I C ACID. o - CARBOXYPHENYLTHIOLACETIC ACID v. PHENYLTHIOGLYCOL-O-CARBOXYLIC ACID.

CARBURINE, Trade name for a variety of petroleum spirit, sp gi 0 68 Used as an 'enricher' of coal gas.

CARCLAZITE or CARCLAZYTE. A name suggested by J. H Collins (The Hensbarrow Grante District, Truro, 1878) for the china-clayrock from which china-clay is prepared Sonamed from the Carclaze mine, near St. Austell, in Cornwall L. J. S.

CARDAMOM OIL v OILS, ESSENTIAL

CARDAMOM SEEDS. Cardamomi Semina, B.P. The dried ripe seeds of Eletteria Cardamomum (Maton)

CARDINAL-RED, Syn Acid magenta (v. Triphenylmethane colouring matters)

**CARDOL.** A non-volatile oil found in the pericarp of the cashew nut, *Anacardium occidentale* (Linn).

CARICIN. An oil found in the seeds of the papaw tree, Carica papaya (Linn.) (Peckolt, Pharm J. [3] 10, 343)

CARMINAPH v Azo- colouring matters CARMINAPH GARNET v Azo- colouring MATTERS.

CARMINAZARIN v Cochineal.

CARMINE. The red colouring matter extracted from the cochineal insect. It is of somewhat variable composition, containing besides the colouring matter some organic matter from the insects and a little of the salt used to precipitate it from solution.

Preparation — The powdered cochineal is boiled with water, the solution filtered or al-

lowed to settle, and the colouring matter precipitated by addition of some salt, such as alum, stannous chloride, cream of tartar, Roman alum, &c. The precipitate is allowed to settle, and collected. Sometimes fish glue or white of egg is added before precipitation. Carmine is somewhat soluble in water and very freely soluble in alkalme solutions. The whole of the colouring matter is immediately precipitated by alum in the form of a lake.

In dyeing with carmine, a mordant of alum or stannous chloride is used. Kielmeyer recommends the following mixture for printing on woollen goods 14 kilos, gum, 17 5 kilos, cochineal lake, and 2 25 kilos, fustic lake, are mixed and warmed with 15 lit water till the gum is dissolved, then I kilo of oxalic acid and 1 75 kilos, acid sodium oxalate added, and the mixture allowed to cool, after which 2 25 kilos, sodium acetate are added. After printing, the fabric must be dried for a day or two, steamed under a slight pressure, and washed in a stream of water.

Detection of carmine in cloth—Alcohol takes no colour from the cloth, but when it is boiled in solution of aluminium sulphate the solution turns red and the colour is unchanged by the addition of acid sodium sulphate.

The absorption spectrum of carmine in alkaline solution is remarkable, showing two dark bands between the lines D and E very similar to those of blood. Murexide and madder show no such bands. When wine is coloured with carmine, the characteristic absorption bands cannot be seen unless more than 12 p.c. of the total colour is due to carmine (Gautier), but the adulteration can be detected by adding alum, when the colour changes to a rose tint.

It is said that the finest shades of carmine can only be obtained by working in direct sunlight (v. COCHINEAL)

CARMINIC ACID v COCHINEAL.

CARMOISINE. Syn for Azorubin (v. Azocolouring matters)

**CARMUFELLIC ACID**  $C_{12}H_{20}O_{16}$  An acid crystallising in thin plates, said to be formed by the action of nitric acid on oil of cloves (Muspratt and Dawson, Phil Mag [4] 2, 293).

CARNALLITE. Hydrated double chloride of potassium (KCl 268 pe) and magnesium, KCl MgCl<sub>2</sub> 6H<sub>2</sub>O. It occurs abundantly in the uppermost layers ('carnallite zone') of the beds of abraum-salts in the Prussian salt deposits, and is the chief commercial source of potassium chloride It is also found at Wittelsheim in Upper Alsace. The mother-liquor, remaining after the potassium chloride has crystallised out, contains traces of bromine and rubidium, the former of which is a valuable by-product. Thallium has also been detected in traces. When pure, the mineral is colourless and water-clear, but it is often coloured yellowish or reddish by intermixed impurities. Red carnallite owes its colour and metallic sheen (resembling that of sunstone) to the presence of minute scales of hæmatite. Crystals are rare; they are orthorhombic, and have the form of six-sided pyramids with numerous well-developed faces. Owing to the extreme deliquescence of the salt, crystals must be preserved in scaled vessels.

Carnallite and bromcarnallite, in which bromine takes the place of chlorine in the above formula, have been prepared artificially from solutions containing a large excess of magnesium chloride and at a temperature of over 25°. L. J. S

CARNAÙBA WAX or CARNAHUBA WAX v. WAXES.

CARNAÜBIC ACID C23H47COOH 18 obtained from Carnauba wax; m.p. 72 5° (Sturcke, Annalen, 1884, 223, 283, Darmstaedter and Lifschutz, Ber. 1896, 619, 2899).

CARNELIAN v. CHALCEDONY.

CARNINE is the name given to a base isolated from meat extract by Voit (Weidel, Annalen, 1871, 158, 353), it occurs also in beer yeast extract (Schutzenberger, Chem. Zentr 1877, 73), in the muscle of some freshwater fish and frog's flesh (Kruckenberg and Wagner, ibid 1884, 107), and in beet juice (v. Lippmann, Ber. 1896, 29, 2645). Carnine crystallises in chalk-white druses, is spaningly soluble in cold, readily so in hot, water, insoluble in alcohol or ether, it forms a crystalline hydrochloride and platinichloride, and a spaningly soluble compound with silver nitrate (Weidel,

Haiser and Wentzel (Monatsh 1908, 29, 157) have shown that carnine is an approximately equal molecular mixture of hypoxanthine and inosine C10H12O5N4, and the two are separated by the action of acetic anhydride which converts the mosme into the acetate ('10H9O2N4(OAc), m p. 236° (decomp) Inosine crystallises in silky needles, m p. 215° (decomp ),  $[a]_{D}^{18°}$  –49 2°, and 100 c c of aqueous solution saturated at  $20^{\circ}$ contains 1 615 grams When mosine is hydrolysed by sulphuric acid it yields hypoxanthine and a pentose, originally termed carnose by Levene and Jacobs (Ber 1909, 42, 2102, 2474) and afterwards identified with a-ribose Inosine is closely related to inosic acid (hypoxanthine phosphoriboside) (Levene and Jacobs, wid 1910, 43, 3147, Proc. Amer. Soc Biol Chem 1910, 25, Haiser and Wentzel, Monatsh 1909, 30, 147), for by heating a dilute aqueous solution of barium mosate at 125°-130° for 6 hours, mosine and barium phosphate are formed relation between the two compounds is probably represented by the following formulæ:

CO-NH

M. A. W. CARNITINE C<sub>7</sub>H<sub>15</sub>O<sub>3</sub>N, a base isolated from Liebig's beef extract and present also in fresh beef, is readily soluble in water yielding an alkaline solution, the nitrate crystallises in needles and has  $[a]_{D}$  about  $-22^{\circ}$ ; the platinchloride C14H32O6N2, PtCl6 forms minute orangered needles, m.p. 214°-218° (decomp) (Gulewitsch and Krimberg, Zeitsch physiol. Chem 1905, 45, 326), the hydrochloride is hygroscopic and lævo-rotatory. The aurichloride

forms pale yellow needles and orange needles or prisms, m p.  $150^{\circ}$ , the mercurichloride

is an oil that slowly crystallises , the compound  $\rm C_7H_{16}O_3N.2HgCl_2$  has m p  $\rm 196^\circ-197^\circ$  (Krimberg, ibid 1907, 50, 361) The monoacetyl derivative of the hydrochloride OAc C7H, O2N, HCl forms a yellow crystalline platinichloride

m p 199° (Engeland, Ber 1909, 42, 2457) The ethyl ester is identical with Kutscher's oblitine (Zeitsch Nahr. Genussm 1905, 10, 534), (Krimberg, Zeitsch physiol Chem 1908, 58, 417; Ber 1909, 42, 3878, Engeland, lc)

Carnitine is identical with Kutscher's novaine (Krimberg, Zeitsch physiol Chem 1908, 55, 466; Engeland, Ber 1909, 42, 2457), and is probably a hydroxy derivative of a betaine, since on reduction with hydrogen rodide and red phosphorus at 130°, it yields a product identical with Willstatter's γ-trimethylbutyrobetaine

$$NMe_{3} < \begin{matrix} CH_{2} & CH_{2} \\ O & -OC \end{matrix} > CH_{2}$$

(Ber 1902, 35, 584, see also Fischer and Goddertz, Ber 1910, 43, 3272–3280) Krimberg (Zeitsch physiol Chem 1907, 53, 514) According to Engeland (Ber 1909, 42, 2457, 43, 2705) and Rollett (Zeitsch physiol Chem. 1910, 69, 60), cainitine hydrochloride is the chloride of ahydroxy-\gamma-trumethylamino-n-butyric acid

the isomeride,  $\beta$ -hydroxy-trimethylamino butyiic acid chloride crystallises in needles, gives a chloroaurate, m.p. 145°, which decomposes 225° (Engeland, ibid. 1910, 43, 2705-2707) V. BETAINES M. A. W.

CARNOSINE C<sub>9</sub>H<sub>14</sub>O<sub>3</sub>N<sub>4</sub>, a base isolated from Liebig's extract of beef (see Smorodinzew, Zeitsch physiol Chem. 1913, 87, 20, ibid 92 19 and 227, for the percentage of carnosine found in muscle extracts) is soluble in water, crystallises in rosettes of large colourless needles, mp  $246^{\circ}$ – $250^{\circ}$  (decomp). It has  $[\alpha]_{D}+21$  independently of the concentration (Gulewitsch (Zeitsch physiol Chem 1913, 87, 6-11), and has a strongly alkaline reaction The vapours of carnosine give the pyrrole reaction with a pine splinter, Gulewitsch (ibid. 87, 8) Carnosine may be precipitated from an aqueous extract of muscle by a saturated solution of mercuric sulphate in 5 p c sulphuric acid, the precipitation is accelerated by the addition of two volumes of alcohol and a little ether (Dietrich, Zeitsch physiol Chem 1914, 92, 212-213). Smorodinzew (101d. 214-220) recommends a 10 pc. solution of mercuric sulphate in 5 pc sulphuric acid solution Carnosine can be isolated by precipitation with a copper salt, Mauthner (Monatsh. 34, 883-900). The nitrate  $C_9H_{15}O_9N_8$  crystallises in large stellate clusters of needles, m.p. 219° (decomp.),  $[\alpha]_D^{19}^{55}+222$ , p=7.603 p.c., c=7.820 p.c. The rotatory power increases slightly on dilution; it falls to about

half its value in presence of nitric acid (Gulewitsch, ibid. 1-6), the copper derivative

C9H14O4N4Cu

is crystalline and decomposes at 220°. The silver derivative and acid double salt with silver nitrate closely resemble the corresponding derivatives of arginine (Gulewitsch and Amiradžibi, Ber. 1900, 33, 1902). Carnosine is probably identical with Kutscher's ignotine (Zeitsch Nahr Genussin 1905, 10, 528, Gulewitsch, Zeitsch physiol Chem 1906, 50, 201, 51, 258; 52, 527). Compare, however, Kutscher (ibid. 1907, 50, 445, 51, 545)

When carnosine is hydrolysed by boiling with barium hydroxide it yields histidine, the other product of hydrolysis being probably

alanıne

$$C_9H_{14}O_3N_4 + H_2O = C_6H_9O_2N_8 + C_3H_7O_2N$$

According to Gulewitsch (Zeitsch physiol Chem. 1911, 73, 434–446) three-quarters of the carnosine nitrogen belongs to instiduce and the rest to  $\beta$ -alanine.

That carnosine is  $\beta$ -alanylhistidine is shown by its synthesia from histidine and  $\beta$ -iodopiopionyl chloride (Baumann and Ingvaldsen, J Biol Chem 1918, 35, 263). This is further proved by its behaviour towards 2 4 5-timitrotoluene. It has been synthesised by condensing  $\beta$ -nitropropionyl chloride with histidine methyl ester and reducing the resulting products with stannous chloride and hydrochloric acid (Barger and Tutin, Bio-Chem J 1918, 12, 402)

The carnosine content of flesh extracts may be estimated by two colormetric methods which control one another, the one depends on the colour yielded by the histidine complex in the carnosine molecule with diazobenzenesulphonic acid (Werss and Sobolev, Biochem Zeitsch 1913, 58, 119–129), the other on the violet colour yielded by carnosine when boiled with cupric hydroxide, Furth and Hryntschak (Biochem Zeitsch 1914, 64, 172–194)

M. A W

CARNOTITE, A hydrated vanadate uranium and potassium, occurring abundantly as a bright canary-yellow powder disseminated in sandstone in western Colorado C Friedel and E Cumenge (Compt rend 1899, 128, 532), by whom it was first described and named (after A Carnot), expressed the composition by the formula  $2U_2O_3V_2O_5K_2O3H_2O$  Detailed analyses were made by W.F. Hillebrand (Amer. J. Sci 1900, 10, 120), and he concluded that the This view mineral is not simple, but a mixture is, however, not borne out by T Crook and G S Blake (Mm. Mag 1910, 15, 271), who find that the material, both from Colorado and from South Australia, is definitely crystallised in minute plates possessing orthorhombic symmetry and with characters analogous to those of autunite [Ca(UO<sub>2</sub>)<sub>2</sub>P<sub>2</sub>O<sub>8</sub>,8H<sub>2</sub>O] The original material was found by M and Mme Curie to contain radium and polonium.

The carnotite-bearing sandstones occur over an extensive tract of country on both sides of the Colorado-Utah boundary, the nichest deposits being in the neighbourhood of La Sal Creek, Roc Creek, and Paradox Valley, in Montrose Co., and in San Miguel and Dolores Cos., in southwest Colorado, other deposits are met with near Coal Creek in Rio Blanco Co. and Skull Creek in

Routt Co in north-west Colorado, and at Mauch Chunk, Pennsylvania. Most of these deposits are in Jurassic (La Plata) sandstone, but some of those in the N.W. are in sandstones of Cretaccous age. They have been mined at several spots, and a plant for the extraction of uranium, radium, and vanadium salts has been erected near Cedar in San Miguel Co., Colorado; but the sandstone ofe is of a low grade, and the difficulties of transport are considerable. An average sample of marketed ore assayed  $U_{\rm s}O_{\rm s}$  11-49 p.c. and  $V_{\rm s}O_{\rm s}$  6-40 p.c. (R. B. Moore and K. L. Kithil, U.S. Bureau of Mines, Bull 70, 1913. C. L. Parsons and others, while Bull 104, 1915)

In South Australia, at Radium Hill, near Olary, carnotite occurs in films on the jointplanes of a large deposit of ilmenite rutile, and

magnetite.

A closely allied mineral, named ferganite (I A Antipov, 1908, Abstr in Jahrb. Min 1909, ii, 38), is a hydrated vanadate of uranium and lithium. It occurs, together with other uranium minerals, in prov. Fergana in Russian Turkestan.

L. J. S.

CARO'S ACID. Monopersulphuric acid (v. Sulphur).

carobs are found to possess the following percentage composition water, 9-13; introgenous compounds, 5-7, cellulose, 8-12, fat, less than 1; ash, 2, sugars (dextrose and sucrose in variable proportions), 30-43; phosphoric oxide, 0·16-0 24 (Balland, J. Pharm Chim. 1904, [6] 19, 569).

CARONE v KETONES.

CARONY BARK v CUSPARIA BARK

CAROTIN. The colouring matter of Daucus carota (Linn) Found also in the leaves of plants to the extent of 0 1 to 0 2 pc (Arnaud, Compt rend 1890, 109, 911) in full vegetation, and to a slight extent in etiolated leaves (Immendorf, Chem Zentr 1890, 1 163), and in the tomato; is present in maize, squash, orange-peel, flax seed, mustaid seed, black sesamé seed, palm oil, but not in rape seed, white sunflower seed, turnip, safflower, cotton seed or turmeric. It forms the colouring matter of yellow and orange pollen, the oily drops on the surface of the pollen of Verbascum thapsiforma contain not less than 66 pc of carotin, and in the animal kingdom it occurs as the colouring matter of milk-fat, hence in butter and the body-fat, corpus luteum and skin secretions of the cow, it constitutes the yellow lipochrome of bloodserum and is found in Diaptomus bacillifer (Blanchard, Compt rend 1890, 110, 292) It can also be extracted from stinging-nettle leaves with light petroleum (Willstatter and Mieg, Annalen, 1907, 355, 1). Carotin crystallises in copper-coloured leaflets which appear red by transmitted light, and have mp 167.8°. It is insoluble in water, readily soluble in carbon disulphide or benzene, sparingly so in alcohol, ether, petroleum, or chloroform Carotin is able to decompose carbon dioxide with formation of oxygen, and this probably explains why etiolated leaves can, under certain conditions, generate

oxygen (Kohl, Chem Zentr 1906, 11 442) Carotin is an unsaturated hydrocarbon having the formula  $C_{40}H_{56}$  (Willstatter and Mieg. lc); it combines with iodine to form the iodide  $C_{40}H_{56}I_2$ , which crystallises in rosettes

of dark-violet prisms, and decomposes at  $140^{\circ}-170^{\circ}$ ; and the  $\mathit{rodide}$   $\mathrm{C_{40}H_{56}I_{3}}$ , which forms dark-violet plates and decomposes at 136°-137°. The bromo-compound C<sub>40</sub>H<sub>36</sub>Br<sub>23</sub> decomposes at 171°-174° (Willstatter and Escher, Zeitsch. physiol Chem 1910, 63, 47) Carotin dissolves slowly in concentrated sulphuric acid, forming an indigo-blue solution from which the hydrocarbon is precipitated in green flakes on the addition of water; it absorbs oxygen to the extent of 34 3 p c of its weight, yielding a colourless substance. When carotin is oxidised with a small amount of acetic acid solution of chromic anhydride, and the temperature kept below  $30^{\circ}$ – $40^{\circ}$ , an oil  $C_{40}H_{66}O_8$ , or  $C_{40}H_{60}O_5$  is obtained, and this does not solidify at  $-104^{\circ}$ (Euler and Nordenson, Zeitsch. physiol. Chem. 1908, 56, 223)

Carotin has no appreciable toxic acid, quantities of 0.2 gram injected into guinea-pigs had no effect on the health of the animals.

CAROUBIN and CAROUBINOSE. According to Effront (Compt rend. 1897, 125, 38), the seeds of Ceratonia siliqua, used as a cattle food in Portugal, contain a carbohydrate, caroubin, which, when purified, forms a white spongy friable mass, giving with water a thick syrup or jelly By hydrolysis, it yields Caroubinose  $(C_6H_{12}O_6)$ , which has about the same reducing power as dextrose Is probably identical with mannose (Van Ekenstein, Compt rend 125, 197) (v Carbohydrates)

**CARPAINE**  $C_{14}H_{25}O_2N$  occurs in papaw leaves (*Carica Papaya* (Linn)) It forms glancing prisms, in p. 121°,  $[\alpha]_p + 21^\circ$  55′, and has a bitter taste (Greshoff, 1890, Med. s'Lands Plantentum, Java, No. 7, 5; van Rijn, Arch Pharm. 1893, 231, 184, 1897, 235, 332). According to Baiger (Chem. Soc. Trans 1910, 97, 466) it is a lactone of carpainic acid C<sub>14</sub>H<sub>27</sub>O<sub>2</sub>N, and may be written

CARRAGEEN. Carageen, Carragheen, or Carragaheen (Fr Gaëmen) Irish or Pearl moss. An alga (Chondrus crispus) found on all the western coasts of Furope and on the east coast In Ireland it is collected, of North America dried, and bleached in the sun Found in commerce as flat, dry, yellow-white pieces 2 to 3 inches long, usually branching, of a very faint odour and mucilaginous taste, swelling in water like gelatine Almost completely soluble in boiling water; the solution sets to a jelly on cooling Contains a high percentage of nitiogen, and is of value as a food in diseases of lungs and general weakness, usually taken dissolved in hot milk. It has also been applied as a size and for stiffening silk, also occasionally in making jellies.

Sometimes found largely adulterated with Gigartina mamillosa, which is less valuable and recognised by its stalked fruits and channelled stem (v. Algæ and Irish Moss Gum, Art Gums.

CARRAWAY OIL v. OILS, ESSENTIAL.

## CARRIAGE VARNISH v. VARNISH.

CARROT, the tap root of Daucus carota (Linn). Several varieties exist, and the roots are employed as cattle food and as a table vegetable Kellner gives as the average composition

According to American analyses, the average composition of the edible portion, as a table vegetable, is

The Belgian white carrot is often used for the field crop, whilst the red carrot is preferred for table purposes The reddish colouring matter of the carrot, carotin, is sometimes used in colouring butter and cheese. It has the composition  $C_{40}H_{56}$ , and can be obtained by rasping the roots, precipitating the juice with lead acetate, with which the carotin forms a lake This is dried and extracted with carbon disulphide The solution is evaporated to dryness and repeatedly extracted with cold petroleum spirit in order to remove the phyto-The residue is then dissolved in carbon disulphide, from which the carotin is precipitated by the addition of alcohol—It can then be recrystallised from benzene in the form of rhombic plates with a metallic lustre, blue by reflected, orange-red by transmitted light The substance which Husemann discovered in the carrot and called hydrocarotin is a mixture of sitosterol and stigmasterol, and consists of the same phytosterols as are found in other plants

(Beschke, Bei 1914, 47, 1853) (v Carotin)
As a farm crop, carrots are particularly valued for horses and dairy stock. They are best grown on light land, free from stones and н 1. weeds and well tilled.

CARROT GUM v Gums CARVACROL v Gamphors, also Oils, Es SENTIAL

CARVEOL v Camphors.

CARVOL v Camphors, also Oils, Essential

CARVONE " KETONES

CASCARA SAGRADA, BP, US.P. The dued bark of the Rhamnus Purshiana (DC), a small species of thorn tree inhabiting the Pacific slopes of the Rocky Mountains (For botanical characters, see Hooker (Flora Bor Amer 43), Prescott (Amer J Pharm 1879, 165), Moller (Pharm J [3] 14, 467), Moss (ibid. [3] 649, 19).) Cascara sagrada had been long known as a cathartic to the Indians and trappers of Califorma, when it was later introduced into American and European medicine Its physiological action is similar to the other and better-known Rhamnus frangula (Linn.), but the Purshiana is the more esteemed (Cf. Kennedy, Amer J Pharm 1885, 496)

Cascara bark has been examined by Prescott (*ibid* 1879, 165), Limousin (J Pharm. Chim. [5] 6, 80); Wenzell (Pharm Rundsch. 1886, 79); Meier and Webber (Pharm J. [3] 18, 804), Schwabe (Arch Pharm. 226, 569); Zeig (Pharm. J. [3] 20, 173); Le Prince (Compt. rend. 115, 286), Dohme and Engelhardt (Amer. Chem. J. 20, 534); but the statements made regarding it

are very conflicting, for a summary of which see Jowett (Rep. Amer. Pharm. Assoc. 1904, 1), who has also made an exhaustive examination of the bark. He has confirmed the presence of emodin together with a small proportion of an isomeric substance which may be identical with the iso-emodin obtained from Rhamnus frangula by Thorpe and Miller (Chem Soc. Trans. 61, 6) Glucose is also present, and a substance which, on treatment with acid, yields syringic acid. No evidence was obtained of the existence of chrysophanic acid or chrysarobin or of glucosider yielding on hydrolysis emodin, chrysophanic acid, or rhamnetin. The cascarine of Le Prince and the purshianin of Dohme and Engelhardt are probably impure products Emodin is not the active principle of the drug, and no clue could be obtained as to the nature of this principle (Compare Nouveaux Remèdes. 1909, 471, where the active principle is stated to be precipitated by lead acetate and ammonia, after treatment with lead acetate alone ) It is a light yellow powder of bitter taste, probably a glucoside According to Tschirch and Monikowski (Arch Pharm. 1912, 250, 92) the patent cascara preparation 'peristaltin' consists of a mixture of glucosides, hydrolysed by steam to rhamnose, chrysophanic acid, emodin methylether and cascarol C15H10O5, yellow needles, m p 218°. Cascara bark contains about 2 p c of fat and a small proportion of a volatile oil The constituents of  $\tilde{R}$ . Californica appear to differ very little from those of R Purshiana For methods for the estimation of the active constituents of Cascara sagrada, see Warin (J Pharm chim. [6] 22, 12), whose coloring method is approved by Tschirch and Pool (Arch Pharm 1908, 246, 315) For differential tests of Cascara and Rhamnus frangula, see Kroeber (Schweiz. Woch Chem Pharm 1910, 48, 377); Beal and Okey (J. Amer Chem Soc. 1917, 39, 716) For standards for the evaluation of liquid extracts of Casacara and Rh frangula, see Apot Zeit 1913, 39, 579).

CASCARILLA BARK, Cascarilla (BP)
The bark of Croton Eluteria (Benn), or 'seaside
balsam,' a cuphorbiaccous tree growing in the

West Indies

Duval obtained from it a substance, cascarilline C<sub>12</sub>H<sub>18</sub>O<sub>4</sub>, in p 205°, which Alessandri (Arch Pharm 220, 690) also isolated by extracting the bark with ovalic acid and precipitating the extract with ammonia. This substance dissolves in hydrochloric acid, producing a rose-coloured solution, which changes through purplish-red, violet, and green to a sky-blue without the addition of water. The drug also contains betaine (Naylor, Pharm. J. 59, 279), and 15 p c. of an essential oil (q.v)

CASCARILLA OIL v OILS, FSSENTIAL.

casein. Casein occurs in the milk of mammals, and is the principal protein of cow's milk, which, up to the present, is its only commercial source. Cow's milk contains, on an average, about 3 pc of casein, which exists in a suspended or colloidal condition, probably as a lime compound in combination with calcium phosphate, giving the milk its opaque appearance. This can easily be proved by allowing the milk to pass through a filter of unglazed porcelain, when a white mass of casein and fat will be found on the surface of the filter, and the

filtrate will contain the lactose, soluble albumen, and soluble salts of milk,

Preparation of pure casein. Separated milk is diluted to about five times its volume with water and sufficient acetic acid added to give 0.1 p.c. of acid in the solution. The casein is precipitated, carrying down with it the fat. The curd is washed about ten times by decantation, collected on a cloth filter, washed, and pressed as dry as possible. It is then dissolved in the least quantity of ammonia, and filtered. The filtrate is precipitated with acetic acid, washed, and redissolved in ammonia. This process is repeated three or four times. The curd is then rubbed up in a mortar several times with 80 p.c. alcohol, and finally with absolute alcohol. It is next similarly treated with ether, and extracted with ether in a Soxhlet apparatus, and the casein finally dired at 100°C.

Properties — Thus prepared, casein is a white, amorphous body without taste or smell. It is very sparingly soluble in water, and quite insoluble in alcohol and ether. Its most probable percentage composition is: C, 53·13; H, 706; N, 15·78, P, 0.86, O, 22·40; S, 0.77.

Casem is a phospho-protein. Its molecular weight is stated to be about 1135. It is a weak dibasic acid, forming acid and neutral salts with alkalis. In dilute alkaline solutions, it has a lawo-rotatory action on polarised light. At the ordinary temperature, I gram of casem combines with 9 c c. of N/10 hydrates or carbonates of alkalis, and with nearly 7 c c. decinormal solutions of common acids. It is soluble in caustic alkalis, and in the carbonates, bicarbonates, phosphates, &c, of the alkali metals, and is precipitated unchanged by addition of sufficient acid to neutralise the alkali

Casein is soluble in strong hydrochloric acid, giving a fine violet colour, and is also soluble in dilute hydrochloric acid containing 0.2 p.c. of the fuming acid, but is insoluble in 2 or 3 p.c. acid

Preparation of industrial casein. There are many methods of preparing casein, but on the large scale these are generally limited to three:

(1) Precipitation by means of acid—usually sulphuric, though acetic, phosphoric, lactic, hydrochloric, &c., are also employed.

(2) Curdling by means of rennet(3) Curdling by 'self-souring.'

A very interesting method of preparation is by means of electricity, but so far this does not seem to have progressed beyond the laboratory stage. The method, however, will be described later. Whichever method be employed, the first essential is a machine-separated milk, containing the smallest possible quantity of fat Milk taken from cows shortly after calving, and containing colostrum, is unsuited for casem manufacture. It gives a soft curd which squeezes through the filter cloths, and holds a great deal of water. It has sometimes been found necessary to throw away a whole batch as the curd has been quite unworkable.

Also, if the milk has been pasteurised at such a high temperature that some of the lact-albumin becomes insoluble, the curd will give a great

deal of trouble.

The milk from the separators is run into a tank of suitable dimensions, and brought to the required temperature by means of steam.

CASEIN 93

Precipitation by sulphuric acid.—The requisite quantity of sulphuric acid, previously diluted with water in a wooden or enamelled iron bucket, is then introduced, and the whole well agitated, when the milk will gradually lose its opaque appearance and become clear with separation of the curd. As soon as the whey is yellow and transparent, the agitation is stopped, and the curd falls to the bottom of the tank. The whey is run off, the curd allowed to drain and washed with cold water until all the whey has been separated. The bulk of the water is then removed by means of a suitable press or by a centrifugal machine.

Curdling with rennet —The amount of rennet required depends upon the acidity of the milk and the temperature. The more acid the milk and the nearer the temperature approaches 106°F., the less is the rennet required. At 118 5°F, and at 86°F, the curdling power of rennet is only about 70 pc of that at 106°F

If the acidity of the milk be 15°, about 2½ oz. rennet should curdle 100 gallons of milk in about I hour. When the acidity reaches 30°, half this quantity will suffice The requisite quantity of rennet is diluted with water and thoroughly stirred into the milk at a temperature of about 106°F The milk is then allowed to rest until it has 'set' like a junket The mass is cut up into small cubes by special knives, and then agitated gently at first, and afterwards vigorously while being very slowly heated up to about 120°F by means of free steam or a steamjacket The curd will then be in hard grains or nodules at the bottom of the tank. The whey is rapidly run off, the cuid washed well with cold water, and pressed

Curdling by 'self-souring.'-When milk is exposed to the air, it becomes acid. This change is effected by the Bacillus acidi lactici, which converts the lactose into lactic acid sufficient acid has been formed, the curd separates from the whev On the large scale, the separated milk is run into a tank, the temperature brought to and maintained at about 98°F, at which point the growth of the bacilli is most After some hours, according to the freshness of the milk, a sour smell will be noticed and the milk will begin to curdle When the right point has been reached, steam is turned on and the mass agitated in order to make the curd firm, and in a suitable condition for washing The whey is then run off and the and pressing curd finished in the usual manner

Drying of the curd —When as much moisture as possible has been removed from the curd, it is broken up and dried. This may be effected either by spreading on trays in heated drying-rooms fitted with ventilating appliances to facilitate the removal of the moisture, or in a drying cylinder heated from inside or outside or both, and rotated to keep the contents in constant motion. Another method of drying the curd is by means of a Passburg vacuum drier, consisting of a cast or wrought-iron case, which can be hermetically closed by doors. A vacuum is produced by an air pump, and the moisture is vapourised at a low temperature so that drying is soon effected.

Whatever process be adopted, the temperature must be carefully watched, as a few degrees too much heat will 'burn' the casem to a rich brown colour, and if the temperature be too low the curd will take a long time to dry, and may become sour and mouldy.

The dry casein is then ground to a fine

powder, when it is ready for the market

Curdling by electricity.—Although this process is so far not a commercial one, it is sufficiently interesting to warrant a short description milk is heated to 120°F, and a porous cell, containing dilute milk, is placed in the centre. A current of electricity of sufficient tension is passed, and the soluble elements, the phosphates in particular, are decomposed anode is placed in the milk and the cathode in the porous cell Phosphorie acid is then liberated in the milk, and causes the precipitation of the casein, while the liquid in the porous cell becomes strongly alkaline. The chief difficulty lies in the choice of electrodes—the cathode in the porous cell can be carbon, but the anode must be of platinum, which makes the original cost very heavy Any of the common metals are attacked in a very short time, and carbon throws off small particles which cause the precipitated cuid to become black. It is, of course, possible to have both electrodes of carbon in porous cells, but it is found that the curd soon fills up the pores and the action becomes very slow

Properties.—Industrial casein, prepared with sulphuric acid, will dissolve in the following quantities of alkalis or alkaline salts, though larger quantities are always used in the various industries—

 Sodium bicarbonate
 3½

 Sodium carbonate
 2¼

 Sodium hydroxide
 1¼

 Sodium silicate 140°Tw.
 10

 Sodium arsenate
 20

 Sodium sulphite
 9

 Strontium hydrate
 20

 Trisodium phosphate
 8

 Ammonia, 880
 2½

Sodium tungstate

Borax

Per cent.

Solutions of the hydroxides of potassium, sodium, lithium, and ammonium of equal concentration, dissolve casein at approximately the same rate. Solutions of the hydroxides of the alkaline earths dissolve casein much more slowly, strontium hydroxide dissolves it most rapidly, calcium hydroxide more slowly, and barium hydroxide more slowly still.

Rennet casein is insoluble in carbonate and bicarbonate of soda, partly soluble in borax and ammonia, and soluble in trisodium phosphate and lime

Its solution in alkalis does not coagulate on heating, but is precipitated by acids and many metallic salts. When acted on by formaldehyde and certain other bodies, casein is rendered insoluble. Advantage is taken of this fact in various industries to give a waterproof coating of casein.

Pure casein contains only a very small amount of ash, but industrial casein may contain up to about 7 p.c. or 8 p.c.—rennet-prepared

 $<sup>^1</sup>$  The 'degree of acidity' of milk is the number of c c, of N/10-NaOH sol, required to neutralise the acid in 100 c.c.

CASEIN.

casein containing the most, and 'self-soured' casein the least.

For the analysis of commercial casein v. Hopfner and Burmeister, Chem. Zeit. 1912, 36, 1053; Analyst, 1912, 493.

Casem is employed in a Uses of casein. large number of industries, and its use is increasing every year, in fact, there are but few industries in which casein, in some shape or form, is not or cannot be employed with advantage

The greater quantity of imported casein comes from the Argentine Large quantities also are received from France and Germany, and smaller quantities from Holland, Belgium, and tne United States

Casein is used in paste and dry distempers, in cardboard manufacture, paper glazing, leather dressing, soap manufacture, cotton sizing, diessing for textiles, finishing, waterproofing, calico printing, cask glazing; imitation bone, ivory, tortoiseshell, amber, ebony, &c picture mouldings, sealing for bottles, cabinet and planoforte making, emulsions, fining wines, boot polishes, waterproot cements, polishes, films, diabetic and protein toods

Alimentary casein. Some of the best-

known food preparations of casein are Plasmon —This is a cream-coloured powder, somewhat coarse in texture It is a soda compound of casein, and is partially soluble in water. It contains nearly 2 pc of lactose (milk-sugar)

and about 70 p c of proteins

Sanagen —A fine white powder, completely soluble in water. It is a compound of casein with about 5 pc of sodium glycerophosphate made by a special process. It contains about 76-78 p.c. proteins, and is practically free from lactose, and so is specially recommended for diabetic patients.

Sanatogen -A white powder soluble in It consists of casein with about 5 pc. of sodium glycerophosphate, and is free from fat and lactose

Dr. Reigel's milk protein —A soda compound of casein, soluble in water. According to the patent specification, the curd is precipitated from the milk by means of ethyl-sulphuric acid

Eucasein - A casein-ammonia compound, soluble in water

Nutrose.—A soda compound of casein.

Casein solids. Galalith —Insoluble treated with formaldehyde and subjected to heavy pressure, gives plates of great hardness, and by blending pigments with the casein, imitations of bone, ivory, ebonite, &c., are obtained. The products are known as Galalith (from the Greek gala, milk, and lithos, stone) It is prepared from a specially pure casein, precipitated by rennet, from sweet milk freed from fat by centrifuging. The curd is drained and dried, forming lumps 8-12 mm. thick These are ground to a gritty meal, which is moistened with water, coloured as required, worked up by steel helices to a plastic mass, which is then hot-pressed in hydraulic presses. The pressed cake is hardened in a solution of formaldehyde, the length of treatment varying from 2 to 30 weeks, according to the thickness of the articles, whereby the casein acquires the qualities of horn (Bonwitt, Zeitsch. angew. Chem. 1914, 27, 2). The following articles are

made from galalith brush-backs, combs, eigar and cigarette holders, penholders, electrical fittings, umbrella and stick handles, buttons, dominoes, &c Being uninflammable, casein possesses great advantages over celluloid. It is a bad conductor of electricity, and is therefore suitable for electrical fittings A plate of galalith 1 inch thick, is impervious to a current of 16,000 volts. The sp gr. is 1317-135 (celluloid 134-140) The hardness is 25 (celluloid 2). It resists only liquids and acids, but is swelled by alkaline liquids — Its chief drawback is its hygroscopic nature, it absorbs 30 pc of its weight in 12 days. It is not elastic like celluloid, and breaks when slightly bent. On account of its tendency to flake it cannot be worked so well as celluloid, and sheets cannot be prepared less than 2 mm. in thickness (Bonwitt, lc).

Lactoform — This is a patented preparation of casein resembling horn, giving, according to the specification, a hard and tough article which does not swell or materially soften when immersed in water. It is prepared by adding various salts to solutions of casein, thus, by adding a solution of lead acctate to a solution of casem, a coagulum is produced from which lactoform, resembling horn, may be obtained, while to manufacture a substitute for ivory, carbonate of soda is added to the casein solution and precipitated by sugar of lead, thus forming carbonate of lead in the mass obtained By the use of chromium compounds, bluish-green, also yellow and orange lactoform, is obtainable; by the action of iron, just-coloured, and in combination with tannic acid, slate and blackcoloured, lactoform may be produced. These are merely intermediate products, and are not claimed as a new invention, but are converted into lactoform by treatment with formaldehyde

For glues and sizes, casem is dissolved in borax, soda, or ammonia, and the resulting coating may afterwards be rendered insoluble by formaldehyde, or the casein may be dissolved in lime, which will give a waterproof coating when dry.

Films.—Schmidt has patented a process for making casein films as follows: 100 grams of casein are dissolved by heating in 1 litre of water containing 8 grams of 25 p c. hydrochloric acid: 10 grams of 40 pc. formaldehyde and a small quantity of glycerol are added, and the solution poured on glass plates to dry. He states that films made from alkaline solutions of casein are not impermeable to moisture.

Schmidt patents the following process: 100 grams of casein and 1.5 grams caustic soda are dissolved in I litre of water, to which are added about 15 grams of a 40 p c. solution of formaldehyde, or 100 grams of casein and 10 c c. of a 10 p c solution of ammonia are dissolved in 2 litres of water, to which are added about 30 grams of a 40 p.c solution of formaldehyde. Either of these he states, gives insoluble films on drying, which may be used for photography, surgical bandages, paper-coating, &c.

Buss, in his patent for photographic films, proceeds as follows 500 to 700 grams dry casein are heated with 9 litres of water to 35° or 50°, mixed with 250 grams of citric acid dissolved in 1 litre of water, and the heating con tinued until the vhole has dissolved. Then 100

CASSIA. 95

or 200 grams of glycerol are added, and the liquid filtered. It is applied by machine to plain or baryta paper, and allowed to dry. It is next rendered insoluble by being drawn through a 5 or 7 p.c. solution of ammonium or sodium chloride, and finally sensitised in a silver nitrate bath, ranging between 4 and 12 p c. in strength

Scaling for bottles.—For this purpose, Joirgnot has patented the following preparation: Casein is treated with a hot aqueous solution of sodium phosphate, ammonium fluoride and glycerol, the whole being then heated in the water bath until it is homogeneous and viscous. This product is employed for sealing bottles of metal, glass, porcelain, &c, by applying it in a warm state to the edge of the vessel or cover, and then treating it with formaldehyde

Fining of wines.—For this purpose, a readily soluble sodium compound of casein is used is dissolved in 10 or 20 parts of water, the 5 p c. solution effecting a somewhat slower though more complete clarification. The casein compound has the advantage that it is very much cheaper than albumen, and is free from all flavour such as is found in even the purest Moreover, if in fining wines with white of egg, albumin or gelatin, too large an excess in proportion to the tannin be used, this excess remains in solution, and may lead to turbidity and alteration in the wine; whereas the casein, even when added in considerable excess, is completely eliminated, being coagulated by the acid in the wine

Casem water-paints —Casem in fine powder is intimately mixed with sifted slaked lime and incorporated with the pigments The proportion of lime to easein is very important if too little lime be used, the mixture will, after a short time, especially if exposed to air, become insoluble in water, and if too much lime be used, adverse results will follow The brushes will be rotted and the paint will be hable to flake off The next matter of importance is the proportion of lime and case to be incorporated with the pigment. If too little be used, the paint will dust off and will not stand washing, and if too much be used, the coating will be brittle and flake off

In selecting the pigments, it is necessary to employ only those which are unaffected by lime. The most suitable are

For White.
,, Yellow.
Whiting, China clay, zinc oxide.
Ochre, Hansa yellow, naphthol
yellow, chrome yellow.

,, Brown. Umber, sienna ,, Black. Carbon blacks. ... Red. Oxides, lithol red.

,, Red. Oxides, lithol red .. Blue. Ultramarine.

Green earth, ultramarine green, lime green.

The ingredients must be thoroughly incorporated and kept in air-tight packages. The paint is prepared by mixing the powder with cold water, and is ready for use in a few minutes. When thoroughly dry, the surface is washable.

L. M. N.

CASHEW GUM v Gums. CASIMIROINE  $C_{22}H_{14}O_6N_2(OMe)_2$ , m.p.  $196^\circ-197^\circ$ , and casimiroedine  $C_{17}H_{24}O_6N_2$ , m.p.  $222^\circ-223^\circ$ , occur in minute amount in the seeds of Casimiroa edulis (La Llave et Lejarza) a reputed

Mexican drug. The alkaloids are physiologically mert (Power and Callan, Chem Soc. Trans. 1911, 99, 1993, who give other references)

CASSAVA. The root of Manshot palmata (Muell), 'sweet cassava;' or of Manshot utilissima (Pohl), 'bitter cassava' These plants are widely cultivated in the tropics for the sake of their starchy edible roots. From them the substance tapicca is prepared. The starch is also known as mandice starch, Brazilian, Bahia, Rio or Para arrowroot, and by other

The giains of starch are grouped in twos, threes, or fours, the separate grains have usually one or two flat surfaces, but, on the whole, are rounded and show delicate concentric strictions and a distinct follows:

Cassava root contains a volatile poisonous substance (Fermin, 1764), which proved to be hydrocyanic acid (Boutron-Chailaid, 1836, and Francis, Analyst, 1878, 2, 4). The latter observer found 0 0168 p.c. of prussic acid in sweet cassava root and 0 0275 p.c. in the bitter root. He found 594 p.c. and 614 p.c. water in the roots, and about 30 p.c. of starch

Collens (Bull. Dept. of Agric. Trimidad and Tobago, 1915, 14, 54) found that the freshly-dug roots of the sweet and bitter varieties contain 0.0048 p.c. and 0.053 p.c. of hydrocyanic acid respectively, the leaves 0.0162 p.c. and 0.041 p.c., and the peelings of the stems 0.043 p.c. and 0.041 p.c. respectively

Leuscher (Zeitsch offentl Chem 1902, 8, 10), as the mean of six analyses of the fiesh root, found:

Water Protein Fat Starch Sugar Crude fibre Ash 70 25 1 12 0 41 21 44 5 13 1 11 0 54 He states that the yield, on well-irrigated land, may amount to 15 or 16 tons per acre

Tapioca is made from the starch of cassava by stiring the damp product on a hot iron plate until it agglomerates into the irregular, spherical masses familiar as tapioca. By this treatment, many of the starch grains become changed, being converted into shapeless, translucent

According to American analyses (Wiley, Bull 13, U.S. Dept. of Agric. Div. of Chem. 1898), tapioca contains—

Water Protein Fat Carbohydrates Fibre Ash 11 23 0 37 0 03 88 16 0 12 0 09

Owing to the exceptionally low demands of cassava for mineral matter, it has been recommended as a crop particularly well adapted for cultivation on poor sandy soils, as it feeds very largely upon air and water (Ewell and Wiley, Amer Chem J 1893, 15, 285)

H. I.

CASSEL EARTH OF VANDYKE BROWN v. PIGMENTS.

CASSEL YELLOW. Lead oxychloride. A pigment also known as Turner's Yellow, Montpelier Yellow, Mineral Yellow, and Patent Yellow.

CASSELMAN'S GREEN. A pigment made by mixing boiling solutions of copper sulphate and an alkaline acetate

CASSIA. The bark of Cinnamomum cassia (Blume), a tree belonging to the order Lauraceæ or true laurels. It is grown in China, Java, &c. C. cassia furnishes a bark which is much like cinnamon [C. zeylanıcum (Nees)], but thicker, coarser, stronger, less deheate in flavour, and

96 CASSIA.

cheaper; hence it is frequently used to adulterate cinnamon. The bark is stripped off the branches, when it rolls up in the form of quills It is said to be preferred to cinnamon itself by some chocolate makers on account of its stronger flavour. The unopened flower-buds are also sold under the name of 'cassia-buds,' and possess similar properties to the bark. Other species of the genus afford aromatic barks, eq C culilawan (Blume), native of Malaya and China, the bark of which has a flavour of cloves, tamala (Nees), which occurs in sub-Himalayan India as far as Burma, and provides a bark that is used as a substitute for or adulterant of true cinnamon: C Burmanni (Blume) is the source of the bark exported from Sumatra The bark known as clove-cassia is obtained from Dicypellium caryophyllatum (Nees), which is also a

tree of the laurel family growing in Brazil

The employment of cassia as a sophisticant of the more esteemed and more costly bark of cinnamon when in the ground state may usually be detected by adding tincture of rodine to a decoction of the powder; when, since cassia contains more staich than cinnamon, the substance will turn blue, but, where inferior qualities of the true cinnamon have been examined, their constitution has approached that of the coarser bark of cassia, and this test has proved uncertain (cf O. Hehner, Analyst,

1879, 225-228)

Cassia bark is thicker, redder, stronger to the taste than cinnamon. The powder under the microscope exhibits a general resemblance to cinnamon, but it will be seen that cassia differs from cinnamon in its coarser structure, and especially in the greater size and number of its starch corpuscles

Cassia yields by distillation an essential oil largely used in perfumery, especially for producing the scent in brown Windsor soap (v Oils, ESSENTIAL)

CASSIA OIL v OILS, ESSENTIAL

CASSIA TORA (Linn ) or TAGEREY-VEREY. This plant produces grains known in the East Indies, Arabia, and Japan as Tora, but which are known in Pondicherry and other parts of Hindostan as Tagerey-verey It is regularly used as a component of the indigo vat in dyeing, apparently serving the same purpose as the bran-madder or molasses used in Europe

The native dyers use the Tagerey-verey in the following manner. To dye about 200 yards of cloth, 111 lbs of the grain are steeped in from 5 to 6 gallons of cold water, and then boiled for four hours The grains are swellen and softened by this treatment, and the water becomes thick and gummy The whole of this is added to the indigo vat, and after standing for fifteen hours, the vat is ready to dye.

CASSITERITE or TIN-STONE. Native tin oxide, SnO2, containing 78.6 pc. of tin, and practically the only ore of the metal It varies much in appearance—the colour ranges from white to black, being usually dark brown-and its most characteristic feature is its high sp.gr. of 6.8-7.1; hardness, 6-7, easily scratching Crystals are tetragonal, with very characteristic knee-shaped twins; their faces are often very brilliant with an adamantine lustre There is an absence of cleavage; and this readily helps to distinguish the mineral from blende.

Crystals are, however, not very common, the mmeral being more frequently found as disseminated grains and water-worn pebbles. Grains of cassiterite in a concentrate panned from a river sand may be readily detected by a very simple characteristic test. When placed on a zinc plate, or in a basin with fragments of zinc, and dilute hydrochloric acid poured over, any grains of tin-stone soon show a shining white coating of metallic tin. Veins of tin ore usually occur at the junction of granitic masses and slates, and the ore also impregnates the rocks bordering the veins. Such ores are known as vein-tin, to distinguish them from the stream-tin of alluvial deposits. A compact variety with a radially fibrous and concentric structure and sometimes a botryoidal surface is known as wood-tin. The ore as prepared for market is known as black-tin, and in South America as barilla. The principal tin-producing countries are the Federated Malay States, the islands of Banka and Billiton in the Dutch East Indies, Siam, China, Australia (Queensland, New South Wales, Western Australia), Tasmania, Bolivia, Cornwall, and more recently South Africa (Transvaal and Swaziland), and Northern Nigeria. The production of tin ore amounts to about 100,000 tons per annum CASSIUS, PURPLE OF, v. GOLD J. L. S.

CASTILE SOAP v SOAP

CASTOR or CASTOREUM. (Bibergeil, Ger) A name given to a secretion of the beaver (Castor fiber), contained in pear-shaped cellular sacs (found near the genital organs of the male and female animal), which are cut off and dried, to prevent the skin being affected by water It is a substance analogous to civet and musk, and is of the consistency of thick honey and unctuous to the touch. The interior substance is solid, of a dark-brown or black colour. It has a faint smell, softens when heated, but becomes brittle When chewed, it sticks to the teeth when cold It has a bitter acrid taste, and a like wax fætid, penetrating smell. Its fracture is shining, and when the substance is genuine it shows fragments of membrane indicating organic structure

Three varieties are said to exist, the Russian, Bavarian, and American or Canadian, the firstnamed variety containing about 2 pc of the volatile oil, while the last contains only 1 p.c.

Wohler obtained phenic and benzoic acids and salicine from it Lehmann found bile in a fresh sample together with alkaline urates, sebates, and an albuminoid substance. Laugier, Brandes, Batka, Bouillon Lagrange, and Hildebrant have also examined it By extracting it with six times its weight of alcohol a colourless substance, castorin, has been isolated Used in medicine and perfumery.

CASTOR OIL is obtained from the seeds of Ricinus communis (Linn ) and R. sanguineus

The former plant appears to be indigenous to Africa (Egypt), but has been transplanted to India at a very early stage of history, so that some writers consider the plant indigenous to India itself The plant grows in enormous quantities in all tropical and sub-tropical countries, and is even found wild to day in South America, notably in Paraguay and the

The most important sources of castor seed

are East India, Java, the Mediterranean counties, Mexico, and the United States of America, whilst the cultivation of the plant has of recent years been developed in the Egyptian Sudan, Fiji, and the East African Protectorate

In 1913 the imports of castor seeds into Marseilles amounted to 25,157 tons, as compared with 16,588 tons in 1912, and 11,509 tons in 1911 (Diplom and Consular Reports, No. 5377 of 1914) By far the largest producer is East India

The total imports of castor oil into this country in 1916 were 1,322,000 gallons, in 1917 they were about 2,000,000 gallons. The imports of castor seed in 1917 were 45,000 tons, the bulk of which went to Hull, and yielded 4,176,000 gallons of oil, the greater part of which was used for aircraft engine lubrication (J Soc Chem Ind 1919, 20 R)

The seeds consist of 20 p c of husks, which are rich in mineral matter, but contain no oil, and 80 p c of kernels, forming a white, softish mass. The whole seeds contain a toxic principle—'ricine' or incinne. The kernels contain a powerful fat-hydrolysing enzyme, which can be made use of on a manufacturing scale for hydrolysing oils and fats for soap-making purposes.

poses

The enzymes lipase and esterase can be extracted from castor beans by means of ether (Falk and Suguira, J. Amer. Chem. Soc. 1915, 37, 217). The lipase is capable of hydrolysing oils at temperatures as low as  $-5^{\circ}$ , a fact which has an important bearing on the question of the enzymic hydrolysis of fats in food preserved in cold storage (Blanchet, Compt. rend. 1914, 158, 895). It is reversible in its action, and is capable of forming mono-di- and tirolein from an emulsion of oleic acid and glycerol (Krausz, Zeitsch angew. Chem. 1911, 24, 829).

The seeds contain 46 to 53 pe of oil oil is produced on a large scale either by expression or extraction For the production of best quality oil -for medicinal purposes—the seeds are decorticated before being expressed press-cake is subjected to a second and third expression, but the oils so obtained are unfit for medicinal use, and are employed for technical purposes (soap-making, Turkey-red oil, and lubricating oils). The cakes retain the poisonous alkaloid, and are therefore unfit for feeding purposes Although it is stated that by washing the cakes with 6 or 7 times their amount of a 10 pc sodium chloride solution, they can be freed from the alkaloid, no castor cake is actually used for feeding cattle, but applied exclusively to manuring the land The extracted oil is only used for technical purposes.

Another method of removing the toxic principle is to grind the seeds with water into a thin paste, which is then poured into boiling water. This coagulates the proteins, and the solid matter when strained off and expressed leaves a meal with the following composition Water, 9·2; asn, 7·3, oil, 2·6; proteins, 7·1·7, crude fibre, 5·0; and carbohydrates, 4·2 p.c (Indian Trade J. 1908, 9, 309; 1909, 11, 273)

A simple physiological method of detecting castor seed in other seeds is described by Lander and Geake (Analyst, 1914, 39, 292)

Crude costor oil is refined by steaming the oil, whereby the albuminous substances are Vol. II.—T.

coagulated, so that they can be removed by filtering.

Castor oil is a colourless or pale-greenish oil, having a taste at first mild, then harsh, this harsh taste is more pronounced in American than in Italian or French oils. The oil differs from all other fatty oils by its high specific gravity, viz up to 0.968. Contrary to older statements, the oil does not div, even when exposed to the atmosphere in thin layers. It has the following analytical characters. Sp.gr. 0.959, 0.968, solidification point, -17° to -18°; nodine value, 84-86°, Reichert-Meissl value, 2-2.5°, saponification value, 175-183°, acetyl value, 150°, and m.p. of fatty acids, 13°.

Castor oil is strongly destro-rotatory; the optical activity is due to the presence of nein-

oleic acid

The chief constituent of castor oil is the triglyceride of ricinoleic acid, both isomerides ricinoleic and isoriemoleic acid being comprised under this term, in addition thereto, castor oil contains a small proportion of glycerides of a natural dihydroxystearic acid and of stearic acid. Palinitin and olem are absent.

Owing to the presence of tririemolein, castor oil is distinguished from all other fatty oils by its high 'acetyl value' (see above and Oils and Fats) Castor oil further differs from all other fatty oils by its ready miscibility with absolute alcohol and glacial acetic acid, as also by its insolubility in large quantities of light petroleum, kerosene, and high-boiling paraffin oils. The oil gives a homogeneous solution with an equal measure of light petroleum, or one volume and a half of kerosene or paraffin oil; if more of the solvents is used, the excess will float on the top of the mixture. If castor oil be mixed with a small quantity of another fatty oil, this characteristic insolubility in petroleum distillates is

On subjecting castor oil to destructive distillation (as in the preparation of 'cognac oil'), undecylenic acid and conanthaldehyde distillation, whilst the residue solidifies to a very bulky, spongy, indiarubber-like mass, for which a solvent has not yet been found. This residue appears to consist of the anhydride of triundecenoic acid. By heating castor oil for 10 hours at a temperature of 260°-300° under a pressure of 4-6 atmospheres, the oil becomes miscible with mineral oils

When heated at 150°-250°, and then neutralised with alkali, castor oil yields soaps which are soluble in hydrocarbons, &c If, after heating, the oil is sulphonated before neutralisation, a substitute for Turkey-red oil is obtained, whilst evaporation of the product leaves a soap which does not readily form insoluble soaps with calcium and magnesium salts (Blumer, D. R. P. 227993, 1907, see also Eng Pat 24868, 1897).

Owing to its viscous character easter oil is particularly suitable for lubricating internal combustion engines, and is now largely used for aviation motors. Oil for this purpose should be colourless, have sp.gr. of 0 964–0 968, and an acid value of less than 0·3, and when heated in a flat dish for an hour at 215° should not darken to only material extent. The chief adulterants of oil supplied for lubrication are sugar and arachis oil (q.v.).

Castor oil is a good solvent for resins in

H

rubber, and may be used in the purification of ; raw rubber The solution of extracted resins may be made into soap (Schopper, Gum. Zeit. 1916, 30, 601).

A process of preparing acidyl derivatives of castor oil, which are free from the objectionable odour and taste of the oil, has been patented (D R P 226111, 1908) J L CASTOR OIL SEEDS or CASTOR BEANS.

The seed of Ricinus communis, a plant which grows as a tree in warm countries, attaining a height of 30 or 40 feet, but which in cooler climates is only an annual shrub

The following is an analysis of Indian seeds (Halenke and Kling, Landw. Versuch St 1906, 64, 51) .-

Sol carbo-Water Protein Oil hydrates Fibre Ash Kernels, 70 p.c. 3 60 23 43 66 02 4 01 0 70 2 24 Husk, 30 pc. . 8 76 4 76 0 98 32 92 48 69 3 89 . 5.14 17 88 46.65 12 61 14.99 2 73 Whole seed

In many varieties the proportion of husk is much lower, sometimes down to 20 p c

Of the total 'protein,' 22 62 pc, 4 18 pc., and 17:13 p.c respectively, are true albuminoids, whilst of the carbohydrates, 089 pc 15.63 p.c, and 5.29 pc respectively, are pentosans

The ash of castor seeds was analysed by

Thoms (1890)-

Sand, Water etc. Fe<sub>2</sub>O<sub>3</sub> CaO. MgO. K<sub>2</sub>O. Na<sub>2</sub>O. P<sub>2</sub>O<sub>5</sub> SO<sub>3</sub>. Cl. 0 **30** 19 59 5 65 17 08 10 53 **14** 30 1 88 23 67 6 01 0 29

Castor see'ls are very poisonous Cornevin found the following were the toxic doses in grammes per kilogram of body-weight Horses. oxen, sheep and dogs, 3, pigs about 5-6, hens about 40.

The poisonous constituent was believed to be ricinine by Soane (Chem Zentr. 1895, 1 853), but is now known to be ricin, an albumin, with two non-toxic substances—a proteose and a globulin-constitute the proteids of the seeds (Osborne and Mendel, Proc Amer Physiol Soc. 1903, 36).

Ricinine, according to Macquenne and Philippe (Compt rend 1904, 138, 506 and 139, 840), has the composition  $C_8H_8O_2N_2$ ; it melts at 201.5° and yields ricinic acid  $C_7H_6O_2N_2$  by saponification. They represent its constitution as

$$N(CH)_3$$
 $C(COOCH_3): C$ 
 $CH = CH \cdot C$ 

Ricin, the poisonous principle, has been examined by Stillmark (Chem. Soc Abstr 1890, 535), Osborne, and other (lc. and Amer Jour Physiol. 1905, 14, 259), and Brieger (Chem. Zentr. 1904, 1. 1286) It can be extracted from the pressed seeds by a 10 pc. solution of common salt, from which it is precipitated by magnesium and sodium sulphates.

Ricin agglutinates the red corpuscles of the Osborne found 0.002 milligram per kilogram of body weight sufficient to kill a rabbit when subcutaneously injected. It can be kept for months without deterioration, but loses its poisonous qualities when coagulated by heat Toxic symptoms do not appear until after a latent period, of about 15 hours Castor seeds also contain an enzyme capable of decomposing fats into glycerol and fatty acids, or, according to Falk and Sugiura (J. Amer Chem. Soc 1915, 37, 217), two enzymes—esterase and lipase, the former active towards ethyl butyrate, the latter towards triacetin. They also contain an enzyme ureasc, capable of hydrolysing urea (Falk, J. Amer Chem. Soc. 1913, 1914, 36, 2166), similar to, but weaker than, the urease of soja

Castor seeds are largely used as source of castor oil (q r) The cake left is rich in protein, and if it were not poisonous, would prove of value as a cattle food It contains-

By subjecting the cake to high-pressure steam, the poisonous qualities are said to be destroyed, and cake so prepared may safely be used as a part of the ration of fattening animals According to Ehrlich, animals may be immunised against the poison by feeding them with very small quantities of the crude cake until they become accustomed to it The serum from such animals may then be used to immunise others

Neither of these methods of utilising castor cake appears to have been successful, and, in England at least, it is rarely or never used for

feeding purposes. The leaves of the castor plant are rich in lime and potash an analysis by Wayne (Pharm. J Trans 1874, 3, 749) gave as the composition of the ash of the leaves

Cl CO<sub>2</sub> SiO<sub>2</sub> H I

CASUMEN. Trade name for a soluble form of casein

CATALASE v. Enzymes, art FERMENTATION. CATALYSIS v CHEMICAL AFFINITY.

CATAPLEIITE. Hydrated silicate and zirconate of sodium and calcium, containing ZiO2 30-40 p c The crystals are yellow and have the form of thin hexagonal plates, sp gr. 28, H. 6. It occurs with other rare minerals in nephelinesyenite on the islands in the Langesund-fjord in south Norway, and at Naisarsuk in Greenland L. J S.

CATECHIN v. CATECHU.

CATECHOL (Pyrocatechol or Pyrocatechin, o-Dihydroxybenzene, 1.2-Phendiol CoH4(OH), occurs in the leaves of the Virginia creeper (Ampelopsis hederacea) (Gorup Besanez, Ber 1871, 4, 905), in the sap of various kinds of the kino plants (Eisfeldt, ibid. 906; Flückiger, Ber 1872, 5, 1, 47), in crude beet sugar (Lippmann, Ber. 20, 3298; 1893, 26, 3061; J. Soc. Chem. Ind. 1893, 535); in the dry external scales of the onion; in Paglia olive oil (Canzoneri, Gazz. chim. ital 27, 3); in the colouring matter of red grapes (Sostegni, Gazz. chim. ital. 32, 17); in Silesian and Westphalian coal (Bornstein, Ber. 1902, 35, 4324); and in the tar water of bituminous shale (D. R. P. 68944, 1892; Ber. 1902, 35, 4325). The catechol complex also exists in certain glucosides (Nedra, J Soc. Chem. Ind. 1900, 686; Schmidt and Waljaschko, Chem. Zentr. 1901, u. 121), in a large number of products of vegetable origin, as the sulphate, catechol occurs in the urine of man and herbivorous animals (Muller, Ber. 1874, 7, 1526; Schmiedeberg, Zeitreh. physiol. Chem. 6, 189); it is probably also contained in the cerebro-spinal fluid (Halliburton, J Phys Chem. 10, 247)

It is formed in the dry distillation of moritannic acid (Wagner, Annalen, 1850, 75, 351, 80, 316); of catechin (Zwenger, Annalen, 1841, 37, 327); of protocatechine acid (Strecker, Annalen, 1860, 118, 285), and of all tannins which give a green colouration with iron chloride (Uloth, Annalen, 1859, 111, 215). It is also formed when filter paper is heated to 200° or when starch or sugar is heated to 280° with water (Hoppe Seyler, Ber. 1871, 4, 15) and in many other reactions.

Catechol is prepared by fusing sodium o-benzene disulphonate (10 parts) with sodium hydroxide (10-15 parts) in an autoclave for 8-10 hours at 280°-300° and pressure of 2-3 atmospheres. The mass is then dissolved in dilute sulphuric acid, neutralised with chalk, evaporated, cooled and filtered. The mother hquor is now heated in an autoclave with 50 p c sulphuric acid (10 parts) to 180°-220° phenol is removed from the solution so obtained by blowing steam through it; it is then decolourised with animal charcoal, after which the catechol is extracted with other (Baum, Eng. Pat. 21853; D. R. P. 80817; J. Soc. Chem. Ind. 1894, 879; Ellis, *ibid*. 1897, 674; Fidl. 1894-7, Catechol is also prepared by heating catechol disulphonic acid or the sodium salt with water or mineral acid to 200°-215° for 10-15 hours under pressure (Tobias, D. R. P. 81209, 1894, Frdl. 1894-7, 117), or by treating commercial catechu with water (Gutknecht, Chem. Zeit 1891, 15, 959), or by diazotising o-aminophenol, treating the product with boiling aqueous copper sulphate, and then extracting the catechol with ether (Fidl 1905-7, 128); or by the distillation of orthobromo- or chlorophenol in vacuo or in steam (D. R. P. 76597; Frdl 1890-4, 845), or by heating with sodium hydroxide (Frdl. 1894-7, 114, D. R. P. 84828) An almost theoretical yield of catechol can be obtained by treating guaracol with fuming hydrodic acid (Perkin, Chem. Soc. Proc. 1890, 90); or by the interaction of o-hydroxybenzaldehyde with aqueous sodium hydroxide in equimolecular proportions; to the sodium sait thus formed, an equimolecular proportion of 2-3 p c hydrogen peroxide is then added and the catechol is extracted with a suitable solvent (Dakin, Amer. Chem J. 1909, 42, 477) Also formed, together with phenol, by acting on benzene with hydrogen peroxide in presence of a terrous salt Other methods of preparing catechol are described by Cross, Bevan, and Heiberg (Ber. 1900, 33, 2018), Kunzkrause (Ber 1897, 30, 1620); Meyer (ibid 2569), Hartmann and Gattermann (Ber 1892, 25, 3532; D. R. P. 70718; Frdl 1890-4, 52); Gillard, Martend Catter (D. R. P. 70700), (Chem. Monnet and Cartier (D. R. P 97099; Chem Zentr. 1898, n. 521); Messel (D. R. PP. 68944, 60637; Frdl 1890-4, 844, 846)

Catechol is readily soluble in water, alcohol. ether, benzene, and crystallises from these solutions in prismatic needles or tablets, m.p. 104°; bp. 245° (Grabe, Annalen, 1889, 254, 296). Its aqueous solution precipitates metallic silver from silver nitrate solution at ordinary temperature, and with ferric chloride it gives a green colouration, which turns violet on addition of ammonia or sodium acetate (Wishcenus,

Annalen, 1896, 291, 174), whilst in the presence of aromatic amino-sulphonic acids the colour changes to red-brown and the test is more delicate than with ferric chloride alone (Bayer, Biochem. Zeitsch. 1909, 20, 178). Unlike its isomeride, resorcinol (metadihydroxybenzene), it is converted into an unstable quinone by the action of silvei oxide, and its calcium salt  $\operatorname{Ca(C_6H_5O_2)_2}$  is precipitated by an ammoniacal solution of calcium chloride (Bottinger, Ber 1895, 28, Ref 327). Catechol can be separated from its isomeride by adding antimony fluoride to their aqueous solution, when only catechol antimonyl fluoride will be precipitated (Causse, Ann Chim Phys 1898, (vii) 14, 526, 538)

99

An alkaline solution of catechol becomes green, then black on exposure to air; catechol is also oxidised to a black insoluble substance by the action of Catecholase, a thermostable substance obtained from the young shoots of Salir purpurea (Weevers, Pioc K. Akad Wetensch, Amsterd 1909, 12, 193). Heated with ammonium carbonate to 140°, catechol yields protocatechuic acid C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub>CO<sub>2</sub>H.

Catechol is used in photography, forming a very efficient developer in the presence of sodium sulphite or tribasic sodium phosphate as accelerator (Eder, J. Chem. Soc. Ind. 1890, 102., Bill. J. Photogr. 1900, 47, 676.) It can also be used as the starting-point in the preparation of a large number of valuable derivatives

Alkalı salts of catechol of the general composition OH  $\rm C_8H_4$  OM,  $\rm C_8H_4$  (OH)<sub>2</sub>, can be obtained by treating catechol with a solution of the theoretical quantity of alkalı hydroxide or of a salt with an alkalıne reaction, such as the sulphites. They form readily crystallisable salts, used in photography, and can be employed for producing pure catechol (Meister, Lucius and Bruning, D. R. P. 164666, 1904; Frdl. 1905–7, 129).

The mono- and di-ethyl others of catechol are prepared by heating under pressure a solution containing catechol, sodium hydroxide, and sodium ethyl sulphate or chloride in equivalent proportions. The ethers are then removed by steam distillation and separated by treatment with sodium hydroxide, in which only the monoether is soluble (J. Soc. Chem. Ind. 1896, 741, Eng. Pat. 16047, 1896). The diethers can be prepared by the action of alkalis on the monoethers (Fid. 1894-7, 123. D. R. P. 92651).

Catechol monoethyl ether forms large clear colourless crystals, mp 26°-27°, bp 215°, which have an odour similar to that of thymol, and are soluble in alcohol and ether, but not m water.

The monoalkyl ethers of catechol, when fused or treated in solution with various proteids, albumoses, and peptones, yield compounds which are applied therapeutically (Fehrlin, D. R. P. 162656, 1906).

Monoethyl catechol ether v Guaiacol. Dimethyl catechol ether v. Veratrol.

Catechol, when fused with aliphatic amines or heated with them in suitable solvents, yields stable amines which form useful photographic developers (D. R. P. 141101, 1903; Frdl. 1900-2, 1218). Other amino- and also nitroderivatives are described by Meldola, Woolcott and Wray (Chem. Soc. Trans. 1896, 1332); Heinisch (Monatsh. 1894, 15, 229); Nietzki and

100 CATECHOL.

Moll (Ber 1893, 26, 2182); Blankswa (Rec. trav. chim. 1905, 24, 40); Grischkewitsch-Trochimowski (J. Russ. Phys. Chem. Soc. 1909,

41, 1324); Dakın (l c.)

Good stable black, greenish, and bluish-black dyes, soluble in carbonates and alkaline sulphides, have been obtained by the action of aliphatic amines on catechol at 180°-222° (D. R. P. 84632; Frdl. 1894-7, 1048).

Methylene catechol ether CH<sub>2</sub>CO<sub>O</sub>C<sub>6</sub>H<sub>4</sub> is a colourless liquid; b p 172°-173° (Moureu, Bull. Soc. chim 1897, 15, 654, Mameli, Atta R. Accad Lincei, 1906, (v) 15, n 101; Perkin, Robinson, and Theorem. Robinson, and Thomas, Chem. Soc Trans 1909, 1977) A number of derivatives have been prepared (Moureu, lc 1898, 19, 507, Medinger, Monatsh, 1906, 27, 237; Barger, Chem. Soc. Trans. 1908, 563, 2081, Barger and Ewins, ibid. 735).

Catechol diphenylmethylene ether

$$CPh_2 < 0 > C_6H_4$$

is prepared by gently heating catechol with benzophenone chloride; colourless prisms, m p 93° (Sachs and Thonet, Ber. 1904, 37, 3327)

The methylene ethers of catechol derivatives can be converted into chlorine delivatives of their cyclic carbonates by being heated with thionyl chloride (Wellcome and Barger, Eng Pat 15987, 1907; J. Soc Chem. Ind 1908, 355)

287, 1907; J. Soc Chem. Ind 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 1906, 19

216° (110°-111°, 31 mm), is readily obtained by heating catechol with potash, ethylenic bromide and water in an atmosphere of hydro-Potassium permanganate, hydrogen iodide, and ammonia have an action on it, but it is oxidised by chromic acid to oxalic and carbonic acids It yields a number of substitution derivatives (Moureu, Compt. rend 1898, 126, 1426; Bull Soc chim 1898, (iii) 19, 507; Gattermann, Annalen, 1907, 357, 313, Lazennec, Bull Soc chim. 1909, (iv.) 5, 501, 509).

prepared by dehydrating orthohydroxyphenoxyaldehyde with phosphoric anhydride in presence of quinoline It is a colourless oil, boiling at 193°, and, when strongly cooled, forms brilliant white crystals, mp 16°-20° It forms a dibromide C<sub>6</sub>H<sub>4</sub>O<sub>2</sub>(CH)<sub>2</sub>Br<sub>2</sub>, mp 103°-104·5°, which, in contact with water, is converted into catechol, glyoxal, and hydrogen bromide (Moureu, Compt rend. 1899, 128, 559)

Catechol forms a number of acetals (Moureu, Compt. rend 1898, 126, 1656); also a vinyl derivative (Kurz Krause, Ber 1897, 30, 1617; Pauly and Neukam, *ibid* 1908, 41, 4151).

Catechol sulphonic acid C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub>SO<sub>3</sub>H is conveniently prepared by stirring together a meta-halogen phenol sulphonic acid with about an equal weight of sodium hydroxide in aqueous solution, and heating the mass to 250° for 8-10 hours The mass is now dissolved in water and neutralised with a mineral acid such as sulphuric. The solution is evaporated and the sodium sulphate filtered off, whilst the concentrate solution of catechol sulphonic acid yields pure catechol when heated with mineral acids

(Frdl. 1897-1900, 150, 1902-4, 100; D. R. P. 9623; Ellis, Eng. Pat. 14931, 1896).

Catechol disulphonic acid is obtained by treating catechol with 5 times the amount of fuming sulphuric acid (Cousin, Compt rend. 1893, 117, 113), or by fusing phenol trisulphonic acid or its salts with alkalis and decomposing the alkalı sulphonate so obtained with dilute sulphuric acid (Frdl 1894-7, 118)

Halogen derivatives (Jackson and Boswell,

Amer Chem. J 1906, 35, 519; Dakin, lc.).

Monochlorcatechol C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub>Cl, m p. 80°-81°, is obtained by the interaction of sulphuryl chloride with catechol in ethereal solution; by using a larger proportion of sulphuryl chloride, the duhlorcatechol, mp 105°-106°, is obtained (Paratoner, Gazz chim ital 1898, 28, ı 197)

Trichlorcatechol C<sub>6</sub>H<sub>3</sub>Cl<sub>3</sub>O<sub>2</sub> is obtained by the action of a chlorine solution in acetic acid or an acetic chloroform solution on catechol It forms colourless odourless prisms, mp. 104°-105°, with a burning taste Excess of chlorine yields tetrachlorocatechol (Cousin, Compt rend 1895, 120, 840) The corresponding bromoderivatives are obtained similarly (Cousin, lc).

The bismuth brominated catechols are recommended as a satisfactory substitute for tribromphenoxide, which is employed medically, under the name of 'xeroform' They are produced by adding an acid solution of bismuth nitrate to alkaline, tetra-, tri-, or di-biomocatechol, or by introducing precipitated bismuth hydroxide into an alcoholic solution of the bromocatechol (D. R. P. 107544, 1909)

Catechol carboxylic acid

$$C_6H_4(OH) O CO_2H, \frac{1}{2}H_2O$$

m p. 240°, is formed by the action of ammonium carbonate on catechol at 130°-140° under pressure, or by heating catechol with glycerol and potassium hydrogen carbonate at 180°-210° in an atmosphere of carbon dioxide (Praxmarer, Monatsh 1906, 27, 1199) The di-carboxylic acid can also be prepared Derivatives of catechol carboxylic acid, used in medicine, are obtained by the interaction of alcohols and also of primary and secondary bases with catechol carbonate; thus.

$$C_6H_4$$
 $O$ 
 $CO+C_2H_5OH$ 
 $=C_6H_4(OH)OCO_2C_2H_5$ 

by using anılıne, the compound C<sub>6</sub>H<sub>4</sub>(OH)O CONHC<sub>6</sub>H<sub>5</sub> is obtained, whilst with ethylene diamine it forms dicatecholdicarbethylenediamide  $C_2H_4(NH\cdot CO_2C_6H_4\cdot OH)_2$ (Einhorn and Lindenberg, Annalen, 1898, 300, 135; D. R. P. 92535; Frdl. 1894-7, 1110). A number of similar compounds are described by Einhorn and Lindenberg (l.c.); Einhorn and Pfeiffer (Annalen, 1898, 310, 218).

Methyl catechol dicarbonate, m.p. 41°, is described by Syniewski (Ber. 1895, 28, 1874).

Carbonyl esters of catechol of the type

CO(OC, H4OR), also used in medicine, are prepared by the action of phosgene on the corresponding alkyl catechol in sodium hydroxide solution (D. R. P 72806, Frdl 1890-4, 854).

¶Catechol monoacetic acid

C6H6(OH)O·CH2·CO2H

CATECHOL. 101

m.p. 131°, is obtained by the action of monochloracetic acid on catechol in aqueous or alkaline solution. When heated to 120°, it forms the anhydride C<sub>6</sub>H<sub>4</sub>CO, m.p. 56° (J Soc. Chem. Ind. 1896, 588), which, when treated with water, yields the acid in a very pure state (D. R. PP. 87336, 87668, 87669, 89593, Frdl 1894-7, 1106, 1107, 1108) The sodium salt of this acid is said to be a better remedy for phthisis and loss of appetite than guaiacol or its carbonate (Frdl. 1894-7, 1106). A large number of derivatives formed by the acid are described by Ludwig (J pr. Chem 1900, n. 61, 345). Catechol diacetates are described by Drezgowski (J. Russ. Phys. Chem Soc. 25, 157), and by Voswinckel (Ber. 1909, 42, 4561)

Catechol reacts with chloracetyl chloride or chloracetic acid, forming chloroacetyl catechol,

 $C_6H_4(OH)_2+ClCH_2COCl=C_6H_3(OH)_2COCH_2Cl$ which, with ammonia or a primary aliphatic amine, yields amino-acetyl compounds of the type RNHCH<sub>2</sub>CO C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub>, and these ketocompounds can be readily reduced to the cor responding alcohol (Imray, Eng Pat 26480, 1903, J. Soc. Chem. Ind 1904, 1043, D R P 13209, Frdl 1890-4, 857) (see also ADRENA-

Aniline and chloroacetyl catechol form an andrde C611 (OH)2CO CH2 NHPh, which, with mitrous acid, yields catecholglycophenyl triazine [CeH3(OH)2]2CO CH2 N Ph N NCH2CO, mp. 115°, with decomposition The corresponding tolyl- derivative melts at 120°, and is prepared in a similar manner (Dzierzgowsky, Ber 1894, 27, 1983, D R P 71312, Frdl 1890-4, 857)

Acetocatechol C  $_{\rm 6}{\rm H}$  ,Ac (OH)  $_{\rm 2},$  m p  $\,$  116° ,  $\,$  dimethoxyacetcatechol C  $_{\rm 6}{\rm H}$  ,Ac (OMe)  $_{\rm 2},$  and catecholqlycothrocyanate C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub>CO CH<sub>2</sub> SCN, m p 147°-150°, have been prepared by Dzierzgowsky (lc). According to Abderhalden and Kautsch, the chloracetyl derivatives of catechol, when treated with ammonia, are resolved into their components (Chem Zentr 1910, n 1356)

Alkyl acetyl catechol alkyl ethers, used in pharmacy, are obtained by the action of the alkyl acetyl chlorides on catechol monoalkyl ethers (J. Soc Chem. Ind 1906, 653, Eng Pat

25571, 1905; US Pat 822339, 1906).

A soluble tasteless product, C<sub>10</sub>H<sub>8</sub>N<sub>2</sub>O<sub>6</sub>, useful for medical purposes and as a substance from which other valuable technical products can be obtained, is formed by the condensation of catechol with alloxan in the presence of zinc chloride or mineral acids It forms prismatic crystals, which decompose above 200° (D. R. P. 107720. Frdl 1897–1900, 864).

Catecholdiantipyrine is obtained by mixing aqueous solutions of its constituents, and forms colourless needles, mp 78°-79° (Patein and

Dufan, Compt. end 1895, 121, 532)

C,H,1N,[C,H4(OH2] Prperidine catechol forms white crystals which rapidly become red and then brown on exposure to air. It melts at 80°-81° (Rosenheim and Schidrowitz, Chem. Soc. Trans. 1898, 140).

Catechol, when treated with phthalic anhydride and sulphuric acid at 150°, yields alizarin,

thus:

$$\begin{array}{c} {\rm C_6H_4} < {\rm OH(1) \atop OH(2)} + {\rm C_8H_4} < {\rm CO} \atop {\rm CO} > {\rm C} \atop {\rm OH(2)} \\ = {\rm C_6H_4} < {\rm CO} \atop {\rm CO} > {\rm C_6H_2} < {\rm OH(1) \atop OH(2)} \end{array}$$

(Baeyer and Caro, Ber. 1874, 7, 972; Liebermann and Hohenemser, ibid. 1902, 35, 1778) (v. ALIZARIN AND ALLIED COLOURING MATTERS)

Catecholphthalein C20H14O6 is prepared by heating phthalic anhydride (3 parts) with catechol (2 parts) and zinc chloride (3 parts) to 140°-150° for 3-4 hours. The aqueous extract of the product is boiled with animal charcoal, and the filtrate, on cooling, deposits yellowishwhite leaf crystals of the substance which sinters 80°-90°, but does not melt (Meyer and Pfotenhauer, Ber. 1907, 40, 1442, Bayer, Chem. Zentr. 1910, n. 1524).

Under certain conditions, catechol combines energetically with diazo- compounds yielding azodyes (Witt and Mayer, Ber. 1893, 26, 1072, Orton and Everatt, Chem Soc Trans. 1908, 1021) Thus Benzencazocatechol C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub>N<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, mp 165°, which dyes cotton mordanted with alumina a golden yellow, is formed by the interaction of an alcoholic solution of catechol and a concentrated solution of diazobenzene chloride, the mixture then being poured on to ice. The nitro- derivative NO<sub>2</sub> C<sub>6</sub>H<sub>4</sub>N<sub>2</sub>C<sub>6</sub>H<sub>3</sub>(OH)<sub>2</sub> colours alumina mordants ied, and iron and chromium mordants brown, the sodium sulphonate  ${\rm SO}_3{\rm NaC}_6H_4{\rm N}_2{\rm C}_6H_3({\rm OH})_2$  colours wool mordanted with alumina, yellow, with chromium, red-brown Similar toluene compounds have also been obtained Catechol also forms triazoor azoimido- compounds (Rupe and Majewski, Ber 1900, 33, 3401)

Catechol glycorsoquinoline

forms glittering orange-coloured crystals which give a red colouration with very dilute ferric chloride and a green colour in presence of excess of the latter, both the base and the chloride dye cotton mordanted with iron, aluminium, or chromium salts, black-yellow or reddish-brown respectively (Nencki, Ber 1894, 27, 1969).

 $Catecholylycotetrahydroquinoline = \mathrm{C_{17}H_{17}NO_3}$ crystallises in yellow lustrous plates, m p 170°

(Nencki,  $l \in$ )

Catechol, when heated with zinc chloride and formic acid, yields aurin dyes which give fast colours with metallic oxide mordants (Caro, Ber 1893, 26, 254)

Catechol tannins are employed in colouring leathers (J Soc Chem Ind. 1907, 423, 882, ibid 1909, 1151)

A number of products obtained by the condensation of catechol with ketones are described by Fabinyi and Széky (Ber. 1905, 38, 2307).

Catechol, when warmed with an alcoholie solution of picryl chloride containing sodium, yields 3 5-dinitro-o-diphenylene oxide,  $\rm C_6H_4O_2$   $\rm C_6H_2(NO_2)_2$ , mp 192° 192 5° (Hillyer, Amer. Chem. J. 1900, 23, 125)

Catechol heated with hippuryl chloride on the water-bath yields o-hydroxyphenyl hippurate NHBz CH<sub>2</sub> CO<sub>2</sub>C<sub>4</sub>H<sub>4</sub> OH, m.p 134°-136°, which with hydrogen chloride gives the anhydride  $C_{16}H_{11}O_{3}N$ , m p. 232°-233° (Fischer, Ber. 1905, 38, 2926). 102 CATECHOL.

When catechol is hydrogenated in the presence of finely divided nickel, cis-cyclohexane 1:2-diol, m.p. 75°-76°, b p. 225°, is formed (Sabatier and Mailhe, Compt. rend. 1908, 146, 1193).

A number of quinone and halogen quinone derivatives of catechol are described by Jackson and Koch (Amer Chem J 1901, 26, 10), Jackson and Porter (ibid. 1903, 30, 518), Jackson and Russe (ibid 1906, 35, 154, Ber. 1905, 38, 419) Other derivatives of catechol have also been prepared (Cousin, Ann Chim Phys 1898, (vii ) 480; 1899, (vii ) 18, 76; Bischoff, Ber 1900, 33, 1669; Wisinger, Monatsh 1900, 21, 1007; Bischoff and Hedenstrom, Ber. 1902, 35, 3452; Bischoff and Frohlich, Ber. 1907, 40, 2779, 2790; Barger and Ewins, Chem Soc. Trans 1909, 552, 1482; Fourneau, J. Pharm Chim 1910, (vii ) 1, 55, 97).

Catechol phosphines and chlorophosphines (Krauer, Ber 1894, 27, 2565); phosphites and sulphites (Anschutz and Posth, Ber 1894, 27, 2751); also a number of antimonyl catechol compounds (Causse, lc), some bismuth derivatives (Richard, Chem. Zentr. 1900, n 629), and a lead salt (Jackson and Koch, Ber. 1898, 31, 1458),

have been prepared.

Homologues of catechol, such as homocatechol (o-dihydroxytoluene), can be prepared by heating the corresponding alcohol with catechol in the presence of zinc chloride in open or sealed vessels (Merck, Frdl. 1894-7, 115, D R P. 78882), or by the interaction of persulphates and p-phenol derivatives in alkaline solution, the product then being warmed with acid (Frdl 1894-7, 121; D R. P. 81298).

$$\begin{split} \mathbf{C_6H_4} & \stackrel{OH(1)}{\underset{R}{\bigcirc}} + \mathbf{K_2S_2O_8} + \mathbf{KOH} \\ & = \mathbf{C_6H_3} \stackrel{OK(1)}{\underset{OSO_2OK(2)}{\bigcirc}} + \mathbf{K_2SO_4} + \mathbf{H_2O} \\ \mathbf{C_6H_3} & \stackrel{OK(1)}{\underset{R(4)}{\bigcirc}} \\ \mathbf{C_6H_3} & \stackrel{OK(2)}{\underset{R(4)}{\bigcirc}} + \mathbf{H_2SO_4} + \mathbf{H_2O} \\ & = \mathbf{C_6H_3} \stackrel{OH(1)}{\underset{R(4)}{\bigcirc}} \\ \end{split}$$

This method can be applied to the production of a large number of various derivatives.

CATECHU or CUTCH. There are several varieties of catechu or cutch bearing different names according to the country or plants from which they are obtained The following are those principally employed by dyers and tanners: Gambier catechu, Bengal or Acacia catechu, Bombay or Areca catechu, and Mangrove cutch

Although catechu has perhaps received greater attention than other natural dyes, the results have been so varied that the chemistry of the subject was for a long time in a most unsatisfactory Some confusion has arisen from the condition non-appreciation of the fact that the main constituents of Gambier and Acacia catechu are not identical, and some uncertainty has also apparently existed as to the botanical origin of the commercial varieties. Bombay catechu 18, for instance, occasionally referred to as originating from the Acacra catechu, and Bengal cutechu from the Areca catechu (Linn.); and in many cases, at the present time, it is impossible

to ascertain the botanical derivation of commercial brown cutch preparations.

Gambier catechu.—Gambier, yellow cutch, cubical cutch, cube gambier, or terra japonica, is obtained from the Uncaria gambier, an extensive scandent bush which is met with, both wild and cultivated, in Malacca, Penang, and Singapore. The catechu is isolated by extracting the leaves and twigs with hot water until the liquid becomes syrupy, the insoluble matter being removed from time to time by means of a strainer. On cooling, the pasty mass is cut into cubes with sides I inch in length and dried on bamboo trays

Catechin C<sub>15</sub>H<sub>14</sub>O<sub>6</sub>,4H<sub>2</sub>O, the crystalline principle, was first described by Nees van Esenbeck (Annalen, 1832, 1, 243), was subsequently examined by Berzehus in 1837 (J 14, 235), and more recently by numerous chemists isolate catechin, Lowe (Zeitsch anal. Chem 13, 113) devised the following method: catechu is washed with cold water to remove catechu tannic acid, well pressed, allowed to dry, and dissolved in hot acetic ester The filtered solution is evaporated and the residue crystal-

hsed from hot water

According to Perkin and Yoshitake (Chem. Soc Trans. 1902, 81, 1162), a combination of this method and that of Berzelius (lc) gives good results

The finely powdered catechu is extracted with 10 times its weight of boiling ethyl acetate, the solution is evaporated, and the residue crystallised from 10 times its weight of water The product is again dissolved in boiling water, and lead acetate solution added, drop by drop, until a coloured precipitate is no longer formed, and the filtrate is almost colourless, the latter, while hot, is treated with sulphuretted hydrogen, the lead sulphide removed, and the crystals, which separate on cooling, are collected, washed, and allowed to dry at the ordinary temperature. The substance is now practically colourless, and the yield greater than if the purification had been effected by frequent crystallisation from water

The formulæ which have been assigned to catechin are very numerous It has been shown by v Kostanecki and Tambor (Ber 1902, 35, 1867), and simultaneously by Perkin and Yoshitake (lc), that it is correctly represented

as C<sub>15</sub>H<sub>14</sub>O<sub>6</sub>,4H<sub>2</sub>O

Catechin forms colourless needles, and when crystallised from water the air-dry product melts at 96° (Clauser, Ber 1903, 36, 101). After standing over sulphuric acid, it possesses the formula C<sub>15</sub>H<sub>14</sub>O<sub>6</sub> H<sub>2</sub>O, and melts at 176°-177°, and this is also the melting-point of the anhydrous substance. Catechin is readily soluble in boiling water and cold alcohol, and gives with lead acetate solution a colourless precipitate, and with ferric chloride a deep-green liquid. With pine wood and hydrochloric acid, it gives the phloroglucinol reaction.

On fusion with alkali, protocatechuic acid, phloroglucinol, and probably acetic acid are produced, and it is interesting to note that catechu has been considerably employed for the commercial preparation of the former compound.

Schutzenberger and Rack (Bull. Soc. chim. 4, 6) have described a dibenzoyleatechin; and Liebermann and Tauchert (Ber. 1880, 13, 695), diacetyl catechin, diacetyl dichloreatechin, and diacetyl dibromeatechin; but their formulæ for these compounds are now known to be incorrect.

Penta - acetyl - catechin  $C_{15}H_{9}O_{6}(C_{2}H_{3}O)_{5}$ ; colourless needles, m.p.  $124^{\circ}-125^{\circ}$  (v. Kostanecki and Tambor); pentabenzoyl catechin,  $C_{15}H_{9}O_{8}(C_{7}H_{8}O)_{5}$ , colourless needles, m.p.  $151^{\circ}-153^{\circ}$ , tetrebenzoyl catechin,  $C_{15}H_{10}O_{6}(C_{7}H_{5}O)_{4}$ , prisms, m.p.  $171^{\circ}-172^{\circ}$  (Perkin and Yoshitake), disazobenzene catechin,  $C_{15}H_{12}O_{8}(C_{5}H_{5}N_{2})_{2}$ , salmon - red needles, m. p.  $193^{\circ}-195^{\circ}$  (Ett. P. and Y.), acetyldisazobenzene catechin, orange red needles, m.p.  $253^{\circ}-255^{\circ}$  (P. and Y.); catechin tetramethylether  $C_{15}H_{10}O_{3}(OCH_{3})_{4}$ , needles, m.p.  $141^{\circ}-146^{\circ}$  (K. and T.), acetyl catechin tetramethylether  $C_{15}H_{9}O_{2}(OCH_{3})_{4}C_{2}H_{3}O$ , needles, m.p.  $92^{\circ}-93^{\circ}$ , and catechin pentamethylether  $C_{15}H_{9}O(OCH_{3})_{5}$  (K. and T.), have been prepared M.

When catechin tetramethylethor, suspended in water, is oxidised with potassium permanganate, it gives veratric acid, and most probably phloroglucinol dimethylether (Chem Soc Trans 1905, 87) As the result of this investigation, Perkin considered that catechin was possibly a reduction product of quercetin

and indeed a small quantity of the latter colouring matter is present in catechi (Lowe Perkin, *ibid* 71, 1135)

More recently, v Kostanecki and Lampe (Ber 1906, 39, 4007) have shown that when catechin tetramethylether is brominated in the presence of sunlight, only a monobromeatechin tetramethylether, mp 173°-174°, is produced, whereas the above constitutional formulæ require the formation of a dibromo-compound. When oxidised with permanganate, bromeatechin tetramethylether gives veratric acid, so that the bromine must have replaced one hydrogen of the phloroglucinol nucleus. When alcoholic solutions of catechin-tetramethylether and of iodine and iodic acid are mixed, long colourless needles of whours:—

$$\begin{array}{c} \text{MeO} \\ \text{MeO} \\ \text{MeO} \\ \end{array} \begin{array}{c} \text{H} \\ \text{CH}_2 \\ \text{OH} \\ \text{OMe} \end{array}$$

m.p.  $192^{\circ}$ - $193^{\circ}$ ; this yields an acetyl derivative  $C_{21}H_{23}O_{,}I$ , m p.  $189^{\circ}$ 

Iodocatechin-tetramethylether when treated with zinc-dust and alkali yields the original catechin-tetramethylether. this elimination of reduced being also noticed with other members of the maclurin type (v Kostanec'i and Lampe, Ber. 1907, 40, 4910). v Kostanecki and Lampe

have assigned the following constitution to catechin:—

and this is in harmony with many of the properties of this substance. When catechin tetramethylether is oxidised with chromic acid, catechone trimethylether, orange-yellow needles, m p 210°, is produced (K and T).

$$\begin{array}{c|c} CH_3O & CH_3O \\ \hline \\ CH_3O & -CH(OH) - \\ \hline \\ \end{array}$$

whereas catechin pentamethylether gives the tetramethyl compound.

and this behaviour is analogous to that of leucomaclurin pentamethylether (see MACLURIN)

Acetyl catechone by reduction with zinc-dust and glacial acetic acid, and hydrolysis of the product, gives hydroxy-catechin, yellow needles, melting at 284°–285°—

The hexamethylether has mp 102° (Nierenstein Annalen 1903, 396, 194).

stein, Annalen, 1903, 396, 194).
When catechone trimethylether is nitrated, a mononitro- derivative (1)

$$(2) \quad \begin{array}{c} \text{CH}_3\text{O} \\ \text{CH}_3\text{O} \end{array} \begin{array}{c} \text{NO}_2 \\ \text{-COOH} \end{array}$$

m p. 141°, is obtained, and on oxidation yields nitroveratric acid (2) Finally, by the reduction of catechin tetramethylether with sodium and alcohol, an oil is produced which, on treatment with methyl sulphate, gives pentamethoxyethyldiphenylmethane

a reaction which points clearly to the presence of the cumaran ring in catechin.

Ryan and Walsh (Proc. Roy. Dub Soc.

1916, 15, 113) have attempted to decide between the chroman structure of Perkin and Yoshitake and the coumaran formula of v Kostanecki and Lampe. To this end they have synthesised  $2.4 \cdot 6:3' \cdot 4'$ -penta-methoxy-3-ethyl-diphenylmethane, in order to ascertain whether this was identical with the deoxy-hydro-catechin-pentamethyl-ether obtained from catechin, as described above by v Kostanecki and Lampe

The synthesis is made clear by the following scheme :-

They were, however, unable to obtain the resulting product in a crystalline conditionthe addition of a crystal of the product obtained catechin (deoxy-hydro-catechin-pentamethyl-ether) failed to cause it to crystallise —hence they were not able to obtain a definite settlement of this point

Catechin is oxidised in aqueous solution by potassium ferricyanide in the presence of potassium acetate with formation of an orangecoloured amorphous compound (Perkin), which dyes mordanted calico orange-brown shades which are fairly fast to soap It is probable that this dyestuff is related to the catechone of v Kostanecki and Tambor, and which was obtained by these authors in the form of its tri- and tetramethyl ethers

When catechin is treated with carbon tetrachloride and aqueous potassium hydroxide, it yields catechin carboxylic acid, colourless needles, m.p. 274°-277° (decomp).

COOH 
$$C_6H_2(OH)_2$$
— $CH(OH)$ — $OH$ 

The annexed constitution has been assigned to it on account of the fact that its pentamethyl-ether methyl-ester C22H26O8, mp 92°, when oxidised with alkaline permanganate, yields hemipinic acid.

Catechin carboxylic acid has been resolved, and the *d*-acid has  $[a]_D^{18} = +76 \, 4^\circ$ , and mp. 273° (decomp), whilst the *l*-acid gave values  $[a]_D^{18} = -68 \, 22^\circ$ , and m.p. 270°-273° (decomp.) (Nierenstein, Annalen, 1913, 396, 194)

Catechutannic acid. In addition to catechin.

gambier catechu contains a small amount of catechutannic acid, and this substance is present in considerable quantity in the browner varieties To isolate it, the following method of cutch has been employed .-

A hot aqueous extract of catechu is allowed to stand until no more catechin is deposited, and the clear liquid is evaporated to dryness. residue is dissolved in alcohol, the solution treated with ether to precipitate impurities and then evaporated to diyness

Catechutannic acid consists of an amorphous reddish-brown powder, readily soluble in water According to and alcohol, insoluble in ether Lowe (Fr. 13, 121), it possesses the formula C21H18O8, and gives a lead salt 3PbO,2C21H18O8,

but this most probably requires revision According to Etti (Annalen, 186, 332), catechutannic acid is an anhydride of catechin, and is derived from this substance by elimination of It is said to be produced when an aqueous solution of catechin is heated to 110° (Lowe, *ibid* 12, 285), or by boiling catechin with solutions of the alkali carbonates Again, catechin is decomposed at its melting-point with evolution of water and formation of a product resembling catechutannic acid, and aqueous solutions of catechin on long standing become brown-coloured with apparent formation of this That the products obtained by these compound methods resemble catechutannic acid is certain, but the subject has not been fully investigated.

Catechutannic acid solution gives a precipitate with lead acetate, and also resembles catechin in giving a green colouration with ferric chloride, and the phloroglucinol reaction with pinewood and hydrochloric acid It is a powerful tanning agent, and appears to differ but little from the so-called 'catechol' tannins

Three other catechin anhydrides have been described, viz the second anhydride ('42H34O15(?), (Etti), which is produced by heating catechutannin to  $162^{\circ}$ , the third anhydride  $C_{21}H_{16}O_{7}(?)$ , formed by digesting catechin with boiling dilute sulphune acid for several hours, and the fourth anhydride, which is obtained by heating catechin with hydrochloric acid at 160°-180°. These compounds consist of reddish-brown powders, and the fourth substance is insoluble, both in alkaline solutions and all solvents A product similar in properties to Etti's fourth anhydride is readily formed by adding sulphuric acid to a hot acetic acid solution of catechin (Perkin). A bright orange powder almost immediately separates, and this, on analysis, gave C=63 26 p c, H= 3 89 рс It is interesting to note that the formation of these red anhydrides, or phlobaphenes is characteristic of all the catechol (phlobo) tannins

According to Perkin and Yoshitake, gambier catechu contains a small quantity of a second catechin, which crystallises in small prisms. devoid of water of crystallisation, and melts at 235°-237° Its general reactions are identical with those of ordinary catechin, and by fusion with alkali it also gives phloroglucinol and protocatechnic acid. The azobenzene compound  $C_{15}H_{12}O_6(C_6H_5N_2)_2,$  orange-red needles, melts at  $215^\circ\text{--}217^\circ$  .

Finally, there are present in catechu, certain brown substances known as rubinic and japonic acids, which, according to some writers, appear to have been formed by the oxidation of the and also of their derivatives, differ widely, as is Their chemical nature is, however, catechin unknown.

Catechin, though largely employed for tanning purposes, does not precipitate a gelatine solution, and is not itself a tannin matter. On the other hand, it is absorbed by the hide, and there gradually passes into catechutannic acid

Dyeing Properties -On cotton, catechu is largely used for the production of the wellknown 'catechu brown,' which is exceedingly fast to light, acid and alkaline solutions, and also to bleaching powder. To obtain this, cotton is steeped in a hot solution of catechu (1-2 pc) to which has been added about 6 pc of copper sulphate, reckoned on the weight of the catechu employed The material is allowed to remain in the bath as it cools, and without washing is then treated in a warm or boiling second bath containing 1 or 2 grams of bichromate of potash per litie According to Hummel and Brown (J Soc Chem Ind 1896, 15, 422), in this operation the copper sulphate probably converts the catechin into catechutannic acid, and this is subsequently oxidised to japonic acid by the action of the bichiomate The colour is apparently intensified by the formation of a basic copper chromate Gambier catechu is also employed in dyeing compound shades with log-wood, fustic, and alizarin in conjunction with bichrome, and with bismarck brown, magenta, In the latter case catechu-tannin forms the mordant for the basic colour

Wool may be dyed with catechu in a similar manner to cotton. In silk-dyeing, catechu is largely used for weighting purposes. The silk is steeped in basic ferrous sulphate solution, then in potassium feriocyanide and hydrochloric acid, which causes the production of Prussian blue on the fibre It is then worked in a strong solution of gambier to which stannous chloride solution has been added.

Bengal or Acacia catechu.—Bengal catechu is derived from the Acacia catechu, a tree 15 to 20 feet high, which is common in most parts of India and Burma To isolate the catechu which is present in the red heartwood, the tree is cut down while it is fullest of sap, and the internal portion is sawn into small sticks and extracted with boiling water The liquid thus obtained is concentrated over a fire, and then allowed to evaporate spontaneously in shallow dishes extract thus obtained comes into the market as Pegu catechu, Brown cutch, and Brown catechu (Crookes, Dyeing and Calico Printing)

A purer substance, kath, or the pale catechu of India, is prepared by suspending twigs in the hot concentrated extract and collecting the

crystals which thus separate

Acacatechin  $C_1$ ,  $H_{14}^{\dagger}O_6$ 3 $H_2O$  can be obtained from the acacia kath by methods identical with those which are employed for the isolation of catechin from gambier catechu. It melts at 204°-205°, is somewhat more sparingly soluble in water than catechin, and when fused with alkalı gives phloroglucinol and protocatechnic acid. The reactions of acacatechin and catechin, in so far as they have been observed, are identical, and there is no difference in the composition of their derivatives when prepared under similar conditions. On the other hand, the melting-points of the substances themselves.

evident from the following table (Perkin and Yoshitake) :-

	Acacatechin	Catechin
Pentaacetyl derivative		124°-125°
Pentabenzoyl ,,	181°-183°	151°-153°
Azobenzene "	198°-200°	193°-195°
Tetramethyl ,,	152°-154°	144°-146°
Acetyl ,,	135°-137°	92°93°

It appears certain, therefore, that catechin and acacatechin are isomerides, and it is possible that the latter may, on the basis of v Kostanecki and Lampe's formula for catechin, be represented thus (Perkin, private communication):

a point which should not be difficult to determine

Bombay or Areca catechu. This variety is obtained from the fruit of the Arcca catechu, or betel-nut palm, a tree which is common in tropical Asia. Though its chief constituent resembles catechutannie acid, catechin itself has not been isolated from this product possesses a bright-chocolate and sometimes an orange-brown colour, and yields, on dyeing, very similar results to the ordinary cutches

Mangrove cutch. Mangrove cutch is obtained from the bark of the mangrove Cerups candolleana, and is of somewhat recent employ-In its preparation it is preferable to extract the fresh bank, which is of a light colour internally, rather than the stored product which has become red, or is said to have 'sweated' The extract, on evaporation, becomes deep-red in colour, and the object of the manufacturer is to prevent this anhydride formation from going too far, otherwise a certain portion of the extract is rendered insoluble in water

By fusion with alkali, it gives protocatechnic acid, but at present no catechin has been isolated from it (Perkin, private communication) dissolved in sodium bicarbonate solution, acetic ester extracts an almost colourless tannin, which possesses the properties of a catechol tannin

For many purposes, mangiove cutch is competing closely with the other varieties of catechu, and this competition will no doubt become keener if its quality can be further improved

Mahogany. According to Caseneuve (Ber. 8, 828), mahogany wood contains a catechin

Kino or Gum kino (see under Kinos).

CATELLAGIC ACID C<sub>14</sub>H<sub>6</sub>O<sub>6</sub>. Schiff (Ber 12, 2590), on heating protocatechnic acid with arsenic acid, obtained a substance which he named catellagic acid, and to this he assigned the formula C<sub>14</sub>H<sub>10</sub>O<sub>7</sub> or C<sub>14</sub>H<sub>8</sub>O<sub>7</sub> Perkin and Nierenstein (Chem Soc. Trans 87, 1417) obtained apparently the same substance by the oxidation of protocatechuic acid, and also parahydroxybenzoic acid with potassium persulphate and sulphuric acid. According to the latter authors, catellagic acid crystallises from pyridine in colourless needles, which melt above 360°, and sublime with but moderate carbonisation at higher temperatures. Solutions of the alkali hydroxides dissolve it with a pale-yellow colour,

and with nitric acid it gives a magenta-coloured hquid Diacetylcatellagic acid, colourless prismatic needles, melts at 322°-324°.

Catellagic acid, by distillation with zinc dust, gives fluorene, and is closely related to ellagic Its constitution (P. and N.) may be expressed as follows -

From the oxidation products of parahydroxybenzoic acid, Perkin and Nierenstein isolated, in addition to catellagic acid, compound,  $C_{14}H_8O_6$ , colourless needles, mp above 360° This, which gives the diacetyl derivative  $C_{14}H_6O_6(C_2H_1O)_2$ , colourless needles, mp 267°–268°, and by distillation with zincdust fluorene, possesses the formula .

and evidently originates from the interaction of one molecule of protocatechuic acid and one of parahydroxybenzoic acid

CATHARTIC ACID v SENNA LEAVES.

CATHARTIN v. SENNA LEAVES.

CATHARTOGENIC ACID v SENNA LEAVES CATHARTOMANNITE v SENNA LEAVES

CAULIFLOWER. A variety of Brassica oleracca, in which numerous imperfect flowers form a compact head, which forms the main edible portion

An American analysis shows the edible portion to contain:

Soluble Water Protein Fat carbohydrates Ash 908 1.6 08 6 Ö 0.8 H I

CAULOPHYLLINE v CYTISINE

CÂY CÂY FAT or COCHIN CHINA WAX v. WAXES

CAYENNE PEPPER v Resins.

CEANOTHUS WAX v Snowbrush WAX, art. Waxes

CEDAR CAMPHOR v CAMPHORS

CEDAR OIL v OILS, ESSENTIAL

CEDRA. (Cedrat, Fr.) The fruit of a species of orange, citron, or lemon The peel is very thick, covered with an epidermis containing a fragrant essential oil, used for flavouring preserves, also used whole for wet comfits, and cut in quarters for dry comfits A liqueur is made from it by gathering the fruit before it is ripe, and grating the peel into brandy

CEDRELA TOONA. The Cedrela toona, the Toon or Indian mahogany tree, is a large tree 50-60 feet, although occasionally reaching 100 feet in height, the wood of which closely resembles mahogany and is imported into England under the name of 'Moulmein Cedar,' and much used for making furniture It occurs in the tropical Himalaya from the Indus eastward, throughout the hilly districts of Central and Southern India to Burma, and is also found in Java and Australia. The flowers constitute one of the less important Indian natural dyestuffs, and are reputed to yield both a red and a yellow dye (known in Bengal as 'Gunari,' which is applied without mordants). Although employed for dyeing purposes alone, the flowers are also used in Burma in conjunction with safflower and together with turmeric to produce the sulphur-yellow colour or 'basanti' of Cawnpore.

To isolate the colouring matters, Perkin (Chem. Soc Trans. 1912, 101, 1539) digested an aqueous extract of the flowers with a little hydrochloric acid at 70°-80°, which caused the separation of a reddish-brown precipitate. This collected and dried was exhausted with boiling alcohol, and the solution on evaporation deposited a small amount of a red microcrystalline powder By re-crystallisation from pyridine and removal of an insoluble amorphous impurity, this separated in large orange-coloured leaflets which contained pyridine, and became red and opaque when washed with benzene or alcohol,

the pyridine being thus eliminated.

It melted at 285°-287°, and with cold concentrated sulphuric acid gave a deep indigo blue liquid, resembling in this property bixin, the colouring matter of annatto, Bixa orellana (l.c.). This compound proved to be identical with the red colouring matter nycanthin isolated by Hill (*ibid.* 1907, 91, 1501) from the flowers of the Nycanthes arbor-tristis, to which the erroneous m p 234°-235° has been assigned. The amount which is present in the Toon flowers is very small and does not seem to average more than 0 1 p c. Though there appears to be little doubt that nycanthin and bixin are closely related, the connection between the two compounds cannot be predicted with any reasonable certainty until chemists are in agreement as to the correct The analyses of Hill and formula of bixin of Perkin indicate that nycanthin contains approximately C=7268, H=775, and the simplest formula to which this corresponds is C<sub>15</sub>H<sub>18</sub>O<sub>3</sub> Among the numerous expressions proposed for bixin is, however, that of van Among the numerous expressions Hasselt (cf Annatto), C<sub>29</sub>H<sub>34</sub>O<sub>5</sub>, and should this eventually prove to be correct, then nycanthin is possibly an oxy- or hydroxy-bixin,  $C_{29}H_{34}O_{6}$  (C=72 8, H=7·1).

Addition of zinc-dust to the hot orangeyellow coloured alkaline solution of nycanthin causes decolorisation From this by acidification a colourless or faintly yellow precipitate separates, and this, as is well known, is a property also possessed by bixin

Cotton on immersion in a boiling dilute solution of nycanthin in sodium carbonate, and subsequently rinsing with acidified water, is dyed an orange shade indistinguishable from that given in these circumstances by annatto, a fact which again points to a relationship between the two colouring matters

The flowers of the Cedrela toona contain in addition to nycanthin a fair amount of quercetin probably as glucoside, and also a second flavone or flavonol dysetuff, the identity of which was not ascertained A sugar, C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>, is also present, and this after purification melted at 165°-168°

Cotton and woollen fabrics can be dyed a dull yellow colour by mere immersion in a boiling extract of Toon flowers. This, however, is not permanent, and is removed, although much more quickly from cotton than wool, by the actior of soap or d.lute alkalı. Better results are obtained by the employment of mordants, and

the following shades were obtained with mor danted woollen cloth :-

Chromium Aluminium Dull brownish- Full golden- Bright Dull yellow. yellow. yellow olive-brown.

During these operations, nycanthin or its glucoside apparently plays no part, and the colours obtained are due to the flavone glucosides present Not only the flowers but the seeds of the Cedrela toona are stated to give a red dye

A. G. P. CEDRIRET. A substance found in woodtar by Reichenbach and recognised by Liebeimann as a quinone-like substance and termed by him cærulignone. Hofmann showed that it was tetramethoxydiphenyl quinone, formed by the oxidation of pyrogallol dimethyl ether (v CERU-LIGNONE and QUINONES)

CELERY. A prum graveoleus The blanched stems are used for food

The composition, as given by American analyses, is .

Water Protein Fat Carbohydrates Ash

A more detailed analysis is given by Church (Foods, 1889)

Starch and Water Protein mucilage Sugar Cellulose Ash 20 16 0.9

Mannitol is one of the chief carbohydrate Asparagme (about 05 pc) and constituents tyrosine, are among the nitrogenous constituents (Bamberger and Landsiedl, Monatsh 1904, 25, 1030).

The characteristic odour and flavour of celery are due to an essential oil, which, according to Ciamician and Silber (Ber 29, 1799, 1897, 30, 492, 1419), consists of a terpene, probably  $C_{13}H_{24}$ , and two acid bodies, sedanonic acid  $CO_2H$   $C_6H_5$  CO  $C_4H_9$  and sedanolic acid

 $CO_2H C_6H_8 CH(OH) C_4H_9$ the anhydride  $C_6H_8 \subset C(C_4H_8) O$  of the former and the lactone  $C_6H_8 < C_6C_4H_9 > 0$ 

of the latter appear to be the constituents upon which the smell of the celery depend acid and a phenolic substance are also present.

CELERY OIL v. OILS, ESSENTIAL CELESTINE BLUE v OXAZINE DYESTUFFS CELESTITE or Celestine. Native strontium sulphate (SrSO<sub>4</sub>), forming orthorhombic crystals isomorphous with barytes (BaSO<sub>4</sub>) The crystals are colourless or yellowish, but sometimes they show a pale tinge of sky-blue, and on this account the mineral receives its name. The mineral is often very like barytes in appearance, and it is also heavy (sp gr 39, of barytes, 4.5), it is most readily distinguished by the colour the powdered mineral moistened with hydrochloric acid imparts to the Bunsen flame. Numerous finely crystallised specimens have been found in the red marls of Triassic age in the neighbourhood of Bristol, and in the sulphur-mines near Girgenti in Sicily. Deposits of celestite occur in beds of Cretaceous age over very wide area in Russian Turkestan; and the mineral is found also in Baluchistan. The

largest crystals, measuring a foot across, are from a cave in limestone on Strontian Island, Lake Erie, Ohio. Workable deposits are mined in Arizona and California. The mineral is dug in numerous shallow pits in Gloucestershire and Somersetshire, the production amounting to about 20,000 tons per annum, valued at about 11 per ton Most of this was exported to Germany, where in the form of strontium hydroxide, it was used in the refining of beet-It is also converted into strontium nitiate for making red fire in pyrotechny. For details respecting the British occurrences, see Special Reports on the Mineral Resources of Great Britain, vol III. (Mem Geol Survey), 1915, 2nd ed 1918 L. J. S.

CELLOIDIN. Trade name for a purified

form of pyroxylm.

CELLOSE (CELLOBIOSE) v. CARBOHY-DRATES.

CELLOTROPIN. Trade name for benzoic ester of arbutin

CELLULASE v ENZYMES

CELLULOID (Pyroxylin Plustic) is a colloid composed of soluble mitrocellulose, approaching  $C_{12}H_{16}(NO_2)_4O_{10}$  in composition, together with camphor—It is obtained by gelatinising nitrocellulose by means of a solution of camphor in ethyl alcohol or a similar menstruum. The word 'celluloid' is a registered trade-mark in the United States The substance was first produced by Alexander Parkes, of Birmingham, in England, under the name of 'Parkesine.' Parkes was succeeded by Daniel Spill, of Hackney, London, who adopted the name of 'Xylonite,' under which name it is still well known in Great Britain Other trade names are Fiberoid, Py-ra-lin, Viscoloid, &c Great improvements in the manufacture, especially in mechanical features, were made by John W. Hyatt, of Newark, N.J., U.S.A., and the material first became a commercial success in the United States.

The introcellulose employed is preferably made from tissue paper by subjecting it for 20 minutes or longer to the action of a bath of mixed sulphune and nitne acids containing 19-20 p c of water, at a temperature of 30° or more The resulting nitrocellulose is thoroughly washed with water to free it from acids, and bleached The surplus water is then removed

The nitrocellulose is then mixed with alcohol, camphor, and if necessary, pigments and fillers, and thoroughly incorporated in a kneading Transparent material receives a small The doughpercentage of urea as a stabilizer like mass is worked and formed into sheets between slightly heated rolls. These sheets are consolidated in hydraulic presses heated to about 75°C, into a solid cake, from which after cooling sheets of the desired thickness are cut, or the mass taken from the rolls is placed in a heated cylindrical hydraulic press, provided with suitable nozzles and dies, from which it emerges in the shape of tubes, rods, or the like, of any desired cross-section Rods may also be made out of the solid cake previously mentioned, by forcing a cutting tool through it. All this material has to be freed from its excess of solvent, and is seasoned in rooms, heated to 30°, where it is kept for a period of days, weeks, or months, according to thickness. The camphor solvent

108 CELLULOID.

usually employed is denatured alcohol, owing to the fiscal policy of the countries concerned, but wood alcohol, pure grain alcohol, and a mixture of one or both of them with fusel oil, or its various constituents, have also been widely used. Owing to the great fluctuations in the price of camphor in recent years, the attempts to make this substance synthetically, and to find a substitute for it, have been pursued with vigoui. Synthetic camphor in commercial quantities is an accomplished fact, and is the equivalent of the natural product. The search for substitutes was begun in the early eighties, the most suitable found so far being acid derivatives of aromatic amines.

Celluloid is coloured by means of mineral colours, lakes, and coal-tar dyes, to reproduce the appearance of some more expensive substances, such as ivory, tortoise-shell, leather, coral, amber, &c, or to replace glass in the manufacture of articles in which brittleness is a serious disadvantage. A special variety of celluloid is used in the manufacture of cinematograph film. This contains a higher proportion of introcellulose than ordinary celluloid, and the nitrocellulose is more highly intrated.

Celluloid, while highly inflammable, is non-explosive under any conditions to which it may be subjected in the work of cutting, turning, or moulding, or indeed under any normal conditions. Its specific gravity varies with its composition. It is about 1.35 for the unpigmented material.

Celluloid is plastic at a temperature of 75°, and in practice it is usually softened in boiling water or on a steam table preparatory to moulding. This property of turning plastic at comparatively low temperatures, and its hardness and elasticity at ordinary temperatures, as well as its resistance to the action of water, dilute acids and many other agents, constitute its great technical value. It is used extensively in the manufacture of combs, and other toilet articles, knife handles, piano keys, collars, toys, and for a great variety of other purposes. A relatively less important branch of the

A relatively less important branch of the celluloid industry is concerned with the manufacture of artificial leather by spreading a solution of celluloid on fabric. Celluloid solutions are also used in the manufacture of patent leather, and for metal lacquers, wood enamels, incandescent mantle dips, and a variety of other purposes where a quick drying water-proof transparent or pigmented coating is required, which may be either hard or flexible

For a method of analysing celluloid, sec Hervé, Caoutchouc et Gutta Percha, 1918, 15, 9601; J Soc. Chem Ind. 1918, 686 A.

CELLULOSE, considered physiologically, is the main product of vegetable life. It is the preponderating constituent of all vegetable tissues, and essential to the cell as the structural unit. For while the synthesising activity of the cell is identified with its protoplasmic contents, its main function is the building up of non-introgenous substances, of which cellulose is the chief, elaboration of cellulose, in fact, is synonymous with growth. As an organic product it is known to us in the specialised and diversified forms of plant structures, which are employed in industries either in their original

form or after treatment for separation of their fibrous components.

Cellulose is the basis of our staple textile and paper-making industries, and, when modified by various chemical treatments, of modern high explosives, 'celluloid' products and 'artificial silk'

Cellulose, as a component of vegetable foodstuffs, and in connection with agriculture, is economically important, and is a dominant factor of the cycle of processes, organically interdependent, of the vegetable and animal worlds.

By cellulose, as a chemical individual, we understand usually the residue from the exhaustive alternate treatment of vegetable tissues with alkaline solvents and weak oxidants. Generally, no cellulose is isolated on the large scale except as the result of a chemical treatment, more or less drastic, of fibious raw materials; and all that we know of cellulose as a chemical individual results from the study of products so The typical cellulose is the substance of bleached cotton, the bleaching process such as above described removing the substances other than cellulose with which it is associated in the plant. So obtained, it is a white substance, translucent when viewed under the microscope, but more or less opaque in the mass. This appearance is a consequence of the peculiar form of the cotton fibre, a flat thin-walled tube, the fibrillæ of which are membranously disposed.

The following brief outline of the special chemistry of cellulose necessarily precedes an account of its industrial applications, so far as they involve chemical principles or processes of treatment

CELULOSE —Generally, the non-nitrogenous skeleton of plant structures Type the fibre substance of cotton, freed from associated 'impurities' by processes of (1) alkaline hydrolysis and oxidation (bleaching), (2) digestion with hydrofluoric and hydrochloric acids, to remove mineral impurity or ash constituents

Composition — Elementary C 444, H, 62; O, 494, whence the empirical formula,  $C_6H_{10}O_5$ 

Constitution.—By solution in sulphuric acid as disulphuric ester  $n [C_6H_8O_3(SO_4H)_2]$  and ultimate hydrolysis, the colloidal aggregate is resolved with formation of dextrose in large proportion (Flechsig Zeitsch physiol Chem. 7, 523) The identity

$$[n C_6 H_{10} O_5] + n H_2 O = n C_6 H_{12} O_6$$

which has been assumed, has not been verified experimentally, a maximum yield of 90–100 p c. dextrose [theory, 111 11], but only indirectly estimated and accompanied by acid products, leaves the problem undetermined (Ost and Wilkening, Chem. Zeit 34, 461) See also recent critical discussions (Chem. Soc. Trans. 1918, M. Cunningham. Cross and Bevan). Cellulose is thus variously regarded.

- 1 As a polyhexose anhydride, (a) aldose, (A. G. Green, Zeitsch. Farb. Chem. 3, 97), (b) ketose (Fenton; Cross and Bevan, Chem. Soc. Trans. 82, 361, 366).
- 2. As a polycyclohexane derivative, this view is based on further considerations of density and molecular volume (Cross and Bevan, Ber. 1909, 42, 2198.
  - 3. As an essentially labile aggregate, that is

of variable constitution or configuration, varying according to the action of reagents (Cross and Bevan, Chem. Soc. Trans. 85; Cellulose Researches, n 1905).

Cellulose and water.—Hydration Constitutional water is retained by cellulose in the airdry condition Cotton holds 6-8 pc, varying with temperature and moisture of atmosphere Cotton which has been hydrated by treatment with alkalis (mercerisation) or solvents (infra) or by prolonged attrition in presence of water, shows an increased capacity for moisture when air-dry, and retains 9-11 pc Conversely, the action of condensing acids (HCl, HBi) produces modified aggregates ('hydrocellulose') with a lower capacity (3-5 pc.) Cellulose separated from its solutions, or regenerated from its derivatives by the action of reagents, shows a maximum hydration capacity; the colloidal or gelatinous hydrates having the composition (in contact with water at ordinary temperatures)

20-10 p c. Cellulose Water 80 - 90

Cellulosesolvents -Insoluble in all andneutral solvent liquids, cellulose is dissolved by .

1. Concentrated solutions of zinc chloride (40-50 p c ZnCl<sub>2</sub>), on heating at 80°-100°, when previously hydrated, at lower temperatures.

2 Solution of zinc chloride in twice its weight of HCl Aq (35 pc HCl) this reagent rapidly attacks and resolves the aggregate. Aqueous hydrochloric acid of maximum concentration (>40 pc HCl) acts similarly (Wilstatter, Berl. Ber 46 (1912) 210) but with fixation of negative groups (Cl) in stable union with the products (M Cunningham, Trans Chem Soc lc).

3. Solutions of cuprammonium hydrate containing .

Cu, as CuO  $n{\rm H}_2{\rm O}$  . . 2 5–3 5 p c NH  $_3$  (as NH  $_4{\rm OH}$ ) . . 15 p c

Ethylene-diamino-cupiic solutions act more powerfully (W Traube, Berl Ber. 44, 3319 (1911)

From solutions 1 and 3, the cellulose may be recovered quantitatively, but as a hydrate and otherwise modified in constitution

4 Concentrated solutions of certain metallic sulphocyanides at high temperatures, 120°-140°. Reactions —The above reactions resulting in solution of the cellulose are characteristic; otherwise it is exceptionally non reactive dilute solutions of iodine, in presence of certain dehydrating agents, it is coloured blue

Cellulose compounds A. Synthetical derivatives. Esters.

(a) Nitrates. By direct reaction with nitric acid, usually in presence of sulphuric acid, in which case unstable mixed esters are formed as a stage in the reaction, the NO<sub>3</sub> displacing the SO<sub>4</sub>H residues (Cross, Bevan and Jenks, Berl. Ber. 34 (1902) 3496) The esters are formed without sensible structural modifications are purified from residual SO<sub>4</sub>H by prolonged boiling with water, and are then 'stable.' A series of these esters is known, the highest approximating to the trinitrate (C<sub>6</sub>) (gun-cotton); the intermediate terms—dinitrate—being soluble in ether-alcohol (collodion cotton), the lowest having physical properties but little different from the original cellulose.

These esters are variously formulated as

nitrates of a reactive unit of C<sub>6</sub>-C<sub>12</sub>-C<sub>24</sub> dimensions.

Solvents.—The special solvents of these esters are acetone, ether-alcohol, nitrobenzene.

A diversified range of solvents is enumerated in a paper by H. dc Mosenthal (J. Soc. Chem. Ind. 1907)

Saponification.—By certain alkaline and reducing agents (alkaline sulphydrates) the nitric groups are eliminated and cellulose regenerated.

(b) Acetates. By reaction with acetic anhydride under various conditions. (1) at 110°. direct formation of monoacetate (C, insoluble in all neutral solvents and in the solvents of cellulose (2) At 140°-160° · formation of higher acctates, attended by solution in the reaction mixture (3) In presence of catalytic agents  $(ZnCl_2-H_2SO_4-H_3PO_4)$  at intermediate temperatures, H<sub>2</sub>SO<sub>4</sub> determines reaction at 25°-30° The extreme products are mixtures of tri- and higher acetates (4) When the reaction mixtures are diluted with hydrocarbon, the fibrous cellulose may be acetylated without solution or visible structural change

Industrially important acetates are the acetone soluble esters C<sub>12</sub>H<sub>15</sub>O<sub>5</sub> OAc<sub>5</sub> produced by acetylation with mixtures of Ac2O and H2SO4. These are the basis of aeroplane varnish-dopes

Solvents of the higher acetates, are chloro-

form, acctone, phenol

For an exhaustive account of the industrially important types, see E C Worden, J Soc Chem, Ind 1919.

Saponification -The acetyl groups may be removed by boiling with alkaline solutions, the cellulose being regenerated In quantitative determinations the saponification may be effected by boiling with normal sodium hydroxide diluted with an equal volume of alcohol

(c) Formates—Cellulose and formic acid interact at 60°-100°, in presence of catalysts the reaction proceeds to complete solution limit of esterification appears to be the diformate (C<sub>6</sub>), but the product is unstable, decomposing spontaneously with liberation of formic acid (1 mol). The reactions are not those of simple ester (anhydride) formation; the celluloses in reaction appear to undergo constitutional change (Cross and Bevan, J. Chem. Soc. Ind. 1911, 99, 1450, Berl and Smith, Berl Ber 1907, 40, 906, Woodbridge, J. Amer Chem Soc 1909, 31, 1067, Dewing, ibid 1911, 33, 1179)

(d) Acid-sulphuric esters By the action of sulphuric acid an extended series of esters is formed, which have been described as cellulose sulphune acids But they are certainly derivatives of products of resolution. The first stage results in the formation of a disulphuric ester  ${\rm C_6H_8O_3(SO_4H)_2},$  but its relationship to the parent complex is doubtful. The ester is soluble in water, the Ca, Ba, and Pb salts are insoluble alcohol By progressive hydrolysis the cellulose is ultimately resolved in the direction of dextrose (see M. Cunningham and Cross and Bevan, l.c).

(e) Aceto-sulphates and mixed esters, con taining the SO<sub>4</sub>H residues associated with acetyl and other negative groups in combination, are obtained when sulphuric acid is allowed to act under regulated conditions simultaneously with other esterifying agents. Thus a mixture of acetic anhydride (50 parts), glacial acetic acid (50 parts), and sulphuric acid (4-6 parts), acts rapidly at 30°-40°. The first product appears to be a neutral body of the empirical formula

$$(4C_6H_7O)SO_4(C_2H_3O_2)_{10}$$

and under the action of water to undergo an internal hydrolysis, the SO<sub>4</sub> group becoming SO4H, which forms a stable combination with bases The Mg, Ca, Zn salts are insoluble in water, but soluble in acctone

(f) Benzoates result from the action of benzoyl chloride in presence of alkaline hydroxides monobenzoate (C6) is obtained by treating cellulose with a solution of sodium hydroxide of 10 pc. (NaOH) strength, and shaking with benzoyl chloride This benzoate is formed with only slight structural change. The dibenzoate (C<sub>6</sub>) is obtained by the interaction of benzoyl chloride and alkali cellulose (mercensed cotton) in presence of sodium hydroxide solution (15 p.c. NaOH) Its formation is attended by structural change; the fibrous cellulose is disintegrated, the dibenzoate being an amorphous substance The dibenzoate is soluble in acetic acid and chloroform

Mixed esters, containing the henzoyl and nitric residues, result from the action of nitric acid upon the benzoates Simultaneously a nitro- group enters the benzoyl residue.

Alkalı cellulose The fibrous cellulose undergoes considerable structural modification under the action of solutions of sodium hydroxide of 12-15 pc NaOH These reactions and phenomena were first studied by Mercer They are the basis of the process of mercenisation (q v)under which cotton-yarn and cloth assume a special lustro and silky finish. There is a definite synthetical reaction in the ratio C<sub>6</sub>H<sub>10</sub>O<sub>5</sub> · 2NaOH, which is a stage in the formation of the dibenzoate (supra)

The compound is completely dissociated by water; by treatment with alcohol an equilibrium is reached when the reagents are associated in the ratio  $C_{12}H_{20}O_{10}\cdot NaOH$ The alkali-cellulose hydrate, of composition  $\cdot$ 

 $\begin{array}{cccc} \text{Cellulose} & . & . & 30\\ \text{Sodium hydrate} & 15\\ \text{Water} & . & . & 55 \end{array} \right) \begin{array}{c} \text{cellulose} \cdot \text{ sodium hydrate} \\ \text{C}_6\text{H}_{10}\text{O}_5 & 2\text{NaOH} \end{array}$ 

is the first stage in the synthesis of

Cellulose xanthogenic acid, which results from the interaction of the alkali cellulose and carbon disulphide at ordinary temperatures sodium salt is soluble in water. It is an unstable compound, the solution undergoing spontaneous progressive change. The solution, which is highly colloidal, finally solidifies Bymeans of the characteristic reaction of the xanthates with iodine

$$\begin{array}{c}
\text{CS} & \text{OX XO} \\
\text{SNaNaS} & \text{CS} + \text{I}_2 \\
= 2\text{NaI} + \text{CS} & \text{S} & \text{CS}
\end{array}$$

the progress of the change may be followed, the essential feature being the elimination of the CS<sub>2</sub> residues with reaggregation of the cellulose units Well-marked stages in the series occur at the points denoted by the empirical formulæ  $C_{12}H_{19}O_9CSSNa$ ,  $C_{24}H_{39}O_{19}CSSNa$ . The former represents an equilibrium attained after the solution has remained for some hours at the solution has remained for some hours at the of the hypochlorites, permanganates; but when ordinary temperature; the latter is reached in the limit is pussed the oxidations which result

from three to four days. The cellulose under the reaction acquires a more acid character, an additional OH group combining with alkali. The lower terms of the series, e.q. C<sub>48</sub>H<sub>79</sub>O<sub>38</sub>. CSSNA, though insoluble in water or dilute saline solutions, are dissolved by the addition of sodium hydroxide The sodium atom in combination with the CSS residue is not attacked by By double weak acids such as acetic acid decomposition with soluble salts of Cu, Zn, &c, the corresponding xanthates are produced as insoluble colloidal precipitates.

In the above reactions the cellulose aggregate is maintained, the solutions of the derivatives are viscous and colloidal, but in the following

B Reactions of decomposition, which are determined by hydrolytic and oxidising agents, the directions of resolution are extremely various, and the relationships of the products to the original aggregate are undetermined

(a) Sulphuric acid dissolves the cellulose as a disulphuric ester, but decomposition attends the reaction, and on diluting and boiling, the hydrolysis is carried to the extreme molecular limit, the final product being dextrose (supra); but the reactions are not simple as generally stated, they are complicated by relatively stable union of the SO4H with the resolved cellulose 'residues,' and by tendency to reversion or condensation

(b) Hydrobromic acid in ethereal solution attacks the cellulose profoundly with production of brom-methyl furfural The formation of this compound indicates a previous or intermediate stage in which the products of resolution are molecular ketonic bodies of carbohydrate constitution (Fenton, Chem Soc Trans 1901, 361).

(c) Hydrochloric acid, in presence of water, dilute sulphuric acid, and acids generally, attacks the cellulose aggregate with production of a variety of derivatives (1) Insoluble: these are generally termed 'hydrocelluloses' They are disintegrated residues of the original fibres, they differ chemically from the parent aggregate in the presence of free aldehydic groups, and in readily yielding to the action of alkalis (2) Soluble molecular products, chiefly dextrines and dextrose.

(d) Alkaline hydroxides and alkalis generally in the form of dilute solutions have little apparent action on cellulose—even when treated at elevated temperatures Sodium hydroxide in solutions of concentrations of 12 p c. NaOH and upwards, combines with the cellulose, producing profound structural modifications (mercerisation), but without resolving the aggregate.

At higher concentration and temperature the cellulose is partially dissolved; but even under the conditions of a 'fusion' at 180° the resolution is limited to the conversion into alkali soluble modifications, which are precipitated in the colloidal form on diluting and acidifying. At higher temperatures (250°) and with larger proportions of the alkaline hydroxides, the cellulose is resolved into acid products of low molecular weight, chiefly acetic acid and oxalic acid

(e) Oxidants. The directions of oxidation of cellulose are likewise extremely diversified. The aggregate manifests considerable resistance to alkaline oxidants in dilute form, eg. solutions

CELLULOSE

are drastic in the sense that the soluble products are of low molecular weight, chiefly carbonic and The insoluble fibrous residues, oxalic acids. more or less disintegrated, are known as oxycelluloses. They contain free aldehydic groups, are easily attacked by hydrolysing agents, and on boiling with hydrochloric acid (106 sp gr) are decomposed with production of some furfural

Resolved by the action of concentrated solutions of the hypochlorites, cellulose yields chloroform and carbon tetrachloudes The hypobromites give the corresponding bromine derivatives Nitric acid (1 25 sp gr ) at 80°-100° converts cellulose into a series of coxycelluloses, which are resolved on boiling with calcium hydroxide into acid products, among which isosaccharinic and dioxybutyric acids have been identified. In the original oxidation small quantities of the higher dibasic acidssacchanic and tantanic acids-are produced, but the main products are oxalic and carbonic acids

With chromic acid an endless series of oxidations may be effected, the degree of action depending upon the proportion of the active oxidant and the associated hydrolytic action of The oxycelluloses produced are mineral acids distinguished by relatively large yields of furfural when decomposed by boiling HClAq (1 06 sp gr) In presence of sulphuric acid there ensues complete combustion, and the reaction is the basis of quantitative analytical methods.

(f) Resolution by ferment actions Under the actions of specific organisms the cellulose complex is totally resolved, the main products being methane, hydrogen, and carbonic and The decomposition may be associfatty acids ated with the action of an enzyme, but a remarkable feature of the process is the absence of intermediate products, at least in the cases hitherto investigated In the digestive tract of the herbivoia, cellulose is resolved, and from the investigation of the process, necessarily by indirect observations, it appears that, in addition to a destructive resolution to ultimate gaseous products, there occurs a resolution to proximate groups of high nutritive value, which are assimilated by the animal organism

(q) Resolution by heat · destructive distillation Cellulose heated at 100°-120° is dehydrated at 150° decomposition is evidenced by change of colour. At this early phase maltol appears to be formed The decompositions of cellulose at temperatures exceeding 250° are necessarily

extremely complex

The groups of products show an average proportion:

Solid 30 p.c.	Liquid 50 p.c.	Gaseous 20 p.c.
Charcoal or pseudo-car- bon	Containing acetic acid(2p c.), methyl alcohol (7 p.c.), acetone, furfural (12 p.c.)	Chiefly CO and CO,

the actual proportions and composition of these mixtures varying with the temperature and duration of their heating

By 'dry' distillation in value of cellulose

(and starch) large yields of lavoglucosan are obtained. Pictet and Sarasin, Compt Rend. 166 (1918) 28.

111

General view of the decompositions of cellulose. It is clear that the cellulose complex breaks down under destructive influences, in directions depending upon the nature of the attacking agent, its concentration, and all the surrounding physical conditions. The study of surrounding physical conditions these decompositions has thrown but little light on the actual nature and constitution of the cellulose aggregate. for the reason, perhaps, that we have endeavoured to maintain a basis of interpretation such as is applicable to ordinary molecular compounds or complexes. Cellulose is a typical amphoteric colloid the reactions with hydrolytic reagents suggest many points of analogy with the state of a complex salt in solution (in water) Its component groups of unknown, but possibly small dimensions, appear to exist under a certain internal stress which determines migrations of H and OH under reaction. While, therefore, we have reduced the reactions of cellulose to a short systematic expression, it is clear that the main aspect of the subject cannot effectually be so treated, because it involves as theoretical basis the actual state of matter and the dimensions and distribution of the reactive unit-groups in the cellulose complex, and this basis is, as yet, largely undetermined (see Cross and Bevan, Cellulose Researches, 1905, 1910)

the typical The cellulose group.  $\mathbf{From}$ cellulose we pass to the diversified group of Then general characteristics are those of the prototype, the variations they present are especially such as involve the unde-termined factors of constitution. With these there are certain correlative variations which afford an empirical basis of classification. These are (a) the degree of resistance to hydrolytic and to oxidising agents, (b) the percentage yield of furfural when decomposed by boiling HClAq, (c) elementary composition, in respect of the ratio C O

The fibrous celluloses are grouped as follows.

Туре	Cotton sub-group A. Bleached cotton.	Wood cellulose sub-group B Juta cellulose.	Cereal cellulose sub-group C Straw cellulose
Hygroscopic moisture Elementary composition C O Furfural Other cha- racteristics	6-8 p c.  { 44 0-44 4   50 0   0 1-0 4   No active CO groups	9-11 p c 43 0-43 5 51 0 3 0-6 0 Some free CO groups	9-12 p c 43 5-44 5 52-50 12 0-15 0 Variable reactivity of CO groups.

See Esparto Cellulose and the Problem of Constitution: Cross and Bevan, Chem. Soc. Trans. 113 (1918) 182.

Of these groups the following points may be noted.

A. Comprises, in addition to cotton, other industrially important celluloses, e.g. flax, hemp, and rhea. They occur in the plant-world in association with compounds easily removed by the action of alkalis. They pass through the

cycle of reactions involved in their solution as xanthates, without hydrolysis to soluble derivatives

B These celluloses are obtained as products of decomposition of a compound cellulose. They may be regarded as partially hydrated or hydrolysed. They are more readily attacked by hydrolysing agents and, in the xanthate reactions, are partially resolved to alkali-soluble, derivatives.

U These celluloses are in most cases a complex of structural elements, and not homogeneous chemically They are still less resistant than the preceding group, and more especially the furfural-yielding components, which are selectively attacked under certain conditions

The cellulose groups, as above, pass by imperceptible gradations into a heterogeneous class of natural products, which, while possessing some of the characteristics of the celluloses proper, are so readily resolved by hydrolytic treatment that they must represent a very different constitutional type or types. To this group of complex carbohydrates the class-name hemicellulose has been assigned. They are structurally different from the fibrous celluloses, occurring mostly in the cellular form (parenchyma, &c.). They differ in physiological function, and in being readily resolved by hydrolysis into the crystalline monoses.

The foregoing exposé will enable us to group the industrial and manufacturing applications of cellulose in reference to first principles as

follows .—

GENERAL.—The uses of the fibrous celluloses, more particularly of cotton and the cotton group, in which we may include flax, hemp, and ramie, depend, apart from the primary condition of length of ultimate fibre, upon the extreme chemical resistance of the fibre substance, which is a function both of the constitution of the ultimate groups and of their mode of union to constitute the aggregate itself

Insensitiveness to oxidation is an obvious condition of such stability as is required for textile fabrics and paper, and, as evidence of resistance to atmospheric oxygen, we have the fact that flax and cotton fabrics and papers composed of these fibres have been handed down to us from antiquity, with little change. Next in order of importance is resistance to water and alkaline hydrolysis. This resistance enables textile fabrics to withstand periodical cleansing or laundrying, in which alkalis are the principal agents. In the purifying of textile fabrics by the bleaching processes and the preparation of paper pulps, the same negative qualities are of first importance.

Textules are subjected to severe processes of alkali treatment, and to the action of bleaching agents, such as hypochlorites, without evident

chemical or structural modification.

On the other hand, there is a limit of the resistance to exidation, and any unduc action of the ordinary bleaching solutions leads to the formation of 'oxycellulose,' and structural weakening as an attendant result.

While extremely resistant to alkalis, the celluloses are sensitive to acids, and here again in bleaching, finishing, and dyeing processes, the incidental acid treatments require careful adaptation to the limit of sensitiveness of the cellulose.

SPECIAL.—Constitutional moisture and hydration effects. These involve a range of phenomena of obvious industrial importance. The normal moisture of cellulose has to be taken into account in the finishing of textiles, and their package for shipment. The condition of the fibres as regards hydration, of which the capacity for combining with atmospheric moisture is an index, directly affects the finishing processes which are chiefly the mechanical operations of mangling, calendering, and beeting.

The further hydration of the celluloses which takes place in the beating operations of the papermaker is of considerable practical importance, and affect the quality and character

of the papers in a very great degree

Cellulose and ash constituents. The normal ash of the cellulose of the cotton group is small in proportion, and we are not able to affirm that these inorganic constituents have any determining relationship to the constitution of celluloses, though this is an accepted probability.

It is a problem for colloidal chemistry to establish this relationship in specific terms. Industrially, the question arises only with regard

to the aggregate.

Thus in the purest form of cellulose required as filter paper for the quantitative separations by chemists, it is of importance that the ash components should be reduced to a minimum

This minimum of 0.05 p.c. by weight, is attained by digesting the cellulose with hydrofluoric acid and hydrochloric acid, and thoroughly

washing the paper in pure water.

Pure unsized cellulose paper thus treated constitutes the well-known 'Swedish' paper, so called from its country of origin, but now produced in this country and other centres

Cellulose as a typical colloid enters into characteristic combinations with inorganic oxides, and this property is an important basis of the processes of the dyer and printer.

Oxides of the polyvalent metals, notably of aluminium, chromium, iron, tin, and lead, are taken up by the celluloses from solutions of their salts, and the cellulose thus mordanted has increased special affinities for colouring matters

A more special industrial application of this property is the method for making the mantle, now in common use for intensitying coal-gas illumination (Auer-Welsbach). The colloidal oxides of thorium and cerum in suitable proportions are fixed upon cellulose fabrics, which are then dried and ignited. The result is an inorganic skeleton of the original fabric, prepared in any desired form, and composed of the oxides in question

The particular form fulfils the conditions of maximum illuminating surface, together with the desired degree of mechanical resistance to stand shock and wear and toai under the particular conditions of employment

Cellulose and solvents It is in the solutions of cellulose or its derivatives that the particular properties of the cellulose aggregate are evident

In the case of the aqueous solvents observation shows that the cellulose, employed as fibre, passes through phases of progressive hydration with considerable distension until the final stage of homogeneous structureless solution is reached. The general character of the solutions is their high viscosity, which imposes a limit of from 7 to 9 p.c. cellulose for solutions which

require filtering.

The technical applications of these solutions depend upon the fact that the cellulose is brought into the plastic homogeneous structureless condition, in which it may be formed or shaped at will, and in finally reverting to the solid state preserves its structural continuity through the successive phases of dehydration.

The solution in zinc chloride has been used for making threads to be carbonised for electric

incandescence lamps.

The solution is forced through narrow orifices into alcohol, which precipitates the cellulose as a transparent hydrated solid, retaining zinc oxide (hydrate) in combination. The solid hydrate is formed and manipulated in continuous length, and, when freed from zinc oxide and dried, is a structureless cylinder of cellulose. Heated out of contact with air, it is converted into a 'pseudocarbon,' which constitutes the filament in question.

The solution of zinc chloride in hydrochloric acid has been employed for converting cellulose papers into a special agglometated product or board, which is used as an insulating material and for structural purposes. The sheets or webs of paper are treated with the solvent by immersion, and the material thus attacked is then welded into the composite fabric by means of the adhesion of the viscous cellulose solution resulting from the attack.

The composite sheets or webs so produced are finally compacted by pressure, and then exhaustively washed to remove the solvents

Cuprammonium There are various industrial applications of this characteristic solvent Textile materials are treated by passing through the solution. The cellulose is attacked superficially, and under the pressure of rolls the gelatinised product is distributed uniformly, the fabric is then dried. The fabric is thus coated with a variush composed of cellulose and the hydrated copper oxide, of greenish colour which renders the fabric impervious to water and rotpicof.

Papers are similarly treated, the sheets of webs being welded together under pressure. These tabries have been known for a number of years as the 'Willesden' waterproof fabries.

The most important industrial application of the solution is in the manufacture of artificial cellulose threads, which are produced in various grades of fineness, and known as 'Artificial Silk,' 'Monofil' (Lustra Cellulose, Soie Artificielle, Glanzstoff, Crin).

The process in its simplest form consists of the production of the solution by treating bleached cotton with the cuprammonium solvent, and filtering through fine metal-wire cloth to remove incidental impurities

The solution is projected through orifices of varying diameter, according to the size of the

threads to be produced, into solutions which

precipitate or coagulate the cellulose.

The setting solutions are of two types, acid and alkaline.

The cellulose is precipitated as a hydrated solid, in continuous length, and is drawn forward at speeds varying from 20 to 50 metres per tures. Vol. II – T.

minute, according to the diameter of the thread and the conditions of the precipitation

In the case of fine filaments, constituting artificial silk, several of the threads so produced are twisted together to form a compound filament or yarn suitable for weaving purposes. The products are dried under special conditions, to secure maximum lustre and elasticity

Cellulose esters The application of the nitrates to high explosives is an application of cellulose that stands apart and depends, of course, upon the fundamental chemistry of the product.

The ratio of oxygen is such that the products contain the elements for their complete conversion into gaseous products when raised to the reactive temperature.

The nitrates in the fibrous form in which they are produced, are explosives of the disruptive or blasting type, and were exclusively

so used for many years

The type of explosion required for the propulsion of projectiles, is a restrained or graduated combustion, which is determined by a change of form of the intrates. These are convertible into plastic modifications under the action of their specific solvents, and in this state they may be formed or fashioned into threads, cylinders, or solids of any desired form or dimension. The combustion of these structureless solids, in its aggregate explosive effect, may be made to fulfill any desired ballistic conditions.

The most interesting case is that of the employment of nitroglycerine, itself a high explosive, and the basis of dynamite, to reduce the cellulose nitrate, acting as a solvent, to the structureless condition.

Mixtures of introcellulose and introglycerine, in suitable proportions, are drawn to threads and constitute with certain admixtures the modein explosives cordite, ballistite, &c. (see Explosives)

The collulose intrates, treated with neutral solvents and subjected to a process of milling, are reduced to a structureless plastic mass. This may be reformed into solid substances which may be produced in any desired form or dimension. In sheets or film they constitute the basis or transparent carrier of photographic emulsions.

In solid form they are known as celluloid, xylonite, &c., and find endless applications both useful and ornamental (v Celluloid).

The solutions of mitrates in neutral solvents, such as ether-alcohol (collodion) or amylacetate, are used as varnishes and lacquers for various purposes

purposes

The cellulose intrates in ether-alcohol solution are employed industrially in the manu-

facture of artificial silk

This, in fact, was the pioneer invention 'of Chardonnet,' and dates from 1883

The process of spinning consists in forcing the solution through capillary glass jets, either into air (Chardonnet) or into water (Lehner).

The thread is thus formed and obtained as a nitrate, in which form it is twisted into compound textile filaments, which are reeled into

skeins of yarn.

A process of saponification for removing the nitric groups follows. This consists in digesting the nitrate yarn in solutions of ammonium-magnesium sulphydrates at ordinary temperatures.

114 CELLULOSE.

As a result of the cycle of operations, the denitrated cellulose is obtained and constitutes the artificial silk of commerce.

The xanthogenic esters of cellulose in aqueous solution (viscose) are the basis of the production of artificial textile fibres, and the viscose artificial silk has displaced the pioneer product as well as to a great extent the cuprammonium product (Glanzstoff) The manufacturing process comprises the following operations:—

(1) Wood cellulose is converted into alkalicellulose (hydrate).

(2) After ageing, the product is treated with carbon disulphide in closed vessels

(3) The xanthate is dissolved and filtered, and after ageing is projected through fine apertures into acid solutions, which cause rapid decomposition and reversion to 'cellulose'

(4) The reprecipitated cellulose hydrate is

dried and dehydrated under strain.

For a full description, see L P. Wilson,

J. Soc Chem Ind. 1917, 817.

Cellulose acetates are the analogues of the nitrates with the radical distinction of being non-explosive, and, in fact, eminently stable compounds.

By treatment with their special solvents, they are dissolved to homogeneous viscous solutions, and they may be fashioned into films and solids as are the nitrates.

So far the industrial applications of the acetates have been limited chiefly by the

prohibitive costs of production

One important application, however, produces results so very much sui generis and in an arena where costs of production are a minor consideration, that it has led to extensive developments of the manufacture. Solutions of the acetate (6–8 pc) in 'organic' solvent liquids constitute a viscous varnish-dope, which when applied to a (stretched) fine textile, cotton or flax cloth, produces a considerable shrinkage, and this under strain, as on the wing planes of the aeroplane, gives a taut, smooth surface, at the same time waterproofing the fabric and thus fulfilling the exacting technical specification of the effect required for these special and highly specialised structures.

The acetate is also used as a varnish covering of fine wire used in electrical construction. The acetate is a high insulator with exceptionally low

inductive capacity.

Cellulose acetate has already been spun to artificial silk and coarser textile threads, but there has been so far no industrial development of the processes

The technical interest of such a product is that it is employable as an ester, whereas the nitrate is entirely unsuitable, as such, for textile

applications.

Cellulose and alkalis (Mercerisation). The action of alkaline hydroxide in strong solution (15-20 p c. NaOH) upon cotton textiles, is the basis of a special industry. By converting the cotton fibre into the modified form, cotton yarns and fabrics are changed in appearance and acquire a silky lustre and touch.

The goods to be 'mercerised' are passed on the nature of the fibre substance. So too through a solution of caustic soda in the cold. This produces the effect of considerable shrinkage accompanying the change of form and treated for the isolation of the fibres. We must

dimensions of the ultimate fibre, from a flat membranous, to a thick-walled cylindrical homogeneous tube.

This shrinkage is resisted by placing the goods under the condition of mechanical strain, either during the process of alkaline treatment, or, after allowing the shrinkage to take place, the goods may be afterwards stretched to their original length.

The fundamental observations which condition this process were made by J. Mercer (1850–1860), but they remained undeveloped until late in the century (1880–1890).

The industry is now a considerable one, and is to a certain extent independent of fashion

(see MERCERISING).

Cellulose and sulphuric acid. Whereas cellulose reacts with the more concentrated acids ( $H_2SO_4-H_2SO_4,H_2O$ ) to form sulphuric esters, the acids of 670 to 780 pc  $H_2SO_4$  (approximately constant) matelyH2SO4,2H2O-H2SO4,3H2O) hydrate, and dissolve the cellulose to a colloidal solution, from which, if immediately diluted, the cellulose is precipitated as a gelatinous hydrate ('Amyloid').
These reactions are industrially applied in the preparation of 'parchment paper,' or 'vegetable parchment' A pure cellulose paper is passed through a bath of acid, and immediately, in continuous length, into water, which arrests the action of the acid and reprecipitates the cellulose superficially dissolved as a structureless colloid. After washing from the last traces of acid, the web of now parchmentised paper is dried under special conditions of strain to control the shrinkage of width (10-13 pc), so as to maintain an even surface and prevent deforma-To overcome the brittleness product, which would result from total drying, it is treated with solutions of neutral hygroscopic substances, such as glycerol or glucose, before passing on to the drying cylinders. A sufficient proportion is retained in the finished product to ensure a normal 'humidity.

The total hydrolysis of cellulose to fermentable 'sugars' is effected by dissolving in the more or less concentrated acid, and, after allowing the acid to act until the solution becomes limplied, diluting and heating the dilute solution. The acid is neutralised with chalk, filtered from the insoluble calcium sulphate, and concentrated. The industrial value of this piocess is obviously limited by the relatively low values of starches and the more economic conversion of these by simpler means and to superior products.

simpler means and to superior products

The compound celluloses (Vegetable fibres).

While the simplest form of vegetable life is
the cell, and the cell is, moreover, the universal originating unit, we observe, as we
ascend the scale of plant organisation, an increasing variety of differentiations of this simplest structural unit graduating into the fibres
and vessels of the stems of the higher flowering
plants. It is to the presence of such elongated
cells as warrant the designation 'fibre' that
these stems mainly owe their powers of bearing
strains and resisting fracture. The uses to which
these fibres are put in the arts depend upon the
same characteristics of form, and obviously also
on the nature of the fibre substance. So too
the anatomy of fibrous stems is necessarily
a main factor of the processes by which they are

CELLULOSE 115

here except from the general anatomical view we are about to take the most important of the vegetable fibres, viz cotton, which as a seed hair is distinct physiologically, and differs structurally from all other fibres employed for textile

purposes.

The 'ultimate' fibre is a hollow tube, the walls being thin and membranous, it is attached at one end to the testa of the seed, the other being free, and there is no adhesion of the fibres to one another. The preparation of the textile fibre involves, therefore, only a mechanical treatment, and that of the simplest kind, for separating it from the seeds. The other vegetable fibres employed in the arts are without exception constituents of complex structures (stems and leaves) which require a more elaborate mechanical operation for their separation, in addition in many cases to a preparatory chemical treatment

The fibres of stems are arranged according to the two main structural types of growth, the exogenous and endogenous In the exogenous stem of annuals (dicotyledonous) they constitute a localised tissue, the bast, which is an annular ring external to the wood and beneath the cortex. In the endogenous stem the fibres proper (bast fibres) are generally aggregated together with vessels into bundles known as fibrovascular bundles, which are scattered more or less irregularly throughout the cellular matrix

In regard to the leaf, the two great divisions of the phanerogams are also contrasted, the fibro-vascular bundles of the former are not only nregularly disposed, but interlaced (reticulate venation), whereas the leaves of the typical monocotyledon are characterised by a parallel arrangement of the bundles, and fulfil, theretore, a necessary condition of separation for tex-Fibres of this class are amongst tile purposes the most useful that we have There is a distinction to be noticed, though rather of a technical than anatomical nature, between the vegetable fibres and 'fibrous substances' former we understand isolated bast fibres or fibro-vascular bundles, such as can be spun into a continuous length or yarn A fibrous substance is any vegetable structure containing fibres Many of these do not admit of being economically treated for the isolation of a textile fibre, but, on the other hand, are resolved by suitable chemical treatment into a fibrous mass or pulp, available for paper making. Many of our most valuable raw materials for this industry are of this nature. We may further distinguish, according to the above classification, (1) evogenous bast tissues, such as those of the Adansonia and Broussonetia (paper mulberry) and entire stems (woods), and (2) endogenous stems (straws and bamboo), and leaves (esparto). In this more miscellaneous group also we should include the refuse from the preparation of textile fibres, such as the root ends of jute (butts), the waste from the flax-scutching process

(infra), &c.
We have already excepted the cotton fibre from the above classification, indicating a structural as well as a physiological distinction from other textile fibres. These are, in fact, always obtained and spun in the form not of ultimate fibres, but of bundles of these. The ultimate bast fibre is also a hollow tube, but more or less

thick walled and with tapering extremities, in section it is seen to be cylindrical or polygonal The dimensions are remarkably uniform for the bast of any given plant or species, but exhibit wide variations from species to species. Thus the average length of the bast cell of jute is 3 mm, and of flax 28 mm. The textile 'fibres' are aggregates of these ultimate fibres, and it is convenient to employ for them the special designation of *filament* This complex spinning unit varies in the number and degree of cohesion of the constituent fibres These variations even in the same plant or species are very considerable, but more so as we pass from species to species. Expressed in another way, the textile fibres vary very much in regard to fineness and divisibility of the filaments. These, it is scarcely necessary to say, determine in a very important sense their spinning qualities. The nature of the fibre substance is a self-evident factor of the mechanical properties of the filaments, and lastly, though less obviously, the length of the ultimate fibre, which greatly influences the strength and durability of the yarn.

To the spinner the chemical composition and properties of the fibre substance may appear of altogether secondary importance; but they are obviously of first importance in regard to the application of the yarn, as of the goods woven from it. Supposing them put into use without further preparation, the durability of the fabrics will depend upon their resisting the disintegrating action of air (oxygen) and water, reinforced by heat and light But a large proportion of the textile fabrics of commerce are supplied to the consumer in the bleached state, the process of bleaching consisting in the removal, by a more or less drastic treatment with reagents of those fibre constituents to which the colour of the raw material is due These processes being the subject of a special article, we shall not require to deal with them at all in detail. The principles underlying them, however, have a close and obvious connection with the properties of cellulose, and the relationships subsisting between the cellulose of the vegetable fibres and the non-cellulose constituents with which each is associated in the plant, and from this point of view we proceed to give a short account of the

## CHEMISTRY OF THE COMPOUND CELLULOSES.

There are many reasons for regarding cellulose and the non-cellulose constituents of vegetable tissues and fibre substances as united chemi-While this is certainly true of the cutocelluloses (infra) there is evidence that in the ligno celluloses the bond of union is rather that of mutual interpenetration or 'adsorption' At the same time the combination is equally resistant to reaction as that of a union of 'residues' in the strictly chemical sense. Taking cellulose as the constant, the basis of our classification of the compound celluloses will be the nature of the non-cellulose, the characteristics of which are as distinct as they are general Physiologists recognise three 'modifications' of cellulose in the plant, and these we term 'compound celluloses.' We shall find the compound celluloses.' distinguishing chemical features of the three groups of compound celluloses as well marked 116 CELLULOSE.

as their functional correlations, and the corresponding classification has a 'natural' in addition to its chemical significance. We shall consider first the group which is of least, because of only incidental, importance in relation to our subject.

Cutocelluloses (Adipocelluloses). The protective tissues of plants, which are, moreover, cellular rather than fibrous, viz cork and cuticle, are composed of a substance of marked water-resisting characteristics. In this property, as well as in empirical composition, the cuticular substance designated cutose by Fremy resembles many of the waxes. The following percentage numbers are the results of the analysis of the cuticle of the apple purified from wax and adventitious matters: C, 73 66, H, 11 37, O, 14 97.

Cork is not a simple substance, but contains a number of products, wax, fatty esters and anhydrides, proteid residues, and tannins, &c, associated with the matrix or skeleton substance, which is of the nature of cutose Both cutose and cork yield a residue of cellulose when treated with special reagents (sulphites) for the resolution of its union with the wax-like non-cellulose The latter 1 is resolved by boiling with strong solutions of the alkalis into two fatty acids, a solid acid designated stearocutu and a liquid called oleo-The investigation of these cuticular substances has not been sufficiently extended to warrant our entering into further detail in regard to their composition and properties We may regard them as compound celluloses resolved by alkaline saponification into cellulose and a mixture of fatty acids of peculiar properties, but altogether unknown constitution They may be conveniently grouped under the term adipocelluloses, which broadly expresses their chemical relationship As regards their distribution, they occur chiefly in the epidermal and cortical tissues of leaves and stems, but the observations of Fremy lead to the conclusion that they are by no means confined to these, and there is evidence of their presence, if only in minute proportions, in the fibres themselves, but their occurrence, chemically speaking, may be regarded as adventitious, the fibre substances belonging to essentially different classes of compounds. These adipocelluloses, moreover, in most of the technical applications of vegetable materials which come within the province of this article, have to be got rid of An instance of an epidermal tissue (cutocellulose) used as such is the Raffia, which consists of the cuticular tissues of Raphia ruffia, which in horticultural work replaces the linden bast or 'bass' of a former age A more exhaustive investigation of Raffia as a type of cuticularised tissue has established the constitution of the cuto-cellulose of the mixed tissue as an ester of an oxidised 'cellulose' and a fatty acid

$$C_{17}H_{32}O_3 = C_{16}H_{30}(OH) COOH$$

But investigations in this field are complicated by the resistance of the mixed tissues to proximate resolution; the true cuto-cellulose when obtained is more or less modified by the chemical treatments required for its separation,

According to Fremy, it is the cutose itself which is so resolved, but the author's experiments have always yielded a residue of cellulose. and the yields are small (Cross and Bevan, J. Soc Dyers, 1919, 35, 70).

When the textile fibres are isolated from compound structures this is accomplished for the most part by the mechanical processes of preparation, scutching, hackling, &c When, however, the tissue intervening between the fibres and the cortex or epidermis is of small extent, the separation is never complete. In flax, for instance, residues of the epidermal tissue remain associated with the fibre, and the relative difficulty of attacking these substances by chemical means is a factor of the laborious process of bleaching linen goods In isolating the so-called New Zealand 'flax' much greater difficulty This fibre consists of the fibrois experienced vascular and other fibre bundles of the swordshaped leaf of the Phormium tenax, and these are in such close and intimate contact with the epidermal tissues that their isolation is always very incomplete In those raw materials of the paper maker which consist of the stems and leaves of monocotyledons (straw and esparto), the epidermal adipocelluloses constitute a sensible proportion. They are resolved in this case by chemical means, which require to be much more drastic in consequence. The fibrous pormore diastic in consequence tions of these materials, on the other hand, are made up of a compound cellulose belonging to the next group which we shall have to consider, an important characteristic of which is the comparative ease with which they are attacked and resolved by alkalis Although, therefore, the connections of these cuticular celluloses with the fibres are for the most part only inducet or incidental, we see in the typical instances above cited that they have to be taken into account in relation to a great number of the treatments of cellulosic raw materials, both mechanical and chemical

Pectocelluloses. This, the second division of the compound celluloses, is characterised by the non-cellulose constituents belonging to the group of pectic substances or pectins. The pectins are carbohydrate-colloidal aggregates or supposed complexes of hexosanes and pentosanes, with a small proportion of acid groups combined as lactone or ester. Sensitive to acid hydrolysis and feriment actions, they occur naturally in the most varied torms. Thus in fruit juices they exist as colloidal ropy solutions. In the more permanent fibrous tissues they are insoluble in water (Tollens, Annalen, 286, 278–292, Cross, Ber 1895, 2609)

In composition they are more complicated than the celluloses, and they contain less carbon and more oxygen. In this and in many of their properties they resemble the oxycelluloses above described. Generally, they yield gelatinous hydrates, and under the action of hydrolytic agents they undergo a succession of changes resulting in the conversion of the comparatively mert members of the series pectose and pectin into highly acid derivatives, such as pectic and metapectic acids, bodies which, in addition to being acids, have properties indicating the presence of aldehydic groups

The typical pectocellulose is the flax fibre. The purified bast of a Russian flax was found by Kolb to contain C, 43.7; H, 5.9; O, 50 4.

On boiling with dilute alkaline solution about 20 p.c. of its substance is readily converted into

soluble derivatives, of which a considerable proportion is a body identical with Fremy's pectic acid (C 42 0 p c., H=48; i.e. C<sub>16</sub>H<sub>22</sub>O<sub>15</sub>) The fibrous residue from an exhaustive treatment with alkaline solutions, amounting in most flaxes to about 75 pc of the original weights, is approximately pure cellulose In this experimental account of its constitution we disregard such adventitious admixtures as cuticular tissue (adipocellulose) and wood or 'sprit' (lignocellulose), the influence of which may be calculated and eliminated from the result, but the presence of these in commercial flax introduces complications of very great moment in regard to the spinning of the fibre and the bleaching processes to which flax goods are subjected In regard to the latter, an important technical difficulty is experienced in washing flax yains after a 'chlorine' (hypochlorite) bleaching treatment 'Active chlorine' remains persistently in the goods. This is now elucidated, as due to the formation of chloroumine delivatives, from residues of chlorophyll and other nitrogeneous products containing amino- groups always present in cuticular tissues (Cross, Bevan, and Buggs, J Soc Chem Ind 27, 1908)

Lignocelluloses. As the name indicates, this group comprises the substance of the ground tissue of the woods. The simplest type of lignocellulose, chemically as it is seructurally, is the jute fibre, the lignified bast of an annual (Corchorus tibiacca) cultivated on an enormous scale in India, and the staple of an important textile industry centred in Dundee.

The elementary composition of the purified fibre is expressed by the following percentage numbers:—

The constitution of a lignocellulose is revealed by the action of chlorine, which combines with the non-cellulose or lignone component to form a quinone-chloride of constant composition,  $C_{19}H_{19}Cl_4O_9$ , with production of hydrochloric acid, also in constant proportion to the lignone, the estimation of the acid affords an indirect measure of the proportion of lignone or noncellulose constituents (Cross and Bevan, Cellulose, Heuser and Sieber) The chlorolignone reacts with sodium sulphite (solution) to form a sulphonated derivative of characteristic magenta-red colour, freely soluble as sodium salt, consequently, the chlorinated fibre is resolved by treatment with solution of sodium sulphite (2 p c Na<sub>2</sub>SO<sub>3</sub>), the lignone-chloride dissolving from the cellulose, The quantitative which is thus isolated constants of this reaction, which is specifically dia

agnostic, ai	e as ionows		
0			P c of
			lignocellulose
Chlorine co	mbining with	lignone	8.0
••	as H	ICI .	. 80

And the lignocellulose may be statistically calculated to be composed of

The cellulose is separated actually as a hydrate of the composition  $3C_5H_{10}O_5,H_2O$ . It further differs from the normal cotton cellulose in being non-homogeneous. It is a mixture of two celluloses, the  $\alpha$ -cellulose (65 p.c.), approximating to the normal, and  $\beta$ -cellulose (15 p.c.); the latter being relatively sensitive to hydrolysis and yielding a large proportion (40–50 p.c. of its weight) of furfural on boiling with condensing acids

Constitution of lignone The characteristic groups of the lignone complex are those which combine with chlorine and which appear to be of diketo r hexene constitution. The reaction with bisulphites is also quantitative and the sulphonated derivatives are formed without resolution Certain decompositions yield products quantitatively related to original groups. thus acetic acid is a product of oxidations both acid and alkaline, and appears to result from hydrolysis of ketene groups By oxidation with CiO3 in presence of hydrolysing acids the lignone is entirely resolved to oxalic and carbonic acids, with liberation of acetic acid. From the formation of volatile pyrone derivatives in the first stages of destruction, distillation (150°-200°), and from the statistics of ultimate composition, the presence of a pyrone group in the lignone complex is inferred.

An integral expression of these quantitative relationships is given in the form of a general statistical constitutional formula for the lignone-cellulose complex

$$\begin{array}{c} \text{CO} \\ \text{CH} \\ \text{CH} \\ \text{CH} \\ \text{CH}_2 \\ \end{array} \begin{array}{c} \text{CH}_2\text{CO}_{12}\text{--HC} \\ \text{CH} \text{-CH} \text{-CH} \text{-OMe} \\ \text{CO} \\ \end{array} \\ \begin{array}{c} \text{CH} \\ \text{CH}_2 \\ \end{array} \begin{array}{c} \text{CH} \\ \text{CH} \text{-CH} \text{-CH} \text{-OH} \\ \text{CH} \\ \end{array} \begin{array}{c} \text{Cellulose} \\ \text{CO} \\ \end{array}$$

(Cross and Bevan, Researches in p 104) This has been critically examined by C Doré in relation to the study of the action of ozone on the typical lignocelluloses (J. Chem. Soc. 103 (1913), 685) of the results of which it affords consistent explanations

The wood substance of perennial woods (dicotyledons), with the same general characteristics, differs in the relation of lignone to cellulose, both in proportion and in mode of union. Whereas the jute fibre is attacked by solvent reagents similarly to the celluloses, the woods are extremely resistant. The following comparison of typical constants for beechwood is noteworthy—

The wood substance of the Conferæ, first investigated by Erdmann (Ann Suppl 5, 223), is a homogeneous complex—lignocellulose—of empirical composition,  $C_{30}H_{42}O_{21}$ , differing in minor respects from the woods of angiosperms chiefly in the lower proportion of 'furfuroid' and acetyl groups. A reaction of fundamental import and general for the entire group of

lignocellulose, is the basis of the 'bisulphite' process of preparing cellulose from woods, applied chiefly to the conferous woods, the insoluble residue of disintegration or cellulose constituting the most important staple of modern paper making. The lignone complex reacts quantitatively with the disulphites to form derivatives of the empirical composition C<sub>24</sub>H<sub>23</sub>(OCH<sub>3</sub>)<sub>2</sub>O<sub>3</sub> SO<sub>3</sub>H, the mechanism of the reaction being (1) a hydrolysis of 'acetal' groupings due to combination of H CO and bisulphite residues, (2) the  $SO_3H$  residues then migrating to unsaturated -CH = CH positions, the lignone complex being thus obtained as lignone sulphonates (Ca or Mg salts), which are gummy deliquescent solids

These reactions take place progressively in the range of temperatures 80°-150°, and industrially are carried out in digestors of special construction (silicate-lined) and enormous

capacity.

118

A third reaction characteristic of the lignone complex as such and taking place in definite quantitative proportions is its condensation with phloroglucinol in presence of HClAq There are two phases of the reaction (1) with formation of a crimson colouring matter. This takes place with a proportion of the phenol less than 1 p.c of its weight. The depth of colour is quantitatively related to the lignone, and is, therefore, an approximate measure of its proportion in a lignocellulose, as of lignocellulose (e g. 'mechanical' wood pulp) in admixture with cellulose (2) A further combination with the phenol (6-7 p c of its weight) to a product which is extremely resistant to hydrolysis (Cross, Bevan, and Briggs, Ber. 40, 3119, Cross and Bevan, J Soc Dyers, [v ] 32 (1916))

Characteristic colour reactions are those with aromatic bases; the anilines giving yellow to orange-colouring matters. Dimethyl-p-phenylenediamine give a deep-red colouration, which is the basis of the well-known quantitative method of estimating lignocelluloses in mixture

(C. Wurster)

Ferric ferricyanide in solution is reduced by the lignocelluloses, and the intermediate ferrosoferric blue cyanides in the highly hydrated colloidal form, are precipitated upon the fibre substance in intimate 'adsorption' combination. Jute fibre will take up from 30 to 50 p.c of its weight of the blue pigments thus formed, without any change of essential characteristics These phenomena have especial interest in relation to theories of dyeing and the colloidal state (Cross and Bevan, J. Soc Chem. Ind. 1893, 12, 104), and are of use in laboratory investigations of plant products (Sherman, J Amer. Chem Soc 1897, 291).

The lignocelluloses have many-sided affinities for colouring matters, due to specific reactivities of the constituent groups of the lignone complex. They are dyed directly by dyestuffs of the most varied constitutional types; and in this respect stand in marked contrast with the celluloses, which present a narrow and selective range of

direct finctorial capacity.

Other less characteristic reactions of the lignocelluloses may be briefly described.

Halogens.—Bromine reacts to form brominated lignone derivatives; but the attack is much feebler than with chlorine, and is complicated

by oxidations (Cross and Bevan, J. Soc. Dyers, vol. 32, 1916). Cellulose is isolated by repeated treatments with bromine water, followed by dilute alkalis. This is a useful quantitative method of cellulose estimation (Hugo Muller, Pflanzenfaser). Iodine is absorbed from its aqueous solutions (in KI) in definite proportions. Jute fibre digested at 18° with 20 times its weight of N/10 solution as ordinarily prepared, absorbs 12.9-13.3 p.c. iodine (Cross and Bevan, Cellulose, 138, 139).

Hydrolytic agents generally attack the lignocelluloses as a whole, that is, without resolving the lignonecellulose combination At elevated temperatures (100°-160°) dilute alkalıs (NaOH, Na<sub>2</sub>S) selectively attack the lignone groups, breaking them down to acid products-acetic acid, and the so-called lignic acids-which dissolve as sodium salts, and cellulose is obtained

as a residue.

Dilute acids at higher temperatures yield acetic acid as a product of hydrolysis (Cross and Bevan, Cellulose, 160, 161; W. E. Cross, Ber. 1910, 43, 1526), and a complex of products of condensation, including furfural

Oxy-hydrolytic agents attack the lignone complex owing to its unsaturated and oxidisable characteristics. By the action of nitric acid even in dilute form (5-10 pc. HNO<sub>3</sub>) and at low temperatures (70°-80°) the lignone is resolved into gaseous and soluble products, the B-cellulose components are also attacked, and a residue of a-cellulose is obtained, but more or less oxidised to oxycellulose

It is noteworthy that in presence of urea the action of nitric acid loses its specific character and resembles that of the simple hydrolysing acids (Cross and Bevan, Cellulose, 141, 145).

Chromic acid in presence of hydrolysing acids oxidises the lignone to oxalic and carbonic acids. Acetic acid appears as a product of hydrolysis (Cross and Bevan, J. Soc Dyers, vol. 32, 1916) The insoluble residue is an oxycellulose soluble in large proportion in dilute solution of sodium hydroxide.

Permanganates in sufficient proportion oxidise the lignone complex to products of low molecular weight, but the action may be regulated and graduated to produce valuable bleach-

ing effects

Hypochlorites are used in industrial bleaching operations (jute yarns and fabrics), but require most careful control to prevent chlorination of the fibre substance, the fixation of chlorine, even in minute proportion, leading to ultimate disintegration of the lignocellulose (Cross, Bull. Mulhouse, 1883)

Ultimate oxidations. By 'fusion' with the alkaline hydroxides at 250°-350° the lignocelluloses are oxidised to oxalic acid, the yields from the wood (sawdust) being large (80 p.c.). The process is of considerable industrial importance, being the main source of the oxalic acid of commerce

By chromic acid in presence of sulphuric acid the lignocelluloses are oxidised to gaseous products ( ${\rm CO}_2$  and some  ${\rm CO}$ ) and water. The reaction is the basis of a quantitative volumetric method of estimation.

Synthetical reactions. The ligno-Esters.celluloses react, as do the celluloses to forr... nitrates, acetatos, and henzoates; these, however, are mainly reactions with the cellulose of | Separate portion taken for each determination. the complex, the lignone groups containing no reactive OH groups: these remain in condensed combination in the resulting ester

The xanthogenic ester which results from the successive action of sodium hydroxide (17.5) p.c. NaOH solution) and carbon disulphide is a mixture or series of products, partly soluble, partly a highly hydrated and distended fibrous colloid which may be separated by filtration There is little resolution or separation of the lignone groups from the cellulose.

Destructive distillation. The highly complex changes which the lignocelluloses undergo when heated out of contact with the air, are only approximately accounted for in terms of the original substance. Typical of an enormously diversified series of volatile products are:

Methyl alcohol Acetone Acetic acid Furfural

Methyl furfural Methyl- and methoxyderivatives of Pyrogallol

which may be regarded as direct products of

resolution of the lignone complex.

Of great industrial importance are the products of the first group. The solid residue (about 30 pc) is wood charcoal, a highly condensed 'pseudo-carbon' of the approximate

elementary composition, C<sub>30</sub>H<sub>18</sub>O<sub>4</sub>.

Autoridation of liquocelluloses. Lignocelluloses in the form of ground wood or 'mechanical' wood pulp are largely used in modern printing Such papers are not only inferior in physical quality—tensile strength and elasticity -but in permanency; they change in colour (to yellow and brown) and in structure, becoming brittle. These are primary effects of oxidation; they are influenced by light, and by catalysts such as iron compounds usually present in papers as an incidental impurity, and resin acids added as sizing agents These effects and defects are generally recognised and lead to the specific exclusion of 'mechanical' pulps from papers intended for documents of permanent value (Report of Society of Aits Committee on Deterioration of Papers, J. S. A. 1898).

An important elucidation if these phenomena is suggested by the researches of the late W. J. Russell, described in the Royal Society papers on The Action of Wood on Photographic Plates in the Dark (Phil Trans (B), 1904, 197, 281, Proc. Roy. Soc. 78, 385, 80, 346) The action in question is a 'photographic' image reproducing the structural details of the wood sections, placed at some distance from ordinary sensitive plates, but with exclusion of light

On the evidence, the effects are due to an 'active emanation' from the wood substance, which appears to be a product having the characteristics of hydrogen peroxide. The general evidence also connects the phenomena with those of autoxidation, and these again with the constitution of the lignone groups of

the wood substance.

#### Assay of Fibrous Materials.

For the investigation of fibrous raw materials in the laboratory the following scheme, devised by the author (Ind. and Col. Exhibition Reports, 1887), has been generally adopted:-

Results calculated in percentage of dry substance.

1. Moisture .

Hygroscopic water, or water of condition, loss on drying at 100°.

2. Ash:

Total residue left on ignition.

3. Hydrolysis:

- (a) Loss of weight on boiling 5 minutes in 1 pc solution of caustic soda,
- (b) Loss of weight on continuing to boil one hour.

#### 4. Cellulose

Isolation and estimation as previously described (chlorination method).1

In the case of raw materials containing cutocelluloses the alkaline saponification is intensified by substituting alcoholic (2 p c NaOH) for the aqueous solution of sodium hydrate.

5. Mercerising:

Loss of weight on treating one hour with strong solution of caustic soda, 15-20 pc NaOH, in the cold.

6. Nitration.

Weight of nitrated product, obtained by treatment with mixture of equal volumes of nitric and sulphuric acids one hour in the cold.

7. Acid purification

Loss of weight after boiling with 20 p.c. acetic acid and washing with water and alcohol

8 Carbon percentage

Determined by combustion

1a Moisture. The mere operation of drying at 100° determines molecular changes in such compounds as the lignocelluloses, changes which affect the subsequent action of reagents. It is necessary, therefore, to weigh separate portions for the particular experiments in the air-dry condition, and calculate the dry contents from the data of the specimen weighed after drying at 100° The dried specimen is treated for 1b Fat, war, and resin These constituents are estimated as an aggregate extract obtained by boiling out with hydrocarbon solvents, or with ether-alcohol

The specimen weighed after extract-2 Ash ing may then be burned for the determination of ash The ash in isolated fibres is low, viz 1-2 p c; in fibre aggregates it is often high, thus in esparto and straw from 3-6 p.c., and should be taken into account in calculations of vields or loss of weight. In such cases, after weighing the product of any given reaction the specimen or a portion of it should be burnt and the ash determined. The ash-free product is calculated in terms of the original substance, also taken ash-free.

The ash constituents are: (1) those of the normal fibre; and (2) in certain cases adventitious mineral matter. These are easily distinguished and separately estimated.

3. Alkaline hydrolysis. This is the first stage

'After lengthened critical investigation and discussion (1895–1918) the method of chlorination is generally adopted as the standard process of estimating cellulose'—with variations in treatment in special cases (dense lignocelluloses—cutocelluloses) and reservation as to the definition of 'cellulose' to include furfural-yielding components For Bibliography, see W. H. Dore, J. Ind. & Eng. Chem. ii (1919), 556

towards the isolation of the cellulose. When the numbers obtained for the short period (a) and for the long digestion (b) show a marked difference, it is an obvious general indication of

low paper-making quality.

4. Cellulose Having isolated a white residue from a raw fibrous material, it may be weighed and then further investigated. An estimation of furfural will establish its position in the classification of the celluloses. For practical purposes it is sufficient to establish the degree of resistance of the cellulose to further alkaline hydrolysis, and to oxidation by alkaline cupric ovide (Barreswil's or Fehling's solution)

5 Mercerising The effect of the action of caustic soda as measured by loss of weight, and also changes in appearance and structure, may be usefully observed on the isolated celluloses, and in certain cases upon the raw fibrous The pectocelluloses are considerably materials gelatinised by the treatment, the fibres often undergoing agglomeration and drying to a harsh wiry condition The lignocelluloses are effected in the contrary direction The cutocelluloses are not attacked.

From all the above treatments the products should be treated with dilute acids by immersion in the cold, and copiously washed before drying The drying is accelerated by treatment with alcohol

6 Nitration, or conversion into nitric esters, affords useful information; but judgment must be exercised as to the applicability of the treatment to the raw material, to the partially purified products (alkaline hydrolysis), or the isolated cellulose It is a general measure of the proportion of reactive alcoholic OH groups, and also of resistance to oxidation

7 Acid purification removes basic mineral matter, often present in some quantity in raw materials. of the pectocellulose class. It may materials, of the pectocellulose class sometimes be required to attack and remove more resistant mineral constituents, such as silica or silicates In this case the material may be digested with a mixture of hydrofluoric and hydrochloric acids in vessels suitably resistant The treatment is to the action of these acids followed by copious washing

8 Carbon percentage. This is only necessary in certain investigations The value of the constant will be recognised from the fact that it varies considerably for the different groups of compound cellulose The most convenient method is that of combustion, with chromic

scid, after solution in sulphuric acid

For special purposes the above scheme is extended by operations on a larger scale, and under conditions which approximate to those industrially employed, as, for instance, in estimating the value of a raw material for papermaking purposes complementary investigation of structural characteristics, including measurements of length and diameter, of ultimate fibres (cellulose) is also required. processes and methods the special text-books must be consulted

The following is a general view of the

# CELLULOSE INDUSTRIES.

Celluloses, fibrous material, used as such: (1) Mechanical with incidental chemical treatments (bleaching, dyeing, mercerising finishing):

Textiles, fine fabrics:

Cotton, flax, hemp, ramie.

Coarse textiles:

Jute (flax and hemp).

Ropes, twines.

Hemp flax, manilla, aloe, and fibres of monocotyledons

Hat-making materials:

Straw, wood.

Paper, boards, and fibrous agglomerates: Cotton and linen rags and textile wastes (wood. straw. esparto pulps)

(2) Chemical industries. Cellulose derivatives, or celluloses, or products of resolution of compound celluloses.

Explosives:

Nitro celluloses.

Artificial fibres:

Nitro celluloses (acetates),

Aqueous solutions of cellulose derivatives.

Zinc chloride, cuprammonium. xanthogenic ester (viscose)

Films or sheets:

Nitrocellulose (acetates), xanthogenic ester.

Massive solids ·

Nitrocellulose (acetates), xanthogenic

Preparation of wood pulp, esparto, straw, and other celluloses for

Treatment of woods. esparto, straw, and textile waste materials.

(3) Products of decomposition:

Oxalic acid:

Fusion of wood wastes with alkaline hydroxides.

spirit, acetone,

Acetic acid, wood

creosote, charcoal: Products of destructive distillation of

## Supplementary Note. (Jan. 1, 1920)

The chemistry of 'cellulose' is obviously a section of the 'chemistry of colloids', the state of development of this extensive and complex subject is set forth in the Reports (I, 1918; II, 1919) of the British Association Committee Cellulose is in many respects the prototype of the organic colloids, as a primary natural product it is only known to us in its natural forms of organised structures, and it is an index of the necessary trend of progressive research in this field that the works of the histologist and chemist are convergent W. L Balls (Proc R. S (B) 1919, 90, 542) demonstrates daily growth rings in the cell wall of cotton by taking advantage of the xanthic ester reactions under limiting conditions, such as induce controlled rehydration of the cell substance without disruptive distortion

Robinson (Prov. Rep. R. S. Proc. Nov. 1919), in investigating the deformation of wood structures under stress, finds positive evidence of assocrated deliquification in the chemical sense.

Cross and Revan (J Soc. Dyers, 1918, 34,

215, 247) have investigated two cases of profound structural change in cotton under mechanical stresses, associated with fundamental changes of chemical constitution.

Of these effects the immediate factor of change may be a particular form of energy, in respect of which the effects of heat, electricity, light, and radio-active 'matter,' have been shown to be definitive (ibid.; and S J Lewis, 34, 167; W Harrison, 1919, 35, 243-250)

It is clear that for exact characterisation of cellulose, as a colloid, it must be investigated as such, and not in terms of products of proximate This postulates physical methods, resolutions of which the most obvious is the measurement of volume and volume changes (Cross and Bevan, Berl Ber 1909, 42, 2198), with temperatures and under reaction Such investigations have already suggested that the cellulose may be made up of reactive unit groups of small dimensions-of cyclic configurationand having the reactive continuity of liquid Research and progress by way of such problems and hypotheses, many of which arise in the technical routine of the cellulose industries, are the leading characteristics of the present-day movement towards a more comprehensive science of the subject

## Bibliography.

General.

'Cellulose,' Cross and Bevan, Longmans, London, 3rd ed (1918). A text-book of the subject.

'Researches on Cellulose,' I (1900), II (1905), III (1912), Cross and Bevan. general account of later investigations and

researches, to respective dates
'Chemie der Zellulose,' by C G Schwalbe,
Berlin, 1910 Full general account of contributions of original workers

haustive bibliography

A Reports on Colloid Chemistry and Industrial Applications, I (1917), and II (1918)—Cellulos, and Derivatives of Col-

loids—with comprehensive bibliography.

'Die Zellulose,' by C Piest, Stuttgart (1901), F Enke General account of subject, but with special reference to explosives

'('ellulose, ('ellulose Produkte,' 1st ed Bersch, Leipzig (1904), Hartleben A useful account of cellulose industries

Special

Pflanzenfaser,' Hugo Muller A W Hofman's Bericht, Entwicklung d Chem Ind Braunschweig (Vienna Exhibition Reports, 1873) Original investigations of cellulose and fibrous materials pioneer technological study.

'Etudes sur les Fibres Vegetales,' Vetillart, Paris (1876) Miscroscopic study of structural characteristics of cellulose and

fibres · a proneer work

'The Textile Fibres,' J. Merritt Matthews,

'Text-book of Papermaking,' Cross and Bevan (Spon), 4th ed (1916). Principles of industrial preparation and application of cellulose, &c.

'Chemistry of Papermaking,' Griffin and Little, (New York) Lockwood. 'Die

Cellulose Fabrication,' Max Schubert, Berlin, (1906) M. Krayn. Full account of wood pulp and processes of treatment.

'Die Mercerisation der Baumwolle,' Gardner, Berlin, (1878) Springer. A full account of mercerising process, and lustre finishes of textiles, with special reference to patent literature

'Kunstliche Seide,' Suvern, Berlin, Springer. A full account of artificial silk processes, with special reference to patent literature

'Die Viskose,' B. M Margosches, Leipzig (1906) An exhaustive account of scientific and technical papers, and patent literature of viscose developments. C F C.

CELTITE v. EXPLOSIVES.

CELTIUM Ct At wt ' An extremely rare element intermediate in characters between luterium and scandium, occurring in gadolinite and first isolated by Urbain Forms a white oxide, and an anhydrous chloride more volatile then lutecium chloride, but less volatile than scandium chloride The hydroxide is a weaker base than lutecium hydroxide, but is stronger than scandium hydroxide Gives a characteristic arc spectrum (Urbain, Compt. rend. 1911, 152, 141)

CEMENT. In this article the word 'cement' is used to mean a calcareous compound employed in building, and does not refer to minor ad-

hesives of the class typified by glue

This material is so Portland cement. greatly more important than any other cement of its class, that it may appropriately be dealt with first The name is trivial, referring merely to the colour of the set product which somewhat resembles that of Portland stone 'he invention of Portland cement is attributed to Aspdin of Leeds, who calcined a mixture of chalk and clay, and found that the product, when ground and mixed with water, would set; but development from this elementary stage to the status of a valuable industry was the work of many minds, which, operating mostly empirically, arrived at a procedure competent to ensure a commercially useful product, although by methods of which the rationale was only dimly understood At the present time, a better state of knowledge exists, and except in a few minor points, such as the adjustment of the setting time, which still cannot always be completely controlled, the manufacture is performed with precision and the product is of uniform and excellent quality Nevertheless, the chemistry, both of the formation of Portland cement from its raw materials and of the changes which it undergoes on setting, is incomplete, many of the reactions supposed to occur remaining obscure and hypothetical

Composition —Portland cement consists of a mixture of compounds of lime and silica and of lime and alumina These are the essential constituents, but it commonly contains in addition loose compounds of lime with ferric oxide, magnesia, and small quantities of mert insoluble silica or silicates, alkalis and sulphates Except when perfectly calcined and absolutely fresh, it also contains a little water and carbonic anhy-The ultimate dride absorbed from the air composition of Portland cement varies with the nature of the raw materials from which it

has been made. The following is a typical analysis of cement made from chalk and clay in the Thames and Medway district :--

	Per cent,
	. 220
	0.7
	6 5
	3 3
	63 0
	10
	1.5
	0 5
	10
	0 5
	100 0

All the constituents may depart a good deal from these proportions The silica may vary from 18 to  $2\overline{7}$  p c, the alumina from 5 to 10 p c, the ferric oxide from ml to 7 pc, the lime from 58 to 67 pc, the magnesia up to 3 pc, the sulphuric anhydride up to 275 pc, the alkalis to 2 p.c, and the water and carbonic anhydride according to the length of time the cement has been made and the extent to which it has been exposed

The limits laid down by the British Standard specification are as follows The proportion of lime to silica and alumina shall be not greater than the maximum nor less than the minimum ratio (calculated in chemical equivalents) re-CaO presented by  $\frac{\text{CaO}}{\text{SiO}_2 + \text{Al}_2\text{O}_3} = 2.85$  or 2.0 respectively. The percentage of insoluble residue shall not exceed 15 pc, that of magnesia shall not exceed 3 p c , and the total sulphur content calculated as sulphuric anhydride (SO<sub>3</sub>) shall not exceed 2.75 pc The total loss on ignition shall not exceed 3 p.c.

Although the ultimate composition of Portland cement is easily and accurately ascertained by the ordinary methods of chemical analysis, its proximate composition, that is, the nature and quantity of the chemical compounds or physical mixtures which constitute it, are not known with exactness and cannot be determined with precision by existing methods of inquiry The view most in favour and best supported by experimental evidence, is that the essential constituent is tricalcium silicate 3CaO SiO2, and that the body next in importance is a calcium aluminate, to which the formula 2CaO Al<sub>2</sub>O<sub>3</sub>, or alternatively

3CaO Al<sub>2</sub>O<sub>3</sub>,

is assigned. Dicalcium silicate 2CaO  ${\rm SiO_2}$  is also present. The oxide of iron occurs as a calcium ferrite, the magnesia possibly as such, the sulphuric anhydride and carbonic anhydride probably as calcium sulphate and carbonate, and the alkalis perhaps as double compounds with lime and silica or alumina As far as is known, the substances of primary importance are the tricalcium silicate and aluminate only, the minor bodies being little more than makeweights with the exception of calcium sulphate, which is of moment, because of its influence on the setting time of the cementitious constituents proper. The best proof that these smaller constituents are not essential is afforded by the fact that Portland cement of the highest quality can be prepared containing only silica, alumina, and lime in proper proportions.

It was formerly held that the calcium aluminate and silicate did not exist separately but as solid solutions, which are recognisable as distinct minerals in Portland cement clinker. The mineral termed alite is a solid solution of tri-(or di-) calcium aluminate in tricalcium silicate, whilst the solid solution of dicalcium aluminate in dicalcium silicate is called celite. As both are solid solutions and not definite compounds, their composition is not constant, but varies within the limits of solubility of the solute in each case Celite appears to be almost mert, and to take little part in the setting of the cement, but alite is decomposed by water, and both the calcium silicate and calcium aluminate are hydrated according to the following equations, in which, for simplicity, the two substances are dealt with separately

(1) 2(3CaO SlO<sub>2</sub>)+9H<sub>2</sub>O=2(CaOSlO<sub>2</sub>)5H<sub>2</sub>O+4Ca(OH<sub>2</sub>). Tricalcium silicate Hydrated monocalcium silicate.

(2)  $3CaO Al_2O_3 + 12H_2O = 3CaO Al_2O_312H_3O$ Hydrated tricalcium Tricalcium aluminate aluminate.

It will be seen that in the process of hydration of the tricalcium silicate, two-thirds of the lime are liberated and appear as calcium hydroxide, and in consequence this substance forms a large proportion of Portland cement when set. although—accepting the tricalcium silicate view—uncombined lime is absent from Portland cement as it comes from the kiln

The belief that tricalcium silicate is the essential constituent of Portland cement has been disputed on the ground that there is no conclusive proof of the existence of this body as a definite chemical individual. The alternative theory is that the essential constituent is dicalcium silicate 2CaO SiO2, associated as a solid solution with lime, which may be present in amount equal to that occurring in the tricalcium compound. On this assumption, the equations written above would still stand with the modification that this part of the lime is hydrated directly on setting, instead of being both liberated and hydrated by the action of water. Whichever view may be nearer the truth, it is certain that set Portland cement contains large quantities of calcium hydroxide, a fact which has an important practical bearing in the use of the material, as will be shown when dealing with puzzuolana.

Since the time when this idea was put forward the older view has been rehabilitated by the preparation of tricalcium silicate as a definite compound, and that it and tricalcium aluminate are the two substances to be considered as the essential constituents of cement It follows that the equations originally put forward by Le Chatelier may be taken as representing the process which cement undergoes on setting:-

 $3\text{CaOSiO}_2 + \text{Aq} = \text{CaOSiO}_2 2\frac{1}{2}\text{H}_2\text{O} + 2\text{Ca(OH)}_2$ and

 $3{\rm CaOAl_2O_3} + {\rm Ca(OH)_2} + {\rm Aq} = 4{\rm CeOAl_2O_3l2H_2O}$ 

It will be seen that the hydration of the tricalcium silicate provides ample calcium hydroxide for the production of tetracalcium aluminate, and as stated above the existence of this calcium hydroxide in set Portland cement is of importance when siliceous materials such as puzzolana are used conjointly with cement.

Insoluble silicious matter

Water, alkalis, and loss

Manufacture.—As would be expected from what has been said concerning the composition of Portland cement, the raw materials from which it can be prepared are of the most varied description Chalk, limestone, or any other approximately pure form of calcium carbonate on the one side, and clay, shale, or any typical argillaceous material on the other, can be mixed and burned together to produce Portland cement, provided that the product has a composition falling within the limits already laid down follows naturally that substances which contain both calcium carbonate and clayey matter, such as calcareous shales and argillaceous limestones, can be similarly used. On this account and because of the low price of the manufactured article and the consequent high proportion which freight bears to the total cost at the point of consumption, the manufacture of Portland cement is not confined to a particular spot, but is successfully undertaken wherever deposits of suitable raw materials occur, provided the cost of fuel is not extravagantly high But although it is perfectly practicable to manufacture Portland cement from a wide range of raw materials, yet certain of these possess qualities which give them a substantial advantage The large deposits of chalk in the Thames and Mcdway district, together with the clay in the form of river mud adjacent thereto, make this region one of the centres of the industry in this country In like manner, the existence of huge quantities of argillaceous limestone in Pennsylvania has led to the establishment of many large works in that State In the former case, the chalk and clay, being soft materials, can be readily handled and intimately mixed, and in the latter the mingling of the calcareous and argillaceous ingredients has already been accomplished, and a correct composition can be attained by the addition of a small amount of pure lime-

The following are analyses of some of the raw materials commonly employed —

### Chalk

			Da	r cent
Moisture .			10.	19 03
Sand and clay .				0 93
Combined silica (SiO,)	•			0 43
Ferric oxide + alumina (Fe <sub>2</sub> 0	) <sub>3</sub> ++	$\mathbf{H}_2\mathbf{O}_3$ )		0 48
Lime (CaO) i				42 90
Magnesia (MgO)				0 42
Carbonic anhydride (CO <sub>2</sub> )				34 16
Alkalis, organic matter, and	loss			1 65
,,		•		
			l	00 00
• Clay				}
Cuty			Do.	cent
Sand				28 42
	•			
Combined silica (SiO <sub>2</sub> )	•	•		30 32
Alumina $(Al_2O_3)$				15 49
Ferric oxide $(Fe_2O_3)$ .				7 74
Lime (CaO) .				2 04
Magnesia (MgO)				1 96
Sulphuric anhydride (SO <sub>3</sub> )	•	•	•	1 96
Combined water and loss	•	•	•	
Combined water and loss	•	•	٠	12 07
			- 14	00 00 🗆

 $<sup>^{\</sup>prime}$  Corresponding with calcium carbonate (CaCO  $_{3})$  76.60 p.c.

Limeston	e.
----------	----

Per cent

0.20

1 59

100 00

Alumina + ferric oxide $(Al_2O_3 + Fe_2O_3)$	. 0 08
Lime (CaO)	. 56 02
Magnesia (MgO)	. 0 24
Carbonic anhydride (CO <sub>2</sub> ) <sup>1</sup>	43 38
Carbonic annivuriue (CO <sub>2</sub> )	
Combined water and loss .	. 018
	100 10
Shale	
	Per cent.
Silica (SiO <sub>2</sub> )	60 22
Ferric oxide+alumina (Fe <sub>2</sub> O <sub>3</sub> +Al <sub>2</sub> O <sub>3</sub> )	. 27 20
Lime (CaO)	trace
Magnesia (MgO)	1 62
Carbonic anhydride+water (CO,+H,O	
Alkalis and loss	642
Arkans and loss	. 0 12
	100 00
Marl	
	Per cent,
Silica (SiO.)	15 10
Ferric`oxide+alumina (Fe <sub>2</sub> O <sub>3</sub> + Al <sub>2</sub> O <sub>3</sub> )	7 30
Lime (CaO)	$42 \ 16$
Magnesia (MgO)	0 34
Carbonic anhydride (CO)	33 51

Whatever may be the raw materials, the first step in the manufacture of Portland cement is to secure their comminution and intimate admixture. It must be understood that as the process of burning cement involves the sintering but not the fusion of the materials, these two conditions are of primary importance. The raw materials are commonly ground so that they leave a residue of less than 10 p c on a 180×180-mesh sieve, and intimacy of contact is secured by mixing the materials together at as carly a stage of the process of comminution as is practicable, and in addition grinding them together

The procedure is as follows In the case of soft, wet raw materials, such as chalk and clay, the two substances in proportions determined from their analysis are weighed into a wash mill, which is a large tank in which a vertical spindle having radial aims from which depend harrows is caused to revolve Enough water is added to allow the materials, when well beaten up, to form a thick cream or slurry, which is mobile The slurry enough to flow and to be pumped passes through screens to retain stones and grit, and is ground either between millstones or in more modern practice in a tube mill, which is a cylinder made of steel lined with a hard material such as quartzite blocks and partly filled with flint pebbles The cylinder is set horizontally and revolved, the slurry being fed in through a trunnion and passing out at the other end. During its passage through the cylinder, it is ground by the pebbles which are continually raised up the periphery of the cylinder and continually drop back through the liquid mass, pounding its particles on the lining of the cylinder and between the pebbles themselves. After being thus ground, the slurry flows into large reservoirs, called dosage tanks, where it is sampled and

 $<sup>^{1}</sup>$  Corresponding with calcium carbonate (CaCO  $_{3})$  98 36 p.c

analysed and, if necessary, its composition is corrected by the addition of more chalk or clay In the early days of the industry, the slurry was prepared with so large a proportion of water as to be a thin cream from which the water was allowed to settle in large tanks, the supernatant water being drawn off and the deposited solids dug out when they had arrived at the consistency of mud, and this method is still occasionally practised Usually, however, the quantity of water used is the least that will allow the slurry to flow (35 to 50 p c of the weight of the slurry, according to the nature of the raw materials), adequate mixture being thus obtained and the delay and cost of double handling The next step is to dry and burn the slurry to clinker The modern method combines these operations, the slurry being pumped into the upper end of a rotatory kiln (see below), where it is dried and burned as it travels down the kiln. Before the invention of the rotatory kiln, drying was effected either by the heat of coke ovens used for manufacturing the coke needed in the subsequent burning or by the waste heat of chamber kilns (see below), the mass thus dried cracking into rough cakes suitable for loading into the kiln Where the kiln was of a type which provided no surplus heat, the slurry was dued independently, crushed, and moistened with a little fresh slurry, moulded into bricks, redried and burnt. The choice among these various procedures depends on the kind of kiln for which the slurry is destined

When dry raw materials such as limestone and shale are used, they are crushed in a stone breaker or between rollers, according to their hardness, any moisture such as quarry water or damp from exposure to weather being removed by passing through rotating cylinders, heated by flue gases, or by an independent fire or in towers through which hot air is blown, and are ground together in the calculated proportions grinding is performed by ball mills, consisting of a drum lined with perforated steel plates and partly filled with steel balls, followed by a tube mill, such as has been described above, or in mills of the Huntingdon class, in which rollers depending from spindles attached to a vertical axle are driven outwards by centrifugal force as the spindle is rotated and bear against a steel grinding ring between which and the rollers the material is ground It will be understood that many different forms of comminuting apparatus can be and are used in cement works, according to the nature of the material. Discussion of their relative merits would be out of place in an article of this kind. The dry raw materials are fed as powder (often slightly damped to prevent dust) direct into a rotatory kiln or are moistened sufficiently to allow them to be moulded into bricks, dried and burnt in one of the kinds of fixed kıln.

Although wet raw materials are naturally ground wet, and dry raw materials are ground dry, yet occasionally dry raw materials are ground wet, it being maintained by some that any extra expenditure of heat to drive off the water added is compensated for by the more intimate admixture of the raw materials and the ease with which the slurry may be conveyed by being pumped or allowed to flow in contrast to

dry raw materials which require elevators, worm conveyors and similar mechanism.

The raw materials finely ground, intimately mixed and in correct proportion (containing in most cases about 75 pc of calcium carbonate calculated on the dry substance, the balance being the clayey constituents) are burnt to cement clinkers in various types of kiln.

The earliest was the bottle kiln, a domeshaped brick structure lined with fire bricks, having a short chimney and a grate lumps of dried slurry were loaded into this, mixed with coke, and the charge fired and allowed to burn out, the products of combustion escaping at the top of the kiln An improvement on this is the chamber kiln shown in Fig 1 a cup-shaped structure, lined with fire brick, in which the dired slurry and coke are placed in layers, communicating with a wide flue, which forms a floor on to which wet slurry can be run, the far end of the flue being connected with a The hot gases from the kiln chimney stack proper pass over the slurry and dry it, so that a charge is ready when the previous one is burnt. An ordinary chamber kiln will produce about 30 tons of clinker per run, which, including




FIG. 1 —CHAMBER KILN.

loading and discharging, usually takes a week, and it will burn 40-50 p c of coke, reckoned on the weight of clinker produced. The coke is generally bought from gas works, though formerly coke ovens were in use at the cement works themselves, the products other than coke As the coke and raw materials being wasted are mixed in the kiln, the ash of the coke appears in the finished cement, raising the proportion of silica appreciably, on this account it is customary to use raw materials containing a higher percentage of calcium carbonate in a kiln of this kind than in one in which the whole of the ash is not retained, so as to compensate for the added silicious matter, but even with this precaution the product is irregular, the ash attaching itself chiefly to the outside of the clinker and not penetrating to the centre of the lumps. It is impracticable to buin the material uniformly. Parts will be almost fused, parts merely lightly calcined and decarbonated, barely sintered and not truly clinkered, while yet other parts will have attacked the silicious liming of the kiln, forming with it a sort of slag. Hence it is necessary to pick the charge when it is drawn, rejecting the underburnt and the slagged material, and on the perfection of this picking the quality of the finished cement largely depends.

The shaft kiln is another type of fixed kiln

n which the fuel is mixed with the raw material. I'here are various forms of shaft kiln, but the runciple is the same in all cases, the raw material and fuel proceeding vertically down a shaft to a soint where combustion occurs, and passing from his down a continuation of the shaft to a trawing hole from which the clinker can be exercted. Loading and drawing proceed reguarly, the kiln working continuously instead of ntermittently, as does the chamber kiln. The working is regenerative in the sense that the products of combustion from the burning zone assign through the unburnt raw material, which is thereby heated and the air for combustion flows.




Fig. 2.—Aalborg Kiln.'

clinker in the lower part of the kiln, and thus is heated before taking part in the Combustion The Schneider and the Aalborg kiln are examples of this type, the latter is shown in Fig 2 Another wellknown shaft kıln with the modification that the upper and lower parts of the shaft are not in the same vertical line, is the Dietsch kıln (Fig 3) In this, the heated raw material is received on a stage between the upper and lower part of the shaft and passes thence the burning zone In the illustration, two such kılns are shown, built back to back There are various modifications of the shaft kiln differing in details of design. some working with forced draught

through the hot

From their regenerative
mode of working, they are
economical of

fuel, but require a good deal of moderately skilled labour and give a product mixed with the ash of the fuel and requiring picking

The Hoffmann or ring kiln (Fig 4), largely used for brick making, is also used in the manufacture of cement. The materials are moulded into rough bricks, and these are stacked in chambers arranged round a chimney which serves them all. Fuel is distributed between the blocks, and the working of the kiln is systematic, continuous, and regenerative Passage from one chamber to another and to the chimney can be controlled, and thus burning can be carried on in one chamber while the air for combestion flows in through hot clinker in chambers in which burning has been finished and the

n which the fuel is mixed with the raw material. bot gases from the burning chamber heat the libere are various forms of shaft kiln, but the raw material in chambers ahead of it on their raw material in chambers ahead of it on their way to the chimney, the chamber representing and fuel proceeding vertically down a shaft to a be combination continually shifting forward step by step all round the ring. As might be expected, this kiln is economical of fuel, trawing hole from which the clinker can be ex-




FIG. 3 -DIETSCH KILN.

the chambers, and yields a product lacking uniformity

In modern practice, these various kinds of fixed kiln have been displaced by the rotatory kiln, which, though originating in this country, was brought to a working success in the




FIG. 4.—HOFFMAN KILN.


United States, and began to be adopted here and on the Continent about 1898. This kiln is a steel cylinder lined with fire brick, set at a small angle with the horizontal, and capable of being slowly rotated round its axis. At the upper end the raw material is fed in, and at the lower is a burner sufficiently powerful to reach a clinkering temperature (about 1400° C.). As the cylinder rotates, the raw material travels

down the cylinder on account of the inclination, and on its journey is gradually raised in temperature until it reaches the lower end, where it is clinkered and drops out. The process is continuous, systematic, and to a considerable extent regenerative. The labour required is small and the output large per unit of plant. A large proportion of the fuel ash passes through the kiln without attaching itself to the raw material, and such portion as remains is incorporated with the raw materials and forms part of the finished product instead of being largely confined to the surface of the clinker, as is the case with the product from fixed kilns. When burning is carried out properly, the whole of the product is uniformly clinkered and needs no picking, but may need quenching.

The earlier rotatory kilns were some 60 feet

The earlier rotatory kilns were some 60 feet long and 6 feet external diameter. At the present time, an ordinary size is  $200 \times 8\frac{1}{2}$  feet and larger have been made and used. The output of a single kiln of this size is as much as 1000 tons a week, according to its size, as compared with 30 to 70 tons a week for a fixed kiln. A rotatory kiln is iun as nearly as possible day and night, including Sundays, throughout the year, stoppages being made only for repairs

or to patch the coating of clinker which forms on the fire-brick lining of the kiln (Fig. 5). kiln was formerly of uniform diameter, but modern designs are of larger diameter at the burning end, so as to provide a wider portion at the place where combustion is at its height. The earliest form of 10tatory kiln which approached success, was fired with producer gas, but many difficulties arose in the use of this fuel, and although the idea has merits and has been revived, it has not reached fruition. Oil burners, like a large blast lamp, having the oil sprayed in by steam or compressed air, are used with complete success wherever oil is cheap enough, but as the cost is too high in most countries, powdered coal is the standard fuel. Coal, preferably bituminous, and containing a moderate amount (10 pc) of ash when well dried and very finely powdered, can be used to form a sort of blowpipe flame much as if it were a fluid, the powder being driven into the kiln by a blast of air and issuing from a jet much as gas might do, and burning there in a supply of secondary air supplied to the space around the jet and preferably pre-heated. In actual working, these conditions are realised in the following way. Small coal is dried in a rotatory drier heated by an independent fire cr



It is powdered very finely by the kiln gases so as to leave not more than 5 pc residue on a sieve having 180×180 meshes per square inch This powdered fuel is fed from a hopper into a pipe through which air is driven by an air compressor or a fan, and issues from the mouth of this pipe, which passes through a shield or screen covering the lower end of the kiln, as a jet of dry finely powdered combustible matter carried by a blast of air, insufficient for its combustion and serving chiefly as a means for its propulsion in the form of a jet This jet is set into the mouth of the kiln and pointing towards the far end. Air for the combustion is drawn in by the draught of the chimney stack, with which the upper end of the kiln communicates, and on its way passes through a rotating cylinder down which is proceeding the red-hot clinker issuing from the kiln, absorbing a portion of that heat, and carrying it to the place of combustion. By these means a powerful jet of flame is produced, the temperature of which is ample for clinkering

Either dry or wet raw materials can be burned in a rotatory kiln. Dry materials are fed in by a screw, the mass being usually slightly damped by a small proportion of water in the trough in which the screw works, so that the fine dusty particles are not carried away by the

draught of the kiln Wet raw materials are simply allowed to flow in from a trough in which the height of the slurry is maintained constant by a device such as a pump more than capable of the maximum flow, the surplus being returned by The water, about 40 pc of the a by-pass weight of the slurry, is evaporated by the hot gases coming from the lower end of the kiln and the dried slurry forms itself into small nut-like masses as it rolls round the inside of the cylinder. These travel slowly downwards in a spiral path, and their temperature rises until the calcium carbonate is decomposed, the clay is dehydrated, and union begins between the acid and basic oxides, the silica, alumina, and ferric oxide acting in the first and the lime, magnesia and alkalis in the second sense, until finally the material becomes slightly plastic, and full combination is effected at a point near the lower end of the kiln, where the temperature is highest white hot clinker passes into a cooler, such as the rotating inclined cylinder already mentioned in connection with the heating of the air necessary for combustion, where it is often sprinkled with water for the double purpose of completing its cooling and of hydrating any lime which may not have been perfectly combined in the kıln.

With dry raw materials the course of operations is the same, save that only the small quantity of water used to prevent dust has to be evaporated.

On account of the fact that cement clinker is a basic material and that fire brick is an acid substance, using those words in the ordinary metallurgical sense, interaction may occur between these at a clinkering temperature, and a slag may be produced. Any overheating will cause the rapid formation of this slag, with the result that the clinker will be spoiled and the lining destroyed. But if the burning is properly conducted, the action is not uncontrollable, and further can be prevented by the device of heating down the slightly plastic clinker on to the firebrick lining, so that it adheres and itself forms the true lining, to which, of course, the rest of the The coating clinker is chemically indifferent breaks away from time to time, and needs to be patched by beating down fresh clinker on the fire brick thus left bare Linings of less acid substances such as bauxite, and basic linings such as magnesia bricks, have been tried, but all present some practical defect which makes them less suitable than fire brick used in the way described

Good well-burnt clinker, whether made in a fixed or rotatory kiln, is a hard rock-like mass of dark colour, varying from brown to almost The pieces of clinker from fixed kilns are chiefly rough lumps of similar size to that of the blocks of raw material, and need a preliminary crushing in a stone breaker before they can be ground. The product from a rotatory kiln is in the form of small rounded masses, varying in size from a pea to a walnut times a number of these will stick together, making a mass a foot or more in diameter, but the bulk of the product is small enough to be

ground without previous crushing

Grinding is effected in machines similar to those used for preparing the raw materials Ball and tube mills and mills of the Griffin and Huntingdon class are in most general use, millstones, such as were used for grinding coin, having been superseded on account of their small output and heavy upkeep The clinker is ground to cement having a residue of not more than 14 pe on a 180×180-mesh sieve, and is conveyed mechanically to warehouses where it is stored and bagged or filled into casks. In a modern works, the whole operation from the reception of the raw materials to the filling of the bags is conducted almost wholly mechanically, manual labour being eliminated as completely as possible, an object easily attained with rotatory kilns, which do not require to be loaded and unloaded by hand

Testing Portland cement. The quality of Portland cement and its suitability for building purposes is ascertained by various mechanical tests and by its chemical analysis The analysis shows whether the constituents fall within the recognised limits stated above, and detects the presence of such adulterants as blast furnace slag or Kentish rag (a silicious limestone), which differ widely from cement in composition or contain some characteristic constituent such as calcium sulphide, which is almost or wholly absent from Portland cement. But as it is im-

Portland cement, the ordinary analytical figures will not indicate whether the raw materials have been sufficiently intimately mixed and sufficiently thoroughly burnt to form those cementitious substances constituting sound and serviceable Portland cement. This question can, however, be satisfactorily settled by physical and mechanical tests:

Codes of testing are in common use in all countries, and the following are the principal requirements of the British Standard Specification .

The fineness to which the cement has been ground is a useful indication of quality, masmuch as the coarser parts hydrate so slowly as to be of little or no cementitious value

The residue on a sieve  $180 \times 180 = 32,400$ meshes per square inch, shall not exceed 14 p c. The residue on a sieve  $76 \times 76 = 5776$  meshes

per square inch, shall not exceed 1 p c

The specific gravity of cement is a useful criterion, and shall not be less than 3 10 below the normal value, say 3 10 to 3 15, according to the degree of aeration to which it has been subjected, it may generally be inferred that the cement has been underburnt

The mechanical strength of cement is determined by ascertaining the tensile strain of briquettes of 1 square inch section made of neat cement, and also of cement mixed with three times its weight of standard sand

The breaking strength of the neat briquettes days after gauging shall not be less than 450 lbs. per square inch, and of the sand

briquettes 200 lbs per square inch The breaking strength of the briquettes at 28 days after gauging shall show an increase on the breaking strength at 7 days, and shall be not less than the number of lbs per square inch of section arrived at from the following formulæ for neat and sand briquettes respectively -

Breaking strength at 7 days

40,000 lbs Breaking strength at 7 days Breaking strength at 7 days

10,000 lbs + Breaking strength at 7 days

The setting time of cement is useful in aiding choice of a cement for a given class of work. Three distinct graduations of time of setting are arranged for, designated as 'quick,' 'medium' and 'slow,' ranging from 10 minutes to 7 hours.

Finally, the soundness of cement, that is, its constancy of volume when set and consequent freedom from tendency to expand in the work of which it forms part, and thus to subject the work to internal stresses sometimes so great as to cause its destruction, can be determined by observing the alteration of volume and condition of test pieces made of the cement and kept under conditions designed to develop and accelerate any latent tendency of the kind. The test is carried out by means of the Le Chatelier apparatus, which consists of a small split cylinder of spring brass or other suitable metal forming a mould 30 mm. diameter and 30 mm. high. On either side of the split is attached an indicator 150 mm. long with pointed end. The mould is filled with neat cement gauged in the usual manner and placed under cold water for practicable to make a proximate analysis of 24 hours. It is then boiled in water for 6 hours;

any alteration in the distance between the points is an indication of the constancy of volume of the coment. The expansion should not exceed 10 mm, when the sample has been aerated for 24 hours, or 5 mm, after the sample has been aerated for 7 days

The specifications in use on the Continent and in America are similar to the British Standard specification as regards their aim, which is to ensure that the cement shall be mechanically strong and chemically stable. The tests themselves are also generally similar, the chief differences concerning the use of compressive as well as tensile tests and variations in the mode of determining the soundness of the cement

Other cements of the Portland class. Roughly it may be said that all cements made by burning a mixture of calcarcous and argillaceous materials are of the Portland cement class, although they may not be true Portland cement Roman cement is an example. It was manufactured before the invention of Portland cement by calcining the nodules of indurated chalk and clay called septaria, found in the Isle of Wight, and dredged upon the Kent and Essex costs. On burning, without any preparation, these nodules give a cement having a composition of which the following analysis is an example—

Silica (SiO <sub>2</sub> )			P	er cent 19 62
Insoluble residue	•			5 86
Alumma (Al <sub>2</sub> O <sub>8</sub> )				10 30
Ferric oxide (Fe <sub>2</sub> O <sub>3</sub> )				7 44
Manganese oxide (MnO)				1 57
Lime (CaO)				44.54
Magnesia (MgO)				292
Sulphuric anhydride (SO <sub>3</sub> )				2 61
Carbonic anhydride (CO <sub>2</sub> )				343
Water $(H_2O)$				0.25
Alkalıs and loss	•	•	•	1 46
			_	100.00

Roman cement has been almost wholly replaced by Portland cement. It is reddish in colour and very quick-setting, this property making it useful for work which has to be exposed to water immediately after it is laid.

Various marls contain chalk and clay in about the proportion necessary to constitute Portland cement. These are burnt without preparation and form what is known as 'natural cement'. Cement of this soit is, of course, irregular in composition and is apt to be unsound. If well slaked before use, it is a serviceable material of its class. Magnesian limestones containing argillaceous matter, are largely used in the United States for making natural cement. The following is an example of their composition:—

_				Pe	er cent.
Silica (SiO°) .					22.77
Alumina + Ferric oxide	(Al	$O_3+$	Fe <sub>2</sub> O <sub>3</sub> )		10 43
Lime (CaO)					34 54
Magnesia (MgO) .					21.85
Sulphuric anhydride (S					1.44
Carbonic anhydride (C	$(0_2)$				2 84
Alkalıs					3.63
Water and loss .					2 50
				_	

Hydraulic limes are cements of the Portland class, and are made by burning limestones containing sufficient clayey matter to yield the calcium silicates and aluminates necessary in hydraulic cement. There are two types, with many gradations between them. The first is Chaux do Teil, made by burning a limestone containing silica so finely divided and so evenly distributed that it unites almost completely with the lime without the aid of any considerable quantity of fluxing constituents, such as alumina and oxide of iron. Its composition is shown below

In a labla and las	Per cent.
Insoluble residue	0 62
Silica (SiO <sub>2</sub> ) .	22 10
Alumina $(Al_2O_3)$	.) 1 00
Ferric oxide (Fe <sub>2</sub> O <sub>3</sub> )	`} 182
Lime (CaO)	. 66 72
Magnesia (MgO)	. 117
Sulphuric anhydride (SO <sub>3</sub> )	. 049
Carbonic anhydride (CO.)	. 0 64
Water (H,O)	. 536
Alkalıs and loss	. 108
	100 00
	100 00

The second instance is afforded by the hydraulic lime prepared from blue has limestone occurring in the Rugby district, near Lyme Regis and by Aberthaw. In this material, the proportion of alumina is much higher relatively to the silica than in the case of the Chaux de Teil. An example of its composition is appended.

appended.	
	Per cent.
Insoluble residue	$2\ 39$
Combined silica (SiO <sub>2</sub> )	. 14 17
Alumina (Al,O3)	6 79
Ferric oxide (Fe,O <sub>3</sub> )	2 34
Lime (CaO) .	63 43
Magnesia (MgO) .	1 54
Sulphuric anhydride (SO <sub>3</sub> )	1 63
Carbonic anhydride (CO.)	3 64
Water (H,O)	2 69
Alkalıs and loss	1 38
	100.00

As it is not designed in the case of hydraulic limes to cause complete union of the acid and basic oxides, they invariably contain free lime which needs to be slaked. This is done systematically by the manufacturer of lime of the Chaux de Teil class, but is not generally practised in this country, the operation being performed by the user. In either case, perfect slaking is necessary because any negligence in this respect will lead to expansion in the work of which the lime forms part, and the slaking must be accomplished in such a way that, whilst the free lime is slaked, the cementitious compounds romain untouched.

There is, however, a material chalk and clav made in Germany and known as Eisen-Portland cement, which consists of Portland cement (often made from blast furnace slag) to which some 25 p c of slag has been added after the manufacture of the Portland cement itself. The slag added has a puzzuolanic function similar to that of other silicious materials which are sometimes added to Portland cement But apart from these uses of slag, various patented products, of which Passow cement and Collos cement are examples prepared by treating liquid blast furnace slag in such a way that it becomes definitely hydraulic and will set and resist the action of water as does Portland cement, are Processes for making these manufactured cements generally consist in suddenly cooling the liquid slag and in adding to it, at the moment of cooling, small quantities of some extraneous material such as salts of magnesium or the alkalis. The cause of the action of the mode of cooling and of the addition is obscure, but the fact remains that in this manner blast-furnace slag, which untreated is relatively mert, is conveited into a true hydraulic cement

Puzzuolanic cements. Except for limes which, being made from a silicious limestone, happened to possess hydraulic properties, the earliest kind of hydraulic cement was made from ordinary lime mixed with a puzzuolana, that is, a silicious material sufficiently active to combine with the lime and form a cementitious silicate Natural puzzuolanas are volcanic in origin, the deposits known as Roman and Neapolitan puzzuolana, santorin earth, and trass being examples.

Analyses are given below of Roman and Neapolitan puzzuolanas and of a similar material from Java

<u></u>	Neapo- litan puzzu- olana	Roman puzzu- olana	Java trass
Insoluble silicious matter Soluble silica ( $\mathrm{SiO_2}$ ) Alumina + Ferric o ide $\mathrm{Al_2O_3}$ + Fe $_2\mathrm{O_3}$ Lime ( $\mathrm{Ca(0)}$ ) Magnesia ( $\mathrm{Mg(0)}$ ) Sulphuric anhydride ( $\mathrm{SO_3}$ ) . Water+carbonic anhydride ( $\mathrm{H_2O+CO_2}$ ) . Moisture, alkalis, loss	p c 35 38 27 80 19 80 5 68 0 35 frace 4 27 6 72	p c 25 94 32 61 22 74 4 06 1 37 trace 8 02 4 33	p c. 50 40 19 32 16 96 nil 0 13 7 57 5 62

Granulated blast-furnace slag, broken brick and burnt clay are artificial puzzuolanas sometimes used to produce hydraulic coments when added to ordinary lime. The quantity to be added can be calculated on the assumption that the liberated lime combines in about the same proportion with the active silica in these as it does with the silica in Portland cement, it being understood that the silicious material is sufficiently finely ground to unite with the lime. In practice, puzzuolanic materials are usually coarsely ground, and only a fraction of the active silica enters into combination, the balance of the puzzuolana acting merely as an aggregate

From what has been said concerning the constitution of Portland cement and the forma-

tion of slaked lime by its hydration in setting, it will be seen that there is a considerable quantity of calcium hydroxide unprovided for by any acid oxide. It follows that this will act in the same way as ordinary slaked lime in combining with any active silicious substance, and hence that puzzuolanas can be employed as additions to Portland cement which are useful in increasing its content of cementitious matter.

Use of Portland cement. The original object of manufacturing Portland cement and similar substances was to obtain a material which would both set and would resist the action of water But it was soon found that the great mechanical strength of Portland cement made its use advantageous, even when it had not to be exposed to water, and at the present day it is the structural cement par excellence, and on account of the great decrease in cost due to modern method of manufacture, it tends to displace all other building cements, including common lime Its largest application is to form concrete by cementing together any cheap meit stony material which may be locally available and by the use of this substance, huge works of all kinds are constructed at a cost much inferior to that of the same work in masonry It has been found that Portland cement, when made into mortar or concrete, will adhere strongly to non and steel if the latter is clean and not too smooth, and on this observation the preparation of the material known as reinforced concrete is based. It is made by constructing a skeleton of mild steel rods, often tied together by steel wire and filled in with Portland cement concrete, the steel being so disposed as to take the tensile stresses in the coimposite structure It fortunately happens that the coefficients of expansion by heat of ordinary mild steel and of concrete are nearly identical, and thus internal stresses, such as might occur where two diverse materials are bonded together and exposed to fluctuations of temperature, are of very moderate magnitude and do not appreciably influence the mechanical stability of the product Although on the whole a permanent material, Portland cement may be injured or destroyed under various conditions incidental to its use Set cement is the product of the action of water, but it is not indifferent to the further action of water The products of the hydration of Portland coment and its congeners are calcium hydroxide, which is soluble in, and hydrated calcium silicates and aluminates which are decomposable by water. Therefore when water has free access to set cement and can flow through the concrete or mortar of which it forms part, the cement is dissolved and destroyed. In practice good concrete is made so nearly impermeable that, except under heavy pressure, no such flow occurs, but if the concrete is loose in texture and the flow of water correspondingly easy, destruction is only a matter of time. When the water contains salts of magnesium or calcium sulphate, which act on the lime in the one case and on the calcium aluminate in the other, forming part of the set cement, the rate of destruction is much Thus it comes about that in all increased under-water work, and more particularly in sea water, the concrete to be permanent must be impervious. There are minor and local causes

of destruction such as contact with waters containing dissolved gases, eg CO<sub>2</sub> and SO<sub>2</sub>, competent to act on the calcareous constituents of cement, but in general the failure of a concrete structure, when due to chemical action and not to mechanical weakness, can be traced to the flow of water, and usually saline water, through the failure.

CEMENTATION v. IRON.
CEMENT COPPER v. COPPER.
CEMENT STEEL v. IRON.

**CEMENTITE**, an iron carbide, Fe<sub>3</sub>C, containing 6 9 p.c. carbon, is an important constituent of cast iron and steel (Abel and Deering, Proc. Inst. Mech. Engineers, 1885, 30., Moissan, Compt. rend. 122, 421). In steels containing 1.5 p.c. or more of carbon, free cementite is found crystallised in large pseudo-hexagonal plates (Groth, Chem. Krystallographie, 1906, 1), and in practically all steels it exists with pure iron (ferite), in the cutectic mixture called pearlite (Ledebur, Stahl u. Eisen, 1898, 742).

It is very brittle and harder than hardened steel or glass (H=6) (Muller, Stahl u Eisen, 1898, 292), and its presence is the cause of the hardness of 'white' or 'chilled' cast iron (Howe, Metallographist, 1901, 177) It is recognised under the microscope by its hardness and resistance to etching.

As cohemite, it is present in meteorites (Weinschenk, Ann Museum, Wien, 1889, 4, 94)

CEPHAELINE v IPECACUANHA. CERADIA v OLEO-RESINS.

CERARGYRITE or Hornsilver. Native silver haloids, Ag(Cl, Br, I), occurring as small holohedral-cubic crystals and as compact masses with a horny appearance and texture material is soft  $(H = 2\frac{1}{2})$ , tough, and sectile, and cuts like horn. On exposure to light, it quickly The name 'hornsilber' was used by C. Gesner in 1565, and the Greek form cerargyrite was first used by F S Beudant in 1832 After the recognition of the bromide and chlorobromide in 1841 and 1842, the name cerargyrite came to be applied more especially to the But as the different members of the group so closely resemble one another that they can often only be distinguished by analysis, it is more expedient to include them as sub-species

under the name cerargyrite (see Prior and

Spencer, Mineralog Mag. 1902, xiii, 174), viz.

Chlorargyrite, AgCl

Bromargyrite, AgBr. Embolite, Ag(Cl,Br) Iodembolite, Ag(Cl,Br,I) Grey or colourless, sp gr 5.556. Greenish-grey. Greenish-grey. Greenish-yellow to orange-yellow

Isomorphous mixtures are the rule, and of those sub-species embolite  $(q\ v\ )$ , with chlorine and bromine in indefinite proportions, is the most abundant. Large quantities of these minerals have been mined as silver ores in Mexico, Chañarcillo in Chile, and Broken Hill in New South Wales. They occur in the upper oxidise I zones of veins of silver ore, and have no doubt been formed by the action of salt waters percolating from above.

Silver include forms a distinct mineral (v. Iodyratte) crystallising in the hexagonal system, but at a temperature of 146° it passes over into a cubic modification. L. J. S.

CERASIN v. Gums. CERASINE, -ORANGE, -RED, v. Azo-COLOURING MATTERS.

CEREALS. The cereals are the fruits or seeds of various species of Gramineæ In all parts of the world they constitute the most important food of mankind, and are utilised not only for bread corn and the feeding of stock, but also for the fabrication of a number of fermented drinks and industrial products like starch

The most widely distributed cereals are wheat and barley, which grow in almost all climates, oats are more confined to nothern latitudes, maize, and still more so rice, do not ripen, as a rule, north of 45°, while the millets, sorghum, and durra are truly tropical or subtropical in their habits.

In view of their importance as food, a very large number of analyses have been made of the various cereals grown under all conditions, and of the products manufactured from them, the most complete sources of information on this point are Konig, Chemie der menschlichen Nahr- und Genussm. 1903, and the various publications of the U.S. Dep. of Agiic, Div of Chemistry, particularly Bull. 13 (1899), 9, 45, and 50

While each cereal grain possesses its own characteristic composition, subject to variations determined by climate, season, variety, &c, they have still a general similarity and possess many features in common The predominant constituent is always starch, which, together with certain closely allied carbohydrates-the amylans—and a little sugar, forms from 50 to 80 p c of the whole grain Various proteins are also present in inverse ratio to the carbohydrates. they fall as low as 6 p c in rice, and rise to 20 p c. in some varieties of wheat A small quantity of fat or oil is also found, most abundantly in maize and oats, where the average percentage is about 5-6 The proportion of fibre and of ash values very greatly according to whether the grain of commerce consists of the naked fruit alone, as in wheat, rve, and maize, or whether, as in barley and oats, the flowering glumes remain adherent and can only be removed by subsequent manufacturing processes The ash of the cereal grains shows certain marked characteristics it is always very rich in phosphoric acid, up to nearly one-half of the ash when the grain is naked; potash and magnesia predominate among the bases, lime and soda being present in but small amounts, while still less chlorine or sulphure acid (derived from organic sulphur compounds) are to be found; silica is almost absent from the naked grains, but may constitute 5-20 pc of the ash of these grains possessing an adherent husk.

The structure of all cereal grains shows the following features in common: in addition to the chaffy membranes of the flower, which may or may not be adherent, the grain proper possesses several fibrous coats or skins. Within these skins at the end of the grain formerly attached to the plant, lies the embryo or germ, constituting about 3 p.c of the whole, but the greater part of the seed is occupied by the starchy endosperm, the outer membrane of the embryo which abuts against the endosperm brung known as the scutellum. The endosperm

CEREALS.

consists of a series of thin-walled cells tightly Vilmorin, Bull Soc Nat. Agr France, 1901, 61, packed with starch grains enclosed within a thin network of protein, but immediately below the skin there exists a row of differently shaped cells known as the aleurone layer, especially rich in protein. It will now be convenient to consider the cereals separately.

Wheat. The average composition of wheat from various sources is given in Table 1, ex tracted from the sources above quoted A number of other analyses may also be found in (6) date of cutting, &c

131

The composition of wheat is subject to considerable variation, and as the value of the wheat for bread-making is intimately connected with the amount of nitrogen it contains, the effect of external conditions upon this constituent has been extensively studied As factors bringing about variation, we may enumerate (1) variety, (2) climate, (3) season, (4) soil, (5) manuing,

TABLE I .- ANALYSES OF WHEAT.

	No of samples	Water	Protein	Fat	Carbo- hydrates	Crude fibre	Ash
English	22	13 37	10 99	1.86	69 21	2 90	1 67
United States .	168	10 62	12 23	1 77	71 18	2 36	1 82
" " New York .	8	10.38	12 34	1 92	71 63	2 07	1 67
" " Oregon	11	11 53	9 19	1 72	73 61	2 25	1 69
" South Dakota	8	9 02	14 89	2 04	69 89	2 49	1 70
Canadian North-West (Fife)	18	$12\ 05$	13 84	1 79	68 53	2 21	1 58
France .	70	$15\ 20$	12 64	141	68 92	2 00	1 66
Germany, S and S-W	1 1						
winter wheat	52	13 37	12 29	171	67 96	2 82	1 85
" " spring "	30	$13\ 37$	14 95	1 56	67	93	2 19
Russian, spring wheat	39	$13\ 37$	17 65	1 58	65 74		1 66
Aigentine	5	9 55	12 78	1 84	71 15	2 69	1 99
American—durum .	2	10 40	15 68	249	76 74	2 61	1 90
"—einkorn .	4	8 34	14 67	2 19	55 67	13 56	5 57
"—emmer	25	8.68	13 24	1 89	60 85	11 27	4 07

(1) High nitrogen content in wheat is usually associated with a horny translucent grain, hardness, and strength (ie capacity to make a large well-piled loaf). These qualities are most markedly seen in the wheats from Manitoba, Kansas, Hungary, and South Russia, where the high nitrogen content is mainly due to environment but partly to the varieties grown England, where the climate tends to produce a soft, starchy wheat, it is found that variety is the chief factor in determining the proportion of nitiogen In the typical English wheats, the variations in the percentage of nitrogen are small, 16 to 25 pc, according to the season, but certain wheats, like Rough Chaff, Red Lammas and Golden Drop, always show higher percentages of nitrogen than others like Rivett, Square Head's Master, and Stand Up, when grown alongside on the same soil and in the same Moreover, a few varieties have been found, notably Fife, a wheat of Galician origin but widely disseminated in Canada and the United States, which retains in England something of the high nitrogen content and strength characteristic of North-Western America, and nearly always shows a nitrogen content of well over 2 pc (see Hall, J. Board Agric 1904, 11, 321, Humphries and Biffen, J. Agric Sci 1907, 2, 1, papers read at the Winnipeg meeting of the British Association, Jour. Board Agric. 17, 1910) Moreover, Biffen (J. Agric Sci. 3, 1909, 86) has shown that high and low nitrogen content are Mendelian characters which segregate and are inherited without blending

new varieties thus raised, e.g. Cortain Burgoyne's Fife and Yeoman, combining the high introgen content and strength of Fife with

the cropping powers of English wheats, are now grown on a commercial scale

(2) Climate is more potent than any other factor in causing variations in the composition Leclerc and Leavett (US Dep of of wheat Agric Div of Chem Bull. 128, 1910) grew the same varieties of wheat in three localities with widely differing climates, and in the next year grew on seed from each locality at all three places, with the following results as regards protein content -

_		Protein	Protein in product grown in			
<b>Y</b> ear		in seed	Kansas	Calı- forma	Texas	
$ \begin{array}{r}     \hline                                $	Kansas Kansas California Texas	16 22 19 13 10 38 12 18	19 13 22 23 22 23 22 81	10 38 11 00 11 33 11 37	12·18 16 97 18 22 18 21	

Thus the seed of Kansas origin and containing 1913 pc. of protein, when grown in Califorms in 1907 yielded grain only containing 11 00 p c of protein, whereas in the same year seed from the same original stock but grown in California in 1906 and only containing 10:38 p.c. protein, when moved back to Kansas, yielded grain containing 22 23 p c. of protein In these cases, the influence of environment is enormous, and in the main the strong high nitrogenous horny wheats of the American North-West, Hungary and Russia, owe these qualities to their environment They are spring sown when the severe winter is over; they receive considerable rainfall during their early growing months, followed by great heat with progressive dryness of the atmosphere, the cycle of growth being completed in four months or less. The soft low-nitrogenous wheats are generally winter sown with an equably distributed rainfall, as in Western Europe, though, as the experience of California, Oregon, Australia, and India shows, great heat and dryness at the ripening stage will not ensure a high nitrogen content. While climates like Kansas impart a high nitrogen content to all varieties, some varieties commonly grown there, like Fife, possess a high nitrogen content congenitally and retain it in climates like the English, where other wheats with a high nitrogen content in Kansas at once deteriorate

(3) Season has a much greater effect than manuring in altering the composition of wheat, change of season being, in this respect, comparable to change of climate and environment. The variations of English seasons, however, are

never sufficient to induce 'strength' comparable to that found in the wheats of the American North-West, &c

The following table shows the variation in the composition of wheat grown on the experimental plots at Rothamsted in 1852, a wet cool year, and 1863, a dry and hot year

(4) Taken by itself, soil has not a very large effect upon the character of the wheat grown; in fact, in practice it is difficult to disengage the variations due to soil from those depending on the changed environment generally, soil being, as it were, a factor in climate

(5) The composition of wheat shows some variation with the character of the manuring, but the variations are not large. The following results were obtained with the wheat grown on the Rothamsted Experimental plots in 1893, the fortieth season in which the plots had been

		2	3	7	10	11
		Dung	Un- manured	N P <sub>2</sub> O <sub>5</sub> K <sub>2</sub> O	N only	P <sub>2</sub> O <sub>5</sub>
Weight per bushel, lb	$\cdot  \cdot \begin{cases} 1852 \\ 1863 \end{cases}$	58 20 63 10	56 60 62 70	56 00 62 60	55·90 62 60	55 60 62 50
Weight of 100 grains, grams	. \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	3 46 5 35	2 88 5 02	$\frac{308}{479}$	3 26 4 51	$\begin{array}{c} 2\ 94 \\ 4\ 76 \end{array}$
Grain to 100 straw .	$\cdot  \cdot \begin{cases} 1852 \\ 1863 \end{cases}$	49 60 67 50	53 90 70 40	41 90 59 40	47 30 74 30	47 80 70 40
Nitrogen in dry grain, p e	$\begin{cases} 1852 \\ 1863 \end{cases}$	2 02 1 52	2 08 1 65	$\frac{2}{1} \frac{29}{53}$	2·48 1·70	1·95 1·79
Nitrogen in dry straw, p c.	$\cdot  \cdot \begin{cases} 1852 \\ 1863 \end{cases}$	0·46 0·25	0 57 0 33	0·87 0 36	$\begin{array}{c c} 0.89 \\ 0.35 \end{array}$	0·46 0·44
Ash in dry grain, p e .	. \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	1 98 1 85	2 03 1 95	1 95 1 73	1 83 1 56	1 96 1 72
Ash ın dry straw, p c	$\begin{cases} 1852 \\ 1863 \end{cases}$	$\begin{array}{c c} 7 & 04 \\ 6 & 42 \end{array}$	7 04 7 12	$555 \\ 522$	5 60 5·40	6 10 5 48

growing wheat with the same manures on the same land, thus setting up much more extreme differences than would occur in practice:—

Plot	Manuring	Weight perbushel lbs	Nitrogen, p c	Ash, p.c
3 5	Unmanured	62 7	1 997	1 84
	Mineral, no N	62 3	2 015	2 04
6	$M_{inerals} + 43 lbs N$	63 1	2 187	1 98
7	" +861bs N		2·401	1 92
8	" +1291bs N		2 352	1 89
10	86 lbs N only	59 1	$\begin{array}{c} 2\ 084 \\ 2\ 264 \end{array}$	1 70
2	Farmyard manure	63· <b>4</b>		1·96

The wheats from these plots have also been ground and the flours resulting baked separately, whereupon they show extreme differences in quality and in the character of the loaf they yield (see Hall,  $l\,c$ )

(6) It will be seen from the table of composition that spring wheat possesses a higher nitrogen content than winter wheat, and it has been supposed that the difference can be correlated with the shorter period of growth of the former. Direct experiment, however, with the same variety of wheat sown at monthly intervals, did not support this view (Humphreys, l.c.). In the climates producing high nitrogen content

and strength, the wheats must be necessarily spring sown,  $\iota$  e spring sowing and high nitrogen content are both results due to the same environment and are not casually related

Similarly, the farmer's idea that early cutting gives rise to strength is not confirmed by experiment, though wheat cut green may appear somewhat horny Again, premature ripening is associated with climates producing strength, though it is not the cause of strength

(7) Various investigations have been made into the change of composition of the grain of wheat during its development (Pierre, Mem Soc Linnéene de Normandie, xv 1869, 1, 220; Déherain, Ann Agron 8, 1882, 23; 20, 1894, 561; Adorjan, J fur Landw. 1902, 50, 193; Teller, Arkansas Expt Sta. Bull. 52, 1899).

Hall and Brenchley (J Agric. Sci 1909, 3,195) obtained the results on wheat grown at Rothamsted, shown in table on opposite page.

It will be seen that the immature grain is richer in nitrogen and ash; it also contains more ready-formed sugar, and possesses a slightly higher diastatic power. Other determinations show that in the early stages about three quarters, but at ripeness practically the whole, of the nitrogen is in the protein form. The figures, however, do not support the idea that in the later stages of filling the grain, only starch

BROADBALK, PLOT 3.-1907.

Date	Green weight of 1000 grains	Dry weight of 1000 grains	Specific gravity	Per cent nitrogenin dry matter		Per cent. P <sub>2</sub> O <sub>5</sub> m ash	Per cent dextrose in dry matter	Maltose produced by 100 of dry matter
July 16	Grams 13 75 21 05 32 04 32 04 39 70 45 95 51 30 56 69 57 91 62 48 63 68 63 19 70 89 66 30 61 01	Grams 3 51 5 43 8 14 11 16 14 05 17 99 21 15 24 97 28 98 32 20 35 09 37 93 38 69 37 96	1·116 1 116 1 113 1 116 1 099 1 116 1 128 1 113 1 1196 1 215 1 218 1 231 1 204 1 271	2 679 2 406 2 458 2 167 2 119 2 055 1 856 1 828 1 801 1 720 1 856 1 787 1-846 1 778	3 70 3 03 3 14 2 80 2 66 2 39 2 38 2 16 2 16 2 09 1 89 1 96 1 94 1 93	33 66 36 91 36 88 38 73 38 86 38 68 40 35 42 54 44 17 44 30 44 06 46 50 46 50 46 30	14 99 11 08 7 36 6 71 6 23 3 70 2 42 2 17 1 86 1 46 1 99 1 91 2 02	339 4 324 7 541 4 650 7 597 1 510 6 442 9 412 0 378 1 277 9 441 6 343 7 322 1

enters, the material entering the grain is of almost uniform composition throughout. In the final ripening piocess, desiccation is the most pionounced change setting in

Carbohydrates of Wheat—The carbohydrates present in wheat grain have been examined by Stone (US Dep. of Agric, Office of Expt Stations, Bull 34, 1896), who found sucrose and reducing sugars, amylans and pentosans, in addition to starch, which is the principal constituent of the endosperm. It has also been shown by Whymper (Int. Cong. App. Chem. London, 1909) that the larger granules of the starch contained in the endosperm are more readily attacked by the diastase secreted during germination.

The sugars are chiefly present in the germ, and among them Schulze and Frankfurter identified raffinose (Bei 27, 1893, 64)

Proteins—The proteins contained in wheat, and particularly in flour, have been the subject of considerable investigation, references to Ritthausen and the other earlier workers will be found in Osborne's papers (Amer Chem J 1893, 15, 392; 1894, 16, 524; Amer J Phys 1905, 13, 35; 1906, 17, 223; Carnegie Inst Washington, 1907, Pub 84), from which the following summary has been extracted—

'The proteins of the wheat keinel are gliadin, insoluble in neutral aqueous solutions, but distinguished from all the others by its ready solubility in neutral 70 pc alcohol, glutenin, a protein having a similar elementary percentage composition to gliadin, soluble in very dilute acid and alkaline solutions, but insoluble in dilute alcohol or neutral aqueous solutions, and yielding a wholly different proportion of decomposition products when boiled with strong acids; leucosin, an albumin-like protein, freely soluble in pure water, and coagulated by heating its solution to 50°-60°; a globulin similar in composition and properties to many globulins found in other seeds, and one or more proteoses which are present in very small quantity It has also been shown that the proteins obtained from the embryo of the wheat are the globulin, albumin, and proteose above mentioned, and that these form nearly all of the protein substance of this part of the seed It thus appears that these three proteins are contained chiefly in the embryo, and that gliadin and glutenin form nearly the whole of the proteins of the endosperm, or over 80 pc of the total protein matter of the seed. It is possible that a part of the albumin, globulin, and perhaps minute quantities of the proteose, are contained also in the endosperm, for these proteins are always found in flour from which, in the milling process, the embryo is very nearly completely separated.

Gluten contains the greater part of the protein matter of the seed, together with a little starch, fat, lecithin, and phytocholesterin, and possibly some carbohydrate substance or substances of as yet unknown character non-protein substances are probably not united with one another in the gluten, but are mechanically mixed. The quantity of staich that remains in the gluten depends on the thoroughness of the washing, while the other substances owe their presence largely to their insolubility ın water The chief constituents of the gluten are the two proteids, gliadin and glutenin, the relative proportions of which vary with the variety of wheat from which the flour is made. The character of the gluten and the commercial value of the flour depend, to a large extent, on the proportion of gliadin to glutenin

In the most gluten, these proteins are present combined with about twice their weight of water, which is gradually lost on exposure to dry air or at an elevated temperature.

dry air or at an elevated temperature

'The gladin and glutenin are present as such in the seed, and are not, as was formerly supposed, derived from other protein substances through the action of an enzyme. This is shown by the fact that they may be obtained directly from the flour by the same treatment as that which yields them from the gluten, and under conditions which preclude the action of an enzyme.

'The characteristics of the different wheat constituents, the products which they yield on hydrolysis, and related questions are also considered, and the nutritive value of wheat gluten and other proteins is discussed on the basis of the kind and amount of the cleavage products which they yield on hydrolysis.

134 CEREALS.

'The amount of glutaminic acid which the gluten proteins yield is far greater than that yielded by any of the other food proteins, with the exception of gliadin from rye and hordein from barley The proteins of the legumes and nuts which are used as food yield from 15 to 20 p c of glutaminic acid, so that the mean amount of this amino- acid from the wheat protein is nearly twice as large. The same also is true of ammonia

'The proportion of arginine from wheat gluten is relatively small compared with that from most other seed proteins, many of which

yield from 10 to 16 p c of this base 'The proportion of lysine is likewise small, especially compared with that obtained from the leguminous seeds. The amount of histidine, however, does not differ very greatly from that of the other seed proteins What significance these differences have in respect to the nutritive value of these different proteins must be determined by future investigation, for it has only very recently been discovered that such differences exist

Fat —Wheat contains about 15 p.c of fat, the highest proportions being found in the wheats from the ('anadian North-West It is mostly present in the germ, in which it may

amount to 15 pc. This oil easily turns rancid, and is characterised by a high iodine number

Ash —Analyses of the ash of wheat may be found in Wolff, Aschenanalysen, Berlin, 1871 and 1880, Lawes and Gilbert, Chem. Soc. Trans. 1884, 45, 305

The table given below, from unpublished Rothamsted records, shows the variations induced by manuring in 1896, the forty-fourth season in which the wheat had been grown on the same plots with the same manures.

Mill products of Wheat -An account of the various products obtained during the conversion of wheat into flour, with analyses, is contained in Part 9, Bull 13 (1898) US Dep of Agric Div of Chem See also Maurizio, Geticide Mehl

u Brot Berlin, 1903. For milling offals, see Cranfield, J. Agric Sci. 1914, vi. 102 Baking qualities of Wheat Flour—The earliest attempt to determine the quality of flour in a baker's sense, i e the power of the flour to produce a large firm loaf, consisted in estimating the weight of gluten that could be kneaded out of a given weight of flour. While this test proved to be of value in comparing flours from similar wheats or from wheats grown in the same locality, so many exceptions occurred that

Broadbalk Wheat Grain—Season 1896

Plot	2b	3 and 4	6	8	10ab	11	12	13	14
Ash in dry matter	1 90	1 84	1 88	$174 \\ 2101$	1 57	1 72	1 78	1 76	1 76
Nitrogen in ,, .	2 115	1 835	1 849		1 94	1 931	1 949	1 875	1 954
Ferne oxide Lime Magnesia Potash Soda	$ \begin{vmatrix} 0 & 67 \\ 2 & 40 \\ 10 & 87 \\ 31 & 75 \\ 0 & 036 \end{vmatrix} $	$egin{array}{c} 0.84 \\ 3.30 \\ 10.28 \\ 33.62 \\ 0.101 \\ \hline \end{array}$	0 81 2 70 10 31 33 08 0 007	0 76 3 00 10 50 33 23 0 084	0 98 4 91 10 10 34 79 0 196	0 735 4 63 9 86 32 33 0 105	0 755 4 00 9 96 32 62 0 115	0 744 3 21 9 93 33 97 0 078	$\begin{array}{c c} 0 \ 695 \\ 4 \ 23 \\ 10 \ 36 \\ 32 \ 21 \\ 0 \ 114 \end{array}$
Phosphone acid	51 70	48 34	49 61	48 97	43 07	48 09	48 30	48 79	47 83
Sulphuric acid	0 89	1 52	1 26	1 28	3 38	1 49	1 39	1 22	1 49
Chlorine	0 020	0 305	0 054	0 205	0 973	0 199	0 138	0 088	0 279
Silica	0 46	0 55	0 53	0 43	0 64	0 46	0 48	0 45	0 45
Sand and charcoal	1 29	1 38	1 86	1 61	1 40	2 30	2 18	1 45	2 39

no absolute value could be attached to the test

Estimates based on the total nitiogen in the flour proved to be subject to the same errors Measurements were then made of the gliadin, the wheat protein soluble in 70 pc alcohol; Fleurent further suggested that the ratio of the gliadin to the glutenin in the gluten determined its quality, but these methods in turn broke down under cuticism (see Hall, lc) T. B. Wood (Jour. Agric. Sci. 1908, 2, 139, 267) regards baking quality as dependent on two factors . the size of the loaf is determined by the amount of sugar contained in the flour, together with that produced in the dough by diastatic action. This determines the evolution of carbon dioxide, under the action of the yeast, both as to rate and total amount Then the quality of the gluten, which determines how the

dough will stretch into a sponge and retain the carbon dioxide, depends on its physical character, which will vary enormously according to the state of equilibrium between the colloid gluten and the salts in the water with which it Under certain conditions the is in contact gluten loses all consistency, its physical character being entirely conditioned by the surrounding liquid.

In practice millers now add malt extract to certain flours in the dough-making process, in order to increase the evolution of carbon dioxide. Soluble phosphates are also introduced whereby the elasticity and gas-retaining power of the gluten is increased. The quantities necessary depend upon the origin of the wheat, and must be determined empirically for each flour mixture (see Wood, The Story of a Loaf of Bread: Cambridge University Press, 1913).

135

CEREALS. Nutritive Value of Wheat.—The nutritive calculated by Chamberlam (U.S. Dep. of Agric value of various samples of wheat has been Div. of Chem. Bull 120, 1909) as follows:—

		1	Digestible o	onstituents			Production	Albu-
Variety	Water	Protein	Fat	Carbo- hydrates	Crude fibro	Ash	calories per 100 lbs.	minoid ratio
North-West spring Soft winter Durum	10 11 10·55 9 77	11 78 9 96 13 02	1 41 1 20 1 72	73 04 75 24 70 77	1 17 1 17 1 33	1.85 2 00 2 02	94,717 94,760 94,457	6 6 7 9 5 8

from the total constituents of the wheat, and in the last column but one the value of the wheat for the production of energy or increased weight is expressed as calories developable from 100 lbs. of the grain, after allowance has been made for the non-digestible portions and the work consumed in the process of digestion. The last column gives the ratio of the non-nitiogenous constituents [digestible carbohydrates and fibre+digestible fat  $\times 2.25$ ] to the digestible proteins.

The question of the digestibility and food value of the different fractions of the wheat grain produced in commercial milling—the comparison between 'white' and 'brown,' 'standard' or 'whole meal' bread, has received greatly increased attention during the war, which forced war bread 'upon nearly all civilised nations Before the war wheat was milled to yield less than 70 pc of white flour and 30 pc or over of offals, during the war the extraction was raised to 80 p.c and eventually to 90 p c The general result of the experience goes to show that the digestibility of the included fractions is higher than had been expected from previous experiments, and that only a comparatively small fraction of the population suffers from digestive

The analyses give the digestible as distinct | troubles in consequence of cating bread from flour containing a marked proportion of husk While the old conclusion that white bread is the most concentrated and digestible of all breads is confirmed, the utilisation of the wheat grain for human food increases up to the 90 pc. extraction, both as regards available energy and protein. See Report of the Food (Wai) Committee of the Royal Society, 1918 The older literature is summarised in Hamill, Local Government Board Reports on Public Health and Medical Subjects, No 55, 1911

Wheat Straw -The composition of wheat straw is more variable than that of the grain: the more complete the ripening process the more thoroughly will the carbohydrates, proteins, and valuable ash constituents of the straw be transferred to the grain. The variations in the composition of the straw in response to different conditions of soil and manufing, are minimised in the grain, which the plant endeavours to make of as constant a composition as possible The following table (Konig) gives a few typical analyses of straw, which are followed by a table of the mean composition of the ash of straw from wheat grown on the Rothamsted plots, to illustrate the variation with manufing

Origin	No of samples	Water	Fat	Protein	N free extract	Crude fibre	Ash	
English winter sown German ,, American spring ,,	$\frac{-}{7}$	13 6 14 3 9 6	1 3 1 2 1 3	3 3 3 0 3 4	3 94 <sup>1</sup> 35 9 43 5	37 1 40 8 38 0	5 3 <sup>2</sup> 4 85 4 2	

Broadbalk Wheat Straw-Mixed Samples, 10 Years, 1882-91.

Plot	2	. 3	5	7	10	11	12	13	14
Ash in dry matter	. 8 13	7 69	7 95	5 89	6 02	5 84	5 69	5 93	5 52
Femc oxide .	0 31	0 94	0 60	0 50	0 54	0.43	0 33	0 34	0 41
Lame.	3 64	$4\ 38$	3 49	5.68	8 36	9 14	7 71	5 38	7 70
Magnesia	1 51	1 51	141	1 76	2 29	2 25	1 92	1 53	2 46
Potash	18 41	13 51	16 30	25 84	13 67	9 92	14 65	23 23	14 88
Soda	0 09	0 10	0 09	0 21	0 41	0 58	0 57	0 03	0 33
Phosphoric acid .	.   3 87	2 97	4 24	3 81	2 26	4 26	3 64	3 38	3 87
Sulphune acid .	3 44	3 81	4 76	5 40	6 72	5 44	5 32	5 06	5 31
Chlorine .	. 2 92	1 81	1 90	6 59	2 72	1 66	2 88	5 60	2 81
Carbonic acid	2	2	2	2	1 25	trace	none	none	trace
Silica	. 64.64	67 70	65.14	49 59_	60 44	65 20	61 79	54.16	61 08
Sand and charcoal	1 41	3 78	2 27	1 93	2 23	1 52	1 62	2 36	181

<sup>1</sup> Carbonydates.

<sup>2</sup> Not determined.

136 CEREALS.

Methods of Analysis. - In the analysis of wheat. the ordinary methods employed for feeding stuffs are followed, the only special determinations required being those of gluten and gliadin

For the determination of gluten, 20 grams of dry flour, or of the ground wheat from which the husk has been sifted, are weighed out on to a piece of bolting silk, about 8 inches square, moistened with water and kneaded up into a paste silk is then gathered together over the dough and kneaded in several changes of water or under a tap until no more starchy liquid flows away, being finally kneaded in the air until it ceases to lose water. The gluten will then generally form an elastic coherent mass which can be removed from the silk without loss, weighed in its wet condition, spiead on a tared sheet of glass or porcelain basin, dried at 100° to a constant The ratio between weight, and weighed again wet and dry gluten is regarded as of some importance

For gliadin, 5 grams of flour are put in a flask with 250 c c. of 70 p.c. alcohol, and shaken in a machine for 6 hours. The mixture is then rapidly filtered, and 100 c c of the solution are placed in a Kjeldahl flask The alcohol is distilled off and the nitrogen in the residue is determined in the usual way Nitrogen  $\times 5$  68= gliadin.

Barley generally shows a smaller Barley. protein content than wheat, this difference is not only characteristic of the two species, but barleys are likely to have been unconsciously selected for low protein content, this being the quality most desired by the brewer, who has always been the chief purchaser of barley The adherent glume and pale further cause an increase in the crude fibre and ash, as compared with wheat

The following analyses of barley are derived, 1-6 from the author, 7 from Konig (lc), and 8 and 9 from Wiley (lc):—

portunee		1					
	Number of analyses	Water	Fat	Proteins	Carbo- hydrates	Crude fibre	Ash
English Chevaliei—  1899—grown after roots 1899—, , , wheat . 1900—prize winners . 1900—poor quality . Chili Chevalier, 1900—prize Asia Minor, four-rowed—prize Germany, Middle and North United States Canada .	21 7 7 4 1 2 120 32 20	16 23 16 44 16 30 16 80 15 6 15 3 14 05 10 8 11 96	1 65 1 31 1 50 2 28 2 06 1 86 1 80 2 13 2 06	9 69 9 12 7 81 10 78 9 44 8 75 9 88 10 69 10 57	66 06 <sup>1</sup> 66 05 <sup>2</sup> 68 36 <sup>3</sup> 64 24 <sup>4</sup> 67 04 <sup>5</sup> 67 47 <sup>6</sup> 66 75 69 89 68 90	4 10 4 64 3 80 3 82 3 61 4 38 4 77 4 05 4 10	2 27 2 44 2 23 2 08 2 25 2 24 2 ·75 2 44 2 41

The composition of barley is affected by climate, environment, manuring, &c, in the same way as the composition of wheat, barley being somewhat more susceptible to change The commercial value of barley varies also to a much greater degree, not only are there two distinct grades-barley for malting and barley for feeding purposes, the price of the former being nearly 50 p.c. higher than that of the latter, though the two kinds grade off into one another-but special prices are paid for bailey of exceptional quality

(I) Variety The number of barley varieties is small compared with that of wheat. The tworowed varieties fall into two types, wide and narrow eared, and are most prized for brewing purposes The four-rowed and six-rowed types are not so much used for brewing comparative analytical data are lacking

(2) The following data, drawn from the Rothamsted records, show the effect of season and manuring upon the character of the barleyin 1893, a summer of extreme heat and dryness, and 1894, a wet and cool season .-

Plot	Manuring	Yield of grain		Weight p	er bushel	Grain to	100 straw	Nitrogen per cent in grain		
1100	Manuing	1893	1894	1893	1894	1893	1894	1893	1894	
10 40 1A 2A 3A 4A	Unmanured Minerals, no N N only N, $P_2O_5$ , no $K_2O$ N, $K_2O$ , no $P_2O_5$ N, $K_2O$ , $P_2O_5$	Bush 8 3 9 9 11 6 18 1 16 8 30 8	Bush 10 0 13 1 10 4 34·9 17 8 41 4	1b. 55 6 56 1 55 1 54 0 55 8 56 3	1b 51 1 52 1 50 4 51 9 51 5 54 1	71 9 74 3 85 3 101 0 85 9 102 2	70 3 79 2 67 5 77 0 73 8 77 7	1 899 2 011 2 188 2 129 2 171 2 081	1 409 1 467 1 646 1 600 1 614 1 440	

The results, as shown in the table at top of next page, also obtained at Rothamsted, show the effect of nitrogenous manuring

These results confirm the general opinion that barley for malting purposes should not be grown on land that is in high condition, either as a result of its previous treatment or through

the recent application of nitiogenous ferti-

- 1 Containing starch, 57 9 p c.
- Starch = 54 2 p.c
   Starch = 61 5 p c
   Starch = 58 0 p c.
- Starch = 612 pc
- 6 Starch = 62'4 p.c.

	Dressed grain. Weight per bushel lbs	Grain to 100 straw	Offal grain to 100 dressed grain	N in dressed grain p.c
No nitrogen 86lbs per acre organic 86 lbs per acre organic N in the previous year	58.0 57.3 55 1	110 4 96 6 72 8	5 9 12 5 34 9	1 61 1 79 2 42

Composition of Barley -(a) Carbohydrates Barley contains soluble sugars, which may amount to 4 p c of the dry grain; sucrose is the chief constituent, but dextrose is also found, and O'Sullivan has shown the presence of raffinose The main carbohydrate present is starch, of which H T Brown and his colleagues found 55.5 pc in the dry material of 10 baileys examined (Trans Guinness Research Lab 1903, I, 91) The same investigators (l c. 1906, 2, 312) proceeded to the determination of certain water soluble polysaccharides, called by O'Sullivan amylans, which yield on acid hydrolysis a racemic acid, glucose, arabinose and xylose

They further showed that the sum of the sugars, starch, and amylans, with a little ash and soluble protein, made up the whole of the barley grain that goes into solution when the barley is extracted with boiling water and hydrolysed with malt extract. They obtained from 95 to 10 p c of amylans on the dry weight of various samples of barley examined.

137

In the 'grains' not rendered soluble by diastase, there are considerable quantities of furfuroids' to be found, besides cellulose and

(b) Osborne and Ritthausen have examined the proteins of barley, and Osborne (J Amei Chem Soc 1895, 17, 539) obtained 4.5 p.c. of an insoluble protein, 4 p c of hordern soluble in dilute alcohol, 03 p c of leucosin, and 195 p c of edestin and proteose soluble in water has also been shown that a small proportion of the nitrogen of barley is not combined as protein

(c) The ether extract of barley is found to contain neutral fats, free fatty acids, and lecithin.

(d) The following analyses show the composition of the ash of barley from the Rothamsted plots in 1871 —

BARLEY-SEASON 1871.

			-									
	Un	1 O 4 O Complete manured minecals		Niti	A Togen nly monia)	A A Nitrogen and nunerals		1 C Nitrogen only (as tapecake)		7 Farmyard dung		
	Grain	Straw	Grain	Straw	Grain	Straw	Gram	Straw	Grain	Straw	Grain	Straw
Ash in dry matter	2 64	6 17	2 77	6 96	2 38	5 51	2 64	6 70	2 84	5 54	2 87	7 55
Ferric oxide Lime .	0 49 2 91	0 78 10 04	$\begin{array}{c} 0 \ 41 \\ 2 \ 67 \end{array}$	0 46 7 21	0 35 3 29		0 31 2 51	0 44 7 56	$0.35 \\ 2.31$	$0.42 \\ 9.52$	0 31 2 22	0 31 6 56
Magnesia Potash	7 70	1 96	7 65	1 44 27 36	7 95	1 97	8 07	1.89	7 91 28 89	2 18	7 75	1 30
Soda .	1 30	5 85	0 50	- 00	2 16	13 24	0 68	2 16	0 56		0 40	
Phosphoric acid Sulphuric acid .	$\begin{vmatrix} 34 & 05 \\ 2 & 24 \end{vmatrix}$	5 53	$\begin{array}{c} 35 \ 95 \\ 2 \ 15 \end{array}$	7 26		4 51	35 64 1 56	3 17 5 54	$\frac{36}{1} \frac{39}{60}$		$\frac{36}{1} \frac{72}{65}$	
Chlorine Silica	1 71	6 96	0 71	7 72		<b>1</b> 5 10	0 86		0 68	4 55		11 99
Sand and charcoal	19 62 4 1 67	3 52	18 99	42 63 2 18	18 57 2 84	37 98 1 64	20 38 0 93	$\begin{vmatrix} 32 & 71 \\ 2 & 13 \end{vmatrix}$	19 85 1 80	42 04 1 87		42 71 1 68

pared from barley is 'pearl barley,' in which the husk and germ have been removed mechanically This material is also sometimes ground into barley flour. Important by-products for cattle

Barley Products -The chief material pre- | feeding are also the 'grains' obtained in the brewing process after the diastatic extraction of the starch contained in malt, and the malt culms or sprouts knocked off the partly germinated barley after malting Analyses are given below'-

		Water	Fat	Proteins	Carbo- hydrates C	ruae fibre	Ash
1 .	et)	14 9 11 8 75 0 10 0	1 0 2·1 2 0 7 0	7 6 28·3 5 5 20 0	75 1 42 8 12 0 42 0	0 6 12 4 4 5 16 0	0 8 7 6 1 0 5 0

Maize. Maize or Indian corn is a cereal grain characterised by a high proportion of carbohydrates and oil but comparatively low protein and ash. Its composition does not vary to the same degree as that of wheat or barley. The average analyses (shown on top of page 138)

are derived from Wiley (US Dep of Agric. Div of Chem Bull 50, 1898) and Konig (l c.).

The only variation in composition worthy of note is that in 'sweet corn,' varieties of which are grown in America to yield immature cobs for boiling as a vegetable, the grains contain a

	Source				Moisture	Crude fat Proteins		Carbo- hydrates	Crude fibre	Ash	
United	States,	Mean .			10 75	4 25	10 00	71 75	1 75	1 50	
,,	,,	Northern	٠.		9 98	5 11	10 64	71 32	1 41	1 54	
,,	,,	Middle W	Vest		$12\ 33$	4 97	10 89	68 16	2 22	1 43	
,,	,,	Pacific sl	ope		9 78	6 40	8 14	72 13	2 07	1.48	
Italy			-		13 13	3 84	10 26	67 72	2 88	1 95	
German	y (1880	))		14	13 00	5 11	8 62	70 54	1 38	1 34	
Hungar	ý (1880	)		38	13 00	3 85	9 06	71 10	1 69	1 30	

higher proportion of sugar than is normal, especially in the immature state, and also a greater amount of oil.

Marze oil is expressed in considerable quantities from the germ, which is removed in preparing meal from the grain. It is a light amber-coloured tasteless oil used for salad oil, as a lubricant, and for soap making

Carbohydrates —Less than 1 pc of sugar is present in maize, the greater part being suciose with a small amount of reducing sugar. Starch is the principal carbohydrate, but a little dextrin or other gum-like body has been found, also furfuroids and pentosans up to about 5 pc

Proteins—Osboine has isolated from maize, myosin, vitellin and another unnamed globulin, small quantities of two albumins, and zein, the characteristic protein which exists in both a soluble and insoluble form. It is noteworthy that zein, on hydrolysis, yields no lysin and no tryptophan, and certain defects in the nutrition of animals fed on maize alone have been set down to the lack of these groups in the zein molecule

Ash—The following table from Wolff's Aschenanalysen gives the average composition of the ash of maize —

	c. ash n dry K <sub>2</sub> O	Na <sub>2</sub> ()	CaO	MgO	Fe <sub>2</sub> O <sub>3</sub>	$P_{2}O_{\mathbf{\delta}}$	$so_3$	SiO <sub>2</sub>
Red German American	1·30 24 33 1 28 26 75	1·50 3 85	3 16 2 56	16 00 15 24	1 88 2 00	49 36 47 47	1 00 1 20	2 77 1 93
German .	1 72   25 73		2 03	17 35	1 51	43 80		4 99

Maze Products—Maize is ground into a meal, 'corn flour,' the germ and the more flinty portions of the grain being removed, and the composition of the meal is set out in the table below, in which is also given the composition of maize 'stover,' i e the stalk, leaves, &c, without the cob

A large number of commercial products are derived from maize, especially starch and glucose; and the manufacturing processes give use to various by-products—maize germ meal, gluten feed, &c, used for feeding cattle, but their composition depends entirely upon the method by which they are prepared.

	Moisture	Oil	Proteins	Carbo- hydrates	Crude fibre	Ash
Corn flour · three samples . Marze stover .	12 57 19 81	$\frac{1}{2} \frac{33}{56}$	7 13 4 19	78 36 42 87	0 87 26 02	0 61 4 55

Oats. From the food point of view, oats form one of the most valuable of the cereals, being exceptionally rich in oil and protein; it is only the adherent husk which prevents them

from being the most concentrated of all the food grains Oats show considerable variations of composition, as the following analyses indicate —

No. of Analyses	Water	Fat	Protein	Carbo- hydrates	Crude fibre	Ash	party of the same
103 196 22 72 10	12 11 12·11 12 11 9·96 16 90	5 30 5 46 4 96 4 42 5 59	10 82 9 52 11 · 26 12 07 11 57	58·23 60·47 59 35 58 28 57 54	10·25 9·18 9 33 11 92 8 25	3 29 3 26 2 99 3 35 2 59	König " Wiley Aitken 1
16	7 30	7 20	16 10	67 50	0 90	1 90	Atwater 2
	103 196 22 72 10	103 12 11 196 12 11 22 12 11 72 9 96 10 16 90	Interest of the control of t	103	Table   Part   Protein   hydrates	Analyses   Water   Fat   Protein   hydrates   fibre	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$

As might be expected, the composition of oats varies with variety, climate, season, and manufing, in much the same way as wheat does. Of these variations, those due to the variety of

oat are of most importance, particularly in the

Trans. H & Ag Soc. 1901, v. 13, 292 and 293.
 U S. Dep. of Agric. Office of Expt. Sta. Bull. 28, 1899.

CERITE. 139

manufacture of oatmeal. The proportion of husk to kernel is of most significance, and the individual Scotch analyses, of which the mean is given in the table above, show variations in the fibre of from 6 to 14 p.c., and in oil, the other variable constituent, of from 4 to over 8 p.c.

Wilson (Trans. H. and Ag. Soc. 1903, v. 15, 183) gives figures relating to the amount and composition of the oatineal obtained from several varieties of oats grown on the same experimental area in Aberdeen, as follows:—

Variet	Variety		Year	Husk, p c	Oatmeal, p c	Oil in oatmeal, p c	Protein in oat- meal, p c.	
Dun			1901	21 98	66 21	7 68	14 37	
,, .		.	1902	21 36	64 13	7 74	15 50	
Potato .		.	1901	$22\ 24$	66 32	9 59	15 44	
,, .		1	1902	$22\ 47$	64 02	10 01	12 31	
Newmarket			1901	21 79	66 65	4 60	14 37	
,,		.	1902	20 46	64 84	6 63	13 50	

See also Hendrick, Trans H. and Agnic Scc 1910, xxii 16

Other Cereal Grains and Products therefrom. most important -

Of the very large number of other cereal grains consumed as food, the following are among the most important —

	No of samples	Water	Crude fat		Carbo hydrates	Crude fibre	Ash	Authority
Rye German American (World's Fair) .	63 18	13 37 10 62	1 84 1 65	11 52 12 43	68 88 71 37	2 55 2 09	1 94 1 92	Konig Wiley
unhulled hulled . polished		10 50 12 00 12 40	1 60 2 00 0 40	7 50 8 00 7 50	67·40 76 00 78 80	9 00 1 00 0 40	4 00 1 00 0 50	Wiley 1
Millet (Panicum miliaceum) . unhulled hulled	6 34 9	12 50 9 00 11 79	3 89 3 25 4 26	10 61 12 83 10 51	61 11 71 50 68 16	8 07 8 84 2 48	3 82 3 58 2 80	Konig Chamberlain Konig
Sorghum saicharatum ,, vulgare Andropogon sorghum	38 12 10	15 17 11 46 —	3 36 3 79 3 25	9 26 8 96 11 7J	67 99 70 25 81 58	2 51 3 59 1 80	171	"," ('hamberlain

Chamberlam (l  $\epsilon$ ) has calculated the feeding value of the cereal grains of American origin as follows —

	Produc- tion value, cala per gram	Starch cquiva- lent	Albumi- noid ratio
Wheat .	2 08	88 1	6.5
Barley	1 99	84.2	80
Oats	1 70	72.0	58
Maize	2 23	94.2	12 3
Emmer .	1.64	69.4	60
Einkoin	1 60	67.8	5.2
Rye	2 08	88 2	6.9
Panicum miliaceum	1.84	78 1	115
$And ropogon\ sorghum$	2 06	87.2	10 5

The production value represents the number of calories of energy available from the gram of food, after deduction has been made for the energy in the excreta and the energy consumed in digestion processes. The starch equivalent gives the effect of 100 lbs. of these foods in the production of fat or external work in terms of lbs of pure starch required to produce the same effect.

A. D. H

CEREBRIN v ENZYMES
CEREBROSE (Galactose) v CARBOHYDRATES
CEREDINE. Trade name for a preparation
of dry powdered yeast.

CERESIN. Ceresin occurs in commerce in the form of thin round masses several inches in diameter, harder than wax, of a dazzling white appearance, inodorous, and transparent at the edges. The fracture is somewhat similar to that of wax, but ceresin does not become plastic in warm water The melting-point is between 75° and 80°. It is not attacked by acids, either in the cold or when heated, nor by alkalis, which do not saponify a trace of it. It is entirely volatilised at a high temperature without alteration Ceresin is apparently a paraffin, it does not appear to be obtained from coal tar, but from Galician 'earth-wax,' i c ozokente. It is used Galician 'carth-wax,' i c ozokerite as a substitute for beeswax (J. Soc. Chem. Ind. 11, 372

CERFLUORITE v. YTTROCERITE. CERIN and CERYL CEROTATE v CORK; WAXES

CERISE. A trade name for a variety of amiline red and acid magenta

CERITE. A hydrous silicate of cerium metals. Rammelsberg's analysis (1859) showed the presence of 64 55 p.c. Ce<sub>2</sub>O<sub>3</sub> with 7·28 p.c. (La,Di)<sub>2</sub>O<sub>3</sub> and approximated to the formula 2Ce<sub>2</sub>O<sub>3</sub> 3SiO<sub>2</sub>:3H<sub>2</sub>O. Later analyses showed a predominance of lanthanum and didymium, suggesting the name lanthanocerite. Crystals are orthorhombic, but of rare occurrence. The mmeral is usually found as reddish granular masses much resembling in appearance certain

1 Type analyses from various data.

140 CERITE.

varieties of corundum, from which, however, it is readily distinguished by its lower hardness (H 5½; sp.gr. 49) It occurs as a bed in gneiss at Bastnas in Sweden, and was the material in which the element cerium was first detected L. J. S.

CERIUM. Symbol Ce. At. wt 140.25 (H Robinson, Proc. Roy. Soc. 1884, 37, 150; Brauner and Baték, Zeitsch. anorg. Chem 1903, 34, 103, 207) The most abundant of the rareearth elements. The metal has been prepared by (1) the electrolysis of the fused chloride with or without an admixture of barium or calcium chloride or alkali chloride, (2) the electrolysis of a mixture of cerium dioxide and cerium fluoride (Borders and Stockem, D R. P. 172529; Muthmann and Weiss, Annalen, 1902, 320, 231; 1904, 331, 1, 1907, 355, 116). By the latter process, the yield corresponds with 57 p.c. of a current of 750 amperes and 7.5 volts with current densities of 9-10 amperes and 3 amperes per sq cm. at cathode and anode respectively.

Metallic cerium has the colour and lustre of non; its hardness is comparable with that of silver or tin, mp 635° (Hirsch), 623° (Muthmann and Weiss); sp gr. of fused product of electrolysis 692 at 25°, sp.heat 005112 (20°-100°), hardness (as determined by the scleroscope), 25 9 and 9 5 for rolled and freshly cut surfaces respectively; ultimate strength (as determined in an Olsen machine, with test bars of 0 212 in diameter), 12,900 lb. per square inch; and no elongation with a constant load at 350 lb. Cerium is paramagnetic, magnetic susceptibility, (K)+182 2 10-6 (Monatsh. 1899, 20, 369) In dry air the metal is moderately stable, but becomes superficially tarnished with yellow dioxide. The temperature of ignition in oxygen is about 150°-180° (Annalen; 1903, 325, 261) The metal is malleable and ductile. it has a fairly high latent heat of fusion and thermal conductivity. The heat of oxidation of the metal is 1740 cals per gram, or 60,900 cals per gram. equiv. (CeO<sub>2</sub>=243,600); the wire burns even more brightly than magnesium, and combines readily with the halogens. Cerium is very slightly attacked by cold water, a slow evolution of hydrogen occurs in boiling water, and the metal is tarnished black, it is readily dissolved by dilute acids, but concentrated nitric acid produces a brown deposit consisting mainly of dioxide By heating the metal to 345° in hydrogen, cerum hydride is produced, without luminescence. At 1000° it combines with nitrogen to form the nitride, and direct combination also occurs when the metal is heated in sulphur vapour (Hirsch, Met and Chem Eng. 1911, 9, 540) Alloys have been prepared by melting cerium with other metals under molten sodium and potassium chlorides, the combination with aluminium, magnesium, bismuth, silicon, or zinc is attended by a generation of heat; liquid and solid cerium amalgams have also been obtained. Cerium silicide (CeSi2) is formed by the reduction of cerium oxide with graphite and silicon. This compound is brittle and when added to cerium in such proportion that the mixture contains about 15 pc. of silicon, forms a good pyrophoric alloy. The magnesium-cerium alloys are excellent for thermal reduction processes since—the combination of the two metals being endothermic-

more heat is developed by the oxidation of the alloy than by that of the two metals in admixture (Hirsch).

When scratched or rubbed with a steel edge, cerium emits brilliant sparks, and this property has been utilised in the preparation of gas lighters. For a description of pyrophoric cerium alloys, see Engineering and Mining Journal, 1917, p. 212, Vogel, Zeitsch. anorg. Chem. 1917, 99, 25.

Separation and purification of ceria.—Of all the rare earth oxides, ceria is most readily isolated in a pure state, and the following methods have been employed in separating this oxide from lanthana, prascodymia and neo-

dymia:-

(1) Treatment of the hydroxides suspended in aqueous caustic alkalis with chlorine, ceric hydroxide iemains undissolved, while the other hydroxides are converted into soluble chlorides and hypochlorites (Mosander, J pr Chem. 1842, 30, 276. Mengel, Zeitsch anorg. Chem. 1899,

19, 67)

(2) The mixed oxides dissolved in hot nitric acid and the solution evaporated down with ammonium nitrate until ceric ammonium nitrate separates. The double salt is crystallised from nitric acid until the oxide left on ignition has a light-yellow colour (Auer von Welsbach, Monatsh. 1881, 5, 508, Schottlander, Ber. 1892, 25, 378); or the mixed oxides are treated with four times their weight of concentrated nitric acid and gradually heated to 60°. The solution, after decantation from any insoluble matter, is evaporated to a syrupy consistency, mixed with water from 10 to 30 times the weight of the oxides taken, and 1 gram of ammonium sulphate added for each litre of The fiquid is boiled for a few minutes, solution when a pale greenish-yellow precipitate of basic ceric nitrate and sulphate is formed, which is separated and washed with a solution of a mixture of ammonium nitrate and sulphate (Wyrouboff and Verneuil, Sterba)

(3) The solution of mixed nitrates neutralised with ammonia and boiled with magnesium or sodium acetate and hydrogen peroxide whereby cerium is completely precipitated as a basic acetate (Meyer and Koss, Ber. 1902, 35, 672).

- (4) The mixed ammonium double nitrates boiled with aqueous ammonium persulphate, calcium carbonate being gradually added to keep the solution neutral. The precipitate contains all the cerium as ceric nitrate and sulphate mixed with calcium sulphate (Witt and Theel, Ber. 1900, 33, 1315, cf. Meyer and Marckwald, ibid 3003)
- (5) The joint action of potassium permanganate and a basic substance (mercuric oxide, zinc oxide, magnesia, sodium carbonate, &c) in neutral solution. The precipitate contains all the cerium as ceric hydroxide (Stolba, Ber. bohm Ges Wiss. 1878, Drossbach, Ber. 1896, 29, 2452, D. R. P 143106; Meyer, Zeitsch. anorg. Chem. 1903, 37, 878).

The last of the foregoing methods is generally applicable to the mixed oxides of the rare earths, and permits of the complete separation of cerium in one operation. These processes require to be repeated before the cerium is completely freed from the last traces of lanthanum; thorium also is removed completely, only with

CERIUM. 141

some difficulty. The solution of nitrate, mixed with an excess of ammonium carbonate, is treated with ammonia when the greater portion of the thorium is precipitated. To remove the remainder, anhydrous cerium sulphate is prepared and added to ice-cold water, when the pure hydrated salt separates (Wyrouboff and Verneurl, Compt. rend 1897, 124, 1230; 125, 950; 128, 1331; Urbain, Ann Chim Phys. 1900, [7] 19, 184; Sterba, *ibid* 1904, [8] 2, 193; Brauner, Zeitsch. anorg. Chem. 1903, 34, 103; Meyer, *ibid*. 37, 378; Orloff, Chem. Zeit. 1906, 30, 733, 1907, 31, 562; Neish, J. Amer. Chem. Soc. 1909, 31, 517).

Only a small proportion of the cerium oxide extracted from monazite is used in the mantle industry, the greater amount is employed in the manufacture of sparking alloys for eigar lighters, &c. A further quantity is used (in the form of fluoride) for impregnating arc-light carbons; in weighting silk, in the dyeing and photographic industries, and to a slight

extent in medicine.

Cerium forms several series of compounds derived from the oxides Ce<sub>2</sub>O<sub>3</sub>, Ce<sub>4</sub>O<sub>7</sub>(?), CeO<sub>2</sub>, and CeO<sub>3</sub>.

#### CEROUS COMPOUNDS.

Cerium hydride CeH<sub>3</sub>(?). Cerium combines readily with hydrogen at 250°-270°, yielding a brownish-black amorphous hydride which not improbably consists to some extent of CeH4. This product is stable in dry air at the ordinary temperature, but on heating it is converted explosively into cerium oxide and nitride (Matignon, Compt. rend 1900, 131, 891; Muthmann and Kraft, Annalen, 1903, 325, 281

Cerous fluoride 2CeF<sub>3</sub>,H<sub>2</sub>O (John, Bull Soc chim 1874, [2] 21, 533), known also in the anhydrous condition, and employed in the electrolytic production of the metal (v. supra, and cf. Brauner, Ber 1881, 1944, 1882, 109, 115,

Monatsh. 1882, 3 1)

Cerous chloride CeCl<sub>3</sub>,7H<sub>2</sub>O or 2CeCl<sub>3</sub>,15H<sub>2</sub>O (J. Amer. Chem. Soc. 1894, 16, 649; Ber. 1902, 35, 2622)

The anhydrous chloride, a fusible deliquescent crystalline mass, is obtained by the following

- (1) Heating the dry oxalate or cerium dioxide in hydrogen chloride (Robinson, Proc. Roy. Soc 1884, 37, 150).
- (2) Decomposing cerium carbide with chlorine or hydrogen chloride
- (3) Reducing anhydrous cerous sulphate to sulphide by heating in hydrogen sulphide and decomposing the latter with hydrogen chloride (Ber. 1899, 32, 3413) Cerous sulphide and hydrogen bromide yield cerous bromide.

(4) The anhydrous sulphate, when heated in a current of chlorine and sulphur chloride is converted directly into cerous vapour,

chloride.

The dry chloride is readily soluble in alcohol and ebullioscepic determinations of the molecular weight in this solvent gave numbers corresponding with CeCl<sub>s</sub> (Muthmann, Ber. 1898, 31, 1829).

It forms double salts with the chlorides of mercury, gold and platinum, and of certain organic bases.

Cerous bromide CeBr<sub>2</sub>, and lodide CeI<sub>2</sub> are

also known, as are the bromate Ce(BrO<sub>2</sub>)<sub>3</sub>,9H<sub>2</sub>O,

and sodute Ce(IO<sub>3</sub>)<sub>3</sub>,2H<sub>2</sub>O.

Cerous hydroxide Ce(OH)<sub>3</sub>, a white precipitate obtained from soluble cerous salts with ammonia or caustic soda, is stable only in absence of oxygen. Exposed to air it becomes slowly oxidised to yellow ceric hydroxide, passing through an intermediate reddish-violet stage xCe(OH)<sub>3</sub> yCe(OH)<sub>4</sub> (J. Amer. Chem. Soc. 1894, 18, 649).

Cerous oxide has not been isolated with certainty, since the ignition of decomposable cerous salts (nitrate, sulphate, oxalate, &c.) leads to the production of cerium dioxide. The reduction of the latter oxide with hydrogen at high temperatures gives rise to an unstable darkblue oxide approximating in composition to Ce<sub>4</sub>O<sub>7</sub>

Cerous sulphide Ce<sub>2</sub>S<sub>3</sub> (v. supra), a reddish purple substance (Ann. Chim. Phys. 1904, [8] 2, 193), is moderately stable in air and slowly

decomposed by boiling water.

Cerium nitride CeN, obtained by heating cerium in nitrogen (Annalen, 1903, 325, 261), or by passing ammonia over heated cerium carbide (Compt. rend. 1900, 131, 865), is decomposed by water, yielding cerium dioxide, ammonia and hydrogen. An explosive cerium azide is precipitated by adding sodium azide to aqueous cerous nitrate (Curtius and Darapsky, J. pr. Chem 1900, [2] 61, 408).

Cerous nitrate Ce(NO<sub>3</sub>)<sub>3</sub>,6H<sub>2</sub>O, prepared by (1) dissolving cerium dioxide in nitric acid in the presence of a reducing agent; (2) decomposing cerous oxalate with strong nitric acid. Deliquescent triclinic crystals. When heated strongly, pure cerous nitrate leaves a residue of pale-yellow cerium dioxide, but when praseodymium nitrate is present, the residual oxide is brown (J. Amer. Chem. Soc. 1909, 31, 517).

Cerous sulphate  $(e_2(SO_4)_3)$ . The anhydrous sulphate is produced by dehydrating the hydrated salt at 400° Above 500° it decomposes, leaving a residue of dioxide A saturated solution of the sulphate at 0° deposits the dodecahydrate Ce<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub>,12H<sub>2</sub>O (asbestos-like needles) Above 5° this hydrate is labile, and at higher temperatures the solution of the sulphate deposits hydrates containing 12, 9, 8, 5, and 4 molecular proportions of water. With the sulphates of ammonium, sodium, and potassium cerous sulphate forms double sulphates isomorphous with the corresponding salts of lanthanum, neodymium, and praseodymium. In moderately strong sulphuric acid, cerous sulphate gives rise to the acid sulphate Cc(HSO<sub>4</sub>)<sub>3</sub> (Brauner and Picek, Zeitsch. anorg. Chem. 1904, 38, 322). This acid salt has been recommended as an oxidising catalyst in the production of aniline black, 03 gram of cerous oxide in this form sufficing to develop the black from 6 kilos of aniline hydrochloride.

Cerous carbonate Ce<sub>2</sub>(CO<sub>3</sub>)<sub>3</sub>,5H<sub>2</sub>O, prismatic crystals, precipitated from solutions of cerous salts by ammonium carbonate, forms soluble crystalline double carbonates with the alkali and ammonium carbonates.

Cerous cobalticy anide  $Ce_2(COC_6N_6)_2, 9H_3O$ . The most soluble of the rare-earth cobaltieyanides in 10 p.c. hydrochloric acid (1000 pts. of saturated solution contain 10.75 parts of the 142 CERIUM.

Cerous oxalate Ce<sub>2</sub>(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub>,10H<sub>2</sub>O, crystalline white precipitate (Power and Shedden, J. Soc. Chem. Ind. 1900, 19, 636), has been used medicinally as a sedative in sickness, particularly that of pregnancy, also in chronic diarrheea, hysteria, epilepsy, and migraine

Cerous phenoxide has been recommended as a non-irritant disinfectant (J Soc. Chem. Ind.

1909, 1221; D. R. P. 214782).

Cerous acetylacetonate Ce(CH<sub>2</sub> CO CH CO CH<sub>3</sub>)<sub>3</sub>,3H<sub>3</sub>O, obtained by the interaction of cerous ammonium nitrate, acetylacetone, and ammonia (Urbain, Ann. Chim. Phys 1900, [7] 19, 184; Biltz, Annalen, 1904, 331, 334)

Cerous dimethyl phosphate Ce2(Me2PO4), H2O white, hexagonal plates, solubility 796 at 25° and about 65 at 95° (Morgan and James, J. Amer. Chem Soc 1914, 36, 10).

For other cerous salts of organic acids, v Biltz, Annalen, 1904, 331, 334; Wolff, Zeitsch anorg Chem 1905, 45, 89; Morgan and Cahen, Chem Soc Trans 1907, 91, 477; Pharm J. 1907, 78, 428; Rimbach and Kilian, Annalen, 1909; 368, 110; Pratt and James, J. Amer Chem. Soc. 1911, 33, 1330, Katz and James, ibid. 1913, 35, 872; Erdmann and Wirth, Annalen, 1908, 361, 190; Rimbach and Schubert, Zeitsch physikal. Chem 1909, 67, 183, Jantsch and Grunkraut, Zeitsch, anorg Chem 1913, 79, 305

For complex cerum salts, v. Wyrouboff and Verneuil, Ann. Chim Phys. 1906, [8] 9, 289, Barbieri, Atti R. Accad. 1908, [5] 17, 1, 540

#### CERIC COMPOUNDS.

Cerium carbide CeC<sub>2</sub>; microscopic crystals, sp gr 5 23, prepared by melting together in the electric furnace cerium dioxide (192 parts) and sugar carbon (48 parts), using a current of 300 amperes and 60 volts (Moissan, Compt. rend. 1896, 122, 357, 1897, 124, 1233)

$$\begin{array}{c} 2\text{CeO}_{2} + \text{C} = \text{Ce}_{2}\text{O}_{3} + \text{CO} \\ \text{Ce}_{2}\text{O}_{3} + 9\text{C} = 2\text{CeC}_{3} + 3\text{CO} \\ \text{CeC}_{3} = \text{CeC}_{3} + \text{C} \end{array}$$

When decomposed with water, the carbide yields 75.5 p.c acetylene, 20.5 p.c. methane, and 4 p.c.

ethylene

Cerium silicide CoS12; small crystals with a steely lustre, produced by heating cerium dioxide with crystalline silicon, stable in air and decomposed by acids (Sterba, Compt. rend. 1902, 135, 170)

Ceric oxide (syn. Cerium dioxide) CeO<sub>2</sub>; a white or light-yellow powder, sp gr 6 405; also in isotropic crystals, sp gr 6 76-7 905. The in isotropic crystals, sp gr 6 76-7 905. amorphous oxide is prepared by (1) the oxidation of the metal in oxygen; (2) the ignition of cerous salts (nitrate, sulphate, carbonate, oxalate) or ceric compounds (hydroxide, nitrate, sulphate) The colour of ceric oxide is pure white when prepared at low temperatures, but becomes pale yellow with decrease in volume on ignition (Spencer) It darkens when heated, but becomes nearly colourless or at most pale yellow when cooled The colour of ordinary ceria is probably due to traces of praseodymium The ignited oxide is insoluble in hydrochloric or nitric acid or in dilute sulphuric acid. The crystalline form, obtained by heating the amorphous variety with sodium chloride, borax or potassium hydrogen sulphate, is quite

unaffected by acids or alkalis. The dioxide is reduced with difficulty by aluminium and magnesium, when the corresponding cerium alloys are produced. For further references to cerium dioxide, v. Compt. rend. 1895, 120, 663; 1897, 124, 618, 1233, 1300; 125, 950; 1901, 133, 221; Ber 1900, 33, 3003, 3506; Ann Chim. Phys. 1904, [8] 2, 193; Zeitsch. anorg. Chem 1903, 34, 103, 207; 37, 378); Hofmann and Höschell, Ber. 1914, 47, 238; Jiede and Birbrauer, Zeitsch anorg Chem 1914, 87, 160; Bekk, Ber. 1913, 46, 2574, Reimer, J. Amer. Chem. Soc. 1915, 37, 1636).

Ceric hydroxide Ce(OH), a yellowish precipitate obtained from soluble ceric salts by means of ammonia or the caustic alkalis. Cerous hydroxide is slowly oxidised to ceric hydroxide by atmospheric oxygen and more rapidly by

mild oxidising agents.

Although ceric hydroxide is more stable than the lower hydroxide, the ceric salts are less stable than the cerous compounds. Ceric chloride itself has not been isolated, but its additive compounds, R2CeCl6, with the salts of organic bases (pyridine, quinoline, triethylamine, &c.) are stable in alcoholic solution.

Colloidal ceric hydroxide is obtained by dialysing an aqueous solution of ceric ammonium nitrate The hydrosol is readily coagulated and dues to a gummy mass, soluble in hot water (Biltz, Ber. 1902, 35, 4431, 1904, 37, 1095).

Ceric sulphate  $Ce(SO_4)_2$  The anhydrous salt is produced by heating the dioxide with strong sulphuric acid With the diluted acid oxygen is evolved, and the resulting solution deposits, on concentration, ceri-cerous hydrogen sulphate  $2\text{Ce}(SO_4)_2, \text{Ce}_2(SO_4)_3, \text{H}_2SO_4, 24\text{H}_2O$ (Brauner. Zeitsch. anorg Chem. 1904, 39, 261), and then yellow hydrated ceric sulphate, Cc(SO<sub>4</sub>)<sub>2</sub>,4H<sub>2</sub>O. The former of these compounds corresponds with the intermediate oxide  $Ce_4O_7$  (=2 $CeO_2$ ,  $Ce_2O_3$ ).

Ceric sulphate forms a series of double sulphates (e.g.  $Ce(SO_4)_2$ ,  $2K_2SO_4$ ,  $2H_2O$ ) with the

alkalı sulphates.

Basic ceric sulphates, 4CeO<sub>2</sub>,3SO<sub>3</sub>,12H<sub>2</sub>O, and 2CeO<sub>2</sub>,3SO<sub>3</sub>,4H<sub>2</sub>O have been described by Hauser and Wirth (Zeitsch anorg Chem. 1908),

and CeO<sub>2</sub>SO<sub>3</sub>,2H<sub>2</sub>O, and 3CeO<sub>2</sub>,4SO<sub>3</sub> by J. F. Spencer (Chem. Soc. Trans 1915, 107, 1265).

Ceric nitrate Ce(NO<sub>3</sub>)<sub>3</sub>·OH,3H<sub>2</sub>O. This basic salt separates from a solution of ceric hydroxide in nitric acid. The normal ceric nitrate has not been isolated, but the double nitrates R<sub>2</sub><sup>I</sup>Cc(NO<sub>3</sub>)<sub>6</sub>, where R<sup>I</sup> is ammonium or an alkali metal, are well-defined crystalline substances, having a bright-red colour; they are very hygroscopic and readily soluble in alcohol or The double nitrates R<sup>II</sup>Ce(NO<sub>3</sub>)<sub>6</sub>,8H<sub>2</sub>O, where R=Mg,Zn,N1,Co,Mn, form a well-defined These cerium double nitrates are quite analogous to the corresponding double nitrates of thorum.

For other ceric salts, v. Barbieri, Atti R. Accad. Lincei, 1907, [5] 16, 644; Ber. 1910, 43, 2214. For double ceric fluorides, v. Brauner, Ber. 1881, 1944; Monatsh. 1882, 3, 1; Chem. Soc. Trans. 1882, 41, 68; Rimbach and Kilian, Annalen, 1909, 368, 101; Pozzi-Escot, Compt. rend. 1913, 156, 1074; Browning and Flora, Amer, J. Sci. 1903 [iv.] 15, 177).

For ceric acetylacetonate, Ce(CHAc<sub>2</sub>)<sub>4</sub>, ese Job and Goissedlt, Compt. rend. 1913, 157, 50.

Cerium peroxide and its derivatives. The addition of ammonia and hydrogen peroxide to a solution of a cerous salt determines the precipitation of reddish-brown hydrated cerium peroxide Ce(O<sub>2</sub>H)(OH)<sub>3</sub>. When cerous nitrate or chloride is added to concentrated aqueous potassium carbonate, the solution takes up atmospheric oxygen and assumes a deep-red The liquid slowly deposits crystalline dark-red potassium per-ceric carbonate  $4K_2CO_3$ ,  $Ce(CO_3)_3O_3$ ,  $12H_2O$  In the presence of glucose, this alkaline solution acts catalytically When shaken in the air, oxygen is taken up and the per-ceric compound produced On standing, this substance is reduced by the glucose to cerous salt, and the colour disappears peated shaking leads to renewed production of per-ceric compound, so that oxidation and reduction proceed alternately to an indefinite extent (Pissarjewski, J. Russ. Phys. Chem Soc 1900, 32, 609, Zeitsch, anorg. Chem 1902, 31, 359, Wyrouboff and Verneun, Comp. 121, 1898, 127, 863, Job, Compt. rend. 1898, 126, 246, 1899, 128, 178, 1089, 1902, 134, 1052; 1903, 136, 45, Engler, Zeitsch. anorg. Chem. 1903, 36, 2642, 1904, 37, 49, 3268).

Applications of Cerium.—Metallic cerium (in the form of 'mischmetal,' v. ('ERIUM METALS), has been used in the reduction of the refractory oxides of columbium, tantalum, molybdenum, &c.

The oxidising action of cerium compounds has led to their use in photography (Lumière, Compt. rend 1893, 116, 574), and as oxidising catalysts in organic preparations (Atti R. Accad. Lincei, 1907 [5] 16, i 395), and as an ingredient of 'driers.' The oxides have been suggested for use in colouring glass (Chem. Ind. 1904, 27). Cerium salts have been employed as mordants in dyeing In the preparation of leather, hide powder decomposes cerum salts, fixing the element in the form of hydrated oxide. hide partially reduces ceric sulphate or ceric ammonium nitrate, absorbing oxygen yielding a good quality of leather which has a yellow tint and resists the action of water (Garelli, Atti R. Accad Lincei, 1907, [5] 16, 1 **532).** 

In addition to cerous oxalate, other salts (e.g. hypophosphate, and double ammonium eitrate, tartrate and lactate) have been utilised medicinally in the treatment of sea-sickness and nervous disorders. Incandescent gas mantles contain about 1 p.c. of ceria and 99 p.c. of thoria (Meyer and Anschütz, Ber. 1907, 40, 2639).

(For the detection and estimation of cerium, v. Analysis.)

7. ANALYSIS.)

For a bibliography of cerium, see Hirsch,
J. Soc Chem. Ind. 1912, 477.

CERIUM METALS AND EARTHS. (The

Rare Earths.) The so-called rare earths include a scries of basic oxides, very similar in physical and chemical properties, which are generally found associated in certain rare minerals of complex composition (silicates, phosphates, titanates, columbates, tantalates, zirconates).

The metals contained in these oxides may be divided roughly into two groups, named after the minerals cerule and ytterbite, in which the rare earths were first discovered.

The cerite metals (cerium group) are cerium, lanthanum, neodymium, praseodymium and samarium.

The ytterbite metals (yttrium group) include dysprosium, holmium, erbium, europium, qadolinium, lulecium, celtium, scandium, terbium, thulium, ytterbium (neoytterbium), and yttrium of the latter section, europium, gadolinium, and terbium form an intermediate series (the terbium group) connecting the cerium metals with the remaining rate earths of the yttrium group proper. Closely associated with these elements is the metal thorium, and zirconium is generally included in the category of the metals of the rare earths.

Historical Summary. In 1794 ytterbrte, obtained from Ytterby near Stockholm, was shown by Gadolin, its discoverer, to contain a new oxide, to which Ekeberg subsequently gave the name yttria. From the Swedish mineral cerite, Klaproth, in 1803, isolated another new oxide which is called ceria by Berzelius and Hisinger, who also examined it Thoria was isolated by Berzelius in 1828 from the cilicate

thorite, found at Brevig in Norway. In 1839 Mosander isolated the oxides lanthana and didymia (Annalen, 1839, 32, 235; 1842, 44, 125) from cerite, and also effected a further separation from ytterbite of the earths erbia and terbia (Phil Mag. 1843, 23, 251, J pr Chem. 1843, 30, 288). In 1878 Marignae separated from crude crbia a less basic coloui less component which he called ytterbia (Compt. rend 1878, 87, 578), and this fractionation was speedily followed by Nilson's discovery of scandia, the least basic of the ytterbite earths, and its identification as the oxide of the hypothetical ekaboron, predicted by Mendeléeff. In the same year Cleve effected a further separation of the old 'erbia' and found two new earths, namely thulia and holmia (Compt rend 1879, 89, 478, 708). The doubt expressed by Mangnac as to the homogeneity of didymia was shared by Delafontaine, on the ground of spectroscopic evidence (Compt. rend. 1878, 87, 634), and substantiated by Lecoq de Boisbaudian (Compt. rend. 1879, 88, 323), who isolated from this earth the less basic oxide samaria. The didymia left after the removal of the last-mentioned oxide was still regarded as a mixture by Brauner (Monatsh 1882, 3, 486; Chem. Soc Trans. 1882, 43, 278), and in 1885 Auer von Welsbach, by fractional crystallisation of the double didymium nitrates of ammonium and potassium, succeeded in separating this earth into two components, which he named neodymia and praseodymia (Monatsh. 1885, 6, 477). Lecoq de Boisbaudran now showed that Cleve's holmia was a mixture containing a new constituent dysprosia (Compt. rend. 1886, 102, 1003), the simple nature of which was afterwards confirmed by Urbain (Compt. rend. 1904, 139, 736, and 1905, 141, 521).

Gadoinna, which had been recognised by Marignac in samarskile (Compt rend. 1880, 90, 899), was isolated in a more definite form by Lecoq de Boisbaudran (Compt. rend. 1889, 108, 165; 1890, 111, 409, 472), and by Bettendorf (Annalen, 1890, 256, 159, 1891, 263, 164; 1892, 270, 376), and further purified by Demarçay and by Urbain and Lacombe (v. GADOLINIUM). Further researches by Demarçay on Cleve's samaria led to the isolation of europia (Compt.

rend. 1896, 122, 728), an earth provisionally symbolised as Z, and afterwards shown to be identical with oxides indicated by the symbols  $Z_e$  and  $Z_t$  (Lecoq de Bosbaudran, Compt. rend. 1892, 114, 575, 1893, 116, 611, 674) and  $S_\delta$  (Crookes, Compt. rend. 1885, 100, 1380, 1495; Proc. Roy. Soc 1885, 38, 414).

Terbia, which occurs only in small amount in ytterbite, has been isolated in a purified state by Urbain (v Terbium), and Marignac's ytterbia (i.c) has been fractionated by Auer von Welsbach (Monatsh. 1906, 27, 935), and separated into two components, neoytterbia and lutecia, by Urbain (Compt rend. 1907, 145, 759) In 1911 Urbain discovered that the lutecium from gadolinite is accompanied by a small quantity of a rare earth metal which he called celtium (Urbain, Compt. rend 1911, 152, 141).

Occurrence (v. Das Vorkommen des seltenen Erden, Schilling, 1904, Phipson, Chem. News, 1896, 73, 145) The following is a short description of the most important sources of the rare

(For details concerning the various minerals, see under their respective headings)

### 1. Minerals containing thiefly Cerium Earths.

Cerite (ochroite, cererite, cerinstein), a hydrated silicate containing 60-70 p c of cerium and its allies, together with smaller amounts of iron, calcium, and the yttrium earths, found in the Bastnas mine at Ryddarhyttan in Sweden.

Orthite (allanite, bodenite, cerine, muromontite, pyrorthite, tautolite, uralorthite, wasite). a double silicate of aluminium and the cerium metals (16-25 pc), together with smaller quantities of thorium, iron, calcium, and the yttrium metals, found in Greenland, Scandinavia (Arendal, Hitteroe, Stockholm), United States (New York, North Carolina, Pennsylvania, Virginia, Texas), Ural Mountains (Miask), and in several localities in Germany

Monazite (cryptolite, edwarsite, eremite, mengite, turnerite), an orthophosphate of the cerium metals (Ce, La, Nd, Pr)III PO<sub>4</sub>, containing 40-70 pc of these elements together with varying amounts of thorium (0-18 pc), it occurs in the plutonic and older volcanic rocks, and in the alluvial sands of the rivers and coasts of Brazil, North and South Carolina, and Travancore, in the Federated Malay States, Ceylon and Nigeria, &c

Eschenite A complex, columbate-titanate, containing approximately equal proportions (13-23 pc) of thorum and the cerum metals, together with a small proportion of the yttrium group, found at Hitteroe (Norway) and Miask (Ural Mountains)

#### 2. Minerals containing chiefly Yttrium Earths.

Ytterbite (gadolinite), a basic silicate contaming yttrium earths (40-48 p.c.) and cerium earths (5-10 pc), together with glucinum and iron; found in Colorado, Germany (Harz Mountains and Silesia), Scandinavia (Fahlun, Hitteröe, Ytterby), and Texas.

Yttrialite, a silicate of the yttrium metals (43-47 pc), thorium (10-12 p.c.), and cerium metals (5-8 p c.), found in Texas.

Xenotime, a phosphate of the yttrium metals

(54-64 p.c.), corresponding with monazite; it contains also smaller amounts of the cerium metals and thorium; found in Brazil, North Carolina, Georgia, Norway, Sweden and Switzer-

The following minerals contain the yttrium earths and smaller quantities of thoria and the cerium earths united with varying proportions of the more acidic oxides of columbium, tantalum, titanium, tungsten, tin, uranium, silicon, and zirconium

Fergusonite (bragite, kochelite), found in Ceylon, Greenland, Arendal, Ytterby, Massa-chusetts, North Carolina, and Texas.

Yttrotantalite, found in Norway, Sweden, and the Ural Mountains.

Samarskite (nohlite, vietinghofite, ytteroilmenite), found in Canada, Connecticut, North Carolina, and Ural Mountains (Miask)

Polycrase, found in Canada, Scandinavia, and North and South Carolina

Euxenite (loranskite), found in North Carolina, Norway Hitteroe, Brevig), Swaziland

Many other rare complex minerals (fluorides. zircono-silicates, titanates, columbates, tantalates, uranates, &c) contain considerable amounts of the rare earths, which are also not infrequently found in small quantities in commoner minerals (e q limestones, apatites, scheelites, &c ) Traces of the rare earth metals have been detected in mineral waters, in the ashes of plants, in urine, and in bone ash.

Extraction of the Rare Earths -The finely powdered mineral, made into a paste with concentrated sulphuric acid, is heated till the mass becomes dry and hard. The product is extracted with water, the solution treated with hydrogen sulphide to remove copper, bismuth, molybdenum, &c., and the metals of the rare earths precipitated as oxalates by the addition of oxalic acid. The oxalates are decomposed at 400°, the residual oxides dissolved in dilute sulphuric acid, and the solution saturated with sodium or potassium sulphate, when the double sulphates of the cerium metals are precipitated, whilst those of the yttrium metals remain

The complex minerals containing columbium. tantalum, titanium, &c., may be decomposed by hydrofluoric acid, when the foregoing elements form soluble fluorides, whilst the fluorides of the rare earth metals are precipitated. Or these minerals may be broken up by heating with strong sulphuric acid or potassium hydrogen sulphate and the rare earths precipitated by ammonia from the solution of their sulphates. These hydrated oxides are freed from the coprecipitated columbium, tantalum, and titanium hydroxides by prolonged boiling with nitric acid, when the latter hydroxides separate in an insoluble form. The rare earths are precipitated as oxalates from their solution in nitric acid, the oxalates ignited, and the resulting oxides subjected again to the nitric acid treatment until a complete separation is effected. Or the minerals may be heated in the vapour of disulphur dichloride S<sub>2</sub>Cl<sub>2</sub>, or carbonyl chloride COCl<sub>2</sub>, when the chlorides of columbium, tantalum, titanium, &c., are volatilised, and the rare earth metals are converted into non-volvtile chlorides (Hicks, J. Amer. Chem. Soc. 1911, 33, 1492; Barlot and Chauvenet, Compt. rend. 1913, 157, 1153).

The methods of separating and purifying the rare earths, which will be further described under the appropriate headings (v. CERIUM, YTTRIUM; TERBIUM; THORIUM, &c.) may be divided into the following two classes.—

1. Separations based on differences of

basicity.

a. The fractional precipitation of the earths with bases of varying strengths: ammonia, caustic alkalıs, magnesium hydroxide, organic bases.

b. The fractional decomposition by heat of the solid nitrates and other salts.

2. Separations based on differences of solu-

bility.

Considerable ingenuity has been shown in separating the metals of the rare earths by taking advantage of the slight differences of solubility exhibited by their salts with acids of widely different types. The following series does not exhaust the list of compounds which have been utilised in these intricate separa-Sulphates, double sulphates, nitrates, double nitrates, chromates, bromates, formates, oxalates, succinates, ethyl-sulphates, acetylacctonates, dimethyl phosphates, glycollates, and sulphanilates (if. James, J. Amer. Chem. Soc. 1912, 34, 757).

The progress of these separations is controlled by chemical and physical methods.

I The chemical control consists in determining the equivalent of any fraction either by a gravimetric analysis of its sulphate (Kruss, Zeitsch. anorg. Chem 1893, 3, 44, Wild, ibid 1904, 38, 191; Brill, ibid 1908, 47, 464, Jones, *ibid.* 1903, 36, 92; Brauner and Pavhček, Chem. Soc. Trans 1902, 81, 1243), or by a volumetric analysis of its oxalate (Feit and Przibylla, Zeitsch. anorg Chem. 1905, 43, 202).

2. The physical methods consist in observing the magnetic susceptibility of the fractions and their spectroscope characters. The rare earths differ considerably in magnetic susceptibility, and determinations of its amount are readily made by the magnetic balance of Curie and Chereveau (Urbain, Compt rend 1908, 146, 406, 922, 1908, 147, 1286, 1909, 149, 37, 1910, 150, 913; 1911, 152, 141, 1914, 159, 323).

The most convenient spectroscopic con-

trols are (1.) the absorption spectra, exhibited by solutions of the salts of many rare earths (e g neodymium, praseodymium, samarium, terbium, erbium, thulium, dyspiosium and europium), (11.) the arc spectra, in which all the rare earths give characteristic lines both in the visible and ultra-violet regions of the spectrum. Carbon electrodes are employed, the lower one containing a hollow in which is placed the salt under examination. The arc spectra are photographed and the lines identified by comparison with the photograph of an iron spectrum taken under similar conditions The method is extremely sensitive, although not equally so for different elements (cf. Spectrum analysis and v. Exner and Hascheck, Die Wellenlangen der Bogenspektra, 1904; and Hagenbach and Konen, Atlas der Emissionspektren der meisten Elemente, 1905). Other emission spectra (the flame, spark, and cathode luminescence spectra) have been found to give useful indications in certain cases

(Crookes, Phil. Trans. 1883, 174, [3] 891; Proc Roy. Soc. 1885, 38, 414; 1886, 40, 77, 236, 502. Chem. News, 1886, 54, 39, 54, 63, 76, 155; 55, 83, 95; 56, 59, 62, 72, 81; Chem Soc. Trans 1889, 55, 255; Proc Roy Soc. 1899, 65, 237; Lecoq de Bosbaudran, Compt. rend 1885, 100, 1437; 101, 552, 588; 1886, 102, 153, 899; 1536, 103, 113, 627; 1887, 105, 258, 301, 343, 784; 1890, 110, 24, 67; Bettendorf, Annalen, 1892, 270 376; Baur and Marc, Ber. 1901, 34, 2460).

PHYSICAL AND CHEMICAL PROPERTIES OF THE RARE EARTHS AND THEIR MORE IMPOR-TANT COMPOUNDS.

Hydrides. The rare earth metals combine directly at 220°-270° with hydrogen to form brittle amorphous solids ranging in colour from bluish-black (Ce and La), indigo blue (Nd), to green (Pr), and readily dissociated at higher temperatures (Muthmann, Kraft and Beck, Annalen, 1902, 325, 261; 1904, 331, 581).

Oxides and hydroxides. The metals of the rare earths were formerly supposed to be diads (cf. Wyrouboff, Bull Soc chim. 1889, [3] 2, 745, 1899, [3] 21, 118; Compt rend 1899, 128, 1573), but they are now regarded as tervalent elements forming oxides of the type R<sub>2</sub>O<sub>3</sub>, although cerium, neodymium, praseodymium, and terbium are capable of yielding higher The more stable oxides of thorium and zircomum are of the form RO2 The present view of the valency of the metals of the rare earths is supported by the following facts:-

(1) The isomorphism of bismuth nitrate and certain nitiates of the raie earth series (Bodman,

Ber 1898, 31, 1237)

(ii) The cryoscopic and ebullioscopic determination of the molecular weights of the chlorides and acetylacetonates (Ber 1898, 31, 1829; Compt rend 1901, 133, 289, Annalen, 1904, 331, 334); these salts are thus found to have the general formula RX<sub>3</sub>

(III ) The electrical conductivity of the chlorides and sulphates in aqueous solution (Zeitsch physikal Chem. 1899, 30, 193; Amer. Chem J 1898, 20, 606).

(iv ) The specific heats of cerium, lanthanum and didymium (Annalen, 1873, 168, 45; Ber 1881, 2821; Bull Soc chim 1882, [2] 38, 139)

The rare earth oxides vary considerably in basic power, lanthana, praseodymia, neodymia, and ceria, Ce,O3, being the strongest; whilst neoytterbia, lutecia, scandia, and cerium dioxide are among the weakest bases of the series. In general, the oxides (R2O3) of the cerium metals are stronger bases than those of the yttrium

The hydroxides are thrown out of solution as gelatinous precipitates by ammonia, ammonium sulphide, and the caustic alkalis, even in the presence of ammonium salts; but this precipitation is prevented by citric or tartaric acid (except in the case of scandium). Unlike thorium and zirconium, the metals of the cerium, terbium, and yttrium groups are not precipitated as hydrated peroxides from neutral solutions by hydrogen peroxide, but their peroxides separate from alkaline solutions (Brauner, Ber. 1881, 14, 1944; Cleve, Bull. Soc. Chim. 1885, [2] 43, 53). These peroxides are very unstable, evolving oxygen at even the ordinary

temperature; they are completely destroyed by acids (v. Cerium; cf. Melikoff and Pissarjewski, Zeitsch. anoig Chem. 1899, 21, 70). Chlorides. The chlorides of the rare earth

metals are deliquescent salts readily soluble in water or alcohol, and crystallising from the former solvent with 6 or 7 molecular proportions of water. The anhydrous chlorides, which are fusible at red heat, but very slightly volatile are obtained either by evaporating their solutions to drvness with ammonium chloride, by heating the corresponding oxides with carbon in a stream of chlorine, or preferably by heating the oxide or dry sulphate in a current of sulphur chloride or carbonyl chloride or carbon tetrachloride (Matignon and Bourion, Compt. rend 1901, 133, 289, 1902, 134, 657, 1308, 1904, 138, 631, 1905, 140, 1181, Muthmann and Stutzel, Ber. 1899, 32, 3413, Petterson, Zeitsch anorg Chem 1893, 4, 1, Chauveret, Compt rend 1911, 152, 87).

The chlorides of the rare earth metals combine additively with ammonia and organic bases (e.g. pyridine, cf. Compt. rend 1905, 140, 141, J. Amer. Chem. Soc. 1902, 24, 540, 1903, 25, 1128, Ber. 1902, 35, 2622), and exhibit a great tendency to form double salts with the less electropositive metals The platinichlorides of the cerium metals are isomorphous and crystallise in regular octahedra (Ber. 1876, 1722,

Annalen, 1878, 191, 331)

Fluorides. The rare earth metals, including thorum, are precipitated completely as gelatinous fluorides by soluble fluorides and by hydrofluoric and fluosilicie acids, in this respect differing from zirconium, which forms a soluble double fluoride.

Nitrides. The rare earth metals when heated combine directly with nitrogen to form nitrides of the type MN They are brittle amorphous solids, decomposed by moist air or by water, torming ammonia and the corresponding hydroxide.

Nitrates. The nitrates of the rare earths dissolve readily in water or alcohol, those of the cerium group crystallising with 6 molecular proportions of water, whilst the corresponding salts of the yttrium group contain 3-6 molecules of water. The cerium metals readily furnish stable crystallisable double nitrates with the univalent and bivalent metals (ammonium,

sodium, magnesium, &c)

Sulphates. The rare earths dissolve in dilute sulphuric acid, and the solutions deposit hydrated sulphates containing frequently 4, 8, or 12 molecules of water The octahydrated sulphates of praseodymium, neodymium, yttrium, gadolinium, and ytterbium are isomorphous (monoclinic); but the cerous salt  $Ce_2(SO_4)_3$ , 8H2O is rhombic. With excess of acid, the normal sulphates of the rare earths are converted into acid sulphates of considerable stability (Bull. Soc. chim 1889, [3] 2, 745, Compt. rend. 1902, 134, 657; Zeitsch anorg Chem. 1904, 38, 322, Chem. Soc. Trans, 1902, 81, 1243; James and Holden, J. Amer. Chem. Soc. 1913, 35, 559).

The sulphates of the rare earths have the characteristic property of combining with the alkalı sulphates; the double sulphates of the cerium group are only sparingly soluble in concentrated solutions of the alkali salts, whilst those of the yttrium group are readily soluble. The double sulphates of the terbium group occupy an intermediate position, being moderately soluble in solutions of alkali sulphates

Carbides of the raic earth metals may be prepared by heating the oxides with carbon in the electric turnace. They have the general formula MC<sub>2</sub> They are brittle, crystalline solids, transparent in thin layers and of a yellow colour. They are decomposed by water, forming a mixture of acetylene, hydrogen, ethane, and ethylene, and the hydroxide of the metal M(OH)3

Carbonates. Ammonium, sodium, and potassium carbonates precipitate the neutral of basic carbonates from solutions of the rare earths, but these precipitates are soluble in excess of the reagents, and from the solutions thus obtained, crystalline double carbonates are deposited on evaporation. The carbonates of the yttrium metals are generally more soluble than those of the cerium group in solutions of the alkali carbonates.

Oxalates. The rare earth oxalates are precipitated by oxalic acid or soluble oxalates in neutral or acid solutions, and the sparing solubility of these compounds in acids is a characteristic property which serves to separate the rare earths from other metallic bases (Ber. 1898, 31, 1718, 1899, 32, 409)

The oxalates of the rare earths proper dissolve only slightly in ammonium oxalate, differing in this respect from the oxalates of thorium and zinconium, the latter being also soluble in free oxalic acid (Chem Soc Trans. 1898, 73, 951) Aqueous ammonium acetate dissolves the oxalates of the yttirum group (Bull Soc. chim 1896, [3] 15, 338)

Formates. The formates of the cerum group are less soluble than those of the yttrium group These salts have proved useful in separating the elements of the terbium group.

Acetates. The acetates of the cerum group are very soluble and crystallise with difficulty; those of the yttrium group are less soluble Boiling with sodium acetate precipitates ziiconium and thorium from solutions of their soluble salts, but not the rare earths proper, excepting cerium from ceric salts (Bei 1902, 35, 672).

Acety lacetonates. Acetylacetone furnishes well-crystallised salts with the rare earths, and the fractionation of these compounds from alcohol has served in the separation of the yttrium group (Urbain, Compt. rend. 1897, 124, 618, Bull. Soc chim 1897, [3] 17, 98, Ann Chim Phys 1900, [7] 19, 184, Biltz, Annalen, 1904, 331, 60, Zeitsch anorg. Chem. 1904, 40, 218, Hantzsch and Desch, Annalen, 1902, 323, 26, Biltz, *ibid.* 1904, 331, 334, James, J. Amer Chem. Soc. 1911, 33, 1332; Morgan and Moss, Chem Soc Trans. 1914, 105, 189)

Dimethyl phosphates. A solution of the rare earths is made in dimethyl phosphoric acid, the temperature raised and the precipitate collected; additional fractions are obtained by evaporating the mother liquor. In the case of a material containing gadolinium with just sufficient terbium to colour the oxide orangebrown, the gadolinium collects in the most soluble fraction, and the terbium in the least soluble. Lanthanum, cerium, praseodymrum, and neodym'um are left in the mother liquor.

Samarium, europium, and gadolinium are much less soluble than these, but more soluble than terbium, dysprosium, and holmium, erbium thulium, yttrium, and ytterbium collect in the least soluble fractions (Morgan and James, J.

Amer. Chem. Soc 1914, 36, 10).

The Metals of the Rare Earths.—The oxides of the rare earths are very refractory substances, which have not been reduced directly; the metals are generally set free from the anhydrous chlorides. This reduction may be effected with sodium or potassium, but is preferably carried out electrolytically. The dry chlorides, mixed with a small proportion of alkali chloride, or barium chloride, are electrolysed in a copper crucible, fitted with earbon electrodes, the electrolysis being carried out with a current of 30-40 amperes and 12-15 volts. The mixed metal collects in a molten state round the cathode, which passes through the bottom of the The anode is movable, so that the resistance and therefore the temperature can be increased until all the separated metal melts together. A yield of 750 grams of metal can be obtained with a current of 120 amperes in 6 hours.

With the mixed chlorides, prepared from the waste exides of the thoria manufacture, the product is the so-called 'mischmetal' (Ce, 45 pc; La, Nd, Pr, 35 pc.; Sm, Ei, Gd, Y, 20 pc), which has been used, like aluminium in the Goldschmidt process ('thermite'), to effect the reduction of the refractory oxides of molybdenum, vanadium, columbium, and tantalum (Muthmann, Hofer, and Weiss, Annalen, 1902, 320, 331; 1904, 331, 1; 1905, 337, 370).

Bibliography of the Rare Earths—Bohm, Darstellung der seltenen Eiden; Biowning, Introduction to the Raier Elements, Abegg, Handbuch der Anorganischen Chemie, vol in part 1, Herzfeld and Korn, Chemie der seltenen Eiden, Wyrouboff and Verneuil, La Chimie des Terres Rares GTM

CEROSIE v SUGAR CANE WAX AND WAXES. CEROSILINE V WAXES.

CEROTENE C<sub>27</sub>H<sub>54</sub> A crystalline hydrocarbon, m p 65°, found by Konig and Kiesow (Ber 6, 500) associated with cholesterol in the fat wax of hay and straw, and detected by Knecht and Hibbert in ordinary household chimney soot (Mem Manchester Phil. Soc 1914, 58, [2 | 1)

CEROTIC ACID C<sub>25</sub>H<sub>51</sub> COOH is found free in beeswax (Brodie, Phil Trans 1848, 1 147, Marie, Ann Chim. Phys 1896 [7], 7, 145) and combined as ceryl cerotato in insect wax (Biodie, l.c., Henriques, Ber. 1897, 30, 1415). though some doubt has been expressed as to the identity of the latter acid with that in

beeswax.

CEROTIN V WAXES.

CEROTINE-ORANGE, -PONCEAU, -SCAR-LETS, -YELLOW v. AZO- COLOURING MATTERS. CEROXYLIN v. WAXES.

CERULEIN v. ALIZARIN AND ALLIED COLOUR-ING MATTERS

CERUSSITE. Native lead carbonate (PbCO<sub>3</sub>), of some importance as an ore of lead (Pb, 77 5 p.c), being known as white lead-ore. It crystallises in the orthorhombic system and 18 isomorphous with aragonite (CaCO<sub>3</sub>) and

witherite (BaCO<sub>3</sub>). Six-rayed stellate groups of twinned crystals are extremely characteristic. The crystals possess an adamantine lustre, are very heavy (sp gr 65), and effervesce with dilute acid, they are very brittle, breaking with a bright conchoidal fracture; hardness, 3-31. The mineral occurs in the upper oxidised zones of veins of lead ore, having been formed by the action of carbonated surface waters on galena Enormous deposits of cerussite intermixed with embolite (silver chloro-bromide) and iodyrite (silver iodide) have been extensively mined at Broken Hill in New South Wales, where little or none of the oxisided ore now remains mately mixed with hemimorphite (hydrated zinc silicate), it is abundant at Broken Hill in North-Western Rhodesia At both of these localities, many finely crystallised specimens have been

CERYL ALCOHOL v WAXES.

CETIACOL. Trade name for pyrocatechol methylcetyl ether

CETINE v. WAXES

CETOSAN. Trade name for a mixture of the higher alcohols of spermaceti, mainly cetyl and octodecyl alcohols, with vaseline. shaken with water, it forms a permanent creamy emulsion (Blatz, J. Soc. Chem. Ind. 1908, 865)

CETYL ALCOHOL, CETYLIC ACID and CETYL-ACETIC ACID (v SPERMACETI).

CETYLMALONIC ACID

 $\mathrm{CH_3(CH_2)_{15}CH(CO_2H)_2}$ 

Prepared from a-cyanstearic acid and potash (Hell and Sadomsky, Ber. 1891, 2781), or diethylmalonic ester, methyl iodide, and cetyl iodide (Guthzeit, Annalen, 206, 357), m p. 121 5°-122°

CEVADINE AND THE ALKALOIDS OF VERATRUM SPP. The sources principally concerned are the seeds of Schoenocaulon officinale (A Gray) [Asagræa officinalis (Lindley)], and the rhizomes of Veratrum viride (Ait.) (green hellebore) and Veratrum album (Linn.) (white hellebore), all of which belong to the N O. Liliaceae, and have nothing in common with the hellebores proper, which belong to the N O. Ranunculace The alkaloids present in these three drugs are as follows .--

Sabadilla sceds ('evadine (crystallised veratrine)

Cevine (sabadinine) Veratridine (amorphous vera-

Sabadiline (cevadiline)

Sabadine

White Hellebore Protoveratime Jervine

Pseudojervine Rubijervine

Protoveratudine

Veratralbine

Green Hellebore Cevadine

Veratridine Jervine

Pseudojervine Veratralbine

It should be noted that there has been considerable confusion in the nomenclature of these alkaloids. Commercially the name 'veratrine' is applied to an amorphous mixture of alkaloids obtained from sabadilla seeds. The name

verafrine is also frequently applied to veratridine, usually, however, with the prefix 'amorphous,' and to cevadine, with the prefix 'crystalline.' The name veratridine was also applied by Robbins (Pharm. J. 1877-1878, [m.] 8, 316) to a crystalline alkaloid contained in green hellebore, and which, in some respects, resembles protoveratrine It is not identical with Bosetti's veratridine isolated from sabadilla seed (Ber. 1883, 16, 1384).

# Alkaloids of Sabadilla Seeds.

[ Cevadine (crystallised veratrine) C32H49O2N 18 the chief alkaloid of sabadilla or cevadilla seeds

(Schoenocaulon officinale [A. Gray]).

Preparation. The finely-powdered seeds are extracted with alcohol, containing 1 p.c. of tartaric acid. The acid liquid is evaporated and diluted with water, to precipitate resin. To the clear solution soda is added in slight excess, and the mixture completely extracted by repeatedly shaking with other. The othereal solution is mixed with light petroleum and the filtrate allowed to evaporate spontaneously. Crystals of cevadine separate from the syrup, and are recrystallised from their solution in alcohol. The remaining syrup contains veratridine (amorphous veratime) and sabadilline (Wright and Luff, Chem. Soc. Trans. 1878, 33,

338). 'Commercial veratrine' is a macroscopically amorphous mixture of cevadine and veratridine (q.v), possibly with small quantities of other alkaloids It was deleted from the BP. 1914, but is retained in the USP. It is used externally, chiefly against lice. Sabadilla seeds contain about 0.7 pc. of total alkaloids, of which about one-seventh can be crystallised (cevadine)

Properties Rhombic prisms, containing 2C<sub>2</sub>H<sub>5</sub>OH when crystallised from alcohol. The alcohol is lost at 130°-140°, and the alkaloid then melts at 205° with decomposition Nearly insoluble in water, soluble in alcohol or ether Optically inactive (Freund and Schwarz, Ber 1899, 32, 800)

The aurichloride, mp 182° (decomp), and the mercurichloride, mp 172° (decomp), are characteristic crystalline compounds (Ahrens, Ber. 1890, 23, 2700) In carbon disulphide solution additive compounds with chloral and bromal are formed (Frankforter and Kritschevsky, J Amer Chem Soc. 1915, 37, 2567)

Cevadine is intensely poisonous; minute

quantities cause violent sneezing.

Reactions.—When warmed with concentrated hydrochloric acid cevadine produces a fine purplish-red colour, which is a somewhat delicate test for the alkaloid Concentrated sulphuric acid produces a carmine-red colouration, or, if the alkaloid is previously mixed with a little sugar, a blue colour.

Cevadine is a monacidic tertiary base. contains no methoxyl groups and no : NMe group; one HO group is present.

When anhydrous cevadine is distilled β-pico-

line is obtained with tiglic acid.

When heated with alcoholic potash, cevadine undergoes hydrolysis furnishing

C27H48O8N

Luff, Chem. Soc. Trans. 1878, 33, 338; 1879 35, 405; Bosetti, Arch. Pharm. 1883, [iii.] 21 81; Ahrens, Ber. 1890, 23, 2700; Freund an collaborators, ibid. 1899, 32, 801; 1904, 3 1964). Cevine crystallises with 31H2O from dilute alcohol, m.p. 195°-200° (anhydrous), an yields derivatives both with metals and acid It is less toxic than cevadine (Freund, l.c.).

According to Hess and Mohr (Ber. 1919 52 B, 1984) cevine is identical with sabadining which Meick (Arch Pharm. 1891, 229, 164 isolated from the mixed alkaloids of sabadill

seeds.

Veratridine (amorphous veratrine) C<sub>37</sub>H<sub>53</sub>O<sub>11</sub>] is found in sabadilla seeds, and is isolated from the amorphous alkaloids precipitated in th earlier stages of the preparation of cevadin (see above) by treating them with ether, whe they separate into insoluble matter (sabadilline The latter 1 and soluble matter (veratridine). purified by conversion into the sparingly solubl Veratridine is sternutatory, yellowish white, amorphous, soluble in 33 parts of water and melts at 180° On hydrolysis by alkalıs ı yields verine C<sub>2</sub>, H<sub>45</sub>O<sub>8</sub>N, which closely resemble cevine (see above) and veratric acid (3: dimethoxybenzoic acid)

Sabadilline (cevadilline) C<sub>34</sub>H<sub>58</sub>O<sub>8</sub>N 18 second amorphous alkaloid occurring in sabadilla seeds. It yields amorphous salts (W and L lc.)

Sabadine  $C_{29}H_{51}O_8N$ , crystals, mp. 238° 240°, also occurs in sabadilla seeds; the salt are crystalline (Merck, Arch. Pharm. 1891, 229 164)

Sabadıllıne and sabadıne resemble cevadını in action, but are less toxic

Cevadine and veratridine, the two principa alkaloids in 'commercial veratrine,' cannot be separated by crystallisation of the former from alcohol, or extraction of the latter by water but they may be estimated by hydrolysing with alcoholic sodium hydroxide, acidifying, and estimating the volatile tiglic acid (from the cevadine), and the non-volatile veratric acid (from the veratridine), the latter acid is extracted by ether (Wright and Luff, lc.).

#### Alkaloids of White Hellebore Rhizome.

Protoveratrine  $\rm C_{32}H_{51}O_{11}N$  is the character istically toxic alkaloid of white hellebore 'roots. It crystallises in thin four-sided tablets from alcohol, m p 245°-250°, 239°-241° (Bredemann) and dissolves in sulphuric acid with a greer colouration, which gradually passes into deep It is readily decomposed by acids yielding sobutyric acid as one product.

Protoveratrine is accompanied in white

hellebore rhizomes by

Jervine C26H37O3N, prismatic crystals, m.p. 238°-242°; toxic.

Pseudojervine C<sub>29</sub>H<sub>43</sub>O<sub>7</sub>N, six-sided tablets.

m.p. 300°-307°; not poisonous.

Rubijervine C<sub>26</sub>H<sub>48</sub>O<sub>2</sub>N, long prisms, m.p.

240°-246°, 234° (Bredemann); not poisonous. Protoveratridine C<sub>24</sub>H<sub>45</sub>O<sub>8</sub>N, four-sided plates

m p. 265°, not poisonous.

The chief papers relating to protoveratrine and its congeners are Wright and Luff (Chem. Soc. Trans. 1879, 35, 405), Salzberger (Inaug. Diss. Erlangen, 1890; Arch. Pharm. 1890, 228, and angelic and tiglic acids  $C_5H_5O_2$  (Wright and Pehkschen (Inaug. Diss. Dorpat, 1800; J. CHALK. 149

Pharm. Chim. 1890, [v.] 22, 265), and Bredemann (Apoth. Zeit. 1906, 21, 41, 53).

Alkaloids of Green Hellebore Rhizome.

The chief alkaloids occurring in this drug (Veratrum viride, U.S.P.) have been described already, viz. cevadine (see under Alkaloids of sabadilla seeds), jervine, and pseudojervine (see under Alkaloids of white hellebore rhizome). Veratridine (see under Alkaloids of sabadilla seeds) occurs only in traces in green hellebore. Robbins isolated from green hellebore rhizomes (Pharm. J. 1877-1878, [iii.] 8, 316) a crystalline alkaloid which he called veratridine, and which, like protoveratrine of white hellebore, gives a green colouration with sulphuric acid. The amorphous alkaloid veratralbine was also obtained by Wright and Luff from this source (Chem. Soc. Trans. 1879, 35, 421).

CEYLON MOSS v. ALGÆ CHAGRAL GUM v. GUMS.

CHAILLETIA TOXICARIA (Don) A plant found in Upper Guinea, Sierra Leone, and Senegambia, the seeds of which contain a poisonous resin used for killing rats and other animals (Power and Tutin, J. Amer. Chem. Soc. 1906, 28, 1170).

1906, 28, 1170). CHALCANTHITE. Native copper sulphate, CuSO<sub>4</sub>,7H<sub>2</sub>O, found as a bright-blue efflorescence, or rarely as distinct crystals, in many copper mines, being especially abundant in some of the Chilean mines. In solution it is always present in the waters issuing from copper-mines; these are allowed to flow over scrap iron, the copper being precipitated as cement-copper. About 100 tons of copper have been so saved annually from the waters of the Rio Tinto mine in Spain, and considerable amounts were formerly obtained from the mine waters in Co. Wicklow.

CHALCEDONY. A native form of silica occurring in reniform, botryoidal, or stalactitic masses lining and filling cavities in rocks. is usually greyish or creamy-yellow in colour, and has a characteristic waxy lustre with a certain degree of translucency. It possesses a minutely fibrous structure, the fibres being arranged perpendicularly to the concentric banding of the material. The optical character and orientation of the fibres is variable; and based on these differences French authors have distinguished several forms of silica under special names (quartzine, lutecite, lussatite, chalcedonite, and pseudochalcedonite). The physical characters of the mineral approach those of quartz, but with slightly lower values, sp.gr. 2.57-2.64; hardness  $6\frac{1}{2}-7$ ;  $n_{Na}$  1.5325, 1.5435. Analyses show, in addition to silica, the presence of variable amounts of impurities (iron oxides, alumina, &c.), and on the average about 1 p.c. of water. The water is probably present as adsorbed water held between the fibres; and its presence supports the idea that the mineral was deposited in a colloidal form as hydrated silica, and that it afterwards intominutely crystalline quartz Depending on the nature of the impurities present (and partly also on the state of aggregation of the material) the mineral may be of various colours, and on these differences are based several trivial varieties which are used as semiprecious stones. Carnelian owes its red colour

to ferric oxide, and sard its brown colour to hydrated ferric oxide; sard passes over into carnelian when ignited. The apple-green colour of chrysoprase is due to mckel silicate and the dark-green of plasma to an iron silicate. Heliotrope shows blood-red spots on a dark-green ground. Smoke-grey flint passes imperceptibly into more typical chalcedony In agate differently coloured layers are banded together in the same as those of agate (q,v) On the artificial colouring of chalcedony, v. AGATE.

L. J. S.

CHALCOCITE v COPPER-GLANCE

CHALCOPYRITE v. Copper-pyrites.

**CHALCOTRICHITE.** Native cuprous oxide (v. Cuprite).

CHALK. A white or greyish, loosely coherent kind of limestone rock, composed almost entirely of the calcareous remains of minute maiine organisms (foraminifera, coccoliths, &c) and fragments of shells. The purest kinds contain up to 99 p c of calcium carbonate in the form of the mineral calcite. Silica is always present in small amounts as the mineral opal, representing the remains of other minute marine organisms (radiolaria, &c ) and sponge spicules, and it is often segregated as the nodular masses of flint so commonly found embedded in chalk Minute grains of quartz, felspar, zircon, rutile, and other minerals are also often present With the admixture of clayey material there may be an insensible gradation from pure chalk to chalk-marl (v MARL). In phosphatic chalk, there is much calcium phosphate (up to 45 p c); in green qlauconitic chalk there is an admixture of grains of glauconite (a hydrated iron potassium silicate); and in red chalk there is some iron hydroxide Not only may there be considerable variations in the composition of chalk, but there may also be wide variations in the colour (snow-white, grey, &c), and texture of the material. It may be soft, incoherent, and porous, or quite hard and crystalline (as the chalk of the Yorkshire coast)

The following analyses of chalk are selected from those quoted by A J Jukes-Browne, The Cretaceous Rocks of Britain, Mem Geol Survey, 1903 (which see for details respecting the English chalk); I, Middle Chalk from Wye, Kent; II, Chalk Rock from Boxmoor, Hertfordshire; III, soft white Upper Chalk from

Farnha	m, S	Surrey	 		
	•		I.	II	Ш.
CaO			52 750	54 900	55 18
MgO			0.211	0.374	0 30
Fe <sub>3</sub> O <sub>3</sub>			0.067)	0 347	0 40
Al <sub>2</sub> O <sub>3</sub>			0 113 }	0 347	0.40
MnO			$0.149^{'}$		
K <sub>2</sub> O			0.098		0.22
Na <sub>2</sub> O			0 013		0.21
CO			37.670	42 101	42.57
$P_2\bar{O}_5$			0 106	0 713	0 08
$SO_{\mathbf{a}}$			0.082	0 465	0.09
Cl.			0 006		
S102 (8	olub	le)	0.035	0 128	
H <sub>2</sub> O Organi	0 m	· . }	5 400		
Insolul	ole	•	2.480	1.003	0.87
CaCO			99·180 85·61	100·031 97·001	99·92 95·76

150 CHALK.

The uses of chalk are numerous. For example, as a writing material in form of white and coloured crayons; for the manufacture of quicklime, mortar, Portland cement, plaster, &c., and as a fertiliser. Whiting is prepared by grinding chalk and collecting the finer sediments from water; this is used for polishing, making putty, and many other purposes. Under the name of 'Paris white,' chalk is used in the manufacture of india-rubber goods, oilcloth, wall-paper, &c. The harder kinds are extensively used as a building stone

Large quantities of chalk are quarried in the counties of Kent, Surrey, Sussex, Cambridge, Lincoln, and Hampshire; Kent being by far the most important. The production of English chalk is not far short of five million tons per annum. A certain amount of this is exported to the United States, though there are extensive beds of chalk in Kansas, Arkansas, and Texas

CHALK, FRENCH. Steatite or soapstone CHALKONES  $(\chi \alpha \lambda \chi \delta s)$ , an ore, particularly copper), the name proposed by v. Kostanecki and Tambor (Ber 1899, 32, 1923) for benzalacetophenone and its derivatives, on account of the fact that by suitable transformations they can be made to furnish reddish-yellow dyestuffs

In order to show the relationships of the various chalkones with one another, the following scheme of notation is adopted —

$$4' \underbrace{\sum_{5'=6'}^{3'-2'}}_{5'-6'} 1' - \text{CO-CH} = \text{CH} - 1 \underbrace{\sum_{6=5}^{2-3}}_{6-5} 4$$

Benzalacetophenone is obtained by the action of sodium hydroxide on an alcoholic solution of benzaldehyde and acetophenone at ordinary temperature For this purpose 21 grams benzaldehyde and 24 grams acetophenone are dissolved in 200 grams alcohol and 20 grams of a 10 pc solution of sodium hydroxide added The mixture is then allowed to stand for 24 hours, and the precipitate collected and recrystallised from alcohol. Benzalacetophenone is a yellow crystalline solid which dissolves in concentrated sulphuric acid with an intense yellow colour By heating 10 grams benzaldehyde with 30 grams acetophenone (dissolved in 100 grams alcohol) for a quarter of an hour at water bath temperature with 25 grams of a 40 p c solution of sodium hydroxide, benzaldiacetophenone  $\mathrm{C_6H_5\ CH(CH_2\ CO\ C_8H_5)_2}$  is obtained This melts at 85° and crystallises in colourless prisms. Its solution in concentrated sulphuric acid is colourless, but gradually shows a blue fluorescence, and on standing for one or two days becomes faint yellow in colour, showing a strong green fluorescence. Similarly by using a mixture of 21 grams benzaldehyde with 36 grams acetophenone in 200 grams alcohol and warming with 50 grams of a 40 p c solution of sodium hydroxide for 12-24 hours at 40°-50°, dibenzaltriacetophenone is formed It is a white crystalline solid, which melts at 198°. It dissolves slowly in concentrated sulphuric acid, the solution after some time developing a red colour and a cinnabar-red fluorescence, which slowly disappears By carrying out the preparation at boiling waterbath temperature, an isomeride melting at 256° is obtained (v. Kostanecki and Rossbach, Ber. 1896, 29, 1488).

Of the various derivatives, the 2'-hydroxy-chalkones possess the greatest importance, on account of their relationship to the naturally occurring dyestuffs of the flavone and flavanol groups, e.g. chrysine, luteoline, apigenine fisctin, &c  $(q\ v)$  The 2'-hydroxychalkones are obtained by

The 2'-hydroxychalkones are obtained by the condensation of o-hydroxyacetophenones with aromatic aldehydes in the presence of sodium hydroxide

2'-Hydroxychalkone

crystallises in yellow needles, melts at 88°-89° dissolves in alkali hydroxides with an orange colour, and in concentrated sulphuric acid with a yellow colour (v. Kostanecki and Tambor, Ber. 1898, 31, 715). It is converted into flavone by the action of alkali hydroxides on the dibromo-addition product of its acetyl derivative (v. Kostanecki and Feuerstein, *ibid* 1758)

OH
$$CO CH : CH C_{6}H_{5}$$

$$CO CH : CH C_{6}H_{5}$$

$$CO C B_{1} H$$

$$CO C B_{1} H$$

Numerous hydroxychalkones are described in the papers of v Kostanecki and his pupils, for which consult Ber. 1898, et seq

CHALMERSITE. A sulphide of copper and iron, CuFe<sub>2</sub>S<sub>3</sub>, first found as small orthorhombic crystals with pyrrhotite in the Morro Velho gold mine, Minas Geraes, Brazil Later it was found in considerable quantity in the Ellamar district and on Knight Island in Prince of Wales Sound, Alaska, where it has been mined as an ore of copper Here the material is massive and intimately intergrown with copper-pyrites. It is pale yellow in colour with metalle lustre and is strongly magnetic, sp gr. 468; hardness 3½ L. J S

CHALYBITE. Native iron carbonate, FcCO<sub>3</sub>, forming rhombohedral crystals isomorphous with calcite (CaCO<sub>3</sub>), and of importance as an ore of iron. It is often known as siderite, but this name had earlier been applied to two other mineral species, and it is also used as a group name for meteoric irons. Sp gr. 3'9; hardness, 3½ Small, brilliant crystals of varying habit are common in many mineral veins, for example, those of Cornwall. Veins of massive sparry material showing cleavage surfaces (Spathic iron-ore) are mined in the Brendon Hills in Somersetshire, in Weardale, Co. Durham, the Siegen district in Westphalia, dep. Isère in France, &c. The fresh, unaltered material is creamy white in colour ('white ore'), but when slightly weathered it is brown ('brown ore'). The Cleveland ore is a pisolitic form of chalybite occurring as beds in the Lias strata;

and the important clay-iron-stone and black-band iron-stone of the Coal-measures consist of compact chalybite, intermixed with clay and carbonaceous matter. A nodular form with an internal radiated structure is known as sphærosiderite.

Analysis I is of pure selected crystals from near Camborne, Cornwall (A. Hutchinson, Min Mag 1903, xiii, 209); II-IV of ores, by A. Dick, in Iron Ores of Great Britain, Mem. Geol. Survey, 1856-1862, II yellowish-grey spathose ore from Weardale; III greenish-grey earthy oblitic ore from Cleveland (also K<sub>4</sub>O 0 27 in the portion soluble in HCl); IV clay-iron-stone from Shelton colliery, Hanley, Staffordshire.

			I	II	III	IV
FeO			61.08	$49\ 47$	$39 \ 92$	46 35
$Fe_2O_3$					3 60	3 00
Al <sub>2</sub> O <sub>3</sub>				trace	7.86	0 30
MnO			1 12	242	0.95	1.61
CaO			0.10	347	7 44	1 93
MgO			0.13	3 15	382	2 24
$CO_2$			$38 \ 19$	37.71	22.85	32 46
$P_2O_5$				trace	1 86	0 67
$S_1O_2$				1 20	7 12	
$SO_3$				trace	trace	trace
$FeS_2$				0.08	0.11	0.12
$H_2O$					2.97	1.43
Organic	mat	ter		trace	trace	2 95
Insoluble	•	•		3 77	1 64	7 29
			100 62	101 27	100 41	100 38
Total Fe				38 56	33 62	38 29

L J. S

CHAMOMILE FLOWERS. Anthemis nobilis An examination of the flowers of Anthemis nobilis (Linn) by Power and Browning (Chem Soc Trans 1914, 105, 1833) has shown that these contain in addition to numerous other substances an angenin glucoside C<sub>21</sub>H<sub>20</sub>O<sub>10</sub>·2H<sub>2</sub>O, faintly yellow microscopic crystals melting at 178°-180°. It dissolves in alkalis with a yellow colour and gives with aqueous ferric chloride a purplish-brown colouration. Dried at 125°-130°

it loses one molecule of water of crystallisation, but the second molecule cannot be eliminated without decomposing the substance. This is evident from the composition of the hexacetyl derivative  $\rm C_{21}H_{14}O_{10}(COCH_3)_6$ , colourless microscopic crystals, mp  $\rm 144^\circ\!-\!146^\circ$ , the molecule of water in question being eliminated in the process of acetylation

By digestion with 5 pc aqueous sulphuric acid for three hours, this glucoside yields apigenin and dextrose according to the equation:

$$C_{21}H_{20}O_{10}, H_{2}O = C_{13}H_{10}O_{5} + C_{6}H_{12}O_{6}$$
  
A G. P

CHAMOMILE OIL v. OILS, ESSENTIAL CHAMOSITE, CHAMOISTE, v. THURINGITE. CHAMOTTE. A mixture of fire clay and burnt pottery used for making fire bricks, crucibles, pipes, &c

CHARCOAL v CARBON and FUEL

CHARICIC ACID CHAVICIN v Pepper resin, art Resins

CHARTREUSE. A famous liqueur formerly prepared at the head-quarters of the Carthusian monks near Grenoble, but now, since the retreat of the monks from France in 1906, made at Tarragona in Spain

The recipe for the original liqueur or 'elixir vita' as it was called, was given by the Marquess d'Estrées to the Carthusian monks of Paris in 1602, but, owing to the difficulty of procuring the necessary alpino plants, they presented it to their brethren of the Grande Chartrouse in 1735

The elixii vitæ was valued as a stimulant for the less robust members of the frateinity, weakened by age or by the privations and hardships entailed by the severity of their religious exercises, and was also dispensed to the feeble poor of the district

After numerous disasters by fire, pillage, and confiscation of their property through religious and political persecutions, culminating in the loss of everything during the French Revolution, the monastery of La Grande Chartreuse was restored in 1816, although in a very impovershed condition

Ingredients		Green	Yellow	White
China cinnamon .	1	15 grams	15 grams	125 grams
Mace		15 ,,	15 ,,	30 ,,
Lemon balm, dried		500 ,,	250 ,,	250 ,,
Hyssop (flower tops)	!	250 ,,	125 ,,	135 ,,
Peppermint (dried)	į	250 ,,	-	
Thyme		30 ,,		_
Balsime (Bal major) .	. !	125 ,,	_	
Genepi		250 ,,	125 ,,	125 ,,
Arnica (flowers) .	1	10 ,,	15 ,,	
Balsam poplar (buds)	.	15 ,,		
Angelica (seeds).	.	125 ,,	• 125 ,,	125 ,
Angelica (roots)	.	62.5 ,,	30 ,,	30 ,,
Coriander	•		1500 ,,	
Cloves	.	-	15 ,,	30 ,,
Aloes (Socotrine)		-	30 ",	
Cardamom (small) .			50 ",	30 ,,
Nutmegs				15 "
Calamus		*****		300 .,
Tonka beans		minimum		15 ,,
Alcohol at 85°		62 5 litres	42 5 litres	52 5 litres
White sugar	•	25.0 kilos	25 0 kilos.	37 5 kilos.

The secret of the manufacture of the 'elixir' had, however, been carefully preserved, and this was made the basis of other varieties of liqueur, by the sale of which the monks hoped to improve their finances. The result was the production of green, yellow, and white 'chartreuse,' the virtues of which were advertised far and wide by the military officers who had been quartered in the monastery about the middle of last century Extensive and modern plant was installed at Fourvoisie in 1863, and, in course of time, the sale of chartreuse increased the revenue of the monastery to a point it had never reached before.

After the passing of the French Religious Associations Act, in 1906, the monks of the Grande Chartreuse migrated across the border to Tarragona in Spain, carrying their secret formula with them

More than 200 ingredients are said to be used in the manufacture of genuine chartreuse, and its spirit basis is rectified grape spirit distilled from sound wine. The usual alcoholic strength of the three varieties of chartreuse is green, 55° of alcohol, or 96 p c of proof spirit; yellow, 42 5° alcohol, or 74 5 p c proof spirit; and white, 30° alcohol, or 52 5 p c of proof spirit

There are innumerable imutations of chartreuse, mostly prepared from plain spirit flavoured with essential oils and coloured by means of the various ingredients described under Cordials and Liqueurs. The formulæ given in the table on p 151 are amongst the best of those suggested for the preparation of imitation chartreuse.

Digest in the alcohol for 24 hours, distil until nearly all the spirit is over, and dilute to 100 litres. Colour green or yellow as described under Cordials and Liqueurs, add the sugar, allow to rest till clear, and filter (see Cordials and Liqueurs).

CHAULMOOGRA OIL GROUP. This group is remarkable on account of the considerable optical activity which these oils exhibit, owing to the presence of the strongly active chaulmoogric and hydnocarpic acids (discovered by Power and his collaborators, Chem Soc Trans. Hitherto, three oils belonging 1910, 97, 1285) to this class have been described, viz. Chaulmoogra oil, obtained from the seeds of Taraktogenos Kurzıı (King); Hydnocarpus oil, obtained from the seeds of Hydnocarpus Wightiana (Blume); and Lukrabo oil, the fat obtained from Hydnocarpus anthelminticus (Pierre). The saponification values of these oils lie between 204 and 213, and their iodine values between 90 The seeds of Taraktogenos contain a and 86 4 cyanogenetic glucoside Chaulmoogra is chiefly used in the treatment of leprosy and various skin diseases in Indo-China, Siam, and to some extent also in Europe A few years ago chaulmoogra oil was imported to the Continent as 'Cardamom oil' and used in margarine with disastrous results. Hydnocarpus oil, also known as kavatel oil, is occasionally sold as genuine chaulmoogra oil, but is inferior in its medicinal properties and contains a larger proportion of free fatty acids. The genuine oil from T. Kurzu is now official in the British Pharmacopæia, which describes it as a brownish-yellow soft fat, with a distinctive odour and acrid taste; sp gr. 0 940 at 45°; m.p 22°-30°; iodine value,

96-104; acid value, 21-27; and saponification value, 198-213. It is partially soluble in cold and almost completely soluble in hot 90 p c. alcohol. It is largely composed of the glycerides of acids of the chaulmoogric series, having the general formula  $C_nH_2-n_1O_2$ , and contains palmitin (Power, Amer. J. Pharm 1915, 87, 493) Gynocardia oil, sometimes confused with chaulmoogra oil, is optically inactive. For an account of the origin of the confusion in the nomenclature of the acids derived from chaulmoogra oil and the oils which have been considered as identical with it, see Pyman, Reports of Progress of Applied Chemistry, 1916, 1, 292.

CHAULMOOGRIC ACID  $C_{18}H_{32}O_2$ . An unsaturated acid found by Power and Gornall among the fatty acids contained in chaulmoogra oil from  $Taraktogenos\ Kurzii$  (King) Optically active  $[a]_p+56^\circ$ ; mp.  $68^\circ$ ; bp.  $247^\circ-248^\circ$  (corr)/20 mm Easily soluble in chloroform and ether Forms readily crystallisable salts with metallic bases, and esters with diphatic radicals (Chem Soc. Trans 1904, 838 and 851).

CHAY ROOT. Chay root or Chay-aver (from chaya = which fixes colours, and ver = root); also called Indian madder, is the root of Oldenlandia umbellata (Linn), Rubiacea. It bears the following Indian vernacular names: turbuli (Bengali); cheri-vello (Telugu); ché or chay, The plant is a small sayawer, imburel (Tamil) bush or herb found on sandy soils, chiefly near the sea-coast It occurs in North Burma and Ceylon, but is most abundant in certain tracts of the Madras Presidency, from Orissa south-It does not appear to be used in Bengal, but on the Malabar and Coromandel coasts, especially the latter, it is or was extensively cultivated, and employed in dycing a colour analogous to Turkey-red The chief market is Madras, where it was sold in small bundles at about 4d a lb The roots are usually about 10-12 ms long and 4 m thick, somewhat straight and stiff, tough and wiry, and with few or no lateral fibres When freshly gathered, they have an orange colour; but when dried and kept, they assume a yellowish-grey hue Boiling water gives merely a pale yellow extract, but if alkalı is added, a blood-red decoction is soon obtained The colouring principles seem to reside chiefly, if not entirely, in the bark of the root

The older literature connected with chay root is extremely meagre, and is for the most part contained in the following publications: Philosophy of Permanent Colours, 2282, 1813, Bancroft; Bulletin de la Société industrielle de Mulhouse, 5302, 1832, E Schwartz and D. Koechlin; L'Art de la Teinture des Laines, 475, 1849, Gonfreville. Quantities of the root were imported into Europe in 1774 and at later periods, and its dyeing properties were examined by the above-mentioned experts. It met with little practical application, however, as it was not found to possess any advantage over madder; indeed, it was considered to have but one-half or even a fourth of the colouring power of madder; further, it was found to contain certain undefined yellow substances of an acid character, which interfered somewhat with its dyeing power. Still, it was recognised as a good dyestuff, giving the usual madder colours, and equally fast to soap It is evident that the yellow substances referred to are not the same as those described in this article, as Schwartz states that they were present in larger quantity in 'nona' (Morinda citrifolia) than in chay root, whereas the yellow substances described later exist in much larger quantity in chay root than in morinda root. Very probably they refer to Rochleder's 'rubichloric acid,' as this is contained in morinda root in larger amount than in chay root. Schutzenberger, in his Traité des Matières colorantes, 2, 291, 1867, states that he found chay root to contain alizarin and chlorogenin (rubichloric acid) and that it was easy to exhaust the root by extraction with alcohol.

In many respects chay root resembles madder, as both contain ruberythric acid, alizarin, rubichloric acid, and cane sugar, but there are very marked differences in the nature of the other constituents of the two roots. Madder, as is well known, contains purpurin, purpurin- and purpuroxanthin-carboxylic acids, &c., and but traces of yellow crystalline substances, forming barium compounds soluble in water, which have not been fully investigated In chay root the former substances are entirely absent, but, on the other hand, although the root contains considerable quantities of yellow crystalline substances, they are quite distinct from those contained in madder

The phenolic constituents of this root have been studied by means of the sulphurous acid extraction method described in detail in connection with Madder, and also by a subsequent exhaustion of the residual-root with boiling lime-water. In this manner, chlororubine, alizarin, and a mixture of non-tinctorial yellow substances were isolated. These latter are present in chay root (probably as glucoside) in comparatively large amount (1 p c), and have been shown to consist principally, if not entirely, of the following substances.

Alizarin a-methylether crystallises from dilute methyl alcohol in long orange-yellow needles, melting at 178°-179°. It has the constitution—

This compound may be synthesised by the action of an ethereal solution of diazomethane on a nitrobenzene solution of monoacetylalizarin (Oosch and Perkin, Chem Soc. Proc 1914, 30), subsequently removing the acetyl group Owing to the fact that the so-called monoacetylalizarin is a mixture of the two monoacetylalizarins, both  $\alpha$ - and  $\beta$ -alizarin methylethers are thus simultaneously produced. Addition of alcoholic potast to a solution of the  $\beta$ -methylether as potassium salt, the  $\alpha$ -methylether thus remaining in solution

The methoxyl group present in this substance is much more readily hydrolysed than is usually the case, for prolonged digestion with boiling baryta water is sufficient for this purpose, a precipitate consisting of barnum alizarate thus separating. This property, therefore, accounts for the difficulty in obtaining either this compound or alizarin dimethylether by means of methyl iodide, for in this process of methylation a prolonged digestion in the presence of free alkali is necessary. On the other hand, whereas

alizarin is only converted with considerable difficulty into alizarin dimethylether, by the action of methyl sulphate, alizarin  $\alpha$ -methylether readily yields this substance when treated with this reagent in the usual manner.

Acetylalizarın a-methylether crystallises in

yellow needles, melting at 212°
Anthragallol dimethylether (A) forms yellow needles, and dissolves in alkaline solutions with a red colouration. It is distinguished from the isomeric compound described below, in that it yields an ammonium salt, crystallising in searlet prisms, which is sparingly soluble in cold alcohol. Heated with sulphuric acid to 180°, or with hydrochloric acid, it is converted into anthragallol, and when methylated by means of methyl sulphate, is readily converted into anthragallol trimethylether. By the action of 10 pc potassium hydroxide in a sealed tube at 180° for 5 hours, it is partially converted into methoxyalizarin:

and a similar reaction occurs when sulphuric acid at 100° is employed. The constitution assigned to this anthragallol dimethylether (A), is therefore as follows—

Anthragallol dimethylether (B) crystallises in yellow leaflets, melting at 230°-232°, and gives a red ammonium salt which is soluble in cold alcohol Heated with 10 pc potassium hydroxide solution to 180° for 5 hours, it is partially converted into a compound having the reactions of methoxypurpuroxanthm

and most probably possesses the following constitution.—

By methylation with methyl sulphate in the usual manner, this compound is readily converted into anthragallol trimethylether

Acetylanthragallol dimethylether (B) consists of long yellow needles, melting at 176°-178°

Hystazarine monomethylether obtained as long prange-yellow needles, melting at 232°, when heated with hydrochloric acid at 180°, is converted into hystazarine. Alkaline solutions dissolve it with a crimson colouration, and its ammonium and potassium salts, which crystallise in garnet-red needles, are insoluble in sold 1800 butylic alcohol. The constitution of hystazarine monomethylether is as follows:—

hystazarine dimethylether, which consists of yellow glistening needles, melting at 235°-236°.

pale - yellow Metahydroxyanthraquinone, needles, melting at 302°, is identical with the well-known artificial compound prepared from anthraquinone-\(\beta\)-monosulphonic acid by digestion with alkalı.

Dyeing Properties -Although chay root contains acid principles which tend to dissolve the mordants, its employment as a dyestuff presents no difficulty. The only precautions presents no difficulty necessary to be observed are to add 2 pc. of chalk to the dye-bath, and to raise the tempera-

ture gradually to the boiling-point

Dyeing experiments on ordinary stripeprinted calico, containing aluminia and iron mordants, have shown that the dyeing power of chay root is equivalent to the presence of a percentage of 0 33-0 35 alizarin Compared with ground madder root of good quality, it seems to have about half its dyeing power when the comparison is made before soaping, but after soaping it appears to be quite equal to madder The reds, pinks, and chocolates have a distinctly bluer shade than those given by madder, and the blacs are much fuller and brighter and very similar to those obtained from alizarin last feature alone ought to have secured a ready market for chay root among the European dyers, previous to the introduction of artificial alizarin, and it is somewhat strange that its marked suitability for lilacs should have escaped the observation of those who formerly made dyeing experiments with this root

On oil-prepared calico, mordanted with alumina, chay root gives an excellent blue shade of Turkey-red, withstanding the operation of clearing with soap and stannous chloride better even than a madder-dyed red, and quite equal to one obtained by means of artificial alizarin Good brown, red, orange, and purple colours are readily obtained on wool, and also on silk, suitably mordanted with chromium, aluminium, tin, and iron, according to the ordinary method usual with dyers. On wool, the colours not being submitted to any soaping operation, chay root appears to possess about

half the dyeing power of madder Boiled with dilute sulphuric acid, chay root yields a 'garancine' of a very dark-green colour and possessing about three times the dyeing

power of the original root

Literature.—A. G Perkin and J. J Hummel, Chem Soc. Trans 1893, 63, 1160, ibid 1895, 817; A.G. Perkin, ibid 1907, 91, 2066, J. J. Hummel and A G. Perkin, J Soc Chem Ind. 1894, 13, 346

CHEDDITE. An explosive consisting of a mixture of ammonium perchlorate, dinitroto-luene and castor oil (" Explosives)

CHEESE. A solid preparation of the chief constituents of milk, which, by the action of certain enzymes, are made digestible and palatable.

The action of enzymes is to a large extent influenced by the reaction or hydrogen ion concentration of the curd, which in turn is brought about by the extent to which the lactic organisms are controlled or encouraged in the process of manufacture

Cheeses may be made from the milk of

By means of methyl sulphate, it yields | various animals, but they are usually prepared from cows' milk, which is either used in its natural condition, or may be first divided into cream and separated milk, from both of which products cheeses are also made

In the case of cream and cottage cheeses, the lactic acid produced by the lactic ferments may be used for coagulating purposes; but in the majority of cheeses, rennet is used to precipitate the cascin The caseinogen of the milk is converted into casein by the rennet ferment, and this, in the presence of the lime salts of the milk, is followed by the separation of insoluble calcium caseinate or curd, which carries with it nearly all the fat, some of the other constituents of the milk, and the enzymes of the rennet and milk. This curd differs from that produced by acids, which throw out caseinogen from the combinations with lime in which it occurs in milk

The condition of the rennet curd, used for making different cheeses, is controlled by the acidity of the milk and by the temperature of renneting If milk is curdled with rennet without the previous development of acidity, a soft curd results, this curd will not shrink so much as a curd produced from an acid milk. Too much acidity, on the other hand, results in a hard curd, which will shrink to a much greater

extent, giving a hard 'acid' curd

Temperature also exerts an important influence in cheese making; the action of the rennet is hastened up to about 100°F, and ceases at about 130°F. The temperature employed, the degree of acidity, and the quantity of rennet used, will vary with different cheeses and also with the nature of the milk, the ash constituents probably exerting a considerable influence The acidity is often controlled by the addition of 'starters,' i.e. pure cultures of suitable lactic ferments or of mixed cultures in milk or whey, to the milk The average temperature used is about 84°F., at which the coagulation takes some 40 to 60 minutes some cheeses, e g Leicester, Dutch cheeses, &c, annatto is added as colouring matter at least 10 minutes before the rennet. Salt is also added in the case of hard pressed cheese, when the curd has been ground prior to being put in the mould for pressing into shape, in the case of soft cheese, about the time it is fit to take out of the mould.

With hard-pressed cheeses such as Cheddar. Cheshire, Derby, Leicester, Gloucester, Ementhaler, Gruyère, and Edam, the curd is cut up by means of curd knives, it is then stirred and heated to acquire the correct stage of dryness, consistency, and acidity, after which the whey is drawn off and the curd allowed to remain till a certain degree of acidity has developed. This is determined by means of a hot iron against which the curd is pressed, the length of the threads which are drawn out on pulling away the curd being taken as an index of the acidity produced; the percentage of acidity at the various stages of manufacture may accurately be determined by means of the acidimeter. The curd is then ground in a mill, and afterwards pressed in a mould into a shape which varies with the different cheeses. After being pressed for 2 or 3 days, the cheese is ripened in a curing room, the humidity and temperature of which should be under control;

CHEESE. 155

a temperature of about 55° to 65°F. being most generally desirable.

To another group of cheeses, in which the process of manufacture varies considerably from the above, belong the moulded cheeses, in the making of which the pressing process is omitted, or in the case of Wensleydale greatly reduced; the resulting cheese being softer and admitting of the growth of blue veins of mould (Penicillium glaucum) in the cracks left between the pieces of curd which extend throughout the cheese. To this group belong Stilton, Roquefort, Gorgonzola, Wensleydale, &c

In the case of the small 'soft' cheeses of the Camembert type, considerable variations are made in the process of manufacture, the resulting cheese being quickly ripened by moulds, or rather by the enzymes produced by moulds, which grow on the outside of the cheeses. Penuellium candidum and Oidium Lactis play an important part in the ripening of Camembert.

Indole is present in Limburger and Camembert cheeses and in Handkase Limburger cheese also contains phenol Cheddar, Swiss, 'Gammalost,' Wick, and Roquefort cheeses do not contain indole or phenol. Scatole is absent in all cases The amount of indole in Limburger cheese increases during the ripening process until it reaches as much as 1 part in 52,800 parts of cheese (Nelson, J. Biol. Chem. 1916, 24, 533) All cheeses contain p-hydroxyphenylethylamine, which is produced by the action of micro-organisms on tyrosine; the amine is a normal constituent of cheese, and is produced before actual decomposition of the latter takes place The organisms which produce the substance are probably identical with the Baccasei-a-Freudenreich (Ehrlich and Lange, Biochem Zeitsch. 1914, 63, 156).

Tryptophane has also been found in ripe Stilton cheese

The chemical analysis of cheese includes the determinations of the water, fat, total nitrogen, ash, and salt. The products of ripening are also sometimes determined, and the fat may be examined for adulteration with animal fats

To determine the water, ash, and salt, 2 or 3 grams of well-sampled cheese, cut into small pieces, are weighed into a flat-bottomed dish and placed on the water-bath at 100°C, the dish being inclined so that the fat runs off the drying curd. Weighings are made every hour, and the water is found to have all been driven off in about 5 or 6 hours

The melted fat is then poured off and the residue well macerated in hot amyl alcohol; the residue may then be burnt and the ash weighed, the salt being afterwards dissolved out with water and the solution titrated with N/10 silver

The water may be determined more rapidly by heating from 8 to 12 grams of the cheese with petroleum in a distilling flask connected with a condenser. In from 30 to 45 minutes the whole of the water will have been expelled and may be measured in a graduated tube (Analyst, 1913, 146)

According to Evans, Hastings, and Hart (J. Agric. Research, 1914, 2, 167), the organisms present in Cheddar cheese in such numbers as to indicate that they must have some function in the ripening process, are included in the four

groups: B lactis acidi, B. casei, Streptococcus, and Micrococcus; each of these groups may be divided into a number of varieties according to their fermentative powers' The B. cases group is apparently responsible for the pungent taste which develops late in the ripening period The action of two or more organisms growing together is not the sum of their individual When growing actions when growing alone together, they may attack substances that neither can attack alone, or they may produce a larger quantity of acid than the sum of the quantities that either can produce alone. volatile acids formed in cheese result from the action of Streptococci and Micrococci on the citric acid, lactose, or protein present the varieties of Streptococcus b forms comparatively large quantities of alcohols and esters which contribate to the flavour of the cheese. The B. cases group of organisms form propionic acid, acetic acid, and lactic acid. Certain members of the B casei group form llactic acid, and others form d-lactic acid, a mixture of these organisms produces racemic lactic acid The racemic lactic acid found in cheese during the curing process may also be produced to a small extent by enzymic acid, but it is more probably due to the combined action of B lactis acidi and the organisms of the B. casei group (J Soc Chem Ind 1915, 34, 97)

To obtain a typical Cheddar cheese from pasteurised milk, the pasteurising temperature should not exceed 88° (190°F) when the milk is heated rapidly, or 77° (170°F) when the time of heating is from 15 to 30 minutes Heating to at least 77° (170°) is needed to reduce the number of organisms; at 93° (200°F) practically all the organisms are destroyed Cheeses pre-pared from 'heated' milk contained. water, 33.61-36.22; fat, 32.36-34.53; insoluble nitrogen (probably unaltered casein), 2:49-2:91 pc (Benson, J. Board Agric 1915, 21, 878)

Ether may be used to extract the fat, and the dried curd and fat afterwards weighed, but a better method is to take a fresh portion of 3 or 4 grams of the cheese and grind with twice its weight of anhydrous copper sulphate in a small mortar, afterwards extracting with ether in a Soxhlet extractor

Modifications of the Werner-Schmid or Gottlieb-Rose methods also give good results.

The total nitrogen is determined by the Kjeldahl method in about I gram of cheese

According to Richmond, the products of ripening may be determined in the following manner: 10 grams of the cheese are placed in a small mortar and ground up with 25 cc of boiling water, the liquid is then decanted through a filter into a 250 c c flask, the process being repeated with nine further similar measures of boiling distilled water The total filtrates are then cooled, made up to the 250 cc. mark, and well mixed The products of ripening are determined by the evaporation in a dish of an aliquot part (50 c.c); this, after drying till constant at 100°C, is burnt, the weight of the dish and ash being subtracted from the weight of the dish and dry residue. The difference between 100 and the sum of the water, fat, ash, and products of ripening, may be taken as unaltered casein.

The primary products of ripening may be

156 CHEESE.

determined in another 50 c.c. of the filtrate by the Ritthausen method as follows: 5 c.c. of copper sulphate solution are added and the solution neutralised with caustic soda solution; the precipitate which forms is collected in a weighed Gooch crucible, it is then washed well and dried in a water-oven, the fat is then extracted with ether, and the residue dried till constant. The crucible and its contents are then ignited and weighed, the difference giving the weight of the primary products of ripening, i.e. soluble proteins, &c

To examine the fat, about 50 grams are dried till the fat runs out: the remaining fat is then extracted from the residue with ether, the ether evaporated and the fat examined, as in the case of butter fat. The soluble and insoluble acids and the Reichert-Wollny figure being generally determined, the addition of animal fats can be thus detected. The removal of fat from milk used in making cheese can be assumed if the fat is less than 45 pc of the dried cheese or less than six times the total nitrogen.

COMPOSITION, OF CHEESE (BELL).

		100 p	arts con	tain		D	Propor-	a u		tage com-
Description	Water	Fat	Casein Free or acid as genous matter		Ash	Propor- tion of fat in 100 parts of dry cheese	tion of fat in 100 parts of casein and fat	Salt per cent in cheese	-	Insoluble acids
Stilton American (red) American (pale) Roquefort Gorgonzola Cheddar (medium) Gruyère Cheshire Single Gloster Dutch	23 57 28 63 31·55 32·26 31 85 35 60 33 66 37 11 35 75 41 30	30 69	29 64 28 83 27 16 27 88 28 16 30 67 26 93 31 10	0 27 1 32 1 35 0 45 0 27	3 51 4 49 3 42 4 88 4 58 4 22 4 71 4 42 4 49 7 10	51 19 53 57 52 49 50 75 20 39 49 02 46 26 48 78 44 12 38 80	52 50 52 12 53 34 54 24 53 08 50 49 47 07 50 84 45 24 42 41	0 67 0 72 0 82 3 04 2 11 1 43 0 81 1 69 1 28 4 45	4 42 4 26 4 81 4 91 4 40 4 55 4 41 5 55 6 68 5 84	88 96 89 66 88 49 88 70 89 18 88 75 88 97 87 76 86 89 87 58

The ripening process in cheese, whereby it usually acquires a more agreeable flavour, is essentially a fermentation of slow character, which is brought about by micro-organisms which are generally present in the milk or in the air of the dairy. They can only be controlled, to a limited extent, by the process of making, temperature, &c It is not possible to destroy undesirable organisms by sterilisation, for heated milk will not make typical cheese as far as is at present known, though heating to 150°F is now practised for soft cheeses. Cleanliness, from the moment the milk leaves the cow, is of the utmost importance

The organisms which (ause most trouble to the cheese maker are those which produce gas, which, when it develops in the curd, produces what is known as heaving or blown cheeses These organisms, to which Bacillus coli communis belongs, often come from manure. Their munis belongs, often come from manure. absence can be demonstrated by what is known as a fermentation test, which consists in placing about 10 c.c of the milk in a sterile test-tube and keeping it at blood heat for about 18 hours, after which time no signs of bubbles of gas should be seen in the tube

Bitterness in cheese is sometimes caused by bacteria, as is the case when the milk is infected with Micrococcus casei amari or Tyrothrix geniculatus. An interesting example of a bitter cheese organism is that of Torula amara, which, after causing much trouble and loss in Canada, was eventually traced to the leaves of the sugar maple, from which source it infected the clean chaurns which were stood under the trees after be ng cleaned.

produced by two groups of bacteria acting upon caseinogen and certain amino acids

Black spots in cheese may be caused by moulds or by particles of iron which get into the milk and are afterwards changed to sulphide by the sulphuretted hydrogen produced in ripening Other organisms which produce colours (moulds, &c), may get into the milk or may be derived from the air or utensils of the dairy at later stages of the manufacture.

The cheese magget is the larva of the fly Piophila case: There are four species of cheese mites. (1) Carpoglyphus anonymus (the Cheddar mite); (2) Tyroglyphus siro; (3) Tyroglyphus longior; (4) Aleurobius farinæ (2, 3, and 4 are Stilton and Cheddar mites) (N. B. Eales, 1917)

It must be remembered that micro-organisms and the enzymes they produce play a most important part in ripening; how far this process is also dependent on the proteolytic enzymes of the rennet and to the natural enzymes of the milk is not yet fully ascertained. As above mentioned, pure cultures of lactic acid organisms are sometimes added to milk to be used for hard-cheese making, and Camembert is often infected with the appropriate mould. Gorgonzola is also sometimes infected with mould to make the desired blue veins. As a general rule, however, the dairy and district where a particular kind of cheese is made contain the organisms which are necessary for the successful manufacture of the cheese in question.

The natural rind on Gorgonzola cheese amounts to from 2 to 4 p.c. of the weight, but ng cleaned.

In artificially coated cheese it may amount to Discolouration of Stilton cheese may be from 16 to 27 p.c. of the whole weight. This artificial coating consists of a mixture of barytes | stitute for butter has led manufacturers, and tallow coloured on the outer side with oxide of iron The importation of coated cheese into France is prohibited (Hinks, Analyst, 1911, 36, 61).

Oleomargarine and Lard Cheese. The success attending the sale of margarine as a sub- seed oil.

especially in America, to introduce foreign fats of animal or vegetable origin in the production of cheese as a means of utilising skim-milk; the cream which has been abstracted being re placed by lard, oleomargarine, and by cotton-

Analyses of Oleomargarine and Lard Cheeses (Bell).

	100	parts of ch	eese contain	1	Dan	100 parts of	Melting-		
	Water	Fat	Casein and free acids	Ash	Per cent. of salt	Insoluble fatty acids	Soluble fatty acids	point of fat	
Oleomargarine Lard .	30 95 31 30	28 80 24 66	36 27 38 87	3·98 5 17	1 14 1 55	92 43 92 88	2 16 1 55	77°F. 92°F.	

In flavour and character these samples closely resembled milk-fat cheese. Comparison of the results of the analysis of the two samples with those yielded by genuine cheeses in the former table conclusively proves the admixture with foreign fat. The low percentage of soluble acids indicates that more than one-half of the fat has not been derived from milk. J. G.

CHEIRININE. See under CHEIROLINE

CHEIROLINE C<sub>5</sub>H<sub>9</sub>O<sub>2</sub>NS<sub>2</sub>, colourless prisms, m.p. 47°-48°, optically mactive, occurs in wallflower seeds (Cheiranthus Cheiri [Linn.]), 1.6-1.7 pc, and in Erysimum species has the constitution CH<sub>3</sub> SO<sub>2</sub> CH<sub>2</sub> CH<sub>2</sub> CH<sub>2</sub> NCS and has been synthesised (Schneider, Annalen, 1910, 375, 207). It is present as a sinigrin-like glucoside, which is hydrolysed by myrosin from white mustard seeds. Wallflower seeds contain an enzyme capable of liberating mustard oil from (myrosin-free) black mustard seeds (Schneider and Lohmann, Ber. 1912, 45, 2954). Cherroline has no basic properties, and is said to be antipyretic (Wagner, Chem. Zeit. 1908, 32, 76). From wallflower leaves or seeds Reeb has obtained cheirinine C1.4H35O17N3, which resembles quinine in physiological action (Arch. expt. Path. Pharm. 1898, 41, 302; 1899, 43, 130) and a glucoside cheiranthin

CHELERYTHRINE (121H17O4N occurs in various Papaveraceous plants, e.g. Chelidonium majus (Linn.), Glaucium flavum (Crantz.), Sanguinaria canadensis (Linn) (Probst, Annalen, 1839, 29, 120, 250), Bocconia spp. Forms colourless rhombohedra with \$\frac{1}{2}H\_2O \text{ or } 1C\_2H\_5OH, m p 203°, [α]<sub>D</sub>=0°. Salts intensely yellow and well crystallised. The free base is colourless. It contains a carbonyl group, and if this is altered the basic properties are lost. Perhaps it forms oxonium salts (Karrer, Ber 1917, 50, 212). Contains two methoxyl groups, and may be a methyl ether of sanguinarine (q.v.). Narcotic (Konig and Tietz, Arch. Pharm. 1893, 231, 145, 161; Murrill and Schlotterbeck, Ber. 1900, 33, 2802; Fischer, Arch. Pharm. 1901, 239, 409, 426, 438)

CHELIDONINE C20H19O5N, H2O occurs with protopine, chelerythrine, sanguinarine, and  $\alpha$ -,  $\beta$ -, and  $\gamma$ -homochelidonines in the root of the common celandine (Chelidonium majus [Linn.]), and with one or more of the same alkaloids in Stylophorum diphyllum (Nutt.) (Selle, Arch. Pharm. 1890, 228, 96), Adlumia-cirrhosa (Rafin.)

(see under Adlumine), and possibly in Sanguinaria canadensis (Linn.) (Konig and Tietz, Arch. Pharm. 1893, 231, 145, 161) It crystallises in monoclinic tablets, m p.  $135^{\circ}-136^{\circ}$ ,  $[a]_{0}+115^{\circ}$ Yields a benzoyl derivative, m p. 217°, and when heated with zinc-dust furnishes ammonia and a substance with a phenanthrene-like odour (Schlotterbeck and Watkins, Pharm. Arch. 1903, 6, 141). Salts crystallise well (Wintgen, *ibid*. 1901, 239, 438). It is slightly toxic, resembling morphine in its action

CHEMICAL AFFINITY. There has been much fruitless speculation about chemical affinity, about the nature of the force which impels two different substances to react with one another During the last half-century, however, the efforts of physical chemists have been devoted rather to the practical investigation of the way in which chemical affinity manifests itself, and of the extent to which the operation of this force is modified by altering the external conditions Quantitative methods have been devised for studying the influence of concentration, of temperature and pressure, and of foreign substances, on the course and products of a reaction The object of this article is to give an outline of the methods employed and the results obtained in this study of the operation The three chief divisions of chemical affinity into which the discussion of the subject naturally falls are (1) velocity of chemical reactions; (2) chemical equilibrium; (3) catalysis.

Velocity of chemical reactions. It is well known that different chemical reactions proceed to completion at very different rates. One change may be so rapid as to be practically instantaneous, whilst in another case the reacting substances may appear almost indifferent to each other. The rate of chemical change depends, however, to a marked extent on the temperature, so that the velocity of a reaction which takes place with extreme rapidity is moderated on lowering the temperature, and becomes amenable to experimental investigation. Suppose now that for such a reaction as that represented by the equation A+B=C+D, a range of temperature has been found over which the change proceeds at a measurable rate; the question next arises, How is the velocity of the change, at a given temperature, dependent on the quantities of the reacting substances A and B? In dealing with this question, we shall assume that the reaction system is homogeneous, and that the change proceeds completely from left to right, i.e. until

either A or B has disappeared

The reaction system being homogeneous, it is permissible to regard matters from the molecular-kinetic point of view, and to conceive the interaction of the substances A and B as taking place only in so far as the molecules of A come in contact with the molecules of B The velocity of the reaction, therefore—that is, the rate at which A and B disappear—will be proportional to the frequency of the collisions between a molecule of A and a molecule of B, even although only a certain proportion of the collisions is followed by chemical interaction Now, on kinetic grounds, the frequency of the collisions between molecules of A and B is proportional to the product of their concentrations, hence the velocity of reaction between A and B is proportional to the product of their concentrations (or their 'active masses,' as it is sometimes put) This proposition is nothing else than the law of mass action, applied to the chemical change under consideration.

The foregoing proposition may be formulated m more definite shape If a is the concentration of the substance A, and b is the concentration of the substance B at the beginning of the reaction, then the initial velocity of the change,  $V_0$ , is given by the equation  $V_0 = k_1$  ab, where  $k_1$  is a proportionality factor, the numerical value of which is independent of the concentrations of the reacting substances, and varies only with temperature and other external conditions If, now, the reaction is allowed to proceed for an interval of time t, and if x represents the extent to which the concentrations of A and B have been diminished by the progress of the change, then the velocity at time t, Vt, is given by the equation  $V_1 = k_1 (a-x)(b-x)$  It is obvious that  $V_t$  must be less than  $V_0$ , and that the velocity of the reaction must diminish continuously from the start onwards . Vt, therefore, represents the velocity of the reaction only for an infinitely short space of time This velocity is defined by the rate at which x is, at that moment, increasing with the time, and if, using the language of the differential calculus, we indicate by dx the extent to which x increases in the infinitely short space of time dt, we may

write 
$$V_{\bullet} = \frac{dx}{dt} = k_1 (a-x)(b-x)$$
.

The applicability of the law of mass action, as expressed in the foregoing formula, is best appreciated by reference to a particular case, such as the inversion of sucrose. Although this change, which may be represented by the equation  $C_{12}H_{12}O_{11}+H_2O=2C_6H_{12}O_6$ , takes place with appreciable velocity only in the presence of a catalytic agent such as hydrochloric acid, yet the latter is found unaltered when the reaction is over, and the inversion of sucrose may be regarded as a particular case of the reaction A+B=C+D. The fact that the inversion is carried out in aqueous solution permits, however, a simplification of the formula  $\frac{dx}{dt}=k_1$  (a-x)(b-x); the water taking part in the reaction is a small fraction of the total water present, so that x may be neglected in comparison

with b, and we may write  $\frac{dx}{dt} = k(a-x)$ , where  $k = k_1 \cdot b$  Integration of this equation leads to the formula  $k = \frac{1}{t} \cdot \log_e \frac{a}{a-x}$ , in which, as already indicated, a is the initial concentration of sucrose, and a-x is its concentration after the inversion has proceeded for t units of time

In order to test the validity of the formula  $k = \frac{1}{t} \log_{e} \frac{a}{a - x}$  as applied to the inversion of sucrose, it is only necessary to ascertain the amount of sucrose still present in the reaction mixture at intervals during the progress of the inversion, and then to see whether the amount of sucrose varies with the time in the manner required by the formula The simplest way of following the course of sugar inversion is to use The reaction mixture, that is, the polarimeter sucrose solution containing a catalytic agent such as hydrochloric acid, is put in the polarimeter tube, which should be provided with a water-jacket in order to secure a constant temperature during the progress of the inversion The angle of rotation is determined at the beginning of the change, at subsequent intervals, and finally when the change is complete. If we indicate by  $\alpha_0$ ,  $\alpha$ , and  $\alpha_{\infty}$ , the initial angle of iotation, the angle at time t from the start, and the final angle respectively, then  $a_0 - a_{\infty}$  is a measure of a, the total sucrose undergoing inversion, and  $\alpha - \alpha_{\infty}$  is similarly a measure of a-x, the sucrose which has still to undergo inversion after time t. Hence  $\frac{a}{a-x} = \frac{a_0 - a_x}{a - a_x}$ and the formula which should represent the and the formula which blocks  $k=\frac{1}{t}\log_e \frac{a_0-a_\infty}{a-a_\infty}$ From the ascertained values of l,  $a_0$ , a, and  $a_{\infty}$ , it is easy to discover whether the expression on the right-hand side of the equation is constant or not The experimental data recorded in the following table show how far in a particular case the actual course of sucrose inversion corresponds with the velocity formula .-

Inversion of Sucrose at 25° by N/2 HCl.

in minutes	Angle of rotation	$k=\frac{1}{t}$	$\log_{10} \frac{a_0 - a_{\infty}}{a - a_{\infty}}$
0	$+25\ 16^{\circ}$		
56	16 95°		0 00218
116	10 38°		0 00218
176	5 46°		0 00219
236	1 85°		0 00219
371	$-3.28^{\circ}$		0 00221
œ	-8.38°		

The expression evaluated in the last column is  $\frac{1}{t}\log_1\frac{a_0-a_\infty}{\alpha-a_\infty}$  mistead of  $\frac{1}{t}\cdot\log_2\frac{a_0-a_\infty}{\alpha-a_\infty}$ , but, obviously, if the value of the former expression is constant, the value of the latter must be so also. The figures in the last column are satisfactorily constant, and the figure 0 00219 may therefore be taken as the velocity coefficient for the inversion of sucrose under the specified conditions, viz at 25° and in presence of N/2 HCl. The variation of the velocity coefficient with temperature and with the concentration of the acid will be discussed later.

Reactions, in which only one molecule of a single substance undergoes change, are termed unimolecular reactions, or reactions of the first order The course of all such reactions, apart from possible disturbing factors, is represented by the formula  $k = \frac{1}{t} \log_e \frac{a}{a - x}$ Examples of strictly unimolecular changes, to which the foregoing formula has been found applicable, are the transformation of acetochloranilide into p-chloracetanilide, and Beckmann rearrangement, and the decomposition of nickel carbonyl. The inversion of sucrose is, strictly speaking, not a unimolecular but a bimolecular reaction, for both water molecules and sugar molecules are involved in the change The reaction takes place, however, in aqueous solution, and, as already indicated, the consequence is that the course of the inversion conforms to that characteristic of a strictly unimolecular change In many other similar cases, a bimolecular reaction proceeds in accordance with the formula  $k = \frac{1}{t} \log_{e_{\alpha}} \frac{a}{-x}$ Examples of this are the hydrolysis of methyl acetate and the

It is desuable here to emphasise certain features which are characteristic of unimolecular reactions, features which can indeed be read out of the velocity formula, but are most clearly seen by the consideration of experimental data In the first place, for all unimolecular reactions the amount of change occurring in a given interval of time is a constant fraction of the unchanged substance still present This is a necessary consequence of the fundamental velocity equation, as is clear when it is written in the form  $\frac{dx}{a-x} = kdt$ , but the proposition is also in harmony with observation, as will be seen by a consideration of the following data bearing on the inversion of sucrose (Armstrong and Caldwell, Proc Roy. Soc. A 1905, 74, 199) A sucrose solution was inverted at 20° with hydrochloric acid, and the angle of rotation (a) was determined from time to time, with the results shown below

decomposition of diazonium salts

Time	$+21^{\circ}55^{\circ}$
15	20 40°
120	13 75°
135	12 95°
225	8 62°
240	8 02°
οc	−7·18°

The decrease in rotation during the first 15 minutes, viz.  $1^{1}15^{\circ}$ , is a measure of the amount of change during that interval. The average rotation of the solution over this interval may be taken as  $\frac{21^{\circ}55^{\circ}+20}{2}=20\,97^{\circ}$ , and a measure of the unchanged sucrose present is given by  $20\,97+7\cdot18=28\,1b$ . The ratio of the amount of change occurring in the first 15 minutes to the amount of unchanged sucrose present is therefore  $\frac{115}{28\cdot15}=0\,041$ . If now the intervals from 120 to 135 minutes, and from 225 to 240 minutes are similarly considered, it will, be found that the ratio of the amount of change occurring in the 15-minute interval to

the amount of unchanged sucrose present is 0 039 in each case, practically the same value as for the first 15 minutes of the inversion. The experimental data, therefore, bear out the statement that for a unimolecular reaction the amount of change occurring in a given interval of time is a constant fraction of the unchanged substance still present

Another feature of unimolecular reactions is that the amount of substance changed in a given time is always the same fiaction of the total amount of substance present at the beginning of the reaction, however this latter amount be varied. An illustration of this is furnished by the following data, referring to the inversion of sucrose in very dilute aqueous solution under the influence of invertase.

Grams sucrose per 100 c c.	Grams sucrose inverted in 60 mins
1 00	0 249
0.50	0.129
0.95	0.060

This experimental result is implied in the formula for a unimolecular reaction, for if  $y = \frac{x}{a}$  is the fraction of the total sucrose which has been inverted up to time t, the formula may be written  $k = \frac{1}{t} \log_e \frac{1}{1-y}$ , and it is plain that the value of y depends merely on t, and is independent of a

For bimolecular reactions, that is, reactions in which two molecules are concerned, the fundamental velocity equation is  $\frac{dx}{dt} = k(a-x)(b-x),$  as indicated in the argument above. Integration of this equation leads to the formula  $k = \frac{1}{(a-b)t} \log_e \frac{b(a-x)}{a(b-)x}$  If, however, the two reacting substances have the same initial concentration, that is, if a=b, the integrated formula is  $k = \frac{1}{t} \frac{x}{a(a-x)}$ .

A typical case of a bimolecular reaction, or reaction of the second order, is the saponification of an ester by an alkali. Under suitable conditions of temperature and concentration, the velocity of saponification is measurable, and the progress of the reaction can be determined by extracting samples of the reaction mixture from time to time and titrating for free alkali. The figures in the following table, referring to the saponification of ethyl acetate by sodium hydroxide at 24 7°, will serve to show how far the experimental data are in harmony with the formula just recorded The reaction mixture in this case was N/40 in relation both to ester and alkalı, and the figures under a-x in the table are the volumes of a standard acid solution required to neutralise 10 c.c. of the reaction mixture -

11 c		
<i>t</i> (min )	a-x	$ka = \frac{1}{t} \frac{x}{a-x}$
0	8.04	
4	5 30	0 129
6	4 58	0 126
8	3 91	0 132
10	3 51	0 129
12	3 12	0 131
15	2 74	0.129
20	2.22	0.131

The figures in the last column are satisfactorily constant, and confirm the application of the law of mass action to a bimolecular reaction.

Reactions of the third and higher orders are known and the formulæ applicable to these are obtained on the lines already indicated Cases of such reactions, however, are comparatively rare, and it is not proposed to discuss them here Investigation has shown that frequently a reaction which a general knowledge of the facts would lead us to regard as a reaction of a high order turns out to be one of a low order result is undoubtedly due to the apparently complex reaction being really a series of successive changes of a simpler kind Under these circumstances, the velocity of the complex re action is determined by those of the component reactions which are comparatively slow for instance, one of the latter is incomparably slower than all the others, then the order of the complex reaction will be the order of this slow component reaction

Influence of Temperature —It is well known that a chemical reaction is markedly accelerated by even a slight rise of temperature, but a definite measure of the influence of temperature on a given reaction can be obtained only by determining the velocity coefficient of the reaction at different temperatures The value of the temperature coefficient of reaction velocity, deduced in this way, is remarkably independent of the nature of the particular reaction under investigation For homogeneous reactions of the most varied character, the factor by which the velocity coefficient at To must be multiplied in order to give the velocity coefficient at  $T+10^{\circ}$ , lies, as a rule, between 2 and 4 Speaking Speaking generally, one may say that the velocity of a chemical reaction is doubled or trebled for every rise of 10° It is a striking fact that the temperature coefficient of various vital processes, such as vegetable respiration, is of the same order as that found for chemical reactions generally.

Chemical equilibrium. The reactions discussed in the foregoing section have all been changes which proceed from left to right until at least one of the reacting substances has disappeared. There are many reactions, however, which do not belong to this type, masmuch as they come to a stop while the reacting substances are still present in the reaction mixture Suppose, for instance, that the reaction between two substances A and B is of this description, and that C and D are the products of the reaction, then the change proceeds until a state of equilibrium is attained at which all four substances are present It is further found that if the products C and D are brought together, they react to form the substances A and B, while the final state of equilibrium reached is the same as in the previous case Such a reaction is known as a balanced or reversible reaction, and in the corresponding equation a double arrow is substituted for the usual sign of equality, thus  $\cdot$  A+B $\geq$ C+D.

The progress of a reversible reaction towards its equilibrium position may be regarded as the

side react to form those on the right side of the equation, and the other the velocity  $(v_{i})$ with which the substances on the right side react to form those on the left side of the equation. If we take  $A+B \ge C+D$  to represent the reversible reaction in question, and suppose that at some particular moment the concentrations of the four substances in the reaction mixture are a, b, c, and d respectively, then at that moment  $v_1 = k_1 ab$  and  $v_2 = k_2 cd$  The observed between  $v_1$  and  $v_2$  If now the change has proceeded at constant temperature until the equilibrium position is reached, and the concentrations of the four substances are ae, be, ce, and  $d_e$  respectively, then  $v_1 = k_1 a_e b_e$ , and  $v_2 = k_2 c_e d_e$ . Since there is equilibrium, the forward reaction is exactly balanced by the back reaction, so that  $v_1 = v_2$  and  $k_1 a_e b_e = k_2 c_e d_e$ , or  $k_1 = \frac{c_e}{a_e} \frac{d_e}{b_e}$  If the ratio  $\frac{k_1}{k_2}$  is replaced by the one constant K, we have  $K = \frac{c_e}{a_e} \frac{d_e}{b_c}$ . K is known as the equilibrium constant, and its numerical value defines the relationship and its numerical value defines the relationship which must exist between the equilibrium concentrations of the substances involved in a reversible reaction The argument employed above might obviously be extended to reversible reactions other than  $A+B \rightleftharpoons C+D$  · the result in all cases would be an equilibrium formula involving a constant ratio between the product of the equilibrium concentrations of the substances on the right side and the product of the equilibrium concentrations of the substances on the left side, however the values of these concentrations might vary individually - It ought, perhaps, to be pointed out here that, in the equilibrium formula, the concentration of any substance, n molecules of which take part in the reaction, must be raised to the nth power.

In discussing the application of the law of mass action to reversible reactions with the help of the equilibrium formula, we shall deal first with cases in which the reaction system is homogeneous As an example, the equilibrium between carbon dioxide, hydrogen, carbon monoxide, and water vapour may be considered, the reaction in question is represented by the equation  $CO_2 + \hat{H}_2 \gtrsim CO + H_2O$ , and is obviously of the type A+B C+D In Hahn's investigation of this equilibrium (Zeitsch physikal. Chem 1903, 44, 513), platinum was used as a catalytic agent, and on this ground the objection might be raised that the reaction as studied by that investigator is not really homogeneous. A true catalytic agent, however, as will be shown in greater detail at a later stage, does not affect the position of equilibrium in a reversible reaction, but merely lessens the time required for the attainment of equilibrium. The reaction which takes place between carbon dioxide and hydrogen, or between carbon monoxide and water vapour in contact with platinum, may therefore be regarded as a homogeneous reaction. In Hahn's experiments, mixtures of the gases were passed through a resultant of two opposed velocities, one the velocity  $(v_1)$  with which the substances on the left spongy platinum. Provided that the rate of

passage was slow enough to allow the four gases | to come into equilibrium with each other, analysis of the issuing gas gave the equilibrium concentrations,  $C_{CO_2}$ ,  $C_{H_2}$ ,  $C_{CO}$ , and  $C_{H_2O}$ . The relationship between these required by the law of mass action is  $K = \begin{pmatrix} C_{OO} & C_{H_2O} \\ C_{OO_2} & C_{H_2} \end{pmatrix}$ , and the validity of the law in the case of this reaction is shown by the fact that the value of K obtained is, for a given temperature, the same when carbon dioxide and hydrogen, in varying proportions, are allowed to react, as when carbon monoxide and water vapour are taken as the initial sub-A few figures obtained by Hahn at stances 986° will serve to support this statement Mixtures of carbon dioxide and hydrogen, in the proportions shown in the following table, were passed through the reaction vessel, and from analysis of the issuing gas the values of K were calculated by the above formula

Initial	mixture	Equilibrium mixture			
$CO_2$	$H_2$	('O <sub>2</sub>	('O=H2	O H <sub>2</sub>	K
10 Ï	89 9	0.70	9 46	80 38	1.59
30.1	69 9	7 18	$23\ 00$	46.82	1.58
49 1	51.9	$21\ 36$	27.88	22.88	1.59
60 9	39.1	34 20	$26 \ 61$	12.58	1 64
70.3	297	47.66	22.79	6.76	1.61

Experiments were made, also at 986°, in which the initial mixture contained carbon monoxide and water vapour the mean value of K, deduced from these experiments, was 155, slightly lower than the average of the figures recorded in the table. The discrepancy, however, is not great, and the numerical data quoted suffice to show that the balance established between the four gases, whatever be their initial concentrations, is adequately defined by the mass action formula

A reversible reaction of great technical importance is the one which forms the basis of the sulphuric acid contact process

and consideration of this case from the standpoint of the mass action law throws light on many points connected with the process. The reaction may be regarded as a homogeneous one, for the catalysts used in the technical process do not affect the final balance between the gases, but only accelerate the attainment of equilibrium

The application of the law of mass action, in

this case, leads to the result  $\frac{\binom{2}{80_3}}{\binom{2}{80_2}\binom{6}{0_2}} = \text{const}$ , in which  $\binom{C}{80_3}$ ,  $\binom{C}{80_2}$ , and  $\binom{C}{0_2}$  are the equilibrium concentrations of the respective substances. It is convenient, however, to work with the re-ciprocal of the foregoing expression, and to regard the equilibrium constant as defined by

 $K = \frac{C_{SO_2}^2 \cdot C_{O_2}}{C_{SO_2}^2}$  The accuracy of this formula may be tested with the help of the experimental data obtained by Bodenstein and Pohl (Zeitsch.

Elektrochem. 1905, 11, 373) These investigators passed a mixture of sulphur dioxide and oxygen over spongy platinum, contained in heated quartz tubes, and determined the equivor. II.—T.

librium concentrations by analysing the issuing gas. The figures in the following table, obtained at 727°, show that the value of K, worked out from the equilibrium concentrations, is independent (1) of the relative proportions of sulphur dioxide and oxygen passed into the apparatus, (2) of the presence or absence of introgen—

Initial gas ratio	
SO <sub>2</sub> O <sub>2</sub> N <sub>2</sub>	K×10
0 21 . 1 . 0	349
0.62:1 0	3 59
1 20 1 0	3.48
168 1 0	3 51
$3.97 \cdot 1 \cdot 0$	3 67
1 23 . 1 3 76	3 60
$1\ 31:1\cdot 3\ 76$	3 54
$1.55 \cdot 1 \cdot 3.76$	3.52

The experiments showed also that the value of K obtained for a particular gas mixture was independent (1) of the direction in which the equilibrium position was approached, (2) of the rate at which the gaseous mixture was passed over the catalyst. The latter result is the guarantee that the equilibrium position was really attained.

Similar results were obtained at various other temperatures, and it was found that the value of K increased rapidly with rising temperature, as shown by the following figures —

Temp C	K×10
$528^{\circ}$	0.013
$579^{\circ}$	0.077
$627^{\circ}$	0.32
680°	1 12
$727^{\circ}$	3 54
$789^{\circ}$	12 6
832°	28 0
897°	81.6

This increase means that, as the temperature rises, the position of equilibrium is shifted in favour of the system  $2SO_2 + O_2$ , in other words, the dissociation of sulphur trioxide becomes greater For each set of conditions, the extent of the dissociation is defined by the corresponding numerical value of K, deduced from a formula based on the foregoing table It is possible, then, for a given temperature and given proportions of sulphur dioxide, oxygen and nitrogen in the initial gaseous mixture, to calculate what will be the relative quantities of trioxide and dioxide in the equilibrium mixture, that is, to calculate the 'yield' obtainable The results of such calculations for various proportions of the gases and for various temperatures, are embodied in the following table numbers recorded in the last four columns are the percentages of sulphur dioxide convertible into sulphur trioxide under the specified conditions :-

Initial gaseous mixture SO<sub>2</sub> O<sub>2</sub> N<sub>2</sub> 400° 500° 10 1 p c 5 05 p c 84 85 p c 96 2 83 2 59 I 319 70 100 83 0 99 3 93.4 73.3 42 5 14.6 814 99.4 94.9 78.3 48.1 40 800 99 5 95 6 80 5 51 3 180

oxygen over spongy platinum, contained in These figures bring out clearly one other point heated quartz tubes, and determined the equi-

the influence of the partial pressure of the oxygen. If the equilibrium formula is written in the form

$$\frac{\mathrm{C_{80_{3}}^{8}}}{\mathrm{C_{80_{3}}^{2}}} = \frac{1}{\mathrm{K}} \cdot \mathrm{C_{0_{2}}} \text{ or } \frac{\mathrm{C_{80_{3}}}}{\mathrm{C_{80_{3}}}} = \frac{1}{\sqrt{\mathrm{K}}} \cdot \sqrt{\mathrm{Co_{2}}}$$

it is plain that the yield at a given temperature can be raised by increasing the partial pressure of the oxygen, and this conclusion is borne out by the foregoing table. The figures show also that the lower the temperature the better is the yield, but it must be remembered that from the technical point of view the time necessary for the attainment of equilibrium is also an important consideration. Fortunately, with platinum as catalyst, the union of sulphur dioxide and oxygen takes place with satisfactory rapidity, even at temperatures as low as 400°-500°, at which the dissociation of the trioxide is comparatively slight. With less active catalysts, however, such as ferric oxide, a higher temperature is required to secure a satisfactory velocity of reaction, and this involves a reduction of the obtainable yield.

Another reaction which is of exceptional interest from the standpoint of chemical equilibrium, and which has assumed great technical importance in recent years, is the synthesis of ammonia from nitrogen and hydrogen Although, as Deville showed 50-60 years ago, even prolonged exposure of ammonia to electric sparks leaves a trace of the gas undecomposed, and a minute quantity of ammonia is formed, when a mixture of nitrogen and hydrogen is passed through a heated tube, it was not until the early years of the present century that the reversibility of the reaction was properly appreciated and an attempt made to apply the principles of chemical equilibrium. The physico-chemical investigation of the equilibrium N<sub>2</sub>+3H<sub>2</sub> 2NH<sub>3</sub> has been carried out mainly by Haber and his collaborators (see Zeitsch. Elektrochem 1914, 20, 597; 1915, 21, 89 128, 191, 206, 228, 241), but this research, conducted originally on purely scientific lines, has developed in a remarkably short space of time into a large scale process for the production of ammonia from its constituent elements

Granted that there is a genuine equilibrium between nitrogen, hydrogen, and ammonia, Le Chatelier's principle indicates clearly that at any given temperature increase of pressure will shift the equilibrium position in favour of ammonia. The formulation of the equilibrium constant gives definite expression to this pressure effect, and for the present purpose the equilibrium concentrations of nitrogen, hydrogen and ammonia, but in terms of their equilibrium partial pressures  $p_{N_2}$ ,  $p_{\Pi_2}$ , and  $p_{NH_3}$ .

On this basis  $\frac{p_{NH_3}^2}{p_{N_3} \cdot p_{H_2}^3}$  has a constant value at a given temperature, or alternatively,

$$\mathbf{K} = \frac{p_{\mathbf{NH_3}}}{p_{\mathbf{N_2}}^{\frac{1}{2}} \cdot p_{\mathbf{H_2}}^{\frac{1}{2}}}$$

Since at high temperatures the equilibrium concentration of ammonia is relatively small no

great error is made, subject to that condition, by writing  $p_{\rm N_2}+p_{\rm H_2}={\rm P}$ , where P is the total pressure. If, further, the nitrogen and hydrogen have been taken in the exact proportions required by the chemical equation,  $p_{\rm H_2}=3p_{\rm N_2}$ . When these relationships are coupled with the expression for the equilibrium constant it follows that  $p_{\rm NH_3}=0.325{\rm K}\times{\rm P}^2$ . Hence, for low concentrations of ammonia, the partial pressure of this gas in the equilibrium mixture is proportional to the square of the total pressure

If x is taken to represent the volume percentage of ammonia in the equilibrium mixture,

 $x=\frac{p_{\rm NH_3}}{P}\times 100=32~5{\rm K}\times {\rm P}$  This relationship, according to which the volume percentage of ammonia, for low concentrations, should be proportional to the total piessure, is confirmed, for example, by some of Haber's experiments at  $800^\circ$ , in which for P=1 atmosphere, x was found to be 0 012, while for P=30 atmospheres x was 0 34

The shift of the ammonia equilibrium with temperature is shown very clearly by the following table, in which the values of K, based on measurements carried out at a pressure of 30 atmospheres, are recorded for a number of temperatures—

$$t^{\circ}$$
 C 561° 620° 700° 722° 801° 901° 952° K  $\times$  104 21 3 12 6 6 80 5 82 3·56 2·13 1·68

From the graph based on the foregoing values it is possible to work out for various even temperatures and even pressures the maximum quantities of ammonia obtainable by the interaction of introgen and hydrogen under various conditions. A selection of such figures, constituting a summary of the influence of pressure and temperature on the ammonia equilibrium, is embodied in the following table (valid for I vol introgen + 3 vols hydrogen):—

## EQUILIBRIUM PERCENTAGE OF AMMONIA

Temp	at	at	at	at
Cent.	1 atmo	30 atmos	100 at mos	200 at mos
400°	0.44	10 7	25 1	36.3
500°	0 129	362	10 4	17 6
$600^{\circ}$	0 049	1 43	4.47	8.25
$700^{\circ}$	0.022	0 66	2 14	4 11
800°	0 012	0 35	1.15	2 24
$900^{\circ}$	0.0069	0.21	0.68	1 34
1000°	0 0044	0 13	0 44	0 87

From the figures just quoted it is clear that the highest percentage conversion of nitrogen and hydrogen into ammonia is secured by working at a high pressure and a low temperature. In large-scale work, however, there is obviously a limit to the practicable pressure, and the pressure commonly employed in the technical process has been 150–200 atmospheres. A lower limit is likewise set to the temperature which can profitably be employed in practice by considerations of velocity. By themselves, nitrogen and hydrogen combine very slowly at temperatures below 1000°, and it is only in the presence of catalysts, such as osmium, uranium, and iron that the reaction is reasonably rapid at temperatures where the yield of ammonia is moderately satisfactory. A big yield and a

high velocity of reaction cannot be achieved simultaneously, and a balance has to be struck at the temperature which gives the best economic results. On the technical scale the temperature employed is generally 500°-600° C., and the mixed compressed gases are circulated over the heated catalyst in such a way that the ammonia formed is continuously removed by refrigeration or absorption It should be pointed out that in the manufacturing operation the rate of passage of the gases over the catalyst may be too rapid to permit the reaching of equilibrium at each stage, it may be economically preferable to mcrease the rate of circulation and to be satisfied with a correspondingly smaller conversion at each contact with the catalyst (see Maxted, J Soc Chem. Ind. 1918, 37,

Many other dissociation equilibria, similar in type to the cases just discussed, have been successfully attacked from the standpoint of the law of mass action. One instance is the equilibrium N<sub>2</sub>O<sub>4</sub> ≥2NO<sub>2</sub>, a case in which a physical method, namely, the determination of the vapour density, must be employed to find the relative proportions of the reacting substances in the equilibrium mixture. Obviously, no direct chemical method of analysis is permissible in any case where the equilibrium shifts rapidly at the ordinary temperature, for such analysis involves the removal of one of the substances involved, and this at once upsets the equilibrium.

Another technically important reaction, studied from the standpoint of the mass action law, is that which forms the basis of the Deacon process (see von Falckenstein, Zeitsch physikal Chem 1907, 59, 313, 1909, 65, 371) When a mixture of hydrogen chloride and air is passed through a heated chamber containing bricks impregnated with cupric chloride, the hydrogen chloride and oxygen react, producing chlorine and water vapour, but the reaction comes to a stop before all the hydrogen chloride disappears; the reaction, in fact, is a reversible one, and may be represented thus

Experiments have shown that the equilibrium is defined by an equilibrium constant

$$K = \frac{C_{Cl_2}^2 \cdot C_{H_2O}^2}{C_{HOl}^4 \cdot C_{O_2}}$$

the value of which, at a given temperature, is the same, whether the initial gaseous mixture taken is hydrogen chloride and oxygen (air), or chlorine and water vapour. The value of K diminishes as the temperature rises, hence the extent to which hydrogen chloride can be converted into chlorine also diminishes with rising temperature. If x represents the fraction of hydrogen chloride convertible into chlorine, then the value of x can be calculated for any given temperature from the corresponding value of K and the partial pressure of the oxygen. For a dry mixture of 76·1 p.c. HCl and 23·9 p.c.  $O_2$ , the value of x is 0.865 at  $352^\circ$ , 0.79 at  $430^\circ$ , and 0.55 at  $650^\circ$ . If air is substituted for oxygen so that the x-ratio of hydrogen chloride and oxygen remains the same, the value of x is 0.71 at  $430^\circ$ . This is approximately the temperature employed

in the technical process, and the yield obtained is about 70 p.c. The use of lower temperatures, which would increase the yield, is precluded practically because of the much longer time required for the attainment of equilibrium

The cases of equilibrium discussed so far have been those in which gases alone are involved. There are, however, many homogeneous reversible reactions in which liquid or dissolved substances take part. A classical example is the reaction between ethyl alcohol and acetic acid, represented thus:

If I mol of acid and I mol of alcohol are heated in a scaled tube at 100° for a sufficient time, examination of the reaction mixture shows that it contains \{ \} mol each of acid and alcohol, and \{ \} mol. each of ester and water. If then I mol of ester and I mol of water are mixed and treated in the same way, the equilibrium mixture finally obtained has the same composition as in the first case. The reaction therefore is truly reversible and the equilibrium constant

is given by the equation  $K = \frac{C_e}{C_{ac}} \frac{C_w}{C_{al}}$ , where  $C_e$ ,  $C_w$ ,  $C_{ac}$ , and  $C_{al}$  are the equilibrium concentrations of the four substances. If v represents the volume of the equilibrium mixture in litres, then for the case where 1 mol. each of acid and alcohol are taken initially  $C_e = C_w = \frac{2}{v}$  and  $C_{ac} = C_{al} = \frac{1}{v}$ , whence K = 4. It should be pointed out that the composition of the

be pointed out that the composition of the equilibrium mixture may safely be determined by titrating the free acetic acid, for the velocity of the reaction between ethyl alcohol and acetic acid is negligibly small at ordinary temperatures

If the law of mass action is strictly applicable to the reaction under consideration, then the value of K obtained in experiments in which alcohol and acid are taken in other than molecular proportions should also be 4. Suppose, for instance, that m mols of alcohol are allowed to act on 1 mol of acid, and that after equilibrium has been reached, the fraction x of a mol of ester has been produced; then  $C_a = C_{ty}$ 

mod of ester has been produced; then 
$$C_e = C_w$$

$$= \frac{x}{v}, \quad C_{ac} = \frac{1-x}{v}, \text{ and } C_{al} = \frac{m-x}{v}; \text{ so that } K$$

 $= \frac{x^{-}}{(1-x)(m-x)}.$  Instead of seeing whether the insertion of the known values of m and x in this formula gives the value 4 for K, the applicability of the mass action law may be tested by taking K=4 and ascertaining whether the value of x calculated by the formula for a given value of x is in agreement with the value of x determined experimentally. This procedure has been adopted in the actual investigation of the case, with the following results:—

m	$\boldsymbol{x}$ found	x calc.
0.08	0.078	0 078
0 28	0 226	0.232
0.50	0.414	0.423
0.67	0.519	0.528
1.2	0.819	0 785
2.24	0 876	0.864
8.0	0.966	0.945

The excellent agreement of the figures in the second and third columns furnishes a convincing proof of the applicability of the law of mass action. The influence of 'mass' or concentration in affecting equilibrium is very clearly illustrated by the numbers quoted in the table, for it will be observed that when the amount of alcohol is great compared with the amount of acid, the latter is almost quantitatively converted into ester, in other words, the 'yield' of ester from acid is nearly theoretical

According to a widely accepted view, an electrolyte dissolved in water is dissociated into positively and negatively charged particles, called ions, and investigation of the conductivity of electrolytic solutions and of the abnormal effect of electrolytes on the freezing-point and boiling-point of water supports the contention that the degree of this ionic dissociation increases with the dilution of the solution, and diminishes when the solution becomes more concentrated. The equilibrium, therefore, between the undissociated molecule of an electrolyte and the ions may be regarded as reversible

and can be represented thus:  $AB \rightleftharpoons A + B$  If this view is correct, then, as Ostwald pointed out, the equilibrium should be governed by the law of mass action, and the relationship between the

concentrations of AB, A, and B at a given temperature must be defined by an equilibrium constant. Suppose, for instance, that acetic acid is the electrolyte in question, then the equilibrium is represented by

$$CH_3 COOH \rightleftharpoons CH_3 COO + \overset{\leftarrow}{H}$$
 and the equilibrium formula will be  $K = \overset{C_1}{C_2} \overset{C_2}{C_3}$ 

where C, C1, and C2 are the concentrations of

the undissociated molecules and of the two ions respectively Since, however, the ions are produced in equivalent quantities, C<sub>1</sub>=C<sub>2</sub> and  $K = \frac{C_1^2}{C}$ . If we are dealing with a solution containing I gram-mol of acetic acid in V litres of solution, and if we take a to indicate the fractional extent to which the molecules are dissociated, then in V litres there is  $1-\alpha$  gram-mol of undissociated acetic acid, and accordingly  $C = \frac{1-\alpha}{V}$ . Similarly,  $C_1 = \frac{\alpha!}{V}$ , so that  $K = \frac{\alpha^2}{(1-\alpha)V}$ . For all binary electrolytes which, like acetic acid, dissociate into two ions, exactly the same formula would be obtained. The equation is the algebraic expression of what is known as Ostwald's Dilution Law, and K is usually termed a dissociation constant or an ionication constant: it is also called an affinity constart

The determination of  $\alpha$  is based on the relationship  $\alpha = \frac{\Lambda}{\Lambda_{\infty}}$ , where  $\Lambda$  is the equivalent conductivity of the electrolyte at the dilution V, and  $\Lambda_{\alpha}$  is the limiting maximum value of the equivalent conductivity. The application of the law of mass action to the equilibrium between acctic acid and its ions and to other similar cases is justified by the fact that the value of  $(1-\alpha)V$  is independent of the dilution at a given temperature. In support of this

statement, the following figures for acetic acid may be quoted:—

v	$K \times 10^6$
16	1.79
32	1 82
64	1 79
128	1 79
256	1.80
512	1.80

An equally satisfactory constancy of the expression  $\frac{\alpha^2}{(1-\alpha)V}$  has been observed in the case of all acids and bases which are dissociated to a comparatively small extent, and for these the validity of Ostwald's Dilution Law may be regarded as established. For some reason, however, which has not yet been adequately explained, highly dissociated electrolytes, such as strong acids and bases and neutral salts, do not obey Ostwald's Dilution Law, aithough in the case of potassium chloride Weiland (J. Amer Chem. Soc. 1918, 40, 131) seems to have established the validity of the law for this compound at extreme dilutions

Keeping in view this limitation to the validity of the formula  $K = \frac{a^2}{(1-a)V}$ , we may ask What is the significance of the constant K? The answer will be plain if two feebly dissociated acids, HA<sub>1</sub> and HA<sub>2</sub>, are compared at the same dilution V Since the acids are feebly dissociated, it is a very close approximation to the truth to put  $1-a_1=1$  and  $1-a_2=1$ , so that  $K_1=\frac{a_1^2}{V}$  and  $K_2=\frac{a_2^2}{V}$  A comparison of the equations shows that the relative value of the dissociation constants is closely related to the degree of dissociation; to put it generally, the dissociation constant is a measure of the ability of an acid to yield hydrogen ions, which, according to the modern view, are responsible for the characteristic properties of acids The value of K for an acid is, therefore, a measure of its strength, of its power to manifest those properties which are associated with acids set of cucumstances in which the relative strength of two acids is an important factor is realised when one gram equivalent of a base, such as sodium hydroxide, is added to a solution containing one gram equivalent of each of the two acids. The base is insufficient to neutralise both acids, and the result is a competition between the two acids, each securing a fraction of the base A theoretical discussion of this problem shows that if x is the fraction of the base appropriated by the first acid  $HA_1$ , and 1-x accordingly the fraction appropriated by the other acid HA2, then, provided the dilu-K, tion is not too great,  $\frac{x}{1-x} = \sqrt{\frac{K_1}{K_2}}$ . If, therefore, the values of  $K_1$  and  $K_2$  have been

HA<sub>1</sub> HA<sub>2</sub> z calc. z found
Dichloracetic acid Lactic acid 0.95 0.76
Formic acid Acetic acid 0.75 0.76
Butyric acid 0.54 0.53

determined in the way already described, it is

possible to calculate the distribution of the available base between the two competing acids.

That this can be done in good agreement with

experiment is shown by the following figures .-

It is evident, then, that a knowledge of the dissociation constant of an acid enables us to calculate relatively the affinity or avidity of this acid for a base. There are other methods for ascertaining the relative affinity or avidity of an acid, methods available in the case of strong acids which do not obey the dilution law, and to which, therefore, no definite value of K can be Among these methods are the determination of the heat effects accompanying neutralisation (Thomsen), the measurement of the density changes (Ostwald), and the study of the influence of acids in accelerating the inversion of sucrose or the hydrolysis of methyl The order of magnitude of the values obtained by these and other methods for the relative affinity of acids is the same, as will be seen by an inspection of the following table -

		deduced from	
Acid	K×100	Sucrose	Density
T160		inversion	changes
HCl		100	100
HNO <sub>3</sub>		100	100
CCI3 COOH	30 (approx	) 75	80
CHCl, COOH	5.14	27	33
CH₂CĬ COOH	0 155	4.8	7
H COOH	0.021	15	3
СН, СООН	0 0018	04	1

The manner in which a dissociation equilibrium is affected by excess of one of the products of dissociation has already been referred to in connection with the sulphuric acid contact process. It was there shown that as the equilibrium concentration of the oxygen is increased, so the extent to which the sulphur dioxide is converted into trioxide is increased also other words, the dissociation of sulphur trioxide into sulphur dioxide and oxygen is diminished in presence of excess of oxygen, one of the products of dissociation An analogous statement may be made in connection with the electrolytic dissociation of an electrolyte, such as acetic acid. If to a solution of this acid a quantity of CH<sub>3</sub> COO ions is added, the dissociation of the acid is diminished to a large extent, ie the concentration of the hydrogen ions is very much reduced, and the acid character of the solution almost disappears. It is true that CH, COO ions cannot be added by themselves, but a solution of the sodium salt of acetic acid serves the purpose, for such a salt, to judge from its conductivity, is highly dissociated into CH<sub>3</sub>·COO ions and sodium ions That the effective acidity of an acetic acid solution almost disappears on the addition of a little sodium acetate solution is very clearly seen by comparing the effects of (1) the acetic acid solution, (2) the acetic acid solution + sodium acetate, on the inversion of sucrose It has been found that the velocity of inversion in presence of

$$\frac{N}{4}$$
CH<sub>2</sub>·COOH+ $\frac{N}{4}$ CH<sub>3</sub> COONa

in about  $\frac{1}{75}$  what it is in the presence of  $\frac{N}{4}$  CH<sub>8</sub>COOH alone. Not only is this influence of a neutral salt on the effective acidity of a weak acid to be expected on the basis of the mass action law, but it has also been found possible, with the help of the equilibrium formula, to

calculate accurately the effect produced by adding any given quantity of the neutral salt.

The cases of equilibrium so far discussed have been those which occur in homogeneous systems, but there are numerous cases also of non-homogeneous or heterogeneous equilibrium. The bearing of the law of mass action on reversible reactions of this character will be most readily appreciated by considering one or two examples. The dissociation of calcium car-If this bonate may be taken in the first place substance is heated to 750° in a closed space, it breaks up to some extent into calcium oxide and carbon dioxide: if, on the other hand, carbon dioxide at atmospheric pressure is passed over calcium oxide, heated to the afore-mentioned temperature, the reverse action takes place and calcium carbonate is formed That is to say, an equilibrium is established which may be represented thus  $CaCO_3 \gtrsim CaO + CO_2$ . applying the law of mass action to this case, we are confronted with the question. What is the concentration or 'active mass' of a solid taking part in a reaction? An answer to this question may be reached by supposing that in the gaseous phase of the reaction system there is equilibrium between the calcium carbonate, the calcium oxide, and the carbon dioxide, that is, each of the solid substances concerned in the reaction is supposed to have a definite, if infinitely small, sublimation pressure at a given temperature, and therefore to be represented in the gaseous phase On this basis, the equilibrium at a given temperature would be governed by the formula  $K_1 = \frac{C_1C}{C_1}$ , where  $C_1$ ,  $C_2$ , and C are the equilibrium concentrations of calcium carbonate, calcium oxide, and carbon dioxide in the gaseous phase, respectively If partial pres-

sures are employed instead of concentrations, the formula would be  $K = \frac{\pi_2 \pi}{\pi_1}$ . Now  $\pi_1$  and  $\pi_2$ ,

being sublimation pressures, have fixed values at a given temperature, and are independent of the absolute amounts of calcium carbonate and calcium oxide present. Hence  $\pi = const$ , and since  $\pi_1$  and  $\pi_2$  are negligibly small,  $\pi$ , the partial pressure of the carbon dioxide, is practically equal to p, the total pressure exerted by the system, so that we may write  $p=\mathrm{const}$  The application, therefore, of the law of mass action to the dissociation of calcium carbonate leads to the result that at each temperature the reaction system exerts a definite pressure—the dissociation pressure, as it is termed. This statement is valid only on the understanding that the substances involved in the reaction are kept in contact with each other; if provision is made for keeping the pressure of the carbon dioxide permanently below or above the equilibrium value, then a reaction proceeds until the calcium carbonate (in the first case) or the calcium oxide (in the second case) has disappeared By artificial alteration of the pressure at constant temperature, therefore, any reaction of the same type as CaCO<sub>3</sub>  $\gtrsim$  CaO+CO, may be made to proceed either from left to right, or vice versa. The reaction 2BaO<sub>2</sub> ≥ 2BaO+O<sub>2</sub> is also a case where two solids and a gas are in equilibrium

with each other, and it may be treated in the same way as the dissociation of calcium carbonate.

Perhaps the chief point that emerges from the foregoing is that the active mass of a solid taking part in a chemical reaction is constant. The correctness of this conclusion has been confirmed by the investigation of various heterogeneous reactions different in type from the dissociation of calcium carbonate One of these may be considered here, viz. the equilibrium  $3\text{Fe}+4\text{H}_2\text{O} \gtrsim \text{Fe}_3\text{O}_4+4\text{H}_2$ by (see Deville, Compt. rend 1870, 70, 1105, 1201, 71, 30; Preuner, Zeitsch physikal. Chem. 1904, 47, 385) On the assumption that the active masses of the iron and the iron oxide are constant, the application of the law of mass action

obviously leads to the result 
$$\frac{p_{\rm H_2O}}{p_{\rm H_0}} = {\rm const}$$
 ,

 $p_{\rm H_{2}O}$  and  $p_{\rm H_{2}}$  being the partial pressures of the water vapour and the hydrogen when equilibrium has been established at a definite temperature. This relationship is verified by the experimental results, some of which are quoted in the following table. The experiments consisted in exposing iron in an electrically heated closed system to the action of water vapour kept up to a constant pressure  $(p_{\rm H_{2}O})$ . From the total pressure reached at equilibrium— $p_{\rm H_{2}O}+p_{\rm H_{2}O}+p_{\rm$ 

$p_{_{\mathrm{H_2O}}}$	$p_{_{ m H_2}}$	$\frac{p_{_{\rm H_2O}}}{\overline{p_{_{\rm H_2}}}}$
8 8	13 5	0 65
12 7	18 0	0.71
21 5	30 6	0.70
35 0	52.0	0 67
49:3	71.8	0.68

When the difficult nature of the experiments is taken into account, the numbers in the last column may be regarded as satisfactorily constant and as demonstrating the applicability of the mass action law to this case. Although in a closed system an equilibrium is reached in the way described, yet if provision is made for removing the hydrogen as fast as it is formed, and thereby keeping its partial pressure permanently below the equilibrium value, the reaction will proceed until the iron is all converted into oxide

There are many reversible reactions involving a solid substance and a liquid, or a solid and a dissolved substance. Here also the active mass of the solid is constant at a given temperature, solubility and sublimation pressure being analogous quantities. The equilibrium formula undergoes, therefore, a similar simplification to that illustrated in the foregoing paragraphs

that illustrated in the foregoing paragraphs.

Influence of Temperature.—In general, a reaction system which is in equilibrium at a given temperature is no longer in equilibrium when the temperature is altered. A measure of this influence of temperature on chemical equilibrium is found in the change of the equilibrium

constant, and an illustration of the extent to which K may change with temperature has already been given in connection with the oxidation of sulphur dioxide to trioxide. The exact relationship between temperature and equilibrium is defined by a formula associated with the name of van't Hoff, according to which  $d(\log_e K)$ 

 $r' = -\frac{1}{RT^2}$ . The symbol K is the equilibrium constant, that is, the ratio between the product of the equilibrium concentrations of the substances on the right side of the equation and the product of the equilibrium concentrations of those on the left side; q is the heat evolved when the reaction goes completely from left to right, T is absolute temperature, and R is the gas constant The theoretical basis of van't Hoff's formula need not be discussed here, but one or two points involved in it may be indicated. It is evident, in the first place, that when the heat effect of a reversible reaction is small, the variation in K is also small, i e. the position of equilibrium is but slightly shifted on altering the temperature. An instance of this is furnished by the reversible reaction between ethyl alcohol

and acetic acid. Again, if q is positive,  $\frac{d}{dT}$  is negative, so that K diminishes with rising temperature, corresponding to a shifting of the equilibrium in favour of the substances on the left side of the equation. If, on the other hand,

 $d(\log_e K)$ 

$$q$$
 is negative,  $\frac{(d\log_e \mathbf{K})}{d\mathbf{T}}$  is positive, and a rise of

temperature, therefore, involves a displacement of the equilibrium in favour of the substances on the right side of the equation. It will be observed that in both cases, whether q is positive or negative, a rise of temperature involves a shifting of the equilibrium in the direction which means absorption of heat

As an illustration of the applicability of van't Hoff's formula in an individual case, the reaction  $2SO_3 \geq 2SO_2 + O_2$  may be taken From the values of the equilibrium constant obtained for this reaction at different temperatures (see p. 161), the values of q over narrow intervals of temperature were calculated by the formula. It was found that q is a linear function of the temperature, and may, in fact, be represented by the equation q = -47300 + 4". If this expression for q is inserted in the formula  $\frac{d(\log_2 K)}{dT} = -\frac{q}{RT_2}$ , and the equation is integrated, the following formula is obtained.—

$$\log_{10}\mathrm{K} \!=\! -\frac{10373}{\mathrm{T}} \!-\! 2\ 222\ \log_{10}\!\mathrm{T} \!+\! 14\ 585$$

The relation between K and T expressed in this formula is almost exactly that observed in the experimental investigation: this appears from the following figures:—

t∘ C	log K found	log K calc.
528	-4.809	-4817
579	$-4\ 126$	4 099
627	3 500	3 502
680	-2914	-2.893
727	-2451	-2453
789	-1900	-1.922
832	-1553	-1573
897	-1089	-1.083

Catalysis. No discussion of reaction velocity or chemical equilibrium would be complete without a reference to the phenomenon of catalysis. It very frequently happens that a chemical reaction, which of itself is extremely sluggish, proceeds with comparative rapidity in presence of some foreign substance—a catalyst, as it is called. Cases of such an acceleration of a chemical reaction have long been known, and in both laboratory and technical processes catalysts are in common use

A rational quantitative study of catalysis is possible only on the basis of the law of mass action. In the velocity coefficient, as already explained, we have a definite measure of the rate of a chemical change under given conditions For a given reaction, therefore, which is catalytically accelerated, the value of the velocity coefficient at a given temperature is a measure of the efficiency of the catalyst, and by comparing the values obtained for the velocity coefficient in different experiments, one can find out how the efficiency of a catalyst varies with the conditions under which it works, and how the efficiency of one catalyst compares with that of another working under the same conditions

General Characteristics of Catalysts -One very common and striking feature of catalysis is that the quantity of the catalyst is so small compared with the quantities of the main reacting sub-As a rule, too, the activity of the catalyst at the end of the reaction which it has accelerated is unimpaired. An illustration of these features of catalysis is furnished by the influence of colloidal platinum in promoting the union of hydrogen and oxygen at the ordinary temperature. In some experiments made by Bredig, 25 c c of colloidal platinum solution (containing 017 milligram of platinum) were shaken with electrolytic gas, with the following results -

Time in minutes 10	Decrease in the volume of gas 17 8 c c	Decrease per minute 1 78
20	358 ,,	1 80
30	548 ,,	1 90
40	72.4 ,,	1 76
50	90 2 ",	1 78

The average of the figures in the last column is a measure of the catalytic efficiency of the colloidal platinum in the early stages of its The same colloidal platinum was activity. then shaken intermittently during 14 days with the mixture of hydrogen and oxygen, about 10 litres of which disappeared in this time. Not only is the contrast between the minute quantity of the catalyst and the extent of the induced change sufficiently striking, but a determination of the actual rate of disappearance of the gas at the end of the 14 days showed that the rate of decrease, measured over successive 10-minute intervals, was 2 02, 1 87, 1 95, 1.97, and 2.01 cc. per minute From these figures it is obvious that the catalytic efficiency of the colloidal platinum is unimpaired. connection with this case, however, the objection might be raised that hydrogen and oxygen do not combine at the ordinary temperature, and that the platinum ought to be regarded, not as accelerating a change which would take place in its absence, but as actually initiating the re-

The answer is that at 500° hydrogen action and oxygen combine with appreciable iapidity in the absence of a catalyst, but that, in view of the large temperature coefficient of reaction velocity, the rate of union naturally becomes insignificant even at 400°, whilst at ordinary temperatures the two gases appear not to react There is, however, no reason to suppose that there is a definite temperature below which absolutely no combination takes place

The mere fact that the amount of a catalyst may be almost infinitely small compared with the amounts of the substances changed under its influence, shows that the final state of the reactive system must be independent of the catalyst, in particular, if the accelerated reaction is reversible, the state of equilibrium finally attained must be the same as when no catalyst is present. The catalyst, that is, influences only the rate at which the condition of equilibrium is reached, not the position of equilibijum itself Experimental evidence in favour of this proposition is supplied by Turbaba's investigation of the relationship between aldehyde and paraldehyde. The conversion of pualdehyde into the equilibrium mixture, which at 50 5° contains 33 9 pc aldehyde, is accompanied by an expansion, and the course of the change may therefore be followed by means of a dilatometer Various substances accelerate the change, but the difference between the initial and final volumes, as shown by the following figures, is independent of the nature and the amount of the catalyst -

Catalyst Sulphur dioxide	Per cent of catalyst 0 08	Percentage increase of volume 8 20
,, ,	0 07	8 34
,, ,,	0.002	8 19
Zinc sulphate	27	8 13
Hydrochloric acid	0.13	8 13
Oxalic acid	0.25	8.27
Phosphoric acid	0.54	8 10

If the position of equilibrium in a reversible reaction is independent of the catalyst, then K, the equilibrium constant, must be similarly Further, since  $K = \frac{k_1}{k_2}$ ,

mdependent.

 $k_1$  and  $k_2$  are the velocity coefficients of the forward and back reactions, it follows that the catalyst must accelerate the forward and the back reaction in the same proportion. This has been demonstrated in connection with the catalytic effect of acids on the action between an acid and an alcohol and the reverse change. It should further be pointed out that the amount of energy transformed during a chemical change, being dependent only on the initial and final states of the system, is independent of the The latter may be compared to the

energy derivable from the fall of the weight. Another point of great interest is the relationship between the value of the velocity coefficient for a given reaction at a given temperature and the concentration of the catalyst. In many cases the relationship is a linear one The rate of inversion of sucrose by acids, for instance, is proportional to the concentration of

oil which facilitates the sliding of a weight down an inclined plane, without affecting the total

the hydrogen ions, provided that this concentration is low, on the basis of this proportionality, in fact, it is possible to calculate the velocity of inversion by dilute acetic acid from the velocity observed with dilute hydrochloric acid. In other cases, however, the relationship between reaction velocity and concentration of catalyst is not a linear one. The decomposition of hydrogen peroxide under the influence of colloidal platinum is an illustration of this The course of the decomposition can be followed by extracting a definite volume of the reaction mixture from time to time and titrating with potassium permanganate. The course of the change is in harmony with the formula for a unimolecular reaction, and for such concentration of platinum a definite value of k, the velocity coefficient, can be obtained The velocity coefficient, can be obtained The following table contains the values of k corresponding to various concentrations of platinum -

Platinum concentration	$\boldsymbol{k}$
$21.0 \times 10^{-6}$	0.072
10 5×10-6	0.024
5 2×10-6	0 0084
$2.6 \times 10^{-6}$	0 0027

From these figures it appears that when the concentration of the catalyst is doubted, the velocity of decomposition is trebled

Catalysis and the law of mass action—The general characteristics of catalysts have been discussed in the foregoing paragraphs, but it must be admitted that the phenomena associated with catalysis are frequently more complex than has been suggested. It appears that the catalyst often works in such a way as to mask the operation of the law of mass action altogether. Although a detailed discussion of these peculiarities of catalytically accelerated reactions is

not possible here, a brief consideration of some typical cases will be found useful

An investigation has been made of the rate at which sulphur dioxide and oxygen unite in presence of platinum (see Bodenstein, Zeitsch physikal Chem 1907, 60, 1), and the results show that the velocity of union is independent of the oxygen concentration (except when it is extremely small), proportional to the concentration of sulphur dioxide, and inversely proportional to the square root of the trioxide concentration This behaviour is obviously quite different from what would be expected under the law of mass action The explanation adopted by Bodenstein is that the factor which really determines the velocity of reaction is the rate of passage of the sulphur dioxide through a layer of sulphur trioxide, absorbed at the surface of the platinum From this point of view, what is measured is a diffusion velocity, not a reaction velocity, for it is supposed that the sulphur dioxide and the oxygen, when they have reached the platinum, react with great rapidity stein has further shown that the rate of union of the two gases under the influence of other catalysts is governed by the same factors. only so, but it has been found that the rate of change in many other heterogeneous catalytic reactions is, in all probability, determined by the physical process of diffusion This is the case with the union of hydrogen and oxygen in contact with heated porous porcelain (see Bone and Wheeler, Phil. Trans. 1906, 206, 1); in this

instance, the rate of union of the gases appears to be determined to a large extent by the rate of occlusion of hydrogen.

Additional support of the view that the velocity of a heterogeneous catalytic reaction may be determined mainly by a diffusion process, is furnished by the relatively low value of the temperature coefficient in these cases As already indicated, the velocity of a homogeneous chemical reaction is doubled or trebled for every

rise of  $10^{\circ}$ , but the value of  $\frac{k_{\mathrm{T}+10}}{k_{\mathrm{m}}}$  for a hetero-

geneous reaction is frequently not much greater than the value (about 1.3) we should expect it the rate of reaction were determined by a diffusion velocity alone

Among the most important catalytic reactions are those which take place under the influence of enzymes These catalysts are able to promote changes in organic matter which, in their absence, can be effected only with great difficulty. There is a close analogy between enzymes and ordinary morganic catalysts, but it is frequently found that, owing to the operation of various factors, the course of a reaction which takes place under the influence of an enzyme, deviates considerably from what we should expect on the basis of the law of mass In the inversion of sucrose by invertase, for example, the velocity coefficient, calculated by the formula for a unimolecular reaction, increases with the time (Brown, Chem Soc Trans 1902, 373) This is shown by the following table, which refers to the inversion of a 9 48 p c sucrose solution at  $30^{\circ}$ , the symbol x stands for the fraction of the total sucrose inverted by time t —

t min.	x	$\frac{1}{t} \log \frac{1}{1-a}$
30	0 265	$t = 1-x \\ 0.00445$
64	ð 509	0 00483
120	0 794	0 00571
180	0.945	0 00698
240	0.983	0 00737

The departure from the law of mass action becomes still clearer when experiments are made in which a constant amount of invertase is allowed to act for a given time on varying amounts of sucrose in a constant volume of solution Such experiments show that the enzyme, instead of inverting a constant fraction, as required by the law of mass action, has inverted a constant weight of sucrose in the given time, only when the quantity of sucrose is very small is the law of mass action obeyed In the hydrolysis of starch by diastase, and of milk sugar by lactuse, it is similarly found that the amount of change induced by the enzyme is, for the initial portion of the change, a linear function of the time (see Brown and Glendinning, Chem Soc Trans. 1902, 81, 388; E. F. Armstrong, Proc Roy. Soc 1904, 73, 500); the later portion of the change, when the amount of catalyst is larger relatively to the amount of carbohydrate, conforms to the law of mass action.

Another peculiarity about enzyme action, which is frequently observed, is that the activity of the enzyme does not remain constant throughout the whole course of the change which it induces. In the hydrolysis of amygdalin by emulsin, and of milk sugar by lactase, the

products of reaction seem to exert a specific influence in putting the enzyme out of action (see Tammann, Zeitsch. physikal Chem 1892, 13, 271; Armstrong, Proc. Roy Soc 1904, 73, 500). This is indicated by the regular falling off in the value of  $\frac{1}{t} \log \frac{1}{1-x}$  as the reaction proceeds, that is, as the products of the reaction,

accumulate

Mechanism of catalysis - Many theories have been advanced as to the way in which a catalyst exerts its influence According to one very commonly accepted view, a catalyst is effective because it combines with the substrate, and this intermediate compound then breaks up into the final products of change, the catalyst being liberated. Obviously, if this account of catalytic change is to give an adequate interpretation of the phenomena, it must be supposed that the formation and decomposition of the intermediate compound together require a shorter time for their occurrence than the direct change itself. There is a good deal of evidence, largely of an inducet kind, in favour of the view that combination takes place between catalyst and substrate. The increased stability of invertase in presence of sucrose, the specificity of enzymes, and the occurrence of a linear portion in the time curve for the hydrolysis of sugars, are facts which contribute to this evidence In the case of the catalytic influence of molybdic acid on the reaction between hydrogen peroxide and hydrogen iodide, it has been definitely shown that combination takes place between the molybdic acid and the hydrogen peroxide Aluminium chloride also, well known as a catalytic agent, forms definite compounds with the substances the reactivity of which promotes But although we may frequently assume the very probable formation of intermediate compounds in enzyme action and catalysis generally, it is quite impossible, in many cases, to specify the nature of these compounds

There are, moreover, many cases in which the efficiency of a catalyst depends almost certainly on something else than the formation of intermediate compounds. The investigation of non-homogeneous catalytic reactions, such as the combination of sulphur dioxide and oxygen under the influence of platinum, or the union of hydrogen and oxygen under the influence of porous porcelain, has led to the conviction that in these cases occlusion or adsorption of the reacting substances occurs at the surface of the catalyst, and that the higher concentration of the reagents thus secured is responsible for the greater velocity of reaction. It is possible to interpret enzyme action also on this basis, for the solution of an enzyme is, strictly speaking, a non-homogeneous system.

non-homogeneous system.

Bibliography. — Chemical Statics and Dynamics, J. W Mellor; Theimodynamics of Technical Gas Reactions, F. Haber, Nature of Enzyme Action, W. M. Bayliss, Cantor Lectures (Royal Society of Arts), 1918, J. C. Philip.

J. C. Philip.

# CATALYSIS IN INDUSTRIAL CHEMISTRY.

Within the last two decades the applications of catalysis in industrial chemistry have undergone remarkable developments. Many catalytic pro-

cesses are now in operation on the large scale for the production of both inorganic and organic substances, and the steadily increasing amount of research work done in this field of investigation will, without doubt, lead to the introduction of many other methods of manufacture based upon the influence of catalytic agents

Methods of preparation of inorganic substances in which catalysts are employed are as yet comparatively few, but amongst them are included processes for the production of such indispensable materials as hydrogen, chlorine, sulphune acid, ammonia, and nitric acid. In order to meet the growing demand for pure hydrogen various methods for the production of that element from water have been worked out, and several of these are based upon the fact that in presence of suitable catalysts steam and carbon monoxide interact at moderately high temperatures with the formation of hydrogen and carbon dioxide, which, of course, can be easily separated Iron or other metals of the non group serve as catalysts, when the former is employed its activity is increased by the addition of small quantities of 'activators,' such as potassium hydioxide, or by using it in the form of an non-copper couple. The reaction between steam and carbon monoxide may also be effected in presence of heated lime, pre-ferably containing a small admixture of iron, or of a mixture of lime and charcoal

Chloring may be prepared catalytically by the well-known Deacon process, in which hydrochloric acid is oxidised by atmospheric oxygen in presence of cupric chloride at a temperature of about 400°, and also by the method due to Hargieaves and Robinson, in which the reaction between sodium chloride, sulphur dioxide, and oxygen, which results in the formation of sodium sulphate and chlorine, is promoted by the presence of a catalyst such as cupric oxide

In the contact process for the manufacture of sulphuric and the primary reaction, viz the formation of sulphuric anhydride by combination of sulphur dioxide with oxygen, is effected through the agency of platinum, although ferric oxide is also used as a catalyst. In the former case very complete preliminary purification of the gases is essential, since various impurities, and especially arsenic, even in minute quantities, quickly 'poison' the catalyst and

impair or destroy its activity

The synthesis of ammonia is now carried out on a very large scale, since it has been discovered that when a highly compressed mixture of nitrogen and hydrogen is passed over a catalyst maintained at a temperature of 500°-600° the proportion of ammonia produced is sufficiently great to make the process remunerative. Even under the most favourable conditions most of the nitrogen and hydrogen remain uncombined, hence it is necessary to adopt a continuous process in which the residual gases are again passed over the catalyst after removal Many different metals and of the ammonia other substances, e.g. metallic nitrides, have the power of catalysing this reaction, but it is understood that iron is employed in practice; the metal is in a fine state of division and rendered active by the addition of a promoter.

When a mixture of nitrogen with oxygen or air is passed over a suitable catalyst at

temperatures of 500° and upwards nitric oxide is formed; excess of oxygen converts this compound into mtric peroxide, which, when dissolved in water, yields nitric acid and nitric oxide. The catalytic method of manufacturing nitric acid from ammonia is based upon these reactions. Originally platinum was used as the catalyst, but it has been found that iron activated with small quantities of bismuth or other substances is also very efficient. The mixed gases are passed over the heated catalyst with high velocity, hence a relatively small contact chamber suffices for the oxidation of comparatively large quantities of ammonia So far as expenditure of power is conceined, this process is much more economical than the arc process, in which nitiogen and oxygen are caused to combine directly at a very high temperature

Catalysts are also utilised in the manufacture of graphite from charcoal or coke in the electric furnace; in the preparation of carbon tetrachloride; in the elimination of carbon disulphide from coal gas, in the production of sulphur by limited combustion of sulphuretted hydrogen in the ('laus kiln; in the preparation of sulphuryl chloride, of hydrazine, and of aluminium nitride, and in the electrolytic processes for the preparation of hypochlorites, chlorates, and persulphates, and for the regeneration of chromic acid. Moreover, it is claimed that they may be used with advantage in making sodamide, and in the manufacture of alkali cyanides by the action of nitrogen on heated mixtures of alkali carbonates and charcoal.

Largely as a result of the proneer work of Sabatier and Senderens catalytic methods are likely to become of great importance in the domain of organic chemistry Thus, for instance, hydrogen has been found available for many operations of reduction and of hydrogenation when used along with various catalysts the former class of reactions nickel is in general very effective, but as it is sensitive to 'poisons,' particularly the halogens, care must be taken that both the hydrogen and the substance to be attacked must be freed from such impurities. Palladium and platinum, especially when in the colloidal state, are very active catalysts of such The operations may be carried out either at the ordinary or at increased pressure, and either at such temperatures that the reacting substances are vaporised, or at the ordinary temperature with the substances in the liquid state or in solution Among reducing actions of importance carried out at high temperatures may be mentioned the production of methane from carbon monoxide or dioxide, the preparation of aromatic amines from nitro compounds, and the reduction of aldehydes and ketones to alcohols. In the reduction of nitio compounds copper, either alone or with the addition of an activator, is an effective catalyst, but may be replaced by iron, nickel, silver, or gold The hydrogen may be used by itself or mixed with steam, and in place of hydrogen mixtures of carbon monoxide and steam may be employed.

The preparation of indigo white may be quoted as an example of reduction at the ordinary temperature. Indigotin, suspended in a solution of sodium hydroxide, is rapidly reduced by hydrogen at 60°-80° in presence of active mckel; the catalyst is easily removed

by filtration, and the solution of indigo white thus produced has the advantage of being free from any excess of the reducing agent. process is applicable to the preparation of leuco derivatives of other vat dyes.

With nickel as catalyst the addition of hydrogen to unsaturated compounds, when in the state of vapour, is easily effected. Ethylene, for example, is readily converted into ethane at 200°, and the preparation of hexahydro derivatives of benzene and other aromatic compounds is carried out with little difficulty In solution, and especially in presence of colloidal platinum or palladium, various alkaloids have been converted into hydro derivatives, which may prove of value medicinally, and a most useful application of the process of hydrogenation is the 'hardening of fats,' ie the conversion of oils or liquid fats into solid fats This hardening process depends essentially upon the production of the solid glyceride tristearin by addition of hydrogen to the liquid glyceride Nickel is usually employed as the tijolein catalyst, but the oxide of nickel has also been recommended, and the use of a number of other metals (particularly those of the platinum group) and of their compounds has been patented

A number of processes, in which reduction is effected by agents other than hydrogen, are also promoted by various catalysts example, in the reduction of azobenzene with sulphurous acid, the presence of a trace of hydriodic acid has a most beneficial effect, and the remarkable discovery has been made that the reduction of certain dyes with hydrosulphites or formaldehyde-sulphoxylates is greatly promoted by the addition of small quantities of certain other dyes Moreover, in a number of electrolytic reduction processes, the presence of catalysts is advantageous; thus the electrolytic reduction of nitro-, nitroso-, or azo- compounds is facilitated by the addition of titanous salts,

in very small quantity, to the liquid

Many processes of oxidation are also induced or accelerated by catalysts A well-known example of direct oxidation is the production of formaldehyde by passing a mixture of the vapour of methyl alcohol and air through a tube containing heated copper or platinum. Acetaldehyde is rapidly and smoothly converted into acetic acid by the action of a current of air if a little manganous acetate is dissolved in the aldehyde. Anthraquinone may be obtained by heating anthracene with oxygen, under pressure, if some cupie oxide or other catalyst is also present. In oxidations with sulphuric acid, as, for instance, the preparation of phthalic acid from naphthalene, the velocity of the reaction is in many cases greatly increased by the addition of small quantities of mercury or copper. If catalysts are present solutions of alkalı chlorates readily act as oxidising agents in many cases where otherwise the reaction takes place slowly or not at all; thus a trace of vanadium pentoxide has a most advantageous effect in the preparation of aniline black by oxidising aniline with these reagents. Hydrogen peroxide is a most valuable oxidising agent for many hydroxy compounds if traces of ferrous salts are also present, and some processes in which oxidation is effected through the agency of nitrobenzene, e.g. the manufacture of magenta,

CHERRY. 171

re promoted by the addition of ferrous salts hich act catalytically as oxygen carriers. foreover, electrolytic oxidation is frequently ceelerated by catalysts; for example, in this ethod of preparing anthraquinone, a little ceric ulphate or chromic acid is added to the sulhuric acid in which the anthracene is suspended.

Catalysts are capable of promoting processes f dehydrogenation as well as of hydrogenation. Or instance, an excellent method of obtaining cetaldehyde consists in passing the vapour of thyl alcohol over copper heated to about 300°, and in a similar manner ketones may be picared from secondary alcohols, e.g. camphor com borneol, with satisfactory results. Hydrormbons also are dehydrogenated at comparatively low temperatures in presence of metals uch as copper, nickel, or iron, and in this onnection it should be noted that a number of atents have been taken out which cover the mployment of those and other catalysts in he 'cracking' of petroleum for the production f motor spirit, benzene, &c , from heavier oils.

Other reactions in which catalysts are ffective include the hydration and the dehydraon of organic compounds An example of ydration is the preparation of acetaldehyde by iducing the combination of acetylene with ater, which is carried out by passing a current f the gas into an aqueous solution of an acid in resence of a mercury salt which acts as the atalyst. The hydrolysis of esters, which may lso be included in this category, finds practical lustration in the hydrolysis of fats into free itty acids and glycerol by the action of water ontaining a small quantity of a catalyst, eg alphophenylstearic acid. Dehydration is exemhed by the catalytic preparation of ethylene and other unsaturated hydrocarbons. Ethyl scohol breaks down into ethylene and water then its vapour is passed over alumina heated ) a temperature of 360°, and isoprene and other ydrocarbons of the same series can be obtained y a similar dehydration of dihydric alcohols he catalytic influence of mineral acids in ccelerating the formation of esters by the iteraction of organic acids and alcohols is ell known, and it has been found that acetaldeyde and ethyl alcohol readily react with formaon of acetal in presence of calcium chloride and

trace of hydrochloric acid. Moreover, in inny of the processes grouped under the heading condensation,' which find such extended pplication, there is no doubt that the conensing agent plays the part of a catalyst.

Much space would be required if an attempt ore made to illustrate all the other kinds of sactions of organic compounds which are accountably influenced by catalysts, and it must affice merely to indicate the employment of these agents in connection with the following perations: The preparation of aromatic ydrocarbons, e.g. of toluene by the action of iethyl chloride on benzene in presence of luminium chloride; the preparation of halogen

derivatives of both aromatic and fatty compounds, when various catalysts known as halogen carriers are of the greatest service, the preparation of sulphonic acids derived from benzene and other hydrocarbons, from pyridine, &c., the preparation of derivatives of amines or of amino- compounds, particularly of those in which aromatic radicals are introduced into the amino-group, eg the production of phenylanthranilic acid, which is easily effected by the addition of a small quantity of copper powder to a boiling solution of o-chlorobenzoic acid in aniline; the preparation of certain diazo- compounds, and the replacement of the diazo- group by the chlorine, bromine, or cyanogen radical; the preparation of aldehydes and ketones, eg of benzaldehyde from benzylidene chloride or directly from benzene, and of acetone from acetic acid; the preparation of certain sulphur compounds, in which category the important methods of vulcanising rubber may be included. Moreover, in many cases, intramolecular rearrangements and polymerisations are accelerated by catalysts. The transformation of hydrazobenzene into benzidine under the influence of a strong acid, is a familiar example of the former class of change, whilst the latter is exemplified in the synthesis of various kinds of rubber by the polymerisation of butadiene and its homologues, which is promoted by the presence of sodium, acetic acid, and other catalysts, and in the conversion of acetaldehyde into para-acctaldehyde, ethyl acetate, or aldol

Finally, it must be noted that a number of very important operations, for example, the manufacture of ethyl alcohol, are carried out through the agency of that group of catalysts produced by hving organisms which are known as enzymes. In many respects the action of enzymes is similar to that of inorganic catalysts, and especially to that of colloidal platinum or palladium, but is specific in the sense that a definite enzyme is required to bring about a particular transformation.

G. G. H.

**CHERRY.** The fruit of *Prunus avrum* (sweet cherry) or *P. cerasus* (Morella cherry). There are many varieties. Konig gives, as the composition of ordinary cherries:

Free Invert Sacch (albo-Water Protein acid sugar arose hydrates Fibre Ash 80 6 1 2 0 7 8 9 0 5 1 8 5 8 0 5

Included in the carbohydrates is 0.6–1.0 p.c. of pentosans (Wittreaun, 1991). A more detailed examination by Keim (Zeitsch, anal. Chem 1891, 30, 401) showed that in the unripe fruit of *P cerasus*, citric, malic, and succinic acids were present; that cane sugar was present, but diminished greatly as the fruit ripened. The ripe fruit contained no succinic acid, but malic and citric acid equal to 0.46 p.c. expressed as malic acid, were present, and the sugars were dextrose, levulose, and traces of inositol. Shaw (Expt. Station Record, 1899) found, as the mean of the analyses of 13 varieties of cherries:

Average	Whole	e fruit Flesh		In the flesh				
weight	Flesh	Stone	Juice	Residue	Water	Protein	Sugai	Ash
5.29 grams .	93.9	61,	86 4	13 6	81:3	0.9	11 2	05

172 CHERRY

Konig gives, as the mean of many analyses of cherry juice: Sp gr. 1 0737; total solids, 194; sugar, 128; protein, 042; free acid, 075, tannin, 009; ash, 055, grammes per 100 c.c.

Konig gives, as the composition of the ash.

Flesh only 50 1 - 7 0 5 2 - 12 9 -

The kernels of cherry stones contain.

Nitrogen-Water Protein Wat. iree extract Fibre 33 4 3.5 Ash 41.0 14 1 6.9 10

The kernels contain about 0.82 p.c. of amygdalın.

Dued chernes contain, according to analyses by Konig and Krauch (flesh only)

Other carbo-

Invert sugar hydrates Protein Fat Water Fibre 49.9 312 143 0.3

Willstatter and Zollinger (Annalen, 1916, 412, 164) find that the colouring matter of the skins of cheries is keracyanin, the chloride of which can be obtained in crystals containing either 4 mols of water (red slender needles) or  $3\rm{H}_2O$  (stout brown prisms). The chloride has the composition  $\rm{C}_{27}\rm{H}_{31}\rm{O}_{15}\rm{C}I$ , and is readily hydrolysed by boiling with hydrochloric acid, yielding cyanidin, dextrose, and rhamnose. Keracyanın is thus a diglucoside of cyanidin C15H10O6, a frequent constituent of the colouring matter of flowers and fruits (v  $H^{-1}$ ANTHOCYANINS.

CHERRY BARK TANNIN v TANNINS CHERRY-LAUREL LEAVES. Laurocci asifo lia (BP), leaves of Prunus Laurocciasus (Linn).

CHERRY LAUREL OIL v OILS, ESSENTIAL CHERRY TREE GUM v GUMS

CHERT. (Feuerstein, Hornstein, Ger) A silicious mineral allied to chalcedony and flint. Similar terms are applied to horistone and to any impure flinty rock, including the jaspers It is worked extensively from carboniferous hmestone quarries in Flintshire, especially at Halkin and Talacie in Denbighshire, Derbyshire, and at Reeth in Yorkshire It is used in potteries for paving the mills in which flints are The following is an analysis of a sample from Vizzin, Sicily, given by Ricciardi (Gazz chim. ital 11, 319)—

Interior S<sub>1</sub>O<sub>2</sub> Al<sub>2</sub>O<sub>3</sub> FeO CaO 96 31 0 30 0 25 0 47 H<sub>2</sub>O Sp gr at 22° 2 81=100 14 2 572 96 31 03 0 025 047 281=100 14 2572 White crust metamorphosed by heat \$10\_2 Al\_2O\_3 FeO CaO 11\_2O Sp gr at 22 97 02 0 32 3 39 0 55 1 68=102 96 2 520 at 22°

CHESSYLITE or CHESSY COPPER. (Kup ferlusur, Ger) A blue oxycarbonate of copper, 2CuCO<sub>3</sub> Cu(OH)<sub>2</sub>, found native at Chessy near Lyons (v COPPER)

CHESTNUT. The sweet chestnut is the seed of Castanea vesca, the horse-chestnut that of Æsculus hippocastanum.

Average composition:

Pro-Carbotein Fat hydrates Fibre Ash Water Sweet chestnuts. Fresh kernels . Dried kernels 69 80  $\frac{1}{2}.7$ 58 10 6 100 70 9 Spanish 68 1 10 0 93 7.5 27 Horse chestnuts 38 9 4.6 46.2 48 68.3 (air-dried)

Sweet chestnuts are largely used as human food; the horse-chestnut has been tried as cattle food, with success (Gay, Ann Agron. 1896, 22, 401; Auld, J Soc. (hem Ind. 1913, 32, 173)

The kernels of an aquatic plant, Trapa bispinosa, are extensively used as food under the name of water-chestnut, in the North West Provinces of India, being roasted and pounded to a flour Hooper (Pharm J. Trans. 1894, 53, 22) found this flour to contain .

Sugar Starch Cellulose Ash Water Protein Fat and gum 4.7 4284 1.0 14.4 638 36

The ash of the flour contained a little manganese, which, however, was abundant in other parts of the plant.

The bark of the horse-chestnut, especially in early spring, contains asculin  $C_{15}H_{16}O_{9}$ , the glucoside of asculetin  $C_{9}H_{6}O_{4}$ . Another glucoside, argynascin  $C_{27}H_{12}O_{12}$ , occurs in the seeds of the hor-e-chestnut (Rochleder, J. pr Chem 1862, 87, 1)

From the fat-free cotyledons two acids are obtained, viz. asculic acid, fine needles, in p 214°-215°, insoluble in water, forming yellow amorphous salts, and wsculinic acid, goldenyellow needles, mp. 230°-231°, Lavoiotatory, soluble in water, forming yellow salts (Masson, Bull Sci. Pharmacol 1918, 25, 65)

Sweet chestnuts contain small quantities of resins and tannin, and included in the carbohydrates are sugar, dextrin, starch, and other Very diverse amounts of these have substances been found by different observers In the horse-chestnut, according to Laves (Zertsch In the angew Chem 1902, 1013), there are about 50 p c starch, 14 pc cane sugar, 13 pc glucose, and 0 2 p c tannın

CHESTNUT EXTRACT. The wood of the Spanish chestnut, Castanea vesca, though it contains only 3-6 pc of tannin, is the source of the much-valued chestnut extract The bark contains more tannin than the wood (17 pc), but is not much used. The tree, which grows to from 60 to 80 feet in height, is abundant in Italy, the South of France, and Coisica, where it forms immense forests, and it is also very common in America.

Trimble (The Tannins), who very carefully examined the tannin, obtained analytical data and reactions which indicated that it was identical, or nearly so, with gallotannin, but it is probable that this wood also contains traces of a catechol tannin, for a certain quantity of a redcolouring matter is also present, which resembles in character a phlobophane. Some writers have suggested that chestnut tannin is a methyl ether of ordinary gallotannin, but there is apparently no definite evidence in support of this theory

Chestnut is employed almost entirely in the form of extract, the strength of which varies, but usually contains from 26 to 32 pc of tannin. The extract is frequently decolourised, and sometimes mixed with quebracho extract and other materials. Chestnut tannin is the tannin which is most largely employed for the dyeing of silk. Castanea vesca appears to be frequently confused with the horse-chestnut, Esculus hippocastanum The tannin derived from this latter is, however, of little or no practical value (cf. Pollak, Collegium, 1913, 6, 291; J. Soc. Chem. Ind. 1913, 32, 706). A. G. P.

CHIAN TURPENTINE v. OLEO-RESINS

CHICA RED AND CARAJURA. These are rare pigments prepared by the Indians of Central America from species of Bignonia which are very similar in appearance and may contain as a basis the same colouring matter. A red pigment of this character, evidently 'Chica red,' is referred to in The Travels and Researches of Baron Humbolt (Macgilivray, Edinburgh, 1836, p. 229). 'Red paint is the ordinary decoration The most common kind is of these tribes obtained from the seeds of the Bixa orellana and is called annotto, achoute, or roucou Another more expensive species is extracted from the leaves of the Bignonia chica' According to Crookes (Dyeing and Calico Piinting, 1874, 388) chica is obtained from the leaves of the B chica which the Indians boil with water, and add some particles of the back known as 'arvane' to the decanted liquid which causes the precipitation of the colouring matter

An interesting résumé of what is known of this subject is given by Holmes (Pharm. J., 1901), who states that preparations very similar to carajura are employed by the South American Indians in Brazil, Bolivia, and Guiana These include chica from the leaves of B. chica, from 'ula' leaves derived from an unknown species of Bignonia in Bolivia, and a pigment prepared from the heartwood of the B tecoma in Minas (Brazil). The latter, according to Lee (Chem. Soc. Trans. 1901, 284), is prepared by mixing the dust and shavings of this tree with slaked lime and heating the mass with water. From the B tecoma Lee isolated a considerable amount of a yellow crystalline compound, tecomin Bignonia Tecoma.

Carajura or ('rajura was apparently first examined by Virey (Journ de Pharmacie et Chime (3) 5, 154) The sample experimented with, which came from Para, was soluble in alkalis and was precipitated therefrom by acid, and is stated to have been prepared by the Galibis by boiling with water the stem and leaves of an unknown species of Bignonia when the latter in the autumn had acquired a purple

Crookes states (loc cit) that chica is soluble in 36 pc alcohol, in ether and in alkalis with a vinous red colour, and when heated in a scaled tube with alkali and glucose gives a bluish-red liquid which becomes brown on exposure to air and then yields on acidification an orange-red precipitate.

As the result of an examination of chica red or carajura said to be derived from the leaves of the Bignonia chica, Erdmann (Jahres. 1857, 487) assigned to the colouring matter which he isolated by extraction with alcohol the formula  $C_8H_3O_3$ . This substance was soluble in caustic alkali but not in alkali carbonate solutions, and when oxidised with chromic acid gave anisic acid, whereas with nitric acid pieric acid was produced.

Perkin (Proc Chem. Soc 1914, 30, 212), who examined a sample of 'carajura' obtained from Messrs. Wright, Layman, and Umney of London, and which is referred to in the paper of Holmes (loc. cit.), found this to contain a small quantity of the calcium lake of two colouring matters which had either been precipitated on, or intermingled with a substance of the nature of ground

bark or peat After treatment with hot dilute hydrochloric acid, alcohol removes the colouring matters in the form of a resin, and from this, by means of boiling benzene, carajurin is isolated This compound, to which the formula C<sub>18</sub>H<sub>16</sub>O<sub>5</sub> has been provisionally assigned, separates in ruby needles, melting at 204°-206°, is soluble in boiling dilute alkali with a red colour, and is nearly devoid of dyeing properties With mineral acids it very readily yields oxonium salts, crystallising in bright, orange needles, of which the sulphate, probably C18H16O5 H2SO4. H2O is the most stable, the hydrobromide and hydrochloride being decomposed at 100° the hydrochloude a platinichloride can be prepared. Cold acetic anhydride with a trace of pyridine, after two days, gives an almost colourless acetyl compound, crystallising in needles, whereas bromine gives an immediate precipitate with carajurin in acetic acid, which when boiled with this solvent, separates in orange needles Hydriodic acid converts carajuin with loss of 2 molecules of methyl iodide and probably also of a molecule of water into a substance provisionally termed caraquietin iodide, bright scarlet needles, stable in the presence of cold water, and from this by means of cold pyridine, carapuretin C<sub>16</sub>H<sub>12</sub>O<sub>5</sub>, scarlet needles, melting above 330°, and soluble in alkalis with a reddish-violet colour, is produced By dry distillation carajurin evolves a trace of aromatic oil, resembling anisaldehyde in odour, and when fused with alkali, p-hydroxybenzoic acid and a colourless substance, melting at 185°-187°, as yet unidentified, are obtained In many respects carajurin resembles the anhydrohydroxybenzopyranol compounds described by Bulow and Wagner (Ber 1901, 34, 1199).

That portion of the alcoholic extract insoluble in benzene yields to other carajurone, isolated as a scarlet powder, which readily assumes a beetle-green lustre, and possesses strong dyeing properties. Analysis indicates the presence of more oxygen in this compound than in carajurin A small amount of a similar, but brighter lake from British Guiana, and obtained from the leaves of a 'bushrope,' gave a colouring matter dyeing alizaim-like shades. This preparation, considered to be 'chica red,' appeared to differ in some respects from the 'carajura' above described

Bignonia Tecoma. Tecomin.  $(q \ r)$ A G,P CHICAGO BLUE, -ORANGE, v AZO-COLOUR-

CHICK PEA Creer arretinum, I. A leguminous annual, growing to a height of about 2 feet, largely cultivated in India, where its seeds are used as food for cattle and horses, and, after the removal of the husks, for man.

ING MATTERS.

In some cases the young leaves, fried in oil, are also eaten. Church (The Food Grains of India, 1886) gives, as the average composition of the seeds

Water Protein Fat hydrates Fibre Ash With husk 11 2 19.5 4.6 53 8 7.8 3.1 Husked . 11.5 21.7 4.2 59.0 1.0 2.6

Analyses of various parts of the fresh plant were made by Passerini (Staz. Sper Agrar. 21, 20; Chem. Soc. Abstr 1893, ii. 226).

The fresh substance of the stems, leaves, and

entire pods contained 27.54 p.c., 24.39 p.c., and 19 85 p.c. of dry matter respectively.

The dry matter of the stems, leaves, and seeds contained:

Cellu-Other carbo-Protein Fat lose hydrates Ash Nitrogen 49.0 7.81 Stems 6.35 18 35 0 1.62 Leaves 14:21 4.1 139 58.9 2 27 8 83 Seeds 26 20 5.2 17 63.6 3 30 4 19

The ash contained:

Stems 48 4 0.5 23 6 4.4 2.1 0.03 5 3 5 8 6.2 3.7 Leaves 29 6 0 9 40 6 3 3 3 0 -604411318 Seeds 24 6 1 3 4 5 20 0 2 4 - 39 6 3 4 0 7 2 9

Traces of boron, lithium, and copper were

also present in the ash

A detailed analysis of the dry matter of the seeds was made by Zlataroff (Zeitsh. Nahr. Genussm. 1916, 31, 180), who found—fat, 6 30 p c, starch 50·32 p c., fibre 3·62 p c, ash 2·87 p c, total nitrogen 3 34 p c, protein nitrogen 2 11 p c, nuclein nitrogen 0.10 p.c., ammonia nitrogen 010 pc, amide nitrogen 001 pc, amino-acid nitrogen 0.12 p.e.; total phosphoric acid 0.998 p c, inorganic phosphoric acid 0.118 p c, organic soluble phosphoric acid 0.244 p c, legithin phosphoric acid 0 142 pc, phosphoric acid 0.486 p c Oxalic acid (0.007 p.c), citric acid, betaine, choline (0 02 p c), adenine, inositol and a phytosterol were also present

A gum imported into the United States for the manufacture of chewing gum. A sample from Yucatan contained: Acetonesoluble matter (resins) 400; gutta and carbohydrates 174; proteins 06; sand and foreign matter 2 3; water 35 0; ash 4 7 p.c.

For analytical methods of determining its value, see Dannerth, J. Ind Eng. Chem. 1917, 9,

679; J Soc Chem Ind. 1917, 971

CHICORY or SUCCORY. Cichorium Intybus A perennial plant belonging to the Compositæ, growing to a height of about 3 feet, bearing blue flowers and having a long tap root. It is sometimes grown as cattle food, the foliage being thus used, but more generally it is cultivated for the sake of its root, which, when dried, roasted, and ground, is largely used for mixing The blanched leaves are also used with coffee as a salad

According to Letheby's analysis (quoted by Wynter Blyth, Foods, 359, 1909), fresh chicory

roots contains:

Bitter Cellulose, Gummy extrac-tive inulin, Water matter Glucose Fat and fibre Ash 0.6 9.0 0.8 77.0 1.1

Analyses by Mayer (Bied. Zentr 1885, 828) gave:

Bitter Water minoids Fat Inulin Fibre Sugar extract Ash 5.614 72 1.1 0.212.0 1.4 05 to to to to to to 17.3 1.8 60 0.12 1.9

Wolff (Chem. Zentr. 1899, ii 211) also found that chicory root contains from 13 to 15 p.c. of inulin, and an optically mactive sugar, synan-throse. He states that drying the roots has little or no effect upon the sugars, but during | Phosphorus pentoxide 10.0

roasting the reducing sugars are increased, a large portion of the inulin is changed, and caramel and dextrin are formed.

Even by storing a portion of the inulin is converted into inulide and some of the less readily fermented inulides undergo further degradation and are converted into the more readily fermented inulides. Wolff (Compt. rend. 1916, 164, 514) finds a substance which he calls inulo-coagulase in the root, which coagulates the mulin in the expressed juice. It is more abundant in the outer than in the central portions of the root and is destroyed by heating to about 60° It does not coagulate starch, milk or pectin In a later paper, Wolff and Geslin (Compt rend. 1917, 165, 651; 1918, 166, 428) state that inulin breaks down, under the influence of an enzyme present in chicory root, to levulose with intermediate formation of inulides

The bitter principle is believed to be a glucoside of levulose and a pyrocatechuic derivative, probably the aldehyde (Grafe, Biochem. Zeitsch 1915, 68, 1) The determination of cupric reducing power affords a means of detecting the presence of chicory in coffee. The extractives of roasted coffee contain from 19 to 26 p.c of reducing sugars, whilst the extractives of roasted chicory contain from 25.2 to 27.7 pc. When the extractives of a coffee decoction yield more than 3 p c. of their weight of reducing sugars the presence of chicory is indicated. The addition of cane sugar to the coffee does not affect the results and its presence may be detected by the polarimeter

The presence of beetroot in ground chicory may be detected by washing the suspected sample with sodium hypochlorite solution until all soluble colouring matter has been removed, when the particles are to be examined under the microscope If beetroot is present the particles will be seen to contain numerous black cells filled with crystals of calcium oxalate. The cells are oval or elongated and are more numerous near the ligneous vessels of the woody portion of the root (Collin Ann. Falsif. 1916, 9, 271)

Petermann (Bied. Zentr 1883, 843) found

pure roasted chicory to contain:

Soluble in water, 74.2 p c. Dextrin Colouring Water Glucose and mulin Protein matter Ash 16 3 26.1 9.632 16.4

> Insoluble in water, 25 8 p.c Fat Cellulose Protein 3.2 5.7 123 4.6

The ash of chicory root and leaves, according to Wolff, contains:

Root Leaves 60.0 0 7 3.2 14 3 — 9.0 9 0 1.0 1.7

Wynter Blyth (Foods, 359, 1909) gives as the main differences between the ashes of coffee and chicory, the following :-

-	Coffee ash	Chicory ash
Silica and sand .		10·7-35·9 p.c.
Carbon dioxide .	14.9	1.8-3.2
	0.44-0.98	3.1-2.3 ,,
	0.26-1.1	3.3-4.9 ,,
Phosphorus pentoxide		5.0-6.0

The soluble ash of chicory may be taken as about 1.74 p.c., that of coffee as about 3 0 p.c.

Chicory has about three times the tinctorial power of ordinary coffee, and this has been proposed as the basis for a rough estimation of the

amount of chicory present in a mixture of coffee and chicory (Leebody, Chem. News, 30, 243).

The sp gr. at 15 5° of a solution, made by boiling a quantity of the substance with ten times its weight of water, affords a ready method of approximately determining the amount of adulteration The following table gives the specific gravities of infusions made in this way from the various substances :-

Spent tan					1.00214
Acorns					1 00730
Peas .					1 00730
Mocha coffe	90				1 00800
Ceylon coffe	ee				1 00870
Java coffee					1 00870
Costa Rica	coffee				1.00900
Native Cey	lon cof	fee			1.00900
Brown mal	ե				1 01090
Black malt					1 02120
Dandelion i	root				1 02190
Red beet					1.02210
Yorkshire c	hicory				1 01910
Foreign chi	cory				1 02260
Guernsey cl		•			1 02326
Maize .					1 02530
Bread raspi	ngs				1 02630
oted by Wŷr		vth.	Foods.	3	60).

(quoted by Wynter Blyth, Foods, 360).

Hehner and Skertchly propose a determination of furfural (a measure of pentosans) as a means of examining suspected coffee Coffee gives about 5-5 5 p c, chicory about 2 5 p c of pentosans (Analyst, 24, 178) H 1.

CHILE SALTPETRE. Sodium nitrate (v.

CHINA BLUE. Syn Water Blue, Cotton Blue, Opal Blue v. Triphenylmethane colouring MATTERS.

### CHINA CLAY or KAOLIN v CLAY.

CHINA GRASS. Rhea Ramie fibre. This substance is the fibre of two (or perhaps more) species or varieties of the genus Boehmeria, nat ord. Urticaceæ, shrubs allied to the nottle (Urtica), but possessing no stings, namely, Boehmeria nivea (Hook and Arn ) with a white hairy under surface to the leaf, and B. utilis [B. tenacissima (Gaud.)] with a green (Blume) under surface (more or less hairy only on the nerns) to the leaf It is the Tchou-ma of China and the Rheea of Assam. The plants grow to a height of from three to five feet, throwing up numerous straight shoots as thick as the little finger, and covered with short soft hairs. Its leaves grow upon long foot-stalks and are broadly heart-shaped, about six inches long by four broad It is a native of China and Sumatra, where, as in India, it has long been cultivated.

Its manufacture into mats used for making sails, foot-mats, &c., is carried on in Southern China, where three crops of Tchou-ma are taken each year, of which the second is the best. For this purpose the grass is sorted into bundles, which are soaked in water for two or three days and then dried in the sun. If a red colour is desired, the grass is steeped in a decoction of sapan wood; if yellow, in a decoction of the seeds and flowers of the common Chinese plant mi-fa, or in one of Sophora japonica, to which a little alum is added as a mordant. Other vegetable dyes are also employed; but of late years aniline colours have in a measure taken their place.

By processes of manufacture an exceedingly fine fibre is produced on separating the filaments of the stalk—a fibre distinguished by its strength, which is nearly three times that of Russian hemp; by its length, which is greater than that of any bast product; and by its lustrous, silky Great efforts have been made, with more or less success, to render the use of the fibre in European textile manufactures commer-It will be seen from the folcially profitable lowing remarks that there are difficulties in the way In the first place, the method of separating the bast by hand was tedious, and could only be performed whilst the stems were quite fresh, thus limiting its production to the immediate neighbourhood of the plant's growth. In China the bark is stripped from the stems and the fibre scraped off with a knife, a process which removes the whole so-called grass if performed while the stem is still quite fresh, but leaves a large portion if it has become at all dry Even under the best conditions a very skilful operator can collect only two pounds of grass per day, the average quantity obtained being only half a pound to one pound. It is further found that, owing to the large amount of pectinic substances contained in these Urticacca, the long steeping or retting processes used with flax and hemp cannot be practised on account of the ensuing excessive fermentation, which injures the fibre Experiments made in India with unskilled workers gave about four ounces per day each as the result of their labour Attention has consequently been given to mechanical processes for separating the fibres from the bark in the green and in the dry state. With this in view, the India Office some years ago submitted samples of the dried article to Dr Forbes Watson, who reported that he did not find it at all difficult to obtain the fibre from stems which were quite The difficulty in the case of India is that the best crops are obtained during the rainy season, when it is all but impossible to dry the stems except by expensive artificial means, while, as above pointed out, the liability of the plant to ferment resulted in the spoiling of the crops when its treatment was delayed. The Government of India has followed up its inquiries and encouraged inventors to competition in producing a machine capable of dealing with the green crop. It may be said that this encouragement has resulted in an approximation to success; thus, in one instance of a machine consisting of a cylinder with beaters revolving at a high speed against the stems upon which meanwhile a jet of water is poured, Dr. Watson estimated the cost of producing the fibre to average 4l. to 5l per ton The yield from the machine was from 1 to 2 tons of green stalks per day, estimated to produce about 100 lbs. of cleaned fibre.

In the treatment of green stalks by hand the Favier's process consists in subjecting the plants to the action of steam for 15 to 30 minutes, after which the bark is easily stripped. Those stems also which have been standing some time after being cut can also be treated | without leaving any appreciable amount of fibre on the stalks.

The operation of scutching the dried stems presents no difficulty when steam power is applicable and when the quantity of material to be treated is sufficiently large. The next process, however, of 'ungumming' has given a little trouble Since ('hina grass contains from 22 to 28 or, in some cases, up to 35 pc of gum, it is necessary to remove this constituent of the fibre before it can be spun into yarn For this purpose many partly mechanical and partly chemical processes have been devised. It may sufficiently describe the lines which these various systems take to say that they resemble generally the processes adopted in the scouring and bleaching of cotton goods—that is to say, by subjecting the material to pressure under the application of steam, and to alternate treatment of alkaline and acid agents

American inventors have devised means by which the fibre can be spun in ordinary cotton machinery. Rossi, of Naples, has patented a method by which China grass can be retted without chemical agents. When chemically degummed name is manufactured in combination with cotton and dyed and printed, the goods possess a silky lustre which enhances the

colours on both fibres

China grass has little or no attraction for dyes, and consequently the methods adopted in the dyeing of cotton or of jute are applicable to it It is found that mineral colours, such as chrome yellow and orange, destroy the peculiar lustre of China grass, the yellow dye obtained with cadmium sulphide, however, is not open to this objection Alizarin mordanted with Turkeyred oil destroys the lustre, whilst aniline colours fixed with the same mordant do not Rosaniline derivatives and azo- colouring matters may be It is recommended that the used satisfactorily grass should be dyed immediately after retting and before spinning, to preserve the characteristic sheen

The textiles formerly associated with China grass were such articles as handkerchiefs and grass cloths,' imported direct from China, but recent efforts to introduce the manufacture as well as the growth of China grass into Europe have led to other developments Thus at the manufactory of Zittau in Saxony, the fibre has found application in tapestry and dress goods, table-cloths, damasks, lace, fancy knitting yarns, It is also used in the manufacture of incandescent gas-mantles

A number of other species of Boehmeria also supply bast fibres

CHINA INK v. INDIAN INK CHINAPHENIN. Trade name for phenetidine-quinine carbonic ester Formed by treating p-phenetidine with phosgene and combining the product with quinine

CHINA-STONE. A granitic rock, having its felspathic constituent more or less decomposed, but not completely kaolinised, and with an absence of coloured minerals, such as biotite and tourmaline. As the felspar still retains part of its alkali, and as some micaceous mineral is generally present, the stone is fusible; hence it is largely used in the manufacture of porcelain, and when free felspar is not introduced

it forms the only vitrifiable constituent of the China-stone is sometimes known as 'Cornish stone' in consequence of its being largely quarried in Cornwall, where it was discovered by W Cookworthy, of Plymouth, about 1750. He first noticed it at Tregonning Hill. near Breage, and described it under the local name of 'growan' or 'moor-stone' It is now obtained principally from the neighbourhood of St. Stephens, near St Austell In this area it occurs as a local modification of the granite, and is sharply marked off from the normal granite and from the china-clay rock. It is too hard to be worked like the china-clay rock, and has to be blasted and quarried like ordinary granite, in fact it has been used locally as a building stone The rock requires no preparation, but is sent direct from the quarry to the potteries (For an account of the occurrence of china-stone in Cornwall, see Mem Geol. Survey, Bodmin and St Austell District, 1909; and the Handbook by J A Howe, quoted below)

The following commercial varieties are

recognised

I 'Hard purple,' a hard, white rock with a faint purplish tinge due to the presence of fluor-spar
2. 'Soft' or 'Mild purple,' similar to the

last, but softer

3 'Dry white stone,' a soft, white variety.

4 'Bufl stone,' similar to the last, but iron stained

The following analyses by W Pollard and E G Radley are given by J A. Howe, A Handbook to the Collection of Kaolin, China-clay and China-stone in the Museum of Practical Geology, London, 1914 I 'Hard purple' china-stone from Goonvean, near St Stephens, 'Buff' china-stone from the same locality; III China-stone from Jersey ('Jersey stone')

		I	II.	III
$S_{1}O_{2}$		. 72 28	$73 \ 18$	77.06
$T_1O_2$		. 0 05	0.06	0 08
$\Lambda l_{\lambda} \tilde{O}_{\lambda}$		. 14 90	16 13	12 22
Fe,O,		. 0 50	0.52	0 47
Mn(O		. 0 01	0 02	0 19
CaO		. 166	0 61	0.16
MgO		0 15	0.14	0.19
K,0		. 5 25	4 41	5 07
Na,O		. 3 01	2 18	3 23
Lı,O		0.02	0.02	trace ?
H,O (a	t 105	°). 013	0 20	0.39
H,O (o			1.81	0.37
P2O5		. 0 53	0 45	0 25
$\mathbf{F}^{2}$ .		. 088	0 23 F	'eS, 0 30
Čl.		. 0 09		0, 0 04
		100 07	99 96	100 02

Jersey stone is a decomposed partly kaolinised granite quarried in Jersey Granite-pegmatite in France, granite-aplite at Meldon, near Okehampton, Devonshire, porphyrite in Saxony; liparite in Japan; and felsite in China are all rocks that are applied for the same purpose

It has been proposed by J. H. Collins (Hensbarrow Granite District, Truro, 1878) to distinguish china-stone under the name petunzite, a word suggested by the Chinese pe-tun-tse. In the early part of the 18th century a Jesuit missionary named D'Entrecolles, residing at King-te-chin, sent to Réaumur, in Paris, samples of the raw materials used in China. These were the first specimens of their kind that had ever reached Europe, and they led to the foundation of the manufacture of porcelain at Sèvres. The china-stone was described by D'Entrecolles as po-tun-tse, and the china-clay as kaolin; but it appears that the former name, if not both, must have been erroneously applied. Ascording to the Chinese scholar, Stanislas Julien (Hist et Fabric, de la Porcel. Chin Paris, 1856, preface xx.), pe-tun signifies 'white paste,' while the isemerely a diminutive applied to the material when worked up into small cakes. Hence, petun-se is really the name of small blocks of white clay or prepared paste, and not of the granitic rock. Nevertheless, custom in this country has justified the use of the term pe-tun-tse as synonymous with china-stone.

CHINEONAL. Trade name for a compound of quinine and veronal Used as a hypnotic

CHINESE BLUE. (Porzellanblau, Ger) Several compounds are sold under this name, of which the following are examples: ultramarine and flake white; cobalt blue and white lead; and a double cyamide of non (Prussian blue) (v. Pigments).

CHINESE GREEN OF LOKAO V LOKAO CHINESE LAKE OF SCARLET LAKE V PIGMENTS.

CHINESE VEGETABLE TALLOW is the hard fat which coats the seeds of the Chinese tallow tree, Stillingia sebifera (Willd.), Stillingia sinensis, Croton sebiferum, Sapium schiferum, (Roxb), Excecaria sebifera (F Mull), Carumbium schiferum), a tree growing wild in China, and largely cultivated there, as also in Indo-China and the North of India, where the tree thrives luxu-In China the stillingia tree is exploited for both its leaves and seeds. In Tonkin the tree is only cultivated for the sake of its leaves, a decoction of which furnishes a colouring matter used for dyeing silk In China the fat is only a by-product. The tree commences to produce at the age of 4-5 years, and when it has reached its full development it yields about 24-30 kilos of seeds a year. The fruit yields about 30 pc of a mixture of vegetable tallow and of stillingia oil, half of this quantity consists of vegetable tallow alone, and the other half of stilling a oil, the yield of the latter from the kernels being about 62 p c. The fruit consists of three oval seeds surrounded by a thick tallowlike mass, the true vegetable tallow. This is recovered by placing the seeds in perforated cylinders and steaming them, so that the fat melts and runs off. Another method is to pass Another method is to pass the seeds between fluted rollers, which scrape off the outer fat coating without breaking the seed proper The scraped-off mass is pressed in the hot primitive wedge presses, and sold under the Chinese name of 'pi-ieou' or 'pi-yu'; it is known in Europe as 'prima' vegetable The seeds which have been bared from their fat coating, are then crushed separately and yield the stillingia oil, a drying oil known in China as 'tsé-tiéou' or 'ting-yu.' In another process, the seeds, together with their coating of fat, are crushed, and thus a mixture of vegetable tallow and stilling is oil is obtained. • The product Vol. II.-T.

thus prepared is naturally much softer than the true vegetable tallow, and has therefore a lower melting-point and a much higher rodine value. This second quality is sold in Europe as 'secunda' vegetable tallow, and represents a mixture of 'pi-yu' and 'ting-yu.' It is sold in China under the name 'mou-feou'

The saponification value of commercial samples of vogetable fat varies, therefore, according to the mode of preparation, from 179 to 207 Similarly, the iodine value varies considerably, and numbers ranging from 19 to 32 have been recorded. The composition of vegetable tallow has not yet been definitely ascertained Stearie acid is absent (Hehner and Mitchell), and it is probable that the fat consists in the main of palmitin and olem.

The oil has an iodine value of about 155, and yields 20 p c of an insoluble bromide (m p 147°), whereas the fat does not form an insoluble compound with bromine

Vegetable tallow plays, in China, an important rôle as an article of commerce, it is largely used in the manufacture of candles. During recent years considerable quantities have been exported. The bulk of the vegetable tallow is used in the candle and soap industries. According to Diedrichs (Zeitsch Nahr Genussm 1914, 27, 132) the fat, if properly purified, can be used for food, but soon acquires a bitter taste and unpleasant odour. It is frequently adulterated by the Chinese.

J L.

CHINESE WAX or INSECT WAX v WAXES CHINESE YELLOW (King's yellow) v Pig-

**CHINOCOL.** Trade name for a compound of hydroxyquinoline and potassium sulphate, used as an antiseptic and disinfectant v

CHINOFORM. Trade name for quinine formate

CHINOLINE v QUINOLINE

**CHINOSOL.** Trade name for the potassium salt of 8-hydroxyqunoline-5-sulphonic acid  $C_9H_6\mathrm{NOSO}_3\mathrm{K},H_2\mathrm{O}$ , a pale-yellow crystalline powder with a saffron-like smell and a burning taste; in p 175°-1775°, readily soluble in water, sparingly soluble in alcohol, insoluble in ether, with ferric chloride gives an intense green colour, yellowish needles with copper salts, and a white precipitate with barium chloride. Used as an antiseptic. At a concentration of 1 in 1000 its solutions possess as great a bactericidal action as mercuric chloride solutions of the same strength.

CHINOTROPIN. Trade name for urotropine quinate.

CHINOVOSE v CARBOHYDRATES.

CHIOLITE. A double fluoride of aluminum and sodium 3AlF<sub>3</sub> 5NaF, closely akin to cryolite AlF<sub>3</sub> 3NaF, but crystallising in the tetragonal system. Found at the Ilmen Mountains in the Southern Urals and with crystals in Greenland.

L. J. S.

CHIRETTA. Chirata, BP Is the plant Swertia chirata (Buch. Ham.) collected when in flower and dried Japanese chiretta is Swertia chinensis (Franchet). Hohn (Archiv. d. Pharm. 1869, 215) found two bitter constituents in Indian chiretta, viz. chiratin and ophelic acid.

CHITENINE. An oxidation product of

•

178 CHITENINE.

quinine, found in the urine after the administration of quinine. Crystallises from dilute alcohol in prisms, m.p.  $281^{\circ}-282^{\circ}$  [a] $_{\bar{\rm D}}^{17}-122$  6°. Chitin, Chitose v. Carbony drates.

CHLOANTHITE. Native nickel arsemide,  $N_1As_2$ , isomorphous with smaltite (CoAs<sub>2</sub>) (q.v), there being no sharp line of demarcation between the two species. Found as cubic crystals and compact masses at Schneeberg in Saxony and Riechelsdorf in Hesse, where it was formerly mined as an ore of nickel. L J. S.

CHLORACETIC ACIDS v. HALOGEN ACETIC ACIDS

CHLORAL Truchloracetaldehyde CCl, CHO

Chloral was first prepared by Liebig, who obtained it by the continuous chlorination of absolute alcohol (Annalen, 1, 189) It was further examined by Dumas (Ann Chim Phys [2] 56, 125), who determined its composition, and by Stadeler (Annalen, 61, 101) Although a chlorinated aldehyde, it cannot be prepared by treating acetaldehyde with chlorine, unless water is present together with calcium carbonate to neutralise the hydrogen chloride formed, since butyric chloral results when dry materials are employed (Pinner, Ber. 4, 256)

Industrial preparation. The process by which chloral is manufactured is divided into three stages the preparation of chloral alcoholate, conversion of this into chloral, and the

formation of chloral hydrate

In a large glass balloon 60 kilos of alcohol, completely dehydrated as possible, are subjected to the action of a current of chlorme, the escaping hydrochloric acid being absorbed by water. The introduction of the chlorino lasts a considerable time, in some works 10 to 14 days At first the balloon must be cooled, for, if the temperature is not properly regulated, the yield is much reduced After a time the reaction becomes less violent, the mixture is then warmed to 60°, and finally to 100° When the liquid is completely soluble in water, the current of chlorine is stopped, and on cooling, a white mass of the alcoholate is obtained. In this operation Springmuhl recommends the use of iodine as a chlorine carrier, Page, the addition of crystallised ferric chloride (Annalen, 1884, 225, 209), and in either case the ethyl halide is easily recovered.

In the ordinary process the chlormated liquid is not allowed to cool, and is transferred to an enamelled still holding 150-200 kilos, where an equal volume of strong sulphuric acid is added in small quantities. The temperature is raised to boiling, the condensed vapours being returned to the still During the chullition there is an evolution of hydrochloric acid gas, the cessation of which indicates the termination of the decomposition of the alcoholate Fractional distillation follows, this ceasing when the thermometer has reached 100°. The distillate is again rectified, the portion coming over above 94° being pure chloral. In some works the final purification is carried out by distilling over chalk or sodium bicarbonate. Ethylene dichloride, ethylidene chloride, and chlorinated ethylene dichloride (b p 115°) are also formed as by-products in this reaction (Kramer, Ber 3,

quantity of water and the product of the reaction at once poured over a smooth surface, when it solidines; in this state it is sent into commerce To obtain crystals the warm mass (heat is disengaged in this reaction) is mixed with one-third of its volume of chloroform, and the mixture allowed to cool in closed vessels; crystallisation is complete in a week mother liquers serve for a second crystallisation, and the crystals are dued at the ordinary Fluckiger temperature (Zeitsch. [2] 6, 432) recommends crystallisation from warm carbon disulphide or turpentine, and Martius from benzene Other methods, which have been described, differ from the pieceding in that the chlorinated mixture, after solidification and pressure, is sublimed or distilled with calcium chloride (Detsényi, Dingl poly J. 209, 224; Trillat, J Pharm Chim 5, 218).

The following method has been devised for the continuous production of chloral and its hydrate are the most volatile products of the action of chlorine on alcohol, and may therefore be removed by continuous distillation. A special apparatus is described; chlorine and chlorinated alcohol are introduced into a vessel heated in a water-bath, from which the vapours pass into a second and then into a third heated vessel The residual chlorine is absorbed in a cylinder provided with bafile plates rectifier, which also acts continuously, is provided with baffle plates, it is fed at the middle with chloral, and at the top with sulphuric acid (66° B) The final rectification is carried out as already described The same apparatus may be used for converting chloral into chloroform (Besson, Eng Pat. 17202, U.S. Pat 774151; J. Soc Chem Ind 1901, 1139)

Chloral is formed when ethyl alcohol is allowed to drop into the anode compartment of a cell in which an alkaline chloride is being

Theories to explain the action of chlorine on alcohol and the subsequent production of chloral have been proposed by Lucben (Ber 3, 910), Wurtz and Vogt (Compt rend 74, 777), Jacobsen (Ber 15, 601), and Brochet (Bull. See chim

17, [3] 228)

Chloral is a colourless pungent liquid, which solidities at -75° (Berthelot, Ann Chim Phys [5], 12, 536), at --57 5° (Van Rossem Zeitsch. physikal Chem 1908, 62, 681), and is readily soluble in water, alcohol, and other Its b p is 97 7° (corr) and sp gr 1 54175 at 0° (Passavant, Chem. Soc. Trans 39, 55, Thorpe, thid 37, 191), bp 96°-97° at 750 mm, and sp gr 1 5121 at 20°/4° (Bruhl, Annalen, 203, 11), bp 97° at 740 mm (Van Rossem, tc.) Three hydrates exist (1) a semihydrate 2C<sub>2</sub>HCl<sub>3</sub>O,H<sub>2</sub>O, true m p is about 49°, the highest observed 80°; (2) the ordinary monohydrate existing in two pseudo-syminetrical modifications: true m p 47 4°, highest observed 54°, (3) a heptahydrate, mp. -14° (Pope, Chem Soc Proc 1896, 142; and Van Rossem, l.c). According to Wolf (J. Phys. Chem. 4, 21); there is only one modification, the difference in melting-points being due to dissociation Nascent hydrogen reduces it to aldehyde (Personne, Annalen, 157, 113), which it closely resembles in reducing ammoniacal silver nitrate solution (Stadeler, Annalen, 106, 253), and in The chloral is shaken with the theoretical forming crystalline compounds with sodium CHLORAL 179

disulphite (ibid); with ainmonia (v. chloral | Weiner Med. Presse, 1899, 1361) ammonium); with hydrogen cyanide (Bischoff and Pinner, Ber 5, 113; Annalen, 179, 77), with amides (Wallach, Ber 5, 251; Sulzbeig, DRP. 198715; J. Soc ('hem Ind 1908, 958), and with amines (Wallach, *lc*, Rugheimer, Ber 1906, 1653; Wheeler, Amer Chem. J 1902, 1063, 1908, 136, 1909, 937).

When heated with aqueous alkalis it decomposes into chloroform and an alkaline formate. and a similar reaction occurs when alcoholic potash or soda is employed (Kekulé, Annalen, 119, 188) Chlorine, bromine, or rodine and aluminium chlorido convert chloral into hexachlorethane (Mouneyrat, Bull Soc chim 17, [3] 796) Furning nitric acid converts it into trichloracetic acid, and, according to Grabowski (Bei 6, 225), fuming sulphuric acid combines with it in the cold, but on heating converts it into chloralide C<sub>5</sub>H<sub>2</sub>Cl<sub>6</sub>O<sub>3</sub> (compare Grabowski, Ber. 8, 1433, Stadeler, Annalen, 61, 114, Wallach, ibid 193, 1) Chloral condenses with succinic acid by means of acetic anhydride to form tuchloromethylparaconic acid (Fittig and Miller, ibid. 255, 43)

Derivatives Chloral ammonium C<sub>2</sub>H<sub>4</sub>Cl<sub>3</sub>NO is prepared by the action of ammonia on a solution of chloral in chloroform (Schiff, Ber. 10, 167) It is a white crystalline solid, resembling chloral hydrate in taste and colour; it melts at 64°, and is slightly soluble in water, more readily soluble in alcohol (Personne, Annalen, 157, 113, Compt rend 71, 228) Nesbit recommends the use of chloral ammonium instead of methane and chloral hydrate, since the action of chloral hydrate on the respiration centres and the heart is retaided by the introduc-(J Soc. Chem Ind 1889, 413, 1890, 889) By heating chloral ammonium on the water-bath, chloroform and ammonia distil off, and the residue consists of formamide and chloralimide C2H2Cl3N, the latter of which is extracted with alcohol Isochloralimide is also formed, and according to Schift (Gazz chim ital 21, 490), dimolecular and trimolecular imides exist Chloralimide can also be prepared by heating chloral and chloral ammonium until chloroform is no longer evolved, and by the reaction between chloral hydrate and ammonium acetate (Schiff,  $\ell\,c$  , Pinner and Fuchs, Ber 40, 1068 , Årends, Chem Ind 16, 78, Béhal and Choaz, Compt rend. 109, 817, 110, 1270). It has remarkable antipyretic and analgesic properties chloralimido compounds have been described by Moschelles (Bei 24, 1803)

Chloralamide or chloral formamide C3H4Cl3NO2 is produced by gently heating chloral and formamide in molecular proportions (J F von Mering, Eng Pat 7391; D R P 50586; J Soc Chem Ind 1890, 413). When recrystallised, it forms colourless crystals melting at 115°-116°, it is soluble in 9 parts of water, or in 11 parts of 96 pe alcohol It is without smell, and has a slightly bitter taste, when heated above 60° it is decomposed; aqueous or alcoholic solutions are unaffected by weak acids or by silver nitrate, but are rapidly decomposed by alkali hydroxides, more slowly by alkalı carbonates. As a hypnotic it has been proved superior to chloral hydrate, and is of most value in cases of nervous insomnia and similar disorders (Hagner and Hubner),

It has no corroding action, does not affect the digestion, nor disturb the heart. It may be taken internally as a powder, prepared by triturating 1-3 grams of the amide with 1 gram of fenneloil sugar (Ellwosaccharum Fæniculi) (Merck's Bull. 1889, 85)

Chloralose C<sub>8</sub>H<sub>11</sub>Cl<sub>3</sub>O<sub>6</sub> is obtained by heating equal parts of chloral and glucose, both anhydrous, on the water-bath. The vitreous mass is boiled with water to remove the chloral, and the concentrated aqueous solution deposits a crystalline compound, from which chloralose is extracted by alcohol. On treating the residue with other, a compound isomeric with chloralose, termed parachloralose, is obtained. Chloralose melts at 187°, and is a powerful hypnotic, whilst parachloratose melts at 227°, and possesses no hypnotic properties. By oxidising these compounds with nitric acid, followed by potassium permanganate, the corresponding chloralic acids permangulate, the corresponding emoralic acids are obtained, in p. 212° and 202° respectively (Hanriot and Richet, Bull Soc chim [3] 11, 37, 12, 303, Compt rend. 116, 63, 122, 1127, 148, 1265; Meunier, ibid. 122, 142, Petit and Polonovski, Bull. Soc. chim. 3, [3] 861)

Chloral-acetonechloroform  $C_nH_n()_{\downarrow}Cl_n$ prepared by heating molecular quantities of chloral and acctonechloroform. It melts at 65°, and has a faint odour and taste resembling camphor (Hoffmann-Laroche, US Pat 778277; J Soc Chem. Ind 1905, 150).

Chloral-bromal-carbamide is obtained by mixing usea with molecular quantities of chloral and bromal, or equivalent quantities of their hydrates in the presence of concentrated hydrochloric or sulphuric acid (Kalle & Co. D. R. P. 128462, Chem Zentr 1902, 1 547). Chloral hydrate in saturated aqueous solution combines with utotropine and caffeine, forming in each case two compounds according to the proportions of the reagents employed

Monochloralurotropine, CaH12N4, C2HOCl3, H2O, crystallising in Thombohedia, and dichloralwrotropine, C6H12N4,(C2HOCl3,H2O),, crystallising in needles, resemble one another in their properties They are colourless and odourless, and very soluble in alcohol and water volatilise at about 100° without melting. They are both neutral substances, which reduce copper solutions and ammoniacal silver nitrate mineral acids they yield formaldehyde, and with alkalis the chloral is attacked, giving chloroform in the cold and carbylamine on heating.

Dichlorakafferne,

$$C_5H_{10}O_2N_4, H_2O_7(C_2HOCl_3, H_2)_2$$
,

mp 72°-73°, slowly loses 1 mol. of chloral and passes into monochloral cafferne,  $C_8H_{10}O_2N_4$ ,  $H_2O_3C_3HOCl_3$ ,  $H_2O$ , in p. 92° 93°. The latter compound dissolves in water, and in solution slowly dissociates, the caffeine separating out. In hot alcoholic solution it is completely dissociated When heated at 100° all the chloral is driven off, leaving pure caffeine (Leulier, J. Pharm Chim 1912 [vii ] 6, 18)

The following condensation products of chloral have been prepared: chloralacetone, chloralacetophenone, and chloraldol (Gigli, Gazz. chim. ital. 28, [2] 83; Konigs, Ber. 1892, 792; 1893, 554; Wislicenus, Kircheisen, and Sattler, 180 CHLORAL.

vorm. F. Bayer & Co. D. R. P. 121223, Chem. Zentr. 1901, n. 69); chloral mirte (Bertom, Gazz. chim ital. 24, n. 20); the compounds with aminoazobenzene (Betti, vbid. 28, 241); with amidooxybenzoic esters (Kalle & Co. D. R. P. 112216; Chem Zentr 1900, n. 791); with aromatic hydrocarbons (Biltz, Ber 1893, 1952; Dinesmann, Compt rend. 1905–141, 201); with hexamethylenetetramine (Lederer, Eng. Pat. 17693, J. Soc Chem Ind. 1897, 1039), with hydroveratine (Frankforter, Amer Chem. J. 20, 358); with methylpropanoldialkamino-acids (Poulene Frères & Fourneau, Fr. Pat. 382787, D. R. P. 203643, J. Soc Chem. Ind. 1908, 246), with menthol and terpinol (Wagmitz, Eng. Pat. 19103; J. Soc Chem Ind. 1899, 856, Schmitt, Compt. rend. Soc. Biol. 1890, 678); and with resoreinol (Causse, Bull. Soc. chim. 3, [3] 861, Hewitt and Pope, Chem. Soc. Trans. 1896, 1265)

Polymerides. Two polymerides of chloral are known, an insoluble and a soluble substance. When pure, chloral is perfectly stable, but traces of impurities, and particularly of sulphuric acid (Byasson, Compt rend 91, 1071) convert it into metachloral, a white amorphous solid It may be prepared by leaving chloral hydrate in contact with half its weight of concentrated sulphune acid for several hours, until a pasty mass is After pouring off the acid, the residue is introduced into a well-cooled solution of hydrochlorie acid, and the product, which now becomes solid, is washed and dired over sulphuric acid. Chloral alcoholate may be substituted for chloral hydrate in this preparation (Gartner, D R P 170534, Chem Soc Abstr. 1906, i. 628) Another method is to add 7 parts of anhydrous aluminium chloride to 100 parts of well-cooled anhydrous chloral, the temperature being kept below 40° In the course of an hour the whole mass solidities In the After 24 hours the hard mass is broken up, extracted with dilute hydrochloric acid, washed with alcohol, and dried at a moderate temperature (Erdmann, D R P 139392, Chem Zeit 27, It is insoluble in alcohol, ether, water, (20) 223) and acids, but soluble in sodium carbonate solution On warming with alcohol, it yields chloroform, and on distillation at 180°-250° is reconverted into chloral (Kolbe, Annalen, 54, 183 [footnote]) It is tasteless, and can be employed as an anæsthetic, unlike chloral hydrate, it has no irritating effect on the mucous membrane of the stomach.

The soluble polymeride is obtained by treating chloral with pyridine or any other amine in the cold, and then acidifying. It is a slightly volatile anæsthetic substance, volatilising without melting, converted by hot water or hot alcohol into chloral hydrate and chloral alcoholate respectively, and decomposed by alkalis into chloroform and formic acid. It is soluble in water (Gartner, D. R. P. 165984; U.S. Pat. 768744, Chem. Soc. Abstr. 1906, 1.

Chloral hydrate CCl<sub>3</sub>·CHO+H<sub>2</sub>O or CCl<sub>3</sub>·CH(OH),.

Properties.—Chloral hydrate crystallises in monoclinic tables (Groth, Ber. 5, 676), and has a peculiar odour and sharp taste Its m.p. is 57° (Meyer and Dulk, Annalen, 171, 75),

47.4° (Van Rossem, Zeitsch. physikal. Chem. 1908, 62, 681); b p 97.5°; and sp gr. 1.818 (m powder) and 1848 (crystallised) (Schroder, Ber. 12, 562). It dissolves readily in water, 1 part of water dissolving 4.7 parts of chloral hydrate at 17.5° (Mauch, Arch Pharm 240, 113), a label of the schloreform turrentum and light alcohol, ether, chloroform, turpentine, and light petroleum, and its solubility in carbon disulphide is given by Fluckiger (l.c.) as 1 in 45 at 156-18° and 1 in 4 5 at the boiling-point Concentrated sulphuric acid decomposes it into chloral and water, and alkalis act upon it as upon chloral A modification of the hydrate, melting at 80°, is obtained by rapidly evaporating an acetic acid solution of anhydrous chloral over sulphuric acid (Meyer and Dulk,  $l\,c$ ), this is probably the semihydrate described by Van Rossom (l.c) Under the influence of light and air, chloral hydrate is decomposed, the decomposition products being hydrochloric acid and carbon dioxide or monoxide, or water, chlorine, and carbon dioxide, the amount of available oxygen determining the products (School and Van den Berg. Ph Weekblad. 43, 42).

Chloral hydrate should be free from alcoholate. which has not the same physiological properties, and when warmed with twice its volume of water it should give a clear solution, free from oily drops and not precipitated by silver nitrate When carefully heated, it should volatilise completely, and the vapours should not be inflammable, when gently warmed with 3 vols of strong sulphuric acid the mixture should remain On warming chloral hydrate with potassium hydroxide, turbidity is caused through the separation of chloroform, the clear portion is decanted, and rodine dissolved in potassium rodide added So soon as the liquid begins to turn yellow, it is cooled, when the presence of alcohol and consequently of chloral alcoholate, is indicated by the formation of a precipitate of 10doform (Trillat, J. Pharm. Chim 5, 218). Chloral alcoholate may also be detected by the Zeisel method for the estimation of ethoxyl (Schmidinge, Monatsh 21, 36) Chloral hydrate is converted into alcoholate when dissolved in alcohol, and similarly, chloral alcoholate is very slightly soluble in water, dissolving slowly to form chloral hydrate Potassium carbonate is a good antidote to chloral poisoning, if it has not reached the blood (Klar, J Soc Chem. Ind. 1896, 555)

Physiological Action — Compare Liebieich (Ber 2, 269); Personne (Compt rend 78, 129), Byasson (whid 78, 649), Tomaszewicz (abstracted, Chem Soc Tians 1874, 814), Tanret (J Pharm Chim [4] 20, 355), and Trillat (lc). Mering, Zeitsch, physiol Chem 6, 1882, 480. According to the last-named authority is reduced to trichlorethyl alcohol (CCl<sub>3</sub> CH<sub>2</sub> OH) in the body and not into chlorofoim, as suggested by Liebieich, who first proposed its use as an hypnotic.

Estimation.—4 grams of the sample are treated with 30 c.c. of normal caustic soda in a stoppered bottle, and shaken vigorously for a few minutes without the application of heat. On titrating back with normal sulphuric acid, using litmus as an indicator, 5.8 c.c. should be required for complete neutralisation (Alcock and Thomas, Pharm. J. 63, 236; and Sargeant, ibid.). Owing to the secondary reaction CHCl<sub>3</sub>+3NaOH = 3NaCl+CO+2H<sub>2</sub>O, this titration with acid is

CHLORAL. 181

slightly inaccurate. Modifications of the method have been devised by Trillat (l.c.); Hinrichs (Pharm. J. 70, 530); Desgrez (Compt. rend. 125, 780; Analyst, 23, 76); Self (Pharm. J. 79, 4); and Garnier (Chem. Zentr. 1908, 1. 1492). Another method is to heat 03 gram of the sample with 1 gram of aluminium powder, or 2 5 grams of zine filings, 15 c c of glacial acetic acid, and 40 cc of water under a reflux condenser for half an hour. The mixture is filtered and washed, and the chlorine estimated volumetrically or gravimetrically as silver chloride (Self, l.c.). Wallis (Pharm J 76, 162) treats the chloral hydrate with caustic soda in a bottle, and then estimates the chlorine as silver chloride It may also be determined indometrically (Rupp, Arch. Pharm. 241, [5] 326)

Chloral hydrate may be detected by means of the green colour formed on the addition of antimony trichloride to a solution in castor oil (Covelli, Chem Zeit 31, 342); or by the colours formed when it reacts with resorcinol in the presence of sulphuric acid (Jaworowski, Zeitsch anal Chem 37, 60) Another method is to add a solution of ammonium sulphide, when a pinkish or yellow precipitate is obtained, and an oil having the odour of walnuts (Lesinsky and Grundlich, Amer Chem. J. 19, 603). It may also be detected by the reducing action of the formic acid into which it may be converted In presence of compounds containing the CCl, group, eq chloroform and trichloracetic acid, the substance dissolved in water and slightly acidulated with sulphuric acid is treated with zinc in the cold When the evolution of hydrogen has ceased, a strip of filter paper, soaked in a solution of sodium nitroprusside and 5 pc piperidine solution is hung over the liquid, which is gently warmed, when the acetaldehyde formed colours the paper blue 1 part of chloral hydrate in 20,000 may thus be detected Chem 1912, 61, 57) (Jona, Giorn Farm Chloral hydrate is valuable for extracting alkaloids from plants containing much resin, essential oils, &c (Schaer, Zeitsch anal Chem. 37, 469), it is used in rubber analysis (Weber, J. Soc. Chem Ind. 1903, 576), in microchemical investigations (Schaer, Chem Soc Abst 1908, 1 62); and in qualitative analysis (Mauch, Arch Pharm 240, 166)

Derivatives - Chloral antipyrine is obtained by the condensation of chloral hydrate with antipyrine (Béthal and Choaz, Ann Chim Phys. 27, 319).

Chloral caffeine is formed when chloral hydrate and caffeine are brought together in aqueous or alcoholic solutions in molecular pro-It is recrystallised from warm water at 30°, and owing to its easy solubility as opposed to the difficult solubility of caffeine, is of medicinal value (Chem Fab auf Act vorm E Schering, D R P. 75847; J. Soc Chem. Ind. 1895, 987)

Chloral tannin is prepared by the condensation of chloral hydrate with tannin in the presence of sulphuric acid. It is a greyish-brown amorphous powder, having valuable dermatological properties (Farb vorm Fried. Bayer & Co. D. R. P. 98273; Eng Pat. 2882; J. Soc Chem Ind. 1899, 172).

Chloraldoxime is prepared by treating chloral hydrate (1 mol.) with hydroxylamine hydrochloride (4 mols) and a little water; mp

39°-40° (Meyer, Annalen, 264, 116; Tarugi, Gazz. chim. ital. 21, ii. 6)

Chloralhydroxylamine is obtained by triturating molecular quantities of chloral hydrate. sodium carbonate, and hydroxylamine hydrochloride, dissolving the mass in a little water, and extracting with other, mp 98° (Hantzsch, Ber 1892, 701)

Compounds with formaldehyde are described by Pinner (Ber 1898, 1926)

Chloral alcoholates. The preparation of chloral ethylalcoholate has already been described It may also be obtained by the action of chloral on absolute alcohol (Martius and Mendelssohn, Ber 3, 444) It crystallises in needles, melts at 44°-46° (Meyer and Dulk, Annalen, 171, 78), at 46° (Lieben, Ber 3, 907), boils at 115°; is sparingly soluble in water, more soluble in absolute alcohol, I part of alcohol dissolving 4 4 parts of chloral alcoholate at 17 5° (Mauch, Arch Pharm 240, 113) Sulphuric acid decomposes it into chloral and alcohol Chloral propyl alcoholate, b p 120°-122°, and chloral \*\*opropylalcoholate, m p 47°, and b p 108°, have been described by Gabutti (Gazz chim ital 31, 1. 86), and Vitoria (Chem Soc Abst 1905, i. 110), chloral amylalcoholate by Gadamer (Arch Pharm 243, 30), and Kalle & Co (D R P 115251; Chem Zentr 1900, 11 1141) Other alcoholates have been prepared by Jacobsen (Annalen, 157, 224), and Kuntze-(Arch Pharm 246, 91)

Butyl chloral CH, CHCl CCl, CHO; more correctly, tri-chlor-butyric-aldehyde; b p 164°-165° at 750 mm; sp gr 1 3956 at 20°/4° (Bruhl, Annalen, 203, 20) It is a colourless oil, having a characteristic chloral odour, formed by the action of chlorine upon ordinary aldehyde or paraldehyde (Kramer and Pinner, Ber 3, 383) Chloracetaldehyde CH\_Cl CHO is first formed, and this condenses with another molecule of acetaldehyde, forming a-chlorerotonaldehyde CH, CH CCI CHO and water, and this, by the action of more chlorine, forms the butyl chloral.

Preparation —Chlorine is led through cooled paraldehyde (or aldehyde cooled by a freezing mixture), water is then added, and the liquid neutralised with calcium carbonate The mixture is then distilled on the oil-bath in a current of steam, and the hydrate is thus obtained. This is then purified by recrystallisation from water, and decomposed by distillation in a stream of hydrochloric acid gas (Pinner, Annalen, 179,

Butyl chloral is converted by fuming nitric acid into aaß-trichlorbutyric acid. Phosphorus pentachloride converts it into a body C4H4Cl4, which boils at 200°. It combines with zincethyl to form a compound which is decomposed by water with formation of trichlorobuytl alcohol

The ammonia compound

## CH, CHCl CCl, CH(OH)NH,

m p 62°, is formed by leading ammonia into a cooled mixture of 1 part of butylchloral, and 1; parts of chloroform (Schiff and Tassmari, Ber 10, 1783). It forms a solid mass, insoluble in water, soluble in ether, alcohol, and chloroform. Is readily decomposed on heating.

Butyl chloral has a strong affinity for water,

and readily forms

182 CHLORAL.

+H<sub>2</sub>O. This compound, which is also known as butyl chloral, is prepared by passing chlor-ine into paracetaldehyde (Freundler, Compt. rend. 143, 682). It forms white crystalline trimetric plates of a peculiar odour, with difficulty soluble in cold, more readily in hot water; very readily soluble in alcohol; mp. 78°, with partial dissociation, and sp gr 1 694 (Schroder, Ber 12, 562) For some years after its discovery in 1870, it was regarded as crotonchloralhydrate It is decomposed by distillation into the chloral and water Alcohol and potassium cyanide convert the hydrate into ethyl a-chlorerotonate Zine and hydrochloric acid, or zinc-dust and water, produce mono-chlorcrotonaldehyde, and finally crotonaldehyde (Sarnow, Annalen, 164, 108) With acetic acid and iron filings, butyraldehyde, normal butyl alcohol, and crotonyl alcohol are produced (Lieben and Zeisel, Monatsh 1,840). It is a valuable anodyne in cases of neuralgia, facial rheumatism, periostitis, &e The following composition is recommended: butyl chloral hydrate, 2-5 grams, alcohol, 10 grams; glycerol, 20 grams, and distilled water, 120 grams (J Soc Chem Ind 1889, 476; 1890, 889) It is frequently adulterated with the cheaper chloral hydrate, from which it may be distinguished by the following reaction: A solution of pyrogallol in pure 65 p c. sulphuric acid gives, when gently and carefully warmed, a blue colour with chloral, a ruby colour with butyl chloral and a more or less violet-to-blue colour with mixtures of the two compounds On addition of a large quantity of water, the blue colour obtained with chloral changes to yellowish-brown, and the ruby colour obtained with butyl chloral to a more or less deep-violet colour (Galbutti, J. Soc. Chem. Ind. 1894, 273)

Derivatives —Butylchloralantipyrine, yellow crystals, mp 70°-71° (Calderato, Chem Zentr 1902, ii 1387); butylchloraldoxime (Tarugi, Gazz chim ital 21, ii 6)
CHLORALAMIDE. Syn for chloral form-

**CHLORALAMIDE.** Syn for chloral formamide CCl<sub>3</sub>—CH NH CHO used as a hypnotic

фн

and sedative

CHLORALUM v DISINFECTANTS

CHLORAMINE-BLUE, -GREEN, v Azo-colouring matters

CHLORAMINE-T (Sodium - p toluenesulphochloramide CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>NNaCl), first obtained by Chattaway (Chem Soc Trans 1905, 87, 153), has been shown by Dakin to be a valuable antiseptic in military surgery for the treatment of infected wounds, and has been employed in cases of cerebrospinal meningitis, diphtheria, &c It may be obtained from p-toluene sulphonic chloride, a by-product in the manufacture of saccharm (q v.), or, directly, from toluene by sulphonation and transformation of the sodium salt into the sulphonic chloride by the action of phosphorus pentachloride or a mixture of the pentachloride and phosphoryl chloride, and conversion of the sulphonic chloride into sulphonamide by treatment with ammonia or ammonium carbonate. To convert the amide to chloramine, one molecular proportion is dissolved in a warm strongly alkaline solution of sodium hypochlorite (1.3 to

Butylchloral hydrate CH<sub>3</sub>·CHCl·CCl<sub>2</sub>·CHO 2 0 N strength), when on cooling the greater 4.0. This compound, which is also known butyl chloral, is prepared by passing chlorinto paracetaldehyde (Freundler, Compt. d. 143, 682). It forms white crystalline metric plates of a peculiar odour, with iculty soluble in cold, more readily in hot ter; very readily soluble in alcohol; mp. total miles of hypochlorite and 1 mol. of sodium hydroxide Known also as chlorazene and total miles of the chloramine separates out, and the part of the chloramine separates out, and the part of the chloramine separates out, and the remainder can be obtained by evaporation. It is important to have an excess of caustic soda; for each mol of amide about 1 05 to 1 1 mols. of hypochlorite and 1 mol. of sodium hydroxide Known also as chlorazene and the chloramine separates out, and the part of the chloramine separates out, and the part of the chloramine separates out, and the remainder can be obtained by evaporation.

Dichloramine T, toluene-p-sulphodichloraamide CH<sub>3</sub>C<sub>4</sub>H<sub>4</sub>SO<sub>2</sub>NCl<sub>2</sub>, is also employed as an antiseptic. It is prepared by dissolving ptoluenesulphonamide in bleaching powder solution or by acting on chloramine T by hypochlorous acid—Insoluble in water but dissolves in chlorinated parafin or chlorinated eucalyptol which is then diluted with parafin oil.

The persistent odour which clings to the hands after immersion in hypochlorite solutions appears to be due to protein chloramine combinations (Briggs)

CHLORAMINE-YELLOW v. PRIMULINE AND ITS DERIVATIVES

CHLORANIL v QUINONES

CHLORATE OF POTASH, SODA v CHLORINE. CHLORAZENE. Syn for Chloramine T (q v)

CHLORAZIDE. N<sub>3</sub>Cl A gas obtained by mixing solutions of sodium azide N<sub>3</sub>Na, and sodium hypochlorite, and acidifying with acetic acid. Chlorazide is highly explosive, slightly soluble in water, and insoluble in caustic alkalis (Raschig, Bot. 1908, 4194).

CHLORAZOL-BROWN, -DEEP BROWN, -BLUE, -VIOLET v Azo- colouring matters

CHLORENE C<sub>48</sub>H<sub>26</sub> A green hydrocarbon obtained together with decacylene and fluorocyclene by heating accomplishment with lead oxide at 350° and extracting with benzene and light petroleum in the cold. It crystallises in small dark green scales. Dilute solutions are pure deep green in colour but become brownishing of exposure to sunlight. Concentrated solutions appear purple-red by a bright transmitted light. (Direwonski and Suknatowski, Ber 1918, 51, 457).

CHLORETONE " ACETONECHLOROFORM CHLORIDE OF LIME, POTASH, SODA "

CHLORINE. Symbol Cl At wt 35 46. HISTORICAL Chlorine was first obtained by Scheele in 1774 by the action of hydrochlorine acid upon manganese dioxide, and was termed by him dephlogisticated manne acid air' In 1785 Berthollet suggested its use for bleaching vogetable fibres, and this application continued until 1789, when this very objectionable gas, and its equally objectionable aqueous solution, were replaced by an unobjectionable solution prepared by dissolving the gas in an alkali; the hypochlorites so prepared have been used for bleaching from then to this day

The idea that it was a compound substance prevailed until 1809, when Gay-Lussac and Thénard suggested that it was probably an elementary body, a supposition confirmed by the investigations of Davy in 1810 (Phil. Trans. 1811, 1, 32) Davy gave it its present name, from χλωρόs, greenish-yellow, in allusion to its characteristic colour.

OCCURRENCE Chlorine is never found free in nature, it always occurs combined with metallic elements Sodium chloride occurs to

the extent of 3 p.c. in sea-water, and as rock salt in large quantities in Galicia, Tyrol, Transylvania, Spain, and in England (particularly in Cheshire) (v Sodium Chloride, art. Sodium) Potassium chloride is found as sylvine (q.v), and in association with magnesium chloride as carnallite KCl MgCl2,6H2O (q.v.) in the saltbeds of Stassfurt, and in a few parts of North France, and in sea-water. Other naturally occurring chlorides are matlockite PbCl. PbO. horn silver AgCl, atacamite Cu2Cl(OH)8, calomel HgCl, &c. Ferrie chloride Fe<sub>2</sub>Cl<sub>6</sub> and sal ammoniae are occasionally found native as products of volcanic action. Magnesium chloride is found in considerable quantity in seawater and in many natural waters Calcium chloride is found in certain brines

Sodium and potassium chlorides are present in most animal secretions, and free hydrochloric acid is met with in the gastric juice. Many plants, growing in proximity to the sea, and many inland plants contain notable quantities

of chlorides

Preparation for Laboratory Purposes The oldest mode of preparing chlorine for laboratory purposes consists in heating manganese dioxide with hydrochloric acid, or by heating a mixture of manganese dioxide, common salt, and sulphuric acid

$$\begin{array}{l} {\rm MnO_2 + 4HCl = Cl_2 + MnCl_2 + 2H_2O} \\ {\rm 2NaCl + MnO_2 + 3H_2SO_4} \\ {\rm = Cl_2 + 2NaHSO_4 + MnSO_4 + 2H_2O} \end{array}$$

It is readily prepared by slowly dropping a clear strong solution of bleaching powder into concentrated hydrochloric acid. Bleaching powder made commercially into small cubes may be used in a Kipps' gas generator containing strong hydrochloric acid. If required in rather large quantity it is most easily obtained from liquid chlorine supplied commercially in steel

cylinders

PROPERTIES OF CHLORINE At ordinary temperatures and pressures chlorine is a greenish-yellow gas, which becomes darker in colour as it is warmed, under strong pressures the colour becomes orange-yellow; its absorption spectrum shows numerous dark lines in addition to complete absorption in the blue and violet. Even when largely diluted with air, the gas possesses a characteristic pungent and irritating smell, it rapidly attacks the nasal and lung membranes, and in such cases the best relief is obtained by breathing alcohol vapour or a little chloroform vapour and keeping in a warm room

The sp gr of chlorine gas shows a gradual diminution up to 240°, when its density becomes normal, viz 2 4502 (Jahns, Ber 15,

1242)

By the application of either suitably increased pressure, or of a suitable low temperature, chlorine gas is readily liquefied to a dark greenish-yellow liquid of sp.gr. 1.33, b.p.  $-34.5^{\circ}$ . The critical temperature is  $144.0^{\circ}$ , the critical pressure 76 l atm, and the critical density 0.573. The density of liquid chlorine may be represented by the formula :

$$d_t = 0.687014 + 0.0002379(144 - t) + 0.0622109 \sqrt{144 - t}$$

The ratio of the critical density to that calculated from the simple gas laws is 3 635, and the value of Trouton's constant, that is the quotient of molecular heat of evaporation and the absolute temperature of the boiling-point, is 20 67, indicating that chlorine behaves as a normal substance. The specific heat between 0° and 24° is 0 2262, and the heat of evaporation at 8° is 62 7. The vapour tension is in atmospheres.

It can be solidified at a low temperature Liquid chloring is not miscible with water, its refractive index is lower than that of water, and it is a non-conductor of electricity

Chlorine gas is readily soluble in water, its absorption coefficient between 10° and 415°, according to Schonfeld (Annalen, 93, 26, 96, 8),

$$3.0361 - 0.046196t + 0.0001107t^2$$

by means of which the following table showing the volume of chlorine at  $0^{\circ}$  and 760 mm absorbed by 1 vol of water at varying temperatures from pure chlorine gas having a pressure of 760 mm is calculated:

The volume of chlorine absorbed from a mixture of that gas with an indifferent gas, as hydrogen or carbon dioxide, is greater than that calculated from the law of partial pressures (Roscoe, Chem Soc Trans 8, 14)

Chlorine-water has the characteristic smell and colour of the gas, in daylight, and especially when exposed to sunshine, it gradually loses its colour owing to the action of the chlorine on the water, the main reaction being

$$H_0O+CI_0=2HCI+O$$

Chlorine Hydrate A solution of chlorine in water below 9° deposits crystals of so-called chlorine hydrate of a light-yellow colour, which become almost white when cooled to --50° By gently heating the crystals in a sealed tube, they are readily resolved into water and free chlorine, which may be thus obtained as a liquid

GENERAL CHEMICAL PROPERTIES—Chlorine is an active chemical agent, and combines with many of the elements even at the ordinary temperature Phosphorus ignites spontaneously in the gas, forming phosphorus trichloride. Finely divided arsenic, antimony, copper, tin, lead, and iron burn in chlorine, forming the respective chlorides, but most of these metals, when presented in bulk, behave quite differently, as the trace of solid chloride formed acts as an impervious layer and the reaction can proceed no further, but metallic tin forming a liquid chloride Sn(14 is entirely converted into that chloride. Chlorine and hydrogen may be mixed together in the dark without combining, but such mixtures explode violently in bright daylight or when heated During any of the above acts of combination much heat is evolved

Certain of these reactions are modified if the chlorine is absolutely free from moisture. Wanklyn (Chem News, 20, 271) found that sodium might be melted in dry chlorine without action, although the metal readily burns in chlorine as ordinarily prepared. Cowper (Chem Soc Trans. 43, 153) found that Dutch metal, zinc, and magnesium were unacted upon by dry chlorine, silver and bismuth were only slowly attacked. The indifference of dry chlorine to steel and brass enables liquid chlorine to be manufactured and shipped in vessels made of those metals.

With many inorganic chemical compounds chlorine also combines directly, thus it unites with sulphur dioxide, carbon monoxide, phosphorus trichloride, ferrous chloride, stannous chloride.

With very many organic non-saturated compounds chlorine also "combines directly, eg with ethylene, acetylene, and benzene (under certain circumstances), and additive compounds are formed; in the case of acetylene the reaction is so energetic that it becomes explosive; lignocellulose also readily combines with chlorine

With saturated organic compounds as maish gas and the paraffins, acetic acid, alcohol, &c, chlorine forms one or more substitution products, and the eliminated hydrogen is simultaneously converted into hydrochloric acid

With such organic compounds as benzene and toluene, chlorine may react either to form addition or substitution products, or both simultaneously, and such reactions are largely influenced by the temperature and by the presence of other substances as ferrous chloride, iodine, &c

With turpentine chlorine reacts so readily that the oil inflames spontaneously, and much of it is totally destroyed with the production of free carbon and hydrochloric acid

In presence of water chlorine acts on many substances, both morganic and organic, as a powerful oxidising agent, and these reactions may be explained by the assumption that the mixture reacts to a small extent to form hypochlorous acid, that is HOH+Cl2=HOCl+HCl3, hypochlorous acid is an intensely active substance, capable of converting manganous oxide into permanganic acid, chromium trioxide into chromic acid The oxidising action of chlorine and water often becomes more energetic in the presence of bases. In this way monoxide of lead may be converted into dioxide, manganous oxide into manganese dioxide, sulphur into sulphuric acid, ferrocyanides into ferricyanides, manganates into permanganates, and many organic substances and colouring matters may be more or less completely oxidised.

## INDUSTRIAL PREPARATION OF CHLORINE

Chlorine gas has been prepared on the small industrial scale by the addition of strong sulphuric acid to a mixture of sodium chloride and manganese ore (which is essentially manganese peroxide) and heating

The commercial processes of preparing chlorine to-day are described in the following account in detail, in the order of their development:—

(a) The action of an aqueous solution of hydrochloric acid on one or other of the higher oxides of manganese, or their compounds at temperatures below 100°:

 $4HCl (solution) + MnO_2 = Cl_2 + MnCl_2 + 2H_2O$ 

(b) The action of hydrochloric acid gas on atmospheric oxygen in the presence of catalysts (as copper compounds) at temperatures just below a red heat:  $4HCl+O_2=2Cl_2+2H_2O$ .

(c) The electrolysis of metallic chloride (as sodium chloride, potassium chloride, zinc chloride), which are occasionally anhydrous and fused, but are much more generally employed in aqueous solution 2NaCl=Cl<sub>2</sub>+2Na, and ZnCl<sub>2</sub>=Cl<sub>2</sub>+Zn; the energy absorbed in these reactions is supplied electrically

A number of proposals or non-important

methods are briefly enumerated

(a) The Aqueous Hydrochloric Acid and Manganese Oxide Method —The solution of hydrochloric acid used is obtained in the manufacture of sodium sulphate that is salt cake. It is known as 'Commercial Hydrochloric Acid,' or as 'Muriatic Acid' The solution used should be as strong as possible, because complete exhaustion of the acid by manganese peroxide is impracticable, and a residual liquor containing a considerable amount of unavailable acid is always left. The stronger the hydrochloric acid solution to start with the smaller the percentage of the total HCl which is lost (for actual figures, see below)

The manganese ore used for the generation of chlorine should contain a high percentage of MnO<sub>2</sub>. The two next lower oxides of manganese also liberate chlorine, but only a smaller proportion of the total chlorine employed as shown by the following equations —

$$\begin{array}{l} MnO_2 + 4HCl = Cl_2 + MnCl_2 + 2H_2O \\ Mn_2O_3 + 6HCl = Cl_2 + 2MnCl_2 + 3H_2O \\ Mn_3O_4 + 8HCl = Cl_2 + 3MnCl_2 + 4H_2O \end{array}$$

Pyrolusite is the manganese ore generally used; it usually occurs in distinctly crystalline masses of a fibrous structure, its sp gr is 47-5, and its colour iron-black to dark steel-grey with a metallic lustre, it also occurs as amorphous friable masses that soil the fingers It has been mined in North Wales, and occurs in quantity in Germany, Bohemia, Transylvania, Russia, India, Australia, Japan, and the United States of America

The chief source of the industrial pyrolusite used in this country is Southern Russia. The ore varies usually from 75 p c. to 85 p c.  $MnO_2$ . An average analysis of the Russian ore is . 80 p c  $MnO_2$ , 1 p c MnO; 3 p.c  $H_2O$ .

It is customary to buy and sell manganese

ore on the percentage of MnO<sub>2</sub> only.

The higher the percentage of MnO<sub>2</sub>, the more valuable the ore, as the impurities, even if substances such as BaSO<sub>4</sub> and SiO<sub>2</sub>, do harm by coating the particles of MnO<sub>2</sub> and preventing the action of the HCl, and also by adding to the amount of waste material to be handled. If the impurities are carbonates (CaCO<sub>3</sub>, BaCO<sub>3</sub>, &c), they not only waste hydrochloric acid, but the CO<sub>2</sub> evolved during their solution harms the bleaching powder into which the bulk of industrial chlorine is made.

The physical condition of the manganese ore is also important, the softer ores being preferable

to the harder ones, which are more slowly acted on by the hydrochloric acid

For generating somewhat small quantities of chlorine the generating vessel, or still, may be made of stoneware Fig 1 shows one of the





Fig 1 shows one of the best-known forms of stoneware stills, intended for heating from the outside in a water-or steam-bath, made of wood or masonry. The manganese is placed in the perforated cylinder, which holds about 1 cwt, and is put in and out by means of specially shaped tongs, through the large open-

ing The small necks serve for introducing the acid and taking away the gas. At the end of the operation the still liquor is drawn off by a siphon, or by a discharge pipe passing through the steam-jacket. Such small stills permit of careful and economical working so far as materials are concerned, since the acid is not diluted by condensed steam.

Fig 2 shows a contrivance which allows the chlorine pipes and the main pipe to be instantly connected or disconnected. There is a Y-shaped lead pipe a, the upper arms of which are converted into hydraulic lutes by 'burning' an outer pipe, bb, on to them. The ends of the gas pipes, both that leading from the stills c, and that leading to the gas main d, are widened out into cups dipping into the water



which fills the annular space at b. The rubber pipe c and its attached cup contain sufficient water to fill and thus stop the gas-way a whenever the pipe and cup are raised

For generating large quantities of chlorine the stills are made of silicious sandstone or other stone

not acted upon by acids, not pervious to them, and not liable to crack by rapid changes of temperature. Such stone is found at Southowram, and is known as 'Yorkshire flag'.

A small old-fashioned stone still is shown in Fig 3, which represents it cut through in the middle. The bottom is formed by a stone, 10 to 12 inches thick, with grooves for the sides The sides are stone slabs, 5 to 6, or even 8 inches thick, and joined at the corners on the 'feather and groove' principle. The end stones project both ways over the sides, which fit into grooves cut in the end stones The joints are made tight by dressing a small groove into the two adjoining faces, and placing a 5-inch india-rubber cord in the channel thus formed. The lid consists of one or more stones laid in a rebate, and the joint is made gastight by partly filling the rebate with a putty of linseed oil and china-clay. The diagram shows how the still is fitted with the following parts. A false bottom a; a pipe b, for the introduction of hydrochloric acid, sealed at the bottom by the acid in a small earthenware bowl; a leaden steam pipe, c, connected at the bottom with a stone or stoneware steam column

d, which is provided with side holes (not shown in the figure) situated a short distance above the bottom slab, for the issue of the steam into the acid, and at the top by the loop e with a tap f (the loop protecting the tap to some extent against the action of the chlorine), the earthenware or leaden gas pipe g, the connection of which with the gas main h can be opened or





Fig 3

shut off at will by putting more or less water into the pot i. The man hole for charging the still and the discharging hole must be assumed to be provided in that half of the still not shown in the diagram. These stills are made  $7 \times 5 \times 3$  feet, inside measurement, and upwards

Fig. 4 shows a still of the bevel-joint type, seen from above, with the cover removed, a are the flags forming the false bottom, one of which, a', is made to stand up; c is the steam stone, but the long central hole and the radial short holes for the passage of the steam are not shown, A, B, C, D are the rods binding the stones together by means of the cast-iron corner



pieces b, b. The joints are made by the compression of the stones on to round rubber cords shown as black dots in each corner joint in the figure. The cord for the bottom joint may be one inch diameter, that for the upright joints may be \(^2\) inch diameter; the junctions of the several cords with each other are made liquortight by boring with a cork borer a comparatively

small hole in one cord, and then inserting the end of the other cord into the broad end of a hollow conical steel needle, which is smeared with glycerin, the second cord may be passed through the hole in the first cord and the small

projecting end neatly cut off

The Process—In order to work the stills, those of smaller sizes, Fig. 1, where the manganese is contained in a sieve, are usually first charged with the requisite quantity of acid. The sieve containing the manganese (about 1 cwt) is then put in, the cover is at once put on and made tight by water-luting or cementing, and the evolution of chlorine begins forthwith. When it slackens, heat is gradually supplied by turning on the steam till the acid is exhausted. The stills are then allowed to cool down, usually till the next morning. They are then opened, the sieve is lifted out, the manganese remaining in it is washed and replenished by more ore, the liquor is siphoned or run out of the still, fiesh acid is run in, and a new start can be made at once.

The large stone stills are charged with from 6 to 10 cwts of manganese broken up into pieces, which must be smaller in the case of hard ores, but should not be too small, lest too much is washed down below the false bottom, The manhole is then closed and acid is run in, at first quickly, afterwards gradually, so that it takes several hours to run in the full charge All this time chloring is given off, much heat being generated in the process, of course much more with concentrated than with somewhat dilute acid. When the evolution of gas becomes slow, steam is injected into the still, but very cautiously, as otherwise the water might be thrown out of the lutes, and only at intervals, for instance for 10 minutes every hour much steaming causes too rapid an evolution of chlorine, and the volatilisation of much acid and water. The temperature of the stills ought never to get above 90°. It is preferable not to steam all the stills of a set at the same time, but in regular rotation, to produce a more continuous current of chlorine. Chlorine stills ought to be kept going at least 24 hours, but large stills are better kept going for 48 hours, as more of the acid will then be utilised

When the waste liquid, 'still liquor,' is run off it emits an intolerable stench of chlorine, so that this operation is most irksome, both for the workmen and those people who live near the The 'still liquor,' even after being works freed from its muddy constituents by settling, is, on account of its acidity, one of the worst nuisances for any watercourse into which it may be run These nuisances have been completely overcome by the introduction of the Weldon process, where the still liquor from the fresh manganese ore is run into the Weldon stills, as described a little later on, and at works where chlorine is made on too small a scale to permit of applying the Weldon process, the still liquor now has to be run into a closed tank, and neutralised by chalk or limestone dust before running it away, even in a very diluted form.

The chlorine evolved in the stills is always contaminated with hydrochloric acid gas, and a considerable amount of aqueous vapour on account of its high temperature. To remove

these impurities the gas is passed through a long string of earthenware or lead pipes, externally cooled by air (or, if needful, by water), so that the water vapour condenses, and the water produced dissolves the hydrochloric acid gas, and both are then drained away by

trapped escape pipes

The Chemistry of the Process.—The first reaction taking place in the chlorine stills is:

MnO<sub>2</sub>+4HCl=MnCl<sub>4</sub>+2H<sub>2</sub>O or (more probably)

2MnO<sub>2</sub>+8HCl=Mn<sub>2</sub>Cl<sub>6</sub>+4H<sub>2</sub>O+Cl<sub>2</sub>. (There is some discrepancy of opinion as to which of these two chlorides of manganese is formed. (Compare W. W Fischer, Chem News, 37, 250;

U. S Pickering, ibid. 39, 225; Berthelot,

Compt rend. 91, 251.)

These chlorides, MnCl<sub>4</sub> and Mn<sub>2</sub>Cl<sub>6</sub>, form a dark-brown solution which quickly decomposes even at the ordinary temperature into MnCl<sub>2</sub> and free chlorine, so that the ultimate result is: MnO<sub>2</sub>+4HCl=MnCl<sub>2</sub>+2H<sub>2</sub>O+Cl<sub>2</sub>, but this reaction is only completed at about 100° and with a certain excess of hydrochloric acid Theoretically for 100 parts by weight of MnO<sub>2</sub>, or an equivalent quantity of manganese ore, almost exactly 170 dry HCl or 530 acid of 32° Tw, ought to be consumed; practically, at least, 10 pc mole is used, and with low-strength acid, hard manganese, direct steaming, &c., twice the theoretical quantity of acid may be required to be present

The still liquor contains chiefly manganous chloride, free hydrochloric acid, the chlorides of other metals present in the ore, and a little

chlorine

Black's analysis of still liquor, in a well-conducted works, showed how incompletely the acid is utilised

MnCl					-10.6
HCl					66
$Al_{s}Cl_{g}$					0.6
Fe,Cla .					0.5
Cl <sub>2</sub> .			not d	leter	mined
H <sub>2</sub> O (by di	ff)	•	•	•	81.7
					100.0

On the assumptions that all the manganese in the ore was in the form of MnO<sub>2</sub>, and that no hydrochloric acid gas was carried away by the chlorine, it follows that out of every 100 parts of HCl added 33 6 parts had been left untouched.

The free acid is tested for in daily practice by an easy but approximate method, namely to a measured volume of still liquor, standardised caustic soda liquor is run in from a burette till the formation of flakes of Fe<sub>2</sub>(OH)<sub>6</sub>, which do not dissolve on shaking, indicates the saturation of the free acid.

TREATMENT OF STILL LIQUOR FOR RECOVERY OF THE MANGANESE AND THE METHOD OF USING THE RECOVERED MANGANESE Many suggestions have been made and tried with this object, but one process has proved superior to all others, and is now universally adopted. The first foundations of this process were laid in 1837, when Gossage precipitated the still liquor with lime, ran off the supernatant liquor, and agitated the precipitate with air; the oxidation was, however, very slow, and was never so much as half complete. In 1866 Walter Weluon made his first attempts, and

by 1869 his new process was in commercial use at Messrs Gamble & Co's works at St Helens, and by 1871–1873 had been so improved as to be adopted by all English and continental alkali works. The apparatus employed has received but few modifications, but the conduction of process, which in the critical part is complicated, and still not thoroughly explained, has been more modified. In the following the apparatus and process are described together, and in the sequence of events occurring when starting up a new plant.

Fig 5 shows the plant in elevation.

The process is commenced in the stone still E, which has been already described in construction and process by Figs 3 and 4 and the accompanying text. The only changes demanded by the Weldon process are. (1) That the acid employed should be as free as possible from sulphuric acid, because when subsequently the liquor is treated with calcium carbonate the voluminous crystalline needles of CaSO<sub>4</sub> 2H<sub>2</sub>O that are formed entail extra labour and expense in filter pressing and removing the filter cakes, and there is an accompanying loss of manganese chloride solution (2) That the run-off pipe for the still liquor, instead of leading into a neutralising well, leads into a mud still D, which is shown in section in Fig 5 The utility of this last change does not arise until a later stage in the process, and the special construction of this large mud still p will therefore not be now described

The still liquor may be at once run from D into one or other of the two neutralising wells G, G by means of pipes or coveted channels not shown in the figure, and charge after charge of native manganese ore is dissolved in E in the same way, filling the wells G,G alternately and repeatedly for the next operation.

The neutralising wells a, a are sunk in the ground on a bed of puddled clay, and after being built are surrounded by puddled clay to prevent possibility of leakage, they are built of large stone slabs of acid-proof Yorkshire flag, as already described for the still E, except that it is much larger and is octagonal in shape, their dimensions are, depth 6 feet, diameter 15 feet. On the top of the wells two large timber beams are seen supporting the agitators composed of wooden gates driven by short vertical shafts (situated above the liquor level), and large crown wheels and pinions situated above the timber beams, the pinions are driven by one horizontal shaft and the small steam engine situated between the two left-hand mud stills D, D. The liquor, while continuously agitated, is slowly treated with crushed chalk or powdered limestone, which is preferably very finely ground. The free HCl is first acted very finely ground upon, yielding calcium chloride, which is an essential component for the remaining processes, and carbonic acid gas, which in escaping carries away the small quantity of dissolved chlorine gas; these two objectionable gases are prevented from escaping freely into the air by covering the wells with strong planks well tarred and by connecting them to the suction of the nearest chimney by an earthenware pipe Towards the end of the operation the additions are made slowly, and with frequent testing of samples, by seeing if they effervesce when


poured on to ground chalk. When the free acid has been neutralised the ferric chloride, and aluminium chlorides shown in the preceding analysis are also converted into calcium chloride and ferric and aluminium hydroxides, and these last carry down as ferric arsenate any arsenious acid originally present in the muriatic acid During this treatment any sul-phuric acid present in the muriatic acid is converted, as already mentioned, into CaSO<sub>4</sub>,2H<sub>2</sub>O. Complete neutralisation is necessary to remove all the impurities named, and also to prevent corrosion of the settling tanks A, A, into which the liquor is next pumped, but the excess of CaCO, employed should be the smallest possible to reduce the cost of the subsequent settling and filtering operations and their loss of manganese liquor

As soon as the neutralised liquor in one of the wells a 13 ready a swivel pipe is lowered from above into the liquid, as shown in Fig. 5, in the left-hand well, and the turbid liquor is then piped to the force pump shown in the figure in front of the third tank p from the left The internal working parts of this pump are made of gun metal. The figure shows the delivery pipe rising to the top of the plant, then running horizontally and discharging through control cocks into one or other of the four chloride of manganese settlers A, A, A, A As soon as the well a is emptied and the pump stopped, a small valve is opened at the base of the pump delivery pipe, and the pipes thus drained back into the well; the swivel pipe taking the liquor from the well is also withdrawn, these things are done to prevent The well is then ready corrosion of the pipes to receive its next charge, and so on

The chloride of manganese settlers are made of 3-inch thick boiler plates, and are 18 feet by 12 feet and 6 feet deep, and therefore are carried by a very strong timber framework that rests upon the strong brick walls that carry the settlers c, c, c As the stock of liquor in A does not fill the tank, and is moreover not sufficient for the next operation, the processes already described must be repeated until at least one settling tank A is quite full. After making the last addition a few hours' settling should allow the liquid to settle quite clear, it is important that the liquor going forward to the next operation should be quite clear, otherwise the charge in the next operation is apt to froth over, and also the recovered manganese will consume an undue amount of acid

As the correct proportioning of the manganese chloride liquor and the milk of lime in the following process are admittedly the key to the rapidity of the subsequent absorption of oxygen, and as no practical method of regulating this proportion is described in Lunge's standard work, "Sulphuric Acid and Alkali," and as it is admitted that milk of lime differs very considerably in its initial action on manganese chloride liquor, according to the source of the lime and the manner of making it into milk, the usual description is here slightly modified so as to contain a means of ascertaining and controlling the proportions of MnCl<sub>2</sub> and CaO used.

Having correctly measured the length and width of the settler A and the position of the



710. 57

surface of the liquor, and recorded these measurements, and having also taken a sample of the liquor for analysis as shortly described, the clear liquor may be drawn off from the settler by lowering a swivel pipe just under the surface of the liquor, where the open end of the pipe can be inspected and the liquor above it can be seen to be quite clear. The bottom end of the swivel pipe is attached to the side of the tank about a foot above the bottom of the tank, so as to leave plenty of room for the mud to collect and to be stored pending its removal Outside the tank the flow of the clear liquor is controlled by a cock, and cocks control its entrance into the oxidising tower B, shown in Fig. 5, or into a second similar tower situate just behind it. The treatment of the mud left in the tank will be described later, as it is first allowed to accumulate.

The oxidising towers is are made of 3-inch thick boiler plates, and they may be 8 feet diameter and 28 feet high, the bottom is flat, and the top is quite open. The pipe bringing the manganese chloride liquor enters the tower about 6 feet from the top of the tower, and as the tower shakes considerably during the subsequent operations, the connection between the tower and the clear liquor main is made through a short piece of indiarubber pipe diameter and height of the tower being carefully measured, its capacity when half full is calculated, and a further reckoning gives the new position the surface of the liquid will occupy in the settler a when the tower has been half tilled, and as the liquor approaches this level the top of the swivel pipe may be set to give exactly the desired amount of liquor. When the liquor ceases to run the cocks are closed, and the residual liquor in the settler is reserved for the finishing of the charge as explained later

The manganese chloride liquor should be tested for MnCl<sub>2</sub> by diluting a measured sample with boiling water, adding zine oxide until the mixture is white as milk, and, lastly, adding standard potassium permanganate with occasional boilings, until the supernatant clear liquor shows a pink flush, the reaction yields a manganite exactly like the Weldon process under consideration ZnO+MnO+O=ZnO MnO2, and the necessary oxygen is obtained from the permanganate similarly

3ZnO+2KMnO<sub>4</sub> =K<sub>2</sub>O·ZnO+2(ZnO MnO<sub>4</sub>)+3O The percentage of manganese chloride having been calculated from the analysis, the lbs. of manganese chloride in the batch is then easily reckoned.

The temperature of the manganese liquor in the tower will be above atmospheric temperature, and in subsequent operations will be still more so, but experience has shown that if it is below 55° C, it should be heated to this point by blowing in steam The steam pipe is not shown in the figure, and because of the shaking of the tower it must pass over the top of the tower, and terminates at the bottom of the tower in several branches containing perfora-tions. The temperature is ascertained by drawing a sample from a sampling cock situated in the cabin there is easy access to all other parts of the plant.

The manganese chloride liquor during the above heating may be agitated, and for the subsequent operation must be well agitated. This agitation is effected by a blast of air The blowing engine-house is shown on the left hand of Fig 5 The right-hand cylinder and fly-wheel constitute the steam engine, the steam supply pipe being shown descending to the centre of the cylinder. The left-hand cylinder is the blower, the air is taken in on the ends of the cylinder through numerous valves of the flap type, faced with rubber, and is delivered through the pipe shown leading to the air pressure regulating vessel m, and thence by the 7-inch diameter pipe d over the top of the tower B, because of the considerable shaking of the tower during the subsequent operations, and thence to the bottom of the tower, where it ends in a cross or a gridion of pipes, marked n in Fig 5, that contain a multiplicity of perforations on their underside, so that the pipes keep themselves clear from the liquor and mud treated in the tower. When used for agitating purposes alone the blast of air required is not great, but the blast can be regulated as desired, and as described later, can be increased very considerably

Milk of hime is required for the next operation, and it is prepared in the upper of the two tanks, J, J, shown in Fig 5 Each of these tanks is made of boiler plate, and is provided with an agitator, the drive being effected by a small steam engine affixed to the right-hand side of the upper tank J, and transmitted by a shaft and two small bevel pinion wheels to either or both of the bevel crown wheels seen on the tops of the two agitator shafts upper tank only contains a lime cage, the bottom of which is indicated by a horizontal line above the third agitator blade, the cage is built of iron rods or perforated metal plates on the bottom and sides, the top of the cage is quite open. The tank is filled with water while steam is being also blown in, until the hot water stands well above the level of the bottom of the cage Quicklime is selected so as to contain less than 1 p c of magnesia, it should be well burnt, but not over-burnt. The magnesia is very objectionable, because it accumulates during the repetition of the following processes, and in the generation of the chlorine it neutralises acid to no purpose, the method of iemoving it is described later. Well-burnt lime does not contain more than 2 pc. carbonic acid, and it slakes quickly and completely, giving a milk that is particularly free from git. Over-burnt lime slakes exceedingly slowly, and the resultant milk contains large quantities of minute grains of unslaked lime, that react only exceedingly slowly with water or the manganese calcium chloride liquor, and so remain mert until the chlorine is generated, when they neutralise acid to no purpose. The selected quaklime in the form of large lumps is put into the cage. it soon slakes, generating thereby much heat, and this addition is continued until the milk registers 50° on Twaddell's hydrometer, the test jar being gently rotated the while; the milk then contains 20 lbs. CaO per cubic testing cabin, shown in the figure, from which foot, and is as strong as can be conveniently

prepared. The milk is then run from a valve at the bottom of the tank on to a wire sieve (not shown in the figure), and thence into the bottom tank J, which is used as a combined store and measuring tank After cleaning out the cage and the wire sieve the upper tank J is then ready for the next operation.

The next operation is the addition of milk of lime to the hot and agitated manganese calcium chloride liquor in B. The agitation of the lower tank J being stopped for a minute, the exact position of the surface of the liquid is noted, and the internal diameter of the tank measured. the agitation being re-started, a sample is taken, from which, while constantly being stirred, a pipette is filled and, after washing the exterior, the contents are titiated with standard hydro-chloric acid till neutral. It is now possible to calculate the position of the surface of the liquid in the tank J when enough milk has been removed to (1) theoretically precipitate all the MnCl2, and to (2) increase the total amount to 16 times the theoretical, as long experience with the process and the due consideration of opposing considerations, namely, the rapidity of oxidation, the completeness of the oxidation, the economy of lime and of hydrochloric acid, the settling quality of the recovered manganese, and the avoidance of 'stiff batches' have shown that this figure gives the best result second addition is called the excess lime, and the amount desired is accordingly 0.6

The pump k is now put into operation, and the milk pumped into the tower by the pipes shown, due regard being had to the means already described of overcoming the effect of the shaking of the tower on the end of the pipe During this operation the utmost vigilance is required in the testing cabin, samples being drawn, filtered, and tested as rapidly as possible The first test is simply that for alkalinity with red litmus paper, but as manganous hydroxide is to some extent soluble in calcium chloride solution, the alkalı test is not proof that all the manganese chloride has been decomposed second test is therefore made by adding to the filtered sample a strong solution of bleaching powder, which produces a brown precipitate or colour if any manganese is still present Should manganese still be present, either more milk of lime must be added, or a little more time given for that already added to react. When sufficient lime has been added to just remove all the manganese from the solution the addition is stopped, and the amount that has been added is measured and recorded.

The amount of lime so far added is always found to exceed that expressed by the simple equation MnCl<sub>2</sub>+Ca(OH)<sub>2</sub>=Mn(OH)<sub>2</sub>+CaCl<sub>2</sub>, because of the solubility of Mn(OH)<sub>2</sub> in CaCl<sub>2</sub> solution already mentioned, and because milk of lime always contains larger particles, or mactive particles, that only react very slowly, and for these reasons the excess lime may vary from 0 10 to 0 45 (the theoretical lime named above being 100); but it is fairly constant for any one source of supply of the quicklime, and for a routine method of working

The amount of calcium chloride present profoundly alters the solubility (and hence the reactiveness) of the calcium hydrate, as is shown by the following figures for 60° C.:

Percentage CaCl2 in solution 100 c c dissolve 09 '11

19

17 .22 38

and whereas in the two weaker solutions the solubility decreases with further rise of temperature, in all the other solutions the solubility increases, and the more so the stronger the solution. The solubility is more than mechanical, as on cooling very long crystalline needles of the oxychloride

### Cl·Ca·O Ca OH+7H<sub>2</sub>O

separate and from strong hot solutions so much that the mixture becomes apparently solid and can be inverted, and this has been assumed to be one of the reasons for the production of stiff batches' mentioned later

The amount of calcium chloride in the batch may be readily determined with sufficient accuracy from the sample drawn when the precipitation of the manganese was found to be complete, by filtering and taking the degrees Twaddell of the filtrate and applying a correction of 0 07° Tw per 1° temperature variation, and the table

Percentage CaCl,

grams of CaO

10 15 20 Twaddell at 186° c 15° c -01 80 165 255 348 446 546

In the present batch from native manganese the liquor will contain about 18 p.c. only, and because of the low solvent power of this strength of calcium chloride on the free lime in the excess lime already added, it is not permissible to add any more excess lime until some of the free lime has been removed by converting it into calcium manganite

For this purpose the blast of air from the blowing engine is increased to its maximum and continued until the end of the operation. The fixation of the oxygen causes the temperature to rise a few degrees, and the reaction CaO+MnO+O=CaO MnO2 reduces the amount of free lime and changes the colour of the precipitate from light yellow to brown, and then to black After about a quarter of an hour a quarter of the remaining excess time desired is added, and these additions repeated after three more periods of blowing

This procedure is necessary to prevent the batch going 'stiff,' ic changing from a freeflowing turbid liquid to a pasty liquid, or even to a paste so thick that it stops the blowing engine, and the whole plant has to be stopped while the paste is dug out The cause of this stiffening is the separation of crystalline compounds said not to be the calcium oxychloride already mentioned, but rather a very basic calcium manganous manganite

## 2CaO 2MnO·MnO<sub>2</sub>

To attempt to cure a stiff batch the excessive bases must be reduced by running in more manganese chloride, and increasing the blast to its utmost.

If, on the other hand, a sufficient supply of free base is not present when the blast is turned on full, then the manganous hydroxide is not converted into a manganite at all, but into the oxide Mn<sub>3</sub>O<sub>4</sub>, which may be written MnO Mn<sub>2</sub>O<sub>3</sub>, and the batch turns 'red' or foxy, and no amount of blowing or of further base will subsequently convert it into a manganite.

During the blowing, samples are taken each half-hour and tested for the amount of MnO by adding to a measured quantity a measured solution of standardised ferrous sulphate and sulphuric acid, so that the manganites are all dissolved, when the excess of ferrous sulphate determined by titiation with standard permanganate solution. By the repetition of this test the time required for the blowing is found, that is the time required for the MnO. per cubic foot to attain its maximum; three hours should suffice During the progress of the blowing filtrates from samples will show a diminution of and then the disappearance of the free lime. The simultaneous oxidations of the mixture (CaO+MnO) and of the excess MnO to form (Ca OMnO<sub>2</sub>) and (MnO MnO<sub>3</sub>) is now complete, and continued blowing has no further offect

But a new reaction is now brought into play by the addition of a second supply of manganese chloride liquor to the blowing tower, namely 2(CaO MnO<sub>2</sub>)+Mn(I<sub>2</sub>

 $= (CaO 2MnO_2) + CaCl_2 + MnO$ 

and by continuing the air blast the MnO is converted into (MnO MnO2), as in the first part of the operation The addition of the manganese chloride is continued only until a sample filtered and tested with hypochlorite gives a reaction for manganese samples show that as a result of the continued blowing this reaction soon ceases, whereupon a little more manganese chloride is added ' blowing out' of the manganese becomes slower with each addition, and when it becomes too slow, no more is added, and the whole operation The effect of this last operation is to ıs ended increase the amount of chlorine that can be manufactured from the mud, without increasing the amount of acid required for that manufacture.

The mud, consisting of calcium dimanganite  ${
m CaO~2MnO_2}$ and manganeso manganite MnO MnO<sub>2</sub>, suspended in a calcium chloride solution is now run from the base of the oxidising tower by the cock and main c, c, Fig 5, into the mud settlers C, C, similar in construction to the settlers a above. After a few hours settling is complete and the clear supernatant calcium chloride liquor may then be run away through the main E, E by opening the cock near the bottom of the tank and gradually lowering the swivel pipe p, until the black mud below is reached

The manganese mud is now ready to be used in the manufacture of chlorine. It differs from native manganese in that it is entirely soluble in hydrochloric acid and that it dissolves very much more readily. The generating tank therefore does not require a false bottom, and it is made very large. Fig. 5 shows four such stills D, D, that may be 12 feet diameter and 12 feet deep. They are octagonal, but otherwise resemble in construction the native manganese still shown by Fig. 4. The right-hand end still is shown in section, the two columns s, s are the perforated stone blocks for blowing in steam. The outlet liquor pipes are not shown, nor are the gas outlet pipes, nor the acid inlet pipes

Into the still is charged any liquor available from the still E, because though it is not strong enough to act further on native manganese, it

will act on recovered manganese mud, and this is one of the advantages of the Weldon process Muriatic acid is then also run into the still till the acid liquors are 2 feet deep. The settled mud in the settler tank c may then be stirred by a hand rake, to make it a little more fluid, and run through a cock q situated in the bottom of the settler, into the main f, f, and thence through another cock into the luted trap h, whence it runs into the still. The chlorine is given off very readily, and the rate of evolution of the chlorine is regulated by regulating the admission of the mud. As the acid becomes exhausted steam is blown in to raise the temperature, and the further additions of mud are made with caution, until samples taken from the still by an earthenware cock, not shown, cease to be clear yellow, and become a clear coffee colour that is not removed by further steaming A sample tested with standard caustic soda solution will then show the presence of about 0.5 p.c. of free acid. The waste manganese liquor is now run, as before described, into a neutralising well a by opening an earthenware cock and the Weldon recovery process is repeated, but with the following alterations

The manganese chloride liquor contains a large quantity of calcium chloride, but is comparatively free from ferric chloride, aluminium chloride, and free hydrochloric acid, and therefore requires much less limestone dust to neutralise and purify it. There is accordingly much less deposit in the settling tanks A, A. The liquor charged into the oxidising tower B contains an increasing amount of calcium chloride, and this may amount to nearly three times as much as the manganese chloride with advantage, as instead of adding the residue of the 'excess' hme in small repeated portions as previously described, the whole of the residue may be added at once, and by adopting the method of testing already described the whole of the lime may be added at once without discriminating between the 'theoretical lime' and the 'excess lime,' thus saving time in blowing and labour in testing

When sufficient mud has accumulated in settlers A, A, a long-handled plug (indicated by a dotted line in each of the tanks) is lifted, and the mud is run through pipes q to filter presses (not shown in the figure), where as much manganese chloride liquor as is possible is recovered, and the cakes of stiff nearly dry mud are then thrown away

When sufficient manganese chloride liquor has been made from native manganese in the still E to sufficiently stock the settlers A and C and to give working charges in G, B, and D, then only so much native manganese requires to be dissolved in E as will counterbalance the unavoidable losses in the whole of the Weldon plant.

(b) The Gascous Hydrochloric Acid and Almospheric Oxygen Method — After many unsuccessful attempts made by Oxland (1840), Jullion (1846), Binks (1860 and 1862), Dufrené (1865), and others, the labours of H Deacon and F Hurter, in the years from 1868 on, produced an entirely successful chlorine process which is founded on the direct oxidation of the H in HCl by atmospheric oxygen, and which utilises much more of the HCl than any process employing

either native or recovered MnO<sub>2</sub> Some processes patented after theirs (Henderson, 1871; Weldon, 1871, Wigg, 1873, &c.) proceed on the same lines, but have not been practically successful.

The Deacon process starts from the wellknown fact that a mixture of hydrochloric acid gas and oxygen at a temperature just below a red heat, especially in contact with porous substances, partially changes into chlorine and water:  $2HCl+O=H_2O+Cl_2$  This decomposition is very much assisted by the employment of substances which serve as oxygencarriers, and of these the salts of copper (previously employed by Vogel, 1855, Gatty and Laurent, 1860, Mallet, 1866), have been found most useful Probably in all cases cupile chloride is formed, which at once splits up into The Cu2Cl2 cuprous chloride and free chlorine combines with oxygen to form oxychloride CuO CuCl<sub>2</sub>, and this acts again upon HCl, yielding water and reforming cupric chloride, so that the reactions can begin over again:

(1) 
$$2CuCl_2 = Cu_2Cl_2 + Cl_2$$
  
(2)  $Cu_2Cl_2 + O = Cl - Cu - O - Cu - Cl$   
(3)  $Cu_2OCl_2 + 2HCl = H_2O + 2CuCl_2$ 

Theoretically, all the HCl would thus be converted into chlorine, but in practice, under the best conditions, about two-thirds of this decomposition is effected. The undecomposed HCl is, however, not lost, but is recovered and can be utilised for other purposes, eg for the Weldon process

From a diagram constructed by Hurter, showing the affinity of technically available elements for oxygen, chlorine, and hydrogen, it appears that no other metal than copper forms two oxides and two chlorides, in which the combination is of so loose a character. It is therefore almost a certainty that a cheaper and equally efficient catalyst cannot be found, and that the Deacon process is the best for the direct conversion of gaseous HCl into free chlorine.

Deacon's process deals with the gaseous hydrochloric acid, as it is evolved in decomposing common salt by sulphure acid without condensing it into liquid acid. This is evidently a great advantage in one way, but when the salt cake is made in the ordinary handworked pot and furnace, it has the drawback that in the beginning of each operation much more HCl, and hence more Cl, is got than afterwards. In practice, only the gases from the decomposing pots, say 68-70 pc. of the total HCl, are used for the Deacon process, those coming from the roasters being too dilute and containing too much sulphuric acid. In order to equalise the operation to a certain extent, the gases from two salt-cake pots can be mixed and the pots worked alternately, so that the strong gas from the early part of one charge is mixed with the weak gas from the latter part of the other charge.

The quantity of air necessary to form chlorine enters through the joints of the doors and dampers. It is regulated by the speed of the aspirator (a Root's blower) or cast-iron fan, which is placed at the end of the whole apparatus and produces the movement of the gases through it. Usually there are 4 vols. of air to 1 vol. of

HCl. An excess of air is less injurious than a deficiency of it, in which case there is a poor decomposition.

The gases leaving the pot are first cooled by a long string of pipes and a scrubber; thus much of the water and about \( \frac{1}{2} \) or \( \frac{1}{2} \) of the HCl is removed. A patent by Deacon and Hurter (Eng Pat 2104, 1888) provides for cooling the gas and drying it in a sulphuric-acid tower before it enters the decomposer, such complete drying being very advantageous for the process. The gases now enter the heating apparatus. This is a furnace, 16 foet \times 10 feet, in which twenty-six vertical pipes, 12 inches wide and 9 feet high, are arranged in two sets of twelve each, connected like the 'breeches pipes' in the blast-heating stoves, formerly used in nonworks and now replaced by brick heaters.

The gases are here heated up to 500°, the waste heat of the furnace is sufficient for heating the decomposer, which does not possess any fire of its own.

The gases now pass into the decomposer (Fig. 6) This is an upright cast-iron cylinder, 12 to 15 feet wide.

It contains a cylindrical ring of broken bricks D, D, supported by two cylindrical cast-iron rings of shutters placed concentrically one within the other The gases enter at the circumference




Fig 6.

at A, pass radially across the decomposing mass n into the inner space, and are led away through the pipe ob. The annular space between the shutters is 3 feet wide, and is divided into six compartments, one of which is emptied every fortnight This is done by means of the discharging doors c, c below the cylinders Fresh decomposing material is then thrown in from above through E, E, this material consists of buint clay broken up into lumps, or ordinary bricks broken, freed from dust, and dipped into a solution of cupric chloride The mass contains about 06-07 pc of metallic copper. After the active substance has served once, that is, for 10 to 12 weeks, it is thrown away Each cwt of clay produces about 10 to 12 cwts of bleach It has been found useless to attempt utilising the contact substance by redipping it in the copper solution or in any other way. The cost of copper for the contact substance does not exceed is per ton of bleach produced.

The temperature, both in the heating apparatus and in the decomposer, must be most carefully regulated, for which purpose special pyrometers have been constructed (cf. Hunter on Pyrometers, J. Soc Chem Ind 1886, 625). In the decomposer, it ought to range from 480° to 500°.

When the gases leave the decomposer, they consist of a mixture of HCl, free Cl<sub>2</sub> and steam. In the best case, two-thirds of the HCl is converted into chlorine, but sometimes only one half. The hot exit gases are now cooled by passing through a long string of earthenware or glass

through a long string of earthenware or glass pipes, and are then freed from HCl by washing with water in the ordinary acid condensers,

consisting of stone towers filled with coke, combined or not combined with acid cisterns, Woulff's jars, &c. By properly managing this process, all the acid can be obtained in the strong state, as required for the Weldon process

and other purposes

The gases can now be employed directly for making bleach liquor or chlorate liquor from milk of lime. If, however, bleaching powder is to be made, they must be freed from water because the ratio of water to chlorine is so great that strong bleaching powder could not be made; the amount of water depends on the temperature, and can be found from tables of the tension of aqueous vapour, the amount of chlorine is at most 10 pc by volume. The water is removed by passing the gases through a lead tower packed with coke, down which sulphuric acid of not less than 140°Tw is kept flowing.

As will be seen from the description, there is theoretically no loss of HCl in the Deacon process, the whole of the acid evolved from the salt being either converted into chlorine or recovered by condensation as liquid hydrochloric acid. In practice, of course, some losses occur, but they are not necessarily large, and their amount depends principally on the efficiency of the draughting arrangements and the avoidance of low-level escapes at the salt-cake pots and furnaces. The fuel used per ton of bleaching powder to heat the decomposing apparatus depends naturally on the efficiency of working of the process, and varies from 6 to 10 cwts per ton of bleaching powder

During the early years of its existence, the Deacon process had many difficulties to contend with, and the working results were disappointing

and unsatisfactory

The fact that the impurities accompanying the gaseous HCl were sufficient to seriously affect process was not realised. The impurities are sulphuric and sulphurous acid, arsenious

acid, water, and carbon dioxide

The sulphure acid and sulphurous acid carried along with the gaseous HCl, especially when attempts were made as they frequently were in the early days, to utilise the roaster or furnace gas as well as the pot gas, converted the CuCl<sub>2</sub> into CuSO<sub>4</sub>, which requires a higher temperature and gives a lower decomposition than CuCl,

The arsenious acid derived from the As<sub>4</sub>O<sub>8</sub> in the sulphuric acid used forms arsenate of copper, which is even less reactive than the CuSO<sub>4</sub> The presence of these impurities necessitated the frequent renewal of the decomposing material. Water is one of the products of the decomposition, and hence, when present in the inlet gases, increases the partial pressure of the products of the reaction and lowers the maximum p.c. decomposition obtainable under the conditions of the process

Carbonic acid derived from the fuel gases, either of the salt-cake furnaces or of the decomposing furnace itself, was a serious obstacle to, and frequently entirely prevented, the production of strong bleaching powder. This difficulty has been overcome by greater care in the construction of the apparatus and testing for leakage.

It will be noted from the above description worked.

Vol. II.-T.

that only the gaseous HCl from the salt-cake pot is dealt with by the Deacon process, and that allowing for the liquid acid collected after the decomposers, about 60 p c of the total acid is obtained in the liquid form, and has to be sold as such or used in some other way such as by the Weldon process.

193

Hasenclever, however (Eng. Pat. 3393, 1883), describes a process for gasifying the HCl in the liquid acid by mixing with strong hot sulphuile acid in a series of earthenware vessels through which air is blown. By this means, a steady continuous stream of gaseous HCl is evolved with an easily regulated supply of air passing through the series of vessels, the vitriol diluted with the water contained in the liquid hydrochloric acid has to be reconcentrated to the original sp gr. The concentration is generally performed in long, lead brick-lined pans, heated by top heat, the fuel gases being in direct contact with the sulphuric acid. This process is used to a considerable extent, but from the large quantity of sulphune acid to be concentraied per ton of bleaching powder, a very material addition is made to the cost of manufacture

Other contact substances—Many substances have been the objects of proposals, some of which are inseparable from the manufacture of chlorine from metallic chlorides, that is, from HCl in the nascent state. A general investigation on the mutual action of HCl and O in the presence of certain metallic compounds (of Cu, Fe, Mn, and Cr) was made by Lamy (Bull Soc. chim 1873, 20, 2). We must refer to the original, and will only mention that copper salts were found to be much more efficient than others, and that with these the decomposition is at its maximum at about 440°

The proposals to employ ferric chloude (Thibierge, 1855, and others) or chromic oxide (Hargreaves and Robinson, 1872) have not had

any practical success

Nickel oxide was proposed by L. Mond (Eng. Pat. 8308, 1886) He passes HCl over heated NiO, when the chlorine combines with the base The product is then in the same apparatus exposed to dried and heated air, whereby NiO is reformed and Cl is given off order to increase the active surface, pumice stone is impregnated with the chloride, and after drying the mass is exposed to the action of the gases in cylinders made of earthenware or enamelled iron The gaseous HCl and air being previously heated, the retorts do not require very much additional heat from without, and therefore can be made of a large diameter is claimed for this process that it converts the HCl completely into chlorine, and that impure HCl, which is unfit for the Deacon process, is here quite suitable

De Wilde and Reychler proposed (Eng. Pat. 17272 and 17659, 1889) to employ a mixture of magnesium and manganese chlorides with magnesium sulphate. The reaction takes place in two stages.

(1)  $Mn_3O_8Mg_3+16HCl$ = $3MgCl_2+3MnCl_2+8H_2O+4Cl$ (2)  $3MgCl_2+3MnCl_3+8O=Mn_3O_8Mg_3+12Cl$ 

The process is not, as far as is known, actually worked.

(c) The electrolysis of metallic chlorides method—Recent years have seen a large development of electrolytic processes for the decomposition of metallic chlorides, chiefly those of sodium and potassium into chlorine and alkali. It was observed as early as 1800 by Cruikshank that on electrolysing a solution of common salt, caustic sodia was formed at the negative pole Berzelius, Hisinger, and Davy all worked on the same subject, but a long time elapsed before their observations were utilised in actual manufacture

The difficulties experienced in the electrolytic decomposition of alkaline chlorides on a manufacturing scale have been first the cost of the electric current which was absolutely prohibitive until the dynamo-electrical machine had been so far developed as to be a commercial reality, viz in 1872 by Gramme, and when it became possible to conduct electrolytic experiments on the large scale over long periods of time, it was soon found that there were many and serious difficulties as the corrosion of the anodes, the evolution of gas from the cathode, the diaphragms were either excessively porous or not sufficiently porous, and they disintegrated rapidly, there were excessive resistances, voltages, and the generation of heat, and there were leactions, so  $\mathbf{that}$ neither chlorine nor the alkali metal product was obtained in the expected purity or quantity

The various electrical processes suggested or tried for the decomposition of alkaline chlorides are all attempts to overcome economically these

difficulties

Theoretically a current of 1 ampere liberates

in 1 hour 1 3236 grams of chlorine.

The number of patents taken out since the year 1885 in connection with electrolytic cells for the production of chlorine, is enormous, embodying every variety of shape, design, and method of work. We shall only deal with a few representative samples, illustrating the broad classes of design which have, up to the present, been used industrially on a large scale

In describing these developments of the electrolytic processes, and their effects upon Leblanc's process, Lepsius in Berliner Berichte (1909, 2915) says, with German pride, that it is no wonder that the electrolytic process was first carried out in Germany, the richest country of the earth in salt and potash, and while he pities the fate of poor Leblanc, yet he concludes with characteristic German military imagination and arrogance, that the bitter international struggle between the old process born in France and developed in England, and its two younger rivals, is now finished after the crippling caused by the Belgian rival, The Solvay Ammonia Soda Process, because there appeared a German rival, the electrolysis of the alkali chlorides, whose victorious conquest of the chlorine market and other successes caused such annihilating wounds that the hundred-year old process was no longer able to stand

Electrolysis of Potassium and Sodium Chlorides

The cells may be roughly divided into four classes

I The cells in which the electrolyte consists of the fused salt

II. The cells in which the electrolyte is a

salt solution, and in which the cathode and anode are separated by a porous partition

III. The cells in which the electrolyte is a salt solution, and in which a moving mercury cathode is used to receive and remove the sodium produced.


IV. The cells in which the electrolyte is a salt solution, and in which the caustic soda solution produced is only kept from mixing with the brino undergoing electrolysis by reason of the greater sp gr of the former

Class I The Cell with the Fused Electrolyte — The best known examples of this cell are the

Vautin and the Acker cell

The Vautin cell (Eng. Pat. 13568 of 1893 and 9878 of 1894) electrolysed fused NaCl or KCl, with or without the addition of fluorides or other chlorides to lower the point of fusion, over a cathode of molten lead or tin with which the sodium or potassium formed an alloy. (See paper by Vautin, J. Soc. Chem. Ind. 1894, 448.)


The Acker cell is designed on the same principle as the Vautin cell and makes use of the lead cathode (see Eng. Pat. 6636 and 6637 of 1898). Below is a sketch (Fig. 7) of the Aker.



cell, taken from a paper by J B Kershaw in Electrician, Oct 25, 1901. The bottom of the cell P is trough-shaped and of cast non serves as electrical connection between the main conductor and the mass of fused lead L resting on it The upper part of the cell Q is constructed of any refractory basic material not attacked by chlorine gas or fused salt anode c is of carbon and dips into the fused The channels and subsidiary vessel R are salt arrangements designed to remove and decompose the lead alloy as it becomes sufficiently saturated with sodium. The decomposition of the alloy is effected by blowing steam into it, and the heat generated by the conversion of the sodium into hydroxide is returned to the cell and assists in maintaining the necessary temperature The voltage actually taken by the Acker cell is 675, and the current efficiency 54 p.c. (Report of Commission on Science and Arts, J. Franklin Inst. 1903, 156). The Acker cell was worked at Niagara Falls on an industrial scale, from about 1900 to 1907, when the works were completely burned down and have not been rebuilt.

Class II. The Porous Draphragm Cells.—The best known of the diaphragm cells is the Griesheim cell, which has been used on a very large scale in Germany, and also in Spain, France, and Russia.

Fig. 8 represents the cell in longitudinal sectional elevation, Fig. 9 represents it in plan,



and Fig. 10 represents it in cross sectional The plate upon which the cell rests is shown by Figs. 8 and 10 to be insulated from




Fig. 9.

the ground by wooden bars, this is so that a number of such cells may be run electrically in The containing bath KK, Fig. 8, KMK,

Fig. 9, M, Fig. 10,




Fig. 10.

is built of sheet iron, and its two long sides are each covered with steam jacket M, the object of which is increase temperature of the electrolytes and so to reduce their electrical resistance. The negative electrical lead is attached to the iron bath as indicated the binding bу screw, lead and -

sign on the right-hand side of Fig 8 The bath is closed by a cover cut into segments and shaped so as to allow of the insertion of the five hanging iron cathode plates B, Fig. 10, and shown but not lettered in Figs. 8 and 10, and the six porous anode compartments A shown in each figure. Water is supplied to the tank K by a pipe (not shown), and by the electrolysis of the brine in the anode compartment is converted into a

caustic soda solution and hydrogen, the caustic soda solution is drawn off by the pipe under the centre of the tank shown in all the figures; the hydrogen gas is led away by a pipe not shown to gasholders for subsequent use The anode to gasholders for subsequent use compartments are composed of iron frames covered inside with cement, in the frames are set and cemented the porous diaphragms in large flat sheets These porous diaphragms were made by mixing cement, common salt and hydrochloric acid in due proportions into a paste, and then forming into the shape required: when the salt subsequently dissolves the diaphragm becomes very porous The anode compartments contain the anodes, which were first the ordinary pressed carbons, then gas retort carbons, and then carbons prepared in a special anode works, where the large pressed plates were roasted at the highest temperature of a porcelain kiln. These special anodes had a life of twelve months The Figs 8 and 10 show them passing through the covers of the anode compartments, and the external electrical Each cover also contains a connections c. long tubular earthenware basket z, Figs. 9 and 10, for the addition of solid salt and a pipe L, Fig 10, for the escape of the chlorine gas The covers on the bath, and on the anode compartments, having been made gas tight by cement, all the anode connections are electrically connected as shown in Fig. 8, all the chlorine pipes are connected into a main LL, Fig 9, and the electrical current may then be forced through the cell.

When the process has proceeded so far that the cathode liquor contains about 8 p.c caustic potash or soda and about 14 pc. of the corresponding chloride, it was found that the chlorine gas contained several per cent of carbonic acid, and as this interferes with the use of the chlorine for making bleaching powder, and as the destruction of the anodes is too costly, the process is interrupted by stopping the current and running off the cathode liquor; the cathode compartment is then refilled with fresh brine and the current restarted The cathode liquor is then evaporated in large double or triple effect vacuum evaporators until it contains 50 pc caustic alkali, during which all but a small quantity of the accompanying chloride crystallises out and is afterwards returned into the process.

The corrosion of carbon anodes led to their replacement by cast anodes of magnetic oxide of iron prepared by fusing ferric oxide in the electric arc furnace, whereby there is also a partial reduction. The anodes are difficult to prepare, but they have a good conductivity, and they last indefinitely With oxide of iron anodes there is no further trouble from carbonic acid, but the anode liquor then contains potassium chlorate which crystallises out and is removed periodically and utilised This output of chlorate can be increased at will by increasing the strength of the caustic alkali in the cathode compartment.


These magnetic anodes were exported by Germany to America, but when war was declared the supply was stopped and a number of attempts were made to produce them in America. Their great drawback is their extreme brittleness, but it has been found that this can be reduced to

one-half by suitable slow cooling or annealing, and that it can also be reduced to one-third by the admixture of about 5 pc of copper oxide, and by combining both modifications rods of 3 9 cm. when supported at 17 8 cm apart require a weight of 119 kg in the middle before they snap (Thompson and Atchison, Trans American Electrochemical Soc 1917, 213)

These cells in Griesheim absorbed from 1884–1892 400 h p, after which it was increased to 800 h p, and soon after to 2000 h p, by 1894 installations began to be erected elsewhere, Bitterfeld in Savony, Westeregeln, Ludwigshafen, Lamotte in France, Flix in Spain, and Slawjansk in South Russia, by 1909 the total h p, absorbed by this cell was 33,000. (Lepsius,

Ber 1909, 2895)

The Hargreaves-Bird Cell—The following illustration shows the Hargreaves-Bud cell (Eng Pat 18871 of 1892: 5197 and 18173 of 1893). The peculiarity of this cell is that the diffusion of the cathode and the anode liquids into each other through the diaphragm, which takes place in ordinary diaphragm cells, is in this cell largely counterbalanced, as far as the diffusion of the cathode liquid is concerned, by employing a very considerable hydrostatic pressure on the anode liquid This electrostatic pressure is, however, not sufficient to force any noteworthy amount of anode liquid into the cathode compartment because the diaphragm is in the ordinary sense nearly liquor tight, but it is yet sufficiently porous to allow the cation, sodium, with its electrical charge to pass through under the driving force of the electric current. The Fig. 11 is a vertical and diagrammatic cross



section of the cell It shows two thick lines marked 'Diaphragm and Cathode'; each of these consists of a sheet of iron wire gaure on to one side only of which is plastered first a layer of asbestos pulp, and then a layer of Portland cement. The frame of the cell consists of a long base shown in the figure and two uprights not shown in the figure, turning the cement sides of the two "diaphragm and cathode" plates towards each other they are so firmly pressed against either side of the base and uprights as to form a watertight cell marked 'anode chamber,' at the same time electrical contact is made

between the iron wire gauzes and the negative electric conductor. Two other sheets of thin iron are then similarly pressed against corresponding ledges provided in the base and uprights, but not shown in the figure, so as to make with the two 'diaphragm and cathode' plates two chambers marked 'cathode chamber' in the Fig Into the anode chamber is placed the anode built up of blocks of gas retort carbon on a central leaden core and also the brine to be electrolysed Into the cathode chamber no liquor as such is put, but steam is blown in, which by heating the anode cell condenses as water on the cathode and so continually converts the sodium deposited by the electric current on the wire gauze cathodes and washes the solution of caustic soda formed down to the bottom of the cathode chamber, from whence it flows by the trapped run-off As it was found that a higher officiency was obtained by passing also carbonic acid gas (the exhaust from gas engines) into the cathode chamber, the manufacture of caustic soda was abandoned and replaced by the manufacture of sodium carbonate

A paper on this cell, by Hargreaves (J. Soc. Chem Ind 1895, p 1011), states the current density as 187 amperes per square foot of diaphragm, the voltage required for the preceding current as 3.4 volts; the current efficiency as 80.3 p.c., and that the resultant soda liquor contains per 100 molecules  $\mathrm{Na}_2\mathrm{CO}_3$ , only 3.0 molecules NaCl if ordinary care is exercised, but that it is possible to reduce this amount to only 0.03 molecule. The present working is understood to give a current efficiency of about 90 p.c., and to require 4 volts per cell.

The Townsend Cell (Eng Pat 18403 of 1904) resembles the Hargreaves-Bird cell with this difference, that the cathode compartment is filled with kerosene oil which squegees the alkali as it is formed off the cathode, and thus makes it possible to produce chlorine and caustic soda in this type of cell without great loss of current efficiency Kershaw (Electrician, April 24, 1908) gives the current efficiency as 90 p c, and the voltage per cell as 47 In the Seventh International Congress, 1909, 10, 36, Backeland describes the result of 31 years' work at Niagara with 1000 kilowatts, the old cells were made for 2000 amperes, but new cells to consume four times this power will be capable of taking 6000 amperes The current density per square foot of anode surface has also been increased from 100 amperes to 144 amperes The attack on the carbon anodes requires the cleaning of the cell in about 30 days, which is a very simple matter, and the graphite anodes are so arranged that they can be repaired, and are ultimately entirely consumed. Aluminium conductors last longer than copper conductors in atmospheres where small amounts of chlorine are occasionally present. The hydrostatic pressure in the anode chamber is regulated so that with a cell using 2500 amperes, the amount of liquor passing through the diaphragm is most advantageously 24 litres per hour, which gives a soda liquor containing per litre 150 grm. NaOH and 200 grm. NaCl, and under these conditions the anode liquor contains but little hypochlorite and chlorate and consequently the chlorino has a satisfactory degree of purity.

The Finley Cell (Eng. Pat. 1716 of 1906) has a double diaphragm dividing the anode and cathode compartments. The diaphragm is very open in texture, and the brine feed for the cell is run into the centre compartment between the two diaphragms, from where it passes outwards on either side to the anode and cathode compartments. In this way, an additional obstacle, depending for its strength on the speed of feed of the brine, is interposed to prevent intermixture of the products of the electrolysis.

The diaphragms are made of asbestos 2-3 mm thick, and the compartments are also only a few millimetres thick, the diaphragms and frames are belted together in any number after the fashion of a filter piess. (Seventh International Congress of Applied Chemistry, 1909,

x. 42)

Class III. The cells in which the electrolyte is a salt solution, and in which a moving mercury cathode is used to receive and remove the sodium

produced

The mercury cathode for the preparation of alkali metals was first used by Davy, it was described in an application for a patent (Eng Pat 4349 of 1882) by Nolf It is mentioned again in Hermite's patent, 3957 of 1886 Further patents describing its use are Hermite and Dubose, Eng Pat 21959 of 1891, Greenwood,

Eng. Pat 5999 of 1891

The general principle of these cells is that the electrolysis takes place between an anode above and a cathode of mercury vertically below. The chlorine liberated naturally rises upwards and out of the solution, and the sodium or potassium freed combines with the mercury to form an amalgam. The amalgam, when sufficiently highly saturated with the alkaline metal, is replaced by fresh mercury and the amalgam itself decomposed by water, with the liberation of H<sub>2</sub> and the formation of a caustic alkaline solution

The current efficiency of the modern mercury cell is stated to be not less than 90 p c, and the

voltage required 4 3 volts

Professor Lepsius (Ber 1909, 2903) enumerates the following advantages and disadvantages of the mercury cell as compared with

the diaphragm cell -

The advantages of the mercury cell are (1) the purity of both the chlorine and the caustic alkali produced, which are very important advantages, (2) the comparatively high concentration of the caustic solution; (3) the high current efficiency.

The disadvantages of the mercury cell are:—(1) the higher voltage, 4 3 volts as against 3 5 volts in the diaphtagm cell; (2) the cost of the mercury, of which 72 tons are required for a plant utilising 6000 h p , (3) the greater cost of installation apart from the question of the mercury

The various patents taken out and cells suggested differ chiefly in the method adopted for the movement of the merculy. To Castner, however, belongs the chief credit of making the

mercury cell technically successful.

Casiner's Rocking Cell—Castner (Eng. Pat No. 16046 of 1892) employs a moving mass of meicury, connecting up the anode and cathode chamber. Patent No. 10584 of 1893 makes the movement of the mercury by a rocking motion of the cell, see Fig 12, which represents the original rocking cell A is the cell divided into




Fig. 12.

three compartments, and resting on the righthand side on adjustable kinfe edges B The lefthand side of the cell rests on an eccentric, c, tevolving on a shaft; this gives the rocking motion The two outer or anode compartments have carbon anodes and are filled with saturated bime, the centre compartment contains iron cathodes and is filled with NaOH solution A layer of mercury, about  $\frac{1}{8}$  inch thick, hes on the bottom of the cell and alternately flows from first one and then the other anode compartment into and through the cathode compartment into the other anode compartment. The chlorine liberated in the anode compartment is carried away by pipes on

Castner, in the Chemical Trade Journal, 1894, 211, stated that the results of 1 week's work on a unit of 28 working cells, arranged in two parallel series of 14, gave a yield of 930 lbs. chloring and 1120 lbs. NaOH per 24 hours, the amperes used averaging 571, and the volts 55 1 or 3 9 per cell

In 1895 the Castner Kellner Alkalı Co was formed to work this cell. Works were erected in Runcorn with 1000 h p, and other similar works were soon erected abroad, viz. at Osternicuburg near Bernburg with 1500 h p by the Deutsche-Solvay Werke, at Augsburg by Merster Lucius & Bruning, at Jemappes near Brussels with 1000 h p, at Liubimow in Russia, at Jaice in Bosma with 1300 h p, and at Niagara Falls, N America, by the Castner Electrolytic Alkalı Co in 1901 with 6000 h p. (Lepsius, Ber 1909, 2903)

in 1901 with 6000 h p (Lepsius, Ber 1909, 2903) Kellner Pressure-displacement Cell—In this cell the chlorine compartment is placed in the




Fig. 13.

centre of the cell and on either side of it is a caustic compartment, each of which is provided on its further side with a recess, DD, which forms the alternative receptacle for the mercury. In these receptacles are immersed pressure vessels open at the bottom and connected at their tops to compressed air supplies which are automatically so regulated that the mercury is forced to move continuously from side to side by compressed air.


Above (Fig 13) is a sketch which shows the construction and working of the cell, taken from the Chemiker Zeitung, Oct. 30, 1909.

The anodes are platinum gauze of which I gram is made to cover four square inches, and are placed in a cement box which has proved more satisfactory than the stoneware jars Each cell contains six electrode formally used boxes, and each box holds 88 pieces of platinum gauze. The cathodes are cast iron, cast in grill shape, and the primary battery, sodium amalgam-caustic solution-iron, acts independently of the main current through the chloring cell, The current used is and also regulates itself 4000 amperes per cell

The great advantage of this cell is the highcurrent density which has resulted in consider-

able reduction of plant

The Kellner-Solvay Scoop Wheel Cell —This cell consists of two slanting troughs communi-



cating with one another at each end, in which the mercury is kept in constant circulation by a rotating scooped wheel placed between the two troughs at the one end. This cell is built; in very large units, taking up to 10,000 amperes, and it is said that a 15,000-ampere cell is well within the bounds of possibility

Fig 14 is a diagrammatic sketch of the chlorine side of the cell, B being the mercury inlet and D the outlet for the amalgam, the scooped wheel is, however, not shown, neither is the caustic

side of the cell

Whiting Cell —A cross sectional view of this cell is shown in Fig 15 The right-hand side shows in the upper portion the tement cover with its attached graphite anodes and connections being lowered into the brine cell below, which consists of a number (say five) of separate compartments

intervals of two minutes, the mercury amalgam is entirely emptied by the automatic lifting of poppet valves, which are operated by the little shaft and cams shown above the cells towards the left, and the liquid amalgam runs over the graphite plates lining the threefold zie-zae iunway shown on the left of the figure; these plates and the amalgam being immersed in water act as a primary cell and generate caustic soda solution, hydrogen and pure mercury. The mercury next reaches the scoop wheel and by arrangements not shown in the figure is lifted and flows to the right side of the figure, whence it enters each cell

The ebb and flow of the mercury in rotation is used to admit fresh brine supplies to each compartment. This complicated arrangement is made to prevent the unequal flow of mercury and the formation of sodium amalgam of excessive strength. The cell was operated at the

Oxford Paper Co., Rumford, U.S.A., and is de scribed in Trans American Electrochem. Soc., 17, 327 (1910)

The new Castner-Kellner Cell differs essentially from the Kellner-Solvay cell only in the method of conveyance of the mercury, which, in this case, is effected by means of an Archi-Taussig, Zeit. für Elektrochemie, median screw 1909, 659

Class IV The Specific Gravity Cell — The best known cell of this type is the so-called 'Glocken Cell,' patented by the Oesterroichischer Verein für Chemische und Metallurgische Produckte, and from 1901 working on a large scale at Aussig in Bohemia, Berlin, and Stassfurt

Fig 16 is a diagrammatic sketch, in which A is the anode and K the cathode, separated by the non-conducting and impervious walls D, E of the bell a The bath w is filled with salt solution, which enters at L and overflows at the pipe F The chlorine escapes at c. These walls are required to keep the ascending current of hydrogen gas arising from K quite distinct and separate from the ascending current of chlorine gas arising from A The anode and cathode are both situated in the upper parts of the electrolyte, and these parts are greatly agitated by the ascending gas bubbles-but the lower portions of the electrolyte are almost entirely stagnant and serve only as a means of conducting the electric current from the anode to the cathode. As the brine around the anode loses a portion of its chlorine by escape of the gas, and loses a corresponding portion of its sodium by electric migration of the cation Na' towards the cathode, this layer of liquid becomes continually lower in specific gravity and would ultimately lose all its dissolved salt were it





Fig 15.


not for the constant feed of fresh brine at L: the quantity of this feed is such that it is only partly able to counterbalance the reduced On the other hand, the sodium gravity liberated on the cathode forms with the adjacent brine an intensely strong film of caustic soda

solution, which falling by its high specific gravity into the lowest part of the cell forms a



heavy layer there which is scarcely disturbed by the slow current of weak brine from the inside of the bell trickling past the edges of the bell to be partially converted into caustic soda and

hydrogen before it passes away by the exit pipe F



Figs 17, 18, and 19 are diagrams of the actual cells, Fig 17 being a cross sectional elevation, Fig 18 a

plan, and Fig 19 on the left-hand side a longitudinal sectional elevation of the bell and its attachments, and on the right-hand side a cross sectional elevation of the same. The





Fig. 18

outer containing vessel is seen to contain two ledges to carry the bells, of which there are a great number in each cell. The junction of the anode with its conducting stem is badly shown



in the figures. The brine supply is different from that shown in Fig 16, for, to prevent the agitation of the electrolyte by the inflow of brine, the supply pipe is made to traverse the whole

length of the anode, its two ends are closed and the brine escapes by a number of small holes along its upper side. The main portions of the cathode are best seen on the right-hand portion of Fig 19 at x. The chlorine passes from bell to bell by the connecting arms shown in Fig. 18, and then makes its exit from some particulai bell by the larger pipe shown in Figs 17 and 19, but not shown in Fig. 18. The electrical connections are indicated diagrammatically in Figs 17 and 19 only Each bell takes about 22 amperes at 4 volts (Lepsius, Ber. 1909, 2904)

Electrolysis of Zinc Chloride—The waste calcium chloride from the ammonia soda process has been utilised since 1897 by Brunner Mond and Co, of Winnington, as a source of By heating the liquor with crude chlorine zinc oxide and carbon dioxide the reaction  $CaCl_2 + ZnO + CO_2 = ZnCl_2 + CaCO_3$  takes place, or by treating it with native zinc carbonate (calamine) the reaction  $CaCl_2 + ZnCO_3 = ZnCl_2 + CaCO_3$  The filtered solution of zine chlorido is freed from its impurities (amongst which is a not inconsiderable amount of cadmium) and the resultant solution is electrolysed with carbon anodes and circular revolving cathodes that are only partially immersed, to produce chlorine gas and an exceedingly pure metallic zinc. Many of the details are due to C. Hopfner (But. Pat. 11724 of 1891, and D R P 85812 of 1895) Fuller details are given in two volumes of Monographien über angewandte Elektrochemie, namely—Die Darstellung des Zinks auf elektrolitischem Wege, Gunther, 1904, 99-180 and Die elektrochemische und elektrometallurgische Industrie Grossbritanniens, Keishaw, 1904. 108, and also in Applied Electro-Chemistry, Allmand, 1912, 286

(d) Proposals and non-important methods for the manufacture of Chlorine—Many processes have been proposed and tried for utilising the residual ammonium chloride, or the residual calcium chloride of the ammonia soda process, but with the exception of Hopfner's zinc chloride method just previously described, none are now in use Many other processes have also been tried for preparing chlorine from various sources and in various ways, but have been abandoned. Some of these old methods are enumerated below, fuller information may be obtained from Lunge's Sulphuric Acid and Alkali

1 Chlorine from ammonium chloride—The most persistent efforts for the direct manufacture of chlorine from NH<sub>4</sub>Cl were made by Mond, who took out the English patents Nos 65, 66, 1049, 3238, 8308, of 1886, 10955 and 17273 of 1887, 2100 and 2575 of 1889 Articles dealing with the Mond process are found in J Soc Chem Ind 1892, 466, 1893, 10, and 1893, 63 There was also a proposition from the Verein Chemische Fabriken of Mainz (Eng Pat 3322 of 1886)

2. Chlorine from CaCl<sub>2</sub>.—Persistent efforts for the direct recovery of the chlorine in the CaCl<sub>2</sub> produced in the ammonia stills of the ammonia-soda process were made by Solvay: Patents 77 and 171 of 1877, 838 of 1880, 7258 and 7259 of 1884; 8724 of 1885; 13389 of 1886; 18574 of 1888 Other patents: Twynam's process (Eng. Pat. 731 of 1885); Bramley's process (Eng. Pat. 8289 of 1887), Lyte and Steinhart (Eng. Pat. 21225 of 1890)

Chlorine from magnesium chloride.—Many efforts were made by Weldon to produce chlorine direct from MgCl<sub>2</sub>; his patents are: Eng Pt 565 of 1868; 967, 968, of 1881; Weldon and Péchiney, 9305, 11035, of 1884; 14653, 14654, of 1887 Detailed description of the Weldon-Péchiney experimental plant at Salmide, is given by Dewar (J. Soc. Chem Ind. 1887, 775) Other patents are Wilson (Eng. Pat 3098 of 1885), Gamble (Eng. Pat 17217 of 1889), Lyte and Tatters (Eng. Pat 17217 of 1889), Schlosing (Eng. Pat. 11821 of 1887, 11469, 11470 of 1891) (See also Eschellmann, J. Soc. Chem. Ind. 1889, 2, comp. ibid. 100)

Chlorine produced by passing HCl gas over heated peroxide of iron on manganese is the subject of a patent by H Burns (No 15448 of

1886).

Chlorine by permanganates has been the object of patents by Tilghman (1847), Condy (1866),

Tessić du Motay (1871 and 1873)

Chlorine from chromates and HCl has been proposed by MacDougal and Rawson (1848), Peligot (Ann Chim Phys. [2] 411, 267), Gentele (1861), Shanks (1858), Claus (1867), Aubertin (1873)

Chlorine by heating metallic chlorides with MnO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> was patented by Binks in 1853


Chlorine by the assistance of nitric and or nitrates - Baggs and Simpson (1864) proposed making chlorine from aqua regia Dunlop (1847) patented the preparation of chloring along with nitious acid, by heating together common salt, sodium nitrate, and sulphuric acid, the nitious acid is absorbed by vitiol and the chlorine is passed on to bleaching-powder This process was working for many chambers years at St Rollov; it was patented again by Roberts and Dale in 1868, and a patent running on similar lines is that of J Taylor (No 13025 of 1884) Other processes employing nitrates or nitric acid, which have not had any practical success, are those of Banks (1839), Tessié du Motay (Bull Soc chim 22, 48), Schlosing (Compt. 1end 55, 284) The later The later patents are those of Donald, Eng Pat 62 of 1887, and Just, Eng Pat 14857 and 14859 of 1888, Davis, Eng Pat. 6416, 6698, and 6831 of 1890, Wallis, Eng Pat 13822 of 1892, and Krause, Eng Pat 16227 of 1894.

Chlorine Liquid — Liquid chlorine was first obtained in 1805 by Northmore — The pure gas can be condensed to a liquid at atmospheric temperature, 15°, by a pressure of six atmospheres, or at atmospheric pressure by reducing its temperature to —34° — If the gas is not pure it requires higher pressures or lower


temperatures to liquefy it

The first to prepare it commercially was Knietsch of the Badische Amlin und Soda Fabrik in 1888 The English Patent 13070 of 1888 records that the ordinary metals withstand the action of chlorine gas provided the gas be dry, so that cast iron, wrought iron, steel, phosphor bronze, brass, copper, zinc and lead may be used for making apparatus for dealing with gaseous and liquid chlorine. The resistance of ordinary metals to attack was evidently not sufficient for the manufacture of a pump with moving metallic piston or plunger, for the patent describes two pumps each with a moving piston of concentrated sulphuric acid to confine

and compress the chlorine gas contained in one limb of a U-tube, while the upper part of the other limb contains another liquid like paraffin oil in which pressure is exerted either by an external pump filled with the same liquid, Fig 20, or an internal plunger A, Fig 21 In the first figure the gas valves D and C are shown



external and are actuated by hand. In the latter figure the valves are internal and are actuated by their own weight and the pressures of the gas, the inlet valve sits on the top of the pipe the outlet valve sits on the top of the pipe opening into the bottom of the chamber in To ensure that the whole of the gas is swept out of the compression chamber a small pipe with an external controlling valve is arianged between the chamber in and the compression



chamber so that on each suction stroke a portion of the acid in D enters the compression chamber along with the gas from F, and on each pressure stroke first chlorine gas and then a little sulphuric acid passes the automatic valve in the bottom of To prevent intermingling of the paraffin and acid the pump limb in Fig 21 is en-

larged in diameter at B. To prevent the possibility of gas being liquefied in the compression chambers, these are heated by an external packet as around B in Fig. 20, and as E.G in Fig. 21. The compressed gas on leaving either form of pump is liquefied in a cooler, as G in Fig. 20, and then runs into the collecting vessel in

In 1909-10 the manufacture was started in the United States of America. With minor

alterations Knietsch's process is still used though it had its difficulties. The Metallurgical and Chemical Engineering of 1916, 1, 215, states that liquid chlorine is now made almost entirely from electrolytic chlorine, and that Deacon chlorine is no good for the purpose unless it is first strengthened, as by extracting the chlorine with stannic chloride as proposed by Goldschmidt Starting with a gas supply of 97 to 98 pc purity the uncondensable blow-off gas still contains 55-65 p c chlorine, and this is absorbed in bleach chambers.

As already stated pure chlorine can be liquefied at atmospheric pressure by reducing its temperature to  $-34^{\circ}$  The Linde Co, of Germany, have introduced the process of simply cooling the chlorine gas to -45° without applying any pressure, but they are said not to liquery so high a percentage of the gas as the older process In 1911 Blucher's Modern Industrial Chemistry states that it was becoming more and more general simply to cool the chlorine gas in refrigerators so that it liquefied at normal

pressures

The liquid is supplied in steel cylinders or in ik cars holding several tons. The physical tank cars holding several tons The physical properties of the liquid were determined by Knietsch (Liebig's Ann 259, 100; the Fifth International Chemical Congress, Zeit f Comp flussige Clase, 1904, 71), and certain of them have been confirmed by Lange (Zeit f Comprimirte und flussige Gase, 1901, 39), who showed that the thermal coefficient of expansion rises steadily to 88° when its value is 0 00366, that is the same value as is given by ordinary gases, and that above that temperature the liquid expands more rapidly than the gases, he also showed that whereas liquid chlorine does not act upon non at any temperature up to 83°, yet at 96° there is a small action, for these several reasons a cylinder of chlorine filled in the ordinary way begins to become unsafe when heated to 90°

Applications of Chlorine.—It is used for the extraction of gold from its ores, and much work has been done on its use for extracting copper, lead, and zinc from mixed ores used for converting ferrocyanide into ferricyanides, manganate of soda into the permanganate, ferrous chloride into ferrie chloride, and was used for converting alumina into aluminium sodium chloride, conveiting tin into stannic chloride either from pure tin or from scrap tinned iron, after which the residual iron can be again smelted for metallic iron, this Goldschmidt process of detinning consumed large quantities of chlorine in Germany and led to the preparation of liquid chlorine in the United States of America when a plant in Chicago was installed in 1911 to treat 40,000 tons of tin scrap per year. It is used along with electrolytic hydrogen for preparing hydrochloric acid in Germany and America It is largely used for purifying drinking water, the liquefied chlorine being extremely handy for this purpose as the supply can be automatically regulated in any proportion to the amount of water passing in the mains, and a summary of the work done on gaseous chlorine as a disinfectant is given by Clayton (Journ Soc Chem Ind., 1896, 320) When used for bleaching it may be absorbed in caustic alkali to make the corresponding hypochlorite, or it may be absorbed in sodium carbonate solution to make hypochlorous acid

 $Na_2CO_3+Cl_2+H_2O=HOCl+NaHCO_3+NaCl$ which acts approximately twice as energetically as the corresponding amount of neutral hypochlorite

It is used for converting earbon disulphide into carbon tetrachloride and sulphur chloride. and for preparing the latter direct from sulphur, for converting benzene into monochlorbenzene, dichlorbenzenes, and trichlorbenzenes, acetic acid into monochloracetic acid, toluene into benzylehloude, benzylene dichloude, and benzenyl trichloride, and alcohol into chloral all these reactions the hydrochloric acid simultaneously formed is condensed in water and a very pure munatic acid is thus obtained

It is combined with acetylene to form tetrachlorethane, and from derivatives of this it is used to form pentachlorethane and tetrachlorethene, with carbon monoxide it is combined to form carbonyl chloride, with sulphur dioxide it is combined to form sulphuryl chloride (Ullmann, Chemische Industrie, 1908, 405) Other uses of chlorine are for chlorinating anthracene, anthraquinone, the cresols, benzoic acid, the xylenes, isoamylchloride (Perkin, Journ See Chem Ind 1912, 616), for the preparation of acetyl chloride, and of acetic anhydride

Many of the above uses of chlorine have been developed because of the necessity of finding new outlets for the enormous quantities of chlorine produced in recent years by the electrolytic processes for preparing caustic potash and caustic soda

The poisonous properties of chlorine gas were first used in waifare by the Germans In April, 1915, near Ypres, a German desetter in the Ypres salient gave warning of the first attempt to use poison gas in modern warfare believed him, but a week later the enemy launched his first attack with chlorine gas against our unprepared and unprotected troops. The gas was liberated from cylinders containing 40 lbs of liquid chlorine The German war of nightfulness was made to include chemical frightfulness Protection against the chlorine gas clouds was obtained by a helmet consisting of a flannel bag soaked in a solution of sodium thiosulphate and sodium carbonate, and contaming a mica window (Bevan, Journ Soc Chem Ind 1916, 651, Self, Journ Soc Chem Ind 1915, 868) Since then the Germans have used a number of gaseous chlorine compounds as poison gases, as phosgene COCI2, chloropictin CCl<sub>3</sub>(NO<sub>2</sub>), trichloromethylchloroformate ClCOOCcl<sub>3</sub>, "mustard gas" dichlorodiethyl-CICOOCCI<sub>3</sub>, "mustard gas uninotical control of the sulphide chloroacetone, chlorosulphonic acid, other, Soc Chem methylchlorosulphonate (Journ Ind. 1918, 127 R, and Tone, Metallurgical and Chemical Engineering, 1918 [2] 357). Shame to their boasted Kultur

In the following pages are described in detail the applications for the manufacture of bleaching powder, liquid chloride of lime, calcium hypochlorite crystal, potassium hypochlorite solution, sodium hypochlorite solution, sodium hypochlorate crystal, potassium chlorate, sodium chlorate and hydrochloric acid.

Chloride of lime or bleaching powder. By far the greater part of the chlorine produced on a manufacturing scale is converted into chloride of lime or bleaching powder.

In 1799 Charles Tennant took out a patent for absorbing chlorine by dry hydrate of lune, and the compound thus formed, 'bleaching powder,' was sold by the St. Rollox Works, established in the same year, at 140l. per ton.

Composition of Bleaching Powder—This substance being originally supposed to be a direct compound of chlorine and lime received its original name of chloride of lime; but in 1835 Balard propounded the theory that bleaching powder was a compound or mixture of equivalent proportions of calcium hypochlorite and chloride

mixed mechanically with a residue of calcium hydroxide In 1842 Gay-Lussac supported these views.

Subsequently Odling, Lunge, Schappi and Naef showed that CaCl, as such does not exist in bleaching powder, as it cannot be extracted by alcohol, etc, and they therefore proposed the constitution Cl Ca OCl, and the full formula for bleaching powder may then be written Ca  $\left\{ \begin{array}{c} \text{Cl} \\ \text{OCl} + \text{H}_2\text{O} \end{array} \right\}$  Free lime is always present in commercial bleaching powder because it is only guaranteed to contain 35 p c of available chlorine, and stops are taken not to let the chlorine exceed this amount essentially, bleaching powder can experimentally be made much stronger than 35 p c. So much so that any attempt to express the free lime as an essential component is evidently fallacious

Preparation of Lime for Bleaching Powder—
The lime required for the manufacture of bleaching powder must be from as pure a limestone as possible and must be well burned. Among the most suitable limes for bleaching powder is that used in Lancashire, and made by burning the Derbyshire limestone in the district round Buxton. The average composition of this limestone is:

CaCO <sub>3</sub> .				98 9
MgCO <sub>3</sub> .				0 29
$Fe_2()_3$ .				0 17
Insol. etc				0.35
$H_2O$ .	•	•	•	0 05
				00.76

The buint lime was until recently slaked by spreading it on a brick floor, sprinkling it with water and turning it over with a spade till it has completely fallen to powder, after which it is allowed to remain in a heap for 10 hours, and the slaked lime is then passed through a wire sieve of 12–18 wires to the lineal inch by the aid of rotating cylindrical sieves. The operation of hand slaking gives rise to a considerable amount of heat, steam and dust which being unconfined are very trying to the workers, who protect themselves with muzzles and goggles.

For this reason slaking apparatus have recently been introduced whereby the operation is entirely mechanical and the apparatus is entirely enclosed and properly ventilated, so

that not only are steam and dust prevented from escaping but labour is also saved.

The slaked lime as used for bleaching-powder manufacture contains about 25 p.c. H<sub>2</sub>O, and should not contain more than 2 5 p.c. CaCO<sub>3</sub>.

The Bleaching-powder Chambers.—The apparatus in which the chlorine is brought into contact with the slaked lime varies with the concentration of the chlorine gas to be treated.

With highly concentrated chlorine gas of 85-90 p c. chlorine chambers of large size are used, namely of about 6½ feet in height, so that a man can work inside them, of a width of from 10 to 20 feet and more, and up to 100 feet long. Their sides and top are usually made of about 5- or 6-lb sheet lead, fastened by external lead straps on an external timber framework, exactly like vitriol chambers.

The floor is made of large stone flags, or of coal-tar asphalt, the joint between the sides and the edge of the floor being specially protected by a skirting and asphalt poured behind it.

The chambers are fitted —with doors at each end, so as to ventilate the chamber when it is being charged or discharged (generally made of non protected by paint), with glass panes in two opposite sides or ends for observing the colour of the gas, with 'lutes' for introducing the gas, with other lutes in the roof of the chambers for the exit of the unabsorbed gases, and with some contrivance for removing the gas remaining at the end of the operation (v infia).

Sometimes the chambers are not erected on the ground-level, but upon pillars, like vitrol chambers, leaving about 10 feet of clear space from the ground. In this case the packing of the finished bleach into casks can be done by means of trap-doors in the chamber floor, with wooden hoppers attached underneath, below which the casks are placed, the bleach being pushed down into them from above by means of wooden rakes.

Sufficient slaked lime is wheeled into the chamber to make a uniform layer of 3 to 4 inches, and the piles of lime are then spread by shovel and a big wooden rake with a plain blade from which stick out below two long gauging spikes and with this the lime is very quickly spread into a perfectly even layer of the desired thickness. In order to assist absorption the surface of the lime is drilled into furrows by a take similar to that described except that the lower edge is provided with deep notches about three inches wide and the same in height; the increased rate of absorption is caused not so much by the extra surface exposed to the chlorine as by the continuous draught of gas through the ridge resulting from the gas in the furrow (chlorine) being heavier than the gas in the ridge (the residual air from the not pure chlorine). The chamber doors are then closed and the joints made gastight with lime putty or with a luting made of

tar and china clay
The 'gassing' of the chambers is done
either singly or in series, the former being the
old method

The Single-chamber Method —A small lute in the top of the chamber being opened, the gas connection is opened, and through the small windows the heavy chlorine could be seen

falling to the bottom of the chamber, and there vanishing. As the formation of the bleaching powder proceeds the gas spreads further, forms a layer all over the lime, the layer increases in thickness and presently chlorine begins to issue from the open lute, which must then be closed and the chlorine main connected to another chamber, without however shutting the gas off from the first chamber, which thus continues to receive a certain amount of chlorine. After a short time the gas inlet to the first chamber is closed, and the chamber is left to itself for 12 to 24 hours, when the gas becomes pale in colour or colourless.

It is difficult to get full-strength bleach by one operation; there is always a superficial crust which greatly hinders the penetration of the gas It is, therefore, necessary to turn the stuff over with spades, and treat again with chloime.

The opening of the chamber doors for this purpose was formerly one of the most awkward operations in an alkali works, as a nuisance was caused not only in the works but sometimes at a distance of a mile or two, and on occasion has been so intolerable as to have produced fatal effects. The matter is now under the Inspector of Alkali, &c, Works. The gas in the chamber must, before any door is opened, be tested by drawing a sample from the exit end at a height of not more than two feet above the chamber floor.

For testing the sample, several kinds of apparatus have been proposed eq that of Ballard (21st Report on Alkalı, &c, Works, 37, 22nd do 55) and of Stark (J Soc Chem. Ind. 1885, 311) Ballard's apparatus consists of a cylindrical glass jar, closed at the top by an indiarubber cork, through which the nozzle of a 4-oz enema ball (finger pump) passes, as well as an elbow tube reaching down to the The lower end of this tube is closed at the lamp, so as to leave an opening fine enough for a needle only to pass through houzontal branch of this tube is inserted in the side of the chamber at a point about 2 feet from the bottom When compressing the ball, the air is driven out by a small hole made in the nozzle just above the cork, this is now closed by the finger, and on relaxing the pressure the ball will expand, and will cause 4-oz measures—that is, 25 of a cubic foot—of chamber air to enter into the jar and bubble through the test solution contained therein This is continued till the end of the reaction is indicated by the test liquor turning blue The liquor consists of 3 485 grains of arsenious acid, dissolved in sodium carbonate and rendered neutral by sulphuric acid, to which is added 250 grains potassium iodide, 50 grains precipitated chalk, 10,000 grains water, 6-10 drops For each test 400 grains are used, ammonia adding a little starch paste at the time of using When 2½ grains of chlorine per cubic foot are present in the chamber air, ten aspirations of the chlorometer will cause a blue colour to appear; if more aspirations are needed, there is less chlorine present, and vice versa. Stark's apparatus attains the same end, not by the finger pump (which is certainly a somewhat rough instrument), but by a gas burette on the Orsat'

Another form of apparatus is Fletcher's in J. Soc. Chem. Ind. 1887, 367.

Collapsing Gas Sampler, an aspirator shaped like a concertina made with ebonite ends and indiarubber coated cloth sides, into which a measured quantity of sodium arsenite is charged It is easily carried to the chamber without fear of breakage An attached rubber tube, a short length of glass tube, both cleaned by recent washing, and a piece of soft, luting enable it to be attached air-tight to the chamber. By fully pulling the bellows open one tenth of a cubic foot of gas is drawn in, and the rubber tube being then closed by kinking, the gas is thoroughly shaken with the solution; a side opening previously closed with a finger is then opened and the gas expelled into the operator's face, it will probably not smell of chlorine, further measured quantities of gas are drawn in and shaken until the expelled gas begins to smell of chlorine.

If the test is above 2½ grains chlorine further time must be given for the chlorine to be absorbed, or it must be otherwise removed most satisfactory way of removing the chlorine is to dust into the chamber through a luted hole in the top, more or less dry slaked lime either by hand or by Brock & Minton's whizzer (Eng Pat 7199, 1886; 23 Report on Alkalı, &c., Works, 58-60) This machine, on being lowered into the hole, rests in the water of the lute and no chlorine escapes. The hopper of the machine is filled with slaked lime powder and the handle being turned the lime drops slowly towards an attached paddle which is revolving at a high speed inside the chamber, and the large air currents which it sets up in the chamber carry the lime dust to a considerable distance in the chamber, and a very energetic absorption of To reach every the chlorine begins instantly part of the chamber the machine is lifted out of the hole and carried to one or two other holes and these also dusted. In this way anything up to 2 cwt of chlorine in a chamber 100 ft long may be absorbed, and the dust settled within 3 hours

When it is proved by any of these tests that the air in the chamber contains less than 2½ grains of chlorine per cubic foot, and it is daylight, then the chamber doors may be opened

After the doors have been opened, and the chamber allowed to ventilate itself a little, the workmen wearing thick flannel muzzles may enter, and with shovels turn the bleach. The doors are then closed, litted, and the chamber gassed again as before described.

Sometimes this must be done even a third time, before the appearance of the product and the laboratory test show it to be finished. To avoid the objectionable turning of the bleach in some British works and generally in Germany and France the layers are made only two or even only one inch

The Series of Chambers Method —This method was described in Lunge's Sulphuric Acid and Alkali, 1880, as the only rational one—It was described in the Inspector under the Alkali Acts Report for 1885 as a most important improvement used in a large English works by which stronger bleach was prepared than previously possible and that without losing chlorine or causing a nuisance—It was again described by Jahne of Petrowitz, and the abstract is found in J. Soc. Chem. Ind. 1887, 367.

The series consists of at least three chambers. better four, or even more, and with the Deacon chambers, which are worked on the same method, seven chambers form a series In the series the gas always enters first into a nearly finished chamber, and the unabsorbed gas passes in succession through other chambers which are further and further removed from being finished until the whole of the chlorine is absorbed and the exit gas is accordingly 'sweet' The thickness of the layer of slaked lime must be such that the whole of it can be sufficiently chlorinated without turning. When it is considered that the first chamber has been sufficiently chlorinated the stream of strong chlorine is directed into the next following chamber and is cut off from the finished chamber. The finished chamber is, of course, full of chlorine nearly as strong as that supplied by the generators and it is absorbed by the bleaching powder only exceedingly slowly The casiest and quickest way of removing this gas is by the use of Brock and Minton's whizzer described in the 'single chamber method' When the chamber is sweetened with all the precautions previously described it may be opened, the bleach may be very carefully sampled, taking great care to take due proportions of the top and bottom layers, and the sample tested in the laboratory. The experience of the workmen generally leads to the obtaining of a satisfactory test, but should it be low the chamber may be turned or not, closed, and gassed again for a certain time, and then treated as before, and finally the bleaching powder removed The empty chamber is at once limed, closed, and put into the working series again, but now as the last chamber of the Each chamber in turn comes up to strength and is removed from the series, packed, limed and inserted again into the series, and so on indefinitely The system of gas pipes and lutes necessary for making these changes is figured in Lunge's Sulphuric Acid and Alkali, 3rd ed vol. 3, p 618, for a set of three chambers. and the same system of pipes are used for the Deacon Chambers next described

The following rules relating to bleaching powder chambers worked in either of the two preceding ways are recommended by the Chief Inspector of Alkali, &c, Works (Mr A E.

Fletcher), in his 21st Report, 12 -

I Bleaching-powder chambers should contain not less than 200 square feet of floor space for every ton of powder made in them per week If there is less space, the work, and especially the

packing, must be hurried too much

II. The chambers should be so arranged and connected together that the residual chlorine from a finished chamber may be drawn into one containing fresh lime, unless some other means be employed for absorbing it (such as Brock and Minton's plan described above) The air finally discharged into the atmosphere or into a flue or chimney must not contain more chlorine than  $2\frac{1}{2}$  grains per cubic foot.

III. No chamber should be opened when the air in it contains more than 21 grains of chlorine per cubic foot. In testing for this, the air should be drawn from a point not more than 2 teet above the chamber floor, and taken from

the draught end

IV. No chamber may be opened until the air

in it has been tested, and the result of such test entered in a book.

V. No chamber may be opened except in

daylıght.

Mechanical Chambers for absorbing Strong Chlorine Gas — Many proposals have been made for mechanical absorbing apparatus for strong chlorine, but none of them seems to have met with permanent success; sometimes the working parts are worn out after a short time, and sometimes the bleach produced in them is too weak. Of course, it cannot be said that the construction of a mechanical chlorine absorber for dry lime is an impossibility, but it must be difficult to construct unless a perfectly regular stream of chlorine gas is at command, and efficient means are employed for artificially removing the heat given off during the absorption

The absorption of weak chlorine, that is, chlorine diluted with very much nitrogen and oxygen, as it is obtained in the Deacon process, and in most of the processes proposed for dealing with calcium and magnesium chlorides, cannot be carried out in the ordinary chambers. The most efficient plants for this purpose are Deacon's shelf chambers, and Hasenclever's

mechanical chambers


Deacon's chambers,—To expedite the late of absorption the furrowed layer of lime is reduced greatly in thickness, viz to § inch As the weak chlorine contains 90 or more p.c. of mert gases, that is, it contains per unit of chlorine nearly 200 times as much mert gas as does Weldon chlorine, the dilution of the incoming Deacon chlorine by the very large volume of air or residual gases in the Weldon chambers is unnecessary to keep down the temperature of the bleaching powder, and the layer of gas as accordingly reduced greatly in thickness, viz to about 6 inches Further, as the escaping mert gases are able to carry away the heat generated, it is unnecessary to have the large heat radiating surfaces which are provided by the Weldon chambers, and it is therefore possible to reduce the ground space occupied by placing as many as sixteen such lime and gas layers as described above over each other Chambers only 6 inches high cannot, of course, be charged with lime and discharged of their bleach by workmen entering the chamber as with the Weldon chambers, but must be operated by the workmen from outside the chambers with long-handled rakes, which, from the hygienic point of view, is a great advantage

Figs. 22 and 23 show the construction of a set of seven Deacon chambers, Fig 22 being a front elevation of 31 chambers, and Fig. 23 being a view of the broken end shown by Fig. 22; the dimensions of each chamber are 8 feet wide and 13 feet high, as shown by Fig. 22, and 21 feet long and 13 feet high, as shown by Fig 23 Each chamber is subdivided into two halves by a central partial partition, as is shown by the right-hand side of Fig. 22, but the figures do not show the gas passage which exists through the bottom of these partial partitions, nor do they indicate that the gases flow downwards in each left-hand half, and flow upwards in each right-hand half, The partitions and partial partitions chamber are made of slate or sandstone flags, and they are carried by the pillars and girders shown in

The partitions and partial partithe figures tions in turn carry the sixteen shelves to support the lime layers, and also carry the top covering shelves. The shelves are made of slate or flagstone, the bottom and top slabs only reach to each end of the chamber, all other slabs end 6 inches short of one or other end of the chamber to allow the gases from the inlet pipe to zigzag down the one half of the chamber, as shown by Fig 23, and then zigzag up again on the other half of the chamber to the outlet pipe; all the shelves have a gap of a foot in the middle of the 21-foot dimension, and these gaps are covered by easily removable boards, as shown by B and C in Fig 23. To lime the chamber the wooden covers B are removed, and the wooden slides c are pushed on one side with the exception of the two on the bottom Sufficient lime to cover one shelf with a layer nearly 1 meh thick is then dropped through the gaps from which the covers B have been removed, and falls down to the two bottom shelves. There it is spread and then furrowed by long-handled rakes from each end

of the chamber The slides c on the bottom shelf but one are then pushed over then gaps, and the lime charges for those shelves dropped and levelled as before The test of the shelves are limed similarly till all are limed, and the top gaps covered with then boards B and made gas-tight with lime or with luting The ends of gas-tight with lime or with luting the chamber are closed by hanging iron plates on the shelves, as shown on the left side of Fig. 22, the joints are made gas-tight with luting. Each of the other six chambers is limed similarly The two top cover flags of each chamber each contain a hole provided with a water lute, the left one serving as inlet, the right one as outlet to the chamber The gas supply main is shown in Fig 22, with a lute for each of the seven chambers, and the movable aim is shown in position which passes the gas into the first The exit gas from the first chamber chamber passes into the second chamber, and all following chambers in turn by movable aims, shown in Fig 22 and in Fig 23, where one arm is marked From the seventh chamber the exit gas passes by an arm into the middle or exit main

205



shown in Fig 23, whence it is sucked by a fan, and being now free from chlorine, is blown into the air, the exit main runs the whole length of the chambers, and is provided with a lute adjacent to the exit lute of each chamber When samples taken through test holes in the end covers show that the bleach is up to strength, the gas arms are changed, so that the gas supply is cut off the first chamber, and then goes to the second, and thence, as before, to the seventh The finished bleaching powder in the first chamber is discharged by removing the end doors, pushing on one side the slides c, and then pushing the bleaching powder to the centre of the chamber, where it falls through the gaps and the hopper shown in the figures into the The chamber is next relimed, cask below. and the ends closed as before described, and it is ready to become the last chamber in the series. To effect this the exit from the seventh chamber is cut off from the evit main, and instead it is joined by a gas arm to the right-hand main shown in Fig. 23, which return main carries the gas the whole length of the chambers, and by a suitable arm is connected to the inlet


of the first chamber The exit from the first chamber is connected to the exit main Each chamber in turn is thus discharged, relimed and remserted into the cycle Such a set of chambers furnishes 4 tons of bleach per day.

Hasenclever's chambers —Another style of chamber which is particularly suitable for absorbing weak chlorine such as is produced by the Deacon process, is the mechanical chamber of Hasenclever (Eng Pat 17012, 1888), employed at Aachen, and strongly recommended by an English Commission as a thorough remedy of the difficulty arising from the contact of the workmen with the chlorine gas and the bleaching powder dust connected with the use of the Weldon chambers. To make these Hasenclever chambers suitable for use with strong chlorine they may be worked intermittently, and it has been proposed to make them suitable by mixing air with the The Figs 24 and 25 show one of the Fig. 25, in its original forms of the apparatus central portion, is an elevation, but in its upperand lower portions is a sectional elevation, and in several respects is very indistinct; Fig 24

shows on the left side an elevation of the back ends of the tubes, while on the right side it shows the front ends of the tubes with the covers removed that carry that end of the shafts. The slaked lime passes from the hopper shown in both figures into the top tube, where it is slowly worked by the blades attached to the revolving shaft towards the back end of the tube, where the first and second tubes are united by a large passage not adequately shown in Fig 25. The direction of iotation and the setting of the blades in the second tube are such that the lime is now slowly moved along the second tube towards its front end, where it drops through another large passage into the third tube, from the back of the third tube it drops again into the back of the fourth tube, and finally at the front end of the fourth tube drops into the hopper, which is closed by a sliding door. The chlorine enters at the front of the fourth tube and passes to the tubes above

by the large passages already mentioned, and the residual gases escape from the front end of the first tube as shown, to the suction fan actuating the whole Deacon plant. Fig. 25 shows four circles at the back ends of the tubes and two circles at the right ends of the tubes, and Fig 24 shows two gas passages on the middle tabes, which are antiquated, and should not have been shown By gently drawing the slide of the hopper the bleach falls direct into the cask with practically no nuisance Means are provided at the back end of the fourth tube of easily starting or stopping the shafts, and doors are provided for inspecting and, when necessary, cleaning the inside of the tubes, as shown by Fig 25, for second and third tubes, and by Fig 24 right-hand side for all four tubes Such a set of tubes makes I ton bleach per

General Remarks on the making of Bleaching Powder—Good strong bleach cannot be made






Fig. 25.

from lime that is badly burnt, slaked, or sitted, nor from chlorine that contains too much carbon dioxide or aqueous vapour or hydro-chloric acid gas, nor in very hot weather, nor if the chlorine is initially passed into the chamber so fast as to allow the temperature to rise too high It ought to consist of pretty heavy flakes, easily crushed, or a heavy but not dusty powder, which can be kneaded with the fingers into a tough paste. The laboratory test of a sample carefully taken from various parts must test at least 35 pc available chlorine, and if it is intended for export it must test still higher.

Packages.—Bleaching powder must be packed in very well-made casks, preferably made of hard wood The ends are sometimes secured by a coat of plaster of Paris. This is done in order to prevent the access of air, which quickly deteriorates the powder both by its moisture and its carbonic acid.

places; both rain and sun would quickly spoil the bleaching powder, even if placed in good casks.

Properties of Bleaching Powder. — Good bleaching powder is a nearly white powder, often mixed with lumps which, however, test as high or even higher than the powder. It has a peculiar smell, quite different from that of chlorine, and not owing to hypochlorous acid, since it is just as perceptible in alkaline solutions If it gets on to the skin a very strong odour is developed, which is highly characteristic; this odour persists even after washing the skin with soap.

In the air bleaching powder attracts moisture and carbonic acid, and gradually changes into a pasty mass. Mixed with a little water, it forms a stiff paste, with a perceptible swelling and rise of temperature; this is probably caused by the transformation of 2CaOCl<sub>2</sub> into The packages must be kept in cool, dry | Ca(OCl)2 and CaCl2, which then acts upon the

free lime to form crystals of calcium oxychloride which causes the mass to set like cement.

Bleaching powder decomposes gradually even in closely stoppered bottles or well-protected casks. The presence of light, heat, moisture, and carbonic acid greatly accelerates this change; so does shaking, e.g. in transit Hence the strength of bleach is usually only guaranteed at the place of shipment, but it ought not to lose above 2 or at most 3 p c by a few weeks' journey.

Experiments on the rate at which bleaching powder loses its available chlorine have been made by J. Pattinson, in 1874 (Chem. News, 29, 143) and 1886 (J. Soc. Chem. Ind. 587). He experimented in the first case with samples kept in loosely corked bottles, sheltered from direct sunlight, and found that these lost on an average 0.33 p.c. monthly from February to April, 0.86 from June to September, 0.28 from

November to January The greatest loss occurred in August, viz 1 4 p c per month

The keeping properties of bleaching powder depend on the quality of the line used, on the temperature conditions under which it has been made, and on its stiength. The more free calcium hydroxide is left in the bleaching powder, the more stable it is. This shows that, when bleaching powder is required for export, especially to tropical countries, it is a mistake to require a high initial test.

A few experiments made by Thummel (Arch Pharm 1834, 22, 1) prove that light has much more action on bleaching powder than exposure to air.

The following complete analyses of the samples of bleaching powder investigated by J Pattinson (v supia) may serve as an example of the best descriptions of the commercial article—

	Made of Irish limestone (from Larne)	B made of Irish limestone by another manufacturer	C made of French cliff
Available chlorine	37 00	38 30	36 00
Chlorine as chloride .	0 35	0 59	0.32
Chlorine as chlorate	0 25	0 08	0 26
Lime	44 49	43 34	44 66
Magnesia	0 40	0 31	0 43
Ferric oxide	0 05	0 04	0.02
Alumina	0 43	0 41	0.33
Oxide of manganese .	trace	trace	trace
Carbonic acid .	0 18	0 30	0 48
Silicious matter	0 40	0 30	0.50
Water and loss .	16 45	16 33	17 00
	100 00	100 00	100 00
Total chlorine	37 60	38 97	36 58

The tollowing are analyses of bleaching powder made by Lunge and Schappi, from perfectly pure lime and with the greatest care, so as to obtain the strongest possible product.—

Available chlorine			43 13
Chlorine as chlorid	е		0.29
Lime .			39.89
CO <sub>2</sub>			0.42
H <sub>2</sub> O (direct estima	tion)		$17\ 00$
Calculated as			
$CaOCl_2 + H_2O$			88 08
CaCO <sub>3</sub> .			0.96
CaCl <sub>2</sub> .			0.45
$(Ca(OH)_2)$			674
H <sub>2</sub> O not combined	•		3 77
		-	.00 00

The small quantity of unchanged calcium hydroxide is proof for the opinion that this is merely lime mechanically protected against the action of chlorine by the bleaching compound CaOCl<sub>2</sub> previously formed

Liquid chloride of lime or Bleach liquor. This substance was first prepared in 1798 by Chas. Tennant by absorbing chlorine gas in milk of lime. Many bleach works have prepared it on quite a small scale for their own use, but it is now prepared by alkali works on

a considerable scale for use in bleach works in the not too distant neighbourhood, packed in carboys or steel tank wagons, as being more convenient than the preparation of clear solutions of bleaching powder by the bleach works.

Absorption of Chlorine by Milk of Lime -As all the processes for preparing chlorine on a commercial scale produce it at approximately atmospheric pressure, the gas has to be absorbed by the surface of the milk of lime, and because of the very small solubility of lime in water agitation of the milk of lime is an absolute necessity to keep the lime in suspension and to accelerate its solution, and to bring the lime solution to the surface, and after it has absorbed its modicum of chlorine, to quickly remove it and again saturate it with lime Various forms of agitators have been used. blades attached to horizontal shafts, or to vertical shafts, or to vertical revolving cones or cylinders, and these last are considered to be the most efficient, as besides setting up a swirl in the liquid, they also set up a very rapid flow of liquor from the bottom to the top, thus very effectually and rapidly renewing the surface of the liquor.

The containing vessels have been made of lead, or flagstone, but cast iron is now employed, as it soon becomes coated with a limey deposit

both below and above the liquor, and any action of the gas or liquor is reduced to a minimum Three or more vessels are coupled up so that the gas passes them in series. Each is filled with water and lime in the proportion of 100 gallons and 10 to 15 lbs, making a milk of 12° to 18°Tw During the absorption of the During the absorption of the chlorine the temperature rises, and the admission of the chlorine should be so regulated that the temperature does not rise above 32°, or at most 37°, and for this same reason the absorbers may be cooled by building them inside a water tank. When the sp gr of the liquor after settling clear has risen to 1 040, or the contents of available chlorine is 7 pc, the gas is passed into the next absorber as first in the series, and the first absorber is allowed to settle the considerable amount of excess lime which is absolutely essential to prevent the conversion of the hypochlorite into chlorate The clear The residual liquid is then run off into carboys lime is either augmented with some fresh lime for the next operation, or if the accumulated impurities in it are too great, it is washed to recover admixed hypochlorite, and is then run away

Dissolving Bleaching Powder to make Bleach Inquor.—When bleach inquor cannot economically be obtained from the source just described, it is always prepared by the bleaching works by extracting bleaching powder with water. To avoid a loss of available chlorine and a waste of time in settling out the insoluble matter, certain conditions have to be observed, so that it cannot form lumps inside the dissolver.

1 The best practical temperature for the water is 21° to 26°.

2 The agitation must be effected by mechanical power so as to show a strong swirl on the surface, say, with a peripheral speed on the agitator of 500 feet per minute for tanks of 5 feet to 9 feet diameter, and the lower limbs of the agitator must be not more than 1 inch from the bottom of the tank

3. The bleaching powder must be emptied a little at a time on to a 1-inch mesh sieve placed outside and above the dissolver, and the passage of the powder assisted by working it with a spade, any lumps then left are to be knocked through with the flat of the spade After completing the bath, agitation for 20 minutes is ample for all strengths of solution up to 16°Tw Further time spent on agitation is wasted, and, by beating the insoluble residue, renders subsequent settling appreciably slower

4 The run-off pipe should be at such a height as to allow 5 cubic feet of sludge per cwt of bleaching powder used per batch, for all strengths of solution up to 16°Tw Unless a sufficient space is allowed, enormous waste of time will arise in waiting for the last few inches to settle clear

5. The residual sludge consists, not of bleaching powder, but of insoluble lime suspended in bleach solution. The agitating tank should be filled again with water, and the contents agitated for 5 minutes only prior to settling, when the clear liquor of about 2°Tw. is run off to be used for making the next batch, and the washed sludge is then to be run away to waste.

When the precautions given above are followed, the loss of bleaching powder in the

washed sludge will not exceed 2 lbs. per cwt. of bleaching powder used.

Properties and Uses of Bleach Liquor -The experiments made by Lunge and Landolt (J. Soc. Dyers and Colorists, 1885, Nov. 25) with bleach liquor have shown that the action of the air is nothing like so injurious to the stability of such solutions as that of the light, which ought to be excluded by all means Solu tions kept in the dark in closed vessels, remained practically unchanged up to the 24th day, and up to the 33rd day, when the experiment was finished, the strength had only decreased from 29 58 to 29 12. Kept in open vessels in the dark, they remained unaltered up to the 12th day, on the 33rd day they had lost 330 But when kept in closed vessels in diffused daylight, the decomposition set in at once, and on the 33rd day only 8 52 remained of 33 02 originally present Solutions of chloride of magnesia behaved similarly to those of chloride of lime, those of zinc or aluminium hypochlorite, as might be expected, were much less stable

The specific gravity (at 15°) can be used to approximately determine the available chloring grams per litre in a good bleach liquor

Sp gr	1 01		5 58
1	1.02		11 41
	1 03		17.36
	104		2375
	1.05		$29\ 60$
	1.06		35.81
	1 07		42 31
	1 08		49 96
	1 09		55.18
	1 10		$61\ 50$

but such a method is utterly useless for an old solution, or one that has been partly used, or is in any other way abnormal, the only way of testing such an abnormal solution is to make a chemical test, the best of which is Penot's method, with sodium arsenite in presence of bicarbonate of soda (Lunge and Hurter's Alkali Maker's Handbook, 2nd ed 126)

The use of bleach liquor for bleaching purposes is described in Cross and Bevan's Cellulose, 2nd ed 284-294. The effect of free lime on bleaching is similar to that of free caustic soda when sodium hypochlorite is employed for bleaching

The germicidal power of bleaching-powder solution is very great, and it has been used successfully for years as a disinfectant. One part of bleaching powder will, in 2-4 hours, destroy the following bacteria: coli, typhosus, and cholera in 2000 parts of ordinary sewage

(See Schumacher, in Gesundheit-Ingenieur of Munich and Berlin, August 10, 19, 30, 1905; Sheridan Dolépine, in Medical Chronicle, May, 1894, and G. C. Clayton, in J. Soc. Chem. Ind 1896, 320)

Calcium hypochlorite crystal. The Chemische Fabrik Griesheim Elektron (D. R. P. 188524, 1906) describes the production of calcium hypochlorite as follows:—

Chlorine is introduced continuously with agitation into milk of lime until the latter is nearly saturated. The solution is then concentrated in vacuo until the calcium hypochlorite crystallises out and can be dried in vacuo. The calcium hypochlorite obtained is said to be more

stable than ordinary bleaching powder, gives a clear solution in water, and contains 80-90 p.c available chlorine.

Potassium hypochlorite solution (Chloride of potash of Eau de Javel) This liquor was first made in 1789 at the Javel Works near Paris, and was the first bleaching compound known It was then made by passing chlorine into a solution of potashes (crude potassium carbonate) in eight parts of water. This liquor is rarely made at the present time, as it has been replaced by the cheaper soda compound, which has now mostly usurped its name

Sodium hypochlorite solution (Eau de Labarraque, usually called Eau de Javel). This liquor was made by passing chlorine into a solution of sodium carbonate in 10 parts of water till the liquid began to effervesce and to bleach litmus

$$Cl_2+2Na_2CO_3+H_2O$$

=NaOCl+NaCl+2NaHCO<sub>a</sub>

any further passage of chlorine produced free hypochlorous acid and CO<sub>2</sub>

The liquors containing free HOCl keep very badly, the HOCl partly volatilising and partly losing its oxygen

Hypochlorite of soda solution is used to day as a bleaching agent for the bleaching of special fabrics, in steam laundries to whiten clothes, and as a disinfectant, and is now prepared by one of the three following methods

- 1 By the precipitation of a solution of bleaching powder by sulphate or carbonate of soda, when a solution is obtained containing about 5 p c of available chlorine, but no free HClO, and which will, therefore, keep unchanged for some time. This operation is usually conducted at the factory or the place where the solution is used.
- 2. By passing chlorine into a solution of caustic soda until nearly all the soda is chlorinated

This solution is made at alkali works, and sold at a strength of 10–15 p c. available chlorine

It has been found necessary to leave a small amount of free soda, about ‡ p c. Na<sub>2</sub>O, in order to make the solution stable, by removing all free hypochlorous acid and by removing traces of iron which would otherwise be converted into sodium ferrate, and aid the decomposition by carrying oxygen from the hypochlorite to the atmosphere A very full statement on the manufacture and stability of sodium hypochlorite solution made in this way is contained in a paper by Muspratt and Smith (J Soc Chem Ind 1898, 1096)

3 By electrolysis of brine solution Under chlorine we have discussed the electrolytic cells devised to liberate chlorine from common salt solution. In the case of the preparation of sodium hypochlorite there is no necessity when the electrolysis starts to attempt to keep the main products, chlorine and caustic soda, apart, but, on the other hand, they must be brought together. But directly the formation of hypochlorites has begun then troubles begin: (1) there is insufficient agitation to bring the caustic soda to the chlorine, and consequently chlorine escapes, and free hypochlorous

acid and chlorate are formed; (2) the hypochlorite, hypochlorous acid and the chlorate are themselves electrolysed with evolution of oxygen, and the output of hypochlorite rapidly falls to nothing, (3) the hydrogen liberated at the cathode reduces the hypochlorites already formed, and again the output coases

The following cells have been introduced (see papers by W. H. Walker in Electro-Chemical Industry, 1, 439).—

The Kellner cell consists of a stoneware vessel divided into a number of small compartments by glass plates fitting into small grooves in its sides. Those glass plates are wound with platinum-indium wire, and form bipolar electrodes. The electrolyte is circulated rapidly through the cell and through a cooling coil.

The Schuckert cell is a large stonewaie trough, separated into 8-10 compartments. In these compartments are bipolar electrodes in the making of which platinum-indium wire may or may not be used. Circulation is effected by means of the hydrogen evolved, and each cell has a cooling coil

The Haas-Octel cell consists of a rectangular box divided into small compartments by a series of vertical bipplar carbon electrodes set in grooves

From the figures given by Walker, the k w h. per ton available chlorine produced in solution can be calculated, and are given in the following table for the Kellner and Schuckeit cells. The Haas and Oettel figure is taken from a paper by Reuss before the Society of Dyers and Colorists, Feb. 1911.

Name of cell		of a stan- zed cell	K W H per ton available
	Amps	Volts	Chlorine
Kellner Schuckert . Haas and Oettel	120 100 80	110 110 110	8800 8500 8400

These cells are adapted for the production of weak solutions only of hypochlorite of soda containing 1–2 pc available chlorine, as only a partial decomposition of the NaCl is possible without an enormous fall in efficiency. The cost of salt is, therefore, high, and actually about 7 tons of salt are required per ton of available chlorine produced

Cells such as these are designed for the purpose of erection at the factory or works where the hypochlorite solution is used, and where neither bleach liquor nor bleaching powder are readily available

Full details of these and other cells are to be found in the series Monographien uber Angewandte Elektrochemie, namely, vol. 8, Engelhardt, Theoretical Constructive Description, 1903, 275; vol. 17, Abel, Theoretical Description; 1905, p. 110, vol. 38, Ebert and Nussbaum, Practical Applications, 1910, p. 367, all under the title, Hypochlorite und Elektrische Bleiche.

Uses of Sodium Hypochlorite Solutions — The uses include the bleaching of such cotton

and linen fabrics as require to have the softest possible feel, and the fullest degree of whiteness, for bleaching paper pulp; for whitening clothes in steam laundries, for disinfectant purposes Hypochlorite of soda solution possesses strong germicidal powers, which Klein estimates for a solution containing 10 p c. by weight of available chlorine at 21 times that of phenol (See Klein, Comparative Bacteriological Report on Chloros, Public Health Engineer, June 9, 1906 further data, see Schumacher, Gesundheit Ingenieur of Berlin and Munich, Aug 10, 19, 30, 1905; Clayton, J Soc Chem Ind 1896, 320) And lastly, the strong solution prepared by method 2 is used in enormous quantities by the Badisch Anilin und Soda Fabrik for the conversion of phthalimide into anthranilic acid, which last is used in the manufacture of artificial indigo (Haller, Les Industries chemique et pharmaceutiques, Exposition Universelle de 1900, Paris, vol 2, pp 128-130).

Sodium hypochlorite crystal. Until recently sodium hypochlorite in the solid form was unknown Muspratt and Smith, in 1898, however, by the systematic production of a more and more concentrated solution, produced crystals of the following composition (J Soc Chem. Ind 1898, 1100)—


NaOCl	37 брс
NaClO <sub>a</sub>	nıl
NaCl.	37,
Water (by diff)	58 7 ,,
	-
	100 0

The crystals dissolve in their water of crystallisation at 18°-21°, and their rapidly decompose. By drying under vacuum part of the water of crystallisation is removed, and a more stable product obtained containing up to 60 p c of available chlorine with a melting-point of over 43° (see Muspratt, J. Soc Chem Ind. 1903, 592)

Magnesium, Zine, and Aluminium Hypochlorites, all of them obtained by adding the sulphates of these bases to a solution of bleaching powder, are sometimes used for special purposes because they have a more tapid action than chloride of lime. Their properties have been fully investigated by Lunge and Landolt in the above-mentioned memoir. They are never manufactured for sale, but prepared by the users themselves.

Potassium chlorate. This salt was originally made by saturating caustic potash solution with chlorine gas. Liebig proposed to saturate a thm paste of three molecules of slaked lime with one molecule of potassium chloride; there is, however, a disadvantage in that some potash is lost along with the insoluble impurities of the lime, and this is obviated by saturating milk of lime only, drawing off the claimed calcium chlorate liquor, evaporating it to the necessary concentration, adding potassium chloride equivalent to the calcium chlorate present, and cooling the solution to crystallise the potassium chlorate This manufacture was started in 1847 at St Helens, at Messrs. Muspratt's works, and was described in detail by Jurisch in 1888, Die Fabrikation von Chlorsaurem Kalı und Anderen Chloraten, and in Lunge's Sulphuric Acid and Alkalı. The absorption of chlorine by milk of

lime begins with the formation of calcium hypochlorite, as has already been described under the heading Liquid Chloride of Lime or Bleach Liquor, and that description applies here, and is supplemented by Fig 26, the sectional elevation of an absorber built up of east-iron plates



with rust cement joints, and 10 feet diameter and 5 feet 6 inches deep; the agitating shaft a with its aims b, b is carried by the foot step, and by the bearing attached to the non guder seen just below the crown wheel, but this girder does not cover the entire top of the tank or prevent access to the manhole f, as the figure suggests; the device of a cup d and a water lute c prevents any escape of chlorine through the shaft hole. The manhole is always open, so that there is no hindrance to the addition of water or lime or the removal of samples times a water-jacket surrounds the absorber to prevent too high a temperature, as described later At least three, better six, absorbers are worked in series on the chlorine, so as to absorb the whole of the chlorine supplied and to obtain an innocuous residual gas, the gas mains for this purpose are not shown in Fig 26, but they are exactly like those indicated in Figs 22 and 23.

The milk of lime is also of approximately the same strength as when making bleach liquor; it should not exceed  $16^{\circ}$  or  $17^{\circ}\text{Tw.}$ , which corresponds to 6.3 to 6.7 lbs CaO per cubic foot.

As the absorption proceeds the temperature slowly rises, and while an excess of lime still remains, the reaction is

$$2Ca(OH)_2+2Cl_2=Ca(OCl)_2+CaCl_2+2H_2O$$

and the temperature should not be allowed to rise above  $32^\circ$  or at most  $37^\circ$ . The reason for keeping the temperature to this limit is that, as the calcium hydrate goes more and more into solution, the impurities that originally accompanied it are more and more released, and become active catalysts, enabling the hypochlorite to decompose  $\text{Ca}(\text{OCl})_2 = \text{CaCl}_2 + \text{O}_2$ , and this catalytic action is the more active the higher the temperature, the principal impurities acting in this way are the iron and the manganese oxides. The evolution of the minute oxygen bubbles in the turbid liquor causes the formation of a very strong and persistent scum which partly or almost completely covering the surface of the liquor forms so effective a

separating layer between the chlorine gas and the alkaline liquid below that absorption of the gas is very greatly reduced or even entirely stopped. When absorption of the chlorine has gone a little further, so that the free ('a(OH), has been acted upon and only remains in small local particles, a series of complicated reactions occur which are not correctly understood, but may possibly be

$$2Ca(OCl)_2 + 4Cl_2 + 4H_2O = 2CaCl_2 + 8HOCl$$

i.e a small amount of free hypochlorous acid is formed which, however, has no effect on the remainder of the hypochlorite, but which when further treated with chlorine yields, it is assumed, chlorine monoxide

$$4HOCl+4Cl_2=4Cl_2O+4HCl$$

and it is assumed that it is this chlorine monoxide which oxidises the hypochlorite to chlorate  $\mathrm{Ca}(\mathrm{OCl})_2 + 4\mathrm{Cl}_2\mathrm{O} = \mathrm{Ca}(\mathrm{O_3Cl})_2 + 4\mathrm{Cl}_2$ , with regeneration of the free chlorine. The sum total of those reactions is

$$3Ca(OCI)_2 = Ca(O_3CI)_2 + 2CaCI_2$$

The last reaction is accompanied by the generation of so much heat that the temperature of the liquor rises exceedingly quickly up to 65°-70°, the scum previously mentioned disappears, as there are no longer any hypochlorites to decompose with evolution of oxygen, and as the reaction of the haud has changed from alkaline to acid the basic oxides in the scum pass into solution, the liquid smells of chlorine, and the dissolved hypochlorous acid bleaches litmus paper instantly; and lastly, the liquid generally has turned from colourless or light yellow to deep pink, due to the conversion of the manganese in the lime into a permanganate By summing the equations representing the formation of the hypochlorite and the conversion of the same into chlorate, the total reaction appears

 $6Ca(OH)_2 + 6Cl_2 = Ca(O_3Cl)_2 + 5CaCl_2 + 6H_3O_3$ which, though expressing the main end result, fails to represent any of the indispensable intermediate stages. The completion of the conversion of the hypochlorite into chlorate may be readily ascertained by adding to a sample of the cooled liquid some strong hydrochloric acid, there should be no evolution of chlorine gas. Full analysis of the liquor always shows that the molecular proportion of chloride is somewhat greater than indicated by the general equation above, namely 50, if much oxygen is evolved the figure may reach 5 8, but with very good work the figure may be 53 The operation lasts from less than 24 hours to 72 hours, according to the speed at which the chlorine can be passed so as to comply with the condition of complete absorption With reference to the possibility of increasing the speed of absorption, and so with efficient cooling reducing the catalytic decomposition of the hypochlorite a description by Lunge (Sulphuric Acid and Alkali) of the absorbing apparatus used by Messrs. Péchiney at Salindres is of considerable interest, the agitation being effected by means of a paddle wheel with horizontal and submerged axis, but each blade in turn projects through the liquid surface as the paddle revolves, and the absorption is excellent and superior to that in all other absorbers.

The finished liquor in the absorber may show, according to the quality and the quantity of the lime used, a strength of from 24 Tw up to 32°Tw, and statements have been made of strengths obtained up to 44°Tw. To obtain the subsequent separation of the potassium chlorate in a proper manner, it is concentrated by evaporation until it shows from 50°Tw. to 70°Tw, and during the evaporation there is considerable corrosion of the pans and also decomposition of the chlorate. For these reasons it would appear that more attention might be given to the direct manufacture of stronger chlorate liquors by an adaptation of the process referred to by Lunge as used by the Griesheim-Elektron Company for preparation of solid crystalline hypochlorite of calcium

The fimshed liquor in the absorber smells of chlorine and of hypochlorous acid, and these objectionable substances for the subsequent treatments may be removed by the addition of small quantities of sodium throsulphate or of tank waste

'The liquor is run from the absorber into a settling tank, whence the clear liquor is drawn through a deep siphon, and the residual mud is run into a filter press, or is diluted and washed by decantation, the washings being used for making fresh milk of lime.

The strong settled liquor is measured and tested for its content of chlorate, and the theoretical quantity of muriate of potash calculated, allowing for the approximately

5 p c. of impurities contained in it

The open boiling-down pans are made either of cast iron or preferably of wrought iron, as it does not crack, and is repairable when damaged by corrosion After charging with the settled liquor the fires are lighted and the muriate of potash is added. The boiling is continued until the hot liquor shows 70°Tw, when the boiling is stopped and the liquor allowed to settle. The mid consists of magnesia originally contained in the lime, or in the muriate of potash, then converted into chloride which, decomposing in the pans into magnesia and hydrochlonic acid, makes a very objectionable mud, and also corrodes the pan from which a sediment of ferric oxide results, the mud also contains gypsum from the original lime, and from the sulphates in the muriate of potash, to prevent this mud from burning fast to the pan it must be removed after each operation, and the soluble salts recovered by washing

The coolers for the first salts are generally made of non, since in no case can a pure product be obtained at this stage. They are best made like those used for soda crystals, with a plughole for running off the mother liquor. They ought to be numerous enough to permit of the liquor remaining in them 9 or 10 days in winter and 14 days in summer. Here the crude chlorate of potash crystallises in long accular crystals, totally unlike in general appearance the thin plates in which the same salt crystallises when pure, but in reality possessing exactly the same crystalline faces, only developed in different fashion. They are easily distinguishable from the large transparent prisms of hydrated calcium chloride which appear if the concentration has been too great. They must be broken up after draining, and the adhering mother liquor washed

off with a little water, and finally drained in a hydro extractor.

The mother liquor is a nearly saturated solution of CaCl<sub>2</sub>, with a little KCl, and with about 30 grams of KClO, per litre With bad work much more chlorate is left in the liquoi. In the very best case 10-12 pc of all chlorate manufactured is lost through this cause, usually much more (25-30 p c.). Usually it is not even attempted to recover the chlorine from these liquors by heating with hydrochloric acid—they are simply run to waste. But, with due care, the mother liquors can be quite well worked for chlorine by taking care to run the hydrochloric acid in pari passu with the liquor. Strictly speaking, it is not the chlorine of the chlorate which is recovered in this way, but chlorine produced by oxidation of the HCl added

Wylde, Hammill, and Auer (Eng Pat. 15097) describe a process for the recovery of about half the chlorate left in the mother liquor by refrigeration The process is described in detail by Hammill (J. Soc. Chem. Ind. 1889, 168), who states that the crude crystals obtained by refrigeration have a finer grain than the ordinary ciude crystals, but when thrown down from mother liquors of not over 72°Tw, are no more impure, and are as readily refined by one recrystallisation. Hammill says that on cooling to  $-12^{\circ}$  only 13 5 grams KClO<sub>3</sub> per litre remain in the liquois. This process has now been worked successfully for many years

Another process for greatly reducing the loss of chlorate in the mother liquors is that of Péchiney (cf Weldon, J Soc Chem. Ind 1882, 41). He concentrates ordinary chlorate liquor made from lime of a density of 42°Tw, up to 100°Tw (hot), and cools it down to at least 12°C., but not below 10°C Of its 5.5 mols of CaCl<sub>2</sub>, 43 are thus caused to crystallise out as CaCl2,2H2O (probably it should be 6H2O), but if the temperature were allowed to fall below 10°, calcium chlorate would crystallise as well The crystals are separated by a hydro extractor, and the mother liquor is treated with the calculated quantity of K(1, 1 there being only 1.2 CaCl2 present to I KICO,, the mother liquous cause a very much smaller loss than in the ordinary process Péchiney's process is specially suited, and was originally invented, for the manufacture of sodium chlorate (cf infra) It is evidently only possible to work it in the cold season, unless artificial cooling is employed

The crude chlorate of potash is always purified by recrystallisation ('finished') For this purpose it is redissolved in a mixture of water and of the mother liquor obtained in the finishing process, which is done in a lead-lined vessel, with the aid of steam, preferably indirect The solution is made at a full boiling heat, sometimes with the addition of very small quantity of sodium carbonate and of sodium sulphide, in order to precipitate any iron and lead present, it is made up to 25°Tw, and, after good settling and straining through canvas, the liquor is run into the coolers. These are made of iron lined with lead (wood lined with lead is not so good); they are either of a deep pattern, with wooden laths placed inside for the crystals to adhere to, or else they are made quite shallow, but covered over to prevent any too rapid cooling The crystallisation lasts from 8 to 14 days, according to the season.

The crystals are drained, washed on the drainer, and then again drained in a hydro extractor. This is imperative, since it is required that chlorate of potash should show little or no trace of chlorides, 005 pc. being a maximum sometimes stipulated. So long as there is PbCl<sub>2</sub> present, this cannot be very easily effected, but the lead can be removed in dissolving, as stated before. The mother liquors and washings are used over again till they become too impure, when they go back to the boiling-down pans. The crystals are dried on a hollow iron plate, covered with sheet lead,

and heated by steam

Chlorate of potash is either sold as 'crystals' or 'ground' For the former purpose the crystals tals, as they come from the drying plates, are sifted through a sieve with eight holes to the linear inch, or less, as the case may be. The powder is best ground while still slightly warm. The mills may be of any description, horizontal, edge runners, rollers, but the chlorate during grinding ought not to come into contact with iron, nor with any organic substance (straw, wood chips, &c.), or dangerous explosions may occur it must be sifted through a revolving sieve, made of miller's gauze, with mechanical brushes. The powder ought to be snow-white, and without any grittiness, like fine flour. The crystals ought to be thin, brilliant transparent plates, excessive drying will make them opaque. Both are usually packed in 1-cwt kegs, laid out with blue paper.

The loss of the chlorine used to amount to about 38 pc, with the new processes the losses are about 30 pc, or at least 25 p.c. The loss

of the potassium amounts to 25 p c

Magnesia Chlorate of Potash Process — The great loss of chlorate in the ordinary process has given rise to the process of Muspratt and Eschellmann, where magnesia is substituted for lime (Eng Pat 3960, 5183, and 5186 of 1883, and 1900 of 1885) A detailed description of this process has been given by Higgins (J. Soc. Chem. Ind. 1887, 248). The magnesia is obtained by burning Greek magnesite at not too high a temperature; when overburnt it is uscless for the absorption of chlorine. The MgO is very finely ground, suspended in water, and exposed to the action of chlorine in the ordinary octagons
The reaction is not so violent as with lime
The temperature of a finished octagon is usually 43°, and it never exceeds 60°. In the finished liquor the proportion of chlorate to chloride is very near to the theoretical, viz 1.5·1 or less, this constitutes an important advantage over the lime process, 7 p.c. less chloride being formed. No pink colour appears with very pure magnesia, but it does when it contains manganese. The of this liquor varies from 1.17 to 1 22. The strength

The octagon liquor is now boiled down to 75°-80°Tw, when on cooling about 50 p.c of the chloride will crystallise out as MgCl<sub>2</sub>,6H<sub>2</sub>O. The mother liquor contains il mol. chlorate to 2 8 chloride. Potassium chloride is now added to it in slight excess, and the potassium chlorate crystallised out. The mother liquors being

<sup>&</sup>lt;sup>1</sup> The intermediate formation of a calcium oxychloride, described in Péchiney's patents and Weldon's paper, has since been given up.

much less in quantity than with lime, and KClO<sub>3</sub> being less soluble in MgCl<sub>2</sub> than in CaCl<sub>2</sub>, the magnesia mother liquors contain much less chlorate than lime liquor (on an average 19 grams down to 10 grams per litre), and the yield of KClO<sub>3</sub> rises to 90 p.c. of the total.

The mother liquor is treated with HCl to destroy the chlorate, with due precautions against explosions. The still liquor is now neutralised with MgO; a solution of kieserite (MgSO<sub>4</sub>) is then run in to decompose the calcium chloride present, and the gypsum, along with iron oxide, is settled out. The liquor is with iron oxide, is settled out. now concentrated in cast-iron pots (wrought iron is acted upon too much); the iron dissolving in the liquor is oxidised to the ferric state, and is precipitated by MgO; the settled liquor is run into barrels, where it sets to a hard mass containing 47 p.c. of  $Mg(\mathfrak{A}_2)$  This magnesium chloride is sold to cotton sizers

This process would, of course, only become general if the magnesium chloride obtained as a by-product could be profitably decomposed for chlorine, as it is not saleable by itself in very large quantities, and could not compete in price with the Stassfurt article Such utilisation of magnesium chloride for the manufacture of chlorine had been intended by Weldon, who first proposed magnesia for the above purpose, and it is again mentioned in a patent of J Wilson (5975 of 1884). Twyman (4397 of 1881) proposes dolomite, under the idea that double chloride of potassium and magnesium (carnallite) will crystallise out of the mother liquor after the chlorate. (This seems quite hopeless, since the principal question is not that of saving KCl, but KClO3)

Electrolytic Chlorate of Potash Processes are the most recent In the preparation of chlorine from potassium chloride solution by the Griesheim process when using magnetite anodes the by-product formed is potassium chlorate, and it crystallises out in the anode cells and is recovered More direct methods are described under Sodium chlorate.

Uses of chlorate of potash are principally for the manufacture of lucifer matches, for pyrotechnical purposes, for tuses, for some descriptions of gunpowder, and other cases where it is a constituent of explosive materials. It is also largely employed, on account of its oxidising properties, in dyeing and calico printing, in the manufacture of alizarin, &c.

Sodium chlorate. This compound, which formerly was only a chemical curiosity, has become of technical importance, since it was found that it is much more suitable for the preparation of aniline black than potassium chlorate, owing to its greater solubility

It was at first thought that it could not be prepared from the calcium-chlorate liquor of the ordinary process, because NaClO3 is very soluble, and cannot be separated from CaCl, by crystallisation, and when a demand for it sprang up, it was made from potassium chlorate by means of hydrofluosilicic acid (cf Lunge's Alkali, 3, 320), and was sold at two or three times the price of the potassium salt

The process of Bottomley and Molesworth (Pat. 3005 of 1881) consists in treating a hot saturated solution of potassium chlorate with one of bitartrate of soda, and separating the precipitated potassium bitartrate from solution of sodium chlorate

The manufacture of chlorate of soda from calcium chlorate liquor was first carried out by Péchiney, by means of his process of removing most of the calcium chloride from the crude liquor by crystallisation. The purified liquor thus accumulated during the colder part of the year is mixed with a solution of ordinary sodium sulphate, that is saltcake, neutralised by lime before it is used, with continuous agitation, until all the calcium soluble salts have been converted into calcium sulphate, and final additions of either liquor are then made until neither is in excess. The CaSO<sub>4</sub>, which, under the proper conditions, is crystalline, is then separated from the liquor by filtration and washed in a hydro extractor The solution of sodium chlorate and sodium chloride is now concentrated by evaporation, during which most of the sodium chloride separates in the solid form and is fished out The remaining solution on cooling yields crystals of sodium chlorate, which are purified by recrystallisation. The purified sodium chlorate crystals contain no more than 0 16 pc NaCl. The mother liquors from the chlorate crystals, and the washings of the sodium chloride, go back into the process, and thus nearly all the chlorate contained in the liquor is obtained

Chlorate of soda is manufactured by the Muspiatt Chlorate Process by means of their magnesium chlorate liquor, freed from nearly half the MgCl, by evaporation and crystallisation (cf supra) To this liquor sodium carbonate is added, till all the magnesia is precipitated as a basic carbonate (magnesia alba), in order to be used over again in the process. The remaining solution of NaClO<sub>3</sub> and NaCl is concentrated by evaporation, with fishing out the NaCl, and after attaining 120°Tw, is allowed to cool down, the sodium chlorate crystallises out, and is purified in the usual manner

Manufacture of Alkaline Chlorates by Electro-The passing of an electric current through a solution of NaCl or KCl liberates Na or K at the cathode, and Cl at the anode The Na or K at once decomposes H O to form NaOH or KOH and hydrogen gas If the electrolyte is heated from 45° to 100°, and the caustic alkali is removed promptly from the cathode to the anode, the CI liberated is absorbed in the caustic alkali

produced to form chlorate There are, however, a number of secondary reactions, both at the anode and the cathode, and the electrolysis of the chloride cannot be carried to completion, because the product, the chlorate, is itself subjected to electrolysis and so destroyed Theoretically, a current of 1 ampere will produce in 1 hour 0.7627 gram KClO<sub>3</sub>, or 0.663

gram NaClO<sub>3</sub>
Gall and Montlaur, in their Eng. Pat. 4686 of 1887, describe the production of alkaline chlorates by the electrolysis of alkaline chlorides in a trough divided into two parts by a porous diaphragm The anodes are platinum-iridium sheets containing 10 p.c. of iridium; the cathodes are iron, though it is stated that nickel would be preferable. The electrolyte consists of a 25-p c solution of K(I, kept at a temperature of about 80°. The liquor from the cathode compartment containing the alkaline hydrate

there produced, is circulated through the anode compartment and combines with the chlorine liberated. A current density of 50 amperes per square decimeter is used, and a pressure of 4.5 to 5.0 volts is required. In the case of potassium chlorate, the anode liquor is run into crystallising vessels, where the KClO3 crystallises out, and the mother liquor, after resaturation with KCl, is passed back to the cathode compartment In the case of sodium chlorate the anode liquor has first to be concentrated to crystallise out the NaCl before crystallising out the NaClO<sub>3</sub>. The crystals obtained require, of course, to be refined by recrystallisation, as in the case of those produced by chemical processes

Hurter (Eng. Pat. 15396 of 1893) electrolyses KCl in a metal vessel which acts as cathode, and which has on its inner side a coating consisting of a mixture of Portland cement, salt and sand, which, after washing out the salt, yields a porous diaphragm. The anode is a sheet of platinum suspended in the centre of the Several vessels are superposed and worked in series, being kept apart by insulating rubber rings A solution of KCI containing a little caustic alkali is run into the top vessel, from which it overflows, through a glass or ebonite pipe, into the cell below. The hydrogen evolved escapes by pipes from the upper part of each compartment

Blumenburg (Eng. Pat 9129 of 1894) describes a cell with a porous diaphragm in which the anode compartment is closed and connected by a pipe with the bottom of the cathode compartment, so that the chlorine liberated at the anode passes into the liquor in the cathode compartment, and there combines with the caustic alkali produced, the tempera-ture being maintained at a temperature of about 49°. In the Ger Pat 89035, and also in the US Pat. 537179, Blumenburg passes the chlorine gases from the anode compartment into a gas holder, from which they pass to a separate combining vessel, where they are absorbed in the alkaline liquor from the cathode compartment. Chlorate is also formed as a by-product in the Griesheim cell described above, when using the oxide of iron anode

In 1899 Imhoff made a great advance in the electrolytic production of chlorate by his discovery of the action of small quantities of chromate of potash or soda in the electrolyte (US Pat 627063 of 1899) The action of the chromate is to form an insoluble film around the cathode, and so to prevent the reduction of hypochlorite and chlorate by the nascent hydrogen evolved at the cathode, and has resulted in a very marked and important improvement in the current efficiency

Details of the electrolytic chlorate processes up to 1903 were collected by J. B. C. Kershaw and published as vol. 19 of the Monographien der angewandte Elektrochemie, under the title, Die elektrolytische Chloratindustrie, 1905,

p 122

Perchlorates of Potassium, Sodium, and Ammonium. That by the electrolysis of chlorates there were formed perchlorates, was first shown in 1816 by Stadion, and in 1847 confirmed by Kolbe With the aid of such electrolytic methods the above-named perchlorates have been manufactured in Mansboe, Sweden, since 1895, in considerable quantities (Carlson, Zeit. für Elektrochemie, 6, 471), and since 1901 in Chedde, France (see Lederlin, D. R. P. 136678). The ammonium salt is used in the manufacture of explosives. Laboratory experiments are described by Foerster (Zeit. fur Elektrochemie, 4, 386) and Winteler (Chem. Zeit. 22, 90); see Kershaw, Elektrolytische Chloratindustrie, p 67.

Hydrochloric Acid. Commercial hydrochloric acid is a more or less pure solution of hydrogen

chloride HCl in water.

Hydrogen chloride is a colourless gas, of sp gr 1 2596 (air=1), and molecular weight 36 457 By weight it consists of 2 74 parts of hydrogen and 97 26 parts of chlorine 1t can be condensed into a liquid by cold and pressure. Its m p is  $-111~4^{\circ}$ , and b p.  $-83~1^{\circ}$ , and its critical constants  $+51~4^{\circ}\pm0~1^{\circ}$  and  $81~55\pm0~15$ atmospheres.

The vapour pressures can be expressed between 10° and 51 4° by the formula:

 $\log p = 1.152048 + 0.010312(t-10.2)$ 

 $-0.000019989(t-10.2)^2$ 

and between  $-24.2^{\circ}$  and  $10^{\circ}$  by the formula.  $\log p = 1\ 10106 + 0\ 013277(t + 24\ 2)$ 

 $-0.000039978(t+24.2)^{2}$ 

(Cardoso and Germann, J. Chem. Phys. 1912, 10, 517; 11, 632).

The heat of formation of a gram-molecule of

HCl is 22001 cals

Hydrogen chloride has a great affinity for water, and is greedily absorbed by it. The solution is the ordinary hydrochloric acid which, when somewhat concentrated, emits fumes of The following HCl on exposure to the air table by H Deicke shows the maximum solubility of HCl in water at varying temperatures. 1 c c dissolves at the temperature  $t^{\delta}$  from an atmosphere of pure HCl at 760 mm pressure —

-	-		•
t°	Vols HClat 0° and 760 mm dissolved in 1 vol H <sub>2</sub> O	Specific gravity of the satu- rated solution	HCl by weight
o°С.	525 2	1 2257	45 148
4	494 7	1.2265	44 361
8	$480\ 3$	1 2185	43 828
12	471.3	1 2148	43 277
14	462 4	1 2074	42-829
18	451 2	1 2064	42 344
23	435 0	1 2014	41 536
	1	1	

The behaviour of aqueous solutions of HCl under varying conditions of temperature and pressure has been studied by Roscoe and Dittmar (Chem Soc Trans 12, 128) They found that concentrated hydrochloric acid on heating loses both gas and water, but as the proportion of HCl to H2O is higher in the vapour than in the original liquid, the residual fiquid gradually becomes weaker, until a sp gr of 1:101 and a percentage of 20 24 HCl is attained at a boiling temperature of 110°; the acid in this state distils without change, provided the atmospheric pressure is 760 mm. At a lower pressure the acid distilling unchanged is stronger, at a higher pressure it is weaker. A similar, but not identical

limit is attained when strong acid is exposed to the air at ordinary temperatures.

The following table gives the percentages of HCl in aqueous hydrochloric acid at various specific gravities and at a temperature of 15°:—

Deg. Twaddell	Spec. gravity	Percentage of	Grams HCl per litre	1 cubic foot weighs lbs	1 cubic foot con- tains lbs of HCl
1	1 005	1 12	11 32	62 66	0 70
2	1 010	2 12	21 45	62 97	1 34
3	1 015	3 12	31 67	63 29	1 97
4	1 020	4 11	41 99	63 60	2 61
5	1 025	511	$52 \ 41$	63 91	3 26
6	1 030	6 11	62 93	64 22	3 92
7	1 035	7 10	73 55	64 53	4 58
8	1 040	8 10	84 27	64 84	5 25
9	1 045	9 10	95.09	65 16	5 93
10	1 050	10 09	106 01	65.47	6 61
11	1 055	11 09	117 02	65 48	7 31
12	1 060	12 09	128 14	66 09	8 00
13	1 065	13 08	139 36	66 40	8 69
14	1 070	14 08	150 68	66 71	9 40
15	1 075	15 08	162 10	67 03	10 11
16	1 080	16 07	$173 \ 62$	67.34	10 83
17	1085	17 07	185 24	$67\ 65$	11 55
18	1 090	18 07	196 96	67.96	12 28
19	1 095	19 07	208 78	$68\ 27$	13 02
20	1 100	20 06	220 70	$68\ 59$	13 76
21	1 105	21 06	232 68	$68 \ 90$	14 51
22	1 110	22 06	244 80	$69\ 21$	15 27
23	1 115	23 05	257 02	$69\ 52$	16 04
24	1 120	24 05	269 34	69.83	16 79
25	1 125	$25\ 05$	281 76	70 14	17 57
26	1 130	26 04	294 28	70 46	$18\ 35$
27	1 135	27 04	306 90	70 77	19 14
28	1 140	28 04	319 62	71 08	-19.93
29	1 145	29 03	332 44	71.39	20.73
30	1 150	$30 \ 03$	345 36	71 70	21.54
31	1 155	$31\ 03$	358 34	72 02	$22\ 35$
32	1 160	$32\ 02$	371 44	$72\ 33$	$23\ 16$
33	1 165	$33 \ 02$	384 64	72.64	23 99
34	1 170	34 02	397 94	72.95	24 82
35	1 175	35 01	411 34	73 26	$25\ 65$
36	1 180	36 01	424 84	73 57	26 49
37	1 185	37 01	438 44	73 89	$27 \ 34$
38	1 190	38 01	452 14	74 20	28 20
39	1 195	39 00	466 00	74 51	29 06
40	1 200	40 00	479 84	74 82	$30\ 00$

Lunge and Hurter's Alkali-Maker's Handbook gives somewhat lower figures, they also give the following table for correcting the hydrometer readings for varying temperatures

# THE MANUFACTURE OF HYDROCHLORIC ACID

In former days small quantities of acid were produced by decomposing common salt with sulphuric acid in east-iron cylinders, and in order to utilise the more expensive sulphuric acid, an excess of common salt was used. The solid product left in the cylinder consisted of sodium sulphate, mixed with 20–30 pc. NaCl and was commercially known as cylinder cake. The production of hydrochloric acid from cylinders has, however, practically ceased.

Some time after the establishment of the Leblanc soda process in this country all the hydrochloric acid of commerce was obtained as a by-product of that process, at first the process produced enormously more hydrochloric acid gas than could be sold as hydrochloric acid liquor,

	ိုင္ပ	10°	12°	200	950	303	350	400	450	:02	0.1.5	600	0.7.0	2.6	1	100	0,0	000	010	1000
		-					3			3	3 ;	3	3	2	2	00	8	an	66	8
1 168	1 165	1 163	1 160	1 157	1 154	1 152	1 1.40	-	1 111	1 149		1 190	1 196	1 100	1 191	190		30	1.09	1.191
1 158	1 155	1 153	1 150	1.147	1 145	1 1 1 2 2	1 130	137	1 134	1 132	130	1 136	061 1	1 199	1 191	1139	177	621 1	611	1.15
1 1 <del>1</del> 8	1 145	1 143	1 140	1 137	1 134	1 132	1 1 2 9	121	195	1 193	200	971	116	1 1 2	171	901	901	#11 T	201	1 000
1 138	1135	1 133	1 130	1.127	1 125	1 122	1119		1 114	113	601	107	152	100	100	1 007	002	1003	60	880
1 128	1 125	1 123	1 120	1-117	1 115	1 112	1 110	108	1 106	103	101	660	960	160	1 091	080	980	1 084	200	1 079
1118	1 115	1 113	1 110	1 107	1 105	1 103	1 101	1 099	1 097	1 094	1 093	060	880	280	1 083	080	220	1 075	1 073	1 070
108	1.105	1 103	1 100	1 097	1095	1 092	1 090	1 088	1 086	1 084	1 082	080	1 078	920	1 073	1 07	640 T	1 066	1 064	1 061
1 098	1 095	1 093	1 090	1 087	1 085	1 082	1 080	1 077	1075	1 073	1071	1 069	1 067	065	1 063	1.00	1 059	1 057	1 055	1 053
1 088	1 085	1 083	1 080	1 077	1 075	1 073	1 070	1 068	1 066	100	1 ()62	1090	1 058	1 056	105	1 053	1 051	070	1 047	1 045
1 078	1 075	1 073	1 070	1 067	1 066	1063	1 061	1 059	1 057	1 055	0.53	103	070	1 078	1 ()46	1 014	1 0.13	1 041	1 039	1 037
1 068	1 065	1 063	1 060	1 057	1 055	1053	1 050	1 048	1046	7	1 042	500	038	1 036	1 034	1 033	1 031	1 090	1 027	1 025
1 058	1055	1 053	1050	1 047	1 045	1 043	1 040	1 038	1 035	1 033	1 031	0.00	1 0 2	1 055	1 023	1 091	1010	201	1 015	1 013
1 048	1 045	1 043	1 040	1 037	1 035	1032	1 030	1 027	1 025	1 022	1 090	300	910	101		100	201	961	000	38
1 038	1035	1 033	1 030	1 027	1 024	1 022	1 019	1 017	101	1 012	200	88	002	100	100	000	0 007	0005	000	0 991
1.028	1.025	1.023	1 020	1 017	1 014	1 012	1 009	1 00 1	100	1.007	200	8660	0 005	0.003	1660	080	0 0 0 0	000	0.083	0 981
1.018	1 015	1 013	1.010	1.007	1 004	1 002	0 0 0 0 0	0 997	766 O	0.992	066 0	0 988	0.985	0 983	0 981	0.07	126.0	0.975	0 973	0.971

and some of the difficulties encountered in consequence are detailed in the following account.

New sources of hydrochloric acid arose when acctic acid began to be chlorinated by the Badische Anilin und Soda Fabrik in great quantities for the manufacture of monochloracetic acid and artificial indigo, and as one after another organic compound came to be chlorinated. Lastly, since the output of chlorine from the electrolysis of the chlorides of potassium and sodium was not sufficiently readily taken up, processes are now proposed of converting chlorine into hydrochloric acid either with the aid of hydrogen obtained in the decomposition of wood or of the hydrogen obtained at the same time as the electrolytic chlorine.

Hydrochloric Acid obtained in manufacturing Solium Sulphate (Salt Cake). When Leblanc decomposed salt in order to obtain sodium sulphate he proposed to convert the hydro-chloric acid gas evolved into ammonium chloride; but when the Leblanc soda process was introduced into this country that part of his suggestions was not carried out, and the acid gas was regarded as possessing no value, and it was simply allowed to escape into the air. This continued to be the case even after that manufacture had attained a comparatively large extension, but the damage caused to the neighbourhood by the acid gases soon brought about the intervention of local authorities and courts of law, and manufacturers were compelled to pay compensation for the damages they had done Repeated penalties compelled the alkali manufacturers to seek means for condensing the acid gas This was first successfully done in 1836 by Gossage's coke towers, but for many years after this there was still considerable damage done by escaping acid gas. The continued disregard by the old alkali manufacturers for the rights of their neighbours led in Belgium, in 1856, to a Parliamentary Commission, and in this country in 1863 to Lord Derby's Alkali Act, and the several subsequent Acts, by means of which British alkalı makers were compelled to adopt thoroughly efficient means for condensing their hydrochloric acid, and this has benefited all other manufacturing countries as well

The reports made by the inspectors under the Alkah Acts form an invaluable record of all the various steps taken for improving the condensation of hydrochloric acid, and have helped to spread a better knowledge of that process and have caused manufacturers to improve it up to the highest standard of efficiency

The Alkah Acts of 1863 made it incumbent upon those decomposing salt to allow no more than 5 pc of the hydrochloric acid to escape into the atmosphere. An additional Act of 1874 prescribed that no more than 0.2 grain of HCl might be present in a cubic foot of the gas escaping from the factory into the air. This corresponds to 0.454 gram HCl per cubic metre, or about three ten-thousandths by volume. The latter clause applies to the chimney gases as well, and not merely to the gases escaping directly from the condensing apparatus. Although these requirements were at first declared by many to be impossible of fulfilment, the actual results proved at once that

they were quite possible, and the escapes registered by the Alkalı Inspectors have averaged far below the limits assigned by the Acts. In 1885 only 2 13 pc of the total hydrochloric acid was allowed to escape, and the chimney and other gases escaping into the air contained only 0 10 grain HCl per cubic foot. No change has been made in the limit fixed in 1874, and the working of that Act remains satisfactory Compare the survey of these Acts by the Chief Inspector under these Acts in J. Soc. Chem. Ind 1892, 120.

Although the various processes and apparatus used for the manufacture of sulphate of soda produce gases differing within wide limits in temperature and concentration of HCl, an efficient condensation, i.e. practically complete absorption of the HCl in water to form a liquid acid of 1 15-1 17 sp gr. is usually obtained

acid of 1 15-1 17 sp gr. is usually obtained

The efficiency of the condensation of gaseous
HCl in water is dependent on three considerations:

l Temperature of the gas and of the liquid.

2 Concentration of HCl in the gases to be scrubbed and of the acid liquid to be produced

3 Time of contact between the gas and the liquid, and the average distance between the gaseous particles and the liquid particles, and also the thickness of the liquid layer, and on its motions internal and as a whole.

On issuing from the sulphate of soda plant the acid gases require cooling. The apparatus

used for this purpose depends on the temperature of the gases.

If the gases issue from the pot or pan of a handworked saltcake furnace, the temperature is comparatively low. In this case the cooling is usually done in a row or rows of earthenware pipes. The pipes used are often unglazed, but are rendered impervious to moisture and gas by boiling in tar In some cases the earthenware pipes are substituted by glass pipes of 12 inches diameter, made slightly taper, as shown in Fig 27. glas pipes are, of course, more liable to breakages by accident, and also through the effect of violent changes of temperature, than the earthenware pipes, but in sheltered positions they last well.

For the hotter gases issuing from the furnace of an ordinary muffle hand sulphate of soda furnace, or from the cylinders of the Hargreaves process, castron pipes are usually used to cool the gases down to a safe temperature before passing them as before into earthenware or

glass pipes The action of the hydrochloric acid gases on cast iron, when at a temperature well above condensing-point of liquid acid, is very slight.

When open furnaces, whether 'hand' or 'mechanical' (such as the Mactear furnace), are used, the acid gases are mixed with the products



Fig 27.

of combustion, and are consequently both much 45°, or even to 25°. The cooling of weak gas is hotter and more dilute. Under these conditions far more important than the cooling of strong gas

extensive series of cast-iron pipes are used for the preliminary cooling before passing the gases into the earthenware pipes for

final cooling.

The earthenware pipes used are usually 12-inch or 15-inch diameter socket pipes, the joints being made with a cement of tar and China clay. The castiron pipes vary from 12 to 20 inches in diameter, according to the quantity of gas, and the arrangement of the cooling system of pipes; they are socket pipes, the joints being made by ramming into the socket iron filings moistened with ammonium chloride solution.

The arrangement of the cooling pipes depends on the relative positions of the salt-cake furnaces and of the acid condensers and the space available

When the salt-cake furnace is a long distance from the condensers, a single direct line of pipes may be a sufficient means of cooling; where the distance apart is small the necessary amount of cooling surface is obtained by adding extra lengths of pipe that run vertically up and down and that start from a small stone eistern and return to another stone cistern for the purpose of collecting any liquid deposited




Fig. 28

The cisterns are usually made in this country of silicious sandstone found in Yorkshire, and known as 'Yorkshire flag.' The cisterns




Fig. 29

of flags 4-6 inches thick, bound together by strong iron ties, and the joints made tight by thick ındıarubber cord laid in a groove between the faces of the

are composed

two stones or by a mixture of tar and fire clay, as in Fig 30 By one or more of the above arrangements of pupes and customs the acid gases are cooled to




Fig. 30.

The gases next enter the apparatus where the hydrochloric acid gas is to be converted into a liquid acid of commercial strength, and that in such a manner that practically hardly

a trace of acid escapes

condensation The apparatus almost universally used are tall stone towers, packed with broken coke specially arranged, down which a stream of water is allowed to trickle. and up which the gases are caused to flow (Fig. 31) These towers were invented and patented Gossage, in 1836.

The coke tower has the merit of simplicity, and, when carefully packed, of efficiency

The comparative efficiency of various modes of treating liquids with gases has been dealt with in a series of papers by Hurter, who also gives some interesting data on cokepacked towers (J Soc Chem Ind 1885, 639; 1887, 707; and 1893, 227 and 989).

Coke towers are erections of considerable height, varying from about 40 feet to 60 feet. The hori-




Fig. 31

zontal sectional area may vary from 25 to 180 square feet. They are provided with a grating a few feet over the bottom, on which

rests the coke which forms the 'packing' of This coke, which must be of as the tower hard a quality as possible (only the best oven-coke is applicable for this purpose), is disposed in such a way that the gas is as much as possible divided in a large number of channels (none of them being so wide that the bulk of the gas can pass up through it), and that the current of gas is constantly changed in direction. For this object long pieces of coke are laid parallel in a row, the next row is made to cross the The size of the pieces lower one, and so forth is gradually diminished from bottom to top. In order to avoid an excessive quantity of the gas passing up the side of the condenser along the wall, it is usual to place a layer of finer coke round the condenser for a few inches from the By this arrangement there is effected both a great division and constant mixture of the gases, and at the same time a suitable



Fig 32

spreading of the feeding water over a very large surface. One great advantage coke has over flints or other packings is that, owing to the roughness of its surface and its porousness, it always retains a large store of liquid, which tends to keep the working of the condensers steady under a fluctuating flow of acid gases, and to tide over difficulties occasioned by any unintentional stoppage of the liquor supply. The water (or sometimes weak acid) is fed in at the top by some contrivance assuring a uniform distribution over the whole area of the tower, and in trickling down it keeps the surfaces of the coke moist, and a very large area of contact between the gases and the liquid is thus pro-These towers are so simple to construct and so effective that of recent years there has been a growing tendency in England to do away with all cooling pipes, cisterns, wash towers, &c, and to use a couple of very large stone towers to do the whole work of cooling and condensation for two or even more salt-cake pots and furnaces. The towers are worked in series, the weak acid from the second or final tower being run down the first tower. In this way both complete absorption of the acid is assured, and the production of a strong liquid acid, i.e. one of over 1.15 sp.gr. To condense the acid from two-hand salt-cake furnaces the two towers may each be 5 feet square and 50 feet high

No difficulty is now experienced in pumping the weak acid made in the second condenser on to the top of the first condenser, either by stoneware ram pumps, chonite ram pumps,




Fig 33.

stoneware acid eggs, or occasionally even yet by the Hazlehurst membrane pump, patented in 1876 The ascension pipe is usually of stoneware or ebonite, occasionally of lead

Brick towers are not to be recommended, as it is practically impossible to keep the numerous joints tight against acid oozing out. Columns of stoneware pipes cannot be made of a large area, and, therefore, are not very well adapted to large works, at least if intended for coke towers, but they do very good service at smaller works, or as supplementary to a large condensing plant of stoneware receivers. The material mostly employed for coke towers is acid proof sand-tone or, in France, volvic lava, just as

for acid tanks. The system of joining the flags together is the same as shown in Figs 28 and 29, or else as in Fig. 30, or as shown in Fig. 32, in which the flags are kept together by means of grooves worked in the stone, and without the aid of any ironwork, which is always a source of anxiety in acid condensers.

The coke towers are generally placed at a high level, and are, if necessary, put upon pillars, for the purpose of running the acid from them by gravitation either into chlorine stills, or into store cisterns for sale They must, of course, have a very secure foundation, in the construction of which not merely the great weight of the tower must be considered, but also the probability that there will be now and then leakages of acid which may make the foundation unsafe. This can be avoided by asphalting the ground in such a way that any acid running down cannot penetrate into the soil, and is conveyed to a safe distance Any damage done to the foundation, even if it amounts to much less




Fig. 34.


than would cause the erection to break down altogether, is extremely injurious, because it will almost unavoidably lead to the tower getting out of level, in which case the liquid will principally descend on the lower side, thus the necessary distribution of the liquid is not attained, and the action of the condenser is most injuriously affected

It has been already stated that the feeding of the towers with water or with weak acid must take place in such manner as to distribute it equally over the whole surface of the coke. Formerly this was generally done by a distributing wheel (shown in Figs 33 and 34) worked by the pressure of the feed water itself 33 ought to show two jets feeding the outer urcle of holes This causes the stream of water to reach successively every one of the holes on the top of the condenser, and thus secures a aniform distribution without depending upon the top of the tower being at a mathematical level.

Recently most manufacturers have done

away with these acid wheels, and have introduced troughs provided with a number of overflows all at the same level, feeding the same number of 'lutes' This arrangement is indicated on Fig 31. This figure represents a tower built as Fig 32, the cross-hatched outer portion representing the timber framework to support the feeding cistern

Another form of apparatus for dissolving the hydrochloric acid gas is the Lunge Rohrmann plate tower, shown in Figs 35, 36, and 37.



Fra 35.

It consists of an earthenware tower packed with a series of perforated plates E, E, the perforations being made to alternate in successive plates, so that each hole corresponds to a solid place in the plates just above and below 36 and 37 show one of the plates in detail. As is to be expected, the evidence for and against the improved packings for the condensing towers is very conflicting, and as the






Fig. 36.

Fig 37

cost of them is naturally higher, the coke-packed tower has fully maintained its position.

In France and Germany a somewhat different system of cooling and condensation is in general use. The earthenware and glass pipes mentioned above as being used almost exclusively in this country for the cooling of the gases are more or less completely substituted by a series of earthenware Woulffe's bottles, known technically as bombonnes or touries (see Fig 38) The figure is drawn too small to show that the vessel on the right is about 1 inch higher than the vessel on the left; the liquids travel down hill from right to left, and consequently the gases are made to travel from left to right The size of the bombonnes varies from 40 to 100

gallons. They are arranged in sets of 40 up to 100, as shown in Fig 39, and, in spite of the relatively high temperature in the leading



PTG. 3N

bombonne, eventually gives a liquid acid of 1.15-1.18 sp gr They are made of the best stoneware, which is able to stand somewhat violent changes of temperature. After leaving the bombonnes, the acid gases are finally

scrubbed with water in a coke-packed tower, similar in design but usually much smaller than those used in this country The weak liquid acid produced in this tower supplies the stream of weak acid required by the bombonnes The figure does not show the means of running off the strong acid from the lowest bombonnes in the series, nor the cisterns, &c, for receiving the acid.

In the condensation of hydrochloric acid gas in water with the apparatus at present used and described above, there is little reason for difficulty in avoiding losses of HCl through in-complete absorption The difficulty more commonly experienced is to combine the complete absorption enforced by law with a high-strength liquid acid

The impurities found in commercial hydrochloric and made by the above processes are, chiefly, sulphuric acid, ferric chloride, arsenic trichloride, and either free chlorine or sulphurous acid

Sulphuric acid is the largest impurity. The hydrochloric acid condensed from the pan or pot gas is always much purer than that from the roaster gas. The quantity of sulphuric acid in

the gases from the salt-cake furnace sometimes amounts to 2 pc. and upwards; it is very troublesome in the Weldon, and fatal in the Deacon, chlorine process, and in some other




Fig 39

applications of hydrochloric acid. The suldenser or dry scrubber before condensation phuric acid in the salt-cake furnace gases can be removed to a large extent by a small confort the purification of the salt-cake furnace

gases, so as to render them suitable for the Deacon process, but the only one of these processes to achieve practical success is the Hasenclever process (Eng Pat 3393 of 1883). (See at end of Deacon Chlorme Process)

Arsenic gets into the hydrochloric acid through the sulphuric acid employed in its manufacture, the arsenic is thereby converted into AsCl<sub>3</sub>, and passes as such into the condensing apparatus. Many plans have been proposed for its removal-most frequently a treatment with sulphuretted hydrogen or with sulphides. Bettendorf (Dingl. poly J. 194, 253) precipitates the arsenic from concentrated hydrochloric acid by stannous chloride, and then distils the decanted acid. Duflos dilutes the acid to sp gr. 1:13, and digests it with strips of sheet copper at 30°C for 24 hours, and repeats this treatment with freshly scoured copper This removes all the arsenic and the free chlorine, and reduces the ferrie chloride to ferrous chloride, which remains behind on Beckurts (Fischer's Jahresb 1884, distillation 348) distils hydrochloric acid with ferrous chloride and removes the first 30 p.c. containing all the arsenic

Selenium is sometimes found in such quantities in hydrochloric acid as to be troublesome in its application (Davis, J. Soc. Chem. Ind.

1883, 157)

Hydrochloric Acid from other Sources than the Attempts have been made to Leblanc Process produce hydrochloric acid from many sources, notably from the residual liquors of the ammoniasoda process A number of suggestions have been made for the direct production of HCl. 1st, from CaCl2, 2nd, from MgCl2, 3rd, from NH Cl In this connection compare references given for the production of chlorine from these bodies earlier in the article. As far as is known, hone of the processes is used. Mond (1883) heats the ammonium chloride with so much sulphuric acid that the acid sulphate is formed and all HCl is expelled. The acid sulphate is converted by means of ammoniacal vapours into the neutral salt of commerce. O. N. Witt (1886) expels the HCl from NH<sub>4</sub>Cl by means of syrupy phosphoric acid, which forms ammonium phosphate On heating this at a higher temperature, the ammonia is driven out as well, and the phosphoric acid is regenerated Jurisch (Dingl. poly J 267, 431) has shown that this process is not placticable, because there is no material known which resists the fusing phosphoric acid, and only 63 to 86 p c. of the NH3 is recovered. Mond passes the vapour of ammonium chloride over nickel oxide, heated to at least 350°, which absorbs the HCl and allows the NH3 to pass on a certain time the process is changed by raising the temperature to 500° or 600° and decomposing the nickel chloride by steam, when HCl is driven off and NiO regenerated (Several patents taken out in 1886)

During recent years, when the chlorination of organic compounds has grown by vast strides, the hydrochloric acid gas produced when a hydrogen atom is substituted by a chlorine atom, has had to be collected and These reactions are referred to in a utilised previous page when describing the uses of chlorine. The hydrochloric acid gas is evolved HYDE

almost pure, is easily condensed, and yields an acid quite free from the usual impurities

Hydrochloric acid has also been produced by passing chlorine over wood at a temperature of 150°-350°, whereby charcoal is also formed (Bosnian Electricity Works at Jacie, D. R. P. 158086)

Lastly, hydrochloric acid has been prepared by processes inverting the practices of the last hundred years, namely, it has been prepared from its elements as a direct and definite process without the formation of by-products, or for any other reasons than its own preparation and the utilisation of the two elements formed by the electrolysis of potassium and sodium chlorides. The gases are mixed and passed over charcoal (D R. P 114219), or they are burnt together at jets (Soc Italiana di Elettro-chimica; J. Soc. Chem Ind 1915, 1142), or they are combined together by Roberts' patent (Electrochemical and Metallurgical Industry, 1910, 704), and by this last process the acid has been manufactured in Niagara since 1914.

The conveyance of hydrochloric acid mostly takes place in glass carboys of 10 or 12 gallons capacity with basket or wire hampers, sometimes in earthenware carboys of the same size, and recently hard white vitrefied earthenware jars have become available At a large French works, some years ago, tank waggons were used, carrying iron cylinders lined with ebonite.

Tank waggons have of late years come into general use, especially on the Continent consist of a waggon platform, carrying a number of stoneware jars of about 1 ton capacity each Fig. 40 shows such a waggon carrying 12




Fig. 40

jars, each of which holds I ton of acid. Specially prepared wooden tanks are also used occasionally.

The applications of hydrochloric acid solu-tion are, first and foremost, for producing chlorine by the Weldon process, for the cleaning of thin sheet iron when the same is to be coated with zinc, tin, or lead; for extracting the phosphates from bones and from bone char-for which purposes it must be quite free from sulphuric acid, for preparing zinc chloride, and zine ammonium chloride from the residual 'flux skimmings' of galvanising works, for souring goods during the process of bleaching; for neutralising the alkaline melts in the manufacture of alizarin, resorein, &c , for the manufacture of aniline hydrochloride, stannous chloride, ferric chloride, for the preparation of pure carbonic acid from limestone; in dyeing and tissue printing; and in the manufacture of many coal tar colours and intermediates, &c

ω-CHLOROACETOPHENONE v KETONES CHLOROBENZALDEHYDE v. BENZALDE- CHLOROBENZENE v. PHENYL δ-CHLOROBENZENEDISULPHONIC ACID

C<sub>6</sub>H<sub>5</sub>Cl(SO<sub>3</sub>)<sub>2</sub>H<sub>2</sub>. Formed by heating at 300° chlorobenzene with five times its volume of fuming sulphure acid containing 20 pe of sulphur trioxide Forms barium, potassium and ammonium salts, a dichloride, mp 105 5°-106° and a diamide, mp 223°-224°. The dichloride heated under pressure at 200°-210° with phosphorus pentachloride yields δ-trichlorobenzene (Olivier, Rec. trav. chim. 1918, 37, 307)

p-CHLOROBENZENE DIAZOCYANIDES v. DIAZO- COMPOUNDS

CHLOROBROM. Trade name for a mixture of equal parts of potassium bromide and chloral formamide dissolved in water.

CHLOROFORM, Trichloromethane CHCl3. Chloroform was discovered by Liebig in 1831

(Pogg. Ann 23, 242), and independently, about the same time, by Souberran (Ann Chim Phys [2] 48, 131). Its actual composition was first determined by Dumas (Ann Chim Phys [2] 56, 115) to whom is also due the name by which it is now known Chloroform is largely employed as an anæsthetic and is an important solvent, finding technical application in the manufacture of fats, rubber, resins, alkaloids, It is also a useful antiseptic

Formation -1 By the action of alkalis on chloral (Liebig, Pogg Ann 23, 442, Annalen, 1, 199; 162, 161)

CCl<sub>2</sub>CHO+NaOH-=CHCl<sub>2</sub>+HCOONa

2. By the action of water, ammonia, or alkalis on trichloracetic acid (Dumas, Annalen, 32, 101, Beckurts and Otto, Ber 14, 589)

$$CCl_3COOH + HOH = CHCl_4 + CO_2 + H_2O$$

3 By the action of bleaching powder (chlorine) on ethyl alcohol or acetone In the case of alcohol the reaction has been represented by the following equations ·

by the Contact (a) CH<sub>3</sub>CH<sub>2</sub>OH+CaOCl<sub>2</sub> = CH<sub>3</sub>CHO+H<sub>2</sub>O+CaCl<sub>2</sub>

(b) 2CH<sub>3</sub>CHO+3CaOCl<sub>2</sub> =2CCl<sub>3</sub>CHO+3Ca(OH)<sub>2</sub>

(c)  $2\text{CCl}_3\text{CHO} + \text{Ca(OH)}_2$  =  $2\text{CHCl}_3 + (\text{H COO})_2\text{Ca}$ 

According to Dott (J Soc Chem Ind 27, 271), however, the reaction is best indicated by the equation

3C<sub>2</sub>H<sub>5</sub>OH+8CaOCl

=2CHCl<sub>3</sub>+3CaCO<sub>3</sub>+CO<sub>2</sub>+H<sub>2</sub>O+5CaCl<sub>2</sub>no appreciable amount of calcium formate being found in the products of reaction

According to Feyer (Zeitsch Elektrochem 1919, 25, 115), the formation of chloroform from alcohol occurs in three stages. first, an exidation of the alcohol to aldehyde; secondly, a formation of hypochlorite, and thirdly, an interaction between the aldehyde and the hypochlorite:

(1) CH<sub>3</sub> CH<sub>2</sub> OH+HOCI

$$=$$
CH<sub>2</sub>CHO+H<sub>2</sub>O+HCl,

(n)  $OH'+Cl_2=H'+Cl'+ClO'$ ;

(III) CH<sub>3</sub> CHO+3HOCI

$$= CHCl3 + HCO2H + 2H2O,$$
(iv) H CO<sub>2</sub>H + HOCl = CO<sub>2</sub> + H<sub>2</sub>O + HCl.

Equation (1) is partly replaced by the reaction

 $CH_3 CH_2 OH + O = CH_3 CHO + H_2O$ 

expressed thus (Orndoff and Jessel, Amer. Chem. J. 10, 363):

 $\begin{array}{l} 2 {\rm CO(CH_3)_2} + 6 {\rm CaOCl_2} \\ = 2 {\rm CHCl_3} + 2 {\rm Ca(OH)_2} + 3 {\rm CaCl_2} + ({\rm CH_3COO})_2 {\rm Ca} \end{array}$ Dott (loc. cit), however, prefers the equation 2CO(CH<sub>3</sub>)<sub>2</sub>+8('aO('l<sub>2</sub>)

 $=2(H(1_3+(CH_3COO)_2Ca+2Ca(OH)_2)$ 

which agrees better with the results of experiment According to Ukita (J Chem Ind Tokyo, 1918, 21, 219) better yields are obtained by passing chlorine into ethyl alcohol until a liquid of D=1 299 is obtained. This is then added to a solution of bleaching powder and lime and distilled

Chloroform is also produced from acetic acid (Liebig), acetophenone (Orndoff and Jessel, loc cit), acetaldehyde turpentine, and ternenes (Chautaid, Jahresb 1851, 501) by the action of bleaching powder or chlorine. (V also Brit Pat 1,6094, 1917)

4 By reactions similar to the foregoing chloroform is also produced by the electrolysis of the chlorides of alkali or alkali earths in presence of alcohol, acctone, &c (Ber 1884, 17, Ref 624, see also Waser (Chem Zentr. 1910, 1, 1122), Feyer, Zeitsch Elektrochem 1919, 25, 115)

5. By the reduction of carbon tetrachloride (Geuther, Annalen, 107, 212, v also Chem Rev 1896, 88, U.S Pat 753325 (1904) and Eng Pat 13733 (1901))

6 By the action of chlorine in sunlight on methyl chloride (Regnault, Annalen, 33, 328)

7 From iodoform by the action of phosphorus pentachloride (Gautier, Bull Soc Chim. [2] 13, 127)

Manufacture - Chloroform was originally almost wholly manufactured from ethyl alcohol or methylated spirit Punfication of chloroform prepared from the latter presents rather more difficulty, but when properly carried out pure chloroform is obtained.

Of late years chloroform has been made for the most part from acetone, and although for some time a prejudice existed against the employment of chloroform made from this source for an esthetic purposes, it is now recogmsed that the product when properly prepared is in every respect equal to that obtained from alcohol.

Chloroform has been manufactured successfully in America for some years past from carbon tetrachloride by reduction with nascent hydrogen This is most efficiently brought about by mechanically agitating a mixture of the tetrachloride and finely divided non in water, a trace of acid being added to initiate the reaction (Smith, U.S. Pat 753325 (1904); see also U.S. Pat 1101025, 1914)


Attempts have also been made to manufacture chloroform by chlorination of the 'natural gas' (methane) produced in enormous quantities in the United States, but so far these attempts have not met with any measure of success

Electrolytic processes for the production of chloroform from alcohol and acetone have formed the subject of a number of patents, but the commercial results do not seem to have been so successful as the original experimental CH<sub>3</sub> CH<sub>2</sub> OH+O=CH<sub>3</sub> CHO+H<sub>2</sub>O work promised, and very little chloroform has In the case of acctone the reaction may be been made by this means (Schering, D. R. P.

29771 (1884), v also Zambelliti, L'Ettricita, 1899, 8, 664; Teeple, J Amer. Chem. Soc. 26, 536; and Waser (Chem. Zentr. 1910, 1, 1122)

Manufacture of Chloroform from Acctone and Bleaching-powder.—This is the process most generally employed. The method differs in minor details with the various manufacturers,

The reaction is carried out in a cast-iron tive. still of about 800 gallons capacity, which is provided with sturing gear, steam and cooling coils, and is connected with a condenser hundred gallons of water are run into the still, and 800 lbs of bleaching powder are added through a manhole, which is then securely but the following may be taken as representa-bolted down During addition of the bleaching



powder the mixture is very thoroughly stirred. (In some processes the mixing is carried out in a separate vessel, and the suspension is strained from the larger unbroken lumps of bleaching powder before being allowed to run into the still) The container (A in the diagram shown) is charged with 70 lbs. of acetone, which is then slowly run into the bottom of the still by means The introduction of the acetone of the valve B is accompanied by a rise of temperature which is not allowed to exceed 110°F., cooling being

effected if necessary by stopping the flow, of acetone and circulating cold water through the cooling coil in the still When all the acetone has been introduced the contents of the still are raised to 134°F. At this temperature At this temperature chloroform begins to distil over. The temperature is then very gradually raised to 150°F., so as to keep the chloroform steadily distilling. Towards the end of the reaction the mixture is stirred and the temperature raised until no more chloroform distils over.

The crude chloroform obtained is separated | and purified first by agitation with concentrated sulphuric acid. This operation is carried out in the vessel shown in the diagram; 1500 lbs of crude chloroform are introduced into the vessel and thoroughly stirred by means of the agitating gear shown, with 600 lbs. of sulphuric acid. The stirring is continued until a sample of the chloroform, when thoroughly shaken with pure concentrated sulphuric acid, does not impart the slightest colour to the latter. The time required for complete purification is usually about 3 hours. The chloroform is next separated from the sulphuric acid and finally distilled over lime.

The yield obtained from the above quantities averaged from over two thousand batches was 124 lbs., the highest yield in any one case being 131 lbs. Variation in yield is attributed to the varying composition of the bleaching powder, though doubtless other factors influence the result. Bleaching powder containing less than 33 p.c. of available chloring gives unsatisfactory results, while samples containing more than 35 p c. of chlorine are also often unsatisfactory The best results appear to be obtained with bleaching powder containing about 34 p.c of available chlorine.

Manufacture of Chloroform from Alcohol -This is carried out in the same plant as that used for the manufacture of chloroform from acetone A mixture of alcohol (150 lbs.) and water (3000 lbs.) is introduced into the still, bleaching powder (750 lbs) added, and the contents of the still gently heated by means of the steam Chloroform begins to distil over at a temperature of about 140°-145°F Further heating is then discontinued, the chloroform distilling off by reason of the heat generated during the reaction When the distillation begins to slacken the heating is resumed until at a temperature of 155°F the distillate consists of a mixture of alcohol and water. This mixture, which contains rather more than half the original alcohol, is used in making up the next batch, sufficient fresh alcohol being added to make up the required 150 lbs. The yield of crude chloroform from each working is about 75 lbs. Purification is effected as described above

Chloroform has also been manufactured by special processes for the purpose of producing directly chloroform sufficiently pure for anæs-Of these the most important thetic purposes

are the following .-

1 Chloroform from Chloral -It was found by Liebig, the discoverer of both compounds, that chloroform could be obtained from chloral by distillation with alkalis, and subsequent washing and drying, and this process has been utilised for the manufacture of chloroform. Upon the same reactions has been based a patent (Liebreich, Eng. Pat 15930, D R P 176063, 1904), whereby a dry mixture of chloral hydrate and alkalı can be prepared and compressed, from this product chloroform may be prepared by treatment with water. Chloroform obtained from chloral is of a high degree of purity. Like all other brands of chloroform it must, however, be preserved by the addition of a small amount of alcohol.

2. Chloroform Pictet.—This product is ob-

tained by fractionally crystallising commercial chloroform The crude chloroform is cooled to -80°, and the solid impurities are filtered off. The filtrate is then cooled to  $-82^{\circ}$ , and the liquid portion is drained off. The crystallised chloroform is then distilled at a very low temperature, and the middle fraction amounting to about 80 pc is collected as chemically pure chloroform (Pictet, Compt rend. 114, 1245, French Pat 215911, 1891, U.S. Pat. 489592, 1893). It was originally claimed that the pure chloroform obtained by this means was stable in the presence of light, and from this Schaet & Bilz (J. Soc. Chem Ind. 1893, 12, 543) deduced that it contained a small amount of alcohol which was subsequently shown to be the case.

3. Chloroform Anschutz ('Saheyhd chloroform').—Anschutz (Annalen, 273, [1] 94) observed that chloroform combined to form well crystalline compounds with the lactide condensation products obtained from o-phenol carbonic acids These compounds break up on gentle heating, giving pure chloroform The compound usually employed in the process is tetrasalicylide  $(C_6H_4 C_0)_4$  m p. 261°-262°, obtained by the action of phosphoryl chloride on This combines with 3 molecules salicylic acid of chloroform which are given up on warming The recovered salicylide can be employed again The product obtained by this process has been extensively used in Russia.

Properties — Chloroform is a heavy colourless liquid having a sweet taste and a pleasant ethereal odour It is not inflammable, but when diluted with alcohol may be made to burn and gives a smoky greenish flame. It is very slightly soluble in water (1 part in 200), but miscible with most organic solvents.

According to the most accurate determinations pure chloroform has sp gr 1 52637 at 0°/4° (Thorpe, Trans Chem Soc 37, 196), 1 52657 (Timmermans, Bull Soc belg chim 24, 244), or 1 4989 at 15°/4° (Baskerville and Hamor, J. Ind and Engin. Chem 1912, 278) According to the same authors the boiling-point is 61 2° (see also Wado and Finnemore, Trans Chem. Soc, 1904, 85, 946) Chloroform melts at -83° (Pictet) Its critical temperature is 262 9°, and critical pressure 53 8 atmos. (Kuenen & Robson)

Reactions and detection of Chloroform —The most important reaction of chloroform is the formation of an isocyanide from any primary amme when the latter is heated in presence of chloroform and alcoholic potassium hydroxide solution

### $CHCl_3+RNH_2+3KOH=R\cdot NC+3KCl+3H,O$

In practice aniline is usually employed as the reacting amine and the characteristic odour of phenylisocyanide is readily recognised. In this way one part of chloroform in 5000-6000 parts of alcohol can be detected. The reaction is also given, however, by bromoform, iodoform, chloral, trichloracetic acid, &c.; but not by ethylidene chloride. Chloroform readily reduces Fehling's solution and the reaction has been used as an indirect method of estimation.

A mixture of chloroform vapour and hydrogen when passed through a red hot tube yields hydrochloric acid This reaction also may be employed for the detection and estimation of chloroform in blood or other (non-chlorinated) liquids (Allen, Comm. Org Anlysis, vol 1 p 275)  $\alpha$  or  $\beta$  naphthol dissolved in concentrated potassium hydroxide solution when heated with a little chloroform give a blue colour which gradually becomes green and finally brown in contact with air

Anæsthetic Chloroform — The use of chloroform as an anæsthetic was first advocated by Simpson in 1848 (Edin Monthly J Med Sc 8, 415), and it is now used in very large quantities for this purpose alone Great care is necessary in its production for this purpose, since the presence of small amounts of impurities have a deleterious and often dangerous influence, and many cases of death under chloroform have been attributed to the employment of impure an esthetic the result of a considerable amount of investigation, it has now been shown fairly conclusively that pure chloroform under the influence of light, air, and moisture decomposes, more or loss rapidly according to the conditions pre-vailing, and that the products of decomposition are carbonyl chloride and hydrochloric acid, a reaction which may be represented thus

$$\begin{array}{c} \mathrm{CHCl_1\!+\!H_2O\!+\!O_2\!=\!COCl_2\!+\!HCl\!+\!H_2O_2} \\ \mathrm{CHCl_3\!+\!H_2O_2\!=\!COCl_2\!+\!HCl\!+\!H_2O} \end{array}$$

It was very early noticed that the tendency of chloroform to decompose was more marked with samples which showed a high specific gravity (1 497 to 1 50) than with those of lower density, and the greater stability of the latter was attributed to the presence in them of small amounts of alcohol It had earlier been suggested by Squibb (Amer Med Monthly, July, 1857), that the best means of preserving chloroform was the addition of a small amount of This practice has since then been alcohol recognised as sound in principle and the addition of alcohol in amounts varying from 0.6 to 1.0 p c by volume is allowed by most of the pharma-The British Pharmacopæia (1914) does not specify the amount of alcohol permissible, but defines the density at 1 483 to 1 487

Considerable diversity of opinion has been expressed with regard to the rôle played by alcohol in preventing the decomposition of chloroform, but it now appears to be fairly well established that it acts by reason of its property as a reducing agent, oxidation of alcohol taking place in pieference to that of chloroform the same time the products of oxidation appear to exert a retarding action which eventually a maximum When chloroform preserved with alcohol the decomposition of the chloroform is entirely prevented until all the alcohol is oxidised, when decomposition of the chloroform immediately commences properly stored and packed so that the action of light and air is reduced to a minimum, chloroform containing from 0.6 to 1 p.c by volume of alcohol will keep for years (Schact and Bilz, Pharm. J. 1893, 1005) The theory of the action of alcohol outlined above is due to Baskerville and Hamor (J. Ind and Chem. Eng. 1912, 368, &c), whose admirable series of papers form the most complete and recent account of the subject of this article Their theory is supported by the fact that a large

number of organic substances soluble in chloro form will inhibit the decomposition of the latter by light and air, and that all such compounds are reducing substances and act solely on account of their capacity for oxidation. That such compounds are not used in practice is merely on account of the special suitability of alcohol for the purpose

The various pharmacopeas also insist on certain tests for the absence of impurities, other than alcohol, which might possibly be present in chloroform through careless manufacture of storage. The details of the tests vary in different countries, but may be generally indicated as follows—

(1) The odour test.—This is usually carried out by evaporating a quantity of chloroform (10–50 c c.) to small bulk, and then allowing the residue to evaporate on a piece of filter paper At no stage of the evaporation should there be any appreciable foreign odour, eg of fusel oil, higher alcohols, or chlorimated derivatives of alcohol or acetone

(2) Residue—There should be no weighable residue after evaporation of 50 to 100 e.e. of chloroform

(3) Sulphuric acid test.—This consists in shaking chloroform (usually 20 c c) with pure concentrated sulphuric acid (usually 15 c c) for a short time and then allowing the mixture to stand. The two layers are required to remain colourless or practically so over a period of time which varies with the different pharmacopenas. The object of the test is to detect the presence of organic impurities which should have been removed in the process of manufacture.

A more delicate variation of this test has been suggested by Langgaard (Therap Monatsh May, 1902) Concentrated sulphuric acid and chloroform are mixed in the proportions required by the German Pharmacopæia and then four or five drops of formalin (40 p c solution) added and the mixture well shaken No coloration should develop on standing The German Pharmacopæia requires the solution to remain colourless for forty-eight hours According to Budde (Veröffentlichungen aus dem Gebiete des Militar-Sanitatswesens, 1913, 6, 113), however, the sensitiveness of the reaction increases with the amount of alcohol present

(4) Acidity —Water shaken with chloroform and separated should remain neutral to litmus

(5) Chlorine—The aqueous layer after shaking chloroform with water should give no reaction when shaken with a 10 pc solution of cadmium rodide and a few drops of starch solution

nodide and a few drops of starch solution
(b) Hydrochloric and—The aqueous layer obtained as above should not give more than a very slight opalescence with a solution of silver nitrate (Cf Analyst, 1919, 407)

(7) Decomposition products—Two c c of the sulphuric acid solution obtained as in test (3) when diluted with 5 c c of water should remain nearly colourless and have no unpleasant odour When further diluted with 10 c c of water the solution should remain colourless and on addition of a little silver nitrate should not give more than a slightly diminished transparency

For the last three tests Budde (loc. cit) proposes to substitute one, namely, a reaction with benzidine, which it is claimed is simpler and at the same time more sensitive. A little

pure benzidme is dissolved in about 10 c c of chloroform. The solution should remain per feetly bright when kept in the dark, out of contact with air for 24 hours. In the presence of chlorinated products or of carbonyl chloride an opalescence or floculent precipitate is produced. If chloral is present, a faint red coloration develops at first and changes to bluish-red after a time

## Other possible Constituents of Anæsthetic Chloroform

Ethyl chloride - Chloroform manufactured from alcohol was at one time considered, for anæsthetic purposes, superior to that obtained from acetone Wade and Finnemore (1904, 85, 938) showed that chloroform prepared from alcohol contained a small quantity (approx 0.05 p.c.) of ethyl chloride, and to this they attributed its superior qualities. According to them pure chloroform prepared from acetone, and to which a small proportion of ethyl chloride was added, was alone clinically equal to that obtained from alcohol Baskerville and Hamor (lor cit p 574), however, state that American anæsthetists have found chloroform from acetone when properly purified to be fully as reliable as They therefore look chloroform from alcohol upon ethyl chloride as an impurity of anæsthetic chloroform which although perhaps not objectionable, has no special value Since the boiling-point of chloroform is lowered to the extent of about 2° by the presence of 0.1 pc of ethyl chloride (Wade & Finnemore, loc cit) its presence is easily detected by fractionation

Acetaldehyde—For the detection of this impurity in chloroform Baskerville and Hamor suggest the following—10 c c of chloroform are shaken with 5 drops of Nessler's solution U.S.P. and allowed to stand for five minutes Anæsthetic chloroform should show no precipitate or colouration under these conditions

For the detection of acetaldehyde chloroform is in practice often tested by allowing to stand over solid KOH. For this purpose the KOH should contain at least 80 p.c. KOH and be free from water, otherwise oxidation of alcohol may take place and a positive reaction be obtained

Acetone —An esthetic chloroform should give a negative result when tested as follows. To 10 c.c. of chloroform are added 5 drops of 0.5 p.c. solution of sodium introprusside and 2 c.c. of ammonia (D=0.925) and the mixture allowed to stand for five minutes in the cold If acetone is present to the extent of 0.2 p.c. an amethyst colour is produced. By saturating the mixture with ammonium sulphate the colour is produced in the presence of 0.1 p.c. of acetone

Carbonyl chloride, which is only likely to be present in badly preserved or stored an esthetic chloroform, may often be detected by its nauseous colour. The following test is also quite satisfactory. To 15 c c of chloroform in a dry glass-stoppered cylinder of 25 c c capacity is added sufficient clear solution of barium hydroxide (0 5 p c.) to completely fill the tube, which is then kept in the dark for 3 hours. In the presence of carbonyl chloride a film of barium carbonate is produced at the junction of the liquids.

Carbonyl chloride is also readily detected by the benzidine test (Budde, *loc cit*), which gives a positive reaction with one part in twenty thousand (1 20,000). The impurities of commercial chloroform may include chloral alcoholate and similar chloral compounds. These may be estimated by treating the chloroform with N/10 alkali and titrating the excess. Pure chloroform is not affected by alkali of this strength. Aldehydic and similar compounds are detected by means of Nessler's reagent, which gives with impure chloroform a red-brown deposit readily changing to greenish black. Pure chloroform gives no such reaction (Enz. Apoth. Zeit. 1913, 28, 672. Analyst, 1913, 497)

Chloroform is comparatively unstable under the influence of ultra-violet radiation (Kailan,

Monatsh 1917, 38, 537)

Detection and Estimation of Alcohol in Chloroform -For the detection of alcohol in chloroform the iodoform reaction provides a very delicate test which, however, is not specific It is best carried out by shaking the chloroform with sulphuric acid, separating and diluting the sulphuric acid layer. This is then distilled and the reaction carried out on the first portions of the distillate To 10 c c of the distillate 6 drops of a 10 p c solution of potassium hydroxide is added, the mixture warmed to about 50°, and then drop by drop a solution of iodine in potassium iodide until a brown colour is produced The solution is then carefully decolourised by addition of potassium hydroxide On standing crystals of iodoform separate

For the estimation of alcohol the density of the chloroform forms an approximate guide provided the chloroform is otherwise pure, but for more exact determinations the method of Nicloux (Bull Soc chim [3] 35, 330, 1906) as modified by Béhal and François (J. Pharm. Chim. [6] 5, 417) is recommended. The alcohol is removed from the chloroform by agitation with sulphuric acid, the latter is diluted and gently The distillate is boiled with sulphurie acid and potassium dichromate solution containing 16 97 grams per litre is added until a green (not greenish yellow) tint is obtained Each 2 cc of solution equals 0.01 gram of absolute alcohol If the solution contains more than 2 p c of alcohol it must be diluted and the estimation repeated A similar modification has been suggested by Budde (loc cit)

1-CHLORO - 8 - NAPHTHOL - 3 6 - DISUL - PHONIC ACID

is prepared by dissolving 1-amino-8-naphthol-3 6-disulphonic acid in a cooled solution of sodium carbonate, carefully acidifying with hydrochloric acid and diazotising with the theoretical amount of sodium nitrite. The separated diazo-compound is warmed with hydrochloric acid and cuprous chloride, the copper removed by sulphuretted hydrogen, and the filtered solution saturated with common salt when the acid sodium salt of the chloroacid is precipitated (Cassella & Co, D R P. 79055). The chloro-acid is also formed by treating 1 8-naphthylene-diamine-3: 6-disulphonic acid with sodium nitrite in acid solution and acting on the azimino-compound with copper paste (Badische Anilin und Soda Fabrik, D R. P. 147852).

CHLOROPHENINE ORANGE v Azo-

CHLOROPHENINE YELLOW v. PRIMULINE AND ITS DERIVATIVES.

CHLORORUBINE v MADDER.

CHLOROXYLONINE C<sub>22</sub>H<sub>23</sub>O<sub>7</sub>N, an alkaloud ccurring in East Indian satinwood (Chloroxylon Inveteria (DC.)). It crystallises from alcohol in olourless prisms, m.p. 182°-183°, [a]<sup>18</sup>-9° 18′, a chloroform, is soluble in excess of ammonia olution, and is neutral to htmus. The salts are rystalline and dissociate in water; aurichloride 3'HCl'AuCl<sub>3</sub> groups of reddish-yellow needles, a.p. 70°. Chloroxylonine contains four 'OMe out no hydroxyl groups (Auld, Chem. Soc. 'rans. 1909, 95, 964). It produces dermantis when applied to the skin and is the cause of the dermatitis sometimes experienced by atinwood workers (Cash, Brit. Med. J. 1911, 9ct. 7; and Bull. Imp. Inst. 1911, 9, 351).

CHLORPICRIN (truchtoron tromethane) CCl<sub>3</sub> VO<sub>2</sub> is prepared by distilling nitro-compounds in particular, pierie acid) with bleaching-powder olutions (Stenhouse, Ann. 1848, 66, 241), by one heating of chloroform with fuming nitric cid (Mills, Ann. 1871, 160, 117), or by the ction of chlorine on nitranilic acid (Levy,

edlicka, Ann 1888, 249, 86)

In Hofmann's method of preparation 45 parts f fresh bleaching powder are made into a paste 7th water and mixed with a saturated solution 4 5 parts of pieric acid in water at 30° As soon s the very violent reaction is finished, the chloronerm is distilled over (Ann. 1866, 139, 111)

Chlorpierin is a mobile liquid which boils at 11 9°C and possesses a specific gravity 1 69225 0/4°) Thorpo. On heating it slightly decomposes nto carbonyl chloride and introsyl chloride Cl<sub>3</sub> NO<sub>2</sub>=COCl<sub>2</sub>+NOCl) (Gardner and Fox, them. Soc. Trans 1919, 1188). On strong coolng it solidifies and then melts at -69 2°C apour acts as a strong irritant to the eyes and aucous membrane For thermal expansion, see 'horpe, Chem. Soc. Trans 1880, 37, 198 eduction with iron and acetic acid it yields vethylamine A similar bromo-nitro-methane is nown. It melts at 10°C, boils at 127° (118 am), and is converted by the action of potassium yanide in cold alcoholic solution into the ipotassium salt of 1122 tetranitroethane Scholl, Ber. 1898, 31, 642, *ibid*. 1902, 35, 4292)

CHLORYL. Syn. for ethyl chloride used as n anæsthetic

CHOCOLATE is a mixture of cacao paste 7th sugar and flavouring matter The paste s prepared from fermented cacao beans (the nfermented kinds being unsuitable for chocoite making) in the manner described under The sugar, which constitutes 50 p c or nore of the mixture, consists usually of white rystals (although the lower-grade coloured ugars may be used for the cheaper chocolates), round to the finest powder and completely ried before admixture with the cacao paste. most common flavouring substance is anilla, but other spices, as cardamoms, cinnaion, cloves, mace, and nutmeg, or the essential ils extracted from them, also artificial flavourags such as vanillin, are frequently employed 'eru balsam and gum benzoin are sometimes sed for a similar purpose. The flavourings are

introduced either in alcoholic solution or incorporated with sugar moistened with a little oil

It is essential that during admixture the paste should be kept in a liquid condition, and to this end the mixing machines are heated to a temperature slightly above the melting-point of cacao butter, say from 35° to 40°, and the added sugar may be pieviously warmed with advantage to obviate a fall in temperature of the liquid cacao paste.

The various characters of French, Spanish, Swiss, and other chocolates depend upon the nature and proportions of the different ingredients used. A number of recipes for the preparation of some of the principal types of

chocolate are given in Zipperer, 156

In addition to sugar, various farmaceous substances are frequently employed in the cheaper forms of chocolate, as wheat and potato flour, rice, arrowroot and dextrin, and oceasionally bailey, oats, 1ye, and chestnut flour. In special cases where sugar is inadmissible, as in chocolate intended for use by diabetic sub jects, the sugar is replaced by saccharin  $(q \ v)$ It is then usual, in order to give the necessary 'body' to the chocolate, to add some form of meal or flour (preferably leguminous), and pea, bean, or linseed meal are prefeired for this Nut chocolate is made by adding nuts, purpose either ground or whole, to the chocolate paste; and milk chocolate by incorporating with it 'milk powder,' i e milk deprived of its moisture and fat The usual composition of milk chocolate, by the best makers, is approximately 10 p c cacao, 25 p c cacao butter, 40 p c sugar, and 25 p c. dued milk The above-mentioned forms of chocolate, after being ground into a more or less homogeneous mass, are placed in moulds to form bars, blocks, cakes, and numerous fancy shapes

An immense variety of fancy preparations of chocolate consist, however, merely of a chocolate envelope containing other substances, as biscuits, creams, dried fruits, nuts, fruit jelly, liqueurs, &c, and in some cases, medicinal preparations, as chocolate-coated pills. The most important of these fancy preparations are the chocolate creams

The so-called 'cream' is made either from powdered loaf sugar heated until it melts, or a mixture of this with glucose, with or without various colouring and flavouring materials. While in a liquid state, it is poured into moulds of the desired shape, made in starch flour on a table, which can be kept cool and which can be agitated by mechanical means On cooling, the creams set into a stiff granular mass, when they are either dipped into a liquid chocolate paste, or the latter is poured over them by hand and spread uniformly by means of the rocking table. The process of covering, as well as most of the other processes in the manufacture of chocolate, are now generally effected by means of ingenious and complicated machinery

An extra amount (about 15 p c ) of cacao butter (or substitute) is added to the chocolate intended for covering purposes, to keep the latter in a sufficiently plastic condition during the process of cooling and to give smoothness to the finished article. The latter, whether in the form of plain chocolate or chocolate confections, should be quickly cooled, and attention to temperature is important throughout all the various operations.

The adulterants commonly used in chocolate are similar to those mentioned under Cocoa,

the most important being the replacement of the cacao butter by other fats and oils (as coco-nut fat, illipé butter, &c), the addition of various farinaceous substances, including dextrin, and the admixture of finely ground cocoa shells (Zipperer, Die Shokoladen Fabrikationen, Vilhers et Collin, Traité des Substances Alimentaires)

CHOCOLATE FATS. Under this name, the writer comprises those vegetable fats which are used in the manufacture of chocolate as cacao butter substitutes. The natural chocolate fat is cacao butter, and best chocolates contain cacao butter only 'Cheap' chocolates are made,' to a larger or smaller extent, if not entirely, with substitutes. At first, animal fats were used, but as they impair an unpleasant flavour ('animal flavour') to the chocolate, at present fats of vegetable origin only are employed, such as coco-nut stearine and palm-nut stearine. During recent years, attempts have been made to produce chocolate fats from Boineo tallow, dika fat, and similar vegetable fats. J. L.

CHOLEIC ACID v. Cholic Acid CHOLESTEROL C<sub>27</sub>H<sub>49</sub>O Discovered by Conradi in gall-stones in 1775 Occurs in all vertebrates and throughout the whole of the animal kingdom with the exception of certain groups of the invertebrata in which it is replaced by closely related substances (Dorée, Biochem J 1909, 4, 72) It is found in bile, blood, the brain, and generally in all animal tissues, being closely associated with the lipins and fat present in the various organs Biliary calculi are composed almost entirely of cholesterol Its most abundant source is, however, wool grease

It may be obtained from animal tissues by the following method the tissue is minced, dried in an oven at 100° and the dried material ground to a powder in a coffee mill and extracted in a Soxhlet apparatus with ether for 9 hours After distilling off the other, the residue is saponified with alcoholic potash. The alcohol is The alcohol is distilled off and the cholesterol separated from the soaps and glycerol by extraction with ether The tendency of the ether and soap solutions to form emulsions makes the extraction difficult and various methods have been introduced to Thus Ritter recomovercome this difficulty mends that the soap residue be mixed with sodium chloride and water, evaporated to dryness and the resultant mass, after drying in an oven at 80°, extracted in a Soxhlet apparatus for 9 hours Kossel and Obermuller (Zeit physiol Chem 1890, 14, 599) recommend saponification of the ethereal solution of fat with an alcoholic solution of sodium ethylate at ordinary temperature. After a short time a compact mass of soaps separates which is easily filterable, the cholesterol being present in the alcohol-ether filtrate By this method wool grease, which is only saponified with difficulty, is completely decomposed in 24 hours at ordinary temperature After filtering off the soap and washing the precipitate with ether, the filtrate is well washed with water, the ethereal solution dried with CaCl2, and the ether distilled off, leaving a residue of cholesterol. It may also be separated from the fatty acids in wool grease by distillation with superheated steam or by direct heating (Mayer, Eng. Pat. 20639; J. Soc. Chem. Ind. 1894, 162).

Cholesterol crystallises from ether or chloroform in silky needles, m.p. 148 4 to 150.8 (corr.). From aqueous alcohol (90 pc.) it crystallises in characteristic plates with one molecule of water of crystallisation. Insoluble in water, sparingly soluble in cold, easily in boiling alcohol (9 parts alcohol, sp.gr. 0.87), and in chloroform, ether, carbon disulphide, less soluble in acetone, petroleum, benzene, and glacial acetic. Soluble in fats, fatty acids and in solutions of bile salts. Sp.gr. 1.046; it is optically active; in ether solution (a) = -31 12, in CHCl<sub>3</sub> (a) = -36 61. It sublimes unchanged at 200°, but is decomposed at higher temperatures, hydrogen being evolved at 300° to 320°. In a vacuum it can be distilled unchanged

Constitution—Cholesterol is an unsaturated secondary alcohol, possessing a complicated terpene-like structure which has not yet been completely elucidated. It has, however, been possible to establish in the molecule the presence of (1) a secondary alcohol group in a reduced ring, (2) a terminal vinyl group, and (3) an isoamyl group. Our present knowledge is best represented by the formula

$$\begin{array}{c} \operatorname{CH}_3 \\ \operatorname{CH}_3 \\ \operatorname{CH} \end{array} \\ \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_{17} \\ \\ \operatorname{CH} \\ \operatorname{CH}_3 \\ \operatorname{CH} \\ \operatorname{CH}_4 \\ \operatorname{CH}_4 \\ \operatorname{CHM}_6 - \operatorname{CH} = \operatorname{CH}_5 \\ \end{array}$$

By applying various reagents under different conditions, a large number of oxidation products have been obtained (Mauthner and Suida, M 24, 489; Windaus, 25 paper in Berichte, 1903 to 1917, Diels and Abderhalden, Ber 1903, 3177, 1904, 3092, 1906, 885, 1371), the formula arrived at for cholesterol being based Oxidation of the secondary on these results alcohol group gives the corresponding ketone, cholestenone, mp 78°, the latter compound has also been obtained by direct oxidation of cholesterol with copper oxide at 300°, more conveniently by oxidising cholesteryl-di-bromide with chromic acid in acetic solution at 70°, and subsequently removing bromine by zine and acetic acid (Windaus, Ber 1906, 518) Cold chromic acid or potassium hypobromite solution oxidises the secondary alcohol to the ketone group and at the same time the vinyl group is oxidised and a new ring formed, the diketone, oxycholestenone is the resulting product

By the addition of two hydrogen atoms to oxycholestenone, cholestanedione, m p 170°-171°, is produced Fuming nitric acid reacts with cholesterol forming a nitro-compound which is reduced to cholestanonol, in.p 140° (Windaus, Ber 1903, 3752, 1917, 133), the latter being oxidised by chromic acid to cholestanedione The relation of these substances to each other is shown by the following formulæ:

Further oxidation of these products has resulted in the isolation of a number of carboxylic acids, respectively mono, di, tri, and tetrabasic, some containing the same as, and some a smaller number of carbon atoms than cholesterol in their molecules

The following compounds have been recognised among the products of the oxidation of cholesterol by nitric acid dinitroisopiopane and acetic, succinic, methylsuccinic and  $\alpha$ methylglutaric acids, acetone, hydroxyisobutyme acid, methylisohexyl ketone, and octane all of which substances may be regarded as derived from an isooctyl side chain in cholesterol On this assumption cholesterol may be represented by the formula

The action of ozone has been investigated by Dorée & Gardner, Molman and by Harries If the above constitution be accepted, only 4 atoms of oxygen should be taken up by the cholesterol molecule, an ozonide containing 7 atoms of oxygen has, however, also been isolated. The balance of chemical evidence being against the presence of two double bonds in the molecule, Dorée and Gaidner regard the addition of the three extra atoms of oxygen, which are only slowly taken up, as probably due to the opening of a bridged ring in the molecule

From the first of the above formulæ, only one dihydro derivative of cholesterol should be expected, several have been described, none of which appears to be identical with coprosterin, the dihydro cholesterol occurring in the intestines The 8 cholestanol, mp 142°, obtained by reducing cholesterol with platinum and hydrogen represents probably the true dihydro derivative (Willstatter & Mayer, Bei 1908, 2199)

Cholesteryl Esters — The cholesteryl esters of the lower fatty acids are prepared by heating cholesterol with the anhydrides of the acids, those of the higher fatty acids by boiling chole sterol with the acid and removing the water formed (Jaeger, Rec. trav. chim. 1907, 311) M p acetate, 114°, propionate, 98° These substances on melting form anisotropic liquids, many of the esters having 3 stable liquid phases. The benzoate melts at 146° to a turbed liquid which examined with the polariscope shows definite crystal structure and at 178° passes to a clear liquid. The laurate is an instance of a compound which can be heated several degrees above its melting-point without melting.

Detection and Estimation —The thin rhombic

plates of cholesterol obtained from 90 p.c. alcohol are very characteristic. • Under the

microscope they often show notched angles and exhibit a curious overlapping arrangement A number of colour reactions have been described of which the following are the most useful. Salkowski's test A small amount of cholesterol is dissolved in 2 c c chloroform and shaken with an equivalent amount of cone H<sub>2</sub>SO<sub>4</sub>, the chloroform first becomes bloodred, then cherry-red, and finally purple, which lasts for several days, poured into a basin, it becomes successively blue, green, and yellow owing to absorption of water The H<sub>2</sub>SO<sub>4</sub> layer is strongly fluorescent

Burchardt-Liebermann Test — Dissolve a little cholesterol in 2 e e chloroform, add 20 drops of acetic anhydride and 1 drop cone  $H_2SO_4^-$  a violet-pink colour appears. This test is also violet-pink colour appears

given by resinic acids Estimation -In animal tissues, the choles-

terol may be estimated in the residue from the ether extract by converting to the acetate by boiling with acetic anhydride. The acetate is washed with water till free from acetic acid, and a weighed amount saponified with a known volume of alcoholic potash. The residual potash is titrated with standard acid and the The residual amount of potash equivalent to the cholesterol thus determined. The iodine value will also give information The most satisfactory method is that in which the cholesterol is precipitated and weighed as its insoluble compound with digi-Ransom, in 1901, found that the hamolytic power of saponin is prevented by the addition of cholesterol Windaws investigated the reaction (Bei 1909, 238) and found that digitonin formed an insoluble compound with cholesterol m alcoholic solution A 1 pc solution of digitonin in 96 p.c. alcohol is added to the ether-alcohol solution of fat, and after standing for some hours the precipitate is collected on a Gooch filter, washed with alcohol and ether, dued at 100° and weighed The precipitate contains 25 pc cholesterol The filtrate is concentrated and after the addition of water extracted with ether, the excess of digitomin remains in the weak alcoholic liquid, and the cholesteryl esters in the ethereal layer. solvent is distilled off and the residue saponified with hot alcoholic potash The cholesterol is then extracted with ether or petroleum and precipitated with digitonin as above described In this way the free and combined cholesterol are separately estimated Various modifications of this method have been introduced, thus Prescher recommends that after saponification, the soap should be decomposed with hydrochloric acid, CHCl<sub>3</sub> added, and the CHCl<sub>4</sub> solution of the fatty acids separated, filtered, and treated with the I pc digitonin solution at 50°, the liquid being stirred during the precipitation (Zeitsch Nahr. Genussm. 1917, 481). In fats where adulteration with vegetable fats may have taken place, it is important to discriminate between phytosterol and cholesterol, for this, the acetates or di-bromides must be prepared, and their melting-points examined Marcusson and Schilling recommend the following method (Chem Zeit 1913, 1001), 50 grams fat are shaken for 15 minutes in a separating funnel with 20 c c 1 p c. alcoholic solution of digitonin. The bottom layer of oil is run off; the alcoholic upper layer containing the flocculent precipitate

of digitonide is shaken with 50 to 100 c.c. ether. The ppt. is washed with ether, dried, powdered and again washed. It is then heated with 1.5 c c. acetic anhydride in a test tube for half an hour. On cooling cholesterol and phytosterol acctates crystallise out. The acctates are re-crystallised and their mps. determined. Cholesteryl acctate melts at 114°, phytosterol acctate at 128°. They may also be separated by crystallising their di-bromides from ether and glacial acetic (Windaus, Chem Zeit 1906, 1011) A mixture of 4 grams cholesterol and 4 grs. phytosterol was dissolved in 80 cc ether and 80 c c 5 p c bromine in glacial acetic solution added After standing for 1 hour at 0°, the ppt is washed with 40c c glacial acetic acid, and with 40 c c 50 p c acetic acid, the washings being added to the filtrate, and a further ppt added to the original ppt, which is then washed with water and dried. The ppt contains the cholesterol bromide, the filtrate the phytosterol compound The ppt is heated with 100 c c glacial acetic acid and 5 grs. zinc dust for 2 hours, the excess of zinc filtered off, and a large quantity of water added A precipitate forms which is boiled for 2 hours with 100 c c. alcoholic potash, and the cholesterol then precipitated by the cautious addition of water. The phytosterol is obtained by similarly treating the solution of the phytosterol brounde

Cholesterol is found in commerce under the name of lanolin and is used mainly for medicinal purposes, and, owing to the ease with which it is absorbed by the skin, as a substitute for vaseline or lard in the composition of plasters, ointments and salves It has also been employed in the manufacture of cosmetics and pomades, and for softening leather Lanolin is obtained from wool oil, which contains about 70 pc of cholesterol and 30 pc fatty acids, by saponification by means of caustic soda, whereby an emulsion is obtained which on dilution with water gives so-called wool-milk On treating in a centrifugal machine impure lanolin separates out as a cream and may be precipitated by lime. The purified product is afterwards kneaded with 30 p.c. of water, in which state it is found in commerce and forms the 'Hydrous Wool Fat' of the B.P. Anhydrous lanolin absorbs about 100 p c of water, does not become rancid, and is antiseptic

Isocholesterol is an alcohol isomeric with cholesterol which occurs only in woolgrease, m p 137-138°,  $\alpha_D = +60$  The mixture of cholesterol and isocholesterol is heated for 30 hours in a sealed tube with four times its weight of benzoic acid. The benzoates are insoluble in boiling rectified spirit and are separated by crystallising from ether Cholesteryl benzoate crystallises in shining plates, its isomer as a light powder, mp 190°-191° Isocholesterol may also be separated from cholesterol by means of oxalic acid, cholesterol forms a crystalline compound with oxalic acid, and the isocholesterol can be estimated in the filtrate as benzoate (Madinaveitia & Gonzalea, Anal Fis Quim. 1916, 398). Isocholesterol gives no colour reaction with chloroform and sulphuric acid, but like cholesterol gives a pale yellow substance when heated with a drop of concentrated HNO<sub>3</sub> and a deep yellowish red colour on the addition I. S M. of ammonia (Schiff's reaction).

CHOLESTROPHAN v. PARABANIC ACID CHOLIC ACID C24H40O5. Occurs in bile, chiefly in combination with taurin and glycocoll as taurocholic and glycocholic acids. Free cholic acid, probably derived from these complex

acids, occurs in small quantity in the intestinal

contents, in the fæces and in the bile.

Preparation -Cholic acid and the closely related isomeric choleic and desoxycholeic acids (C24H40O4) are prepared from bile after first treating it with alkali to decompose the compounds with glycocoll and taurin Five kg ox bile are boiled for 30 hours with 1 kg 30 pc. NaOH solution under a reflux condenser, 5 litres of water added and the mixture precipitated with hydrochloric acid at 50°. The precipitate is well rubbed up with water to remove hydrochloric acid and dried on a waterbath (Bondi and Muller, Zeitsch physiol. Chem. 47, 501) The residue consists of a mixture of choice acid with choice and desoxycholeic acids, which may be separated in various ways, depending on (a) the sparing solubility of the barrum salt of choleic acid, this being pre-cipitated when barrum chloride solution is added to a solution of the ammonium salts of the mixed acids: and (b) the relative insolubility of cholic acid in alcohol Langheld (Ber. 1908, 378) recommends the following method The crude dried acids are shaken with twice their weight of alcohol and the remaining cholic acid twice recrystallised from absolute alcohol Excess of soda is added to the mother liquor, and the reaction mass warmed on the waterbath; a nearly quantitative separation of sodium cholate takes place This is filtered whilst hot, washed with boiling alcohol and recrystallised The mother liquors are concentrated in racuô and cooled the sodium salts of the fatty acids present now separate The remaining acids are regenerated from the salts and esternfied with 02 pc. alcoholic H(I. The esters are boiled with much ligroin which dissolves 4 pc On saponifying the ligioin-insoluble esters with alcoholic potash, a further separation of cholic acid as sodium salt takes place. The residue is extracted with boiling benzene and ether, and then recrystallised from ethyl acetate. It consists of the two isomeric desoxycholeic acids the existence of which has been described by Mylius and by Vahlen By fractionation from ethyl acetato these were obtained as substances melting respectively at 172° and 187°. The latter appears to be identical with the choleic acid obtained by Mylius (Ber 1886, 369 and 2000, 1887, 683 and 1968)

Properties - Cholic acid is a monobasic acid containing two primary and one secondary alcohol groups. It crystallises from aqueous ether or from dilute acetic acid with 1 molecule of water, from alcohol with one molecule of alcohol: it loses both water and alcohol at 120°. It is soluble in acetone and acetic acid, almost insoluble in cold water, more soluble in hot Its solubility in cold 70 pc alcohol (1 m 75)is 1 in 20 Specific rotation:  $a_D=37\,0$  (Vahlen, Zeitsch physiol. Chem 21, 265). It behaves to bromine and potassium permanganate as a saturated substance, but forms an ozonide on treatment with ozone, suggesting the presence of an unsaturated cyclic substance (Langheld, Ber. 1908, 1023; von Furth and Ishihara,

Brochem. Zeitsch. 43, 323). A large number of oxidation products have been obtained (Pregl. Zeitsch physiol Chem 1910, 167). On distilling under 12 mm. pressure, a heavy oil passes oil passes at 200° to 300°, solidifying to the triply unsaturated cholatrienecarboxylic acid  $\rm C_{24}H_{34}O_{2}$ , mp 163°. This acid gives the fluorescence with  $\rm H_2SO_4$  characteristic of cholic acid, but this property is lost on reducing it with hydrogen and palladium black (Wieland and Woil, Zeitsch physiol Chem 80, 287)

Colour Reactions—(1) Choice acid dissolves in conc. H<sub>2</sub>SO<sub>4</sub> with a reddish yellow colour showing a characteristic green fluorescence. (2) Pettenkofer's Test. An aqueous solution gives a red colour with a small amount of cane sugar on addition of conc. H<sub>2</sub>SO<sub>4</sub> or H<sub>3</sub>PO<sub>4</sub>. (3) Myllius' iodine leaction (Zeitsch physiol Chem 1887, 306). To 0.04 gr. m. 1 c. c. alcohol, add in the cold. 2 c. c. N/10 iodine solution and slowly dilute with water; the mixture suddenly solidifies to a paste of crystals which appear intensely blue by transmitted light. This test differentiates choice from choleic and desoxycholeic acids.

Compounds u th Glycocoll and Taurin.—Both choice and choice acids occur in combination with glycocoll and taurin forming the important bile-acids. In man the ratio glycocholate taurocholate varies from 5-1 to 10-1. Taurin is present in the bile of most mammals, reptiles, birds and fishes. Glycocholate is absent in the bile of the polar bear, dog, goose, and in certain fishes. Choice acid is formed in the liver and there combined with taurin and glycocoll It is probable that it is derived from the choicesterol of the red blood corpuscles.

The physical properties of these acids play an exceedingly important part in the digestive processes. They act as solvents for lipins and fat and so increase the velocity of resorption of the fatty acids in the intestine, and act as

activators to certain enzymes

Glycocholic Acut C<sub>16</sub>H<sub>43</sub>O<sub>6</sub>N occurs in quantity in bile, especially in that of herbivora It is extracted from fresh bile by saturating with other and decomposing with from 3 to 4 p c cone, HCl; on standing a crystalline paste of glycocholic acid separates. The bile may also be dried on the water-bath and extracted with 90 p c alcohol. After distilling off the alcohol the residue is taken up with water, neutralised with milk of lime and filtered. On acutifying with sulphuric acid the acid separates. It is easily soluble in acetone and acetic acid, soluble with difficulty in ether and insoluble in chloroform and benzene. According to Knoop, it crystallises with 1½ mol. water of crystallisation and melts at 138°

Taurocholic Acid C<sub>26</sub>H<sub>46</sub>NSO<sub>7</sub>. This is the chief constituent of dog's bile. Alcohol is added to dog's bile and the filtered liquid concentrated to dryness on the water-bath. The residue is dissolved in alcohol, precipitated with ether and dissolved in water. Ferric chloride solution is added to the neutral solution until further addition produces no precipitate. After removing the iron, the filtrate is evaporated to dryness. The sodium salt is rubbed up with alcohol containing 2–3 p.c. hydrochloric acid; after filtering off sodium chloride ether is gradually added until crystallisation sets in. •

Properties — Soluble in water, alcohol, ethyl acetate, insoluble in ether and chloroform. The crystals are obtained anhydrous on drying invacuô. It melts with decomposition, the mp gradually changing from 100° to 180°. It precipitates protein from solution.

1. S. M

CHOLÎNE AND RELATED SUBSTANCES. Choline, timethyl-\beta-hydroxyethyl-ammonium hydroxide, CH2(OH) CH2 N(CH3)3OH, is formed by the hydrolysis of lecithin and hence probably occurs in all living cells, its isolation from animal or vegetable tissues has been recorded very many times, but has no special significance, for even when not originally present in the free state, some is generally formed from phosphatides during the process of isolation Kauffmann (Zeitsch physiol Chem. 1911, 74, 175) has shown, however, that if perfectly fresh ox brain is worked up rapidly, with precautions, hydrolysis may be avoided and then no choline is obtained The best natural source is crude lecithin Hard egg-yolk is extracted with ether and then with alcohol, the extracts are mixed and evaporated and the residue is boiled for one hour with 50 parts of cold saturated baryta solution After removal of the baryta and evaporation, the residue is extracted with absolute alcohol and to the solution alcoholic mercuric chloride or platinic chloride is added (Compare D R P 193449, Moruzzi, Zeitsch Physiol Chem 1908, 55, 352; MacLean, ibid. 55, 360) The mercurichloride is decomposed by hydrogen sulphide, the platinichloride by calculated quantity  $\mathbf{of}$ potassium chloride, in either case the choline chloride is extracted by absolute alcohol from the residue after evaporation Most other natural sources yield very little (seeds and animal viscera something like 0.02 p.c.). The isolation of choline (and betaine) from vegetable sources has been described by Schulze (Zeitsch. physiol Chem. 1909, The aqueous plant extract is purified with lead acetate, and then, after removal of the lead and evaporation, the residue is extracted with absolute alcohol. Choline and betaine are precipitated with alcoholic mercuric chloride After reconversion into the chlorides and evaporation, cold absolute alcohol dissolves choline chloride, and leaves betaine chloride behind

Cholme is best prepared synthetically. Gaseous trimethylamine is made to act (a) on ethylene oxide (Wurtz, Compt rend 1867, 65, 1015, 'névrine'=cholme), (b) on ethylene dibromide; the resulting trimethylamino-bromethylium bromide is best converted into choline by heating with water (Kruger and Bergell, Ber 1903, 36, 2901), (c) on ethylene chlorohydrin (Renshaw, J Amer Chem Societ (a) 1910, 32, 128) Cholme is also obtained (d) by methylating amino-ethyl alcohol (see below and Trier, Zeitsch physiol. Chem 1912, 80, 409). Of these methods (c) is the best, rather more than one equivalent of trimethylamine gas is passed mto ethylene chlorohydrin in a tube cooled to —12° to —20°, which is subsequently heated to 80°–90°. The yield is almost quantitative.

The estimation of choline (and betaine) has been worked out by Stanck (Zeitsch. physiol Chem 1905, 46, 280; 1906, 47, 83, 48, 334), and depends on the precipitation in sodium bicarbonate solution of choline periodide, by

a concentrated solution of iodine in potassium iodide. The weak base betaine is similarly precipitated from the filtrate after acidification. quantities of choline may be estimated physiologically after conversion into the acetyl derivative (see below, also Guggenheim and Loffler, Biochem. Zeitsch 1916, 74, 208; Fuhner, ibid. 1916, 77, 408)

Qualitatively choline is recognised as the dichroitic crystalline periodide (Florence's test for semen; Bocanus, Zeitsch physiol. Chem. 1901, 34, 339, Rosenheim, J. Physiol. 1905-6, 33, 220), and by conversion into other salts. The mercuri-chloride C<sub>5</sub>H<sub>14</sub>ONCl,6HgCl<sub>2</sub>,H<sub>2</sub>O, crossed hexagonal prisms, mp 249°-251° dissolves in about 50 parts of water at room temperature, but in alcoholic solution the limit of precipitation is given as 1 2,000,000. The platinichloride  $(C_5H_{14}(N))_2PtCl_6$  dissolves in 5 82 parts of water at 21° and is dimorphous, the conversion of the regular isotropic form into the anisotropic monoclinic one is the surest means of detecting choline without analysis (Kauff-mann and Voilander, Ber 1910, 43, 2735). The auruchloride dissolves in 75 parts of water at 21° and in hot alcohol. Neither this nor the platimehloride has a characteristic melting-The free base is a syrup, many of the salts are deliquescent (for details see Gulewitsch, Zeitsch physiol, Chem. 1898, 24, 513) soluble non-deliquescent salts, see Ger Pats Nos. 290523 and 290740 (Chem. Soc. Abstr. 1916, 1 468 and 548) The physiological action of choline is in general antagonistic to that of atropine, and not very intense, intravenous injection lowers the blood pressure (e.g. Mendel, Underhill and Renshaw, J. Pharm exp. Therap. 1912, 3, 649) Esterification of the alcoholic hydroxyl enormously increases the activity Acetylcholine is 100,000 times as depressent as choline (Hunt and Taveau, US Hyg Lab. Bull No 73, and physiological estimation above) Acetylcholine occurs in ergot  $(q \ v)$ , but is very readily hydrolysed and is absent from most extracts. The nitrous acid ester, long regarded as 'synthetic, or pseudo-muscarme, is formed by the action of concentrated nitric acid on choline The demonstration by Ewins (Biochem J. 1914, 8, 209) that the reaction product is not an aldehyde, as was commonly supposed, explains the differences in physiological action between it and muscarine from Amanita muscaria, the Fly This natural product requires further investigation, for it is apparently not identical with betaine aldehyde as was supposed by its discoverers. The alkaloid sinapin in black mustard seeds is the choline ester of sinapic acid Many other choline esters and related derivatives have been prepared by Hunt and Taveau (Lc), Menge (J Biol Chem. 1911, 10, 399, and U.S. Hyg Lab Bull No 96), Ewins (Biochem J. 1914, 8, 366), and by Fourneau and Page (Bull See chim 1914 [iv] 15, 544) latter, in connection with the physiological action of snake venom, found that palmityl and stearyl choline are strongly hamolytic

Other bases related to choline are neurine or vinyltrimethyl ammonium hydroxide

# CH<sub>2</sub> CH·N(CH<sub>3</sub>)<sub>3</sub>OH

formed from choline by putiefaction and with a similar physiological action but 10-

20 times as toxic Amino cthylalcohol or hydroxy ethylamine ('colamine') CH2OH CH2NH2 was obtained by Thudichum in 1884 by the hydrolysis of the brain phosphatide kephalin, in which it entirely replaces choline It has been more recently recognised in other phosphatides, Trier (Zeitsch physiol Chem. 1913, 86, 1, 141, 153, 407) For a full account of choline and allied substances, see Barger, The Simpler Natural Bases, Longmans, 1914

CHOLINE ESTERS v MUSCARINE and ERGOT CHONDRIN. A substance allied to and much resembling gelatin Obtained by boiling cartilage with water When dried it is a hard, horny substance which softens to a jelly in cold water, and is entirely dissolved by boiling water It is precipitated from its aqueous solution by nearly all acids, in excess of certain of which it redissolves, and by alum, lead acetate, copper and iron sulphates, and mercuric and mercurous These reactions are said to distinguish nitrates it from gelatin.

By Morochowitz, chondin is regarded as a mixture of gelatin, mucin, and salts (v Glue)

CHROMATOMETERS v COLORIMETERS CHROMAZONE BLUE, -RED v AZO- COLOUR-ING MATTERS

CHROME BLACKS, -DEEP BLACK, -BROWNS, -FAST BLACK, -FAST CYANINE, -FAST YELLOWS, -YELLOW, -ORANGE, -PATENT GREEN, " AZO- COLOURING MATTERS -ORANGE.

CHROME VIOLET. Syn Mauveine, Perkin's violet (v. Azines and colouring matters) The term is also applied to the sodium or ammonium salt of aurintmearboxylie acid (v TRIPHENYLMETHANE COLOURING MATTERS)

CHROMITE or Chrome-iron-ore A member of the spinel group of minerals, consisting essentially of ferrous oxide and chromic oxide FeO  $\text{Cr}_2 \vec{O}_3$ , or  $\text{Fe}(\vec{r}_2 O_4)$ , analogous to the aluminates. The ferrous oxide is, however, often partly replaced by magnesia, and the chromic oxide by alumina, so that there is a passage to chrome-spinel or picotite; on the other hand, with a replacement of chromic oxide by ferric oxide, there may be a transition to magnetite (Fe<sub>3</sub>O<sub>4</sub> or FeFe<sub>2</sub>O<sub>4</sub>) All these minerals are cubic in crystallisation and belong to the same isomorphous group—the spinel group. The actual amount of chromic oxide varies considerably, as shown by the following analyses, the ordinary ore, as mined, contains on an average 45 p c Cr<sub>2</sub>O<sub>3</sub> .-

	Cr <sub>2</sub> O <sub>2</sub>	$Al_2O_3$	Fe <sub>2</sub> O <sub>3</sub>	FeO	MgO
Ι	67.9			32.1	
$\Pi$	44 15	$22\ 41$	5.78	11 76	$15\ 67$
Ш	$59\ 20$	7.15	n d	$25\ 02$	4 42
IV	41.23	24 58	$2\ 28$	16.99	14 77
V	$56\ 54$	12 13		18 01	14 08

I Calculated for FeCr<sub>2</sub>CrO<sub>4</sub>
11. Franklin, Macon co<sup>2</sup>, North Carolina
111. Price's Creek, Yancey co<sup>2</sup>, North Carolina (J. H. Pratt, Amer J. Sci. 1899, vii. 281 Also
SiO<sub>2</sub> 2 80, MnO 0 69
IV Tampadel, Lower Silesia (H. Traube, Zeits
Deutsch Geol Ges 1894, xlvi. 50)
V. Dun Mountain, New Zealand.
(For analyses of chromite isolated from meteorites

(For analyses of chromite isolated from meteorites, see W Tassin, Proc. U S. Nat. Museum, 1908, xxxiv 685)

In general appearance, chrome-iron-ore is very like magnetite, but it is readily distinguished from this by the dark-brown colour of its streak CHROMIUM 233

or powder, and by the fact that it is only slightly, if at all, magnetic Sp gr about 45 Crystals are of rare occurrence, the mineral being usually found as grains disseminated in basic rocks of igneous origin, especially those rich in olivine (viz the peridotites) Sometimes these grains are segregated into nodular masses of considerable size and with a granular to compact texture Rocks of this kind, when subjected to weathering processes, become hydrated and altered into serpentine, and for this reason most of the workable deposits of chrome-non-ore are in serpentine rocks. With the denudation and breaking down of these tocks, the heavy grains of the chemically resisting chromite collect in the beds of streams and rivers, and some of the deposits worked in the Ural Mountains are of this nature

Chromite is the only commercial source of chromium and its compounds It is used for the preparation of the oxides and chromates employed as pigments, and in dyeing, calicoprinting and tanning, in the manufacture of chrome-steel and feriochrome, and for chromebricks and furnace linings. It is mined in European and Asiatic Turkey, Greece, the Ural Mountains, New Caledonia, India, Baluchistan, Rhodesia, Quebec, California and Oregon, and formerly, to a considerable extent, at Bare Hills, near Baltimore in Maryland, and in Lancaster co in Pennsylvania Quarries were, at one time, worked on Unst, one of the Shetland Isles (Special Reports on the Mineral Resources of Great Butain, vol v Mem Geol Survey, 1916)

On the Califorman deposits, see J. S. Diller, Mineral Resources of the United States, U.S. Geol Survey, for 1915, 1916, those of Canada, see F. Cirkel, Dept. of Mines, Canada, No. 29, 1909, and those of Rhodesia, A. E. V. Zealley, Trans. Geol. Soc. S. Africa, 1914, xvii. 60

LJS CHROMIUM. (Fr Chrome; Ger Chrom) Symbol Cr At wt 520 Chromium occurs principally as chrome-non-ore, or chromite (qv) Found largely in Rhodesia, New Caledonia, Russia, Greece, Asiatic Turkey, and India Chromite deposits are also worked to a small extent in Bosnia-Herzegovina, Japan, California, Oregon, Transvaal, and Newfoundland principal producers in 1913 were New Caledonia 62,000 tons, Rhodesia 63,000 tons, Russia 21,000, these countries furnishing about 90 pc of the world's total As sesquioxide C12O3, it is found in small quantities in chrome ochre, generally mixed with clay and associated with chromite, in France and Siberia As lead chromate it occurs in crocoisite and malanchioite PbCrO4, and as a basic lead chromate in the rare mineral phænicite 3PbO 2CiO3. It is usually present in the form of *chromite* in meteorites, but rarely to the extent of more than 1 pc. The rarely to the extent of more than 1 pc greenish colour of the emerald, verd-antique marble, serpentine, sapphire, and many other minerals, is due to the presence of chromium sesquioxide.

Preparation.—Metallic chromium may be prepared by several methods Deville produced it in combination with a small quantity of carbon, in ingots weighing as much as 100 grams, by the ignition of a mixture of chromium sesquioxide and sugar in a lime

crucible, at the highest temperature of a blast furnace

Wohler (Annalen, 111, 230) prepared chromum by fusing a mixture of the violet chromium scsquichloride with twice its weight of zine under a layer of equal parts of potassium and sodium chlorides. The mixture of zine and chromium so produced was treated with dilute nitric acid to remove the zine, and the chromium remained as a light-greenish powder consisting of minute octahedra belonging to the quadratic system (Bolley), of sp. gr. 6-81 according to Wohler, 7-3 according to Bunsen

Chromium may be prepared by the electrolysis of its salts, but the purity and condition of the deposit vary greatly with current strength and solution concentration. A bright deposit, capable of receiving a polish, can be obtained by the use of a hot 25 p c acid solution of chromic chloride, with a current density of about 40–50 amperes per square foot. As in the case of nickel, the deposits have a strong tendency to peel (Cowper Coles, Chem. News, 81, 16).

It is most easily obtained in small quantities by heating equivalent parts of chromic oxide (or mixture of the oxide and anhydride) with aluminium powder. When started, the heat of the reaction causes it to continue and a fused mass of chromium of a high degree of purity is obtained (Goldschmidt, Annalen, 301, 19)

In large quantities it is best prepared by heating the oxide with carbon in the electric furnace. The crude metal thus obtained always contains carbules, of which three definite compounds appear to exist. Cr<sub>3</sub>C<sub>2</sub>, Cr<sub>4</sub>C, and Cr<sub>5</sub>C<sub>2</sub>. It may be refined by further heating in the electric furnace on a bed of lime mixed with chromic oxide. The refined metal crystallises in cubes or octahedra and still contains about 1.5 p.c. carbon. By further refining, however, with a double oxide of chromium and calcium, the pure metal is obtained (Moissan, The Electric Furnace).

Pure chromium is a steel-grey metal, slightly harder than glass, the carbides, however, are harder than quartz, so that a metal with between 1 5 and 3 p e carbon can only be cut and polished by diamond dust Its density is 6 92, and it oxidises superficially in moist air, the meltingpoint is probably above 2000° (1510°, Johnston, J. Ind. Eng. Chem. 1917, 9, 873). The pure finely divided metal burns rapidly when heated in the blowpipe flame, and a pyrophonic powder may be prepared by distillation of chromium amalgam in vacuó (v in/ra) It also unites with nitrogen when heated in nitrogen or ammonia, forming a very stable nitride Ciude chromium does not burn until heated to above It combines with sulphur vapour at 700°, and when heated to redness in hydrogen chloride, forms chromous chloride Electrolytic chromium occludes 250 times its volume of Aqueous hydrochloric acid acts hydrogen slowly on the metal, but the action is accelerated if the chromium is made the anode in electrolytic hydrochloric acid Heating with sulphuric acid causes evolution of sulphur dioxide, concentrated nitric acid and aqua regia have no As in the case of iron, active and passive action varieties of chromium exist; in general, the metal is rendered passive by oxidising solutions, whilst

234 CHROMIUM.

hydrogen ions make it active. In the electromotive series, the active chromium immediately follows zinc, the mactive metal is near platinum. The cause of the passive state is usually ascribed to the existence of a superficial film of oxide, but

this view has been disputed

Chromium carbide boils at 2270° C., and the vapour consists entirely of chromium. Between 2050° and 2250° the fused mass remains constant in composition and corresponds with  ${\rm Cr_3C_2}$  Below 2050° down to 1875° graphite and the carbide  ${\rm Cr_3C_2}$  exist side by side as solid phases  ${\rm Cr_6C_2}$  is also shown to exist (Zeitsch. Elec. Chem 18, 24–157)

Alloys. Various alloys of chromium have been prepared, chiefly by means of the electric furnace, by the 'thermit' method, or by the

mixture of the molten metals

Molten zinc dissolves but little chromium, a hard and brittle alloy has been obtained in the form of hexagonal lamellæ Aluminium and chromium mixtures, containing between 5 and 55 pc chromium, separate into two liquid layers, and probably contain a compound, Cr<sub>3</sub>Al The alloys with a low percentage of chromium are brittle, one containing 13 pc chromium can be powdered in a mortar.

Alloys with antimony are brittle, and compounds represented by SbCr and Sb<sub>2</sub>Cr are known Chromium alloys with copper with difficulty, but an alloy can be obtained by stirring in copper oxide into a molten alloy of aluminium and chromium (Moissan) The liquid metals are miscible only to a small

Cobalt and chromium are miscible in all proportions, both in the liquid and the solid states, and yield a mixture of minimum melting-point 1320° with 47 pc Cr Homogeneous alloys, containing above 30 p.c Cr, when cooled, separate

into two sets of crystals, distinguishable by electrolytic etching.

extent

Nickel chromium alloys with less than 90 p c

nickel are non-magnetic

Silver and chromium, although partially miscible in the liquid state, form no solid solution Molten cadmium does not dissolve any chromium. Molten mixtures of lead and chromium, containing more than 27 pc lead, separate into this mixture and pure lead

Chromium also forms silicides and bondes in the electric furnace, of the former, compounds represented by SiCr<sub>3</sub>, SiCr<sub>2</sub>, Si<sub>2</sub>Cr<sub>3</sub>, and Si<sub>3</sub>Cr, have been described. They are very hard— SiCr<sub>2</sub> is harder than corundum—and are not attacked by ordinary acids, though attacked readily by hydrofluoric acid and by aqua regia. The bordes Cr<sub>3</sub>B<sub>2</sub> and CrB are metallic looking crystalline solids, not attacked even by a mixture of aqua regia and hydrofluoric acid The compound CrB is stated to be weakly magnetic The phosphide CrP, obtained from the metal and copper phosphide, forms dull-grey crystals, sp.gr. 571, which burn in oxygen, forming chromium phosphate, but are unattacked by all acids except a mixture of hydrofluoric acid and aqua regia

By electrolysis of a strongly acid solution of chromic chloride, using a platinum anode and a merculy cathode, an amalgam Hg<sub>3</sub>Cr is obtained, soft and brilliant, and altering but slightly in air. By pressure, a harder amalgam

HgCr as obtained, which alters in air more rapidly. By distillation in vacuó, both lose mercury and yield pyrophoric chromium (v. supra). An amalgam may also be obtained by the action of sodium amalgam on a chromium salt.

With *iron* the alloys of chromium are of great interest. The presence of chromium in iron or steel produces a much finer texture, greater hardness, tenacity, and elasticity, and greater smoothness of fracture. Chromium cannot be used to replace carbon in steel, as has

been asserted (Boussingault)

When crystalline, the tendency of these alloys is to produce needles instead of plates, as in the case of manganese. Chromium lowers the Ar<sub>3</sub> point eventually causing its disappearance, whilst it raises the Ar<sub>2</sub> point of the steel, the proportion of chromium required being less the higher the carbon, eg with 0.2 pc. carbon 5 p c chromium is required Hence chiomium steels are self-hardening (i e. air-hardening). Steels containing 15 p c chromium have been used for tools and have to be cast to shape With 05 pc chromium and 05 pc carbon the steel is used for well bits and just and has no equal in the hardened state. With 1 pc chromium and high carbon the steel is used in bearing balls, cones, roller bearings and crushing machinery. These low chromium steels when well annealed can be machined Its principal use is in conjunction with other alloy steels, especially nickel steels, to which it imparts greater toughness and hardness As such it is used extensively in automobile parts, and when used in armour plates they are case-hardened and quenched. Chrome steel is also used in the manufacture of special kinds of files The alloy is usually prepared of the requisite composition by the addition of a definite amount of 'ferrochrome,' containing from 40 to 85 p c chromium to the molten steel. Ferrochrome was formerly prepared from 11ch chrome-1ron ores in blast furnaces by the use of coke and hot highpressure blast, or in crucibles. In 1890 the manufacture was commenced in the electric furnace and has gradually superseded the other methods. In the reduction of chromium from chromite to form ferrochiome, the iron being reduced, reduction of chromic oxide begins about 1185° (' Thirty parts of pure carbon are theoretically necessary for every 100 parts of non and chromium reduced On the basis of the double carbide Fc<sub>1</sub>C,Cr<sub>2</sub>C<sub>2</sub> being formed in reduction of the chromite the following reaction occurs:

 $\begin{array}{l} 9 \mathrm{FeO}, 9 \mathrm{Cr_2O_3} + 50 \mathrm{C} \\ = [2 (\mathrm{Fe_3C}, 3 \mathrm{Cr_3C_2}) + 3 \mathrm{Fe}] + 36 \mathrm{CO} \end{array}$ 

The ferrochrome resulting would contain theoretically 10 4 pc carbon, 31 4 p.c. iron, and 58 2 pc chromium. The ore has to be finely ground and is then mixed with anthracite coal in the proportion of 37 parts of carbon to 100 parts of ore which forms the above alloy and charged into the furnace. The power consumption in a 750 kilowatt furnace of the Alby carbide type at Kapperaen, Norway, was 3.0 kw. hours per pound, or 0.68 kw. year per, short ton for 5 pc carbon and 65 pc chromium ferrochrome. The percentage of carbon in the ferrochrome cannot be kept low by regulating the coal

CHROMIUM.

235

charged into the furnace, but decarburisation can be effected with an oxide slag of iron or chromium after tapping the slag from the first reduction. (Amer. Electrochem. Soc. 1913, 176.)

The addition of chromium decreases the magnetic properties, but all alloys, to 80 p.c. chromium, are magnetic. The magnetic property of chromium is not increased by cooling in solid hydrogen (Compt rend. 150, 687). A carbide of chromium and tungsten of great hardness and of sp gr 8.41, has been produced in the electric furnace, and to it the special properties of chrome tungsten steels are probably due. These latter (as well as chrome molybdenum steels), containing up to 3 p.c. chromium and 16 p c tungsten, are used for the manufacture of machine tools. Cobalt is frequently added to both classes of tool steel.

Detection of Chromium. When heated strongly, all compounds of chromium impart a green colour to a borax bead in both the reducing and oxidising flames On ignition with potassium nitrate, all chromium compounds produce a yellow colour, due to the presence of potassium chromate. If this is dissolved in water, the addition of a solution of lead acctate produces a precipitate of chrome yellow similar oxidation occurs when chromium compounds are fused with sodium peroxide, or when solutions of chromium salts are boiled with sodium peroxide. A delicate test for a soluble chromate is to add excess of hydrogen peroxide to a little of the solution in a test tube and shake up with a few cc of ether, a blue colouration indicates chromium (See under Perchromic

Solutions of chromic salts of salts of sesquioxide of chromium have an acid reaction With caustic alkalis they produce a green precipitate of hydrated oxide partially soluble in excess of the reagent, but reprecipitated on boiling the solution. With carbonates, a green precipitate is produced, likewise soluble in excess.

Chromates (in which the chromium exists as an acid) are usually strongly coloured. Soluble chromates are reduced, when warmed with sulphuric acid and a reducing agent such as alcohol, with the production of a green colour. The chromium is then present as a base, and may be precipitated, as already mentioned. When heated with hydrochloric acid, they are partially reduced, with evolution of chlorine and formation of chromium chloride and chloride of the metal present as chromate, together with the formation of chromyl chloride CrO<sub>2</sub>Cl<sub>2</sub>

Solutions of chromates containing no free acid except acetic acid give a yellow precipitate with a salt of lead, a red precipitate with silver nitrate, and a yellow precipitate with a salt of barium

Estimation —When present as a base, chromium is usually estimated as sesquioxide  $Cr_2O_3$ . For this purpose, the solution is heated nearly to boiling, ammonia solution added in slight excess, and the temperature maintained until the liquid is perfectly colourless, indicating that the hydroxide, which is slightly soluble in excess of ammonia, is completely precipitated. The precipitate is well washed by decantation and transferred to a filter, washed thoroughly with hot water, dried, and ignited. The oxide produced

contains 68.63 p.c. of chromium The precipita. tion is not complete in presence of organic matter, and when a glass vessel is used, the precipitate always contains a small percentage of silica Chromium, when present as a chromate, cannot be estimated directly by that method quires to be first reduced to a base For this purpose, hydrochloric acid and a small quantity of alcohol are added, and the solution heated until the odour of alcohol is dispelled chromium may then be precipitated as above For the direct estimation of chromium in chromates, the precipitation with lead acetate is most satisfactory The solution is mixed with sodium acetate and acidified strongly with acetic A solution of normal lead acetate is then added, and the precipitate of PbCiO<sub>4</sub> is collected on a weighed filter, washed, dried at 100°, and weighed It contains 16 19 p c of chromium, or 31 06 pc of chromic anhydride CrO<sub>3</sub> Or the precipitate may be dissolved in well-cooled dilute hydrochloric acid (1 of HCl to 3H<sub>2</sub>O) potassium iodide added and the liberated iodine titrated with N/10 sodium thiosulphate. The following scheme may be adopted for the estimation of chromium in chrome iron ore A few grams of the carefully sampled mineral are ground in an agate mortar and passed through a fine muslin The dust so produced is further ground in small portions until all grittiness, on passing a little between the fingers, has disappeared and the ore cakes as an impalpable powder round the About 0 5 gram of the powder is weighed into a platinum crucible of about 50 c c capacity, covered with twelve times its weight of recently fused hydrogen potassium sulphate (potassium bisulphate), and gently heated to just fuse the After keeping at this temperature for 15 or 20 minutes, the heat is gradually increased until the crucible bottom becomes red hot. The fused mass should not be allowed to rise above half-way up the crucible The mixture soon fuses quietly, and evolves dense fumes of sulphuric acid, the heat is gradually increased to bright redness In about half an hour, 6 parts of powdered anhydrous sodium carbonate are added and the mixture again fused for an hour at a red heat, 6 parts of nitre being added in small portions. The temperature is then raised to a full red heat for about 20 minutes, the crucible cooled and transferred to a porcelain basin where the mass is boiled out with water The solution is filtered, and the residue washed with hot water until the filtrate comes through The filter and its contents are dired, the precipitate detached and placed with that still remaining in the basin, the filter paper burned, and the ash also added. To ascertain if the fusion has been satisfactory, this residue is digested with moderately strong hydrochloric acid, when the whole should dissolve Any insoluble black residue indicates either imperfect grinding or insufficient fusion It must be collected on a filter, dried, and the whole ignited in a crucible and treated with potassium bisulphate, &c, as in the first instance. The aqueous extract, after fusion, is mixed with the main solu-To the liquid, a few grams of ammonium nitrate are added, and the whole evaporated to dryness, taken up with water, and filtered from the alumina, silica, &c., into a porcelain basin. An excess of sulphurous acid solution is then 236 CHROM1UM

added, and the solution heated until that gas is nearly expelled. The chromate, having thus become reduced to chromium sulphate, a slight excess of ammona is added, and the solution boiled until colourless. The precipitated chromium hydroxide is washed by decantation with hot water, transferred to the filter, and washed The use of the suction with hot water six times pump is of great assistance in this operation. The precipitate and filter are then diied, transferred to a weighed platinum crucible, and heated sently until the paper is charred. The crucible gently until the paper is charred. The crucible lid is then placed at the edge of the crucible so that the flame may reverberate into the crucible, and the whole ignited strongly for 10 or 15 The weighed precipitate should show no yellowish colour on treatment with a few drops of water, if such a colour is produced it indicates imperfect washing of the precipitate The oxide contains 68 63 p c from alkalı salts of chromium

The oxidation may also be effected by means of sodium peroxide, and the chromate produced, instead of being reduced to chromic oxide, may be estimated as chromic acid volumetrically. For this, various methods are available, eg(1) boiling with standard assenious oxide and estimating the excess by standard rodine, (2) adding potassium rodide and titrating rodine liberated by standard throsulphate; (3) adding standard ferrous ammonium sulphate solution and titrating excess with standard potassium dichromate

A speedy and accurate method for the estimation of chromium in ferrochrome or chromite, is the following 1 part of the ore is finely powdered, as described above, and fused with a mixture of 5 parts of sodium peroxide and 8 parts of caustic soda in a silver crucible The aqueous extract is filtered and again similarly treated, three or four such fusions are necessary to dissolve all the ore, when the only residue is a little silver extracted from the crucible The filtrate is acidified with sulphuric acid, boiled for 40 minutes and diluted to a known volume aliquot part is then titrated with potassium iodide and standard sodium thiosulphate

Chromates may also be readily estimated in acid solution by hydrazine sulphate, whereby nutrogen is liberated, 100 c c of nitrogen at N.T P. being equivalent to 0.59576 gram Ci

#### COMPOUNDS OF CHROMIUM.

Chromium forms a number of oxides, of these, chromous oxide ('r(), and chromic oxide or chromium sesquioxide Cr<sub>2</sub>O<sub>3</sub>, are definite basic oxides, yielding with acids salts in which chromium is present as a divalent or trivalent cation, forming the chromous and the chromic salts. Chromic oxide can also combine with bases to form salts, the chromites, of the type MCrO<sub>2</sub>, in which the chromium is present in the monovalent anion CrO'2. Chromic anhydride CrO3 is an acid anhydride forming, with bases, salts of the type M<sub>2</sub>CrO<sub>4</sub>, the chromates, or M<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>, the dichromates, in which the chromium exists as the divalent anion CrO"4 or Cr2O"7 Other oxides have been prepared and are usually regarded as compounds of the basic and acid oxides, e.g. Cr<sub>2</sub>O<sub>3</sub> CrO<sub>3</sub>=chromic chromate. The chromous salts are of no industrial importance at present.

The salts of chromium are coloured shades of

violet or green; all solutions of violet chromium salts show similar absorption towards light, spectrum due therefore to the chromium ion in solutions of green complex salts the absorption band and limit of complete absorption are both shifted towards the red end of the spectrum.

Chromous oxide, CrO, is most conveniently prepared by the action of dilute nitric acid on chronium amalgam which dissolves the mercury leaving chromous oxide as a black powder.

Chromous salts. Traube & Goodson (Ber. 1916, 1679) have prepared chromous salts by the electrolytic reduction of the corresponding chromic salts as a cathode of pure lead. The most favourable conditions for the reduction of violet chromic salts are in moderately acid concentrated solutions with a current density of 2.5 amps per square decimetre. The more common green salts can be used in more concentrated solutions, but require a greater density of current.

Chromous salts are gradually oxidised in acid solution with the liberation of hydrogen, thus  $2 {\rm CrO} + {\rm H_2O} = {\rm Cr_2O_3} + {\rm H_2}$ . They are on this account able to reduce compounds with double or triple linkings in the presence of water. Acetylene is thus reduced to ethylene, but not to ethane. Nitrous oxide in the presence of alkali hydroxide is reduced to integen. Nitric acid and hydroxylamme are reduced quantitatively to ammonia (Ber 1916, 1692). Chromous salts from stable double compounds with hydrazine salts and may be precipitated from chromous acetate dissolved in air free water, covered with petroleum ether. Chromous di-hydrazine chloride  ${\rm CrCl}_22{\rm N}_2{\rm H}_4$  is a blue-coloured powder sparingly soluble in water.

Chromic oxide, Chromium sesquioxide  $({\rm Gr}_2{\rm O}_3, {\rm This})$  compound is produced by the oxidation of metallic chromium and by ignition of chromic hydroxide, chromic anhydride, and ceitain chomates

Wohler prepared this oxide in fine, small rhombohedral crystals by passing the vapour of chromyl dichloride through a tube heated to redness. The crystals are isomorphous with corundum and of equal hardness, their sp gi is 5.21. Crystalline chromic oxide is also obtained when a mixture of potassium dichromate and sodium chloride is heated to redness (Ditte, Compt. rend. 134, 336). The finest coloured amorphous chromate Hg<sub>2</sub>CrO<sub>4</sub> in a covered crucible, mercury and oxygen escape, and the oxide remains as a green powder. For the preparation of this substance on the large scale, a great number of methods are recommended

Boil a solution of potassium dichromate with half its weight of flowers of sulphur so long as the green hydroxide is precipitated. The addition of a little potash solution, by forming potassium sulphide, accelerates the decomposition. The precipitate is filtered from the solution containing potassium sulphate, and washed. The sulphur retained in the precipitate may be removed by heating. Instead of performing the reaction in the wet way, the mixture may be ignified in a crucible, and the resultant mass digested with water (Lassaigne, Ann. Chim. Phys. [3] 14, 299). Wohler (Pogg. Ann. 10, 46) heats a mixture of potassium

dichromate with its own weight of ammonium chloride and a small quantity of sodium carbonate, and purifies the residue by washing

According to Barian (Rev Scient. 20, 425), a very pure colour, suitable for colouring fine porcelain, is produced by igniting in a crucible a mixture of 4 parts of potassium chromate and 1 part of starch. The mass is washed free from potassium carbonate and re-ignited

Chromic sesquioxide is a green pigment of great permanence. It is not acted upon by chlorine or sulphur gases or by an intense heat It melts at 1990°, and crystallises on cooling

Chromium hydroxide  $(r_4(OH)_5O_2, 10H_2O)$  or  $(r_4(OH)_{10}O, 9H_2O)$  (Jovischitsch Heid Chim Acta, 1920, 3, 46) absorbs carbon dioxide from the atmosphere until saturation limit is reached. The compound appears to be  $(CO_3(Cr(OH)_6)_2, 8H_2O)$  or  $(Cr_2(OH)_5O)$   $(CO_1OH_2O)$  (Jovischitsch) It can be dried at (100)0 without losing  $(CO_2)$ 0, which is, however, liberated by acids (Compt. red. 158, 782)

Hydrogen and carbon monoxide are without action, but it is reduced to the metallic state on heating with carbon at a temperature of 1185°

or higher.

This oxide is largely used under the names chrome green (mixtures of chrome yellow and Prussian blue are also frequently called chrome greens and must not be confounded with the true chrome green), and ultramaine green for imparting a green colour to glass, porcelain, &c It is used for producing the chrome red glaze in ceramic ware, as a pigment, in oil and water colours and in printing, and as a mordant in calico-printing and dyeing. Certain hydrated oxides are also used under various names, their colours are, generally speaking, brighter than that of the anhydrous oxide, but they usually contain small quantities of other substances besides the oxide and water

Guignet's Green; Pennettier's Green, Emerald Green; Veridian Cr4O3(OH)6 This pigment appears to be identical with that formerly manufactured in secret by Pennettier According to Guignet's method, 3 parts of bone acid and I part of potassium dichromate are heated to dull redness in a reverberatory furnace mass swells up, evolves oxygen, and becomes of a fine green colour; it contains the borates of potassium and chromium, or a double borate of those two metals. It is boiled with water, whereby the borate of chromium is decomposed into boric acid and hydrated chromium sesquioxide, potassium borate also remaining in solu-tion. The precipitate is well washed, dried, and finely ground. It usually contains boric acid even after thorough washing. The washings and mother liquors are evaporated for the recovery of the boric acid.

According to the Chem. Zeit 9, 851, the process now used on the large scale is as follows: The boric acid is first purified by solution in how water, treated with animal charcoal, and recrystallised. The crystals are dried in a centrifugal machine. The mother liquor may be employed three or four times for recrystallisation of fresh portions of acid, but becomes then too impure for further use. Eight parts of potassium dichromate and 3 parts of purified boric acid are ground into a stiff paste with water, and the mixture in charges of about 1 cwt. is heated

to dull redness for about 4 hours in a reverberatory furnace. The fused mass is thrown into water, and washed repeatedly by decantation, the pigment is ground while wet, again washed, filtered, and dried. The first two washings contain considerable quantities of potassium borate, which is recovered by evaporating in leaden pans, adding hydrochloric acid and allowing the boric acid formed to crystallise. The mother liquor is further evaporated and crystallised. The crystals are redissolved, the lead (from the pans) precipitated by hydrogen sulphide, and the filtrate recrystallised. In this manner, from 70 to 75 p.c. of the boric acid is recovered.

Guignet's green is the most permanent green pigment known, it is unacted upon by light of concentrated boiling alkalis; it is not affected by acids in the cold, but hot hydrochloric acid slowly dissolves it. When heated to 200° it becomes anhydrous. It is a fine green pigment largely used for the same purposes as the dangerous arsenical greens. It possesses good covering power, and can be mixed with other pigments without alteration. When mixed with lead chromate, it is employed for the pioduction of a pale-green colour in landscape painting.

Other chrome greens are obtained by the use These always contain of sodium phosphate some phosphoric acid They are, however, not as brilliant as the oxide chrome greens. Various methods of preparation are employed Arnaudon dissolves ammonium phosphate with slight evcess of potassium dichiomate in a little boiling water, evaporates until mass solidities on cooling, heats at 80° and later at 200°, washes with hot water, dries and powders Another method consists in boiling a solution of 10 lbs potassium dichromate with 18 lbs sodium phosphate, adding 10 lbs. sodium thiosulphate and a little hydrochloric acid, and boiling until precipitation is complete. precipitate is washed, ground, and dried

According to Bong (Ber 29, 201), a blue pigment, retaining its colour at high temperatures, suitable for colouring porcelain, &c, is produced by igniting strongly, in the air, an intimate mixture of 5 parts borie acid, 15 parts alumina, 20 parts magnesium carbonate, and 2 parts

barium chormate

Chromyl dichloride CrO<sub>2</sub>Cl<sub>2</sub>. Mossan (Compt. rend 98, 1581) prepared chromyl chloride by treating dry chiomic anhydride with hydrochloric acid gas It forms at the ordinary temperature but the action is accelerated by heating It may also be made by distilling potassium bichromate (3 parts), common salt (3 parts) with sulphuric acid (9 parts) in an atmosphere of carbon-dioxide. The corresponding bromide and iodide cannot be made by this method, but the fluoride can, the formula for the latter being CrO<sub>2</sub>F<sub>2</sub> (Gazz chim ital 16, 218) Perkin (Chem Soc Trans 1907, 191) found that the best way of preparing this compound is to dissolve chromic anhydride (50 grams) in strong hydrochloric acid (170 c c) to which is added 100 cc of strong sulphuric acid, 20 cc at a Chromyl chloride sinks as a red liquid and may be separated in a tap funnel. B p at 760 mm. is  $116^{\circ}63^{\circ}$  m p.  $-96^{\circ}5^{\circ}$ ,  $\pm 0.5^{\circ}$ . It is a deep red liquid resembling bromme,  $D_4^{-47}$  2 0528–2 0515,  $D_4^{\circ}$  1 9591–1 9582,  $D_4^{25}$  1 9124–1 9113.

Its vapour is not dissociated at 181° C heated in a closed tube to  $180^{\circ}$ – $190^{\circ}$  trichromyl chloride (CrO<sub>2</sub>)<sub>3</sub>Cl<sub>2</sub> is formed. It is a black non-crystalline powder deliquescing in air Chromyl dichloride combines with phosphorus trichloride in carbon tetrachloride solution to

form CrOCLPOCL

Chromic chloride CrCl3 or Cr2Cl6. anhydrous chloride is obtained by heating a mixture of carbon and chromium sesquioxide in a current of chlorine It can be also conveniently prepared by Burion's method (Compt rend. 148, 170) by passing the vapour of sulphur chloride, b p. 137-90, over the oxide gradually raised to a red heat It forms pale-violet scales of sp.gr 3 03; it is almost insoluble in water, but dissolves readily, being transformed to the green variety (v. infra), if only a minute trace of chromous chloride (1 in 20,000) is present Chromic chloride may be prepared in solution by dissolving the hydroxide in hydrochloric acid It forms two isomorphous hexahydrated salts (CrCl<sub>3</sub>,6H<sub>2</sub>O), one blue and the other green, and also a tetrahydrate and a decahydrate The also a tetrahydrate and a decahydrate blue hexahydrate, when dissolved in water, appears to give the ions  $[Cr(OH_2)_6]'''$  and 3Cl', all the chlorine being precipitated by silver nitrate; the green salt, however, at first forms the ions  $[Cr(OH_2)_4Cl_2,2II_2O]$  and Cl', only  $\frac{1}{3}$  its chlorine being immediately precipitated (Weiner and Gubser, Ber 1901, 34, 1579). In calico-printing a neutral solution of chromic chloride, prepared from potassium dichromate, is frequently used The following method is recommended by De Karrur Dissolve 9 lbs. of potassium dichromate in 5 gallons of boiling water Prepare also a solution of 9 to 11 lbs. of white arsenic in 24 gallons of boiling water, and mix, with constant stirring, with the solution of dichromate greenish precipitate of chromic hydroxide soon separates, it is cooled, filtered, and washed with boiling water The precipitate is added in portions to a solution of hydrochloric acid (diluted until it no longer fumes), with constant stirring, until a portion remains undissolved solution of soda is then added and well mixed. until the hydroxide begins to precipitate dark-green solution, being thus neutralised, is evaporated to 46°B and is ready for use A basic chloride CrCl(OH), is also employed as a mordant for silk and cotton

Chromic fluoride CrF<sub>3</sub>,4H<sub>2</sub>O. This product is used in the printing and dyeing of woollen goods. It is a fine crystalline green powder, very soluble in water It has a corroding action on glass, and is therefore best kept in lead or wooden vessels It may be used as a mordant with both vegetable and animal fabrics, and is often employed in the place of acetate or nitrate of chromium for printing, and in general gives Alizarın dyes, corulein, gallein, finer results &c, and logwood extracts, are fixed as perfectly as with acetate of chromium. One advantage in using it is that it is solid. It has no injurious effect on the pile, and is often used in place of chromates where the oxidising action of these is injurious, especially with wools. Chromium silicofluoride, prepared by dissolving the oxide in hydrofluosilicic acid, has also been used in dyeing and printing in place of the fluoride, and is stated to give better results.

Chromium nitrate. By dissolving strongly

heated chromic oxide in hot concentrated nitric acid of D. 14 a solution is obtained from which the hydrate  $\text{Cr}_{\lambda}(\text{NO}_3)_{\bullet}, 15\text{H}_3\text{O}$  crystallises on cooling in the form of dark brown prisms  $[a \cdot b \cdot c = 1 \cdot 4250 \cdot 1 : 1 \cdot 1158, \beta = 93^{\circ} \cdot 10']$  In contact with dry air the crystals lose 6H2O with the formation of the grey coloured hydrate

Cr<sub>2</sub>(NO<sub>3</sub>)<sub>8</sub>,9H<sub>2</sub>O (Monatsh. 33, 9).

Chromium phosphate, formed by mixing cold solutions of chromic alum and disodium hydrogen phosphate, is a violet amorphous solid, which in contact with water becomes crystalline and has the composition CrPO4,6H2O. On boiling with water it is converted into the green crystalline hydrate ('rPO4,4H2O On further heating it loses two molecules and is converted into the dihydrate. When heated to low redness the phosphate becomes anhydrous and forms a black powder (Joseph & Rae, Chem. Soc Trans 1917, 111, 196).

Chromium phosphides. When finely divided chromium, prepared from amalgam, is heated with red phosphorus in a scaled exhausted tube the phosphide  ${\rm Cr_2P_3}$  is obtained. This on heating in hydrogen to  $440^\circ$  is converted to CrP. Both phosphides are grey and insoluble in acids

Chromium arsenides. When finely divided chromium is heated with an excess of arsenic in a sealed exhausted tube, the product powdered and again heated, the grey arsenide Cr<sub>2</sub>As<sub>3</sub>, D<sub>4</sub><sup>22</sup> 6 2 is obtained When heated in hydrogen at 480°-500° a second arsenide CrAs is obtained Both compounds are insoluble in acids

ds (Zertsch anorg Chem 86, 291)
Chromium sulphates. These are prepared by the solution of the hydroxide in sulphuric It yields a bluish-violet solution, which, on boiling, becomes green. The experiments of Recoura have shown that numbers of compounds of different constitution exist both in the violet and green varieties, but they are not of commercial interest. (f Colson, Compt. rend. 1907, 144, 79, 206, Sénéchal, ibid 1914, 159, 243 The double sulphates of chromium and potassium, sodium and ammonium, are, however, of These are known as chrome alums, importance they crystallise in fine octahedral crystals containing 24 molecules of water isomorphous with those of ordinary alum, and of corresponding composition.

Potassium chrome alum, Chrome alum Cr<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>·K<sub>2</sub>SO<sub>4</sub>,24H<sub>2</sub>O

This substance may be prepared by mixing solutions of chromic sulphate and potassium sulphate in molecular proportions, or more conveniently by the reduction of potassium dichromate in sulphuric acid solution

The best method is to add 5 parts of strong sulphuric acid to a solution of 3 parts potassium dichromate in 30 parts of water with the addition of a reducing agent, preferably a current of sulphur dioxide.

 $\begin{array}{l} {\rm K_2Cr_2O_7 + H_2SO_4 + 3SO_2 + 24H_2O} \\ {\rm = K_2SO_4 \cdot Cr_2(SO_4)_3, 24H_2O + H_2O} \end{array}$ 

When the sulphur dioxide is substituted by other reducing agents, such as alcohol, a larger quantity of sulphuric acid is required.

Chrome alum is obtained in large quantities as a by-product from the oxidation of organic substances by a mixture of potassium dichromate and sulphuric acid, as in the oxidation of anthracene to alizarin, in the manufacture of aniline violet, &c.

Pure crystallisable violet chrome alum is prepared by the electrolysis of a solution of potassium dichromate mixed with sulphuric acid. The liquor is circulated in the cathode compartment which is separated from the anode by a porous vessel. The cathode is preferably of carbon. Dilute sulphuric acid is circulated in the anode compartment and the anode may be copper with copper turnings to utilise the SO<sub>3</sub> ions. (Inter Cong. 1912.)

It crystallises in fine dark-purple octahedra, especially in presence of excess of acid, is soluble in 7 parts of cold water, forming a reddish-blue solution, which, when heated to about 70°, becomes green. It returns to its original colour

after standing some weeks

When heated to 25°-30°, the crystals lose half their water and become hlac-coloured, at 100° a further quantity of water escapes and the crystals turn green, at 350° they become anhydrous, but are still soluble in water; at a temperature slightly above 350° they become greenish-yellow, insoluble in water (Lowe), and only slowly soluble in acids. The green salt only contains \(\frac{1}{2}\) of the SO<sub>4</sub> as an acid ion, and is probably a sulphate of a sulphochromyl hydroxide—For these compounds, reference should be made to the extensive researches of Recoura (Ann Chim Phys [7], 4, 494)

Chrome alum is used in dyeing, calico print-

ing, and tanning

Sodium chromealum Na<sub>2</sub>SO<sub>4</sub> Cr<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>,24 H<sub>2</sub>O corresponds to the potassium salt, but is more soluble in water

### Ammonium chrome alum

is prepared like the preceding alums. It is less soluble in water than potassium chrome alum. It melts at  $100^{\circ}$ 

Chromium chlorate  $\operatorname{Cr}_2(\operatorname{ClO}_3)_3$  is used to some extent in calico printing (Despierres and others, Dingl poly J 228, 260-263), and for increasing the permanence of aniline black may be prepared by the addition of a solution of potassium chlorate to chiome alum or by the addition of barium chlorate to a solution of chromium sulphate, the barium sulphate being precipitated According to Lauber and Weinreb (Dingl poly J 257, 290), it is best prepared by the following process: Dissolve 60 parts of chrome alum in 80 parts (by weight) of hot water; add 20 parts of soda dissolved in 60 parts of water, and wash the precipitated hy-Redissolve the precipitate in 10 parts of cold sulphuric acid of 66°B, filter if necessary and add 22 parts of potassium chlorate dissolved in 50 parts of water The solution is evaporated slightly and the potassium sulphate crystallised out; the more soluble chromium chlorate remains in solution. When heated at 100° it gives off quantities of chlorine and forms a yellow-red liquid containing chromic acid and other compounds.

Chromium perchlorate is obtained by dissolving the hydroxide in perchloric acid and evaporating the solution. It forms short green deliquescent needles and is very soluble in alcohol (Bull. Soc. chim. 1913, 103). Two

perchlorates are known [Cr(H<sub>2</sub>O)<sub>6</sub>](ClO<sub>4</sub>)<sub>3</sub> and [Cr(H<sub>2</sub>O)<sub>6</sub>](ClO<sub>4</sub>)<sub>3</sub>,3H<sub>2</sub>O Both salts are green No sodium-chromi-perchlorate exists,

Chromic anhydride, Chromic acid Cr()3 This important oxide is best prepared by the decomposition of a chromate by sulphuric acid. A cold saturated solution of potassium chromate is added to from 1 to 11 parts of concentrated sulphuric acid, with constant stirring. On cooling, if the acid used be strong enough, crimson crystals of chromic anhydride separate, if necessary, the solution is evaporated until the crystals are formed on cooling. The liquid is poured off and the crystals drained on porous tiles or pressed between porous plates They may be recrystallised by dissolving in water and evapo-The excess of sulphure acid rating to a syrup may be removed by the cautious addition of barium chromate to the solution of the crystals, or the crystals may be gently fused, when the heavier oxide sinks to the bottom of the vessel and the sulphuric acid above slowly evaporates. The mother liquors from the chromic anhydride are used for a fresh preparation until the percentage of potassium sulphate becomes excessive, they are then precipitated by the addition of lead acetate with the production of a light-coloured 'chrome yellow' containing much lead sulphate

According to Zettnow (Pogg Ann 143, 468), the largest yield is obtained as follows: 300 grams of potassium dichromate are dissolved in 500 c c of water, and 400 c c of strong sulphure acid are added. The mixture is allowed to stand for 10 or 12 hours, the solution decanted from the deposited crystals of potassium sulphate, and heated to 80° or 90°, 150 c c of sulphuric acid are then added, followed by sufficient water to redissolve the chromic anhydride at first deposited, and the solution evaporated until the The mixture oxide commences to crystallise is allowed to stand for 12 hours, decanted from the crystals, and evaporated with the production of a second and third crop of crystals. The crystals are drained, introduced into a crucible, mixed thoroughly with 50 cc of nitric acid of not less than 1 46 sp gr (weaker acid would dissolve the crystals) and drained for some hours The operation is repeated if necessary e c of nitric acid. The whole of the on a tile with 25 e.e. of nitric acid sulphuric acid and sulphate having been thus removed, the crystals are heated on a sand-bath and the nitric acid evaporated, or they may be heated to 80° in a current of air In this manner, Zettnow obtained from 85 to 90 pc of the

theoretical yield of pure dry chromic anhydride Duvilliers (Compt rend 75, 711) prepares this oxide by dissolving 100 parts of barium chromate in 100 parts of water and 140 parts of nitric acid of sp gr 1 38 (40°B.) and heating the solution until it becomes red 200 parts of water are then added, and the whole boiled for 10 minutes; cooled, and the barium nitrate allowed to crystallise out The solution is decanted, evaporated to the bulk occupied by the original amount of acid used, and a further crop of crystals of barium nitrate removed The solution, which contains only about I part of barium nitrate for 200 parts of the anhydride, is evaporated nearly to dryness, a little water added, and the evaporation repeated until all the nitric acid is driven off. The anhydride is then crystallised out and dried as usual.

240 CHROMIUM.

Barium chromate is digested for some time with about half the amount of dilute sulphuric acid required for complete precipitation, with the production of chromic anhydride and barium dichromate and precipitation of barium sulphate. Sufficient sulphuric acid is then added to just precipitate the barium, and the solution, which only contains chromic anhydride, is decanted from the precipitate and the solution evaporated (Meissner). According to C Watts, the barium chromate is best treated with excess of strong intre acid, the precipitated chromic oxide filtered through asbestos, and the intricacid driven off by heat

Chromic anhydride is a powerfully acid substance crystallising in scarlet rhombic prismatic needles of sp gr 2 788 (Zettnow). They melt at 193° without decomposition, and the melted mass exhibits supercooling; at higher temperatures oxygen is evolved together with red vapours.

probably CrO.

When heated to 250° they decompose into chromium chromate (°rO<sub>3</sub>·(°r<sub>2</sub>O<sub>3</sub>(=3CrO<sub>2</sub>, also known as chromium dioxide) with evolution of oxygen Above that temperature, the whole is converted into chromium sesquioxide. The perfectly dry anhydride, when heated to 120° in vacuo, slowly volatilises and condenses in red needles (Arctowski) By the action of light on a solution of this oxide, oxygen is evolved and chromium chromate precipitated as a brown powder

Chromic anhydride is very soluble in water 100 parts of water, at 26°, dissolve 62 parts of the oxide. The aqueous solution contains  $\mathrm{Cr}_2\mathrm{O}_7$ "ions, with a small proportion of  $\mathrm{Cr}\mathrm{O}_4$ " ions These ions are capable of transformation by the action of acids or alkalis, as indicated by the equations:

(1) 
$$Cr_2O_7'' + 2HO' \ge 2CrO'' + H_2O$$
  
(2)  $2CrO_4'' + 2H \ge Cr_2O_7'' + H_2O$ 

Datta & Dhar (J Amer Chem Soc 1916, 38, 1303), from a study of the molecular solution volumes and molecular refractivities of chromic acid, potassium chromate and potassium dichromate, conclude that the formula of chromic acid is  $\rm H_2Cr_2O_7$  and not  $\rm H_2Cr_2O_7$ .

It dissolves in acetic acid or ether without decomposition, is almost insoluble in sulphuric acid of 177 sp gr, but dissolves in stronger or weaker acid, is insoluble in nitric acid of sp gr 146

Chromic anhydride is a powerful oxidising agent. Warm anhydrous alcohol inflames when dropped upon it, and even dry ammonia gas is oxidised with production of water and nitrogen Addition of oxalic acid is said to increase its oxidising power. The solution in water, sulphuric acid, or acetic acid, is largely used as an oxidising agent in organic work In most of these oxidations, precautions must be taken to moderate the velocity of the reaction acetic acid is used as the solvent, the substance to be oxidised is usually dissolved in the same solvent and the oxidising solution slowly added, the temperature being kept sufficiently low, i e seldom above 100° Chromic acid, as an oxidising agent, has the advantage that the close of the reaction is frequently indicated by a pure green colour in place of the brownish colour.

It has the property of producing a permanent yellow colour with silk or wool, but not

with cotton. The chromium in such material may be converted into lead chromate upon the fibre or turther dyed by logwood, &c Chromic anhydride usually contains sulphuric acid and sulphates or nitrates, Moissan found 24 pc. of sulphuric acid in samples of the commercial article.

Chromates. All chromates are prepared, directly or indirectly, by the action of oxidising agents upon chrome iron ore or chromite FeO Cr<sub>2</sub>O<sub>3</sub>. This important mineral is widely distributed and extensively mined. Large deposits are found in New Caledonia, containing sometimes 5 pc of chromium sesquioxide in a form soluble in acid, probably as chrome ochre

Chromite belongs to the spinel group of minerals. It is isomorphous with magnetic iron ore FeO Fe<sub>2</sub>O<sub>3</sub>, and may be considered as that substance in which the sesquioxide of iron is replaced by sesquioxide of chromium. A portion of the chromium is usually replaced by aluminium, and some of the iron by magnesium, though this is sometimes absent. The composition of chromite varies extremely, an ore, containing 50 p.c. of the sesquioxide, is of very good quality. Of seven samples from varying localities analysed by Clouet, the chromium sesquioxide present averaged. 44 p.c. For analyses v. Chromite.

Chromite is black or brownish-black, and infusible before the blowpipe. It will scratch glass, but is itself scratched by a good steel kinfe.

A scheme for the analysis of chromite has been given under the estimation of chromium

The chief use of chromite is as a refractory lining in steel or copper furnaces. Smaller amounts are employed in the manufacture of chrome-steel alloys and of chromium compounds

Potassium chromates. Both the normal and acid chromates are of importance in the arts. The acid or dichromate, containing the larger proportions of the active constituent CrO<sub>3</sub>, is generally prepared on the large scale.

A great number of processes have been adopted for the treatment of chromite for this purpose, they all depend upon the oxidation of the chromium sesquioxide into chromic anhydride

The ore is heated to redness and plunged into cold water to facilitate the powdering, stamped, ground in a mill, and finely sifted

In former times, the ore was mixed with about two-fifths of its weight of potassium nitrate and ignited on the bed of a reverberatory furnace, the oxygen required for the oxidation of the sesquioxide into the anhydride and the potassium for its conversion into potassium chromate, being both provided by the nitrate A great saving was introduced by the substitution of atmospheric oxygen as the oxidising agent in place of the nitre, the potassium being supplied by potassium carbonate or, as proposed by Watt, by potassium sulphate The latter salt is largely used where, as in Russia, the demand for potassium sulphate is small, and hence this salt, produced during the later conversion of the chromate to dichromate, is used to replace some of the potassium carbonate. The mass, however, was very liable to fuse and the heavy particles of chromite to sink through the liquid and escape the action of the blast This difficulty was overcome by Stromeyer, who introduced lime into the mixture, thus rendering the mass porous and non-fusible and much increasing the rapidity and completeness of the action.

Tilghmann (Patent 1847) makes use of felspar as the source of alkalı. For this purpose, I part of chrome ore is mixed with 4 parts of felspar and 4 parts of lime or 7 parts of chalk. The mixture is heated to bright redness, but not to fusion, in a reverberatory furnace for about 20 hours, with constant stirring. The chromate thus produced is removed as described later.

The ore may also be mixed with potassium chloride, and heated to whiteness with frequent stirring while a current of superheated steam is passed over the mass; the iron is stated to pass away almost entirely as chloride (Swindells,

Chem. Gazz 1851, 419).

Jacquelain (Ann. Chim. Phys [3] 21, 478) prefers to produce calcium chromate in the furnace and to convert it subsequently into the required chromate by double decomposition. He thus avoids the loss of alkali always experienced in the ordinary process. The ore is finely powdered, preferably levigated and diled before use, mixed ultimately with chalk in the required proportions, and heated to bright redness on the bed of a reverberatory furnace for 9 or 10 hours, with occasional stirring (at least once an hour). The product is ground in a mill, suspended in hot water, and treated with sulphune acid until slightly acid, calcium dichromate is thus produced A little chalk is stirred into the solution to precipitate the iron as oxide, and a solution of potassium carbonate or other salt, according to the chromate icquired, is added. The calcium is thus precipitated as carbonate and the dichromate required remains in solution The solution may also be used directly for the production of chrome yellow, &c.

According to Booth's method (Pat 1852, Dingl poly J 131, 137), the ore is mixed with one-fifth of its weight of charcoal or coal and ignited in a reverberatory furnace like that used for puddling iron, out of contact with the air The product is thrown into vats, and the reduced iron dissolved out by dilute sulphune acid, the residue being treated in the usual manner for the production of chromates The weight of oil of vitual used, amounts to about two-thirds that of the ore It is stated that the value of the ferrous sulphate obtained repays the extra labour

involved

Atcherley (Hoffman's Ber u Entw. Chem. Ind [1] 725) gives a description of the manufacture of potassium dichromate as usually practised on the large scale, of which the followıng ıs an abstract

The ore is stamped, ground in a mill, and passed through a sieve of 80 meshes to the

Potash lye is prepared by soaking 7 cwts. of burned limestone (containing but little silica and magnesia) in a solution of 21 p.c. of potassium carbonate free from chloride (which would hinder the crystallisation of the product), dried, ignited,

and powdered

It is then mixed with 41 cwts of the ore, and spread in a layer 2 inches deep upon the previously heated bed of a reverberatory furnace, at the end nearest the fire. The furnace is about 9 feet long and 6 feet broad. The heat is maintained at bright redness, and the mixture turned over every 15 minutes; in about 2 hours it is gradually pushed towards the farther end

of the furnace, and its place supplied by a fresh charge. The operation is completed in about 4 hours, when the mass is greenish-yellow and contains chromates of potassium and calcium, lime, potassium silicate, and ferric oxide It is broken into coarse powder and lixiviated in wooden pans, placed steplike one above another, with a hot solution of potassium sulphate, which converts the calcium chromate into potassium chromate with precipitation of calcium sulphate. The liquid is allowed to run from one vat to another, until at last a saturated solution of potassium chromate is produced This is run into a depositing vat lined with lead and treated with sufficient sulphuric acid and water to convert the normal salt into the dichromate, and the solution cooled.

As the solution of normal chromate was saturated and contained about 1 part of salt in 2 parts of solution, the dichromate solution formed is supersaturated, that salt requiring nearly five times as much water for solution as the normal chromate, about three-quarters of the total amount in solution is thus deposited on cooling. The crystals are redissolved in sufficient water to form a solution of 1:120 sp gr. and crystallised in deep non pans at a temperature of 21°.

The production of chromates is said to be rendered easier by the addition of oxygencarriers, such as manganates or oxides of copper or iron (D R P 163814). A method dependent on this, in which ferric oxide obtained by heating pyrites is used, is the following: 240 parts chiome non ore of 50 p.c., 170 parts sodium carbonate, and 150 parts ferric oxide, with, if desired, 50 to 80 parts lime, are intimately mixed, roasted, and lixiviated. The residue contains all the original ferric oxide and may be used over again. The solution is concentrated and treated as usual The action of the ferric oxide is regarded as due to the production of an alkaline feirite which becomes converted to the chromate and ferric oxide (J. Soc. Chem. Ind. 1895, 865)

Chromates are also produced by mixing finely divided chrome iron ore with alkaline carbonates and roasting in an oxidising atmosphere in a revolving furnace at a temperature sufficient to keep the mass fluid (National Electrolytic ('o New York, J Soc Chem Ind. 1908)

Potassium chromate K2C1O4 is prepared, as already described, from chrome ore It may be produced by adding potassium hydroxide to a solution of the dichromate. It crystallises in anhydrous yellow rhombic prisms isomorphous with those of potassium sulphate, sp gr 2 71 (Kopp) 100 parts of water dissolve of potassium chromate

105.8° being the boiling-point of the saturated The solution has a fine yellow colour of great tinctorial power; a distinct yellow tinge is imparted by 1 part in 100,000 parts of water. It is insoluble in alcohol and ether, and is very poisonous, producing a condition resembling diabetes when administered by mouth or subcutaneously It also produces slow-healing sores, especially among workers in chromate factories. For these, a sodium bisulphite wash is efficacious. Such workers should pay attention to constant washing of face, nose, and hands, and packers are advised to wear respirators

When heated, it reddens and fuses without decomposition; heated with reducing agents, such as sulphur, it is reduced to chromium sesquioxide. The addition of an acid, even carbonic acid, determines the decomposition into potassium dichromate, as is indicated in the previous equations, and it is evident further that, even when excess of acid is present, the solution still contains CrO47 ions. The equations also indicate that a solution of potassium chromate possesses an alkaline reaction.

Owing to the much smaller solubility of the dichromate (4.64 parts at 0°), when the chromate is crystallised some dichromate also separates

It forms double salts with the chromates of

the alkaline earths and lead

Potassium chromate is not largely used, the dichromate, on account of its greater richness in chromic acid, being generally preferred. The general uses and oxidising properties, however, described under potassium dichromate, also

apply to the normal chromate

Potassium dichromate, generally known as Bichromate, or 'Bichrome,' K<sub>2</sub>(1<sub>2</sub>O<sub>7</sub>. This important salt is prepared in large quantities by the methods already described. It may be made electrolytically by the use of a solution of potassium hydroxide, a cathode of porous copper oxide, and an anode of ferrochrome (50 pc), which has an E M F. of 2 volts ferric oxide is also formed and precipitated (Lorenz, Zeitschanoig. Chem 1896, 12, 396). Considerably over 10,000 tons are annually produced in Great Britain.

It crystallises in anhydrous, fine, red, square tables, derived from triclinic prisms, of sp gr 2 692, and of intensely metallic, bitter taste. They melt at about  $400^{\circ}$  (Tilden and Shenstone), and resolidify on cooling without change. At bright redness, decomposition ensues, and the normal chromate and chromic sesquioxide result. Other chromates of composition,  $K_2Cr_4O_{10}$  and  $K_2Cr_4O_{11}$ , appear to exist in equilibrium with the solution with certain limits of concentration.

100 parts of water dissolve at 0°, 4 64 parts, at 30°, 18·13 parts, at 60°, 45·44 parts and at 104 8°, 108 2 parts, 104 8° being the boiling-point of the saturated solution (Koppel and

Blumenthal, lc)

Potassium dichromate is used in the preparation of all chrome pigments, of 'discharge' for Turkey red, &c, for the production of a large variety of colours in calico printing and dyeing, in the manufacture of safety matches, &c. In solution with sulphuric acid, it is used as a bleaching agent for tallow, palm oil, &c; in the oxidation of anthracene to alizarin, and in the manufacture of aniline violet and in many other organic oxidations. It is also used for tanning leather (v. Leather).

When mixed with organic substances it is reduced on exposure to light; gelatine under such circumstances is rendered insoluble. This reaction is taken advantage of in the 'Carbon' process of photography. The gelatine is mixed with a pigment of any colour,

and the paper carrying this film is sensitised by floating on a solution of potassium dichiomate On exposure under a negative, the gelatine in those portions exposed to the light becomes insoluble, and retains the pigment, while the portions protected by the darker parts of the negative are almost unacted upon, and may be dissolved in warm water. In this manner, photographs of any desired colour, and of great beauty and permanence, may be pro-Contact of the chromated gelatine with the reduced silver of an ordinary biomide print has a similar effect, and bromide photographs may hence be copied by squeezing into contact with a sensitised pigmented gelatine After a few minutes' contact, the pigmented paper is developed like a carbon print The bromide has become bleached and reduced to silver, but can be redeveloped. The insoluble gelatine does not absorb water, but will take oil, which may be mixed with any desired pigment, and this forms the basis of the cil and bromoil processes Gum. mixed with dichromate, is also made insoluble by exposure to light, and this is made use of in photography in the gum dichromate process (v PHOTOGRAPHY).

Sodium chromates. The normal chromate Na<sub>2</sub>(r<sub>2</sub>O<sub>4</sub>, 10H<sub>2</sub>O and the dichromate Na<sub>2</sub>(r<sub>2</sub>O<sub>7</sub>, 2H<sub>2</sub>O are prepared by methods corresponding to those used for the potassium salts. According to Nic. Walberg (Dingl. poly. J. 259, 188), they are prepared on the large scale as follows: 6 parts of powdered chrome ore (44 p.c. (1<sub>2</sub>O<sub>3</sub>) are mixed with 3 parts soda ash (92 p.c. Na<sub>2</sub>CO<sub>3</sub>) and 3 parts of chalk, and heated in charges of 1 ton in a reverberatory furnace for 8 hours. The mass is himited to produce a solution of 45°B, boiled down to 52°B, and allowed to crystallise in leaden pans. The crystals are first dired by a centifugal machine, and finally heated to 30° m. a drying chamber, where they crumble to a yellow anhydrous powder containing about 96 p.e. of the normal chromate

For the production of dichromate, these crystals are dissolved to a solution of 40°B and treated with sufficient chamber acid to determine the conversion into the dichromate, the right point being found by potassium iodide and starch paper Sufficient neutral sodium chromate is then added to bring the percentage of CrO, to about 72.5. The liquor is cooled in lead tanks about 72 5 to 1°, artificially if necessary, and the sodium The liquor is drained sulphate crystallised out off, filtered if necessary, and evaporated to dryoff, filtered it necessary, and the result of the ness in an iron pot, with constant stirring. The ness in an iron pot, while still hot. If the amount of CrO, be allowed to exceed 72 5 p c, the product is damp and cannot be stored in wooden casks An analysis of this substance showed CrO<sub>3</sub>, 72 3, Na<sub>2</sub>O, 26·20, SO<sub>3</sub>,

Newer methods of manufacture are the treatment of the monochromate with ammonia, followed by carbonation and deposition of sodium bicarbonate, the use of caustic alkali in the place of alkali carbonate; and the employment of only half the necessary sodium carbonate required to decompose the ore, the remainder being replaced by sodium sulphate, formed in the later stages of manufacture (Ulm, Chem. Zeit. 1914, 38, 670; J. Soc. Chem. Ind. 1914, 33, 947).

CHROMIUM.

Normal sodium chromate forms fine large fflorescent crystals of the composition

### Na,CrO4,10H2O

somorphous with Glauber s salt Unlike potasium chromate, it is less soluble than the hehromate, it is, however, more soluble than iotassium chiomate, 100 parts of water at 30° hs solving  $87\ 36$  parts of sodium chromate

Other hydrates with 6 and 4 mols of H,O re known. Their transition temperatures have peen suggested as convenient fixed points in hermometry (Richards & Kelley, J Amer

Jhem Soc 1911, 33, 847)

Sodium dichromate Na<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>,2H<sub>2</sub>O crystalises in thin, six-sided prisms with bevelled edges of a fine hyacmth-red colour, and is very oluble in water, 100 parts of water at 30° dis-olving 197 6 parts of the salt—It is hence more han 10 times as soluble as potassium dichromate which frequently makes its use preferable to that of the potassium salt

Ammonium chromate and dichromate are prepared by mixing in proper proportions solutions of ammonia and of chromic acid chromate, on heating, yields nitrogen, water, and chromic oxide, and forms a convenient method for the preparation of nitrogen free from argon

Lead chromates. The neutral lead chromate PbCrO<sub>4</sub> is best produced by the action of a solution of lead acetate on a solution of a chromate Thus produced, it is of a fine lemonyellow colour, insoluble in water and dilute acids When heated at 250°, it becomes reddishbrown, at a higher temperature it fuses, and finally evolves oxygen with the formation of chromic sesquioxide and a basic lead chromate On account of this evolution of oxygen on heating, this substance is frequently used in organic analysis.

Lead chromate is soluble in cold lime water or caustic alkali, but is unattacked by hot lime water For this reason, in immersing calico coloured with chrome yellow in lime water, for the production of an orange colour, care should

be taken that the solution is hot

When heated with caustic alkali, a portion of the chromic oxide is removed, with the production of a basic lead chromate of colour

varying from orange to vermilion

The normal chromate is largely used as a pigment, and for calico printing, and is the essential constituent of various chrome pigments as Chrome yellow, Paris yellow, Leipsic yellow The name Cologne yellow was originally given to chrome yellow containing a mixture of lead chromate and lead sulphate, but the latter compound is usually present in pale yellows known commercially as 'puie' chrome yellows Chrome yellows. The finest chrome yellow

is produced by precipitation, the lead acetate used should not be basic, or the product will be of an orange colour; the solution should be dilute, and the lead salt should be in excess, the presence of excess of chromate is liable to 'tuin' the yellow-ie to produce an orange tinge The substances used should be comparatively

For the lighter yellows, the solution of chromate is mixed with an appropriate quantity of sulphuric acid, Glauber's salts or alum (American chrome yellow), before precipitation;

the latter yields good colours but considerable care is required in the precipitation. The lead chromate is thus precipitated, together with any desired amount of sulphate, and the colour correspondingly lowered. For cheaper chromes, the colour is toned down by the admixture of kaolin, barytes, gypsum or whiting, the finely powdered solid being ground with water and well mixed with water in the tank into which the precipitating liquids are iun (vide infia) Bouthron Chatard gives the composition of ordinary Cologne yellow as CaSO<sub>4</sub>, 60 pc, PbSO<sub>4</sub>, 15 p c , PbCrO<sub>4</sub>, 25 p c.

Biot and Delisse first precipitate lead sulphate from a solution of the acetate, and digest 3 parts of the washed precipitate with 1 part of normal potassium chromate dissolved in hot The chromate is thus wholly converted into chrome yellow, mixed or perhaps combined with the lead sulphate, and a 'Cologne yellow' of good colour is produced. The colour is said to be equal to that obtained when double the ing power is, however, stated to be inferior to

that of the precipitated yellow

Milbauer & Kohn (Zeitsch, physikal Chem 1916, 91, 410) show that the feaction PbSO<sub>4</sub>  $+K_2SO_4 \rightleftharpoons PbCrO_4 + K_2SO_4$  proceeds practieally completely in the direction of left to right From determination of the stability it is shown that the solid phase PbSO4 cannot exist in the presence of K,C1O4K,SO4,PbC1O4, and that the existence of double salts is unlikely general results show that the manufacture of chrome yellows can be carried out from lead sulphate and that various shades can be obtained by varying conditions, temperature, concen-It follows also that an excess tration and time of lead acetate should be used in the manufacture since this will secure the absence of K, CrO, which would otherwise rapidly convert the lead sulphate necessary for the particular shade into chromate

Liebig uses for this purpose the lead sulphate obtained in dye works, &c, as a by-product The pigment, on account of the basic character of the sulphate, is of an orange colour

Anthon substitutes lead chloride for sulphate, treating 100 parts of freshly precipitated chloride with 27 parts of dichromate.

The method usually adopted for production of chrome vellow is the following The two precipitating solutions are placed separately in wooden tanks, and heated by steam pipes The solutions are either filtered or allowed to settle in the tanks and drawn off from above. They are run together into the precipitating tank, which contains about 200 gallons chrome yellow is allowed to settle and the clear liquid drawn off, more water is added, the whole thoroughly stirred and the process is repeated three or four times. The pigment is then put in a filter press and dried at as low a temperature as possible

According to the Textile Colourist, the preparation of chrome yellow is as follows. For the preparation of the lead acetate, four wooden tubs, 3 feet in diameter and 11 feet high, are arranged one above another, so that the liquid may pass downwards from one to the other. They are filled with granulated lead, and strong wood vinegar poured on the uppermost; after 244 CHROMIUM

remaining there for a few minutes, it is run into the second, third, and fourth for an equal time. The surface of the lead is thus thoroughly acted upon, and becomes heated and oxidised by the air. The vinegar is then passed through a second time, remaining in each tub for an hour, on leaving the fourth tub, a saturated solution of acetate is produced.

Bichrome is dissolved in 10 or 12 parts of water in steam-cased copper pans, and sufficient acetate added to just precipitate the chromate completely. The precipitate is settled, washed by decantation, filtered, and dried slowly on boards The filtration should be performed as rapidly as possible, because the pigment swells considerably while drying, and this should take place on the boards

By mixing some lead chromate with the lead acetate solution, a dark lemon colour is produced, and by the addition of caustic potash to the precipitate, a much redder pigment may be obtained

The chrome yellows, by mixture with Prussian blue, form greens known as 'Brunswick greens' and also as 'chrome greens' (not the true chrome green, vide supra) These are valuable and cheap pigments, but not as stable as the corresponding zinc greens (vide infra)

By removal of some of the acid from normal lead chromate, a basic lead chromate of a deep red colour is obtained, PbO PbCiO4, which is the base of a number of chrome reds. Between these and the various yellows a large number of shades of chrome orange may be obtained chrome reds are known by a variety of names, e g Austrian cinnabar, Persian red, Chinese red, Victoria red, Derby red, American vermilion, &c

The pigment is usually prepared by boiling chrome yellow with caustic alkali or by boiling together the lead salt, potassium dichromate, and an alkalı One such method is-20 parts of white lead are mixed into a fine paste with water, 10 parts of potassium dichromate and 3 of caustic soda, are dissolved in a little water, added, and the whole boiled until the desired colour is obtained It is then washed and driea

Runge mixes 45 parts of litharge, 6 parts of salt, and 50 parts of water, the mass becomes white, and swells considerably, more water is added to prevent it from hardening. In three or four days, the whole has been converted into lead oxychloride, a solution of 15 parts of potassium dichromate is then poured on, and the mass well stirred and washed A corresponding process is used with carbonate of lead, the colour produced being preferable to that when chloride, sulphate, or oxide of lead are used

For the production of a fine 'Persian red,' Prinvolt digests 25 parts of lead carbonate in a cold solution of 10 parts of potassium chromate for 2 days. A red crystalline precipitate of basic lead chromate is thus produced, while potassium bicarbonate remains in solution The mixture is boiled for half an hour, whereby a portion of the red precipitate becomes decomposed, with the formation of normal lead chromate and of potassium chromate, turning a violet-red colour. This is filtered off, washed, and digested with one part of sulphuric acid dissolved in 100 parts of water, with the production of 'Persian red.'

For the production of a fine vermilion pig-

ment, Liebig and Wohler's method may be used A mixture of equal parts of potassium and sodium nitrates is just heated to fusion, and chrome yellow, in small fragments, is dropped in, ebullition takes place, and the mass becomes black The heating should be continued until ebullition ceases. The chrome yellow has thus become converted into basic lead chromate and potassium chromate. If the temperature used be too high, the product becomes brown and is spoiled. The supernatant yellow liquid is poured from the heavy basic lead salt, which is then well washed in water. The washings should be poured off as quickly as possible, or the action is liable to be reversed and the normal chromate reproduced in part, with a corresponding reduction of colour.

The colour of the varieties of chrome red is due principally to the size of the crystals, those pigments of which the crystals are largest being

of the most intense colour.

For the production of chrome orange, the chrome vellow is boiled with lime or alkalı (in smaller quantity than that used for production of chrome red), until the shade required results. The cheaper chrome orange pigments, as in the case of the yellows, contain admixed kaolin,

gypsum, &c
The lead chromates all possess good covering power and brightness of colour; they are durable, but blacken under the action of sulphuretted hydrogen. They mix well with other pigments, but should not be used with

sulphides

Calcium chromate CaCrO4,4H2O is largely produced in the manufacture of chromates from chrome non ore It is soluble in water and It becomes anhydrous at 200°, and is then almost insoluble in water

Barium chromate Ba('rO4 is a canary-yellow powder, known as Yellow ultramarine or Lemon yellow It is produced by the addition of a chromate to a solution of a salt of banum.

Barium chiomate is insoluble in water and acetic acid, soluble in hydrochloric and nitric On treatment with a boiling solution of chromium anhydride, it dissolves, and crystallises on cooling as a yellowish-red powder, consisting of the dichromate BaCr<sub>2</sub>O<sub>7</sub>, 2H<sub>2</sub>O.

Both barium and calcium chromates have been employed as pigments, but are now little used on account of their lack of brightness and

covering power

Bismuth chromates. The addition of potassium dichromate to a neutral solution of bismuth nitiate gives a yellow flocculent precipitate, 3B12O3 2CrO3, soluble in acids (Lowe) heating, it becomes partly decomposed and turns greenish. From an acid solution of bismuth nitrate, the precipitate formed consists of Bi<sub>2</sub>O<sub>3</sub>·2CrO<sub>3</sub> (Lowe, Muir), and is quite insoluble in water. Bismuth chromate is used as a pigment and in calico printing

Zinc chromate, Zinc yellow, or Buttercup yellow ZnCrO4, is a beautiful, stable, yellow pigment, produced by the addition of hot neutral solution of zinc sulphate to potassium chromate, or by adding a cream of zinc oxide in water to a boiling solution of potassium dichromate.

Zinc chromate can be obtained in various tints and has a good colour. Though not equal to chrome yellow in covering power or body, it

has the advantage of not blackening by exposure to sulphides. It is largely used for mixing with Prussian blue to form the zinc greens, which are more stable and faster to light

than the corresponding chrome greens.

Perchromic acid. On adding hydrogen peroxide to an aqueous solution of chromic acid. or to an aqueous solution of a chromate acidified with sulphuric acid, a deep blue colouration is produced, which on shaking the solution with ether passes into that liquid. The colour of this substance is so intense that it constitutes a delicate test for both chromic acid and hydrogen peroxide. It is extremely unstable, evolves oxygen, and is gradually transformed into chromium trioxide. It has been regarded as Cr<sub>2</sub>O<sub>7</sub>xH<sub>2</sub>O (Barreswil), CrO<sub>4</sub>3H<sub>2</sub>O (Fairley), CrO<sub>3</sub>,H<sub>2</sub>O<sub>2</sub> (Moissan), and 2HCrO<sub>4</sub>,H<sub>2</sub>O<sub>2</sub> (Berthelot) By adding pyridine, quinoline, and other organic bases to the ethereal solution of the blue compound, Wiede obtained salts of blue salts (NH<sub>4</sub>)C<sub>1</sub>O<sub>5</sub>,H<sub>2</sub>O<sub>2</sub> and KCrO<sub>5</sub>,H<sub>2</sub>O<sub>2</sub> According to Riesenfeld (Ber 1905, 38, 4068, 1908, 41, 2826, 1911, 44, 147), these salts are acid salts of perchromic acid H3C1O8, of the type MH<sub>2</sub>CrO<sub>s</sub>, corresponding with the red per-chromates. Pure perchromic acid H<sub>3</sub>CrO<sub>s</sub>,2H<sub>2</sub>O may be obtained by the interaction of chromic trioxide and 97 pc hydrogen peroxide in methyl ether solution at -30°.

$$2CrO_3 + 7H_2O_2 = 2H_3CrO_8 + 4H_2O$$

After the reaction is complete, the blue solution is poured off, dried with phosphoric oxide, and evaporated in vacuo at -30° It forms a dark blue crystalline mass, which decomposes at above -30°. The water is regarded as water of constitution, since the blue colour is only produced in presence of water The red perchromates are considered as the anhydro salts of the blue perchromic acid (Riesenfeld and Mau, Ber. 1914, 47, 548)

Esters of chromic acid have been prepared by Wienhaus (Ber 1914, 47, 332) by shaking a tertiary alcohol, dissolved in light petroleum or carbon tetrachloride with an excess of solid chromium trioxide. The ester remains dissolved or suspended in the solvent, from which

it can be obtained by evaporation

### CHROMAMMONIUM COMPOUNDS AND DERIVATIVES

Salts of chromium readily combine with ammonia or substituted ammonias, to form complex compounds in which the chromium atom may be combined with six molecules of ammonia, as, for example, in hexamine chromium chloride  $[Cr(NH_3)_6]Cl_3$ . According to Werner's theory, the chlorine of this compound is united by primary valencies, whereas the ammonia is co-ordinated with the chromium atom by supplementary valencies (Ber 40, 20) The ammonia may be substituted by acid groups, the valency and character of the complex radical being determined by the acid and basic radicals in the molecule, as will be evident from the following description of some of the simpler compounds.

Hexamine chromium salts (luteo salts), e g. the nitrate [Cr(NH<sub>3</sub>)<sub>6</sub>](NO<sub>3</sub>)<sub>3</sub>, are prepared by treating the corresponding purpureo salts with

ammonia in cold concentrated solution (Jorgensen, J. pr. Chem. 23, 229; Mills, Phil. Mag.

[4] 35, 245)

Pentamine chromium salts (purpureo salts) Jorgensen (J. pr Chem 2, 20, 105) prepared the chloride [CrCl(NH<sub>3</sub>), ]Cl<sub>2</sub> by reducing violet chromic chloride in a stream of pure dry hydrogen at a red heat, and adding to it a solution of NH<sub>4</sub>Cl in strong ammonia. Air was passed through, HCl added, and the mixture boiled. when a carmine coloured powder is precipitated. It crystallises in octahedra, sp gr 1 687, dissolves in 154 parts of water at 16°. Representing the compound Cl<sub>2</sub>Cr<sub>2</sub>(NH<sub>3</sub>)<sub>10</sub>Cl<sub>4</sub>, Jorgensen found that only 4 atoms of chlorine are precipitated by silver nitrate, which confirms Werner's formula above given.

Triamine chromium salts. These compounds have been investigated by Werner (Ber 1910, 43, 2286), who uses triamine chromium tetroxide in their preparation To a well-cooled solution of 30 grams of chromic acid in 300 c c of water are added 300 c c of pyridine. After remaining half an hour m a freezing mixture, 750 c c 3 p c  $H_2O_2$  are added. The precipitated pyridine perchromate is collected, washed, and added to 90 c c of well-cooled 25 p c ammonia The precipitated triamine chromium tetroxide is collected after 10 minutes and washed with water, alcohol, and ether

Trichlorotriamine chromium  $[(NH_3)_3CrCl_3]$ results when triamine chronium tetroxide is added to cold concentrated hydrochloric acid A blush-grey precipitate is formed, and the filtrate from this deposits the compound after standing two days, in dark blue crystals with a greenish tinge The triaquo derivative

 $[\mathrm{Cr}(\mathrm{OH_2})_3(\mathrm{NH_3})_3]\mathrm{Cl_3}$ 

is obtained by treating one gram of the dichloroaquotriamine chromium chloride (contained in the grey precipitate above mentioned) with 8 c c of water covered with 5 c c of pyridine The resulting solution is filtered, and the hydroxoiodide precipitated by the addition of solid potassium iodide The chloride is obtained from this by triturating with hydrochloric It forms brownish-red hygroscopic crysaeid tals The corresponding bromine compounds have been prepared

The diamine chromium compounds were among the earliest known of these complexes Morland (Chem Soc Trans 13, 252) obtained a compound by the action of K2Cr2O7 on fused KCNS, which was subsequently examined by Reinecke (Annalen, 126, 113) and by Norden-skill (Zeitsch anorg Chem 1, 126), who prepared it by heating NH4CNS to incipient fusion and dropping in one-third of its weight of (NH<sub>4</sub>)<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>. The compound

 $[Cr(NH_3)_2(SCN)_4]NH_4$ 

is known as Reinecke's salt, and according to Werner is the ammonium salt of tetrathiocyanodiamine chromite, the complex molecule having an acidcharacter and being mono-valent It forms small red crystals fairly stable towards acids The potassium salt is obtained by adding KCl to its solution, and the acid [Cr(NH<sub>3</sub>)<sub>2</sub>(SCN)<sub>4</sub>]H by decomposing the barium salt with sulphuric acid. It commences to decompose at 80°. Werner and Klein (Ber. 35, 277), by oxidation of 'Reinecke's salt' with bromine, completely 246 CHROMIUM.

climinated the four thiocyano groups with the formation of dibromodiaquodiamino chromic

bromide  $[(NH_3)_2(\operatorname{Br}_2(H_2O))_2|\operatorname{Br}.$ 

The last member of the series in which all the ammonia groups are replaced by acid radicals may be represented by the salt [Cr(CN<sub>6</sub>)]K<sub>3</sub> potassium chromcyanide or hexacyanochromite It is obtained in light yellow monoclinic crystals by adding a solution of chromic acetate to a boiling solution of potassium

CHROMOCYANINES v OXAZINE COLOURING MATTERS.

CHROMOFORM. Trade name for an orangered crystalline powder formed by the union of dichromic acid and hexamethylenetetiamine (C<sub>6</sub>H<sub>1,</sub>N<sub>4</sub>CH<sub>1</sub>), Cr<sub>2</sub>O<sub>7</sub> Slightly soluble in alcohol and cold water, readily soluble in hot water. The aqueous solution on boiling, or by treatment with acids or alkalis, evolves formaldehyde

CHROMOGEN AND CHROMOPHORE v.COLOUR AND CHEMICAL CONSTITUTION

CHROMOMETERS v COLORIMETERS

CHROMONE. Colouring matters derived from chromone or pheno  $\gamma$ -pyrone (1), the analogue of coumarın (2)

or pheno α-pyrone, have not as yet been found in nature, though an hydroxy chromone (1) analogous to daphnetin (2) will no doubt possess dyeing properties and yield a yellow shade on aluminium mordant

Chromone is specially interesting owing to the isolation of the 3 hydroxy-chromonol

by Schall and Dialle (Ber 1881, 21, 3009) as one of the products of the alkaline oxidation of brazilein Benzo γ-pyrone derivatives were first synthesised by v Kostanecki (Bei. 1900, 1998), but the synthesis of chromone itself is due to Ruhemann and Stapleton (Chem. Soc. Trans 1900, 1179) When phenoxyfumaric acid (1) is treated with sulphuric acid benzo γ-pyrone carboxylic acid is produced (2)

and this when heated in a vacuum yields benzo  $\gamma$ -pyrone:

and its yellow solution in cold sulphuric acid possesses a blue fluorescence A. G. P

CHROMOTROPE or CHROMOTROPIC ACID, 1 8-Dihydroxynaphthalene-3 6-Disulphonic acid.

CHRYSANILINE. Diamidophenylacridine C<sub>19</sub>H<sub>16</sub>N<sub>3</sub>, is obtained as a by-product in the manufacture of rosaniline by the arsenic acid method, and was isolated by Nicholson (Dingl. poly. J 168, 133) by treating the residue with steam, whereby chrysaniline passes into solution and can be precipitated therefrom by addition of nitric acid It was further examined by Hofmann (Ber. 1869, 378), who prepared its methyl-, ethyl-, and phenyl- derivatives, and showed that it possessed the empirical formula ('20H17N3 Its composition and constitution were first established by Fischer and Körner (Ber 1884, 203, Annalen, 226, 175), who showed that a second lower homologue of composition ('19H<sub>15</sub>N', was present (compare also Anschutz, Bei 1884, 433), and who refer its formation in the rosaniline melt to a condensation and subsequent oxidation of orthotoluidine and aniline, since they have effected its synthesis by heating with aisenic acid at 180°-200° the orthodipalatrianimotriphenylmethane obtained by Renouf (Ber 1883, 1304) by the action of dehydrating agents on a mixture of orthonitrobenzaldehyde and aniline (v. also Ostrogovich and Silbermann, Chem Zentr 1907, 1 1195) The commercial product is a mixture of the two derivatives, and in the form of the nitrates or hydrochlorides appears on the market as phosphine

Properties—Chrysaniline crystallises with 2 mols of H<sub>2</sub>O from 50 p c alcohol in goldenyellow needles, and from benzene with I mol of benzene of crystallisation in golden-yellow scales, the latter melt, when placed in a bath heated at 150°-160°, but when slowly heated lose the associated benzene, and melt, like the former, at 267°-270° It is insoluble in water, and sparingly soluble in alcohol. On oxidation with chronic acid, it yields acridine and when diazotised and boiled with alcohol is converted into phenylacridine (mp 164°). Tetrazo compounds have been prepared, but then couple products with amines, phenols, &c, are not as good dyes as those obtained from chrysaniline and its salts (Trillat and de Raczkowski, Compt 1end 114, 1024) Formaldehyde reacts with chrysaniline, yielding an orange dyestuff similar to chrysaniline in its tinctorial properties (T and de R Compt rend 116, 1382) Chrysophenol (19 H<sub>14</sub>N<sub>2</sub>O, a yellow dye, is obtained when it is heated at 160°-180° with concentrated hydrochloric acid, and crystallises from dilute alcohol with 2 mols. of  $\rm H_2O$  in ruby-red needles. Chrysamline forms two series of salts; of these, the sparingly soluble intrate is the most characteristic: it crystallises in ruby-red needles, and is employed under various names as a valuable golden-yellow dve for silk, wool, and leather (v ACRIDINE DYESTUFFS)

CHRYSANTHEMIN v ANTHOCYANINS.
CHRYSAROBIN (B.P., USP) is a yellow crystalline mixture of substances obtained from Avaroba or Goa powder The latter varies in CO CH colour from primrose-yellow to bronze and purple, and collects, possibly as the result of oxidation of the resin in the cavities of the

stems and branches of Andira Araroba (Aguar) (N.O. Lequminsæ), a tree inhabiting the forests of Bahia (Aguar, Pharm. J. [in.] 10, 42; cf Greenish, ibid. [in.] 10, 814) The drug was formerly exported by the Portuguese to Goa, but its origin for a long time remained secret Goa powder is usually mixed with woody fragments from which it is freed by sifting, it is sometimes termed crude chrysarobin. After extraction of Goa powder with organic solvents (benzene, chloroform) there is left behind on evaporation a yellow crystalline powder, introduced in 1875 by a Bombay firm under the name Chrysarobin, which has since been included under this name in the British and United States.

Attheld (Pharm J 1875, [3] 5, 721) obtained chrysarobin in a yield of 80-84 p c of the crude drug, using benzene Tutin and Clewer (Chem Soc. Trans 1912, 101, 290), by successive exhaustion with light petroleum, ether, chloroform, ethyl acetate, and alcohol extracted only 57 5 p c The residue consists of woody fibre, inorganic and resinous constituents, a little

sugar, &c.

Chrysarobin was originally thought to be a pure, or nearly pure, substance, in reality it is a complex mixture, consisting chiefly of reduced di- and trihydroxymethylanthraquinones, which crystallise together and are difficult to separate by purely physical means. Later the name chrysarobin was applied to one of the pure constituents, but it is better to reserve this name for the commercial mixture. Tutin and Clewer found the following percentage composition for three samples of commercial chrysarobin.—

1	Chrysophanic acid	1		
	(=chrysophanol) 1	63	$7\cdot 2$	8.9
2.	Emodin monomethylether 1	}		
3	Emodin	trace	nıl	nıl
4.	Crude anthranol of chryso-			
	phanic acid .	26	46	62
5.	Anthranol of emodin mono-			
	methyl ether	small	amou	nts
6	Monomethyl ether of de-			
	hydroemodinanthianol	41 l	13 4	177
7.	Ararobinol	4 2	4 1	nıl
8.	Inseparable mixture of			
	partly amorphous sub-			
	stances .	23	30	12

The first three of these substances are anthraquinone derivatives (for their properties and constitution, see Rhubarb) Nos 4 and 5 result from Nos. 1 and 2 respectively, by reduction, they are derivatives of anthranol (9-hydroxyanthracene). No. 6 is a less strongly reduced anthraquinone. No. 7, of unknown constitution, is also an anthraquinone derivative, related to No. 1

Chrysophanic acid anthranol C<sub>15</sub>H<sub>12</sub>O<sub>3</sub> has

$$C_6H_3(OH)$$
 $C_{CH}$ 
 $C_6H_2(OH)CH_3$ 

and was called chrysarobin by Jowett and Potter (Chem Soc Trans 1902, 81, 1578). It is obtained from commercial chrysarobin by extraction with light petroleum; by shaking

About 3 or 4 parts of the former to 1 of the latter.

its chloroform solution with sodium carbonate traces of emodin are removed, and after evaporation of the chloroform the residue is crystallised from much ethyl acetate, when it forms lemon-yellow scales, m.p. 202° (corr) (regenerated from its acetyl derivative the substance melts at 204°). It is insoluble in sodium carbonate, soluble in caustic alkalis to a yellow solution, which becomes red by passing air through it, whereby it is oxidised to chrysophanic acid. The change is complete in 2 hours, as shown by measurement of the oxygen absorbed (Tutin and Clewer). The reverse change is brought about by hydriodic acid at 130°-140°. Chrysophanic acid anthranol dissolves in sulphuric acid with a yellow colour; it forms a diacetyl compound  $C_{15}H_{10}O_3Ac_2$ , m.p. 193°, and a triacetyl compound  $C_{15}H_{9}O_3Ac_3$ , m.p. 238°.

The anthranol of cmodin monomethyl other was not isolated by Tutin and Clewer as such, but its presence was inferred from aerial oxidation experiments, which converted it into emodin monomethyl other

The monomethyl other of dehydroemodinanthranol  $C_{16}H_{12}O_4$  is characteristic of chrysarobin. According to Eder (v infra), its constitution is most probably

$$\begin{array}{c|c} CH^2O & CH^2 \\ \hline \\ HO & C \\ \hline \\ OH \\ \end{array}$$

Tutin and Clewer converted it by boiling hydriodic acid into emodin anthranol, a substance containing a methyl group less and 2 hydrogen atoms more, by oxidation with chromic acid they converted it into emodin methyl other

Of the substances 1-6 chrysophanic acid had also been found by Attfield, its anthranol by Jowett and Potter, and emodin monomethylether by Oesterle and Johann (Arch Pharm 1910, 248, 476), but this was the full extent of the agreement among the various investigators of chrysarobin until Eder (Arch. Pharm 1915, 253, 1; 1916, 254, 1) confirmed all the principal results of Tutin and Clewer He oxidised the commercial substance with air in alkaline solution, and considers that hydrogen peroxide is formed in this reaction, and is responsible for further changes. He thus obtained the following percentages. emodin 02, emodin methylether 93, chrysophanic acid 227, dehydro-emodin anthranol monomethyl ether 18 (=chrysarobol of Hesse?), amorphous products 26 It should be remembered that most of the chrysophanic acid and some of the emodin methyl ether were originally present as anthranols. In his second paper, Eder acetylated and benzoylated chrysarobin and obtained by subsequent hydrolysis up to 14 pc. of chrysophanic acid anthranol, up to 9 pc. of emodin anthranol monomethyl ether, and up to 4 pc of emodin monomethyl ether, but no chrysophanic acid The chief discrepancy is that Eder did not find the last of Tutin and Clewer's substances, ararobinol. This, according to the latter authors, has the composition  $\Omega_{23}H_{16}O_5$ , and forms yellow crystals, decomposing without melting at about

225°. It dissolves in alkali with a yellow colour, but, like the methyl ether of dehydro-emodin anthranol, and unlike chrysophanic acid anthranol, it is not readily oxidised by air. It gives a delicate and characteristic reaction with sulphuric acid, a mere trace forms an orange solution, which on gentle agitation gives intense blue streaks, the solution then becomes green, and finally dull grey. Ararobinol gives a triacetyl derivative (decomp at 225°), is reduced by hydriodic acid to dehydroararobinol, and oxidised by chromic acid to chrysophanic acid, but not quantitatively.

With the exception of chrysophanic acid anthranol, Tutin and Clewer could find none of the substances described by Jowett and Potter They had access to the latter's specimens of dichrysarobin' and its 'methylether;' the former they separated into substance 4, and emodin anthranol, the latter into the substances 4 and 5. A mixture of the latter substances is similar in composition and melting-point to 'dichrysarobin methyl ether,' which illustrates the difficulty of separating the constituents of chrysarobin by mere crystallisation

Chrysarobin is occasionally employed intern-y as a purgative Léger (J. Pharm Chim ally as a purgative 1912, [vn ] 5, 588) has suggested that for this purpose chrysarobin should be substituted for chrysophanic acid, since they both have the same purgative effect, and the latter is difficult to obtain in a state of purity. Probably the chief constituent of chrysarobin, chrysophanic acid anthranol, does not exert a purgative action until it has been oxidised in the alimentary canal (to chrysophanic acid), as is the case with aloin (Meyer and Gottlieb, Experimentelle Pharmacologic, Berlin and Vienna, 1914, 198) The chief use of chrysarobin is, however, as an ointment in skin diseases (psoriasis), and the active constituent appears to be the anthranol of chrysophanic acid According to Unna's experiments (Baudisch, Ber 1916, 49, 179), the only active synthetic substitutes are the anthranols of 1-hydroxy and of 1 8-dihydroxy anthraquinone The latter, eignolin

is more active in psoriasis than the anthranol of chrysophanic acid (chrysaiobin), because it has not the dystherapeutic methyl group

Baudisch explains the effect of the 1 8hydroxyls in accordance with Pfeiffer's views by the formula

Various other substitutes have been suggested, e.g. acetyl chrysarobin (Knoll & Co D. R. P. 105871; Chem Zentr. 1900, 1 524), 'lenirobin'; eurobin is said to be triacetylchrysophanic acid (?)

chrysophanic acid (?)
'Dermol' is prepared by precipitating a bismuth nitrate solution by a solution of

chrysophanic acid in sodium hydroxide. 'Anthrarobin' is the anthranol of alizarin.

The different methods of extraction cause considerable variation in the composition of commercial chrysarobin and of its physiological action (Pharm J 1911, [1v] 87, 630) Hence, Tutin and Clewer consider it fallacious to ask for compliance with exact requirements as to solubility, etc (The USP states that 1 gram of chrysarobin dissolves in 385 cc of alcohol, 12 5 c c. of chloroform, 16 c c. of ether, 30 cc of benzene, and 180 cc. of carbon disulphide at 25°) Pure chrysarobin dissolves in concentrated sulphuric acid with a yellow colour, and is insoluble in dilute sodium hydroxide; concentrated sodium hydroxide dissolves it with a green colour Chrysophanic acid, on the other hand, yields a red solution with concentrated sulphuric acid and dilute sodium hydroxide (Liebermann and Seidler, Ber 11, Another method to distinguish chrysarobin from chrysophanic acid is to mix 0 05 gram with 0 2-0 3 gram sodium peroxide in a porcelain capsule, add 5 c c of alcohol, and after 5 minutes 15 c c of water Chrysaiobin yields a wine-lees colour, not destroyed by dilution with water, changing to yellow on the addition of acetic Chrysophanic acid yields a cherry-red colour, becoming brighter on dilution (Alvarez, Ann Chim anal 1907, 9) G. B.

CHRYSATROPIC ACID v LACTONES CHRYSAZIN v ALIZARIN AND ALLIED COLOURING MATTERS.

This hydrocarbon occurs in the highest boiling portions of coal tar, and in the tarry products obtained in the dry distillation of wood, fats, and oils (Laurent, Ann Chim Phys. [2] 66, 136) Synthetically, it can be prepared by passing through a red-hot tube the vapour of benzylnaphthylmethane (Graebe and Bungener, Ber 12, 1078), or those of cumarone and naphthalene, which condense with the loss of a molecule of water (Kraemer and Spilker, *ibid.* 23, 84), or by strongly heating indene. For a complete synthesis, see Beschke, Winograd-Finkel and Kohres, Annalen, 384, 143

Purification—(Chrysene is extracted from the mixture of solid hydrocarbons consisting chiefly of pyrene and chrysene, which boil at a higher temperature than anthracene, and constitute the last portions of the distillate from coal tar, by repeated digestion with carbon disulphide in the cold. This treatment removes all the hydrocarbons in the mixture except chrysene (present to the extent of 18–19 p c.), and this is further purified by crystallisation from coal oil (b p. 150°), from which it separates in golden-yellow scales. The yellow colour, due to an impurity, cannot be removed by repeated crystallisation, but is destroyed by boiling with alcohol and a small quantity of nitric acid (Liebermann, Annalen, 158, 299), or most simply by the action of perchloric acid (K. A. Hofmann. Ber 43, 1086)

A second method for obtaining chrysene from crude greenish-yellow anthracene (m.p. 207°-208°) has been described by Schmidt (J. pr. Chem. [2] 9, 250, 270). For this purpose, 40-50 grams of the anthracene are dissolved in 5 litres of strong (95 p.c.) alcohol, and the

solution, after filtration, is boiled with 30 grams of nitric acid (sp gr. 1·4) in a reflux apparatus for some time. On cooling, red needles of a sparingly soluble compound of chrysene and dimitroanthraquinone.  $C_{18}H_{12},C_{14}H_6(NO_2)_2O_2$  separate, which, when treated with tin and hydrochloric acid, decompose into chrysene and reduction compounds of dinitroanthraquinone. The hydrocarbon is then obtained pure and colourless by crystallisation from benzene

Properties - Chrysene crystallises in scales, or in well-defined, colourless rhombic plates, which show a violet fluorescence, melts at 250° and boils at 448° (Schweitzer, Annalen, 1891, 193) It is a sparingly soluble substance, for example, 100 parts of absolute alcohol dissolve 0 097 part at 16°, and 0 17 part at 18°, and 100 parts of toluene dissolve 0 24 part at 18°, and 5 39 parts at 100° (v Bechi, Ber. 12, 1978), whilst ether, carbon disulphide, benzene, and acetic acid dissolve it only to a slight extent at the ordinary temperature, but more readily on boiling The solutions, like the pure hydrocarbon, show a deep reddish-violet On adding chrysene to a solution fluorescence of formaldehyde and sulphuric acid, a reddishviolet colour is produced, this is destroyed by the addition of water (Ditz, Chem. Zeit 31, 445) The absorption spectrum of chrysene has been studied by Baly and Tuck (Chem Soc Trans 1902, 14) Chrysene combines with dinitroanthraquinone and with pierie acid (trinitrophenol) to form characteristic compounds; the merate  $C_{18}H_{12}, C_{6}H_{2}(NO_{2})_{3}OH$ , obtained by adding chrysene to pierie acid in benzene solution, crystallises in long red needles, and is decomposed by alcohol (Liebermann, Schmidt) When heated with concentrated sulphuric acid, chrysene yields a sulphonic acid, and when treated with nitric acid, under varying conditions, yields mono-, di-, and tetra-nitro derivatives (ibid), whilst chlorine and bromine convert it into substitution-derivatives (Schmidt) Ammochrysene is obtained by the reduction of nitrochrysene with phosphorus and hydriodic acid (Abegg, Ber 1893, 792), or with tin and hydrochlone acid (Bamberger and Burgdorf, ibid. 1893, 1006) On oxidation with chromic acid in acetic acid solution, chrysene is converted into chrysoquinone  $C_{18}H_{10}O_2$ , which crystallises in yellowish-red needles, melts at 235°, yields azines on treatment with ortho-diamines (Liebermann and Witt, *ibid* 20, 2242), and is characterised by dissolving in concentrated sulphuric acid with a blue colour On oxidation with lead peroxide in alkaline solution the quinone is converted into β-phenyl-a-naphthoic acid, from which by elimination of carbon dioxide, \$\beta-phenylnaphthalene is obtained.

CHRYSEOLIN v AZO- COLOURING MATTERS. CHRYSIN, 1 3-dioxyflavone C<sub>15</sub>H<sub>10</sub>O<sub>4</sub>, is obtained from poplar buds (Piccard, Ber 1874, 88; and Darier, ibid 1894, 21) It has been synthesised by treating phloracetophenone with ethyl benzoate in the presence of sodium, when 2:4:6-trimethoxybenzoylacetophenone is formed; on boiling with hydriodic acid, the methyl- groups are eliminated, condensation takes place with the production of 1:3-dioxyflavones (Emilewicz, v. Kostanecki and Tambor, ibid. 1899, 2448; 1904, 3167). It is a yellow crystalline solid, melting at 275°. By heating

it on the water-bath with methyl iodide and potassium methoxide, tectochrysin, which also occurs in poplar buds, is formed (Emilewicz, Kostanecki and Tambor, lc) Chrysin is a member of the quercitin series (Perkin, Chem Soc. Trans. 1896, 1443), and dyes cloth mordanted with aluminium a pure yellow, with chromium, an orange-yellow; and with iron, a pale chocolate (Perkin, ibid. 1897, 818)

CHRYSOBERYL. Aluminate of beryllium, BeAl<sub>2</sub>O<sub>4</sub>, crystallising in the orthorhombic system, and used as a gem stone It is usually met with in well-formed crystals, which as a result of twinning often simulate hexagonal forms, or as rolled pebbles in river gravels Sp gr 3 65-3 75; hardness 81, taking the third place amongst minerals, being exceeded only by diamond and corundum. Few minerals are so resistant to chemical reagents as chrysoberyl, it is not attacked by acids nor fusible before the blowpipe, and is decomposed only by fused alkalis The colour ranges in various shades of yellow and green The variety alexandrite is emerald-green in daylight, but violet-red in lamp-light Another gem variety, known as cumophane or 'oriental cat's-eye,' shows a blush milky band of reflected light which travels across the convex polished surface as the stone is moved about ('hrysoberyl occurs embedded in granite and crystalline schists Material of gem quality comes from Ceylon, Brazil, and the Urals By reason of its high degree of hardness it has occasionally been used for jewelling watch-bearings

CHRYSOCOLLA. A hydrated silicate of copper sometimes of importance as an ore of copper. The formula is usually given as CuSiO, 2H2O, but the composition is variable, the mineral often being mixed with various impurities-silica, alumina, copper carbonate or phosphate, limonite, &c On these differences in composition several trivial varieties have been distinguished by special names The mineral is of secondary origin, occurring in the upper portions of deposits of copper ores It forms encrusting or botryoidal masses, which may be viticous and translucent, resembling opal or enamel in appearance, or dull and earthy. The colour ranges from bright green to bright blue, and there is a wide range in the sp gr 2 0-2 8 Under the microscope it sometimes shows a minutely fibrous structure, and small accoular crystals from Idaho have been described (J B Umpleby, 1914) Such material is optically uniaxial and positive with refractive indices 1 57 and 1 46, sp gr 2 4 Interbanded with the microcrystalline material is often some optically isotropic material, and for this amorphous equivalent of chrysocolla the name cornuite has been proposed (A. F. Rogers, 1917)

A variety known as pitchy copper-ore (Ger Kupferpecherz), dark brown or black in colour and with the appearance of pitch, is of some importance as an ore of copper, particularly in Chile. It consists of a mixture of chrysocolla and limonite Chrysocolla of the more ordinary type has been mined for copper in South Anstralia, Belgian Congo, Chile, Arizona, Idaho, &c. The name chrysocolla, from  $\chi \rho \bar{\nu} \sigma \delta s$ , gold, and  $\kappa \delta \lambda \lambda \alpha$ , glue, was applied by the ancients to materials used for soldering gold, and included, besides certain copper minerals, also borax.

L. J. S.

CHRYSOFORM. Trade name for dibromoduodohexamethylenetetramine.

CHRYSOGEŇ. An orange-coloured stance, containing more than 94 p c. of carbon, and said to be a hydrocarbon, discovered by Fritzsche (Zeitsch Chem 1866, 139) in crude anthracene. It is sparingly soluble in the ordinary organic solvents. In sunlight, its solutions become rapidly bleached, and yield a colourless crystalline compound which, on fusion, becomes orange-yellow

CHRYSOIDINES v AZO- COLOURING MATTERS. CHRYSOIN v Azo- colouring matters

CHRYSOLITE v OLIVINE

CHRYSONAPHTHAZINE " AZINES.

CHRYSOPHANIC ACID (Chrysophanol) v. CHRYSAROBIN, RHUBARB

CHRYSOPHENIN v AZO- COLOURING MAT-TERS

CHRYSOPRASE v CHALCEDONY

CHRYSOTILE v ASBESTOS AND SERPENTINE

CHRYSOTOLUAZINE v AZINES

CHRYSOTOLUIDINE C21H21N3 A yellow crystalline base of unknown constitution (cf Hofmann, Bei. 2, 380), occurring in the residue obtained in fuchsin manufacture by the aisenic acid method (Guard, De Laue, and Chapoteaud, Compt rend 63, 964, 64, 416) The hydrochloride is a yellow dyestuff

CHRYSYL. Trade name for zine boropiciate obtained by boiling together solutions of boric and pieric acids with zine oxide, used as an

antiseptic

CHURCHITE. A hydrated phosphate of cerium (('e<sub>2</sub>O<sub>3</sub> 51 87 p c ) and calcium, occurring as grey fan-like aggregates of acicular crystals encrusting quartz and slate in a Cornish copper Unfortunately nothing is known of the exact locality of the mineral CHYMASE v FERMENTATION

CICATRICINE. Trade name for a solution

of thiosinamine, antipyrine, and lucame lactate CICUTOXIN  $C_{19}II_{26}O_3$ , the poisonous principle in water-hemlock (Cicuta ragans) is contained in the tubers to the extent of 0.3-0.4pc It is found also in C virosa It is a yellow resin-like substance which readily decomposes and polymerises above 50° D<sup>22</sup> 0 9659  $h_{\mathrm{D}}^{25}$  1 5885. Unites energetically with bromine and ignites spontaneously in contact with strong On dry distillation it yields an oil nitric acid  $\mathrm{C_{13}H_{21}O_{2}}$  of sp gr 0 936 at 20° Distilled in a current of steam it gives an oil C14H22O, of a pleasant odour, it gradually polymerises to a transparent semi-solid resin On oxidation with nitric acid cicutoxin yields carbon dioxide, oxalic acid, hydrogen cyanide, isobutyric acid, and acetyl-2-cyclopentanone, b p  $735^{\circ}/15$  mm , Compounds of cicutoxin with lead, barium hydrogen chloride, ammonia, bromine, and iodine are known Cicutoxin appears to be a derivative of 4-pyrone (Jacobson, J. Amer. Chem Soc 1915, 37, 916)

CIDER or CYDER is a vinous beverage produced by expressing and fermenting the juice of Perry is a similar article made from pears The two are very much alike in character, and, the principles of manufacture being generally the same for each, reference will be made to eider only in this article, except when occasion requires the special mention of perry.

The largest cider-producing country of the world is France, followed by England, Germany, the United States and Canada, and it is now also produced in increasing quantities in other countries, notably Ireland and Australia England the principal eider-growing districts are the counties of Somerset, Devon, and Hereford, smaller quantities being made in Gloucester, Worcester, Monmouth, Kent, and Norfolk The mean annual production of the United Kingdom is estimated at 60,000,000 gallons, of a value of about 3,000,000l, including the Excise Duty of 4d per gallon.

The ordinary cider of commerce in England is a beverage of widely varying character. It may contain as little as 2 pe or as much as 8 pc of alcohol by weight, depending on the quality of the apples from which it is made, and the process of its manufacture, particularly the stage at which the fermentation has been stopped Apart from its characteristics of flavour, body, and colour, cider may be roughly divided into three classes or grades, viz: sweet, which has been only slightly fermented; dry, well fermented, and medium, between the two Apple juice of good average quality has a sp gr of about 1050 to 1055, corresponding to 10-12 pc of sugar As the fermentation produces, roughly, I p c of alcohol from every 2 p c of sugar, the following table shows the approximate comparative composition of apple juice, and the three grades mentioned —

	Sp gr	Sugar	Alcohol
Apple juice	1050-1055	10-12 p c	-
Sweet cider .	1025-1035	6-7	2-3 p c.
Medium eider	1015-1025	4 -5	3-4
Dry cider .	1000-1010	0-2	5-6

It must be understood that these figures are only approximate, and show the proportions of alcohol and sugar in cider made from apples of average quality, when the fermentation is stopped at the sp grs mentioned and no sugar is added

The rough cider of the farmhouse, found in the south-western counties, is of a very dry grade, being fully fermented, no attempt being made to check the 'working,' while the ordinary cider (bottled and draught), found in the average hotel in cities, is a beverage which has been but slightly fermented, and is of the sweet to medium grade, the greatest public demand being for cider as a summer beverage, light, sweet, and refreshing It must not be assumed, however, that all bottled eider is sweet, for it is obtainable from several firms of repute in four welldefined grades-sweet, medium, dry, and extra

The raw material from which eider is made varies in composition and character as much as the cider itself. The majority of English orchards were originally stocked with almost innumerable varieties of apples and pears, most of them of inferior quality, for either table use or for the manufacture of eider or perry, but great improvements have of late years been made in the planting and cultivation of the orchards, with a corresponding improvement in the quality of the beverages produced. Speaking generally, it may be said that good cider cannot be made from any but essentially vintage varieties of apples, most of which are quite

unsuited for table use because of their astringent flavour, due to the large proportion of tannin they contain The majority of table varieties yield a juice deficient in sugar and tannin, with an excess of acidity, as compared with the vintage sorts, which are cultivated specially for the manufacture of cider This is particularly applicable to perry pears, which are almost invariably very astringent in taste, although it is curious to note that the percentage of tannin they contain is not exceptionally high. To make good cider, the sp gr of the juice should not be under 1045, it should contain at least 0 15 pc of tannin, which gives keeping and clearing properties, with an acidity not exceeding 05 or 0 b pc of malic acid for sweet cider, and 03-04 p.c. for dry, while those substances which constitute 'body,' consisting of mucilage, albumen, salts, colouring matter, &c, are also of importance; but, having been very little investigated, it is impossible to state definitely what proportions of each are necessary high sp gr with a correspondingly high sugar content, is generally considered the most important quality, but some apples are known to yield a juice high in sp gr but deficient in colour and body (the two are usually associated), while others of a low sp gr but rich in colour and body, may produce cider of better quality than the first-mentioned Too much reliance, however, must not be placed on the mere chemical composition of the juice as indicative of the value of an apple for eider making, the characteristic flavour of the variety and the rate of fermentation, which varies greatly with different juices, being at least equally important

Vintage apples are roughly divided into three classes—sharp, containing normally not less than 0 45 p c malic acid; sweet, containing normally less than 0.45 pc of malic acid, and 02 pc of tannin, bitter sweet, containing normally less than 0.45 p.c. of malic acid and more than 02 pc of tannin The following table shows typical analyses of the juice of several of the best of each class, together with that of a hitherto unknown and unnamed variety recently found in Someiset ('Butleigh No 14'), which is of outstanding merit -

No	Variety	Sp gr.	Malic acid	Tannın	Total sugar
ı	KingstonBlack	1060	0 54	0 126	12 66
2.	Foxwhelp .	1059	1 04	0 270	$13\ 25$
3	Cowaine Red	1054	0.65	0 230	12.05
4	Sweet Alford	1057	0 14	0 120	12 19
5	White Jersey	1060	0.25	0 190	13 33
6	Cherry Norman	1055	0 21	0.354	12 25
7.	Horner	1053	0 15	0 240	11 89
8	Royal Jersey	1065	0 16	0 380	15 00
9.	Barland .	1054	0 62	0 119	11.32
10	Butt .	1053	0 66	0.270	11.75
11.	Oldfield .	1068	0 80	0.128	14 92
12.	'Butleigh No 14'	1104	0.24	0.258	22.22

Nos. 1, 2, and 3 are sharp varieties; 4 and 5, sweet; 6, 7, and 8, bitter-sweet; 9, 10, and 11, perry pears; 12, a bitter-sweet of unusually high quality.

ancient origin, but modern scientific eider making may be said to date from the year 1893, when the Bath and West of England Agricultural Society instituted some chemical and bacteriological research into the subject, a work which was carried on by Mr. F J. Lloyd for some seven years in Somerset, and has subsequently been continued on a more ambitious scale at the National Fruit and Cider Institute, associated with the University of Bustol, under the direction of Prof B T P. Barker The results of these investigations form the foundation of modern cider making, and annual reports thereon constitute the most trustworthy literature on the subject

Whereas the quality of the cider depends primarily on that of the apples used, the nature of the product is largely influenced by the methods of manufacture and fermentation, and general management To ensure good quality, care of the apples in the orchard is essential, for no good beverage can be made from fruit which has been carelessly gathered or stored It should be shaken from the trees when just ripe, bruising being avoided as far as possible, and stored in heaps (preferably under cover if the weather be wet), until it becomes soft and mellow, but not over-ripe, care being taken to avoid either washing by rain, which results in the loss of sugar and yeast, or heating, which produces fustiness in the cider, as well as a great When thotendency to acetic fermentation roughly mellow, the fruit is selected and blended in the proportions necessary to give the degree of sweetness, acidity, and astringency required for the variety of cider it is intended Sweet cider is best made from a to make blend of sweet, sharp, and bitter-sweet apples; while dry cider should be made from a blend giving a minimum of acidity, with only a moderate proportion of tannin Very few varieties of apples are individually capable of producing the finest cider, an appropriate This does not apply blending being necessary pears and perry making: single varieties, unblended, usually giving best results In the actual processes of grinding the fruit and expressing the juice, steam or hydraulic power is generally used in modern factory practice, but, apart from economic reasons, this has no advantage over the old-fashioned hand-power machinery of former days. The fruit is ground to a fine pulp, called 'pomace,' which falls upon the press, is held together in thin squares by manilla or hempen cloths, and is usually pressed In some places, however, the pomace is allowed to remain in tubs for 12 to 24 hours before pressing, which is said to improve the quality of the resulting liquor, but there seems to be no scientific reason for this custom, and investigations have failed to show that sufficient advantages are gained to compensate for the work it entails. With efficient presses, about 75-80 pc of the weight of apples is obtained as juice, and about 80 p c in the case of pears. The most profitable means of utilising the drypressed pomace is as cattle food In France the juice is sometimes extracted by the 'diffusion' process in use in the beet-sugar industry, but this does not meet with general approval, and has not been adopted in this country. The process of cider making is of very fresh juice is run into casks or open vats called

'keeves,' to undergo the important process of fermentation, which commences spontaneously in a few days, the time depending on the character of the juice and the temperature An important question here arises. Shall the cider be allowed to undergo a natural fermentation, or shall a pure culture of yeast be added to dominate it? At the present time this is a debatable point While in France and Germany it is generally acknowledged that the use of pure cultures improves the quality of the liquor, and they are largely used, in this country and America it seems doubtful whether the advantages of their use are sufficient to compensate for the extra expense and trouble involved Pure yeasts, isolated from the best growths of cider and wine, have been cultivated and added in sufficient quantity to dominate the fermentation and prevent the various wild races from exercising much influence (Owing to the difficulty of avoiding a cooked taste, sterilisation of the apple juice, even at comparatively low temperatures, is rather impracticable) The general results tend to show that fermentation by suitable yeasts give a somewhat superior product to that obtained by natural fermentation, but the full advantages are not apparent until the cider is about a year old, whereas, in general practice, it is usual for it to be consumed before it attains that age. The use of sumed before it attains that age pure cultures in the eider industry in this country is, for that reason, far from general, and it does not seem likely to make much headway in the present state of our knowledge of the subject Little is known of the various species of yeasts naturally occurring on the skin of the apple, which subsequently develop in the cider, forming the 'lees' at the bottom, and conducting the fermentation, but the largest number consist of forms allied to the Saccharomyces Ellipsoideus, S Pastorianus, and S apiculatus types A fairly slow fermentation is generally desirable, but, on the other hand, it is one of the most important points in the whole process that the primary alcoholic fermentation be vigorous enough, and is not checked too soon, otherwise the eider is laid open to the attacks of numerous undesirable organisms, which will show their effects in various 'diseases' during the following summer The 'keeving' system of fermentation, in open vats with loose covers, is preferable to that in closed casks, as the 'head,' which is soon thrown up, consisting of yeasts and light portions of tissue, can be skimmed off from time to time It is not advisable, however, to keep the order in the keeves longer than about a week, lest incipient acetic fermentation set in, and, as soon as no further head is thrown up, it is racked into closed casks to complete the primary fermentation. The subsequent conduct of this important process is governed by the grade of eider it is desired to make, the hydrometer being in regular use for watching the fall of the The rate of fermentation depends sp gr partly on the character and composition of the juice, and partly on the temperature; but these conditions are practically beyond control. Blending the apples to ensure the requisite degrees of sweetness and acidity in the juice will comply with one condition of a moderate rate, but as the proportion of nitrogenous matter is

of still greater importance, and is a subject on which, strangely enough, little is known, and no reliable figures obtainable, blending to ensure the proper percentage cannot be scientifically done at present Regulating the temperature is also impracticable, for a large quantity of cider would have to be handled in a very short space of time, and this would be too expensive a process. A large plant would have to be fixed up, which could only be in use a few weeks each year, and possibly it might not be required at all Whenever the spgr. approaches the point at which it is desired to stop the fermentation, the eider is racked off into clean casks, with or without sulphuring or the use of other This alone, however, is not preservative. usually sufficient to stop fermentation, unless an excessive quantity of preservative be used, or racking into sulphured casks be repeated several The best known means, which has the advantage of adding no materials of a questionable nature to the liquor, is to filter it through one of the many useful cotton-pulp filters now on the market Even the use of a filter is open to some objections, for it removes some of the body (probably of a colloid nature) and flavour from the cider The clarification also presents great difficulty with many ciders, neither isinglass finings nor boiled apple pulp, which are commonly used for the purpose, having any effect, so that the filter, which clarifies the cider, and at the same time stops the fermentation, is preferred, in the absence of anything better, for this purpose Depending on various circumstances, the cider is filtered and stored away to mature in casks, at any time from two to four or even six months from the time of making If filtered perfectly bright, and stored in clean casks, kept well filled, it will 'keep its sweets' for a whole season or longer, with the loss of only a few degrees of spgr, even though it may contain as much as 6 or 7 pc of fermentmay contain as much as 0 of 1 p o 3. Assume that a ble sugar. During this period of storage, it mellows and develops its fragrant aroma and delicate flavour, and also loses a certain proportion of its acidity and astringency. The strongest and best cider will keep in cask for some years, but that of average quality is at its best within the first year It keeps best and improves most in bottle, but only eider of good and sound quality should be bottled; it should have an original gravity of at least 1050, should undergo a vigorous primary fermentation, and be perfectly brilliant when bottled The best age for bottling is from four to six months, and if not more than a year old at the time, it will develop sparkling condition like champagne after a few weeks' storage. The carbonating process of bottling is becoming more general, and there seems to be no objection whatever to its use, some of the best brands on the market having been treated in this way

Matured cider is liable to several disorders or 'diseases,' the principal being acetification, 'sickness,' ropiness, and blackening Like all liquors of low alcoholic strength, it is particularly apt to acetify on very slight provocation, and for this reason, exposure to air at all stages of manufacture is to be avoided. 'Sickness' is a recommencement of alcoholic fermentation, which is set up during the summer following its manufacture, and may occur in cask or

bottle. The cider becomes very turbid, and the production of alcohol is accompanied by the development of a most disagreeable smell and flavour, due, it is thought, to formation of acetaldehyde, together with an increase in acidity, partly lactic and partly acetic, which renders the liquor quite undrinkable. It is a source of great loss, for it usually attacks the richest and most fruity ciders, and is due to the action of several special organisms, contributing causes being a deficiency of malic acid in the juice, and a weak primary fermentation Ropiness is not so frequently met with, and little is known of its causes in eider, but it is generally found where there is a deficiency of tannin. Some ciders turn dark very quickly on exposure to air, some almost black in less than half an This may be due either to an excess of iron acquired from the soil on which the apples are grown, but more often from the utensils used in the making, or to enzymic action, the use of a little preservative often prevents its development

A standard of chemical composition for what is sold as eider has been formulated in some countries, but not in Great Britain to the extreme variations in the character and composition of genuine cider, it is almost, if not quite, impossible to fix a definite chemical composition and say that it represents what should be considered a standard cider. For this reason, the standard fixed in foreign countries is necessarily very low, but while it probably admits of a certain amount of adulteration, it no doubt acts as a check. It can hardly be contended that nothing should be sold as eider which is not the pure juice of apples Brewers, wine makers, and distillers are not restricted by law to the use of malt and hops, grape juice and malt respectively, in the making of their products, so that a law to limit cider makers to the use of apples only would operate unfairly against them as compared with the other trades It has been stated that many mentioned varieties of cider apples grown in this country are not capable of making first-class cider, and the better sorts are in too small a minority to meet the demand In poor vintage seasons, also, the general quality is low, sugar being deficient and acidity excessive It is also deficient and acidity excessive It is also undoubtedly true that cider made from the poorer varieties of apples, and in poor vintage years, is capable of great improvement by the addition of a certain proportion of cane-sugar syrup, as is done in the process of 'gallisising' wines, to obtain the correct proportion of acidity to sugar It is difficult to say what maximum proportion of this syrup should be allowed, but there is seldom any occasion for the use of more than 10 p c to make the cider suitable for the English market. The fact that there are so-called 'ciders' on the market, which are innocent of a drop of apple juice, shows the necessity for some sort of standard Generally speaking, it may be concluded that no cider can be considered genuine which does not contain at least 2 p.c. of alcohol by weight, or which does not show an original gravity of 1045 or upwards. Another vexed question is that of the use of preservatives, some people contending that it is impossible to do without them, and others that they are not necessary if Andes, stretching from Columbia in the north,

the cider be well and carefully made There is much to be said on both sides Some makers sulphur their casks, thus adding a certain quantity of sulphurous acid to the cider, but it is generally considered, though very strangely, that this does not constitute the addition of any chemicals.' Others use sulphites or salicylic acid, and justify their use by the argument that all eider, no matter how carefully made, is peculiarly susceptible to the various 'diseases' previously mentioned, and has often to be kept under conditions which render them unpreventable by any other known means, for example, on draught in a warm bar, perhaps for some weeks. There is certainly a great deal of reason in this, and it would probably be a hardship to absolutely prohibit the use of any preservative, but, on the other hand, they are sometimes used in needlessly excessive quanti-ties, which, in the interest of the public health, should be avoided The consumption of eider is rapidly increasing, due partly to its being recommended by medical men, and if it is to hold its own as a popular, and above all as a healthful, drink, it must be kept free from any ingredients of doubtful physiological effects.

Appended are analyses of a few typical ciders of well-known makes -

Description	Sp gi	Alcohol by weight	Total	Total solids	Ash
	1006 1015	4 90 3 90	$\begin{array}{c} 0.31 \\ 0.53 \end{array}$	3·25 6 00	0 245 0 314 0 360 0 390

W D. McC

CIGNOLIN. Trade name for 18-dihydroxyanthranol Used as a remedy for psorrasis Has a more energetic action upon the skin than its 3-methyl derivative, chrysarobin (q v), Wochenschr (Unna, Dermatol 1916, Nos 6-8)

CINCHOMERONIC ACID & BONE OIL

CINCHONA ALKALOIDS. The Cinchona Barks. 1 Their Discovery—The febrifuge properties of the back of the einchona trees growing on the slopes of the Andes was probably known to the native Incas long before its curative properties became known to their Spanish conquerous The earliest mention of a 'cure' is by Jussieu in 1600. The introduction of the bark to European countries was due to the Countess of Chinchon, wife of the Viceroy of Peru, who having been cured of an attack of fever at Lima in 1638, on her return to Madrid in 1640, brought with her some of the precious bark, and administered it to fever-stricken patients on her Spanish estates It is in her honour that Linnaeus named the species 'cinchona' For nearly 200 years the drug was administered in the form of a decoction of powdered bark, and the alkaloids in it were not isolated till 1816, when Gomez discovered the alkaloid cinchonine, followed by Pelletier and Caventou, who isolated quinine

2 The natural Cinchona Trees occur in a long but very narrow belt on the slopes of the through Ecuador and Peru to Belivia on the south. For the greater part of the last century the trees were ruthlessly cut down and the bark stripped from them by the Indian collectors, no attempt being made to replace them or cultivate seedlings, and consequently the species was being rapidly exterminated when more scientific methods were gradually introduced

The trees were first classified botanically by Weddell, in 1849, and in 1860 John Eliot Howard, F.R.S., brought up-to-date and translated the notes of the Spanish botanist Pavon (compiled in 1800), under the title of the

'Nueva Quinologia of Pavon,' and these two books still maintain their position as the standard works on the subject. In 1880 Sir Clements Markham published a most comprehensive work entitled 'Peruvian Bark,' and it is from this book that the compilers of these notes have obtained much of their information. The following table shows the locality and the botanical species of the various commercial barks on the London market in about 1870, and the analyses given, except when otherwise stated, were made by the present authors, of typical commercial samples which had been preserved in a collection of that date

TABLE I -THE NATURAL BARKS OF COMMERCE OF 1870

	-	,							-
				1	ercent:	Typical age of a	analyse: Ikaloids	s present	,
Commercial name	Botanical name	Locality	D15c0ve1€1	Quinine	Cinchoni- dine	Quimqine	Cinchonine	Amorphous alkaloid	Total alkaloids
, Crown Barks	Cinchona officinalis	Loxa (S Ecuador)	Caldas, 1803	3 09 J E	1 34 H (1		0 32	0 59	5 48
Red Barks	Cinchona succirubia	Chimborazo (Central Ecuadoi)	Condamine, 1738		honine	ne and	umme quinidi	ine 1 :	90 22 25 —
1					Tota	ıl alkal	oids	4 3	37 рс.
Columbian Barks Hard	Cinchona pitayensis	Pitayo (Columbia)	Mutis, 1772	2 42	 0 19	 0 82	0 98		 6 66
Pitayo Soft	,,	,,		1 91	0 16	0 77	1 28	1 50	5 62
Pitayo Soft Colum- bian W Coast	Cinchona laneifolia	Santa Fe and New Grenada		1 42	0.36	0 09	0 35	0 65	2 87
Carthagena		Carthagena and Gui- aquil		131	041	0 03	0 07	0 40	2 23
Grev Barks	Cinchona nitida Cinchona micrantha Cinchona Peruviana	Huanuco (N Peru)	Renquifo, 1776		H (1	862) Ci otal alk 862) Qi	' Nearly cincl nchonn caloid 2 uinine ( inchoni	honine ne 2 p ? 7 p c ) 45 p c	e c
Yellow Barks	Cinchona calisaya	Bolivia and S Peru	Haenke and Celis 1776	2 00	0 46	0.20	0 56	0 90	4 12

Note.—The analyses marked J E H are from records of John Eliot Howard The results must be accepted with reserve, as at the date mentioned, the methods of separation of the different alkaloids were very rudimentary.

3 Transplantation and Cultivation—In the middle of the last century the necessity of dealing adequately with the devastations of malaria in India and the diminishing supply of natural cinchonas, directed attention to the

importance of their cultivation in suitable districts in that country.

In 1859 Sir Clements Markham organised expeditions to South America for the purpose of securing seeds of all the cinchonas then of

commercial importance. He himself headed a search for calisaya seeds in Peru and Bolivia Spruce collected succirubra in Ecuador, while Pritchett was responsible for the grey barks of Huanuco in Peru. In order to complete the series further expeditions under Cross were carried out in 1861 for officinalis back from Loxa (S. Ecuador), in 1868 for Pitayo bark from Columbia, and a final search in 1877 for soft Columbian and Carthagena barks.

Markham's narrative of these expeditions tells of the immense difficulties which were surmounted by these public-spirited pioneers, not only from the actual hardships of the expeditions, but also from the suspicious and often actively hostile attitude of the Indians, and to crown all, the problem of transporting the collections to the coast, and carrying them alive to England, and subsequently through the

Red Sea to India

All these difficulties were, however, overcome, and a large proportion of the seedlings survived the ordeal and were successfully planted in the Nilgiri Hills in 1861, under the superintendence of MacIvor. The cultivation of cinchona in India from MacIvor's seedlings rapidly spread, and plantations were established by private owners in the Nilgins, and in the Palnai Hills of Travancore In 1865 plantations established at Hakgala in Ceylon and in Coorg, followed shortly after by further enterprises at Wainad, Mysore, Darpling in Sikkim, and the Karen Hills in Burmah Maclyoi introduced two methods of harvesting the bank .-

1. Coppicing and allowing the tree to shoot

again from the same trunk, and
2 'Renewing' by mossing In the latter
process the back was removed in longitudinal strips about an inch wide, leaving a space untouched between each strip denuded of bark The tree was then bound round with moss and the bark allowed to 'renew' and cover the This method, although more laborious than coppicing, largely increased the yield of each tree David Howard (J Soc Chem Ind 1906, 26, 97) states that renewed bark is richer in alkaloids than the original, and that the proportion of quinine to other alkaloids is

In 1880, when Markham's book was published, the Indian plantations gave every promise of becoming the main source of supply for the markets of the world The fact that India does not find herself to-day in that fortunate position is shown in the following

table, and the cause in the next paragraph

A summary of sources of bark sold in European markets -

In 1880	In 1911
6,000,000 lbs )	
, }	35,200 lbs.
960,000 lbs	
1,170,000 lbs.	458,600 lbs.
70,000 lbs	19,778,000 lbs.
21,000 lbs.	nıl.
nıl.	25,400 lbs.
	6,000,000 lbs 960,000 lbs 1,170,000 lbs 70,000 lbs 21,000 lbs.

The explanation obviously rests on the rise of Java as the greatest cinchona producer, and it is therefore necessary to sketch briefly the history of this wonderful success of the Dutch in their principal colony.

The first attempt at cultivation in Java in 1852 ended in complete failure through the poor alkaloid-producing characteristics of the varieties of cinchonas of which seedlings were secured from South America, the only trees producing a tolerable yield being a small number obtained from MacIvor's plantations in the Nilgiris In 1855, however, Charles Ledger secured some seeds of Cinchona calisava, half of which were planted in India and half in Java. Those in India failed, owing either to the chmatic conditions or the mismanagement of the plantations after MacIvor's death. Those in Java, however, flourished exceedingly, and have produced a wonderful strain of trees, which, by skilful scientific cultivation, have produced, and are producing, back containing up to 10 pc of quinine, and which have, therefore, far surpassed any other variety in its market value from a manufacturing standpoint

4. Hybridisation and Grafting — David Howard (J Soc Chem Ind 1906, 25, 97) deals Grafting — David with the effect of hybridisation (partly deliberate and partly accidental) on Java barks, and mentions a very common hybrid of commerce known as Robusta, a hardy growth, which nearly rivals Ledgeriana in its alkaloid value Regarding grafting, he states that the 'Enten Baast' of commerce, from Java, is the result of grafting Ledgeriana on to a Succirubra stock, the bark from branches giving the characteristics of Ledgenana, while the root back retains the proportion of alkaloids common to Succirubra.

From a large experience of analysis of samples from different parts of the same cinchona trees, he found that the elaboration of alkaloid takes place solely in the downward course of the sap, and that the percentage of alkaloid increases in the bark from the twigs down the stem to the ground, in the less flourishing tices the root back being richest in alkaloid value, but of this alkaloid, the proportion of quinine and cinchonidine is lower, while that of cinchonine and quinidine is higher

5 The Commercial Aspect -In the same paper David Howard shows that the conquest of the world's markets by the Java planters was due to the careful and scientific work of Moens and others in the cultivation of the phenomenally 11ch Ledgeriana to the exclusion of the lower alkaloid-producing varieties, which enabled the Dutch growers to hold on during a period of about 20 years of bitter competition and low prices, which gradually ousted a large number of the older established, but less scientifically managed, Indian plantations from the market.

To emphasise the effect of successful Java cultivation the table given below shows two A, typical of the very best natural analyses Calisaya bark produced from wild trees in about 1865; and B, of a typical good quality cultivated Ledgeriana bark sold in Amsterdam ın May, 1914.

Quini- Cincho-Cincho-Quinine nidine dine nine Amor-Total phous alkaloids рс 0 165 рс 0.34 3 615 рс 075 5.800.937 92 0 105 0.08 0.0850 91

This shows the triumph of skilful cultivation in producing an enormous alkaloid yield of almost pure quinine, over the best unaided

efforts of nature.

6. Other Bark-producing countries only require a passing mention. The barks formerly produced by Jamaica and Mexico have disappeared from the market. The Portuguese possessions in West Africa produce bark regularly, the best-known source being the island of St Thomé South America still produces a considerable amount of bark, presumably now entirely from cultivated and not natural sources. A start was made shortly before the war in German East Africa, and although the plantations are at present small, the analysis of the samples shown below, if they can be regarded as typical, indicate very promising conditions, being in quality well up to the best Java production.

### PERCENTAGES OF ALKALOIDS PRESENT

Samples from German East Africa	Quimme	Cinchonidine	Cinchonine	Quandine	Amorphous	Total alkaloids
Ledgeriana Hybrid . Succirubra Robusta .	2 54	$\begin{array}{c} 0.00 \\ 2.05 \end{array}$	$\begin{array}{c} 0.97 \\ 2.14 \end{array}$	0 00 0 00 0 00 0 00	1 92 1 59	5 00 11 30 8 32 7 61

7. Other Trees producing Cinchona Alkaloids -Two other varieties of trees must be mentioned which, although belonging to the Remijia species, and not being botanically cinchonas at all, yet contain alkaloids of the cinchona group, which will be mentioned below

Bark of the Remijia pedunculata, known as 'Cuprea Bark,' was sold in considerable quantities on the London market in the early eighties, and its analysis is interesting, as being the only natural bark known which contains quinine and other alkaloids (including the rare alkaloid cupreme), but absolutely no einchonidine

The Remijia Purdieana contains interesting but rare members of the family of cinchona alkaloids, viz cinchonamine, chairamine, con-chairamine, chairamidine, and conchairamidine, the peculiar properties of which will be mentioned

REMIJIA PEDUNCULATA ('CUPREA') BARK.

	Sample A	В	C
Quinine alkaloid Cupreino . Quinidine Cinchonine Cinchonidine Amorphous .	. 1 38 0 11 . 0 40 . 0 22 . mil . 1 31	1 27 0 05 0 41 0 26 nil 1 01	1 20 0 25 0 27 nil 1·17
Total alkaloid	. 3 42	3 00	2 89

#### REMIJIA PURDIEANA BARK.

Cinchonamine nitrate Nitrate of other paricine bases 3 30,

There is one variety of cinchona which, although valueless from a commercial point of view, since it contains none of the four alkaloids of commerce, has yet provided a considerable field for research, as several rare members of the cinchona alkaloid family have been extracted from it—It is described by Markham under the name of Pubescens, but was also called Pelleteriana (after the discoverer of the quinine base) and Cusco bank in commerce.

It would appear to have been widely distributed over the forests of Peru, and was found to be growing freely by Markham in the Cara-The alkaloids isolated from it vava district are cusconine, cusconidine, cuscamine, cuscamidine and aricine (called after Arica, the coast

town from which it was exported).

Before leaving the subject of barks, the tree Aspidosperma Quebracho, a member of the Apocynacex, must be mentioned. This was considered by Hesse (Annalen, 1870, 154, 287) to be identical with the Quina Blanca of Mutis.

Two alkaloids have been extracted from it, Paytine and Paytamine (so called from the port Payta in North Peru), which although not actually classified under any group of cinchona alkaloids, have a very similar empirical formula, and possess certain febrifuge properties which justify their inclusion under this heading

Principles of Cinchona Barks.—The following are the names and formulæ of the alkaloids occurring naturally in these barks, isomers being bracketed together -

Cinchonidine Cinchonicine (cinchotoxin) 3 4 Hydrocinchonine (cinchotine)
5. Hydrocinchonidine (cinchami) dine) Cinchonamine Cupreine C<sub>19</sub>H<sub>22</sub>O<sub>2</sub>N<sub>2</sub> Quinamine Conquinamine C19H24O2N2 10. Quinine Quinidine Quinicine (quinotoxin)

13. Hydroquinine C<sub>20</sub>H<sub>26</sub>O<sub>2</sub>N<sub>2</sub>

15. Chairamine Conchairamine 16 17 Chairamidine

18 Conchairamidine 19. Aricine

1. Cinchonine

20 Cusconine 21 Concusconine

22 Dieinchonieine C38H42ON4 Diquinicine (diconquinine) C40H46O3N4

24. Javanine

25. Cusconidine formulæ not determined

27 Cuscamidine

Paricine C<sub>16</sub>H<sub>18</sub>ON<sub>2</sub> 29 Paytine

29 Paytine  $\left\{ C_{21}H_{24}ON_{2}\right\}$ 

These are combined in the bark mainly as quinates and cinchotannates with quinic acid  $C_7H_{12}O_6$ , and cinchotannic acid  $C_{14}H_{16}O_9$ ; the former occurring to the extent of 5-8 pc., and the latter 3-4 pc in the barks. Quinovic acid C<sub>32</sub>H<sub>48</sub>O<sub>6</sub> and oxalic acid (as the calcium salt) are also present in small amounts. The

quinates are soluble in water, but the cinchotannates only slightly soluble, dilute acids, however, decomposing them and effecting solution The colouring matter in the bark is cinchonared C<sub>12</sub>H<sub>14</sub>O<sub>7</sub>, occurring to as much as 10 p c It is soluble in alkalis, being reprecipitated by Two glucosides,  $\alpha$ -quinovin and  $\beta$ quinovin, are usually present in all cinchonas, and give on hydrolysis quinovic acid and quinovose, a sugar isomeric with rhamnose (Liebermann and Giesel, Ber 1883, 16, 926) The neutral bodies, cinchocerotine, and the isomers, cupreol and cinchol  $C_{20}H_{14}O+H_{2}O$  (Hosse, Annalen, 1884, 228, 288; 1886, 234) 375), together with cholestol (Thoms, Arch. Pharm 1897, 235, 39), appear to be constant constituents of cinchona barks

The ash of these barks falls usually between 3 pc and 4 pc, a curious fact being that the greater the altitude at which the bark is grown the higher is the ash. In Indian backs the highest ash is found in Crown barks grown at 7000-8000 feet above the sea, falling in red barks grown at 5000-6000 feet, a minimum ash being found in Ledger bark, which grows at an altitude of 3000-5000 feet (Hooper, Pharm J 1886, 17, 545)

Constitution of the Cinchona Alkaloids -Only a short account of the great amount of work done on this subject can be given here, for fuller details, consult 'Vegetable Alkaloids,' by Pictet (Biddle, 1904), 'Die Konstitution der Chinaalkaloide,' by Comanduci (Sammlung chem u chem Vottage, 1911, 16, Bande, 4-7), and Rabe's and Kaufmann's articles in the Berichte and Annalen since 1907

The constitution of cinchonine has been the most fully investigated of these alkaloids, and its relation to the other common alkaloids being simple, it is an easy matter to deduce their constitution from that of cinchonine

Cinchonine  $C_{19}H_{22}ON_2$  is a ditertiary base forming  $d\iota$ -alkyl iodide addition products. On distillation with potassium hydroxide it gives, as the main product, quinoline, and a smaller amount of lepidine (4-methylquinoline), showing that cinchonine has a methylquinoline nucleus Cinchonine may therefore be written

On treatment with the halogens, or haloid acids, it forms addition products, taking up 2 atoms of halogen or I molecule of acid, a fact pointing to the presence of a vinyl group By the action of permanganate in the cold, a body, cinchotenine, which does not contain the vinyl group, is formed, and an atom of carbon is split off as formic acid, thus

$$C_9H_6NC_9H_{13}ON(CH\cdot CH_2)+4O$$
  
Cinchonine

$$\begin{array}{c} \longrightarrow \mathrm{C_9H_6N \cdot C_8H_{13}ON \cdot COOH} + \mathrm{H \cdot COOH} \\ \mathrm{Cinchotenine} & \mathrm{Formic\ acid} \end{array}$$

The vinyl group probably exists, therefore, in a side chain. Cinchonine and cinchotenine both contain a hydroxyl group, giving mono-acetyl derivatives. This completes the knowledge of the simpler groups in cinchonine, which can now be written C<sub>9</sub>H<sub>6</sub>N·C<sub>8</sub>H<sub>12</sub>(OH)(CH: CH<sub>2</sub>)N There remains the constitution of the C<sub>8</sub>H<sub>12</sub> portion to be elucidated, together with the positions of the hydroxyl and vinyl groups.

Now, on oxidising einchonine with chromic acid, the molecule is split in two, yielding about one-half its weight of cinchonic acid (10H2O2N (4-carboxyquinoline), and about half its weight of a syrupy mass consisting of a primary body, meroquinene  $\rm C_8H_{14}$  COOH N, and its secondary oxidation products, eincholoiponic acid

and loponic acid C<sub>5</sub>H<sub>9</sub>(COOH)<sub>2</sub>N Cinchonine acid is evidently the product of the methylquinoline nucleus, and the other three acids the product of the 'second' half of cinchonine The primary oxidation product of the 'second' half, meroquinene, throws much light on the constitution of this portion of the cinchonine molecule, and is known to have the formula

Now meroquinene is a secondary base, but as emehonine is ditertiary, meroquinene must be derived from a portion of the 'second' half, having the formula

$$\begin{array}{c} \operatorname{CH}_{2}\operatorname{CH}_{2}\operatorname{C} = \\ \operatorname{CH}_{2} \\ \operatorname{CH}_{2} \\ \operatorname{CH}_{2} \end{array}$$

This and the quinoline portion account for  $C_{18}H_{19}N_2$  of the einchonine molecule, and there remains but  $CH_2(OH)$ , for which positions must be found.

On heating cinchonine with acetic acid it undergoes isometic transformation into einchonicine, a molecular rearrangement in which the hydroxyl group vanishes, and a carbonyl group appears, the tertiary character of the 'second' half becoming secondary at the same time This is a change which Rabe's researches have shown to be effected by this grouping -

$$-CH(OH)-('H-N-\rightarrow -('I)-CH_2 NH$$

This supplies the last link required in the cinchonine molecule, which may now be formulated as the basis of the cinchona alkaloids .-

In cinchonine, cinchonidine, cinchonicine, quinine, quinidine, and quinicine R=-CH CH2. In hydrocinchonine, hydrocinchonidine, hy-

droquinine, and hydroquinidine R=-CH<sub>2</sub> CH<sub>3</sub> In cinchonine, cinchonidine, cinchonicine,

hydrocinchonine, and hydrocinchonidine X=H. In quinine, quinidine, quinicine, hydroquinine, and hydroquinidine X=OCH<sub>3</sub>.

In cupreine X=OH (with phenolic proper-

ties).

VOL. II.-T

Cinchonicine and quinicine have the ketonic grouping at carbon 4, as indicated above

The numbered carbon atoms are asymmetric, and the isomerism among these alkaloids is frequently due to the spatial arrangement around the carbon atom 3, since the asymmetry of the other three carbon atoms 1, 2, and 4 may be destroyed without interfering with the isomerism of the products (Rabe, Annalen, 1910, 373, 85)

It may be noted here that in the ditertiary bases fluorescence only occurs in those in which  $X = OCH_{\eta}$ , or a higher alkyl group. The toxicity of these alkaloids also appears to increase as the alkyl group in the X position increases from methyl to octyl

mediyi to octyi

The known conversions of one alkaloid into

another may be given

Cinchonine or cinchonidine on heating with moderately strong sulphuric acid is converted to cinchonicine; quinine or quinidine yielding on similar treatment quinicine (Pasteur, Compt rend 1853, 37, 110, 166)

By heating cinchonine in amyl alcohol solution with potassium hydroxide part is converted to cinchonidine (Königs and Husman,

Ber 1896, 29, 2185)

Cupreine is converted to quinine by heating at 100°C under pressure with sodium and methyl chloride in methyl alcohol solution (Grimaux and Arnaud, Compt rend 1891, 112, 776, 1364) Similarly, higher alkylated bodies may be prepared. Hydroquinine, when heated under pressure to 140° with hydrochloric acid, sp.gr 1 125, or boiled with hydrobromic acid sp.gr 1 49, or hydrodic acid sp.gr 1 7 is converted to hydrocupreine, its methoxyl group being replaced by —OH (Hesse, Annalen, 1887, 241, 255; Pum, Monatsh. 1895, 16, 73).

73).

The hydro- bases (in which R=-CH<sub>2</sub> CH<sub>3</sub>) may be prepared from their respective unsaturated alkaloids, dissolved in dilute sulphuricand, by the action of hydrogen under pressure in the presence of palladium black (Eng. Pat. 3948, Feb. 16, 1912). Cinchonieme has been converted to einchonine by Rabe (Ber. 1908, 41,

67; 1911, 44, 2088)

Kaufmann and Huber have converted hydrocinchonicine into a mixture of hydrocinchonine and hydrocinchonidine (Ber 1913,

46, 2913)

No complete synthesis of any of the cinchona alkaloids has been effected, although bodies very closely related to them both in structural formulæ and medicinal properties have been obtained (Ber 1913, 46, 1032 and 1823; 1917, 50, 144) Robinson has suggested by an ingenious method a possible phytosynthesis of those bases from formaldehyde, ammonia, and an acotone derivative. It is interesting to note that Perkin discovered the aniline dyes whilst trying to synthesise these bases.

Therapeutic Value—The first work of importance on the comparative therapeutic effect of the various alkaloids was in 1868, when an Indian medical commission reported on the relative efficiency of the four principal alkaloids then commercially obtainable in the treatment of malaria. This commission placed the alkaloids in the following order of

efficiency:

1st | Quinine | Quinidine.
3rd Cinchonidine.
4th Cinchonine.

At about the same time, J E Howard reported, as a result of his administration to patients suffering from the mild English malaria or ague still common at that time in the Essex marshes, that he considered quinine and cinchonine equally effective, and einchonidine and quinidine somewhat less satisfactory Of late years the researches establishing the constitutional formulæ of these alkaloids has led pharmacologists to study the particular arrangement of the groups in the molecule which has the greatest effect in malaria prophylaxis and other uses of quinine, and these researches, though at present incomplete, are being actively pursued in various directions.

The modern uses of the cinchona alkaloids may be divided roughly under four headings

1 In small doses as a tonic and stomachic

2 As an antimalarial remedy, acting as an agent that will destroy the malaria parasite. The most comprehensive work hitherto published is that of Major A C. MacGilchrist (Indian Journal of Medical Research, 1914, 2, 315 and 516; 1915, 3, 1)

About one-third of the quinne ingested undergoes cleavage of the molecule in the body, but whether it is this decomposed portion or the remaining two-thirds of the dosage which possesses its antimalarial properties appears not as yet to have been determined. The cleavage would appear to be the breaking off of the 'loiponic ring' (or second half) from the quinoline ring portion of the molecule, the former yielding some such nucleus, as is described by Koenigs under the name of ethyl-quinuchidine

$$\begin{array}{c|c} CH \\ H_2C & CH_2 & CHC_2H_5 \\ & CH_2 & \\ H_2C & CH_2 \\ \end{array}$$

It would appear that the methoxy group or  $-\mathrm{OCH_3}$  side chain of quinine and hydroquinine render the specific action greater than those alkaloids, such as einchonine and cupreine, which do not possess it following up this side chain theory, efforts have been made to prove that by increasing the value of this group as methyl hydrocupreine, or its more highly alkylated homologues, the antimalarial effect is still further increased. Results have not so far been very conclusive, and the point remains to be established in the future.

Another channel in which much investigation has recently been conducted is in the quinotoxin group of compounds. These have for many years been administered as a febrifuge under the name of quinoidine, which is really a mixture of quincine, cinchonicine, diquinicine, and dicinchonicine. The present researches, which are being carried out with the individual bases obtained in as pure a state as possible, depend upon the theory that the quinicine or quinotoxin molecule, for instance, owing to the opening up of the 'loiponic ring,' possesses

a greater specific action than the corresponding 'saturated' molecule

These researches are still incomplete, but if the fact were established that, owing to its increased specific action, quinotoxin could be administered in a very much reduced desage as compared with quinne, then owing to the comparative simplicity of its preparation, it might well form a formidable rival to the natural alkaloids of commerce

3. In treatment of Pneumonia —Most of the work in this direction appears to have followed the 'side-chain theory,' and to have been carried on with ethyl hydrocupreine which was before the war manufactured in Germany and sold in small quantities on this market under the registered name of 'Optochin' No work has, up to the present, been published to show whether a further increase in the side chain to the higher homologues renders this treatment more effective

Apparently the results obtained up to August, 1918, do not show any marked superiority over the ordinary commercial alkaloids.

4 As an Antiseptic in the treatment of Wounds—A considerable amount of work has been done in Germany on the use of the higher alkylated homologues of the hydrocupreme series as a wound treatment in war surgery Prof Klapp, following on the work of Morgenroth and Tugendreich (Pharm Zeit Nov 7, 1917, 603; and Deut Med Woch 1917, 44), claims that, by lengthening the side chain as far as the isoctyl member of the hydrocupreme series, he has obtained an antiseptic superior to any known wound dressing. He claims that this compound has 20 times the power of quinine, and 10 times that of ethyl-hydrocupreme, in the destruction of streptococci.

These comparative results are quoted as having been obtained 'in vitro,' and must therefore be accepted with reserve, and time alone will show whether this treatment will really be found to be superior to the far simpler and cheaper Dakin's solution or the highly satisfactory 'chloramine-T' treatment.

Although the precise relation between the constitution of the alkaloids and their therapeutic value has clearly not been established, enough work has been done to indicate a possible future for derivatives of einchona alkaloids either of the 'unsaturated' (quinotoxin) or of the 'increased side chain' type, which from the purely scientific and medical point of view, might be of even greater importance than the present universal use of the parent alkaloids as one of our few standard remedies

General Properties of the Cinchona Alkaloids.—These bases are mostly white crystalline nonodorous bodies, with a bitter flavour and strong basic properties, some being sufficiently powerful to displace ammonia from its salts. On heating strongly they decompose, evolving fumes of a colour resembling iodine vapour, and with a quinolinic odour. They are very slightly soluble in water, but are fairly readily soluble in organic solvents, there being, however, notable exceptions. Acids dissolve these alkaloids forming two series of salts, neutral and acid, of the general types B<sub>2</sub>H<sub>2</sub>SO<sub>4</sub> and B·H<sub>2</sub>SO<sub>4</sub>. Even more acid salts of the type B·2H<sub>2</sub>SO<sub>4</sub> are

obtainable in some cases. These salts are generally white crystalline bodies, of very varying solubility in solvents generally, and with a very bitter taste. Solutions of the salts in water when treated with alkalis precipitate the alkaloid, with the exception of cupreme, which, being phenolic in character, is soluble in alkalis.

A strong blue fluorescent solution is obtained with many of these alkaloids when dissolved in dilute sulphuric acid, viz quinnine, quindine, hydroquinne, hydroquinneme. This is a very delicate test which is best seen by reflected light. The haloid salts do not show this fluorescence, which is also destroyed by adding a haloid salt or hyposulphite to the sulphuric acid solution.

All the cinchona alkaloids have, in solution, a marked rotatory influence on polarised light, which is affected by the temperature of the solution, the solvent employed, the concentration, and the amount of acid present With mineral acids the maximum rotation is observed when the acid is present in little more than the amount to form the acid salt, but with organic acids a much greater amount of free acid is required.

When certain of these bases are treated in dilute solution with chlorine- or bromine-water, followed by ammonia added drop by drop, a deep green colouration or precipitate is obtained. This is known as the thalleroquin test (thallos, a young twig), and is given by quinine, quinidine, quinicine, hydroquinidine, cupreine, and diquinicine.

The emchona bases are di-tertiary amines forming mono- and di-alkyl rodide additive compounds

The hydrochlorides of the bases form double compounds with several metallic chlorides, of which the chief are. (a) the chlorozincates of the general formula B,2HCl,ZnCl<sub>2</sub>; (b) the chloromereurates B,2HCl,HgCl<sub>2</sub>, (c) the chloro-aurates B,2HCl,AuCl<sub>3</sub>; (d) the platinochlorides, of which there are two types, B,2HCl,PtCl<sub>4</sub> and B<sub>22</sub>2HCl,PtCl<sub>4</sub> Of these (c) and (d) have been much used in the determination of the molecular weights of the bases

The cinchona alkaloids are nearly completely precipitated from solution by tannic, pieric, or phosphotungstic acid, potassium mercuric iodide (Mayer's reagent), and potassium bismuth iodide (Dragendoiff's reagent)

Methods of Extracting the Alkaloids of Cinchona Barks —Before extracting the bark it must be treated in such a manner as will ensure the alkaloids being in a suitable condition for extraction For this purpose the back must be in a finely powdered condition, the finer the powder the more rapid and thorough being the The whole of the bark should be extraction ground to pass a sieve with at least forty meshes to the linear inch This point is of special importance in assaying, as cases are known in which the harder portions of the bark (which might be rejected as gruffs if the whole were not finely ground) are richer in alkaloids than the softer parts. The next stage in the process is that of liberating the alkaloids from their combinations with the natural acids of the bark. in order to render them soluble in the solvent employed. This is effected by moistening the

bark with a dilute alkali, or by mixing it with lime and damping with water, and allowing to stand for some hours The whole may then be dried, powdered, and extracted with chloroform, ether, petroleum spirit, or a mixture of these; or the damp bark may be extracted direct with petroleum, amyl alcohol, benzene, naphtha, or mixtures of these Thus Landrin (Compt. rend. 1889, 108, 750) and Zebel (Chem. Zeit. 1891, 15, 735) describe processes using petroleum; the Brit Pharm. 1914, directs the use of a mixture of 3 volumes of benzene and 1 of amyl alcohol; while the U.S Pharm 1916, employs a mixture of 2 volumes of ether and 1 volume of chloroform On the large scale the solvent employed will naturally be the cheapest, while on the small scale, where an efficient reflux extraction apparatus can be used, chloroform or ether are convenient

Zebel (1 c) describes the commercial process thus: The finely powdered bark is ground to a thin paste with lime, sodium hydroxide, or sodium carbonate, and extracted with warm paraffin oil On standing the oil separates, when it is run off and shaken with sulphuric acid: this acid solution is boiled, and while boiling is neutralised (to litmus paper) with sodium carbonate and allowed to cool Quining sulphate crystallises out on cooling, while cinchonidine, einchonine, and quinidine remain in solution as sulphates The quinine sulphate is purified by recrystallisation from water after treating with animal charcoal The motherliquor containing the other alkaloids is treated with sodium hydroxide and the precipitate treated with dilute alcohol Of the three bases precipitated, quinidine and cinchonidine are dissolved by the alcohol, while cinchonine is left behind: the two former can then be separated by means of their neutral tartrates, that of quinidine being considerably the more soluble

Landrin (lc) gives this description of the small scale extraction 300 grams of bank are powdered and passed through a No 40 sieve, then mixed thoroughly with 1 litre of milk of soda-line containing 75 grams of quickline and 75 grams of solution of sodium hydroxide (sp gr 1 375). To this mixture, diluted if necessary by adding a little water, 2 litres of petroleum are added, heated to 100° for 20 minutes, with constant agitation; the oil is then decanted, and the extraction repeated with a second 2 litres. The 4 litres of extract are washed twice with 75 cc of 10 pc sulphuric acid added to 150 cc of water, and a third time with one-third of these quantities.

The British Pharmacopera, 1914, method is as follows: 10 grams of the bark in No. 60 powder are mixed with 6 grams of slaked lime, and the whole moistened with 22 c.c. of water and allowed to stand for 2 hours. This is transferred to a flask provided with a reflux condenser and boiled with 130 c.c. of benzolated amylic alcohol (benzene, 3 volumes; amyl alcohol, 1 volume) during 30 minutes. The liquid is filtered off and the treatment repeated twice, the mixture of bark and lime being finally transferred to the filter and washed with the solvent till exhausted

The U.S. Pharmacopecia (9th revision, 1916) process is: Introduce 5 grams of einchona in No 40 powder into a 500 c.c. flask, and add

200 c e of a mixture of chloroform and ether (1:2) Stopper the flask, shake it well, and let it stand 10 minutes. Then add 5 c c of ammonia water, shake the flask frequently for 1 hour, and let it stand from 8 to 10 hours. Now add 10 c c of distilled water, shake the mixture thoroughly, and when the drug has settled decant 160 c c of the solution, representing 4 grams of cinchona. Filter through a pledget of purified cotton into a separator, and rinse both cylinder and cotton with ether. Completely extract the alkaloids from the chloroform-ether solution by shaking out repeatedly with weak sulphuric acid.

An older method of extracting the bark, due to de Vrij, was that of treating the powdered bark with a large excess of milk of lime, diying down and extracting with alcohol But this has been entirely given up owing to the great amount of colouring matter extracted with the alkaloids, and the consequent difficulty in

purification

Assay of Cinchona Barks - The accurate assay of a bark is much more difficult than appears at first sight owing to the obstinacy with which one alkaloid will form more or less definite compounds with the others when crystallising as alkaloid or salt, and also on account of the changes in its solubility in solvents when varying quantities of other alkaloids are present For these reasons a process of assay giving good results when, for instance, a Ledger bark, in which quinine is the main constituent is tested, will not be accurate for a Succirubia bark in which the cinchonidine and cinchonine usually exceed the quinine in amount knowledge of the type of bark being assayed, and of its alkaloidal contents, is therefore of great importance, the experienced analyst being able to foretell the state of purity of his various precipitates and so regulate his analysis

A 'full' analysis of a back usually consists of the estimation of the quinine, quinidine, cinchonidine, cinchonine, and total amorphous alkaloids To obtain satisfactory results a quantity of alkaloid of about 5 grams should be available It is desirable to start the assay with the estimation of the total alkaloids, as the subsequent work is simplified by so doing and the result is a useful check on the estimation of the individual alkaloids. For this purpose it is only necessary to evaporate the extract obtained by any of the methods described under the previous heading to constant weight; or, m the case of the paraffin and sulphure acid extraction, to make alkalme with sodium hydroxide, wash out with chloroform, and evaporate to dryness Dry to constant weight at 100°-110°. Having obtained the weight of total alkaloids, add 3 cc of 10 pc sulphuric acid for every giam of alkaloid, and warm on the steam-bath till completely dissolved. Then add 80 c c of distilled water for every gram of alkaloid, boil, and while boiling neutralise exactly to litmus paper with dilute ammonia solution (10 pc) On cooling for 2 or 3 hours with occasional stirring practically the whole of the quinine will crystallise out as sulphate, owing to it being almost insoluble in ammonium sulphate solution. Filter through counterpoised filter papers on the filter-pump, press the crystals down well, pumping as dry as possible,

and following with two washes of cold water each about equal in volume to the bulk of the precipitate. The filter papers and crystals are now dried to constant weight at 100°. The weight of anhydrous quinine sulphate found multiplied by I 18 gives the amount of rough hydrated quinine sulphate, or by 0 868 the amount of rough quinine alkaloid. This quinine sulphate is neither white nor pure, being usually contaminated with colouring matter and cinchonidine sulphate, and requires further treatment, for which see under 'Quinine Sulphate.' But with the present-day pure Ledgeriana barks the weight may be taken as representing the true amount of quinine sulphate, as in some thousands of analyses of this back by the present writers the amount of quinine sulphate left in the mother liquor almost exactly compensates for the impurities in the crystals weighed. It is, however, a safeguard in all cases to test for any undue amount of emchandine as described under 'Quinine Sulphate

To the mother liquor and washings from the crystals add about 20 grams of Rochelle salt, dissolved in 30 cc of hot water, and still occasionally during 2 or 3 hours Cinchonidine is completely precipitated as tartrate collected on counterpoised filter papers at the pump, washed with three lots of cold water each about equal in bulk to the precipitate, which is then dried and weighed. This weight multiplied by 0.797 gives the cinchonidine alkaloid Should the bark contain very little cinchonidine, and no tartrate be obtained in the 3 hours' standing, it is best to leave the solution for a full 24 hours to ensure complete precipitation. The filtrate from the cinchonidine tartrate is now evaporated to one-half, half its volume of methylated spirit and 5 grams of neutral potassium iodide dissolved in 5 c c of water are added, and the liquid left for at least 2 hours with frequent stirring The quinidine hydrodide is filtered off at the pump on a counterpoised filter paper, washed with a little 50 p c methylated spirit, dired and weighed 0 0008 gram is added to this weight for each 1 cc of filtrate and washings, which total multiplied by 0.7186 gives the quinidine alkaloid filtrate is made alkaline with sodium hydroxide and shaken with other (fice from alcohol) in a separator, the aqueous portion drawn off and twice again shaken with small quantities of ether, the ether extracts bulked, allowed to stand I hour, and the emchanine filtered off on counterpoised papers at the pump, two small washes with ether being given. The cinchonine is dried and weighed, and to its weight is added 0 0027 gram for each 1 cc of filtrate and washings The ether evaporated to dryness and constant weight gives the total amorphous alkaloids, from which should be subtracted the weights added on as corrections for quinidine and cinchonine.

The above method is made up of the best points of the many published methods of assay, and will be found to give good results in inexperienced hands For other methods, see Allen's Commercial Organic Analysis, 1912, vol. vi 490-495.

The British Pharmacopæia, 1914, specifies that the bark (C. succirubra) 'when used for official purposes other than the preparation of

alkaloids or their salts, shall yield 5-6 p.c. of total alkaloids, of which not less than half shall consist of quinne and cinchondine. The method used is that of extracting the amylated benzene solution of alkaloids (see above) with dilute hydrochloric acid, neutralising the acid solution with ammonia, and precipitating the quinne and cinchonidine together as tartrates by the addition of Rochelle salt. The remaining alkaloids are then precipitated from the filtrate by ammonia.

The United States Pharmacopora, 1905, prescribes 5 p.c. of total alkaloids and 4 p.c. of ether-soluble alkaloids (i.e. quimine, quimine and cinchonidine). This is carried out by dividing the acid solution of the alkaloids into two equal portions, shaking one with ammonia and chloroform-ether mixture to obtain total alkaloids, and the other with ammonia and ether for other-soluble alkaloids. The USP, 1916, omits the 'ether-soluble alkaloid' test

The German Pharmacopara (1910) method consists of estimating the total alkaloids by extracting the chloroform-ether mixture (1-3) with decinormal acid in excess, and titiating back the excess of acid over the neutral point with decinormal alkali, using alcoholic hamatoxylin solution as indicator. 65 p.e. of total alkaloids are required

Titiation methods are hable to senious errors and cannot be recommended, but can be employed as useful rough checks on analysis Generally speaking, the free alkaloids are neutral to phenol-phthalem, the neutral sulphates (B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub>) are neutral to litmus, brazil wood, lacmoid, cochineal, and hiematoxylin, but alkaline to methyl-orange, which roughly indicates the bisulphate point (B H<sub>2</sub>SO<sub>4</sub>) (Allen, Analyst, 1896, 21, 85; Farr and Wright, Pharm J. 1894, 25, 124, Messner, Zeitschangew. Chem. 1903, 16, 444, Engelhardt and Jones, Pharm J. 1910, 30, 236, Katz, Bei Deut, pharm Ges. 1910, 20, 316).

Oudemans (Annalen, 1876, 182, 33) has strongly recommended the polarimetric assay of the einchona alkaloids, and a good deal of controversy has raged around the subject, the majority of the evidence appears to be against the method, owing to the difficulty of obtaining the bases in a pure enough condition for accurate polarimetric reading (Hesse, Annalen, 1875, 176, 205, 1876, 182, 128, Koppeschar, Zeitsch anal. Chem 1885, 24, 362; Léger, J Pharm Chim 1904, 19, 427).

Quinine and its Derivatives and Salts—Quinine alkaloid is usually prepared by precipitating a solution of quinine hydrochloride with an alkali or ammonia, washing free from chloride and drying at a low temperature. The commercial article is a soft white granular odourless powder with micro-crystalline structure, and contains up to 11 pc H<sub>2</sub>O (dihydrate). It crystallises from ethereal solution below 10° with 3H<sub>2</sub>O (14.3 pc H<sub>2</sub>O), but the ethereal solution evaporated gives a resmoid mass containing 1H<sub>2</sub>O (most other cinchona alkaloids give crystalline residues from ether). Alcoholic or ammoniacal solutions deposit needles with 3H<sub>2</sub>O, which rapidly efficience in the air above 15° giving a dihydrate. This is the composition of the commercial article, although most pharmacopoeias require the trihydrate. The

trihydrate loses all its water over sulphuric acid. It melts at 57°, and in estimating the moisture it is therefore best to dry for some hours at 40°-50°, then gradually raise the temperature to 100°. In this way no melting takes place, and the dehydration is rapid. Other hydrates containing 1, 8, and 9H2O have been described Anhydrous pure quinine melts at 174.9° It is a strong base displacing ammonia from its salts in hot solution, and neutralising the strongest acids. The pure base has a bitter taste, which develops slowly; the commercial article is more bitter on account of the presence of impurities which give more rapid solution Ammonium and calcium salts notably increase the solubility of the alkaloid in water, while the alkalıs render it less soluble. The anhydrous alkaloid is soluble in 1750 parts of water, in 3450 parts of potassium hydioxide (5 p c), and in 1890 parts of ammonia (10 pc) at 25°C The trihydrate is soluble in 0.8 parts of alcohol, 11 parts of chloroform, 19 parts of ether, 212 parts of glycerm, and 166 parts of benzene at 25°C Quinine is Levorotatory, the trihydrate having in 97 p c alcohol  $|a|_{D}^{15} = -145 \ 2^{\circ} + 0 \ 657c$ The anhydrous alkaloid in ether has  $[\alpha]_{D}^{15}$  $=-1587^{\circ}+1911c$ , where  $\epsilon=15-60$ 

The chief impurities likely to be met with in the commercial article are chlorides and sulphates of the alkalis or ammonia, lime salts, hydroquinine, emehonidine, and emehonine I gram should dissolve in a mixture of 6 cc. absolute alcohol and 3 c c ether to give a solution remaining bright on cooling (absence of einchonidine and cinchonine, and inorganic salts) Hydroquinine, which is liable to occur up to 2 pc, may be tested for as given under 'Quinine Sulphate'

Quinine is distinguished from the other alka loids by its ready solubility in ether, fluorescent solution in dilute sulphuric acid (detectable to the extent of 1 part in 500,000), by the thallerogum test, its insoluble neutral sulphate, taitrate and chromate, and by its iotation

The following are the more important derivatives of quinine — Quitenine  $C_{19}H_{22}O_4N_2$  or

C<sub>17</sub>H<sub>18</sub>O OCH<sub>3</sub> COOHN<sub>2</sub>

(Skraup, Annalen, 1879, 199, 348). This feeble base, together with formic acid, is obtained when quimme is oxidised with permanganate in acid solution at 0°, and bears the same relation to quinine as einchotenine does to einchonine (see 'Constitution') The quitenine is carried down by the oxide of manganese, from which it may be extracted with 40 pc alcohol concentration and filtration quitenine comes out in thombic prisms containing 4H2O, slightly soluble in water, insoluble in ether, but dissolves in acids and alkalis. Quitenine is identical with Kerner's dihydroxyquinine (Zeitsch Chem 1869, 5, 593) It gives the thalleroquin reaction, and dissolves in dilute alcoholic sulphuric acid with fluorescence It is laworotatory, giving  $[\alpha]_D = -142^\circ$ ; mp of anhydrous base 228° It gives a sulphate B<sub>3</sub>·2H<sub>2</sub>SO<sub>4</sub>,15H<sub>2</sub>O in fine needles Heated with sulphuric acid it is converted into an isomeric amorphous dark red base, quitenicine, very soluble in water. Its

constitution is that of quinine in which the vinyl is replaced by a carboxyl group. Heated with hydrodic acid (sp.gr. 1.7) its methyl group is

eliminated, and a new base, quitenol, is obtained

Quitenol C<sub>17</sub>H<sub>18</sub>O OH COOH N<sub>2</sub> crystallises in fine needles containing 1H2O, very slightly soluble in water, alcohol, or ether. It dissolves readily in acids and alkalis, being precipitated from the latter by CO2. It gives the thalleloquin reaction, and a red colouration in hydrochloric acid solution with ferric chloride Quitenol bears the same relation to quitenine, that cupreme does to quinine (Bucher, Monatsh. 1893, 14, 598).

Quininic acid C<sub>2</sub>H<sub>5</sub> OCH<sub>3</sub> COOH N (6-methoxy-4-carboxyquinoline) This acid is obtained on oxidising quinine with chromic acid, being the p-methoxy derivative of emchanine acid derived from emchanine under similar treatment (Skraup, Ber 1879, 12, 1104, Comstock and Komgs, Ber 1884, 17, 1984, 1886, 19, 2853) It forms yellowish prisms, almost insoluble in water, very soluble in acids and alkaloids. It is slightly soluble in absolute alcohol, giving a beautiful blue fluorescence, which is destroyed by water or acids with fuming hydrochloric acid under pressure it

loses its methyl group, yielding a phenolic body xanthoquinine acid C<sub>9</sub>H<sub>5</sub> OH COOH N

Meroquinene C<sub>6</sub>H<sub>10</sub> CH CH<sub>2</sub> COOH NH (v

Constitution (Skraup, Monatsh 1903, 24, This body forms small colourless needles, m p 223°-224°, very soluble in water and acids, almost insoluble in alcohol It is an interesting example of a body in which the acidity of the carboxyl group is neutralised by the alkaline properties of the remainder of the molecule  $[\alpha]_D$  in 10 pc aqueous solution  $+27.9^{\circ}$ . It readily forms esters On oxidation with chromic acid it gives cincholorponic and lorponic acids.

Cincholoiponic acid C<sub>6</sub>H<sub>10</sub>(COOH), NH This acid is derived from meroquinene by the oxidaacids and alkalis It forms rhombic prisms containing 1H<sub>2</sub>O, melting at 126°-127°, and when anhydrous, at 225°-226° It is a dibasic acid and a secondary base (Skraup, Monatsh 1896, 17, 372, 1900, 21, 879) When the hydrochloride of this acid is heated with resorein and concentrated sulphure acid, or zinc chloride, a fluorescein is formed, which indicates that the two carboxyls are in an ortho position to each other (Muller and Rohde, Ber. 1894, 27, 1188). A great deal of work has been done on the constitution of this acid, for upon it depends the proof of the constitution of the 'second' half of the molecule of the cinchona alkaloid later work on its stereoisomerides, see Konigs, Ber 1901, 34, 4336; Wohl, Ber 1907, 40, 4693, 4711, 1909, 42, 627 Loiponic acid  $C_5H_8(COOH)_2NH$  (hexahydro-

pyridine-3 4-dicarboxylic acid) v. 'Constitution' Prismatic crystals, m p 259°-260°, fairly soluble in water, easily soluble in alkalis and mineral

n water, cashy soluble in alkalis and mineral acids (Skraup, Monatsh 1896, 17, 366)

Quinine chloride C<sub>20</sub>H<sub>23</sub>(N)<sub>2</sub>Cl This body is formed when dry quinine hydrochloride is treated with phosphorus pentachloride in chloroform solution, the hydroxyl group of the quinine being replaced by chlorine It has basic properties, melts at 151°, gives the thalleoquin reaction, but does not fluoresce in acid

solution (Comstock and Konigs, Ber. 1884, 17, 1984; 1885, 18, 1219, 1892, 25, 1545) On treatment with alcoholic potash hydrochloric acid is eliminated with the formation of quinene.

Quinene  $C_{20}H_{22}ON_2,2H_2O$  forms large thombre crystals melting at  $81^\circ-82^\circ$ . On hydrolysis with 25 p c phosphoric acid it yields meroquinene and p-methoxylepidine (Konigs, Ber 1890, 23, 2669, 1894, 27, 900; 1895, 28, 3143, 1896, 29, 372). On heating with HBi (sp.gr 149) quinene yields apoquinene (p-oxyapocimeheno), thus.

 $\begin{array}{c} {\rm C_{20}H_{22}ON_2 + HBr + H_2O} \\ {\rm = C_{10}H_{19}O_2N + CH_3Br + NH_3} \end{array}$ 

Apoquinene  $\rm C_{19}H_{19}O_2N$  forms crystals melting at 246°, very slightly soluble in water, benzene, or ether, readily soluble in alcohol its solutions in sodium hydroxide or dilute hydrochloric acid are yellow (Konigs and Comstock, Bei 1885, 18, 1226 , 1886, 20, 2675)

Quininone C<sub>20</sub>H<sub>22</sub>O<sub>2</sub>N<sub>2</sub> (p-methoxyem hommon) is obtained by carefully oxidising quinine (or quinidine) with chromic acid in the presence of acetic acid. Treated with introus acid it yields quininic acid, and an oxime which splits up on hydrolysis into meroquimene and hydroxylamine (Rabe, Ber 1907, 40, 3655, 1908, 41, 62, Annalen, 1909, 365, 353, 1910, 373, 85, 1911, 382, 365). Upon this hydrolysis Rabe has based his formula for the cinchona alkaloids (see p. 89).

Dhydroquinine (1,0H2,0U2N). Schutzenberger (Annalen, 1858, 108, 348) obtained an only body by the action of zinc and sulphuric acid on quinine, which he named dihydroquinine. This is not the dihydroquinine of commerce, v. Hydroquinine (Hesse, Ber. 1882, 15, 854). Pum, Monatsh. 1895, 16, 72)

Tetrahydroquinine C<sub>20</sub>H<sub>28</sub>O<sub>2</sub>N<sub>2</sub> is obtained by the action of sodium and alcohol on quinine (Lippmann and Fleissner, Monatsh 1895, 16, 630), also by electrolytic reduction (Freund and Bredenberg, Annalen, 1914, 407, 43)

Dichlorquinine  $C_{20}H_{24}N_2O_2$   $Cl_2$  is formed when quinne hydrochloride, dissolved in a mixture of acetic and hydrochloric acids, is treated with 1 molecule of chlorine. It is a colourless amorphous compound, melting at 97°, and closely resembles quinne in its chemical behaviour. Treated with alcoholic potash it loses 2HCl, yielding dehydroquinine  $C_{20}H_{22}O_2N_2$ . By the further action of chlorine on dichlorquinine, a tetra-chlorimated body  $C_{19}H_{20}O_2N_2Cl_4$  is obtained (Christensen, Ber. Deut. pharm. Ges. 1915, 25, 256)

Hydroxychlorquinine  $C_{20}H_{24}O_2N_2ClOH$  is obtained as an amorphous colourless powder, mp 110°, when quinine dissolved in cold dilute hydrochloric acid is treated with 1 molecule of chlorine dissolved in water (Christensen, bid) When 3 molecules of chlorine are employed a tri-chlorinated body  $C_{19}H_{21}O_3N_2$   $Cl_3$  is obtained

Tetrabromquinine  $\rm C_{20}H_{24}O_2N_2$  Br<sub>4</sub>, obtained by treating quinine dissolved in sulphuric acid with excess of bromine-water. It is a yellow amorphous body with an odour of bromine.

Tribromquinine ('20 H24O2N2, Br3, obtained by treating an alcoholic solution of tetrabromquinine with H<sub>2</sub>S.

Dibromquinine  $C_{20}H_{24}O_2N_2\cdot Br_2+2H_2O$  is

obtained from the mother liquor of tribiomquinine by precipitation with ammonia. Another anhydrous dibromquinine is obtained by acting on a chloroform-alcohol solution of quinine hydrochloride with bromine dissolved in chloroform (Comstock and Konigs, Ber 1892, 25, 1500)

Iodoquinines. By the action of rodine dissolved in alcohol or potassium rodide many badly defined rodoquinines have been obtained, e.g. 4B 3I, 4B 5I, B HI I, B 2HI, B HI I4 (Bauer, Arch. Pharm. 1874, 5, 392)

Hydrochloroquinine C<sub>20</sub>H<sub>25</sub>ClO<sub>2</sub>N<sub>2</sub>, obtained by treating quinine at low temperature with HCl saturated at - 17°, mp 186°-187°. Insoluble in water, soluble in ether or methyl alcohol. It shows fluorescence and gives the thalleroquin reaction. Quinine is regenerated on treatment with alcoholic potash (Comstock and Konigs, Ber. 1887, 20, 2510, 1892, 25, 1539). Hesse obtained this body by heating quinine with HCl (sp.gr. 1189) at 85° (Annalen, 1868, 147, 241).

Hydrobromoquinine C<sub>20</sub>H<sub>26</sub>BrO<sub>2</sub>N<sub>2</sub> is obtained by treating quinne hydrobromide with HBi saturated at - 17°C, at a low temperature for 5 days. It is soluble in ether, but its sulphuric acid solution does not fluoresce (Connected and Konge, Rev. 1896, 20, 203).

(Comstock and Komgs, Ber. 1896, 29, 803) Hydroiodoquinine ( $^{\prime}_{20}$ H $_{28}$ IO $_2$ N $_2$ , obtained by treating quinne with H1 (sp gr. 19) on the water-bath for 1 hour Crystals melt at 150°–155° Soluble in ether, from which crystals containing I(C $_2$ H $_8$ ) $_2$ O are obtained. Its solution in sulphure acid is fluorescent, and gives the thalleroquin test. Heated with alcoholic potash it yields a mixture of quinne, pseudoquinne, isoquinne, and inquine (Skraup, Monatsh 1893, 14, 431, 446, Lippmann and Fleissner, ibid. 1891, 12, 327, 372, 1892, 13, 429, 1893, 14, 553)

Apoquinine  $C_{10}H_{20}(OH)_2N_2$  is obtained by heating quinine under pressure to 145° with HI (sp gr 1 25-135). It is an isomeride of cupteme, which is also converted to apoquinine by heating under pressure with HCl (sp gr 1 125). It crystallises in large needles of m p 210°, and  $\lfloor \alpha \rfloor_D^{15} = -178$  1° for a 2 p c solution in 97 p c alcohol. The acid oxalate  $B_2 \, 3C_2H_2O_4$  and iodohydrate  $B_2 \, 2H_1$  crystallise readily. It gives similar addition products to quinine with the haloid acids, hydroiodoapoquinine yielding on treatment with alcoholic potash the isomer isoapoquinine melting at 176° (Lippmann and Fleissner, Monatsh 1891, 12, 332, 1893, 14, 553; 1895, 16, 33, Mauther, Ber. 1895, 28, 1972)

Derivatives Isomeric with Quinine — Niquine  $C_{19}H_{24}O_2N_2$  (see Hydroiodoquinine) This alkaloid behaves very similarly to quinine, forming similar salts, addition products, &c. Its sulphuric acid solution fluoresces, and it gives the thalleloquin reaction. It is soluble in alcohol, ether, chloroform, or benzene, slightly soluble in boiling water, from which crystals containing  $2H_2O$  are obtained,  $\lfloor \alpha \rfloor_D = -129~2^\circ$  for a 1 p c solution in 98 p.c. alcohol (Skraup, Ber. 1892, 25, 2911, Monatsh. 1893, 14, 428, Lippmann and Fleissner, Monatsh. 1893, 14, 426, 553) By treating the bihydrioidde of hydroiodo-niquine with alcoholic potash inquine is

in part reformed together with an isomer isoniquine.

**Isoniquine**  $C_{19}H_{24}O_2N_2$  crystallises from alcohol-in the form of benzoic acid and melts at  $208^\circ-209^\circ$ . It shows fluorescence in sulphuric acid, and gives the thalleroquin reaction (Skraup, Lippmann and Fleissner,  $l\,c$ )

**Pseudoquinine**  $C_{20}H_{24}O_{2}N_{2}$ , an isomer of quinine ( $\theta$  hydriodoquinine), forms prisms melting at  $190^{\circ}-191^{\circ}$  It is not very soluble in other, readily soluble in alcohol,  $[\alpha]_{D} = -164 \ 4^{\circ}$ 

for a 1 p c solution in 98 p c alcohol

Isoquinine  $C_{20}H_{24}O_2N_2$ , an isomer of quinine (v hydroiodoquinine), crystallises in needles, melting at  $185^\circ$ – $186^\circ$ , and having  $[\alpha]_D^{27}=-186^\circ$ 8° for 0.9 p.c. solution in 97 p.c. alcohol. It is readily soluble in ether and benzene, very slightly soluble in boiling water, and reacts generally slike quinine (Skraup, Monatsh. 1893, 15, 433, 1894, 16, 37). Hesse has also named a base isoquinine, which he obtained by the action of concentrated  $H_2SO_4$  on quinine at  $120^\circ$ – $130^\circ$ , or by heating with glycerin at  $180^\circ$ – $210^\circ$  (Annalen, 1874, 174, 340, 1875, 178, 253, 1888, 243, 149, 1893, 276, 103)

Quinicine (qumotoxin) C<sub>20</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub>, an isomer of qumine, was first prepared by Pasteur in 1853 by heating quinine bisulphate with a little water at 160°, and afterwards isolated from quinine tails by Howard (Chem. Soc Trans. 1871, 24, 61, 1872, 25, 1011). It forms a reddish-yellow amorphous mass melting at about 50°, nearly insoluble in water, soluble in chloroform, ether, and alcohol, the latter solution rapidly absorbing CO<sub>2</sub> from the air. It forms salts of which the oxalate B<sub>2</sub> C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>,9H<sub>2</sub>O bitartiate B C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>,6H<sub>2</sub>O, sulphocyanide

B HCNS, 1H,O

hydriodide B HI, H2O, chloroplatmate

B H, PtCl, 2H, ()

and aurichloride are crystalline. Hesse also obtained a crystalline sulphate  $B_2$   $H_2SO_4$ ,8 $H_2O$ . It is completely precipitated from its solutions by excess of potassium thiocyanate as an oil rapidly becoming crystalline.  $[a]_D^{15} = +44$   $1^\circ$  for a 2 p c. solution in CHCl $_3$ . It gives the thalle-ioquin reaction, but does not show fluorescence in sulphuric acid solution. It is soluble in ammoniacal salt solutions, from which ether extracts it (separation from other cinchona alkaloids). (Hesse, Annalen, 1875, 178, 213; Miller and Rohde, Bei 1895, 28, 1058; Skraup, Monatsh 1899, 20, 573, Howard and Chick, J. Soc. Chem. Ind. 1909, 28, 53; Pharm. J. 1917, 99, 143)

Quinine Salts — Quinine sulphate

 $B_2 H_2 SO_4, 7\frac{1}{2} H_2 O$ 

This salt occurs as white light silky needles, odourless, but with an intensely bitter taste. The hydration is rather indefinite, being recognised as  $8\mathrm{H}_2\mathrm{O}$  by the French Pharmacoperu (=16 17 p.c.),  $7\frac{1}{2}\mathrm{H}_2\mathrm{O}$  by the B P. (=15·3 p.c.), and  $7\mathrm{H}_2\mathrm{O}$  by the USP. (=14·5 p.c.) It is completely dehydrated at  $100^\circ$ . Exposed to the air it loses water until  $2\mathrm{H}_2\mathrm{O}$  remain (=4·8 p.c.), the same amount being absorbed if the fully exsiccated salt is exposed to moist air. The chemically pure salt crystallises in large heavy

needles, the lighter character of the commercial article being due to the presence of a little hydroquinine and cinchonidine, or by being crystallised slightly alkaline or from solutions containing ammonium sulphate. It is soluble m 740 parts of water at 15°, in 30 parts at 100°; in 65 parts of alcohol at 15°, in 6 parts at boiling-point, in 24 parts of glyceim at 15°, 1000 parts of chloroform at 15°, leadily soluble (1 m 7) in a mixture of chloroform, 2 parts, absolute alcohol, 1 part. The sulphates of the other cinchona bases and morganic impurities are insoluble in this mixture, which is therefore a good test for the purity. It is much less soluble in water containing magnesium, sodium, or ammonium sulphate, but more soluble if calcium, sodium, or ammonium chloride, or potassium nitiate or chlorate be present. Quinine sulphate has  $[\alpha]_D^{17} = -157 \ 4^\circ$  in absolute alcohol at a concentration of 22 pc of the 7½H2O salt Alpc solution of the anhydrous sulphate m excess of dilute  $H_2SO_4$  has  $[a]_0^{15} = -235^\circ$ . A dilute  $H_2SO_4$  solution shows a strong blue fluorescence, detectable with about 1 part of quinine in 500,000 On treating a neutral solution of the sulphate in cold water with a slight excess of chlorine- or bromine-water added drop by drop, agitating for 20 seconds, and then adding ammonia drop by drop until faintly alkaline, a deep green colour is produced, or if more than I in 1000 of quimine is present a green precipitate is obtained (thalle-loquin reaction). On neutralising the alkaline solution with dilute acid the colour changes to blue, which with more acid becomes red, reverting to green on again making alkaline By a modified procedure Ramsden and Lipkin (Annals Trop Med 1918, 11, 443) are able to detect 1 part of quinine in 400,000 Neutral solutions of quimine sulphate in water give the following almost insoluble precipitates with potassium chromate  $B_2 H_2 Cr O_4, 2H_2 O$ , soluble 1 in 3150, with potassium oxalate

 $B_2, C_2H_2O_4, 6H_2O$ 

soluble 1 in 1400, with potassium picrate  $B_2$ ,  $C_8H_3N_2O_7$ , soluble 1 in 3400, with sodium salicylate B,  $C_7H_6O_3$ ,  $H_2O$ , soluble 1 in 2100, with tannic acid B,  $(C_{14}H_{10}O_9)_3$ ,  $8H_2O$ , soluble 1 in 2000, with Rochelle salt  $B_2$ ,  $C_4H_6O_8$ ,  $H_2O$ , soluble 1 in 950, all at 25° These are the solubilities of the respective precipitates in water; the solubility is less in excess of the reagent

On adding alcoholic iodine solution to an alcoholic solution of quinine acidined with sulphuric acid an iodosulphate of quinine  $B_4(H_2SO_4)_3(HI)_2I_4,6H_2O$ , called after its discoverer, Herepathite, is thrown down as a reddish precipitate (Ann Chim Phys 1854, 40, 249) It crystallises from strong alcohol in tabular crystals, green by reflected light, and red by transmitted light, almost insoluble in cold water or dilute alcohol, soluble I in 1000 of hot water, but much more soluble in acidified alcohol. By a special method of concentration, Ramsden and Lipkin have been enabled to identify as little as I part of quinine in 4,000,000 of urine, or in 20,000,000 of water by the herepathite reaction under the micropolariscope (Anpals, Trop. Med. 1918, 11, 444). Jorgensen

(J pr. Chem. 3, 145; 14, 213, 256; 15, 65) has prepared whole series of salts similar to herepathite, having varying quantities of acid, iodine, and water.

Most of the above-mentioned insoluble salts have been proposed as means for the estimation of quinine in mixtures of the cinchona alkaloids, but none have stood the test of time on account of the separations not being at all perfect, others of the alkaloids being co-precipitated

Estimation—Two classes of products will be dealt with here (a) the rough quinine sulphate obtained as described under 'Assay of Cinchona Barks,' and (b) commercial quinine sulphate The only alkaloidal impurities likely to occur in either are cinchonidine and hydrogumine. Of these two, hydroquinine is easily and rapidly estimated, but the estimation of cinchonidine is a much more difficult matter For instance, it might be thought that advantage might be taken of the different solubilities of quinine sulphate (1 in 740) and einchonidine sulphate (1 in 92) in water at 15° But it is found that on dissolving a quinine sulphate containing cinchonidine in water at 100°, and allowing the solution to cool to 15°, the majority of the emchanidine will crystallise out again with the quinne. As many as four recrystallisations will be necessary to remove the whole of the emchanidine from the quinine. If now the mother liquois be combined and evaporated to dryness on the steam-bath a product is obtained containing about one-fifth of the original quinine together with all its impurities. If this salt be now shaken into purified ether and aqueous ammonia (sp gr 0 959) and cooled down to 0° for 24 hours, most of the anchonidine will crystallise out at the layer of the two liquids, and may be filtered off, dried, and weighed. A modified form of this test, due to Prunier (J Pharm Chim. 1891, [v.] 23, 163 and 265), was the official method of the BP. 1898, but being a very tedious process is no longer employed. Another method consists in pre-cipitating a neutral solution of the quinine sulphate with Rochelle salt, drying and weighing the combined quinine and cinchonidine tartrates, and taking the polarimetric reading still employed in some laboratories, and usually gives fairly accurate results in experienced hands. But it must be remembered that as the optical relations of the alkaloids are so diverse that should even a small percentage of hydroquinine taitrate be co-precipitated it will lead to inaccurate results. The temperature of the reading is also very important, as well as the concentration of the solution (Oudemans, Annalen, 1876, 182, 33, Hesse, ibid. 128; Koppeschar, Zeitsch anal. Chem 1885, 24, 362; Hooper, Pharm. J. 1886, 17, 61; Jung-fleisch, J. Pharm. Chim. 1887, 15, 5, Léger, ibid. 1904, 19, 427; Montemartini and Bovini, Gazz. chim ital. 1916, 46, 153)

De Vrij proposed a method of precipitating the quinine as neutral chromate (Arch Pharm 1887, 24, 1073), which was severely criticised by Hesse (Pharm J. 1887, 17, 585, 665; see also Paul, ibid 585, footnote).

Schaefer (Arch Pharm. 1887, 25, 64) proposed a separation by means of the insoluble neutral oxalate of quinine, but the separation is not complete.

A better method is that proposed by De Vrij (Chem Zeit. 1885, 968), and recommended by Hesse (Pharm J. 1886, 17, 486) and D. Howard (Pharm J. 1896, 3, 505), which takes advantage of the fact that when 5 grams of quinine sulphate are dissolved in 12 cc, of N/1H<sub>2</sub>SO<sub>4</sub>, and allowed to crystallise, 80-90 p c. of the quimine crystallises out as bisulphate, the whole of the emehomdine remaining in the mother liquor. The crystal is filter-pumped and washed with 3 e.c. of cold water, and the filtrate shaken into 16 c c ether (sp gr. 0721-0.728) and 3 cc of ammonia (sp gr 0.959), and left to stand for 24 hours, preferably at 0° The cinchonidine crystallises out and may be filtered off, washed with a little water saturated with ether, dried, and weighed. Hesse points out that this einchonidine is not pure, but consists of a double compound of quinne and conchonidine  $(C_{20}H_{24}O_2N_2\cdot 2C_{19}H_{22}ON_2)$ , and a portion of the cinchonidine is also soluble in the ethereal quinine liquor, these errors almost exactly compensating each other (Howard, l.c.). However, should quite accurate results be desired, the process may be modified by evaporating the first bisulphate mother liquor until it will again crystallise (sp gr 1 100), filter-pump the crystals and wash as before, and shake this more concentrated cinchondine liquor into ether and ammonia solution. The impure cinchonidine obtained is dired and weighed, then dissolved in hot absolute alcohol, and 2.1 c c of 50 p c H2SO4 added for each gram, and allowed to cool and stand, when the cinchonidine crystallises out entirely as tetrasulphate is filtered off on the pump, washed with a little alcohol, dissolved in water, precipitated with sodium hydroxide, filtered, washed with water, dried, and weighed as alkaloid. The alcohol mother liquor is diluted with water, the alcohol evaporated off on the water-bath, the two lots of bisulphate crystals added, and the solution diluted and neutralised with ammonia, when, on cooling, the pure quinine sulphate crystallises out and may be filtered off, dried, and weighed. For rough quinine sulphate products (a) it

is best to dissolve them in 40 times their weight of boiling water in a flask, add 10 p c of animal black and a few drops of dilute sulphune acid, boil for 20 minutes, filter, and allow to crystal-Filter off the quinine sulphate and return the mother liquor to the flask, together with the filter-paper and animal black, boil for 20 minutes, filter at the pump, wash the filter-paper with two or three lots of boiling water, evaporate down to about one-sixth of its bulk and neutralise with ammonia to litmus paper Allow to cool and crystallise, filter off the quinine sulphate, and add this to the first crystals Dry and weigh the decolourised quinine sulphate. This should then be put through one of the tests for cinchonidine given above, for preference De Vrij's bisulphate test. The mother liquor from the decofoursed sulphate is added to the mother liquor from which the original impure sulphate was obtained, and is put through tests for other alkaloids described under the 'Assay of Cinchona Barks.' It should be noted here that neutral quinine sulphate is absorbed from its solutions by animal black (up to 15 p.c. of the weight of black used), and for that reason 'blacking in neutral solution is not permissible.

For commercial quinine sulphates (b) similar methods to the above may be employed, but as no (or infinitesimal) colouring matter is present the first decolourising process is not But as the above methods take considerable time to carry out the phaimacopæias adopt a qualitative test, in various forms, suggested by Keiner (Zeitsch anal Chem. 1862, 150, Arch Pharm 1878, 14, 438; ibid. 1887, 25, 712, 749) This test depends upon the fact that quinine sulphate is the least soluble of the sulphates of the cinchona bases, and that quinine is the most soluble of the common bases in dilute ammonia. When 2 grams of pure quinine sulphate is shaken with 20 ce of water at 60° for some time, and then cooled to 15° for at least half an hour (to prevent supersaturation), it gives a filtrate which precipitates with aqueous ammonia (sp gr. 0 959), the precipitate just redissolving when 4 3 c c ammonia have been added Now, quinine sulphate containing any of the other cinchona sulphates when treated in this manner takes more ammonia to give a clear solution; and by this means some idea of the purity of the quinine sulphate This test has been universally is obtained adopted by the pharmacopeeas, to which reference should be made for full details, as the directions vary in the different pharmacopæias Quinine sulphate containing about 5 p c total of cinchonidine and hydrogumine sulphates will require between 6 and 7 cc of ammonia, while one containing about 10 p c of these impurities will require about 10 c c Points to be noticed in applying the test are that the weights of quinine sulphate and water taken do not, within reasonable limits, affect the result, the solution should be cooled gradually to 15° (never below), and retained at that temperature for not less than 30 minutes, the filtration should be rapid and the mixing very carefully done by gently rotating and tilting the testing tube, otherwise clotting of the precipitate may take place and a false result obtained test is greatly influenced should the quimine sulphate have an alkaline reaction (presence of free quinine), when the result is much in excess of that required for a neutral sulphate of the same degree of purity (D. Howard, Pharm. J. 1896, iv. 3, 505); and also by the presence of alkaline sulphates, which make it appear purer than it is (Tutin, Pharm J 1909, iv. 29, 606) These points should be ascertained by testing the filtrate for neutrality with litinus paper, and by evaporating 5 c c to dryness and constant weight, when the residue should not weigh more than 0 008 gram, showing the absence of soluble morganic salts (Biginelli, Monit Scient 1908, 22, 175) Should the cooling of the mixture be carried out above 15° a larger amount of ammonia will be required, and between 15° and 20° a correction of 0.5 c c of ammonia for each degree above 15° may be applied

Hydroquinine is best tested for by dissolving the quinine sulphate in dilute  $H_2SO_4$ , cooling to  $0^\circ$ , and adding 4 p.c. permanganate solution with constant stirring, so long as it is rapidly decolourised. The solution is then warmed, the coagulated manganese dioxide filtered off, the filtrate made alkaline with ammonia and shaken out with ether. The ether

solution filtered and dried to constant weight will give the amount of hydroquinine present

Quinidine, einchonine, and amorphous alkaloid, which are seldom met with except in very rough quinine sulphate, are best estimated by recrystallising the salt and putting the mother liquor through the process given under the 'Assay of Cinchona Barks.'

Cupreine, never present now in commercial barks, may be estimated by shaking the recrystallised quimine sulphate into ether and sodium hydroxide, running off the aqueous portion, and washing it twice with more ether. The aqueous liquor is then run into a basin, acidified with sulphuric acid, and neutralised with ammonia. On cooling any cupreine sulphate present crystallises out.

For the estimation and examination of quinine sulphate, see Hille, Arch Pharm 1903, 241, 54, and Lenz, Zeitsch anal Chem 1888, 27, 549.

Kerner's test is recommended also for the other salts of quinine by all the pharmacopenas, but Tutin ( $l\,\epsilon$ ) has pointed out that the results obtained are quite useless, this is confirmed by the present writers' experience. The best method of examination of quinine salts, other than sulphate, is that of obtaining the alkaloid by putting through ether and ammonia, and following the procedure given under 'Assay of Cinchona Barks'

Quinine disulphate B H<sub>2</sub>SO<sub>4</sub>,7H<sub>2</sub>O crystallises out in white rhombic prisms when the normal sulphate is dissolved in an equivalent of 5 p c H<sub>2</sub>SO<sub>4</sub>. The salt effloresces readily, losing 6H<sub>2</sub>O over sulphune acid, and becoming anhydrous at 100°. When fully hydrated the salt softens at 60°, and as it melts gradually decomposes into quinicine bisulphate (Howard and Chick, Pharm J. 1917, 99, 143). The dehydrated salt melts at 160°. It has an intensely bitter taste and the aqueous solution is strongly fluorescent. It is soluble in 10 parts of water and m 45 parts of alcohol at 15°, in 18 parts of glycerin, 920 parts of chloroform, and in 1770 parts of ether at 25°, in 1 part of water, and 0.5 part of alcohol at 60°. On exposure to light it turns yellow to brown on the surface of the crystals. This salt is usually very pure on account of the bi-sulphates of the other bases being much more soluble.

Quinine tetrasulphate B2H<sub>2</sub>SO<sub>4</sub>,7H<sub>2</sub>O is obtained by dissolving the bisulphate in excess of strong sulphuric acid and concentrating over H<sub>2</sub>SO<sub>4</sub>. It is very soluble in water, less so in alcohol (Hesse, Annalen, 1873, 166, 222).

Quinine hydrochloride B HCl,2H<sub>2</sub>O is obtained by decomposing a solution of quinine sulphate with barium chloride 1t forms long needles, which become anhydrous at 100°, and melt at 158°-160° without decomposition. The commercial salt contains about 8 pc H<sub>2</sub>O, and is usually very pure 1t is soluble in 40 parts of water, in 9 parts of chloroform, and in 3 parts of alcohol at 15°, in 1 part of bolling water or 90 pc alcohol It is largely replacing the sulphate in medicine on account of its superior solubility

Quinine dihydrochloride B.2HCl, made by adding an equivalent of hydrochloric acid to the normal hydrochloride and carefully drying. It is a white crystalline powder containing

2-3 p.c H<sub>2</sub>O. Soluble in 0.75 part of water, 5 parts of alcohol, or 7 parts of chloroform, at 15°; insoluble in ether.

Quinine hydrobromide B·HBr,H<sub>2</sub>O, made similarly to the hydrochloride, or by dissolving quinine alkaloid in one equivalent of HBr Long white silky needles, soluble in 50 parts of water, 7 parts of alcohol, or 10 parts of chloroform at 15°, in 1 part of boiling water. It fuses at 152°, melting completely at 200°.

Quinine dihydrobromide H 2HBr, 3H<sub>2</sub>O, made by adding one equivalent of HBr to the normal hydrobromide Large yellowish prisms, melting at 81°-82°, and soluble in 7 parts of water at 15°. Very soluble in hot water or alcohol, insoluble in other

Other salts of quinine in use, with their solubilities in water, and the amount of alkaloid they contain are—

Acetate B CH<sub>1</sub>COOH, 1 m 30 at 15° (84 4 p.c.), arsenate B<sub>2</sub>H<sub>3</sub>AsO<sub>4</sub>,8H<sub>2</sub>O, 1 m 650 at 25° (69 4 p.c.); bi-arsenate B H<sub>3</sub>AsO<sub>4</sub>,2H<sub>2</sub>O, 664 5 p.c.); arsenite B<sub>3</sub> H<sub>3</sub>AsO<sub>3</sub>,4H<sub>2</sub>O, 1 m 150 hot (83 1 p.c.); arihenalate B<sub>2</sub> AsCH<sub>3</sub>O (86 4 p.c.); arsanilate (p-amidophenylarsenate) B NH<sub>2</sub>·Cl<sub>8</sub>H<sub>4</sub>·AsO<sub>2</sub>H, 1 m 635 at 15° (61 8 p.c.), benzoate B CH<sub>4</sub>O<sub>2</sub>, 1 m 350 at 25° (72 6 p.c.), bimurcarbamide B CO(NH<sub>2</sub>), 2H(1,5H<sub>2</sub>O, 1 m 1 at 15° (59 3 p.c.), camphorate

msoluble (70 4 p c.); carbonate B  $\rm H_2(O_3, \rm H_2O, soluble$  (73 7 p c.), carbolate B  $\rm C_6H_6OH$  (77 5 p c.); chlorate (B  $^\circ \rm HClO_3)_4, 7H_2O$  (73 6 p c.), chloronereurate B 2  $\rm HClC_3, 4H_2O$  (73 6 p c.), chloronereurate B 2  $\rm H_2ClC_6, 3H_2O$  (58 3 p c.); bichloroplatinate B  $\rm H_2PtCl_6, 3H_2O$  (58 3 p c.); bichloroplatinate B  $\rm H_2PtCl_6, 3H_2O$  (43 1 p c.), chromate B  $\rm H_2CrO_4, 8H_2O$  (55 3 p c.); cinnamate B  $\rm C_9H_8O_2$  (68 6 p c.), citrate (basic) B  $_2$   $\rm C_6H_8O_7, 3H_2O$ , 1 in 1200 at 15° (72 5 p c.), citrate (neutral) B  $_3(\rm C_6H_8O_7)_2$ , 1 in 890 at 17° (71 7 p c.); citrate (acid) B  $\rm C_6H_8O_7$ , 1 in 640 at 17° (62 8 p c.), ethylsulphate B H  $\rm C_2H_6SO_4$  (71 2 p c.); bi-ethylsulphate B (H  $\rm C_2H_5SO_4$  (71 2 p c.); bi-ethylsulphate B (H  $\rm C_2H_5SO_4$  (71 2 p c.); formate (quinoform) B  $\rm C_6H_3O_2, H_2O$ , 1 in 19 at 15° (83 5 p.c.), bi-formate B (CH\_1O\_2)\_2 (77 9 p c.), glycerophosphate (\*Kinculme\*)

1~m 850 at 25° (72.6 p.c.), bi-glycerophosphate B C<sub>3</sub>H<sub>2</sub>O<sub>3</sub>H<sub>2</sub>O<sub>4</sub>, 10H<sub>2</sub>O (46.8 p.c.), hydrochlorophosphate B HCl (H<sub>1</sub>PO<sub>4</sub>)<sub>2</sub>,3H<sub>2</sub>O, 1 m. 2 at 15° (53.0 p.c.), hydrochlorosulphate

1 m 2 at  $15^{\circ}$  (74 4 p c.), hydrofluonde B HF (msoluble) (94 2 p c.), hydrodde B :Hl, 1 m 205 at  $25^{\circ}$  (71 7 p c.), br-hydrodde B :2Hl,5H $_2$ O 1 m 20 at  $15^{\circ}$  (48 4 p c.), hypophosphite B:H, $_3$ PO $_2$ , 1 m 250 at  $15^{\circ}$  (83 1 p c.), isobutylformate B  $C_5H_{10}O_2$  (76 1 p c.); isovalerianate B  $C_5H_{10}O_2$  (76 1 p c.), lactate B: $C_3H_6O_3$ , 1 m 6 at  $15^{\circ}$  (78 3 p c.), methylsulphate B:HC: $_3$ RO $_4$  (74 3 p c.), mtrate B:HNO $_1$ H $_2$ O, 1 m 70 at  $25^{\circ}$  (80 0 p c.), oxalate B $_2$ C $_2$ H $_2$ O $_4$ ,6H $_2$ O, 1 m 1400 at  $25^{\circ}$  (76 6 p c.); binoxalate

(75 0 p c), perchlorate B 2 HClO  $_4$ ,7H,O (49 8 p.c.); periodide B HI  $_3$  (insoluble) (45 9 p.c.), phosphate B  $_2$  H  $_3$  PO  $_4$ ,8H  $_2$ O, 1 in 800 at 25  $^\circ$ 

(72 8 p.c.); quinate B  $^{\circ}_{7}H_{12}O_{8},2H_{2}O_{1}$  1 m 7 at 15° (58 7 p.c.), salicylate B  $^{\circ}_{7}H_{6}O_{1}$ , 1 m 230 at 15° (70 2 p.c.), bi-salicylate B( $^{\circ}_{7}H_{6}O_{1},2J_{2}^{\circ}H_{2}O_{1}^{\circ}$ ) acetylsalicylate (aspirin-quinnie of xaxaquin.) B  $^{\circ}_{6}H_{4}OCO_{7}(H_{3}COOH,1$  in 33 at 15° (64 3 p.c.), dibromsalicylate ('biomochinal') B  $^{\circ}_{6}H_{4}O_{8}H_{2}^{\circ}$  (71 2 p.c.); sucemate B<sub>2</sub>  $^{\circ}_{2}H_{6}O_{4},8H_{2}O$  (71 2 p.c.); bi-sulphocyanide B HCNS, $^{\circ}_{4}H_{2}O$  (80 8 p.c.), tannate (very varied composition), about 1 in 2000 at 15°; (20–30 p.c.); taitrate B $^{\circ}_{2}C_{4}H_{6}O_{8},H_{2}O$ , 1 in 1000 at 15° (79 4 p.c.), bitaitrate B  $^{\circ}_{4}H_{6}O_{6},H_{2}O$  (65 8° p.c.), urethane-hydrochloride

very soluble (60 l  $\,\mathrm{p}\,\,\mathrm{c}$  ) ,  $\,$  valemanate

1 in 120 at 15° (73 0 p c). There are three general methods employed in making quinine salts: (1) to decompose as aqueous solution of quinine sulphate with the barium salt of the acid, (2) to dissolve quinine alkaloid in a dilute aqueous solution of the acid, and (3) to dissolve quinine alkaloid and the acid in separate portions of an organic solvent by heating, mix the solutions, and allow to cool and crystallise, or evaporate off the solvent.

lise, or evaporate off the solvent

\*Quinine Esters\*\*—The following esters of quinine are also in use \*—\*

Acetylquinine  $C_{20}H_{23}O_2N_2$  COCH<sub>3</sub>, made by heating quinine alkaloid with acetic anhydride at  $60^\circ$ – $80^\circ$  for several hours. The product is diluted with water, piccipitated with ammonia, the acetylquinine collected, diied, and recrystallised from petroleum ether, in p. 116°–117°

Anisylquinine

made by heating quinine alkaloid with phenylanisate at  $120^{\circ}-130^{\circ}$  for several hours. The warm melted mass is treated with benzene and the free phenol removed with dilute NaOH. The ester is shaken out with dilute acid. Fine white needles. Mp. 87°-88° Benzoyl-quinineC $_{20}H_{43}ON_{2}$  O CO C $_{6}H_{5}$ , made

Benzoyl-quinineC<sub>20</sub>H<sub>23</sub>ON<sub>2</sub>O CO C<sub>6</sub>H<sub>5</sub>, made in a similar way to the above, by heating together quinine alkaloid and phenyl benzoate for several hours at 130°–140°

 $\begin{array}{ccc} \textbf{Cinnamyl-quinine} & C_{20}H_{24}ON_2 \ O \ COC_8H_7, \\ \text{made by treating quinine suspended in benzene} \\ \text{with a solution of cinnamyl chloride in benzene.} \\ \text{Tasteless needles, soluble in hot, sparingly soluble in cold, water} & M p. 235°-236° \\ \end{array}$ 

Diquinine carbonate ('Aristoquinine')

made by adding quinine alkaloid to a solution of phosgene in benzol, washing the resulting mass with water, and recrystallising the rough aristoquinine from alcohol—It forms crystalline groups of needles of m p—189°—It is an important ester on account of its tastelessness and its high percentage of quinine—96 I p c

Quinine ethyl carbonate (Euquinine)

$$\mathrm{C}_{20}\mathrm{H}_{23}\mathrm{ON}_2\mathrm{O},\mathrm{CO}|\mathrm{OC}_2\mathrm{H}_5$$

is similarly made by adding quimine to a benzol solution of ethylchloroformic ester. It occurs as white needles, tasteless; mp 95°. Easily soluble in alcohol, ether, and chloroform, not in water. It gives the thalleroquin reaction and fluoresces in dilute H<sub>2</sub>SO<sub>4</sub>. It forms similar

salts to quinine, but does not give the herepathite reaction. Its salicylate, a white tasteless crystalline powder, m.p. 195°, is the only commercial salt

Salicylquinine ('Saloquinine')

 $\mathrm{C}_{20}\mathrm{H}_{23}\mathrm{ON}_2\mathrm{O}\mathrm{COC}_6\mathrm{H}_4\mathrm{OH}$ 

made by heating quinine alkaloid with salol for several hours at 140°-150°. It crystallises from ether in large colourless crystals insoluble in water, mp 140°. Its salicylate is known as 'Rheumatine'

Quinine carbophenetidide ('Chmaphenme')  $C_{20}H_{23}ON_2O$  CO  $NHC_6H_4OC_2H_6$  occurs as a white tasteless powder, slightly soluble in water, easily soluble in organic solvents

Hydroquinine, its derivatives and salts.

**Hydroquinine**  $C_{20}H_{26}O_2N_2$  is invariably present in commercial quinine sulphate to the extent of 1–2 p c. It may be prepared (and estimated) by converting the quinine sulphate into bisulphate by the addition of 12 ce of N/IH2SO4 for each 5 grams of salt taken, filtering off the crystals, evaporating the mother liquoi again to crystallising-point, filtering off the second crystals, adding ice to the mother hquor till cooled to 0° and then adding ice-cold permanganate solution drop by drop with constant sturing until the colour of the permanganate is no longer rapidly discharged The liquor is now warmed until the manganese oxide coagulates, when it is filtered and shaken with ammonia and ether The aqueous portion is given two more washes with ether, the ethereal solutions bulked and evaporated to dryness, the hydroquimine alkaloid being obtained as a pale yellow crystalline mass

Commercially, hydroquinine is now prepared by the hydrogenisation of quinine sulphate in dilute acid solution in the presence of (a) spongy metals of the platinum group (D R P 252136, 1912), (b) palladium black (Eng. Pat 3948, 1912), (c) palladium coated on barium sulphate (D R P 234137, 1911), or (d) by the action of 2 p c formic acid in the presence of palladium (D R P 267306, 1913) In the last process the formic acid decomposes into carbon dioxide, which escapes, and hydrogen, which enters the quinine molecule at the vinyl

group

Hydroquinine crystallises in fine anhydrous needles from chloroform or ether, readily soluble in most organic solvents, sparingly in water, but more soluble in dilute ammonia,  $[a]_D^{20^\circ} = -142 \cdot 2^\circ$ in 95 pc alcohol (c=2.4 p.e.), m.p. 172° physical and chemical properties it closely resembles quinine, forming similar crystalline

salts of like solubility

The chief salts of hydroquinine are: acetate
B C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>,5H<sub>2</sub>O, needles, soluble in water and arsenate B<sub>2</sub> H<sub>3</sub>AsO<sub>4</sub>,10H<sub>2</sub>O, zoate B C<sub>1</sub>H<sub>6</sub>O<sub>2</sub>, soluble in alcohol, chromate B<sub>2</sub>H<sub>2</sub>UrO<sub>4</sub>,6H<sub>2</sub>O, yellow needles, soluble 1 in 663 of water at 15°; citrate B<sub>2</sub> C<sub>6</sub>H<sub>6</sub>O<sub>7</sub>,10H<sub>2</sub>O, sparingly soluble in water, hydrobromide B HBr,2H<sub>2</sub>O, needles, freely soluble in water dihydrobromide B.2HBr,3H2O, needles, very soluble in water; hydrochloride B HCl,2H,O, long flat prisms, freely soluble in water and alcohol; dihydrochloride B 2HCl, very soluble in water; herepathite, 4B 3H<sub>2</sub>SO<sub>4</sub>·2HI·I<sub>4</sub>,6H<sub>2</sub>O, insoluble; hydriodide B·HI, amorphous, freely soluble in water; dihydriodide B 2HI,4H2O, yellow crystals, soluble in hot water; iodo-hydriodide B 2HI I4, dichroic needles; hyposulphite B<sub>2</sub> H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>,2H<sub>2</sub>O, small prisms, sparingly soluble in water; oxalate B<sub>2</sub> C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>,6H<sub>2</sub>O, sparingly soluble in water; phosphate B<sub>2</sub> H<sub>3</sub>PO<sub>4</sub>,7H<sub>2</sub>O, needles, sparingly soluble in water, platinochloride B<sub>2</sub> H<sub>2</sub>PtCl<sub>8</sub>,3H<sub>2</sub>O, amorther than the solution of t phous, spanngly soluble in water, diplatinochloride B H, PtCl<sub>6</sub>, 2H<sub>2</sub>O, amorphous, sparingly soluble in water, salicylate B  $C_7H_6O_3$ , needles, soluble in alcohol, sulphate B<sub>2</sub>  $H_2SO_4$ , 6 or 8H<sub>2</sub>O, needles, soluble in 348 of water at 15°, disulphate B  $\rm H_2SO_4, 3H_2O$ , long thin needles, very soluble in water, tetrasulphate B  $\rm 2H_2SO_4$ , amorphous, very soluble in water; tartiate B<sub>2</sub> C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>,2H<sub>2</sub>O, prisms, soluble in 545 parts of water at 17°

Hydroquinine unites with I molecule of cupreine, quinidine, or hydroquinidine, and with 2 and 3 molecules of cinchonidine or hydrocinchonidine to form crystalline compounds (Hesse, Ber 1882, 15, 854, Annalen, 1887, 241, 255, Pum, Monatsh 1895, 16, 72)

Hydroquinicine. On heating hydroquinine disulphate to 140° it is converted into its isomer hydroquinicine The pure base is obtained by pouring the incited mass into water, and shaking the solution with NaOH and ether The residue left on evaporation when dissolved in alcohol and neutralised with H,SO, yields the crystalline sulphate, which is filtered off from the thick brown mother liquor and recrystallised to give the pure salt, in delicate acicular colourless needles readily soluble in water and in alcohol A solution of the sulphate in water is pale yellow, which on acidification assumes an intense yellowish-green colour Alkaline hydroxides precipitate the free alkaloid as a resmoid mass which gradually dissolves in ammonia or ammonium salts solutions, but not The hydrochloride is in the fixed alkalis amorphous and gives  $[\alpha]_D^{15} = -17^\circ$ , c=3 in excess of dilute HCl It yields a platinochloride B H2PtCl<sub>8</sub>, H2O, as a pale-yellow flocculent precipitate changing to orange-coloured crystals, insoluble in water and spannigly soluble ın dılute HCl An ethereal solution of oxalic acid produces in an ethereal solution of the alkaloid an amorphous brown precipitate, soluble in chloroform This distinguishes it from quinicine, which yields a voluminous precipitate of minute needles under similar conditions.

Hydrocupreine C<sub>19</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub> is derived from hydroquinine by heating it with 8 parts of HCl (sp gr 1 125) at 150° in scaled tubes. The unchanged hydroquinine is removed by shaking the acid solution with NaOH and ether, the hydrocupreine being obtained as sulphate by neutralising the alkaline layer with dilute H<sub>2</sub>SO<sub>4</sub>. The free alkaloid forms a crystalline powder C<sub>19</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub>,2H<sub>2</sub>O, freely soluble in organic solvents, slightly soluble in ammonia. It melts at 168°-170°, is strongly basic, forming crystalline salts, but is also phenolic, dissolving m alkalis. Its sulphate B<sub>2</sub>·H<sub>2</sub>SO<sub>4</sub> is sparingly soluble in water and in alcohol, the dihydrochloride B 2HCl,H<sub>2</sub>O being freely soluble. It forms a sparingly soluble crystalline tartrate  $\rm B_2\cdot C_4H_6O_6, 2H_2O,$  and an orange-red crystalline platinochloride B  $\rm H_2PtCl_2,$  insoluble in water and dilute HCl. Hydrocuprene is important as it is the source of the alkyl-hydrocuprenes of which ethyl-hydrocuprene hydrochloride  $\rm C_{21}H_{28}O_2N_2\cdot HCl$  (optochin) has been tried recently as a specific for pneumonia (Morgenroth, Berl. Klin Woch 1911, 34, 44; ibid 1914, 47–48; Sir Almroth Wright, Lancet, 1912, 1633, 1705; Pyman, Chem Soc Trans 1917, 662, 1127); and the isoctyl derivatives as a wound disinfectant (Morgenroth and Tugendreich, Biochem Zeitsch 1917, 79, 257; Schaeffer, ibid 1917, 83, 269, Bieling, ibid 1918, 85, 188)

Ethyl-hydrocuprēine hydrochloride occurs as a white crystalline powder, m p 242°, soluble in 10 parts of water, giving a blue fluorescence with dilute H<sub>2</sub>SO<sub>4</sub>, and having a bitter flavour Soluble in 90 p c alcohol The free alkaloid is precipitated by NaOH, and is soluble in chloroform

The hydrochlorides of the higher alkyl-hydrocuprenes are insoluble in water, the dihydrochlorides, which are soluble in 30-40 parts of water, being the salts prepared for use Iso-octyl-hydrocupreine dihydrochloride ('Vuzin') occurs as a greenish-yellow amorphous powder giving a milky solution in 30-40 parts of water, which at higher dilution is merely

opalescent.

Quinidine, its Derivatives and Salts -Quinidine (originally called 'conquinine') (C20H24O2N2), the only known natural stereo-isomer of quinine, occurs in most cinchona barks, especially C pitayensis, and also in cuprea bark. It was not prepared commercially at one time, but was precipitated with the amorphous alkaloids in the quinine tails, the whole being termed 'quinoidine' This was then a convenient source of quinidine, which could be extracted from the powdered quinoidine by means of ether (Hesse, Annalen, 1868, 146, 357, 1873, 166, 232) However, quinidine is now extracted and purified on the commercial scale, only a small amount being precipitated with the amorphous quinoidine. It is most conveniently isolated from the quinine mother liquor as hydriodide, as given under the 'Assay of Cinchona Barks,' being the least soluble of the cinchona hydriodides in neutral aqueous solution It crystallises from dilute alcohol in large prismatic crystals containing 2½H<sub>2</sub>O, losing ½H<sub>2</sub>O on exposure to air From ether rhombohedra containing 2H2O are obtained, and from boiling water plates containing 11H2O From benzene anhydrous needles separate which melt with decomposition at 171.5° (corr) [ $\alpha$ ]<sub>D</sub><sup>15°</sup> = +236 S°-3.0c in 97 pc alcohol, c=1-3. It is soluble in 6900 parts of water at 25°, 750 parts at 100°, 26 parts of alcohol at 20°, and in 22 parts of ether at 20°; slightly soluble in chloroform, benzene, amyl alcohol, and carbon disulphide.

Chemically, quinidine behaves similarly to quinine, yielding salts which, however, with the exception of the hydriodide, are more soluble than those of quinine. Its decomposition products are also the same as, or isomeric with, those of quinine. The disulphate on heating with acid gives quinicine. Treated with fuming sulphuric acid the alkaloid yields quinidine isosulphonic acid  $C_{20}H_{22}O_2N_2$ :HSO<sub>3</sub>, identical with the product obtained from quinine under

similar treatment (Hesse, Annalen, 1892, 267, 141). By the action of permanganate in the cold a base quitenidine, isomeric with quitenine  $(q \ v)$  is obtained

Quitenidine C<sub>19</sub>H<sub>22</sub>O<sub>4</sub>N<sub>2</sub> forms thin plates from dilute alcohol, mp. 246°, soluble in ammonia, from which solution it crystallises in large prisms on evaporation. It is readily soluble in boiling water, acids, and alkalis, less so in alcohol. Its H<sub>2</sub>SO<sub>4</sub> solution is fluorescent and it gives the thalleioquin reaction, which on adding ferrocyanide turns violet-black. It yields a sulphate B·H<sub>2</sub>SO<sub>4</sub>,3H<sub>2</sub>O, colourless prisms, and a platinochloride B H<sub>2</sub>Pt(I<sub>6</sub>,3H<sub>2</sub>O,

large orange-vellow crystals.

Apoquinidine  $C_{19}H_{22}O_2N_2$ , isomeric with apoquinine, is formed similarly by heating the base in a sealed tube to 140° with HCl (sp gr 1125). It is a white amorphous powder, readily soluble in alcohol and ether, giving the thalleloquin reaction, but not showing fluorescence with  $H_2SO_4$ . It melts at 137° (uncorr), and has  $[\alpha]_0^{15} = +153$  3° in 97 p.c. alcohol, c=2. It yields a hydrochloride in needles easily soluble in water. Its acetyl derivative, unlike the base, fluoresces in  $H_2SO_4$  solution

Chloroquinidine  $\rm C_{20}H_{23}ON_2Cl$  is obtained similarly to its isomer, chloroquinine. It is readily soluble in alcohol, benzene or chloroform; m.p. 131°-132°. On treatment with alcoholic potash it gives the same product as chloro-

quinine, i e quinene

Hydrochloroquinidine  $C_{20}H_{20}ClO_2N_2$  is obtained similarly to its isomer, hydrochloroquinine. It forms large crystals soluble in alcohol and in excess of ammonia. It gives the thalleroquin reaction. By longer heating methyl chloride is eliminated and hydrochlorapoquinidine results.

a-Isoquinidine  $C_{20}H_{14}O_{2}N_{2}$ ,  $2H_{3}O$  is formed together with hydrochloroquinidine by heating quinidine with HCl (sp.gr. 1189) at 85°. It forms fine rhombohedra soluble in ether.

Hydroiodoquinidine  $C_{20}H_{2\delta}IO_2N_2$  is obtained when quinidine is dissolved in 10 times its weight of HI (sp gr 1 9) and left for 3 days in the dark Its dihydriodide crystallises in large yellow tables, mp 230°, soluble in alcohol Ammonia liberates the base, which is recrystallised from alcohol as colourless transparent piisms, mp 205°–206°, slightly soluble in water. It gives the thalleioquin reaction and yields a crystalline sulphate, nitiate, hydrochloride, hydrodide, and platinochloride

Quantume Salts—Sulphate  $B_2 H_2SO_4, 2H_2O$  crystallises as white needles or long hard prisms, soluble in 100 parts of water at 15°, or 7 parts at 100°, 20 of chloroform at 15°, insoluble in ether. The salt requires a temperature of 120° for dehydration, and reabsorbs  $2H_2O$  on exposure to air.  $[a]_D^{17} = +2115°$  in absolute alcohol, c=194 p c of hydrated salt. A chloroform solution gradually assumes a yellow colour with slight green fluorescence. On shaking this solution with water the aqueous portion acquires a beautiful green fluorescence.

Commercial quinidine sulphate is usually fairly pure; cinchonine is the most frequent impurity, cinchonidine and quinine being rarely found. To test for other alkaloids dissolve

0.5 gram in 20 c.c. of warm water, add 0.5 gram of KI, and stir vigorously for a few minutes. After half an hour filter off the insoluble hydriodide and add ammonia to the filtrate any decided precipitate occurs (none with the pure salt) shake with chloroform, evaporate the extract to dryness, and weigh Test the residue for quinine, einchonidine, and einchonine Should the two latter be present the precipitated hydriodide will tend to be resinous instead of sandy (De Viij, Pharm J. 1877, 8, 745)

Disulphate BH,SO4,4H,O occurs in long asbestos-like prisms, soluble in 7 parts of water at 15°, giving a beautiful blue fluorescence

met with are Other salts benzoate B C, H, O, H, O, soluble in alcohol, chromate B·H<sub>2</sub>CrO<sub>4</sub>,6H<sub>2</sub>(), large yellow tables; hydrobromide B HBr, soluble in 200 of water at 14°, hydrochloride B HCl, H<sub>2</sub>O, asbestos-like needles, soluble in 75 parts of water at 15°, [a]<sub>D</sub>=+212° -2 56c in alcohol, dihydrochloride B 2HCl,H,O, very soluble in water, hydrodide B HI, short prisms, soluble in 1250 of water at 15°, very slightly soluble in alcohol, dihydriodide B·2HI,3H<sub>2</sub>O, large golden prisms, soluble in 90 of water at 15°; hyposulphite

short prisms, soluble in 415 of water at 10°, oxalate B, C,H<sub>2</sub>O<sub>4</sub>,H<sub>2</sub>O, soluble in 151 of water at 15°, nitrate B HNO<sub>3</sub>, small prisms, soluble in 85 of water at 15°, phosphate B H<sub>2</sub>PO<sub>4</sub>, short prisms, soluble in 31 of water at 10°, platinochloride B<sub>2</sub> H<sub>2</sub>Pt(l<sub>8</sub>,3H<sub>2</sub>O, orange needles, diplatinochloride B H<sub>2</sub>Pt(l<sub>8</sub>,H<sub>2</sub>O, yellow amorphous precipitate, insoluble in water; salicylate B  $C_7H_6O_7$ , soluble in 1650 of water at 25°; succinate B  $C_4H_6O_4$ ,2 $H_2O$ , fine prisms, soluble in 41 of water at 10°, very soluble in alcohol, sulphocyanide B HCNS, colourless prisms, soluble in 1477 of water at 20°, disulphocyanide, B 2HCNS, H2O, long yellow prisms, tannate, indefinite composition, soluble in about I in 2000 of water at 20°; tartrate

# $B_2 C_4 H_6 O_6, H_9 O$

silky prisms, soluble in 39 of water at 15°; bitartrate B C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>,3H<sub>2</sub>O, short prisms, soluble in 350 of water at 15°.

Hydroquinidine  $C_{20}H_{26}O_2N_2$  usually occurs in commercial quinidine sulphate sometimes to the extent of 20 pc It may be separated by repeated recrystallisation from water, but it is preferably obtained by oxidising quinidine sulphate mother liquors with permanganate at

0° (v Hydroquinine).

Quinidine may also be hydrogenated in the presence of palladium black yielding hydro-Forst and Bohringer's statement quinidine that this base is formed by the oxidation of quinidine is inaccurate, being due to their using quinidine containing hydroquinidine (Ber 1881, 14, 1954; 1882, 15, 520, 854, Hesse, *ibid.* 1882, 15, 3010; Annalen, 1888, 243, 146) It forms efflorescent prismatic needles containing 2½H<sub>2</sub>O, m.p. 167°, readily soluble in alcohol and chloroform, slightly soluble in ether Its rotatory power is almost the same as that of quinidine It gives the thalleroquin reaction and behaves generally like quinidine. Oxidation with chromic acid yields quininic

Salts.—Its sulphate B, H,SO4,2H,O occurs as fine needles, soluble in 92.3 of water at 16°, or B2·H2SO4,8H2O, thick efflorescent prisms (Hesse) Its H,SO<sub>4</sub> solution shows blue fluorescence. Other salts are . benzoate B C7H6O2, colourless plates, hydrobromide, fragile scales, slightly soluble in water; hydrochloride B HCl, short needles or thick prismatic plates, readily soluble in water; hydrochide B HI, large colourless needles, very slightly soluble in water, dihydriodide B.2HI,3H2O, orange-yellow prisms, fairly soluble in water, platinochloride

## B·H, PtCl, 2H, O

small orange-yellow needles, salicylate B.C. H.O., six-sided plates; tartrate  $B_2C_4H_6O_6.2H_2O$ , brilliant prisms, readily soluble in water, bitartrate  $B_2C_4H_6O_6.3H_2O$ , needles, slightly soluble in water.

Cinchonine, its Derivatives and Salts .- Cinchonine (C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub>) occurs in all cinchona barks It is readily prepared from quinine mother liquors, after removal of the cinchonidine as tartrate, by rendering alkaline and shaking with a limited quantity of washed ether. The cinchonine crystallises out from the ether in small hard pusms, and may be filtered off and recrystallised from dilute alcohol The crystals melt at 264°, subliming with some decomposition at a higher temperature as monoclinic prisms Ammonia precipitates the alkaloid from acid solutions in white flocks gradually becoming crystalline; but from hot solutions of ammonium salts cinchonine displaces ammonia The freshly precipitated alkaloid is much more soluble in solvents generally than crystalline cinchonine. The crystals dissolve in 3670 of water at 20°, in 2500 of water at 100°, in 115 of absolute alcohol at 17°, in 356 of chloroform at 17°, in 371 of ether (sp gr 0 730) at 10°, in 40 of amyl alcohol at 17°, and in 18 parts of a mixture of alcohol and chloroform (4 · 1) at 17°. Readily soluble in acids, but the salts of cinchonine are hable to hydrolyse with water and deposit alkaloid Cinchonine is easily distinguished from the other alkaloids by giving neither fluorescent solutions nor an insoluble tartrate, and by being insoluble in ether  $[a]_D^{15^{\circ}} = +226 5^{\circ}$ in 97 p.c. alcohol, c=0.5Cmchonine does not give the thalleioquin reaction. On oxidation it yields the same products as quinine, except that the quinolinic products lack the p-methoxy group

Cinchotenine  $C_{18}H_{20}O_3N_2$  is obtained on oxidising einchonine with permanganate in acid solution (Skraup, Annalen, 1879, 197, 374) It is purified similarly to quitenine (q v) It forms efflorescent needles or plates containing 3H<sub>2</sub>O, m p 198°.  $[a]_D^{15} = +135^\circ$  in alcohol solution, c=2 of hydrated base. A 2 p.c. solution in dilute H<sub>2</sub>SO<sub>4</sub> gives  $[a]_D^{15} = +175$ 5°. It gives a platinochloride B H<sub>2</sub>PtCl<sub>6</sub>, large orange prisms, and an aurichloride B H<sub>2</sub>Au<sub>2</sub>Cl<sub>8</sub>, yellow needles; both soluble in water and in alcohol.

Cinchotenicine  $C_{18}H_{20}O_3N_2$ , isomeric with cinchotenine, is formed from it by heating to 140° with H2SO4. A brown amorphous body soluble in water, giving amorphous platino-and auri-chlorides.  $[\alpha]_D^{15^\circ} = +0.9^\circ$ .

Cinchoninic acid  $C_9H_6$  COOH N (4-carboxy-quinoline), obtained by oxidising cinchonine with chromic acid (v Quinnine acid) Forms caffeine-like needles containing 1H<sub>2</sub>O, or tables or prisms with 2H<sub>2</sub>O; m p. 253°-254°, subliming with decomposition (Caventou and Willin, Annalen, Supl 1870, 7, 247; Skraup, Annalen, 1880, 201, 294) It is only slightly soluble in water, more so in acids and alkalis, but its salts dissociate in water Heated with lime it yields quinoline Permanganate oxidises it to a-carbocinchomeronic acid (Ramsay and Dobbie, Chem Soe Trans 1884, 35, 189), nitric acid at 120°-140° yields einchomeromic acid (Weidel, Annalen, 1874, 173, 76), and chromic acid gives kynurine (Skraup, Monatsh 1888, 9, 783; 1896, 17, 366) On fusion with potassium hydroxide oxyemchommic acid sublimes (Komgs and Kerner, Ber 1883, 16, 2152) Cinchommic acid may also be prepared by oxidation of lepidine and 4-ethylquinoline

Cinchene C<sub>19</sub>H<sub>20</sub>N<sub>2</sub> is obtained by refluxing cinchonine chloride with alcoholic potash (v. Quinene), as trimetric tables, m.p. 123°-125° It is a ditertiary base, soluble in ether, dextrorotatory, and yields crystalline salts hydrolysis with 25 pc. phosphoric acid it yields lepidine and meroquinene. Chromic acid oxidises it to cinchoninic acid Treated with bromine in chloroform solution two isomeric dibromides ( $\alpha$  and  $\beta$ ) of cinchene are formed, both of which are converted into a new base, dehydrocinchene ( $C_{19}H_{18}N_2$ ), with loss of 2HBr (Comstock and Konigs, Ber. 1886, 19, 2857) Chlorine reacts similarly to bromine with cinchene On heating cinchene in a sealed tube with HBr (sp gr 1 49) to 190° it loses a molecule of ammonia and adds a molecule of

water to form apocinchene (v Apoquinene).

Apocinchene C<sub>19</sub>H<sub>19</sub>ON, colourless needles, 209°, volatilising without decomposing, soluble in alcohol, acids, and alkalis (Comstock and Konigs, Ber. 1881, 14, 1854; 1884, 17, 1989). Its salts give yellow solutions which readily hydrolyse, and its solutions in alkalis are precipitated by CO. It readily forms ethers, which with dilute HNO<sub>3</sub> oxidise to alkylapocinehenic acids. Chromic acid oxidises it to cinehoninic acid, and fusion with potassium hydroxide gives oxyapocinchene C<sub>19</sub>H<sub>19</sub>O<sub>2</sub>N, a base isomeric with apoquinene.

### Ethylapocinchenic acid

#### C<sub>17</sub>H<sub>13</sub> OC<sub>2</sub>H<sub>5</sub> COOH N

on refluxing with HBr (sp gr 1 49) gives rise to a lower homologue of appenihene named homopocinchene  $C_{17}H_{16}ON$ . This body melts at 184°-185°, is very soluble in ether, benzene, or hot alcohol, insoluble in water

Dihydrocinchene  $C_{19}H_{22}N_2$  was found by Konigs in the mother liquors from appointhene, and was derived from the hydrocinchonine present in the cinchonine from which the latter was prepared.

Cinchonine chloride  $C_{19}H_{21}N_2Cl$ , obtained similarly to quinine chloride (qv.). It crystallises in prisms, melting at 72°, and is dextrorotatory On reduction with iron and H<sub>2</sub>SO<sub>4</sub> it gives desoxycinchonine C<sub>19</sub>H<sub>22</sub>N<sub>2</sub>, m p 90°-92°, dextrorotatory. With alcoholic potash cinchene is obtained (q.v.).

Cinchonine dichloride C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub>Cl<sub>2</sub> is made from cinchonine dibromide (q.v). It crystallises

by passing chlorine into a cooled HCl solution of cinchonine. It is very similar to dichlorocinchonine, and may be identical with it.

Dichlorocinchonine C<sub>19</sub>H<sub>20</sub>OCl<sub>2</sub>N<sub>2</sub> is obtained by passing chlorine into a hot concentrated alcoholic solution of cinchonine hydrochloride It crystallises as microscopic needles from alcohol, and gives crystalline salts with 2HCl, 2HBr, and  $H_{\star}$ PtCl<sub>8</sub> (Laurent, Annal Chim Phys (3) 24, 302)

Bromocinchonine C19H21OBrN2 is formed by adding bromine to an alcoholic solution of cinchonine hydrochloride (Laurent, lc, Kopp, Arch Pharm (3), 9, 34) Its dihydrochloride crystallises in rhombic tables. Treated with alcoholic potash it exchanges Br for OH, and

gives oxycinchonine  $C_{19}H_{22}O_3N_2$ Sesquibromocinchonine  $C_{38}H_{41}O_2Br_3N_4$ , obtained by the action of excess of bromine on an alcoholic solution of cinchonine It crystallises in fine needles and forms salts with 4 molecules of monobasic acids Treated with alcoholic

 $\begin{array}{c} \text{potash it yields sesquioxycmchonine } C_{19}H_{20}\mathrm{OBr}_2N, \quad \text{formed} \end{array}$ by adding excess of biomine to an aqueous solution of cinchonine hydrochloride (Comstock and Konigs, Ber 1884, 17, 1995) It gives colourless crystals containing 1H2O from alcohol, msoluble in water Alcoholic potash yields dioxycinchonine C<sub>19</sub>H<sub>22</sub>O<sub>3</sub>N<sub>2</sub>.

Cinchonine  $\alpha$ - and  $\beta$ -dibromides

$$C_{19}H_{22}OBr_2N_2$$

formed by adding biomine to a solution of cinchonine in chloroform and alcohol hydrobromide of the a-compound being less soluble than the β-compound crystallises out, the latter remaining in the mother liquor free α-base crystallises out from algohol (90 p c) and chloroform (1.3) with 1H2O, the B-base being anhydrous. Both behave as true bases Heated with alcoholic KOH forming salts they lose 2HBr, yielding a new base, dehydrocinchonine (Comstock and Konigs, Ber 1886, 19, 2854, 1887, 20, 2510).

Hydrochlorocinchonine  $\rm C_{19}H_{23}CION_2$ , formed by allowing 10 parts of HCl saturated at  $-17^{\circ}$ to act on cinchonine at a low temperature for some weeks. The base crystallises from alcohol in white needles, mp 212°-213° Heated with alcoholic KOH an isomer of cinchonine, (Comstock and a-isocinchonine, is formed

Konigs, Ber 1887, 20, 2519)

Hydrobromocinchonine C<sub>19</sub>H<sub>23</sub>BrON<sub>2</sub> formed similarly to the last compound, but the reaction is more rapid and works equally well at higher temperatures The base crystallises from alcohol in white scales Heated with 85 p c alcohol it is converted into a mixture of apocinchonine, α-isocinchonine and δ-cinchonine Heated with alcoholic KOH a mixture of apocinehonine a-isocinehonine and cinchonifine (? hydrocinchonine) is formed

Hydroiodocinchonine C<sub>19</sub>H<sub>23</sub>ION<sub>2</sub> is obtained by heating dry einchonine with 5 parts of HI (sp gr 17) on the steam-bath The free base crystallises from alcohol in silky needles melting at 158°-160° with decomposition It readily combines with 2 molecules of monobasic acids to form crystalline salts

C19H20ON2, Dehydrocinchonine

from alcohol in needles melting at 202°-203°, subliming at a higher temperature; very soluble in alcohol, acetone, and chloroform, less so in ether or benzene It behaves very similarly to cinchonine, giving salts and halogen derivatives containing H<sub>2</sub> less

Derivatives Isomeric with Cinchonine—

About 20 isomers of cinchonine have been described by various workers, but Skraup's investigations have shown that many of these bodies are identical, and several others are mixtures with cinchonine Hydrocinchonine has also been mistaken for an isomer of cinchonine (Monatsh 1897, 18, 411, 1899, 20, 571, 1901, 22, 1103, 1903, 24, 294, Langer, ibid 1901, 22, 151)

Cinchonicine (cinchotoxin) C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub> occurs in commercial quinoidine, but may be prepared by heating the acid sulphate or tartrate of einchonine or cinchonidine to 130°-140° (Pasteur, Compt rend 1853, 37, 110; Hesse, Annalen, 1875, 178, 213, D Howard, Chem Soc Trans, 1872, 25, 102) It forms a viscid mass, yellow to red, melting at about 50° Roques obtained it in long prismatic needles, mp 49°-50°, by evaporating an ethereal solution at 45°, in a current of hydrogen.  $[\alpha]_{\rm D}^{15}$ =47.2° for 1 pc. solution in chloroform or absolute alcohol It is very soluble in alcohol or ether, slightly soluble in water Its solutions absorb CO2 from the air and expel ammonia from its salts gives the same oxidation products as einchonine Like quinicine, it gives a precipitate with hypochlorites, but does not give the thalleroquin reaction (distinction from quinicine) Many of its salts are amorphous, but the following are ciystalline. hydrodide B.HI, prisms, slightly soluble in water; oxalate B<sub>2</sub> C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>,7H<sub>2</sub>O, BH<sub>2</sub>PtCl<sub>8</sub>,H<sub>2</sub>O, bitartrate B (¹<sub>4</sub>H<sub>6</sub>O<sub>6</sub>,H<sub>2</sub>O Allocinchonine (pseudocinchonine)C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub>

is obtained by heating the hydriodide of hydroiodocinehonne  $C_{19}H_{21}TON_2HI$  with 10 parts of water at  $150^{\circ}-160^{\circ}$  It melts at  $214^{\circ}-216^{\circ}$ , slightly soluble in alcohol or ether, and gives  $[\alpha]_{D}^{20} = +164.8^{\circ}$  for 3 pc solution in absolute alcohol. Its sulphate B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub> is less soluble in water than cinchonine sulphate. On oxidation it yields einchonine acid and allometo-quinene (Lippman and Fleissner, Monatsh 1893, 14, 371)

Tautocinchonine  $C_{19}H_{22}ON_2$  is obtained by the action of alcoholic KOH and silver nitrate on hydrobromocinchonine hydrobromide

# C<sub>19</sub>H<sub>23</sub>BrON<sub>2</sub> HBr

It is soluble in alcohol, slightly soluble in ether, m p  $~252~5^{\circ},$  and gives [a]  $_{D}^{20}\!=\!+209~4^{\circ}$  for 3 p c solution in absolute alcohol (Lowenhaupt, Monatsh. 1898, 19, 461)

α-Isocinchonine (cinchoniline) C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub> is obtained by heating cinchonine with sulphuric acid or HCl (sp gr 1·125); or by heating hydro-bromocinchonine or its salts with alcoholic KOH (Jungfleisch and Leger, Bull. Soc chim 1888, 49, 747; Compt. rend 1892, 113, 651; *ibid* 1894, 118, 536; Hesse, Annalen, 1888, 243, 147; *ibid.* 1890, 260, 213; 1893, 276, 91, Comstock and Königs, Ber. 1887, 20, 2510, 1892, 25, 1539; Skraup, Monatsh 1899, 20,

581; 1902, 22, 1097; Pum, *ibid.* 1892, 13, 676) It crystallises in silky needles from ether containing 3H2O; from petroleum ether in large prisms, volatile, m.p 126 5°.  $[a]_{n}^{15} = +51 6°$ for 3 pc. solution in absolute alcohol. It is readily soluble in ether, alcohol, or benzene, little soluble in water. It forms crystalline salts, all of which except the dhydriodide are very soluble in water Heated with HCl (sp gr 1·125) at 140°-150° it is converted into apoisoeinchonine (apocinchonigine), mp 216°, and with concentrated  $H_2SO_4$  at  $60^\circ-80^\circ$  it changes into  $\beta$ -isoeinchonine. On oxidation it yields cinchoninic acid and a-isomeroquinene.

 $\beta$ -Isocinchonine (cinchonigine)  $C_{19}H_{22}ON_2$ , references under Isocinchonine. It crystallises from alcohol in colourless prisms, volatile, 126°-127°.  $[a]_{\rm D}^{18} = -59.5$ ° in absolute l Very soluble in alcohol, ether, or m.p benzol. Its salts, including the dihydriodide, are very soluble in water. Heated with HCl (sp gr I 125) it yields a poisocinchonine (apocinchinigine), m p.  $216^{\circ}$ . On oxidation it gives cinchoninic acid and B-isomeroquinene.

δ-Cinchonine C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub> is formed at the same time as  $\alpha$ -isocinchonine (q v) when hydrobromocinchonine dihydrobromide is heated with alcohol or alcoholic KOH Long prisms, mp 150°, very soluble in alcohol, benzene, or chloroform, slightly soluble in ether  $[a]_{\rm D}^{17} = +125.2^{\circ}$  for 1 p c solution in 97 p c alcohol. Its salts are very soluble with the exception of the hydrochloride, hydrobromide, and oxalate (Jung-

fleisch and Leger, Compt rend 1894, 118, 29)

ε-Cinehonine C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub> was isolated by
Lowenhaupt from tautocinchonine (q v) It is a base melting at 150°-152°, giving crystalline salts  $[\alpha]_{\rm p}^{20} = +67^{\circ}$  for 1 p c solution in alcohol

Cinchonidine  $C_{19}H_{12}ON_2$  was obtained in 5 pc yield by Konigs and Hussman from emchanine by heating it for 15 hours with an amyl alcohol solution of KOH (Ber 1896, 29 2185)

Hydro-derivatives of Cinchonine -On reducing einchonine with zine and H2SO4, Schutzenberger obtained a resinous body which he termed a hydrate of cinchonine, assigning to it the formula  $C_{19}H_{22}ON_{2},2H_{2}O$ . This body is stable up to 120°, but at 140° it loses  $\frac{1}{2}H_{2}O$ , and at 150° retains 1H<sub>2</sub>O (Annalen, 1858, 108, 348) Zorn reducing cinchonine with sodium amalgam in acetic acid obtained an oily mass containing two bases separated by ether The one insoluble in ether he named hydrocinchonine (the dihydrodicinchonine of Skraup) C<sub>19</sub>H<sub>24</sub>ON<sub>2</sub>, the other soluble in ether he called amorphous Skraup) C<sub>19</sub>H<sub>26</sub>ON<sub>2</sub> The first crystallises from alcohol in small colourless anhydrous plates or scales, mp 257°-258° It yields a crystalline sulphate B·H<sub>2</sub>SO<sub>4</sub>,2H<sub>2</sub>O (or anhydrous) in long needles It is readily attacked by permanga-nate, but on further treatment with sodiumamalgam is unattacked The second dissolves in ether with a fine violet fluorescence, and is obtained as a yellowish amorphous mass on evaporation. It yields no crystalline salts or derivatives. It is a stronger base than cinchonine, dissolving in acids with elevation of

iperature, and is readily attacked by peraganate (Zorn, J pr Chem. 1873, 8, 279, ; Skraup, Ber. 1878, 11, 312, Howard, m. Soc. Trans. 1873, 26, 1180). By the ion of sodium and anhydrous amyl alcohol cinchonine Norwall obtained an oily base  $\rm H_{26}ON_2$ , which he termed tetrahydrocinnine (Monatsh 1895, 16, 321)

None of the above is identical with the lrocinchonine which occurs naturally in shona barks (Caventou and Wilm, Annalen, l. 1870, 7, 247, Forst and Bohringer, Ber. 1, 14, 436; Hosse, Ber. 1895, 28, 1421)

Hydrocinchonine (Cinchotine) C<sub>19</sub>H<sub>24</sub>ON<sub>2</sub> is onstant associate of cinchonine, which may tain up to 10 pc It may be prepared alarly to hydroquinine (qv.), by oxidation the emchanine with permanganate; or by urating an alcoholic solution of cinchonine trochloride with HCl gas, when the dihydrooride of cinchonine separates out, leaving the frocinchonine in solution (Hesse, Annalen, 3, 276, 88). It forms slender anhydrous sms or scales, m p 277°, subliming at higher iperature Soluble in 136 parts of 90 p cohol, or 187 of ether at 20°, slightly soluble boiling water It is not attacked by cold manganate, but chromic acid oxidises it to chonine and cincholoiponic acids With sphorus pentachloride it yields hydrochonine chloride C<sub>19</sub>H<sub>23</sub>ON,Cl, m p 85°-87°, iverted by alcoholic KOH to dihydrochene C<sub>19</sub>H<sub>22</sub>N Its chief salts are . benzoate 17H6O2, small needles, slightly soluble in ter: hydrobromide B HBr,2H2O, dihydromide B 2HBi, prismatic needles, very soluble water, insoluble in alcohol, hydrochloride ICl,2H<sub>2</sub>O, needles, soluble in 47 of water, ydrochloride B 2HCl, very soluble in water, htly soluble in alcohol; hydriodide B·HI,H<sub>2</sub>O, builess prisms, readily soluble in water and shol, nitrate B'HNO<sub>3</sub>,H<sub>2</sub>O, transparent les, oxalate B C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>,H<sub>2</sub>O, fine needles, y slightly soluble in water; platinochloride  $1.PtCl_6$ , orange-yellow crystals, sulphate  $H_2SO_4, 12H_2O$ , fine needles or prisms, efflorent, soluble in 30 5 of water at 13°, sulphoand B HCNS, long needles, slightly soluble water; tartrate  $C_4H_6O_6, 2H_2O, B_2$ , needles ible in 57 of water at 16°; bitartrate L4H6O6,4H2O, needles, soluble in 78 of water

Salts of Cinchonine—Cinchonine sulphate H<sub>2</sub>SO<sub>4</sub>,2H<sub>2</sub>O forms short haid prisms or dles, melting at 198 5° when dehydrated, omposing at 200° It is soluble in 70 parts water at 15°, in 13 parts at 100°, in 9 parts at ing-point, in 2300 of ether at 25° As the phates of quinne and cinchonidine are insole in chloroform, this solvent forms a good test the purity of cinchonine sulphate —A soluroscient of this base in dilute H<sub>2</sub>SO<sub>4</sub> does not presce (although the commercial salt usually as so slightly owing to the presence of a little nine or quindine), nor does it give the illeioquin reaction. With iodine it yields ducts analogous to herepathite, the principal ing 8B·6H<sub>2</sub>SO<sub>4</sub>·6HI I<sub>10</sub>·12H<sub>2</sub>O,

 $4B H_2SO_4\cdot 4HI\cdot I_{10}$ ,  $2B\cdot H_2SO_4\cdot 2HI\cdot I_2$  rgensen). Vol. 1I.-T.

Cinchonine disuiphate B  $\rm H_2SO_4, 3$  or  $\rm 4H_2O$ , large colourless prisms, very soluble in water, and crystallising with difficulty.

Cinchonine hydrochloride B HCl,2H,O occurs as prismatic needles, melting at 130° It is soluble in 24 parts of water at 10°, the solution dissociating slightly and depositing alkaloid. Soluble in 2 parts of 85 p.c. alcohol and 290 parts of ether at 15°.

Other salts are . benzoate B C2H6O2, soluble in 166 of water at 15°, citrate B<sub>7</sub> C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>,4H<sub>2</sub>O, soluble in 48 of water at 12°, and citrate B<sub>2</sub> C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>,4H<sub>2</sub>O, soluble in 55 of water at 15°; hydrobiomide B HB1,H2O, long needles, soluble in 33 5 of water at 15°, more soluble in alcohol; dihydrobromide B 2HBr, large rhombohedra, soluble in 1.4 of water at 15°, less soluble in alcohol, dihydrochloride B 2HCl, rhombic tables, very soluble in water, insoluble in alcohol; hydriodide B HI,H2O, white crystals, dihydriodide B 2H1, H2O, golden-yellow prisms, hydrodate B HIO, crystals, exploding at 120°, intrate B HNO, H,O, long monoclinic crystals, soluble in 26 4 of water at 12°, oxalate of water at 10°, platmochloride B H<sub>2</sub>Pt(1<sub>6</sub>, pale yellow amorphous precipitate, succinate B C<sub>4</sub>H<sub>6</sub>O<sub>4</sub>,1 or 1½H<sub>2</sub>O, long needles or thick prisms, readily soluble in water, sulphocarbolate B C. H. HSO, OH, reddish-white needles, sulphocyanide B HCNS, prisms, soluble in 474 of water at 20°, salicylate B C<sub>2</sub>H<sub>6</sub>O<sub>3</sub>, soluble in 590 of water at 25°, taitrate B<sub>2</sub> C<sub>4</sub>H<sub>6</sub>O<sub>4</sub>, 2H<sub>2</sub>O, crystals, soluble in 33 of water at 16°, bitaitrate (lievo-) B (4H6O6, H2O, soluble in 100 of water at 16°, bitaitrate (dextro-), soluble in 101 of water at 16° (Pasteur)

Cinchonidine and its Derivatives and Salts.

Cinchonidine C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub>, a stereoisomer of emchonine, occurs in most emchona barks, especially in C succirubia. It is prepared from quinine sulphate mother liquoi by precipitation as neutral tartrate with Rochelle salt This tartrate is washed with water and decomposed with ammonia, the rough base being dried carefully at a low temperature On dissolving this in 21 cc of 50 pe H,SO4 and 11 cc of absolute alcohol for every gram of base, and allowing to cool, pure einchonidine tetrasulphate crystallises out nearly quantita-This is filtered off, washed with absolute alcohol, dissolved in water, and precipitated as alkaloid with NaOH, being finally recrystalhised from alcohol It forms short anhydrous prisms or thin plates, soluble in 16 parts of 97 p c alcohol at  $15^{\circ}$ , in 188 parts of ether, or in 4800 parts of water at  $25^{\circ}$ , readily soluble in chloroform or amyl alcohol. It melts at 210 5°, and has  $[a]_D^{15} = -10748^{\circ} + 03c \text{ in } 97 \text{ p c.}$ alcohol, c=1-5Like einchonine, its solutions do not fluoresce, nor give the thalleroquin reaction (distinction from quinine and quinidine). It differs from cinchonine in giving an insoluble tartrate, by its greater solubility in ether, and by being laworotatory For its estimation in cinchona barks, see under 'Quinine Sulphate.' It gives the same oxidation products as cinchonine with chromic or nitric acid, and cinchonicine, identical with that from cinchonine, is obtained by heating with glycerin to 200°,

or with strong H<sub>2</sub>SO<sub>4</sub> to 130°. The following solution in 97 pc. alcohol. It forms very are the most important derivatives of einchoni-

Cinchotenidine C<sub>18</sub>H<sub>20</sub>O<sub>3</sub>N<sub>2</sub>, obtained similarly to quitenine by oxidation of cinchonidine with permanganate at 0°. It forms filiform or thick prismatic crystals containing 3H<sub>2</sub>O, slightly efflorescent. It melts at 256°, and has  $[\alpha]_D^{15} = -189^{\circ}$  in alcohol — Its acid solutions do not fluoresce Its salts are very soluble and difficult to crystallise It forms a sulphate B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub>,2½H<sub>2</sub>O, very soluble in water, and a platinochloride B, H, PtCl, orange-yellow tables Its solutions in alkalis are readily decomposed by CO<sub>2</sub> (Skraup, Annalen, 1880, 201, 300, Hesse, Ber. 1881, 14, 1892).

Cinchonidine chloride C<sub>19</sub>H<sub>21</sub>N<sub>2</sub>Cl is obtained by the action of phosphorus pentachloride on cinchonidine It forms large crystals melting at 108°-109°, soluble in other and possessing basic properties On heating with alcoholic KOH it yields cinchene (similarly to cinchonine chloride), which on treatment with HCl is converted to apocinchene

**Dibromocinchonidine** C<sub>19</sub>H<sub>20</sub>ON<sub>2</sub>Br<sub>2</sub> is formed by adding 4 atoms of bromine to 1 molecule of cinchonidine suspended in CS<sub>2</sub>. Its dihydro-bromide crystallises in fine yellow needles readily soluble in alcohol (Skalweit, Annalen, 1874, 172, 103). On heating with alcoholic KOH it yields dioxycinchonidine

**Dioxycinchonidine**  $C_{19}H_{22}(OH)_2ON_2$ tallises from alcohol in ramified crystals gives two sulphates, B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub>,2H<sub>2</sub>O and B H<sub>2</sub>SO<sub>4</sub>, and a platinochloride B H<sub>2</sub>PtCl<sub>6</sub>

Hydrochlorocinchonidine C<sub>19</sub>H<sub>23</sub>ClON<sub>2</sub> formed by heating cinchonidine with HCl (sp gr 1 189) at 85° for two days. It is a base indistinguishable from hydrochloroapocinchonidine and is probably identical with it

Hydrochloroapocinchonidine C<sub>18</sub>H<sub>23</sub>ClON<sub>2</sub> is obtained together with apprinchonidine by heating cinchonidine in a sealed tube at 150° with fuming HCl for 6 hours - It crystallises in plates from alcohol, m p 200°,  $\left[\alpha\right]_{\rm D}^{15} = -142.2^{\circ}$ in dilute HCl, slightly soluble in ether, chloro-form, or alcohol. It forms a dihydrochloride B 2HCl, very soluble in water, and a platino-chloride B H<sub>2</sub>PtCl<sub>6</sub>,2H<sub>2</sub>O (Hesse, Annalen, 1880, 205, 327)

Hydroiodocinchonidine  $C_{19}H_{23}ION_2$  is obtained by heating cinchonidine on the waterbath with 5 parts of HI (sp gr. 17) for 21 hours It is a base very slightly soluble in hot alcohol, from which it falls on cooling as a crystalline precipitate, mp 166°. It gives a sulphate B H<sub>2</sub>SO<sub>4</sub>, soluble m 25 p c alcohol Heated with alcoholic KOH it yields an isomeride of cinchonidine, \beta-cinchonidine, and with silver nitrate another isomeride, \gamma-cinchonidine (Neumann, Monatsh. 1892, 13, 651)

Derivatives Isomeric with Cinchonidine -Apocinchonidine C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub> is formed by the action of HCl on cinchonidine (v Hydrochloroapocinchonidine) Slightly soluble in strong alcohol, very slightly soluble in dilute alcohol, ether, or chloroform, almost insoluble in water. It forms small plates, m p. 225°, with decomposition, and has  $\left[\alpha\right]_{\mathbf{p}}^{15} = -129 \cdot 2^{\circ}$  for 0.8 p.c. soluble amorphous salts, whose acid solutions do not fluoresce, and is markedly different from cinchonidine in giving a soluble tartrate.

Isocinchonidine C<sub>19</sub>H<sub>22</sub>ON<sub>2</sub> is obtained as its sulphonate when cinchonidine is dissolved in fuming H<sub>2</sub>SO<sub>4</sub> The base melts at 235°, and is very soluble in alcohol or chloroform, very slightly soluble in ether (Hesse, Annalen, 1888, 243, 149)

 $\beta$ -Cinchonidine  $C_{19}H_{22}ON_2$  is obtained by heating hydroidocinchonidine for 4 hours with alcoholic KOH, and is probably identical with Hesse's homocinchonidine It forms plates or prisms, m p 207°,  $[\alpha]_D^{1.0} = -181^{\circ}$  for 1 25 p.c. solution in dilute HCl. On prolonged heating with HCl it is converted into apocinchonidine, from which it may be separated by means of its insoluble taitrate (Hesse, Annalen, 1880, 205, 327)

Hydro-derivatives of Cinchonidine — On reduction with zine and H<sub>2</sub>SO<sub>4</sub>, or sodium and alcohol, cinchonidine yields products very similar, chemically and physically, to those obtained from einchonine Norwall describes a tetrahydrocinchonidine C<sub>19</sub>H<sub>26</sub>ON<sub>2</sub>, obtained with sodium and pure amyl alcohol (Norwall, Ber 1896, 29, 801)

 $Hvdrocinehonidine(cinchamidine)C_{19}H_{24}ON_{2}$ , usually accompanies circhonidine in its commercial salts, and may be obtained from cinchonidine sulphate mother liquors by treatment with permanganate at 0° (v Hydroquinine) The base crystallises from dilute alcohol in hexagonal plates, and from strong alcohol in short anhydrous prisms, melting at 230°,  $\left[\alpha\right]_{D}^{15} = -98.4^{\circ}$  for 2 pc solution in 97 pc alcohol It is almost insoluble in ether, chloroform, or water, and less soluble in alcohol than cinchonidine. Its salts crystallise readily and fluoresce in sulphuric acid solution (Hesse, Bei 1881, 14, 1683, Forst and Bohunger, Ber. 1882, 15, 520)

Salts of Cinchonidine —Cinchonidine sulphate B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub>,3H<sub>2</sub>O occurs as prismatic needles. From dilute aqueous solutions needles contaming 6H<sub>2</sub>O crystallise, and from alcohol fine prisms with 2H<sub>2</sub>O. These salts are completely dehydrated at 100°, but absorb 2H<sub>2</sub>O again on exposure to air. The anhydrous salt melts at 205° with decomposition It is soluble in 100 parts of water, in 60 of alcohol, or 1000 of chloroform, at 15°, soluble in 21 of water at 80°, insoluble in ether or benzene. The commercial salt usually gives the thalleroquin reaction and fluoresces with dilute H,SO4 owing to the presence of quinine which may occur to even 60 p c Cinchonine and quinidine are also frequent impurities A strong hot aqueous solution of the salt stirred with half its weight of Rochelle salt and allowed to cool and filtered, should not show more than a slight turbidity when the filtrate is treated with ammonia (Hesse, Zeitsch anal Chem 1876, 15, 464).

Cinchonidine disulphate B·H<sub>2</sub>SO<sub>4</sub>,5H<sub>2</sub>O occurs in long colourless efflorescent crystals, very soluble in water and alcohol (separation from

quinine, see 'Assay').

Cinchonidine tetrasulphate B·2H<sub>2</sub>SO<sub>4</sub>,2H<sub>2</sub>O<sub>4</sub>
for the preparation see under 'Quinine Sulphate.'

Short colourless prisms, slowly soluble in cold water, almost insoluble in absolute alcohol.

Cinchonidine hydrochloride B HCl, H2O occurs as fine monoclinic prisms, soluble in 30 of water or 300 of ether at 12°; soluble in alcohol or chloroform. From alcohol fine needles containing 2H,O are obtained,

Other salts are: benzoate B C, H,O, soluble in 333 of water at 10°; hydrobromide B HBr,H2O slightly soluble in water, more soluble in alcohol; dihydrochloride B.2HCl,H<sub>2</sub>O, large prisms, very soluble in water and alcohol; hydrodide B  $HI, H_2O$ , colourless needles, slightly soluble in water; dihydriodide B2HI,H2O, lemon-yellow prisms, intrate B  $\overline{\rm HNO_3, H_2O}$ , needles or large prisms, soluble in 71 of water at 10°, oxalate B<sub>2</sub> H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>,6H<sub>2</sub>O, prisms, soluble in 250 of water at 10°, platinochloride B<sub>2</sub> H<sub>2</sub>PtCl<sub>6</sub>,2H<sub>2</sub>O, small orange prisms, diplatinochloride B H2Pt(I6, H2O, orange-yellow crystalline powder, insoluble in water, salicylate Succinate B<sub>2</sub> C<sub>4</sub>H<sub>6</sub>O<sub>3</sub>, soluble in 77 of water at 18°, succinate B<sub>2</sub> C<sub>4</sub>H<sub>6</sub>O<sub>4</sub>,2H<sub>2</sub>O, small prisms, soluble in 583 of water at 10°, tartrate B<sub>2</sub> C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>,2H<sub>2</sub>O, fine needles, soluble in 1265 of water at 10°, insoluble in Rochelle salt solution, acid tartrate B 2C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>,3H<sub>2</sub>O, long prisms

Quinamine and its Derivatives and Salts—Quinamine C<sub>19</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub> occurs in nearly all cinchona barks to a very small extent (Hesse, Annalen, 1873, 106, 266, Ber 1877, 10, 2157) It accumulates in quinine sulphate mother liquois, from which it may be obtained by precipitating quinine and cinchonidine with Rochelfe salt, precipitating the filtrate with ammonia and extracting the precipitate with ether The ether is washed out with acetic acid, the acid solution neutralised, and while still warm treated with thiocyanate, until on cooling emchanine can no longer be detected. The filtrate is precipitated with NaOH, and the resinous precipitate dissolved in a minimum of boiling 80 pe alcohol, from which quinamine crystallises on cooling One gallon of quinine sulphate mother liquor will yield about 3 grams of quinamine The base crystallises in delicate white anhydrous needles, melting at 172°, and has  $\left[\alpha\right]_{D}^{15} = +94^{\circ}$  for 1.5 p c solution in chloroform. It is nearly insoluble in cold water, soluble in 105 of 80 p c alcohol at 20°, and in 50 of ether (Oudemans, Annalen, 1879, 197, 150), also soluble in petioleum, ether, or benzene. Quinamine differs from the other cinchona bases in being monobasic, and therefore giving but one series of salts It is coloured yellow by warm sulphuric acid, and

orange by nitric acid The chief derivatives of quinamine are .—

Quinamicine C<sub>19</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub>, formed by heating quinamine for a few minutes at 100° with concentrated H2SO4. The product is treated with sodium bicarbonate, which precipitates the quinamicine in flocks, gradually becoming crystalline It melts at 109°, has  $[\alpha]_D^{15} = +3.8^\circ$ for 2 p c. solution in alcohol, and is soluble in alcohol, chloroform, or ether (Hesse, Annalen, 1881, 207, 303) Its salts are amorphous.

Protoquinamicine  $C_{17}H_{20}O_2N_2$  is formed by heating quinamine to  $130^\circ$  with concentrated  $H_2SO_4$ . It is an amorphous brown base.

Quinamidine C<sub>19</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub> is formed when quinamine is heated with dilute mineral acids or with tartaric acid. Ammonia does not precipitate the base from its solution, but NaOH does so It crystallises from alcohol in small white tufts of needles, not very soluble in ether or chloroform. It melts at 93°, and has  $[\alpha]_{\rm p} = +4.5^{\circ}$  for 2 p.c solution in alcohol Heated with concentrated HCl it gives a yellow solution turning to brown, which on dilution turns 10se, with a strong green fluorescence Quinamidine yields a crystalline hydrochloride B  $HCl_1H_2O$ , hydrobromide B  $HBr_1H_2O$ ; oxalate  $B_2H_2C_2O_4AH_2O$ , and a platinochloride  $B_2H_2PtCl_6AH_2O$ . Its hydrochloride gives a purple colour with AuCl3 in aqueous solution

Apoquinamine  $C_{19}H_{22}ON_2$  is obtained by boiling quinamine or quinamicine with HCl (sp.gr. 1 125), or with 25 p.c.  $H_2SO_4$  for 3 minutes (Hesse, Annalen, 1881, 207, 294). It crystallises in colourless scales, melting at 114°, soluble in alcohol, chloroform, or ether alcoholic solution is optically mactive and neutral to litmus. It is a feeble base, but gives a crystalline hydrochloride readily B HCl, 1H2O (lavorotatory), mitrate B HNO3, oxalate B, H2C2O4, H2O, tartrate

$$B_2 C_4 H_6 O_6, \lambda H_2 O$$
;

and platmochloride B<sub>2</sub> H<sub>2</sub>PtCl<sub>6</sub>,2H<sub>2</sub>O

Salts of Quinamine—Quinamine gives a crystalline chlorate B HClO3, perchlorate crystallino chlorate B HClO<sub>3</sub>, perchlorate B HClO<sub>4</sub>, hydrobromide B HBr,H<sub>2</sub>O, thick prisms, very soluble in water or alcohol, hydrochloude B HCl, H2O, colourless pusms, readily soluble in water, hydrodide BH, small colourless prisms, soluble in 71 of water at 16°, nitiate B HNO<sub>3</sub>, prisms, soluble in 16.5 of water at 15°, sulphate, hexagonal plates or short prisms, very soluble in water, and an amorphous yellow platmochloride

decomposing in the light

#### Conquinamine and its Salts

Conquinamine, C<sub>19</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub>, an isomeride of quinamine is found in the alcoholic mother liquor from which quinamine is crystallised, and is separated from it by fractional crystallisation of their nitiates, oxalates, or hydrobromides, the conquinamine salts being the more soluble (Oudemans, Annalen, 1881, 209, 38, Hesse, ibid 62) It crystallises from alcohol in long pusms or pyramids, melting at 123°, and has  $[\alpha]_D^{15} = +204.4^{\circ}$  for 1 pc solution in absolute alcohol. It is soluble in 74 parts of 91 pc. alcohol at 19°; in 74 parts of ether at 15°; m 4·1 parts of benzene at 18°, almost insoluble in 50 p.e. alcohol of in water. It behaves very like quinamine and is monobasic, but its salts are generally the more soluble. Its oxalate, heated to 105°, melts and changes to quinamiene oxalate and then to apoquinamine oxalate. Heated with HCl (sp gr 1 125) it is converted into apoquinamine Its acid solution with AuCl<sub>3</sub> gives a golden precipitate followed by a purple colouration

Salts —Acetate B·C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>, dimetric crystals, soluble in 10.1 of water at 13.5°; chlorate B·HClO<sub>3</sub>, needles, soluble in 104 of water at 16°; perchlorate B HClO<sub>4</sub>, long needles or short prisms, soluble in 396 of water at 16°; formate B·CH<sub>2</sub>O<sub>2</sub>, monoclinic crystals, soluble in 10·8 of water at 15°; hydrochloride B HCl, large octahedra, moderately soluble in water, hydrobromide B HBr, monoclinic prisms, soluble in 25·8 of water at 16°, hydrodide B HI, plates, soluble in 106 of water at 16°; initiate B·HNO<sub>3</sub>, rhombic crystals, soluble in 76 of water at 15°; oxalate B<sub>2</sub> C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>,3H<sub>2</sub>O, rhombic crystals, soluble in 82·3 of water at 17°, quinate B C<sub>2</sub>H<sub>1</sub>O<sub>4</sub>,2H<sub>2</sub>O, prisms, platinochloride B<sub>2</sub>·H<sub>2</sub>PtCl<sub>6</sub>,2H<sub>2</sub>O, amorphous orange-yellow precipitate, sulphate B<sub>2</sub>·H<sub>2</sub>SO<sub>4</sub>, long prisms, readily soluble in water or alcohol

#### Amorphous Bases.

Diquinicine (diconquinine) C<sub>40</sub>H<sub>46</sub>O<sub>3</sub>N<sub>4</sub> Thus amorphous base is the chief constituent of commercial quinoidine, and its constitution is very doubtful. Hesse found the molecular weight by the cryoscopic method to be about one-half of that represented by the above formula (Ber. 1877, 10, 2155) It is prepared by neutralising quinoidine with oxalic acid. thoroughly dehydrating, powdering the oxalate and extracting with cold chloroform Quinicine oxalate remains undissolved, and on adding a little water to the chloroform solution and allowing to stand a short time, cinchonicine oxalate crystallises out, and is filtered off evaporating the chloroform solution to dryness diquinicine remains as an amorphous red mass It is dextrorotatory, gives a fluorescent solution in dilute H2SO4, and gives the thalleroquin re-No crystalline salts have been obtained

Dicinchonicine (dicinchonine) C<sub>38</sub>H<sub>44</sub>O<sub>2</sub>N<sub>4</sub> This base occurs in the bark of C rosulenta and C succirubra, together with the principal cinchona bases It is prepared from the mother liquor, after precipitating the einchonidine as tartrate, by rendering alkaline and extracting The ethereal solution is extracted with ether with dilute acetic acid and the acid solution exactly neutralised with NaOH and fractionally precipitated with potassium sulphocyanide The dicinchonine thrown down, mainly in the middle fractions, is again put through ether and crystallised as hydrochloride (Hesse, Annalen, 1885, 227, 153) The free base is a yellow amorphous body with a strong alkaline reaction, m p  $40^{\circ}$ ,  $[\alpha]_{D}^{15} = +91.7^{\circ}$  for a 1.52 p c solution in 97 pc alcohol It is very soluble in ether, acetone, alcohol, chloroform, or benzene, ınsoluble ın alkalıs The molecular weight determined by the cryoscopic method indicates half that of the above formula It gives a crystalline hydrochloride B 2HCl, easily soluble platinochloride B 2H<sub>2</sub>PtCl<sub>8</sub>,4H<sub>2</sub>O, orange flakes, hydrodide, compact crystals, readily soluble in water, insoluble in NaCl or KI solution, and oxalate, easily soluble prisms, but its other salts are amorphous On heating with HCl to 140°-160° it is converted into an isomeride, diapocinchoninine, an amorphous base soluble in ether, alcohol, or chloroform, having  $[a]_{D}^{15} = +20^{\circ}$  in alcoholic solution (Hesse, Annalen, 1880, 205, 333)

Paricine C<sub>16</sub>H<sub>18</sub>ON<sub>2</sub> was discovered by Winckler (Jahresber. 1845, 27, 338) in the barks

of Buena hexandra and C. lutea. Later, Hesse found it in C. succirubra of Darjeeling (Pharm J. 1870, [3] 1, 344; Annalen, 1873, 166, 263). It is separated from the other cinchona bases by intric acid, paricine nitrate being insoluble. The free base is obtained by suspending the nitrate in alcohol, decomposing with ammonia, and crystallising from petioleum ether. It is purified by dissolving in dilute HCl, decolourising with animal black and precipitating with ammonia. The pure paricine thus obtained is a yellow amorphous powder, soluble in water, alcohol, ether, and petroleum ether; in p. 116°. Its salts are amorphous

Javanine was found by Hesse among the amorphous bases of C. Calisaya Javanica. It separates from water in rhombic scales, very soluble in ether, and gives an intense yellow colour with dilute H<sub>2</sub>SO<sub>4</sub>—It yields a crystalline oxalate (Ber. 1878, 10, 2162).

# REMIJIA BASES.

Cupreine and its Derivatives and Salts -Cuproine  $C_{19}H_{22}O_2N_2$  occurs in cuprea bark (Remijia pedunculata) together with the other cinchona alkaloids, except cinchonidine Its molecular combination with quinine was at first mistaken for a new alkaloid, and called homoquinine (Howard and Hodgkin, Chem. Soc Trans 1882, 41, 66, Paul and Cownley, Pharm J 1881, 12, 497) Three years later, Paul and Cownley isolated the new alkaloid cupreme from homoquinine (Pharm J 1884, 15, 221, 401, Hesse, Annalen, 1884, 225, 98, ibid 226, 242, 1885, 230, 72) Cupreine completely crystallises out with the quinine when a sulphuric acid solution of the bases is neutralised It is separated from the quinine by dissolving this rough sulphate in acid, precipitating the solution with eacess of NaOH, and extracting the quinine with ether. The alkaline aqueous hquor on neutralising with H<sub>2</sub>SO<sub>4</sub> throws out cupreinc sulphate. The base crystallises from ether in concentric prisms containing 2H2O, or from alcohol as a crystalline powder containing  $\frac{1}{3}H_2O$ , dehydrating at  $120^\circ$  It is spaningly soluble in ether or chloroform, easily soluble in alcohol, m p 198°,  $[a]_{\rm p}^{16} = -175 \ 3^{\circ}$  for 1 5 p c. solution in 97 p.c. alcohol Its solution in dilute  $\rm H_2SO_4$  does not fluoresce, but gives the thalleroquin reaction, and with ferric chloride a reddish-brown colour is obtained is the lower homologue of quinine having a hydroxyl group replacing the methoxy group of quinine in the para-position of the quinoline ring. This —OH group has phenolic properties which enables cupreme to form salts with the alkalis, and strong compounds with the other cinchona bases. It is also dibasic, forming readily crystalline salts with acids Heated with HCl (sp gr 1 125) to 150° it is converted into an isomer, apoquinine, the same body being obtained from quinine under similar treatment (q v)When sodium cupreinate is treated in methyl alcohol solution with methyl chloride or methyl nitrate, quinine (methyl-cupreine) is obtained (Grimaux and Arnaud, Compt rend. 1891, 112, 374, 766; 114, 548, 612) By this reaction the following higher homologues of quinine were obtained by Grimaux and Arnaud :--

Quinethyline  $C_{19}H_{20}N_2(OH)OC_2H_5$ , a white powder, m.p.  $160^\circ$ ,  $[a]_D = -169 \cdot 4^\circ$ ; sulphate  $B_2 \cdot H_2SO_4, H_2O$ , plates, soluble in 397 of water at  $15^\circ$ .

**Quinpropyline**  $C_{19}H_{20}N_2(OH)OC_3H_7$ , a white powder, m.p.  $164^\circ$ , sulphate  $B_2 \cdot H_2SO_4$ ,  $1\frac{1}{2}H_2O$ , needles, soluble in 454 of water at 13°.

needles, soluble in 454 of water at 13°.

Quinisopropyline C<sub>19</sub>H<sub>20</sub>N<sub>2</sub>(OH)OC<sub>3</sub>H<sub>7</sub>, a white powder, m.p 154°; sulphate B<sub>2</sub>·H<sub>2</sub>SO<sub>4</sub>, H<sub>2</sub>O, needles, soluble in 367 of water at 10°

Quinisoamyline  $C_{19}H_{20}N_2(OH)OC_5H_{11}$ , amorphous, mp 167°, sulphate  $B_2$   $H_2SO_4$ , needles, soluble in 4170 of water at 115°.

These ethers all give fluorescent solutions in dilute  $\rm H_2SO_4$ , whereas cupreine does not — They are strongly febrifugal, but more toxic than quinine

Homoquinine. The molecular compound of quinine and cupreine named homoquinine

C<sub>20</sub>H<sub>24</sub>O<sub>2</sub>N<sub>2</sub> C<sub>19</sub>H<sub>22</sub>O<sub>2</sub>N<sub>2</sub>,1, 2, or 4H<sub>2</sub>O may be obtained by dissolving equi-molecular proportions of the two alkaloids in alcohol and evaporating to dryness, or by mixing 1 molecule of quintine hydrochloride with 1 molecule of sodium cupremate (Hesse, *lc*) It behaves like a simple alkaloid, alkalis alone breaking it up, and forms well-defined salts. It melts at 177°, is lævorotatory, and dissolves with difficulty in ether, but readily in alcohol

For Hydrocupreme, see 'Hydroquinne.' Salts of Cupreine —Acetate B C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>,2H<sub>2</sub>O, fine needles, soluble in 85 of water at 17°; hydrobromide B HBr,H<sub>2</sub>O, needles, soluble in 122 of water at 16°, dhydrobromide B 2HBr, soluble in 125 of water at 16°; hydrochloride B·HCl,H<sub>2</sub>O, small needles, soluble in 53°5 of water at 16°; dhydrochloride B·2HCl, short prisms, readily soluble in cold water, hydrochde B HI, soluble in 106 6 of water at 16°; dhydrochloride B 2HI,H<sub>2</sub>O, orange crystals, soluble in 15 of water at 16°, intrate B HNO<sub>3</sub>,2H<sub>2</sub>O, needles, soluble in 86 of water at 16°; dintrate B 2HNO<sub>3</sub>,H<sub>2</sub>O, yellow transparent crystals, soluble in 12 of water at 17°; oxalate B C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>,2H<sub>2</sub>O, colouiless crystals, soluble in 407 of water at 18°; platinochloride

B2H2PtCl6,4H2O

yellow amorphous precipitate; diplatinochloride  $B\cdot H_2PtCl_6, H_2O$ , orange needles; sulphate  $B_2H_2SO_4, 6H_2O$  (Hesse), anhydrous (Howard and Chick, J. Soc Chem Ind 1909, 28, 53), needles, soluble in 813 of water at  $17^\circ$ ; disulphate  $B\cdot H_2SO_4, H_2O$ , thick prisms, soluble in  $70\cdot 4$  of water at  $16^\circ$ ; tetrasulphate

## B·2H<sub>2</sub>SO<sub>4</sub>,3H<sub>2</sub>O

short silky needles, very soluble in water, less soluble in alcohol; tartrate  $B_2$   $C_4H_6O_6,H_2O$ , white needles, soluble in 517 of water at 16°.

Crachonamine and its Salts—Cinchonamine C<sub>19</sub>H<sub>24</sub>ON<sub>2</sub> occurs in the bark of Remijia Purdicana, together with cinchonine, concusconine, chairamine, chairamidine, conchairamine, and conchairamidine. It is obtained by extracting the powdered bark with hot dilute H<sub>2</sub>SO<sub>4</sub>, the acid extract is precipitated by ammonia, and the precipitated bases filtered and dried. This dried mass is then extracted with boiling alcohol, the alcohol extract concentrated, cooled, and treated with HNO<sub>3</sub> (sp gr. 1 200) until decidedly acid. Cinchonamen intrate is

precipitated and is purified by recrystallisation from boiling water (Arnaud, Compt rend 1881, 93, 593; 1885, 97, 174; Hesse, Annalen, 1884, 225, 218).

The base crystallises in needles or orthorhombic prisms, melting at 185°, and subliming,  $[\mathfrak{a}]_D^{15} = +121\cdot1^\circ$  for 2 pc solution in 97 pc alcohol. It dissolves in alcohol or chloroform, and in 100 of ether at 17°; slightly soluble in benzene, petroleum ether, CS2, or water. It is a strong base, giving well-defined salts, and differs from its isomerides, hydrocinchonine and hydrocinchonidine, in being readily attacked by permanganate at 0° in  $H_2SO_4$  solution. It differs from the cinchona bases in the great solubility of its sulphate and the insolubility of its nitrate. Arnaud proposed the use of this alkaloid for the estimation of nitrates, but the method is limited in its application (Ann. Chim. Phys 1890, 19, 93, Howard and Chick, J. Soc. Chem. Ind. 1909, 28, 53). Treated with excess of nitric acid (sp.gr. 106) a yellow amorphous explosive body, dinitrocinchonamine.

 $C_{1y}H_{22}(NO_2)_2ON_2$ 

is formed, soluble in alcohol, chloroform, or ether, and melting at 118°. This base is insoluble in strong HCl, but soluble in the dilute acid; and on heating with strong HCl does not give methyl chloride. It is strongly febrifugal, but also very toxic

Salts—Citrate B<sub>2</sub> C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>, prisms, soluble in 51 of water at 15°, hydrobromide B·HBr, long needles, soluble in 600 of water at 15° hydrochloride B HCl, soft laminæ, or B HCl, H, O, transparent cubes, both soluble in 200 of water at 27° (Howard and Perry, J Soc Chem Ind. 1905, 24, 1281), hydrodide B HI, long prisms or plates, soluble in 950 of water at 15° hyposulphite B·H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, prisms, slightly soluble in water; malate B<sub>2</sub> C<sub>4</sub>H<sub>6</sub>O<sub>5</sub>, H<sub>2</sub>O, flaky crystals, soluble in 100 of water at 15°; intrate B HNO<sub>3</sub>, short prisms, soluble in 500 of water at 15° platinochloride B<sub>2</sub> H<sub>2</sub>Pt('l<sub>6</sub>, semi-crystalline yellow precipitate, insoluble in water, picrate B  $C_6\dot{H}_2(N\dot{O}_2)_3$  OH,  $\frac{1}{2}H_2O$ , yellow flocks, insoluble in water, "sulphate B. H. SO4, transparent prisms, very soluble in water, less soluble in alcohol, disulphate BH2SO4, octahedra or prisms, very soluble in water, sulphocyanide B HCNS, laminæ or short prisms, insoluble in water, taitrate  $B_2$  ( ${}^t_4H_6O_8$ , crystalline powder or thick prisms, soluble in 87 of water at 15°.

Concusconine  $C_{23}H_{26}O_4N_{\circ}$ . To produce this base Hosse extracted Remijia Purdicana with alcohol, evaporated the extract to dryness, added a large excess of NaOH, and extracted with ether. On shaking the ethereal extract with excess of dilute H2SO4 the concusconine is precipitated as sulphate together with the chairamine isomerides, the cinchonine and cinchonamine remaining in the acid solution rough sulphate is boiled with Na<sub>2</sub>CO<sub>3</sub> solution. and the freed bases washed, dired, weighed, dissolved in boiling alcohol, and I part of H<sub>2</sub>SO<sub>4</sub> added for every 8 parts of alkaloid. Concusconine sulphate separates out completely on cooling (Annalen, 1884, 225, 234). The base cooling (Annalen, 1884, 225, 234). forms colourless monoclinic prisms containing 1H<sub>2</sub>O; mp, after drying, 206°-208° with decomposition;  $[a]_{p}^{15} = +40.8^{\circ}$  for 2 p.c. solution

in 97 p.c. alcohol. It is slightly soluble in strong alcohol, less so in dilute, very soluble in chloroform, ether, or benzene. A solution in acetic acid or HCl gives a dark-green colour with concentrated HNO3. This reaction is also given with the chairamine isomerides. Concentrated H<sub>2</sub>SO<sub>4</sub> gives a blush-green solution, turning olive-green on warming. It is a weak base, its salts are mostly gelatinous, but it gives a crystalline sulphate B<sub>2</sub>·H<sub>2</sub>SO<sub>4</sub> in small white prisms almost insoluble in water or alcohol; an oxalate B<sub>2</sub>·C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>; and a platinochloride B·H<sub>2</sub>Pt(l<sub>8</sub>. Concusconine is isomeric with brucine It contains two methyoxy groups

Chairamine C<sub>22</sub>H<sub>26</sub>O<sub>4</sub>N<sub>2</sub>. On adding a little concentrated HCl to the alcoholic mother liquor of the concusconine sulphate (above), chairamine hydrochloride crystallises out in small needles. The base crystallises from dilute alcohol in slender white needles or from strong alcohol in thick prisms containing 1H2O, which melt at 140°, and, after drying, at  $233^{\circ}$ ,  $[\alpha]_D = \text{about}$ + 100 in 97 p.c. alcohol It is readily soluble in chloroform or ether, and in 540 of 97 p c alcohol at 11° Its acctic acid solution gives a dark-green colour with concentrated HNO<sub>3</sub> In concentrated H2SO4 the base gives a colourless solution, slowly turning dark green It gives a hydrochloride B·HCl, H2O, needles, soluble in water, slightly soluble in alcohol—a platino-chloride  $B\cdot H_2PtCl_6, 2H_2O$ , and a sulphate B<sub>2</sub>·H<sub>2</sub>SO<sub>4</sub>,8H<sub>2</sub>O

Conchairamine C<sub>22</sub>H<sub>26</sub>O<sub>4</sub>N<sub>2</sub>. On adding a little solution of pota-sium sulphocyanide to the chairamine hydrochloride mother liquor (above), conchairamine sulphocyanide 18 obtained The base is liberated with NaOH and re-Thick crystallised from alcohol. colourless prisms BH,OC,H,OH are obtained, melting at 82°-86°, and losing their water and alcohol of crystallisation at 120°, being at the same time converted into an amorphous form An acetic acid solution of the base precipitated with ammonia yields white crystalline flocks containing  $1\rm{H}_2O$ , melting and dehydrating at  $110^\circ$  The anhydrous base melts at 120°, and has  $\lceil \alpha \rceil_D^{15} = +68.4^\circ$  for 2 p c solution in 97 p c alcohol. It is soluble in chloroform or ether and in boiling alcohol In concentrated H<sub>2</sub>SO<sub>4</sub> the base gives a brown solution turning dark green It gives a dark-green coloui in acetic acid solution with concentrated HNO3 A solution of the base in alcohol is neutral to litmus Its salts are readily crystalline, the chief being hydrochloride B HCl; hydriodide B HI, $H_2O$ , platinochloride B  $H_2$ PtCl $_6$ 5 $H_2O$ , sulphocyanide B HCNS, H<sub>2</sub>O; sulphate

### B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub>,9H<sub>2</sub>O

Chairamidine ('22H26O4N2. When more potassium sulphocyamde is added to the filtrate from the conchairamine sulphocyamde until the dark colour of the solution becomes light brown a pitch-like mass separates. The mother liquor is now treated with excess of ammonia and shaken out with benzene; the benzene is extracted with a saturated solution of ammonium sulphate, when a mixture of chairamidine and conchairamidine sulphate is precipitated. The mixed

sulphates are dissolved in boiling water and cooled, a gelatinous mass precipitating, part of which becomes crystalline after standing some days. On now warming to 40°, the gelatinous part redissolves, leaving the crystals of conchair-amidine sulphate. On re-cooling chairamidine sulphate falls again as a jelly, which is drained off, re dissolved in water, and the base precipitated with ammonia.

Chairamidine is a white amorphous powder containing IH<sub>2</sub>O, melting, when dry, at  $126^\circ$ -128° with decomposition, and has  $[a]_D^{15} = +7.3^\circ$  for 3 pc solution in 97 pc alcohol. It is insoluble in water, readily soluble in alcohol, ether, chloroform, or benzene. It is a feeble base, its alcoholic solution being neutral to lithius. It gives a yellow solution in concentrated H<sub>2</sub>SO<sub>4</sub>, slowly turning green. Animal black completely removes it from its solution in acetic acid. Its salts, which are amorphous, have been little studied

Conchairamidine  $C_{22}H_{26}O_4N_2$ . This base is obtained as crystalline sulphate in the preparation of chairamidine (above). The free base crystalline, contains  $1H_2O$ , melts at  $114^\circ-115^\circ$  after drying, and has  $\lfloor \alpha \rfloor_D^{15} = -60^\circ$  for 3 p c solution of the anhydrous base in 97 p c alcohol it is very soluble in alcohol, ether, chloroform, benzene, or acctone. It is a feeble base and neutral to litmus. It gives a dark-green colour in concentrated  $H_2SO_4$ . It gives crystalline salts, the chief being hydrochloride B  $HC_13H_2O$ , platinochloride. B  $H_2PtCl_6,5H_2O$ , sulphate  $B_2H_2SO_4,14H_2O$ 

# Cusco Bases

Aricine C<sub>23</sub>H<sub>26</sub>O<sub>4</sub>N<sub>2</sub>. This alkaloid was discovered in Cusco bank by Pelletier and Corriol (J. Pharm. Chim. 1829, 15, 575), and is identical with Manzini's cinchovatine obtained this base by extracting the powdered bark with alcohol, treating the alcoholic extract with excess of NaOH, and extracting with ether. The ethereal solution was shaken out with acctic acid, and the acid solution partially neutralised with ammonia, when aricine acetate separates out The filtrate mixed with a strong solution of ammonium sulphate, yielded cusconine sulphate (Annalen, 1878, 185, 296) Moissan and Landrin, after powdering the bank, treating it with lime and 40 pc NaOH and re-drying, extract the alkaloids with ether. On washing out the ethereal solution with dilute H<sub>2</sub>SO<sub>4</sub> aricine sulphate falls out, is decomposed with ammonia, and the free base crystallised from alcohol (J. Pharm Chim 1890, [5] 21, 337).

Ariene forms large colouriess prisms, insoluble in water, soluble in 100 of 90 p c alcohol, or in 33 of ether, at 15°; very soluble in chloroform. It melts at 188° and has  $\lceil \alpha \rceil_D^{15} = -58$  2° for 2 p.c solution in 97 p c alcohol. In account in 3 dextrorotatory,  $\lceil \alpha \rceil_D = +14$  5°. It gives a yellow-green colour in strong  $H_2SO_4$ , turning a fine deep blue on addition of ammonium molybdate, which on warming turns olive-green, reverting to blue on cooling. This reaction is characteristic for arieine and cusconino. Nitric acid colours arieine deep green, and dissolves

CINNABAR 279

it to a greenish-yellow solution. Aricine is not | (uncorr), with decomposition, very soluble in is a feeble base, its salts partly dissociating in water

The chief salts are: acetate

# B·C,H,O,3H,O

small white granular crystals, almost insoluble in cold water, slightly soluble in hot; citrate B  $\rm C_6H_7O_6$ , colourless needles, slightly soluble in water, hydrochloride B HCl,2H<sub>2</sub>O, colourless prisms, slightly soluble in cold water: hydriodide B·Hl, colourless prisms, almost insoluble in water; oxalate, granular white crystalline powder; binoxalate B C<sub>2</sub>H<sub>2</sub>O<sub>4</sub>2H<sub>2</sub>O, prisms, rapidly changing to ihombohedra, soluble in 2025 of water at 18°, platinochloride B<sub>2</sub> H<sub>2</sub>PtCl<sub>6</sub>,5H<sub>2</sub>O, orange-yellow precipitate, sparingly soluble in water, salicylate

B C, H, O, 2H, O

pale yellow pulverulent precipitate, sparingly soluble in water, easily soluble in alcohol sulphate B<sub>2</sub> H<sub>2</sub>SO<sub>4</sub>, fine needles, fairly soluble or cold water, disulphate BH, SO<sub>4</sub>, small crystalline groups, very slightly soluble in cold water, sulphocyanide B HCNS, small colourless

prisms, very slightly soluble in water Cusconine  $C_{23}H_{26}O_4H_2$ . The rough cusconine sulphate, obtained from the aricine acetate mother liquid (above), on precipitation with ammonia as the base and crystallising from ether, is obtained as white lamine. From alcohol or acetone larger crystals may be obtained The crystallised base contains 2H,O, which it loses at 80°, the anhydrous base melting at 110°. It has  $[a]_{D}^{15} = -54.3^{\circ}$  for a 2 p c solution in 97 p.c. alcohol The base dissolves in 35 parts of ether at 15°, more easily in alcohol or acetone, is very soluble in chlorotorm, but is insoluble in water. It gives the same colour reactions with H.SO4 and ammonium molybdate as arreine It is a feeble base, its salts having an acid reaction Most of the salts are gelatinous or amorphous Hesse (Annalen, 1878, 185, 296) prepared the following salts. aurichloride, dirty yellow amorphous precipitate readily decomposing; hydrobromide, white ın water, amorphous precipitate, soluble hydrochloride, uncrystallisable, but forms with mercuric chloride a pulverulent precipitate B·HCl  ${\rm HgCl_2,2H_2O}$ , hydrodide, pale yellow amorphous precipitate, freely soluble in water, platinochloride B2, H2PtCl6,5H2O, amorphous flocculent dark yellow precipitate; sulphate  $B_2H_2SO_4$ , lamine from alcohol, disulphate, gelatinous and uncrystallisable; thiocyanate B HCNS,2H2O, pale yellow amorphous powder

Cusconidine, an amorphous base precipitated from cusconine mother liquors with alkalis; it falls as pale yellow flakes which, after washing, mass together, and on drying form an amorphous mass (Hesse, Annalen, 1880, 200, 303)

Cuscamine. From an acetic acid solution of the total alkaloids of Cusco bark nitric acid precipitates insoluble cuscamine and cuscamidine nitrates. On conversion of the nitrates to oxalates, and washing with water, cuscamidine oxalate dissolves, leaving cuscamine oxalate The free base is obtained by treating with NaOH, and crystallising from ether and alcohol Cuscamine forms colourless prisms, m.p. 218°

bitter in flavour, but slightly astringent. It ether, moderately soluble in alcohol. It gives a vellow colour with concentrated H, SO4, turning brown on warming; on adding molybdic acid a bluish-green colour is obtained, turning brown on heating, and violet-brown on re-cooling With nitric acid a yellow colouration is obtained It readily yields crystalline salts (Annalen, 1880, 200, 304).

Cuscamidine is obtained by concentration of its oxalate solution obtained as above closely resembles cuscamine in chemical and physical properties, but its salts are more soluble

#### ASPIDOSPERMA BASES.

Paytine  $\rm C_{21}H_{24}ON_2$  is extracted from the white bark of Payta (Peru) by means of alcohol The alcoholic extract is diled, mixed with NaOH, shaken with ether, the ethereal solution extracted with dilute H<sub>2</sub>SO<sub>4</sub>, the acid liquor treated with animal charcoal and neutralised On adding KI solution ammonia until no more precipitation occurs, paytine hydrodide falls as a white precipitate, becoming dense, yellow, and crystalline. The free base crystallises in fine prisms containing IH,O, and is soluble in alcohol, ether, benzol, ligroin, and chloroform, slightly soluble in water; mp  $156^{\circ}$ ,  $[\alpha]_{j} = -495^{\circ}$  in 0 45 pc alcoholic solution Án alcoholic solution is alkaline to litmus
bitter flavour, but is not toxic

It gives curious characteristic colour reactions. With centrated  $\mathrm{HNO_3}$  it gives a colourless solution passing through gainet-red to yellow. Its hydrochloride yields with platinum chloride a dark yellow precipitate, dissolving in HCl to a reddish-brown solution, turning blue and giving a blue precipitate Bleaching powder produces in acid solution a dark red colour, turning blue and then pale yellow with precipitation Gold chloride gives a purple colour and precipitate; mercuric chloride a yellow amorphous precipi-It gives several crystalline salts, the chief being the hydrochloride B HCl, prisms, soluble in 16 6 of water at 15°. Heated with soda-lime it yields a non-nitrogenous body, paytone (Hesse, Annalen, 1870, 154, 287, 1873, 166, 259, 1882, 211, 280, Ber 1881, 13, 2308, Wulfsberg, Pharm Zeit 1880, 546, Arata, Gazz. chim. ital 1881, 11, 246).

Paytamine  $C_{21}H_{24}ON_2$  is an amorphous alkaloid precipitated from the iodide mother liquor of paytine hydriodide (see above) by alkalis It is readily soluble in other, gives a purple colour with gold chloride, but differs from paytine in yielding a soluble hydriodide, and by not giving paytone on heating with soda-lime B F H & O. C.

CINEOL v CAMPHORS, SANTONICA. For method of estimating cineol in volatile oils, see Dodge, Analyst, 1912, 461 For its solubility m water, see Earle, J. Soc Chem. Ind. 1918, 274 T.

CINNABAR. Native mercury sulphide (HgS), and the only ore of this metal. Its crystals are rhombohedral, with perfect cleavages parallel to the faces of the hexagonal prism; they are of interest crystallographically in presenting the same type of trapezohedral symmetry as quartz, 280 CINNABAR.

and, like this, they rotate the plane of planepolarised light. They are bright-red in colour, often transparent, and have a brilliant adamantine to sub-metallic lustre; the scarlet streak is characteristic. Sp gr. 8:1; hardness, 2-21. The refractive indices are high . for the ordinary ray 2.854, and for the extraordinary ray 3.201 for red light. The massive ore, which may be compact or earthy in texture, is of the same colour, but sometimes it is liver-brown or black. owing to the admixture of clay or organic matter, as in the varieties called respectively hepatic cinnabar and coral-ore (German, Korallenerz). Cinnabar is very sporadic in its distribution, and no mineral containing mercury has yet been found in the British Isles It is mined at Almaden in Spain, Monte Amiata in Tuscany, Idria in Carniola, New Almaden in California, in the provinces of Kwei-chou and Hunan, in central China, and in Mexico and

Cinnabar and two other crystalline modifications of mercuric sulphide have been obtained artificially by E T Allen and J L (Trenshaw (1912) Cinnabar is the stable form at all temperatures up to its sublimation point, about 580°.

L J. S.

CINNABAR, AUSTRIAN, v CHRQMIUM

CINNAMALDEHYDE, Cunamic aldchyde C.H. CH CH CHO, is contained, together with a hydrocarbon, in oil of cinnamon and oil of cassia. It may be extracted from oil of cinnamon by shaking the oil with a concentrated solution of sodium bisulphite, filtering off the crystalline bisulphite compound, washing the latter with alcohol and distilling it with dilute sulphuric acid, when the cinnamaldehyde passes over with the steam The cinnamaldehyde is extracted from the aqueous distillate with other, and, after expelling the ether, is purified by distillation under a pressure of from 40 to 50 mm (Peine, Ber 1884, 2109) It may be extracted from cassia oil by adding to it the barium salt of sulphanilic acid, filtering off the crystalline compound thus formed and steam distilling, whereby the compound is resolved into its components and cinnamaldehyde distils over (Ch. Fab. von Heyden Akt -Gesel D. R. P. 124229, Chem Zentr 1901, 11 903) It is chained by allowing a mixture of 10 parts of benzaldehyde, 15 of ordinary aldehyde, 900 of water, and 10 of a 10 pc solution of caustic soda to stand at a temperature of about 30° for from 8 to 10 days, shaking from time to

### $C_6H_5$ ·CHO+CH<sub>3</sub>·CHO= $C_6H_5$ ·CH CH CHO+ $H_2O$

The solution is extracted with ether and the ethereal extract treated as in the foregoing method (Krszysica, Ber 1884, 2117). It is best prepared by the condensation of benzaldehyde and ordinary aldehyde at a low temperature by means of a concentrated alkali solution. The alcoholic solution of the aldehydes, cooled to  $-10^{\circ}$ , is agitated with a 25–30 p c solution of caustic soda, previously cooled to the same temperature. Under these conditions the formation of resins is avoided (Bochringer and Sohne, Eng. Pat. 10003; J. Soc. Chem. Ind. 1897, 463). Also formed when a mixture of calcium cinnamate and calcium formate is distilled (Piria, Annalen, 100, 105).

It is a colourless oil with a pleasant aromatic smell of cinnamon. It decomposes when distilled under ordinary pressures, especially with access of air, but may be distilled without decomposition in a current of steam or under reduced pressure. It boils at 120°-130° under a pressure of 20 mm (Peine, Ber. 1884, 2110); at 209 5° under 250 mm. (Perkin, Chem. Sec. Trans. 1896, 1247) and solidifies at  $-7.5^{\circ}$  (Pictet, Compt rend. 119, 955); sp.gr. 1 0497  $20^{\circ}/4^{\circ}$  (Bruhl, Annalen, 235, 18).

By oxidation it yields first cinnamic acid and afterwards benzaldehyde and benzoic acid. Direct reduction does not lead to cinnamyl alcohol as the aldehyde polymerises. Reduction with the zinc-copper couple gives rise to hydrocinnamoin C<sub>2</sub>H<sub>2</sub>(OH)<sub>2</sub>(CH·CHPh)<sub>2</sub> (Thiele, Ber. 1899, 1296) It gives the usual reactions of the aldehydes The diacetyl derivative of the aldehyde, obtained by the action of acetic anhydride, can, however, be easily reduced by iron in acetic acid solution, and by subsequent saponification the alcohol is obtained (Barbier and Leser, Bull Soc chim 1905, 858)

Estimation.—To estimate cinnamic aldehyde in oil of cinnamon or oil of cassia, 10 c c. of the oil are heated on the water-bath, and a solution of sodium bisulphite added in small quantities, time being allowed for the solid to liquefy between each addition. The precipitate is filtered off and weighed (Schimmel, Chem. Zentr. 1892, 1. 92). Hanus (J. Soc. Chem. Ind. 1903, 1154) estimates cinnamaldehyde by means of its semioxamazone. See also-Burgess, Analyst, 29, 78.

It may be estimated colorimetrically by the reaction it develops on treatment with sulphuric acid and isobutyl alcohol (Jellenberg)

CINNAMEIŃ v Balsam of Peru, art

CINNAMIC ACID  $\mathrm{C_6H_5}$  CH.CH COOH (Acide cinnamique, Fr., Zimmtsaure, Ger.)  $\beta$ -Phenylacrylic acid. The separation of a solid acid from oil of cinnamon on keeping was first observed towards the end of the 18th century, but the acid was confounded with benzoic acid, until Bizio, in 1826, showed that it was a distinct acid. It was first investigated by Dumas and Peligot (Annalen, 14, 56).

Occurrence—In liquid storax, partly free, partly as cunnamyl cunnamate (styracin), in tolu and Peru balsams together with benzoic acid and benzyl cunnamate; in some sorts of gum benzoin, in Barbados aloes; and in the leaves and stalks of Globularia aliquim and G. vulgaris, in the leaves of Enkunthus, japonicus, and as esters in many varieties of guttapercha.

and as esters in many varieties of guttapercha Formation and Preparation—It is formed by heating benzaldehyde with acetyl chloride at 120°-130° (Bertagnini, Annalen, 100, 126):

 $C_6H_5$  CHO+CH<sub>3</sub>·COCl= $C_6H_5$ ·CH CH CO<sub>2</sub>H+HCl

Perkin showed that it might be more readily obtained by heating a mixture of benzaldehyde, acetic anhydride, and anhydrous sodium acetate

 $C_8H_5$  CHO+(CH $_3$ ·CO) $_2$ O= $C_6H_5$ ·CH CH CO $_2$ H+ $C_2$ H $_4$ O $_2$  (Chem. News, 32, 258, Chem. Soc. Trans. 1877, 388) Edeleano and Budishteano (Bull. Soc. chim 3, [3] 191) state that by heating 1 part of benzaldehyde, 1 part of acetyl chloride, and 3 parts of sodium acetate for 24 hours at 160°, and

subsequently extracting with alkali and precipitating the cinnamic acid with hydrochloric acid, the cinnamic acid is obtained in quantitative yield. The cinnamic acid employed in the preparation of artificial indigo was at first manufactured by this method; but the process was afterwards improved by Caro (D. R. P. 17467 and 18232; Frdl. 1. 26), who showed that cinnamic acid could be prepared by heating benzal chloride with dry sodium acetate, thus avoiding the use of the expensive acetic anhydride

 $\begin{array}{l} {\rm C_6H_5~CHCl_2 + 2CH_3~CO_2Na} \\ {\rm = C_6H_5~CH:CH~CO_2H + 2NaCl + C_2H_4O_2} \end{array}$ 

1 part benzal chloride, and from 2 to 3 parts of finely powdered fused sodium acetate (for which potassium acetate or a mixture of the potassium and sodium salts may be substituted) are heated at 180°-200° for from 10 to 20 hours in an autoclave furnished with a mechanical The melt is mixed with water, rendered alkaline with caustic soda, and steam-distilled to remove volatile oils, after which the solution in the retort is filtered hot, and the cinnamic acid is precipitated with hydrochloric acid, filtered off after allowing the solution to cool, pressed, and finally purified by recrystallisation from boiling water or alcohol A method patented by the Farbwerke vorm. Meister, Lucius und Bruning (D. R. P. 18064), in which lead acetate is substituted for sodium acetate, offers no advantages over the foregoing

Cinnamic acid may be prepared from benzylideneacetone C<sub>6</sub>H<sub>5</sub> CH CH CO CH<sub>3</sub>, which is readily obtained by the condensation of benzaldehyde with acetone under the influence of dilute caustic soda (Claisen, Ber 14, 2471) When this compound is oxidised by warming it with sodium hypobromite, it is converted into cinnamic acid (Farbwerke vorm Meister, Lucius und Bruning, D. R. P 21162; Frdl. 1 28)

By employing substituted benzaldehydes, benzal chlorides, or benzylidene-acetones in the foregoing reactions, substituted cinnamic acids may be obtained.

It may also be prepared from benzaldehyde and malonic acid in the presence of ammonia or amines; in the place of benzaldehyde the benzylidene compounds of certain amines, such as hydrobenzamide, may be used (Knœvenagel, Ber 1898, 2692, Chem Zentr 1898, ii 695; D R P 97735); or by heating benzylidene diacetate with glacial acetic acid and sodium acetate for 10 hours at 160°-180° (Nef, Annalen, 1897, 298, 309)

Cinnamic acid may be extracted from storax by boiling in a retort 15 kilos of storax with 2 litres of caustic soda of 24°B until no more oily drops distil over with the steam. In this process the styracın is hydrolysed, forming sodium cinnamate and cinnamyl alcohol, the latter being volatile with the steam. The aqueous liquid in the retort is separated and the residue repeatedly extracted with boiling water, after which the cinnamic acid is precipitated from the united aqueous solutions by hydrochloric acid and purified by recrystallisation (Beilstein and Kuhlberg, Annalen, 163, 123); Claassen (J. Soc. Chem Ind 1897, 932) uses sodium carbonate instead of caustic soda Precipitated cinnamic acid may be readily freed from adhering resin by recrystallising it from hot light petroleum.

Properties and Reactions -Forms slender needles or large transparent prisms melting at 133°. Sublimes in absolute vacuum at 108° (Liebermann, Ber. 1900, 33, 2402) Boils at 300° (corr) If rapidly distilled, it scarcely suffers any decomposition, but by slow heating it is broken up into stryolene C.H. CH. and carbon dioxide Sparingly soluble in cold, more readily in boiling water; easily soluble in alcohol. Forms crystalline salts closely resembling those of benzoic acid Oxidising agents convert it first into benzaldehyde (distinction from benzoic acid) and afterwards into benzoic acid, the final products when potassium permanganate is used being phenylglyceric acid, benzaldehyde, and benzoic and oxalic acids (Fittig and Ruer, When taken internally it Annalen, 268, 27) is oxidised to benzoic acid, and appears in the urine as hippuric acid. By fusion with caustic potash it yields a mixture of benzoate and It unites with nascent hydrogen to form hydrocinnamic (β-phenyl-propionic) acid; with bromine to form a dibromide, and with the hydracids to form 3-halogen-hydrocinnamic acids—thus \(\beta\)-chlor-hydrocinnamic acid C<sub>\(\beta\)</sub>H<sub>\(\beta\)</sub> CHCl CH, CO,H

Cinnamic acid condenses with hydrocarbons; thus with benzene and sulphuric acid, it forms phenylhydrindone (Liebermann and Hartmann, Ber 1892, 2124), with phenols, phenol itself yielding hydroxyphenylcoumarin (L and Hibid 1892, 957), and with sulphur to form diphenylthiophene For its detection, see Schenk & Burmeister, Pharm Zeit 1915, 60, 213; Analyst, 1915, 409

Esters. Methyl cinnamate White crystalline solid, m p 36°; b p 263° Ethyl cinnamate may be obtained direct

Ethyl cinnumate may be obtained direct from benzaldehyde by means of ethyl acetate and sodium (Farb vorm Meister, Lucius und Bruning, D R P 53671, Eng Pat 4969, J Soc Chem Ind 1891, 358; Claisen, Ber 1890, 976) It is a colourless liquid, boiling at 226° (corr) under 250 mm (Perkin, Chem Soc Trans. 1896, 1228) By heating 2 parts of ethyl cinnamate with 1 part of sulphur at 150°-160° for 8 hours, a yellow odourless compound C<sub>9</sub>H<sub>6</sub>S<sub>2</sub>O is formed, and is suitable for therapeutical purposes (Farb vorm F Bayer and Co Eng Pat 8425, D R P 87931; J Soc Chem Ind 1896, 293; Kuster, Ber 1897, 115)

Benzyl connamate v Benzyl benzoate Isomeric acids. Allocunnamic acid, m p 68°, dimorphous \*\*socinnamic acid, m p 42°/58°, have been described, but a discussion as to their constitution cannot be entered into here (Bulmann (Ber 1909, 182 and 1443) and Liebermann (\*\*shift 1027) Cf Stobbe (Ber 1911, 44, 2739), who has studied the transformations of allocand \*\*socinnamic acids in the fused and crystalline states

Substituted cinnamic acids. Of the substitution derivatives of cinnamic acid, the only one of industrial importance is orthontrocinnamic acid, which is prepared in the manufacture of orthontrophenylpropiole acid (v Indigo, Artificial). On a large scale cinnamic acid is first converted by heating with alcohol and concentrated sulphuric acid into ethyl cinnamate. The latter substance, which, being a liquid, is more easily manipulated in the nitration process than the solid cinnamic acid, is run in a thin stream into the calculated quantity of cold

nitrosulphuric acid. The mixture of ortho- and para-nitrocinnamic esters thus formed is dissolved in warm alcohol. On cooling, the paracompound crystallises out, whilst the ortho compound remains in solution. The two ethyl salts are hydrolysed by heating with sulphuric acid. The ortho- acid has also been prepared from o-nitro-benzylidene acetone by warming it with a 3 p c solution of sodium hypochlorite (Hochster, Farb D R P 21162, Frdl 1 29) In this process the two isomerides are formed in approximately equal quantity.

Orthonitrocinnamic acid

## CaHa(NO))CH CH CO2H

is insoluble in water, sparingly soluble in cold, more readily soluble in boiling alcohol, from which it is deposited in crystals melting at 240°

The para-compound melts at 285°-286°, and is very sparingly soluble even in boiling alcohol It cannot be used in the preparation of artificial indigo, and the attempt made to utilise it in the rosaniline manufacture has not proved industrially successful

CINNAMON. The inner bank of shoots of the Cinnamomum zeylanicum (Nees), nat ord Lauraceæ, or true laurels According to Percival, it is imported principally from Ceylon, but in part also from Madras, Tellicherry, and, rarely, from Java The bark-peelers select such branches as are 3 years old, and not more than 2 or 3 inches in diameter nor less than half an meh The back in drying rolls up into quills, many layers being enclosed in one quill It is used as a condiment, and yields a highly perfumed essential oil (q v), employed in the preparation of chocolate and for purposes of perfumery Both the back itself in its powdered state and the oil derived from it are frequently adulterated by cassia (r. Cassia).

The value of cinnamon depends upon its content of cinnamaldehyde rather than on the essential oil A method for the estimation of cinnamaldehyde depends on the colouration developed by treatment with sulphuric acid and isobutyl alcohol (v Fellenberg, Analyst,

1916, 274) CINNAMON BROWN v. Azo- COLOURING

MATTERS CINNAMON OIL v OILS. ESSENTIAL.

CINNAMON-STONE or Hessonite. A gemvariety of gainet of a warm reddish-brown colour (hence the name) and transparent It is essentially an orthosilicate of calcium and aluminium Ca3Al2Si3O12, containing, however, isomorphous replacements of iron, manganese, and magnesium Bulliant crystals are found in veins in serpentine at Ala in Piedmont, but material of the best gem-quality is found as pebbles in the gem-gravels of Ceylon

L J. S

CINNAMYL COCAINE v COCAINE AND THE COCA ALKALOIDS

CITARIN. Trade name for sodium anhydromethylene citrate

CITRAL v Terpenes CITRAMALIC ACID v Malic acid.

CITRIC ACID C6H8O, Occurrence -This acid commonly occurs in the juice of many fruits, and in the sap of many plants
Synthesis —The acid has been prepared

artificially by Grimaux and Adam (Compt rend.

90, 1252). A saturated solution of dichloracetonic acid was neutralised with sodium carbonate, and heated with two molecules of potassium cyanide. The resulting solution of dicyanoacetates was saturated with hydrochloric acid gas, and heated on a water-bath for 15 hours, the citric acid was then separated as calcium citrate by neutralisation with milk of lime Another synthesis has been effected by Lawrence (Chem Soc. Proc 1897, 65), who obtained ethyl citrate by heating together ethyl bromacetate and ethyl oxalylacetate in the presence of zinc

Properties -The crystallised acid of commerce has the formula C6H8O7, H2O. The crystals are large prisms belonging to the trimetric system. Their sp gr, according to Buignet, is system. Their sp gr, according ar, and in 1553 They deliquesce in damp air, and in perfectly dry air slowly lose the whole of their Different crystals (probably differently prepared) may lose water at very different rates (Warington, Chem Soc Trans. 28, 928, Gros-According to Marchand (J. jean, ibid, 43, 331) pr Chem 1841, 23, 60), crystals obtained from a saturated boiling solution have the formula 2(C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>),H<sub>2</sub>O According to a later investigation by Sarandinaki (Ber 5, 110), the crystals from a solution long boiled are anhydrous Citric acid is optically mactive. The ordinary crystallised acid dissolves in about half its weight of boiling water Aqueous solutions of various strengths have, according to Gerlach (Zeitsch anal Chem 1869, 295), the following sp gr at

Acid p c	Sp gr	Acid p c	Sp gr
1	1 0037	34	1 1422
2	1 0074	35	1 1467
3	1 0111	36	1 1515
4	1 0149	37	1 1564
5	1 0186	38	1 1612
6 7 8 9	1 0227 1 0268 1 0309 1 0350 1 0392	39 40 41 42 43	1 1661 1 1709 1 1756 1 1814 1 1851
11	1 0431	44	1 1899
12	1 0470	45	1 1947
13	1 0509	46	1 1998
14	1 0549	47	1 2050
15	1 0588	48	1 2103
16 17 18 19 20	1 0632 1 0675 1 0718 1 0762 1 0805	50 51 52 53	1 2153 1 2204 1 2257 1 2307 1 2359
21	1 0848	54	1·2410
22	1 0889	55	1 2462
23	1 0930	56	1 2514
24	1 0972	57	1 2572
25	1 1014	58	1 2627
26	1:1060	59	1·2683
27	1 1106	60	1 2738
28	1 1152	61	1 2794
29	1 1198	62	1 2849
30	1 1244	63	1 2904
31	1 1288	64	1 2960
32	1 1333	65	1 3015
33	1·1378	66	1 3071

A 25 p.c. solution boils at 101 8°, a 50 pc. solution at 105 8° (Gerlach, J 1859, 48)

At 15° the solubility of the crystallised acid in alcohol is as follows.

> 100 of 80 p.c. alcohol dissolve 87 90 52 85 100 75 90

100 parts of anhydrous ether dissolve 9 l of the crystallised acid

According to Buchner (Ber. 1892, 1159), the ordinary hydrated acid melts between 135° and 152°, and the anhydrous acid at 153° Saltzer (Arch Pharm. 231, 514) finds that the finely powdered hydrated acid, if slowly heated, becomes anhydrous at 55°, and melts at 160° in this condition.

Decompositions —When heated to 175°, acetone and carbonic oxide are evolved, the residue is a conitic acid  $C_6H_6O_6$  , the same acid is formed in small quantity when a solution of citic acid is long boiled (Dessaignes) When heat is continued beyond 175°, an only distillate appears, which yields crystals of itaconic acid C5H6O4. By further heating, itaconic acid is transformed into an uncrystallisable oil, citraconic anhydride C<sub>5</sub>H<sub>4</sub>O, Heated with sulphure acid, carbonic oxide and acetone are given off, the intermediate stage in the reaction being the production of acetone dicarboxylic acid (v. Pechmann, Ber 1884, 17, 2542) This acctone dicarboxylic acid is also formed in the oxidation of citric acid by potassium permanganate and by the photochemical decomposition of citric acid in the presence of 1 p c uranium oxide

Solutions of citue acid are rapidly destroyed by fungi According to F Watts (J Soc Chem Ind 1886, 215), the action of Saccharomyces mycoderma on lime juice is to convert the citric acid into carbonic acid and water cording to I Macagno (Gazz chim ital 11, 443), an alcoholic fermentation in lemon juice does not affect the citiic acid, but under the influence of bacteria the citric acid disappears, and acetic

and propionic acids are formed

Salts — Citric acid is tribasic, and forms in consequence three classes of salts The salts with the alkali metals are readily soluble in water, the neutral salts of the alkaline earths are only sparingly soluble

Calcium citrate ('a<sub>3</sub>(C<sub>6</sub>H<sub>6</sub>O<sub>7</sub>)<sub>2</sub>,4H<sub>2</sub>O does not lose water at 100° When precipitated in a erystalline state its solubility in water is 1 in 1180 at 14°, and 1 in 1730 at 90°-100° When thrown down in an amorphous condition, the solubility is greater, 1 in 707 at 18°, and 1 in 1123 at 100° (Chem Soc Trans 28, 939)

When a solution of citric acid contains a ferrous, ferric, or aluminium salt, in not too great proportion, it may be made strongly alkaline with potash or soda without producing a picci-Such a solution may also be boiled with chalk without precipitating iron or aluminium, although a large part of the citric acid falls as calcium citrate. (With tartaric acid in the place of citric, iron is to a small extent precipitated by chalk in the cold, and entirely on boiling) If, however, the ferric salt bears a large proportion to the citric acid, the whole is precipitated by chalk, even in the cold (Chem. Soc Trans. 28,

Derivatives .- Methylenecitric acid is obtained

by heating citric acid with four times its weight of polymeric formaldehyde at 140°-160°, and allowing the melt to crystallise It is tolerably stable towards acids, but is decomposed by alkalis with elimination of formaldehyde Its di-sodium It is used as a salt is known as 'citarin.' disinfectant for the urinary passages (Ch. Fab auf Act vorm E Schering, US Pat 699422; J Soc Chem Ind 1902, 791).

Monophenetidenecitric acid is obtained by heating 42 parts of citric acid with 27 5 parts of p-aminophenetol for some hours at 100°. The product is dissolved in caustic soda, and the precipitated with hydrochloric acid Diphenetidenecitric acid is obtained similarly. They are medicinal substances, acting energetically as precipitants without producing secondary symptoms (Farb von Heyden-Nachfolger, Eng Pat 1254, J Soc Chem Ind 1895, 385)

Detection - ('itrie acid is best recognised when in a pure state by its yielding on heating an oily sublimate, which afterwards crystallises (itaconic acid) A solution of citiic acid, neutralised with ammonia, gives no precipitate in the cold with calcium chloride, except after long standing, but on boiling, calcium citrate is pre-cipitated. Calcium citrate, precipitated in the cold, is freely soluble in solution of ammonium chloride, and is precipitated therefrom on boiling Alkalı citiates do not reduce silver salts on boiling, as tartrates do, or only to a trifling extent

Kammerer (Zeitsch anal Chem. 8, 298) recommends the detection of citric acid by the To the formation of a crystalline barium salt free acid, or to the alkali salt, excess of barium acctate is added, and the whole heated on a water bath for several hours, the banum citrate

will then be found in microscopic monoclinic prisms.

Dèniges (Compt. rend 130, 32) adds to the solution to of its volume of a reagent obtained by dissolving 5 grams of mercuic oxide in 20 cc of cone, sulphuric acid, and 100 cc of water. The solution is heated to boiling, and, if citric acid is present, a white precipitate is obtained on the addition of a few drops of a 2 p c solution of potassium permanganate

Another method is to add potassium permanganate to the solution until it is pink. If a precipitate is formed, or a brown colour produced, ammonium ovalate is added until it is destroyed On adding bromine water a white precipitate is obtained, which, on the addition of alkalı, gives the characteristic smell of bromoform (Stahre, J. Soc Chem Ind 1896, 53, Wohlk, Zeitsch anal Chem 1902, 77).

The tests for adulteration will be given at the close of this article

Estimation -When the quantity of citric acid in a solution is determined by alkalimetry, phenolphthalem should be used as the indicator; in coloured solutions red-brown turmeric paper (Thomson, J Soc. Chem Ind 1887, 195) has been recommended. If neutral litmus paper is used, the alkalı must be standardised with citric The quantitative determination of citric acid by precipitation as calcium citrate will be described under the analysis of lemon juice.

Citric acid in milk and wine. Citric acid is

present in milk and wine, and as it occurs in small quantities in most wines, adulteration with a small quantity of citric acid cannot be detected (Hubert, Ann. Chim. anal. 1908, 139). It may be detected in milk by taking 10 c c. of milk, 2 c c ot soulum mecaphosphilis, mercuric sulphate solution, and filtering Half of the filtrate is mixed with 0 5 c.c of 10 p c.

To milk, 2 c c of sodium metaphosphate, 3 c.c. of this 4 drops of a 2 p c solution of potassium permanganate is added, and the solution again This is repeated, and the liquid decolourised by a drop of hydrogen peroxide precipitate or an opalescence indicates the presence of citric acid. This method may be used quantitatively (Denigès, Chem Soc Abstr. 1902, in 365, v also Scheibe, ibid 1891, 1276) It may be detected in wine by the method of Spica (Gazz chim ital 31, [2] 61) A portion of the wine is evaporated to a syrup and extracted with alcohol. This is nearly neutralised with alcoholic potash, and the potassium bitaitrate filtered off. The filtrate is then neutralised with more potash, and the precipitate is filtered off, dried, and heated in a small tube with sulphuric acid, until it goes It is then diluted, made alkaline with caustic potash, and sodium introprusside added A red colour indicates the presence of citric Favrel (J. Soc Chem. Ind 1908, 637) evaporates 100 c c. of wine to dryness, dissolves the residue in 7 cc of boiling water, and adds 35 e.e. of 90 p.e. alcohol A solution of calcium acetate (0 4 gram in 5 cc of water) is added, and the precipitate filtered When dry it is transferred to a test-tube, 3 cc of sulphuric acid added, and the whole heated to 80°-90° for 2 minutes The solution is cooled and extracted with ether White crystals are obtained, giving a violet-red colour with ferme chloride, the colour being discharged by mineral acids other methods and quantitative estimation, vWilliams, Analyst, 14, 25, Devarda, J. Soc. Chem. Ind. 1904, 273, Klinger and Bujard, Chem. Soc. Abst. 1893, 11–54; Schindler, Chem. Zentr 1902, II 1016; and Robin, Ann Chim anal 1904, 453)

Gowing-Scopes finds that a mixture of equal weights of mercuric nitrate, manganese nitrate and nitric acid reacts with citric acid, yielding a precipitate after prolonged boiling which is exactly six times the weight of citric acid present. The method gives good results in presence of tartane, succinic and other organic acids, but with malic and lactic acids the results are high (Analyst, 1913, 12)

3 , , ,

### RAW MATERIAL.

Citric acid is prepared from the fruit jurce of three species of Citrus—the lemon, bergamot, and lime, the first of these is the principal source of citric acid. Concentrated lemon jurce is chiefly imported from Sicily, a very little comes from Naples or Sorrento. Concentrated bergamot jurce is prepared in Calabria, and exported from Messina. Concentrated lime jurce is imported from Montserrat and Dominica. The lemon jurce from Sicily is prepared by pressing the inferior fruit, from which the rind has previously been removed for the manufacture of essence. The so-called 'single' jurce.

is then boiled down till its sp.gr. equals 60° on the citrometer (sp.gr. 1.24); it then forms a dark-brown, rather syrupy liquid.

Lemon juce has its greatest acidity early in the season (November). The juce of the fine fruit, exported as lemons, has a sp.gr. 103-104; it contains free acid equal to 11-13 oz. of citric acid per gallon. The inferior fruit pressed in Sicily yields a juice containing at the beginning of the season about 9 oz. of free acid per gallon, and at the end of the season a good deal less.

There are other differences between the juice of fine lemons pressed in England, and the Sicilian juice employed for the manufacture of eitric acid. The former contains very little combined organic acid, only about 2.5 p.c. of the total organic acid (free plus combined) present. In unconcentrated Sicilian juice the combined acid is about 7–9 p.c. of the total Again, the English pressed juice contains hardly any organic acid save citric, only about 1 p.c. of the total organic acid being unprecipitable as calcium citrate. In Sicilian juice about 8 p.c. of the total organic acid is unprecipitable as calcium salt.

The concentrated lemon juce from Siely is reckoned of standard quality when its sp gr is 124, and its acidity is equal to 64 oz per gallon of nominal citric acid. In dealing with trade analyses it must, however, be borne in mind that the 'crystallised citric acid.' of a trade certificate is not the crystallised acid of commerce, but an acid containing only half the actual amount of water, an acid, in fact, having the atomic weight 201 instead of 210:

## CaHaOz,HaO

No reason can be given for this practice, which should certainly be abolished. In the present article all quantities of citric acid will be expressed in terms of the common crystallised acid.

The concentrated lemon juice from Sicily contains pretty uniformly 7–8 oz per gallon of combined organic acid, equal to about 10 pc. of the total organic acid (reckoned as citric) present. Of the total organic acid about 10 pc. is not precipitable as calcium salt, and is therefore not citric acid; this proportion is, however, by no means constant. Purchases of lemon juice are unfortunately still made on the basis of acidity, and not on the amount of precipitable acid present. Analyses of 895 pipes of concentrated lemon juice by Grosjean have been published (Chem Soc Trans 43, 333). The average proportion of precipitable acid was 99 2 pc of the free acid, but the range of variation was considerable, individual parcels of juice giving 81 1, 85 %, and 103 6 pc.

giving 81 1, 85.8, and 103 6 p c.

Concentrated Bergam of juice has a similar spigr. to lemon juice, but a lower acidity; it generally contains free acid equal to about 51 oz of citric acid per gallon. The quantity of combined organic acid is apparently similar to that in lemon juice, namely, 7-8 oz. per gallon; but the proportion of combined to total is higher, 12-13 p.c.. The proportion of unprecipitable acid is about 13 p.c. of the total. Grosjean's analyses of 90 pipes of bergamot juice show a mean of 98.4 of precipitable acid for 100 of acidity, the extremes being 95.4 and 101.4.

The unconcentrated Lime juice of Montserrat has a mean sp.gr. of 1 036; it contains, according to Conroy (Pharm. J. 1883, 606), an average of 7 84 pc, or 12.54 oz per gallon of free acid. The extremes observed wore 6 70–10 05 pc., equal to 10 7–16 l oz. per gallon. The juice yields 0 43 pc of ash. It contains only a trace of sugar. Warington found the combined acid in two samples 5 pc of the total, 10 pc of the total acid was not precipitable

Concentrated lime juice is a viscid liquid, sp.gr 1 32, and with an acidity averaging about 94 oz of eitric acid per gallon. The combined acid is about 8-9 oz per gallon. The unprecipitable acid is about 10-14 p c of the total. The precipitable acid bears a lower proportion to the free than is the case with either lemon or bergamot juice, the published analyses showing a mean of 93 8 of precipitable acid for 100 of free.

The nature of the organic acids, other than citric, present in lemon, bergamot, and lime juce has not been determined. A little formic acid and acetic acid have been detected in concentrated juice, but the principal acids other than citric are clearly non-volatile and have soluble calcium salts. The acids most probably present are malic and acountic

Besides lemon juice, some clude Calcium citrate, prepared in Sicily by precipitating lemon juice with chalk, is exported into England.

It contains about 64 p c of citric acid

## METHODS OF ANALYSING JUICE AND CITRATE

1 Aculity —The commercial analysis juice is confined to the determination of acidity It is most important that the alkalı (sodium hydroxide) used should be actually standardised with citric acid, and the same indicator employed in standardising as in subsequent analyses. Powdered crystals of citric acid are taken, water carefully determined in one portion, while another is used for standardising the alkali determining water, the powdered acid should be heated for some hours at about 60° before exposing it to 100°, if this is not done, the acid will melt, and lose the rest of its water with great difficulty Warington recommended delicate litmus paper for determining the neutral point F Watts has since employed films of turmeric tincture on a white tile (J Soc Chem Ind. 1886, 214) Thomson recommends red-

brown turmeric paper
2. Combined organic acids—The juice is neutralised with a known quantity of standard alkalı, evaporated to dryness in a platinum basin, the residue gently ignited, the black ash treated with a known quantity (excess) of standard sulphuric acid, the whole boiled and filtered The amount of unneutralised acid is then determined with alkali We have now the neutralising power of the ash in terms of alkali; by subtracting the alkali added when neutralising the juice, we obtain the amount of alkali corresponding to the bases of the organic salts, which became carbonates on ignition the amount of these bases their equivalent in combined citric acid can be calculated.

3 Precipitable citric acid—This is for the manufacturer the only trustworthy method of analysis, but is, in fact, seldom adopæd. War-

ington proceeds as follows: 15-20 e.c. of unconcentrated juice, or about 3 cc. of concentrated juice, are exactly neutralised with sodium hydroxide, the solution (about 50 c c ) is brought to boiling in a salt-bath, and a measured quantity of calcium chloride solution, known to be rather more than sufficient for all the organic acids present added. After boiling for half an hour, the precipitate is collected on a small filter, and washed with hot water. The filtrate and washings are then concentrated to 10 cc and the solution finally neutralised with a drop of am-The second precipitate is collected on a monia very small filter, the filtrate being used for getting the precipitate on to the paper, which is finally washed about five times with a little hot water. The precipitates in their papers are then burnt at a low heat in a platinum basin, and the neutralising power of the ash determined with standard hydrochloric acid and alkali The amount of citric acid which the base is equivalent to can then be calculated, three molecules of base being reckoned as equivalent to one molecule of citric acid

4 Analysis of calcium citrate—The excess of chalk present is first determined by boiling about 4 grams with dilute standard hydrochloric acid in a covered beaker, and then adding standard alkalı tili feebly alkalıne Two grams of the citrate are then gently ignited in a covered platinum crucible, and the neutralising power of the ash determined by solution in standard hydrochloric acid and titiation with alkali By deducting from the neutralising power of the ash that due to the chalk previously determined, we find the amount of base which has to be calculated as citrate. This method presupposes that citrates are the only organic salts present If it is preferred, the citrate may be dissolved in hydrochloric acid, the solution boiled, neutralised with soda, the calcium citrate precipitated by boiling in the salt-bath, and its quantity determined as in the case of juice

The information here given respecting juice, calcium citrate, and the methods of their analysis, will be found in greater detail in Chem Soc Trans 28, 925, 43, 331, v also Ulpiani and Pairozzani (Chem Soc Abstr 1907, n 57), Shica (Analyst, 1910, 35, 519), Barboni (ibid. 1913, 65).

### PROCESS OF MANUFACTURE.

The manufacture of citic acid from concentrated lemon juice is extremely simple A proper quantity of whiting (levigated chalk) is mixed with water, and heated by steam in a wooden vat provided with a revolving agitator, the concentrated juice is then slowly pumped in, care being, of course, taken that the whiting is finally in small excess. The liquor never becomes neutral, however long boiling may be continued, or however great is the excess of whiting present, the adjustment of juice and whiting is therefore effected by ascertaining if the liquor effervesces with more whiting, or the precipitate effervesces with more juice. The amount of unneutralised acid is about 1-21 p.c of the original acidity of the juice. Pure citric acid is readily neutralised by whiting, malic and aconitic acids are not; the final acidity is thus possibly due to the presence of these acids. Citric acid, however, is not

neutralised by chalk if phosphates, and especially ferric phosphates, be present; this fact will also explain the result—It is not advisable to neutralise completely by the use of lime, as vegetable impurities are then thrown down which are afterwards difficult to separate

The precipitated calcium citrate is washed with hot water on a liter. It is next brought by the addition of water to the state of thin cream, and decomposed, with constant agitation, by the addition of a small excess of sulphunic acid (sp.gr. 1.7). The occurrence of an excess of sulphunic acid is known by the liquor affording a precipitate with a strong solution of calcium

chloride after some minutes' standing

The citric acid liquor is then separated from the gypsum, which is washed on a filter liquor is evaporated in shallow leaden baths by steam heat Much gypsum is at first deposited, from this the clear houor is run off and further concentrated When strong enough to crystallise, the hot liquor is run into a wooden tub provided with an agitator, and the liquor is kept in constant motion while cooling, by this process, known as 'granulation,' the citric acid is obtained as a crystalline powder. The mother liquor is again concentrated, and 'salt' again obtained by granulation. The process may be repeated a third time. The liquor is then too dark and impute for further crystallisation, and is known as 'old liquoi' The granulated citic acid when drained, and, if necessary, slightly washed, is redissolved, decolourised by heating with animal charcoal (previously freed from phosphates by hydrochloric acid), again concentrated to the crystallising point, and poured into leaden trays about 3 inches deep, the crystals here formed are the citric acid of commerce Citric acid thus prepared always contains a trace of lead, and often particles of metallic lead, probably due to the plumbers repairing the vessels. To free the and from lead, the process usually adopted is to pass sulphuretted hydrogen through the solution after the first crystallisation, filter, and then recrystallise the citie acid in lead-free vessels

The 'old liquor' is diluted with water, and the citic acid it contains precipitated with an excess of whiting, exactly as in the case of the examination. If iron or copper is present in original juice. The liquor is never neutralised any quantity, a drop of solution of potassium by the whiting, this may be either due to examination. If iron or copper is present in original juice is never neutralised any quantity, a drop of solution of potassium by the whiting, this may be either due to examination. If iron or copper is present in original juice is added before the ammonium sulphide acontic acid liquors, or to the presence of ferric reaction. The standard of purity suggested is or aluminic phosphate derived from the whiting is that lead is not to be present to a greater extent.

Any considerable excess of sulphuric acid in the liquors, or any overheating, must be avoided, as occasioning decomposition of citric acid The presence of non or aluminium in the whiting also occasions loss, as citie acid holding iron or aluminium in solution is not precipitated by calcium carbonate In consequence of the nonprecipitation of iron or aluminium from citrie solutions by whiting, the citric liquors of the factory remain nearly pure, however long the work may have been continued, a result very different from what happens in the case of tartaric acid; the purity of citric-acid liquors is however obtained at the expense of some loss of citric acid In a well-conducted factory the total loss during manufacture will amount to 12-15 p c of the citric acid in the juice.

A method has been patented (Fancheux and terated with tartaric acid. To detect adulteration,

Boissière, Eng. Pat. 11991; Fr. Pat. 338735, J Soc Chem. Ind. 1904, 680) by which the ciude calcium citrate is purified by passing hot gases over it, so as to roast it superficially

Citric acid is also produced by the fermentation of carbohydrates, glycerol, and analogous substances under the influence of certain fungi, named *ettromycetes* (Wehmer, Eng. Pat. 5620, D. R. P. 72957, J. Soc. Chem. Ind. 1894, 275). To prepare the organisms, a sugar solution containing 2 5 p.c. of citie acid is allowed to stand for a few days The growth formed on this solution is introduced into sterilised sugar where it develops. The pure culture prepared in this way is then introduced into large vessels containing sugar solution, in which are dissolved small quantities of ammonium nitrate, dipotassium phosphate, and magnesium sulphate, the liquid is then allowed to remain at the ordinary temperature for 8-14 days. At the expiration of that time, the liquid contains a certain amount of cities acid, so that it can be directly employed for lemon juice. The acid can be isolated, preferably as calcium citrate by adding calcium carbonate to the liquid Other moulds, such as Mucor proforms, which grows on putretying fruit, specially on pears and apples, may be employed (D R P 91891, v also Maze and Perner, Compt rend 139, 311).

Estimation of lead in eitric acid Forty grams of the sample are dissolved in water, pure strong ammonia added in slight excess, the hquid cooled and diluted to 120 cc. For a preliminary experiment 10 cc are taken, diluted to 50 c c. in a Nessler tube, and one drop of ammonium sulphide added, the colour determines what volume should be taken, it varies from 5 to 50 cc. Having repeated this with the volume of liquid thus indicated, the tint has to be matched with that from the pure solution. A volume of pure ammonium citrate solution identical with that taken of the acid under examination is mixed with a measured quantity of slightly acid lead nitrate solution, containing 01 gram metallic lead per litre A drop of ammonium sulphide is added, and various solutions are made up until the colour obtained matches that from the acid under examination If iron or copper is present in any quantity, a drop of solution of potassium which prevents them interfering with the reaction The standard of purity suggested is that lead is not to be present to a greater extent than 5 parts per million (Warington, J Soc Chem Ind 1893, 97, v also Tatlock and Thomson, Analyst, 1908, 173)

Another method is that of Buchet (J Soc. Chem Ind 1892, 848) To estimate the metallic lead, 200 grams of acid are dissolved in three times its weight of water, and a slight excess of ammonia added After 24 hours the liquid is docanted, and the residue collected on a filter, carefully washed and dissolved in nitric acid. The lead is then precipitated as sulphate, and estimated as such. To determine the combined lead, the ammoniacal solution is first acidified with hydrochloric acid, the lead precipitated as sulphide, converted into sulphate, and weighed as such.

Adulteration.—Citric acid is at times adulterated with tertaric acid. To detect adulteration

CLAY 287

Cailletet (Chem. Soc. Trans 36, 674) adds | I gram of the powdered acid to 10 c c. of a cold saturated solution of potassium dichromate the solution remains unchanged in colour for 10 minutes, tartaric acid is absent If tartaric is present, the solution becomes brown Pusch (J. Soc. Chem. Ind 1885, 552) places 1 gram of the acid in a test-tube with 10 grams of pure oil of vitriol. The test-tube is placed in a water-bath, and kept at 100° for I hour citric acid only is present, the colour is lemon-If tartaric acid is present, the colour brown Both these tests are very becomes brown delicate, and readily show 1 or 1 p c of adulteration Piñerua (Compt. rend 124, 291) puts 0 05 gram of the substance in a porcelain basin, and adds 10-15 drops of a reagent made by dissolving 0 02 gram of \beta-naphthol in 1 cc. of concentrated sulphure and 1f eithe acid alone is present, a blue colour is formed, unchanged on heating, which on dilution gives a nearly colourless or a slightly yellow liquid tartaric acid is present, the colour on heating changes to green, and on dilution to a reddishvellow Deniges adds a small quantity of the acid to a solution of resorcinol in dilute sulphuric acid, and heats to 115°-140°. If tartain acid is present, a violet-red colour is formed sugar is present, or other bodies blackened by sulphuric acid, the acid is precipitated as its lead salt, and then tested (J. Pharm Chim 1895, 1 586) Crismei (Chem Soc Abstr 1892, 546) adds 1 gram of the acid to 1 cc of ammonium molybdate solution (10 pc) and 2 to 3 drops of hydrogen peroxide solution (1 to 1 pc), and heats at 100° for 3 minutes. If citric acid alone is present, the yellow colour given by the molybdate is unaltered, whilst if tartain acid is present, a beautiful blue colour results test detects 1 mgm of tartaric acid in 1 gram of citric acid Spindler has modified Deniges' mercuric sulphate method (q v), and uses potassium dichiomate instead of permanganate If extric acid alone is present, a yellow precipitate is obtained; the presence of tartaile acid is indicated by a change of colour to green, due to reduction. Taitaile acid may also be detected by adding a little solution of potassium acetate to a strong solution of the citric acid, and stirring

The quantity of taitaric acid in an adulterated sample may be determined by adding 10 p c of potassium chloride to a strong solution of the acid, and precipitating the tartaric acid with citrate or acctate of potassium, with the precautions described under estimation of tartame acid

CITRIC ACID FERMENTATION v. FERMEN-TATION

CITROCOLL. Trade name for neutral aminophenacetin citrate

CITRON, OIL OF, v OILS, ESSENTIAL CITRONELLA OIL v OILS, ESSENTIAL

CITRONINES v Azo- colouring matters. CIVET. An odorrferous substance resembling musk or ambergris, obtained from the pouches situated between the anus and genital organs of Viverra civetta of N Africa, V Zibetha of Asia, and V. Malaccensis of Java. Is of the consistence of honey, and of a yellowish or brown

Pharm. 1824, 537) (For analyses of civet, see Charabol and Hébert, Bull Soc. chim 1910. 7, 687.) Civet melts at about 36°, and contains from 50 to 70 p c of fatty acids, and should have a saponification value of about 180°, and should have not more than 1 p c of ash.

It has been inferred that the musk aroma of ervet is due to a ketone, muskone, present in musk, but according to Sack (Chem Zeit 1915, 39, 538), it is caused by a specific ketone, Zibetone C<sub>17</sub>H<sub>30</sub>O, of a pleasant musk-like odour, which may be obtained pure by decomposition of the crystallised semicarbazone. Its oxime crystallises in white needles, in p 92°. See Musk

CIVETONE of ZIBETONE v MUSK. CLAUSTHALITE v SELENIUM. CLAVICEPSIN v ERGOT. CLAVIN v ERGOT

CLAY. (Argde, Fr; Thon, Ger) The term 'clay' is ordinarily used to denote certain earths, the chief characteristic of which is that of plasticity when wet, and which have the property of baking to a haid, stone-like mass when heated to full redness Clay is a heterogeneous mixture of certain hydrated aluminium silicates, with particles of quartz, felspar, mica, &c , resulting from the decomposition of various silicious minerals, the chief source being the felspars of granite rocks These hydrated aluminium silicates form the essential basis of all clays, and are known as clay substance or clay base. They are all decomposed by hot concentrated sulphuric acid (infra), whilst the other ingredients mentioned above are affected only slightly or not at all. The method of so-called 'rational analysis' is based on this action, and it has been found that when so treated, the great majority of clays give a clay substance approximately of the composition represented by the formula Al<sub>2</sub>O<sub>3</sub> 2SiO<sub>2</sub>,2H<sub>2</sub>O This corresponds with the composition of the rare crystalline mineral kaolinite, and, consequently, it has often been assumed that all clays contain kaolinite as their clay base It has, however, been shown that the clay substance, in some cases, does not correspond in composition with the above formula, and it is now generally accepted that other hydrated aluminium silicates may be present in clays, forming part, and in some cases the whole, of the clay base Various investigators have attempted to define these substances, and to show that they correspond in composition with certain silicates, such as allophane  $(Al_2O_3 \cdot SiO_2, 5H_2O)$ , pyrophyllite  $(Al_2O_1 \cdot SiO_2, 5H_2O)$ , halloysite  $(Al_2O_3 \cdot 2SiO_2, 2H_2O + Aq)$ , pholerite  $(2Al_2O_3 \cdot 3SiO_2, 4H_2O)$ , &c

It has been suggested by J. W. Mellor (Trans, Eng. Ceramic Soc. pt. 1, 8) that the term kaohnite should be restricted to the comparatively rare crystalline mineral, and that the amorphous clay substance of similar composition (Al<sub>2</sub>O<sub>3</sub>·2SiO<sub>2</sub>,2H<sub>2</sub>O) should be called clayite.

The same investigator points out that this latter substance occurs in clays in non-crystalline or amorphous granules, which, however, sometimes show faint polarisation phenomena. Practically all clays contain more or less free colour. Contains stearm, olem, mucus, resm, silica, undecomposed or partially decomposed volatile oil, and yellow colouring matter (J. felspar, and mica, and they vary in composition

through a wide range, differences in composition being brought about partly as a result of different origins, and partly by differences of deposition, &c. The commoner varieties of clay generally contain a considerable percentage of oxide or carbonate of iron, and small percentages of calcium, magnesium, and titanium salts.

According to Washington (J. Amer Ceram. Soc. 1918, 1, 405) clays are composed of four minerals corresponding to generalised formulas, viz kaolinite (Al<sub>2</sub>O<sub>3</sub>,2SiO<sub>2</sub>,2H<sub>2</sub>O), orthoclase (K<sub>2</sub>O,Al<sub>2</sub>O<sub>3</sub>,6SiO<sub>2</sub>), albite (Na<sub>2</sub>O,Al<sub>2</sub>O<sub>3</sub>,6SiO<sub>2</sub>), anorthite (CaO,Al<sub>2</sub>O<sub>3</sub>,2SiO<sub>2</sub>), limonite (Fe<sub>2</sub>O<sub>3</sub>, H<sub>2</sub>O) with, in some cases, muscovite (K<sub>2</sub>O,2H<sub>2</sub>O, 3Al<sub>2</sub>O<sub>3</sub>,6SiO<sub>2</sub>). The proportions of the several minerals may be calculated from the ultimate chemical analysis. The lime, soda, and potash are first calculated to the felspars, and the alumina and silica required for those are deducted from the total. The iron oxide is calculated as limonite. Any surplus alumina is calculated as kaolinite, and any surplus silica is regarded as quartz (see below)

Clays may be divided broadly into two classes: residual clays and sedimentary clays Residual clays are the insoluble residue left from the breaking down and decomposition of 10ck masses containing certain aluminium silicates (chiefly felspathic minerals), and they are found overlying or adjacent to the rock mass from which they are formed. These residual clays usually present a gradual change in mineral character from the finer or puter portion nearer the surface to the mixture of clay and undecomposed rock which lies in contact with the rock The decomposition of the mineral silicates may be brought about by carbonic acid in the surface waters; but many of them are acted on by water alone, and although this action proceeds very slowly, it may account for the gradual decomposition of the felspathic rocks and the formation of clay. The alkalis of the felspars combine with the free silica formed in the decomposition, and with the carbon dioxide contained in the percolating water, forming soluble salts which are more or less completely washed away. In some cases the decomposition of the mineral silicates may have been brought about by the action of gases, and it has been shown by J. H. Collins (Min Mag 1887, 7) that when felspar is exposed to the action of hydrofluoric acid, hydrated aluminium silicate is formed, mixed with soluble potassium fluoride, whilst silica is deposited on the sides of the tube in which the reaction takes place Residual clays, as a rule, contain a smaller percentage of oxide of non and organic matter than sedimentary clays, and are consequently lighter in colour both before and after baking

Sedimentary clays are formed by the transportation and redeposition in water of residual clays, and clays so formed may be again transported and redeposited in the same way, so that they may have no direct relationship to the rock formation on which they lie. Sedimentary clays belong to all geological ages, from the present to the Cambrian, and are of varying degrees of hardness, from the soft drift which may be dug with a spade to the hard clay shales of the older formations, which require to be blasted. These clays are always roughly

stratified in structure, and show considerable variations in general character and composition throughout their mass, owing to the varying conditions under which they were deposited. The harder varieties of clay are readily converted into the plastic form by grinding with water, or by weathering, and they are now largely used for the making of bricks, &c, where formerly only the softer surface clays were used. When breathed upon or moistened, clays usually emit a peculiar odour known as argillaceous. This characteristic is more marked in the sedimentary than in the residual clays, and may possibly be connected with the contained organic matter.

Plasticity is an essential property of all clays in the moist condition, but this quality The presence varies greatly in different clays of organic matter, finer grinding of the particles of the clay, 'ageing' in the most condition and 'weathering,' all tend to increase the plasticity, but no entirely satisfactory explanation of the cause of plasticity has yet been put forward. Clay loses practically the whole of its hygroscopic water when dried at 100°, and it contracts and loses its plasticity at the same time, becoming hard and brittle On being soaked or ground in water, however, it becomes soft and plastic again The chemically combined water is not driven off until the clay is heated to redness, the temperature of the dehydration, in the case of ordinary clay, being probably between 480° and 600°. Clay that has been heated to this temperature will not again become plastic on being ground with water, so that it is evident that the plasticity of a clay is dependent upon some physical or chemical condition, which is destroyed on heating to about 500° When clays are heated to a higher temperature (generally about 900°), they contract more or less and become hard, compact, and stone-like, and at higher temperatures still they become vitieous and ultimately fuse. The purest varieties of china clay (which approximate to the composition of kaolinite) fuse at a temperature of about 1830°, and other clays fuse at lower temperatures, according to the nature and percentage of the various impurities contained. The chemical impurities which chiefly affect the fusibility of a clay are the salts of magnesium, calcium, iron, sodium, and potassium first pointed out that chemically equivalent amounts of the oxides of these substances exert equal influences on the fusibility of a clay presence of free silica (quartz sand) tends to raise the fusion point of clays containing a considerable percentage of these bases, but it has an opposite effect on the purer china clays the manufacture of pottery ware of all kinds, the firing temperature depends upon the chemical composition of the clay (which determines the fusibility) and upon the result required. For example, in the manufacture of porcelain, where the resultant ware must be non-absorbent or vitreous, the firing temperature ranges from about 1250° to 1550°, according to the chemical composition of the clay mixture or body used, whereas in the manufacture of ordinary white earthenware, where a porous or non-vitreous ware is desirable, the firing temperature ranges from about 1150° to 1250°.

The colour of fired clay, i.e. clay that has

been heated to a temperature of over 900°, depends upon the impurities present, and upon the character of the firing gases. The purer the character of the firing gases. kinds of residual clays fire to a whitish mass. but all other clays become more or less strongly coloured, the tint being determined generally by the percentage of iron oxide present. Iron may be present in clays in various combinations, such as ferric oxide (Fe<sub>2</sub>O<sub>3</sub>), hydrated ferric oxide (2Fe<sub>2</sub>O<sub>3</sub>, H<sub>2</sub>O), ferrous carbonate (FeCO<sub>3</sub>), and more rarely magnetic oxide (Fe<sub>3</sub>O<sub>4</sub>), but in firing in the ordinary way all these salts are converted into ferric oxide Some clays contain a considerable amount of iron pyrites disseminated through the mass in small grains, and these clays are often troublesome in use, as the sulphide is not so readily converted into oxide in the ordinary firing process. Moreover, the sulphide on weathering with the clay gives rise to soluble sulphates which are detrimental. In the yellow and brown surface clays and red-firing shales, the iron is present in the form of ferrous carbonate, which, together with the organic matter present, produces the natural grey colour of these shales. In ordinary clays, and with an oxidising atmosphere, 1-2 pc of iron oxide produces a buff colour, 2-4 pc a salmon colour, and above 4 pc a red colour, which becomes darker as the percentage of iron increases presence of other impurities, however, modifies the colour produced by iron oxide considerably, and this is especially the case when a large percentage of lime or magnesia is present

Some clays (commonly called marls) contain from 15 to 30 pc of lime, and although they may also contain as much as 6-8 p.c of iron oxide, they fire to a light-yellow or buff colour. Such clays are largely used for making the wellknown yellow facing bricks, which were at first made chiefly from the marks of the Thames basin, the Paris basin, &c, and are now made from artificially prepared mixtures of clay or chalk and limestone The presence of certain soluble salts (chiefly sulphates of calcium or magnesium) in clays often produces serious discolouration on firing, owing to the deposition of these salts on the surface during the drying and firing of the ware To prevent this action, it is usual to mix with the clay barium carbonate, to combine with the whole of the sulphuric acid present, and so convert the soluble salts into When clays are fired in a insoluble ones reducing atmosphere, the red ferric oxide is converted into blue ferrous oxide, and the resultant ware is consequently bluish in colour. Blue bricks are made from certain clays, containing generally from 6 to 10 pc of iron oxide

by firing in this way

The presence of clay in soils exerts an important action on the growth of plants Ammonia is retained by clay in such a manner that it cannot be removed except by an excessive amount of washing; in fact, clay will remove ammonia and many organic substances from manure, and still retain them in a form readily available for plants.

Kaolinite Al<sub>2</sub>O<sub>3</sub> 2SiO<sub>2</sub>,2H<sub>2</sub>O A comparatively rare crystalline mineral which occurs in Anglesey, in Colorado, and a few other localities in the form of minute hexagonal plates, having a perfect basal cleavage, and belonging to the monoclinic system. Kaolinite was formerly

supposed to form the base of all clays, but this view is no longer held, and the term should be restricted to the crystalline mineral described

Kao-lin, China clay The Chinese term kao-lin' (literally meaning 'high ridge') was applied to the white earth from which porcelain was made, and it corresponds to the English china clay, so called because it is used for the making of 'china,' or porcelain. These terms are now applied to all the purer residual clays which contain a high percentage of hydrated aluminium silicate (Al<sub>2</sub>O<sub>3</sub>·2SiO<sub>2</sub>,2H<sub>2</sub>O), and which consequently give on analysis 33-40 p c. alumina (Al<sub>2</sub>O<sub>3</sub>), 45-60 p c. silica (SiO<sub>2</sub>), 11-14 p.c. water, with very small percentages of iron oxide, soda, potash, lime, magnesia, &c. Clays of this class are worked in Cornwall and Dovon, at St Yrieix near Limoges, Zettlitz in Bohemia, in the United States, in South Africa, As mined, the clay generally contains a large percentage of quartz-sand and mica, and it is refined by washing In Cornwall the china clay is usually the result of the weathering of pegmatite, and the felspar of the parent rock is often decomposed to a depth of 60 to 100 feet. In this condition it is known as china clay rock, and the workman breaks down the crumbling surface with a pick and then directs a stream of water over this to wash down the material, which is passed into tanks where the suspended matter settles out, the quartz, clay, and mica being deposited according to the sizes of the particles, and so separated from each other The china clay is allowed to dry to a stift paste in the tanks, and is then removed to the drying kilns, where it dies until it can be cut into blocks ready for shipment It is a soft, friable, seetile, white substance, showing irregularly shaped granules under the microscope, and possessing only a moderate plasticity when mixed with water Cf Coon, Trans Eng Coramic Soc 1910-1911, 10, 81 China clay is used in the manufacture of porcelain and white earthenware, in paper making, and the filling of calico, and also in the manufacture of alum, aluminium sulphate, and ultramarine The fine mica separated from the china clay in the washing is used for weighting coarse papers, paper boards, &c

Ball clay, Potter's clay The name given to certain sedimentary clays which fire to whitish or light-buff colour, and which possess high plasticity and tensile strength when in a moist condition. They are worked in Devon-shire and Dorset, and in some parts of the United States, &c., and are chiefly used in the manufacture of white earthenware. They are generally finer in grain than china clay, contain more organic matter and other impurities, and are consequently more plastic in a most state, and fire to a vitreous mass at a lower temperature. They usually also contain a higher percentage of silica than china

Pipe clay A rather loose term, which is applied to white clays which contain a relatively high percentage of silica, and which have been used without admixture for several hundred years for the making of tobacco pipes and also for common pottery. These clays are of various geological origins, and in some localities the

highly silicious clay found in pockets in lime stone formations, and resulting from the decomposition of argillaceous limestones and chert rock, is used in this way. Ball clay is also used for the same purposes, and is sometimes consequently

spoken of as pipe clay.

Plastic clay. This term has been specially applied to those beds of the Lower Tertiary or Eocene strata, which occur between the London clay and the chalk, and which may be recognised at Woolwich, Reading, Blackheath, Bognor, and in the neighbourhood of Paris, &c This clay is mottled red or purple, and fires to a red colour Some of our earliest pottery was manufactured from this clay, and it is still worked in Hampshire, in the lile of Wight, and elsewhere.

Slate clay occurs in Devon, Cornwall, &c., differing considerably in different districts. is massive, interspersed with mica, and of a greenish or greyish colour It is sometimes hard and slatey, being readily cleavable into slabs, whilst in another form, the killas of the Cornish mines, it has an earthy fracture, is soft and frangible, and weathers to a more or less plastic clay It is chiefly used for the manufacture of common bricks and tiles, and, when fairly free from lime, magnesia, and iron, for fire-

Common clay is the term often applied to the ordinary surface and drift clays which have been used for centuries for making common

pottery and bricks

Fireclays. As the name implies, these clays are of a very refractory nature, and although, strictly speaking, practically all the pure residual clays come under this head, the term 'fireclay' has been restricted in use to certain coal-measure shales which have a high fusion point, and which are used mainly for the manufacture of firebricks, gas retorts, glass pots,

furnace linings, &c

These fireclays usually underlie the coal seams, and are often worked in conjunction with Certain districts are particularly the coal noted for the quality of their fireclays, and in England the Stourbridge coal-measure shales have long been considered of the highest quality, and fireclay-mines are extensively worked in the Stourbridge district Good fireclays are also found and worked in the Lancashire and Yorkshire, Derbyshire, and Glasgow coal-fields. The fireclay shales are mostly greenish grey in colour, compact, dense, and of varying degrees of hardness On exposure to the weather they usually crumble down, and in time become converted into plastic clay

The weathering process may be greatly accelerated by grinding the shale, but this is only done when the clay is to be used for the more expensive fireclay goods Only those coal-measure shales which contain small percentages of oxide of iron, lime, and magnesia. are sufficiently refractory to be used as fireclays. and although no definite fusion point has been fixed, it seems advisable that no clay having a fusion-point lower than 1500° should be classed as a fireclay. In the manufacture of fireclay goods, the partially weathered or ground fire-clay is usually mixed with a certain proportion of 'ganister,' coarse sand, burnt clay, or other refractory material which produces in the fired ware a certain open texture, which enables it

better to withstand rapid changes of temperature. Certain silicious fireclays and other coalmeasure clays are used in the manufacture of salt-glazed drain pipes, and also for making common culinary ware, buff-coloured tiles, terracotta, glazed bricks, &c.

Chemical analysis of clay. From 1-2 grams of finely powdered clay, dried at 120°, are weighed into a platinum crucible, and mixed with about six times its weight of a fusion mixture consisting of potassium and sodium carbonates in equimolecular proportions mixing is best performed by first stirring together the clay with about the same bulk of fusion mixture by means of a dry glass rod, and then adding more and more fusion mixture with repeated stirring, until the whole is thoroughly mixed. The glass rod should then be rubbed in a little fusion mixture, which is added to the mass in the crucible. The covered crucible is gradually heated over a Bunsen burner (a Teclu burner may be advantageously used) until the whole mass is in a state of tranquil fusion crucible is allowed to cool, and afterwards the bottom is raised just to dull redness, twice successively, in order to facilitate the removal of the mass from the crucible. After cooling, a few drops of distilled water are added, and the crucible very gently warmed until the mass detaches itself. The contents of the crucible are then washed into a large platinum dish (a porcelain dish may be used without seriously impairing the accuracy of the estimation), covered with water, and then warmed until the fused mass is completely softened. Hydrochloric acid is then added, a few drops at a time, until all effervescence ceases The contents of the dish are evaporated on a water-bath, with constant stirring at the later stages, to ensure a fine powdery residue from which the last traces of hydrochloric acid are readily driven off. When cold, the contents of the dish are moistened with moderately strong hydrochloric acid, and allowed to stand for one hour, water is then added, and the dish again heated on the water-The clear liquid is poured through a filter, and the residue in the basin, which consists of silicic acid, is washed repeatedly by hot water until it remains white when moistened with hydrochloric acid. The residue is then transferred to the filter, and washed with hot water until the filtrate no longer gives a precipitate with silver nitrate solution The filter and contents are then placed in a platinum crucible and dried, and afterwards gradually heated, and finally ignited strongly until the paper is completely incinerated, and the weight constant. In order to ensure the complete separation of the silicic acid, it is necessary to evaporate the filtrate down to dryness, and treat exactly as before, and this process may be repeated again if very great accuracy is required. After filtering, the filter paper and residue are dried and added to the original silicic acid residue before ignition. The contents of the crucible are then weighed as silicon dioxide (SiO2) If the clay contains titanic acid (as many clays do), this will accompany the silica. In order to determine the former, the contents of the crucible are covered with strong hydrofluoric acid, a few drops of sulphuric acid added, and the crucible warmed on the water-bath until all the silicon

fluoride and the sulphuric acid have been driven off. The residue is then weighed as titanic oxide (TiO<sub>2</sub>), which should be identified by the violet-coloured bead with microcosmic salt.

The filtrate is made up to definite volume, and divided into two parts, one for the determination of the iron, and the other for the determination of the alumina, calcium, and magnesium. The iron is estimated as ferric oxide, and determined either by reduction with zine and titration with potassium permanganate solution,

or by the iodimetric method

The other portion of the filtrate is heated on the water-bath, and ammonia added drop by drop, with repeated stirrings, until there is a slight excess of ammonia present The liquid is then boiled for a moment and filtered. The precipitate is washed twice with hot water; washed back into a dish, and dissolved in hydrochloric acid; reprecipitated and filtered as before, and finally washed with hot water until the filtrate gives no precipitate with silver nitrate solution. The washing should be repeated immediately the previous washing has disappeared from the filter, and a jet of hot water should be directed into the precipitate so as to break it up as much as possible. The filter as to break it up as much as possible paper and contents are then dried, the residue removed into a platinum crucible, the paper separately incinerated, and then placed in the crucible, and the whole strongly ignited until the The contents are then weight is constant weighed as alumina (Al2O3) and ferric oxide (Fe<sub>2</sub>O<sub>3</sub>), the amount of ferrie oxide separately determined as above being deducted from the total, leaving alumina only (v art Aluminium) The filtrate from the alumina and ferric oxide is concentrated, and ammonium oxalate solution added while hot. The precipitate is filtered off, thoroughly washed with hot water, dried, strongly heated, and weighed as CaO filtrate from the above is then concentrated, and the magnesium precipitated by sodium phosphate and ammonia. The precipitate is filtered off, dried, and ignited until the weight is constant The ignited residue consists of Mg<sub>2</sub>P<sub>2</sub>O<sub>7</sub>, and the weight must be multiplied by 0 3624 to give the amount of magnesia (MgO)

Determination of the alkalis. The method introduced by Lawrence Smith (Amer. J. Sci. [2] 50, 269) is the most trustworthy, and is -based on the decomposition of silicates by fusion with calcium carbonate. In practice, the dried and finely powdered clay is mixed with its own weight of ammonium chloride, and four times its weight of calcium carbonate. These latter materials must be as free as possible from alkali The ammonium chloride used is prepared by subliming the commercial salt, and the calcium carbonate is prepared by dissolving the purest calcite obtainable in hydrochloric acid, and precipitating with ammonia and ammonium carbonate. The precipitation is carried out in a large porcelain dish, and, after settling, the clear liquid is poured off, and the precipitate washed by decantation until free from chlorides. A special platinum crucible is used for the fusion, about 8 cm. long, and 2 cm. diameter at the top, tapering to  $1\frac{1}{2}$  cm. at the bottom. About 0.5 gram clay is weighed out, and mixed by rubbing in an agate mortar

with the same weight of ammonium chloride, and afterwards 3 grams of calcium carbonate are added, and the whole thoroughly mixed The mixture is then transferred to the crucible. and the mortar is cleaned out with about 1 gram of calcium carbonate, which is added to the mix-ture in the crucible. The covered crucible is supported in an inclined position, and very gradually heated over a small Bunsen flame for about 15 minutes, until no more ammonia is given off, and then the temperature is raised until the lower three-fourths of the crucible is a dull-red, and this temperature is maintained for I hour The crucible is cooled and the contents removed (by tapping the inverted crucible) into a porcelain or platinum dish. If the sintered mass sticks to the crucible, a little water should be added, and the crucible gently warmed until the mass detaches itself, and the whole can be washed into the dish About 60 c c of distilled water are added, and the covered dish heated on a water-bath for 30 minutes, the water lost by evaporation being replaced, and the larger particles reduced by rubbing with an agate pestle. The clear liquid is decanted through a filter, and the residue washed by decantation four times, and then transferred to the filter and washed until the filtrate gives no precipitate with silver nitrate. The filtrate contains the alkalis as chlorides

The calcium is precipitated by the addition of ammonia and ammonium carbonate to the filtrate, which is heated and filtered. The residue, which contains some alkali, is dissolved in hydrochloric acid, again precipitated by ammonia and ammonium carbonate, and filtered. The combined filtrates are then evaporated to dryness in a platinum dish, and the ammonium salts iemoved by careful heating over a moving flame. The residue is dissolved in water, and any traces of calcium left are precipitated by the addition of ammonia and ammonium oxalate After standing 12 hours, the liquid is filtered off, evaporated to dryness in a weighed platinum dish, and gently heated. The alkali chlorides remaining in the dish are weighed, and the potassium determined by dissolving the chlorides in water and precipitating the potassium as platino-chloride The sodium is potassium determined by difference

Rational analysis of clay. To determine the relative amounts of clay substance, quartz, and undecomposed felspar, dry the clay to constant weight, and weigh about 5 grams of the dried and finely powdered clay into a porcelain dish, add 50 c c. of concentrated sulphuric acid and 100 c.c of water, and then heat the mixture for 3 hours on a water-bath, and afterwards on a sand-bath, until the acid commences to fume. After cooling add water, and decant the clear liquid into a 2-litre beaker To the residue in the dish add 10 c c. of Lunge's solution (100 grams crystallised sodium carbonate, and 10 grams sodium hydroxide, dissolved in 1 litre of water), boil for 5 minutes, dilute with water, and decant into the beaker. Treat the residue in the dish similarly with concentrated hydrochloric acid, and repeat the treatment with Lunge's solution and hydrochloric acid, diluting with water after each treatment, and decanting the clear liquid into the beaker. Then wash the residue on to a filter with

dilute hydrochloric acid. Stir up the contents in the beaker, and allow to stand for an hour. Decant off the clear liquid, taking care to retain the residue in the beaker. Wash the residue into the dish, and repeat the treatment with Lunge's solution and acid as above, and pass through the filter. Wash with dilute hydrochloric acid, dry and ignite the filter paper and contents in a weighed platinum crucible. This residue is reckoned as felspar and quartz, and the loss suffered by the clay in this treatment as clay substance.

To determine the quartz and felspar, add about 5 cc of water, 4 drops of sulphune acid, and 15 cc. of hydrofluoric acid to the residue in the crucible, and heat the whole carefully on a sand-bath until nearly dry. Add more hydrofluoric acid, and again heat as before. After cooling, dissolve the mass in water, wash into a dish, and add ammonium chloride and ammonium hydrate to precipitate the hydroxides of aluminium and iron. Filter, wash with hot water, heat and weigh. This weight is multiplied by 5 451 to obtain its equivalent in potash felspar, and the difference between this and the weight of the original residue is taken as quartz.

Many ceramic chemists calculate the rational constitution of the clay from the ultimate analysis, taking the potash as derived entirely from undecomposed felspar (reckoned as  $K_2O \cdot Al_2O_3 6SiO_2$ ), the alumina and iron as derived from felspar, and the clay substance (reckoned as  $Al_2O_1 \cdot 2SiO_2 \cdot 2H_2O$ ) and the silica as derived from the clay substance and quartz.

CLAYITE. A name suggested by J W Mellor in 1909 for the non-crystalline variety of kaolimte H<sub>4</sub>Al<sub>2</sub>Sl<sub>2</sub>O<sub>3</sub>, of which china-clay and most other clays are largely composed. The same name had been earlier used by W J Taylor, in 1859, for an uncertain alteration product (of tetrahedrite?) composed of sulphur, arsenic, antimony, lead, and copper.

L J. S

CLAY IRONSTONE v IRON, ORES OF

CLAY SLATE v SLATE

CLAYTON CLOTH RED v Azo- COLOURING

CLAYTON YELLOW v PRIMULINE AND ITS DERIVATIVES.

CLEVEITE. A variety of uraninite, consisting of a uranate of uranyl, and oxides of lead and the rare earths Usually obtained from Norway and Sweden Hillebrand (Bull. U.S. Geol Survey, 78, 43, 83) found the percentage of rare earths to be about 13, and that on heating with sulphuric acid, over 2 p c of nitrogen was evolved. Ramsay (Chem. Soc. Trans 1895, 1107) proved that the gas evolved was not nitrogen, but helium (q v) Only half the helium is evolved by heat alone, and, moreover, in some cases the disengagement of the gas is accompanied by considerable evolution of heat (Proc Roy. Soc 1898, 64, 140). V BROGGERITE and PITCHBLENDE.

CLEVE'S ACIDS. 1-Naphthylamine-6 and 7-

Sulphonic Acids, v NAPHTHALENE.

CLICHY WHITE. A white lead manufactured at Clichy, in France (v. White lead, art. Lead.)

CLOTHBROWN, ORANGE, RED, SCARLET, v. Azo-colouring matters.

CLOVE OIL v. OII S, ESSENTIAL.

clover. The name by which several leguminous forage plants are known. The chief species are white clover (Trifolium repens, Linn); red clover (T. pratense, L.); alsike clover (T. hybridum, L.); crimson clover (T. incarnatum, L.), yellow suckling clover (T. dubium, Sibth.); mammoth red, or zigzag clover (T' medium, L.); hop clover, or hop trefoil (T. procumbens, L.) All the above belong to the true clovers, to the genus Trifolium. Also included as clover are yellow clover, or trefoil (Medicago lupulina, L.); Bokhara clover, or melilot (Melilotus alba, Desn.); Japanese clover (Lespedeza striata, Hook et Arn.); and Soola clover, or sulla (Hedysarum coronarium, L.). The following table gives the average composition of various clovers at the time of flowering (Kellner):—

			Sol carbo-							
	,	Wate	er Pro	tein :	Fat	hydr	ates	Fibre	Ash	
T	repens .	81	5 4	ŀ4	0 8	8 6	9	43	2 1	
	pratense .	79	0 3	4	0 '	7 9	4	59	16	
T	hybridum .	81	8 2	8 2	0 '	7 7	0	62	15	
T.	incarnatum.	81	5 2	28	0 '	7 7	0	$6 \cdot 1$	19	
T'	procumbens	80	0 3	5	0.8	8	4	57	16	
$M\epsilon$	Ārlotus alba .	79	7 4	.1	0.8	3 7	4	57	$2 \cdot 3$	

The following are analyses of hay from various clovers .—

Sol. carbo-Water Protein Fat hydrates Fibre Ash T. repens 97 15.7 29 393 24.1 8.3 . 16 5 13 5 2 9 37 1 matense 24 0 6 0 T hybridum . 160 136 31 345 257 71 T. incarnatum. 16 7 12 0 2 4 35 5 26.27 2 T. procumbens 16 0 15 4 3 4 33 2 Melilotus alba. 15·7 16·7 2 8 27 9 24.575 30 3 80 Lespedeza striata 9:1 13 7 4 0 47 5 21 6 4.1

White or Dutch clover is a low, creeping perennial plant, valuable for grazing Its growth in pastures is encouraged by phosphatic and potash manuring, but discouraged by introgenous manures. In America, it is said that its blossoms salivate horses, and that consequently much clover in pastures is an objection where horses graze. White clover furnishes most excellent pasturage for bees, the honey which it yields being colourless and of high quality.

Red clover or broad clover is usually grown as a bennial, but will sometimes stand longer. It is often grown as a hay crop, or for cutting for green forage. It is the variety most susceptible to 'clover sickness,' and on many soils can only be grown safely once in 10 or 12 years.

A variety of red clover, T pratense var. perenne, known as 'cow grass,' is not so prone to clover sickness, and lasts longer. It is, however, of slower growth, and usually yields but one crop per year of hay.

Alsike clover has whitish and pinkish flowers, and is a perennial; it is hardly susceptible to clover sickness.

Crimson clover, or 'trifolium,' is also a somewhat slow-growing crop, and is an annual.

Yellow suckling clover is a smaller plant than the other clovers, and is usually only grown in mixtures with rye-grass, &c. It is an annual.

Zigzag clover, known in America as mammoth red clover, ripens later than red clover, and is taller and ranker. Bokhara clover, or large white clover, is a biennial, with small white flowers. This plant is remarkable for its strong odour, more pronounced when dried, of coumarin; animals do not eat it readily until they become accustomed to it.

Soola clover is a perennial resembling red clover, but is richer in soluble carbohydrates

(v Scurti, Chem Zentr 1910, 1 1632).

As the table of analyses shows, clovers are remarkable for the large amounts of nitrogenous matter and of ash which they contain. Of the former, however, a considerable proportion is of a non-protein nature, particularly in the immature plants. In the case of red clover, for example, it has been found that about 40 p.c. of the total nitrogenous matter is present as amides in very young plants, and that the proportion diminishes with growth, until, when the plants are in full bloom, it is about 20 p.c., when fully ripe it sinks to about 12 p.c.

The ash of clovers is remarkable for its richness in lime and its poverty in silica (in which grasses and cereal straws are so rich).

The following are Wolff's analyses of various clover ashes:—

The very high ratio of lime to phosphorus pentoxide in the ash is noticeable.

Clover hay has proved very successful as a constituent of the rations of fattening oxen and

sheep

Out of about 200 feeding experiments with nearly 1000 cattle, conducted in Britain between 1833 and 1908, 17 included clover hay in the daily ration. The average daily increase in live weight per head of the whole was 1 803 lbs., while the consumption of total digestible matter per 1 lb. live weight gained averaged 9 lbs. In the 17 cases where clover hay formed a portion of the food, the figures are 2 1 lbs., and 7 47 lbs respectively (Ingle, Journ High and Agri Soc of Scotland, 1909)

With sheep, the results are equally significant 2390 sheep fed in various ways, between 1844 and 1909, gave an average weekly increase per head of 2 06 lbs, and consumed an average of 7 43 lbs total digestible matter per 1 lb increase in live weight; while 384 sheep receiving clover hay as part of their ration, gave an average weekly increase of 2.58 lbs. per head, and consumed only 6 20 lbs of total digestible matter per 1 lb. increase in live weight (ibid 1910)

CLOVER FLOWERS. It has long been known that clover flowers dye a yellow colour on aluminium mordanted fabrics, and in the past they have been employed to a minor extent for dyeing purposes. Three varieties have been chemically examined, viz. those derived from the Trifolium pratense, the T. incarnatum, and the T. repens.

Trifolium pratense—The flowers known as the 'common red clover,' according to Power and Salway (Chem. Soc. Trans. 1910, 97, 231), contain in addition to isorhamnetin (quercetin, monomethyl ether) and a glucoside of quercetin, melting-point 235°, numerous other phenolic substances, which, judging by their chemical

properties, appear to be closely allied to the colouring matters of the flavone group. These are described below.

 $Pratol\ C_{15}H_8O_2(OH)$  OCH 3, colourless needles, melting-point 253°, readily soluble in hot aqueous sodium carbonate and codium hydroxide with a pale yellow colouration, yields the acetyl compound,  $C_{15}H_8O_2(OC_2H_3O)$  OCH 3, and is probably a hydroxymethoxyflavone.

A yellow compound,  $C_{18}H_{10}O_{7}$ , thin yellow plates, melting-point about 280°, is soluble in alkalis with a yellow colour, and its solution in sulphuric acid exhibits a brilliant green fluorescence. It contains a methoxy group and gives an acetyl compound,  $C_{18}H_{6}O_{7}(C_{2}H_{2}O)_{4}$ , melting-

point 145°-147°.

Pratensol, C<sub>17</sub>H<sub>9</sub>O<sub>2</sub>(OH)<sub>3</sub>, is very readily soluble in alcohol, dissolves in alkali carbonates yielding yellow solutions, and its alcoholic solution gives with ferric chloride a greenish-black colouration Triacetylpratensol C<sub>17</sub>H<sub>9</sub>O<sub>5</sub> (C<sub>2</sub>H<sub>3</sub>O)<sub>3</sub>, colourless slender needles, melts at 189°.

A phenolic substance, C<sub>15</sub>H<sub>7</sub>O<sub>3</sub>(OH)<sub>3</sub>, colourless needles, melting-point 225°, is soluble in alkali hydroxides, and gives with alcoholic ferric chloride a dark green colouration. The acetyl derivative, silky needles, has melting-point 209°.

The glucoside trifolin  $C_{12}H_{12}O_{11}$ ,  $H_{2}O$ , pale yellow needles, melts and decomposes at about 260°. It is soluble in alkalis with an intense yellow colouration, and dissolves in sulphuric acid, forming a yellow solution, which rapidly develops a brilliant green fluorescence. When hydrolysed it yields thamnose and trifolitin  $C_{16}H_{10}O_{6}$ , melting-point about 275°, readily soluble in alcohol,

$$C_{22}H_{22}O_{11}=C_{16}H_{10}O_6+C_6H_{12}O_5$$

Alkalis dissolve trifolitin with an intense yellow colour, alcoholic ferric chloride gives a dark green colouration, and alcoholic lead acetate an orange-yellow lead salt. It contains no methoxy group, and is unaltered when heated for several hours with 30 p.c. aqueous potassium hydroxide. It does not appear to belong to the flavone group, and differs from the flavone compounds by the fact that it does not give an oxonium salt with sulphune acid, and only with difficulty a potassium compound by means of alcoholic potassium acetate. It may possibly consist of a tetrahydroxyphenylnaphthoquinone. The acetyl compound when rapidly heated melted at 116°, resolidified at a higher temperature, and finally melted at 182°.1

The glucoside isotrifolin  $C_{22}H_{22}O_{11}$ , isomeric with trifolin, consists of pale yellow needles, melting-point about 250°, and when hydrolysed yields similarly to the latter trifoliin,  $C_{16}H_{10}O_{6}$ , melting-point 275°. Though in general behaviour it is very similar to trifolin, it is much more soluble in alcohol, and does not appear to be identical with this glucoside.

identical with this glucoside.

In addition to these compounds the flowers contain salicylic acid, commaric acid, myricyl alcohol C<sub>31</sub>H<sub>63</sub>OH, heptacosane C<sub>27</sub>H<sub>56</sub>, hentriacontaine C<sub>31</sub>H<sub>64</sub>, sitosterol C<sub>27</sub>H<sub>46</sub>O, trifolianol C<sub>21</sub>H<sub>34</sub>O<sub>2</sub>(OH)<sub>2</sub>, palmitic, stearie, linolic, oleic, linolene, and isolinolenic acids.

Both in its melting-point and that of its acetyl derivative there is a marked resemblance between trifolitin and kaempferol.

Trifolium incarnatum.—A considerable difference is exhibited between the constituents of the 'carnation or crimson clover flowers' and those of the T. pratense or 'common red clover.'

According to Rogerson (Chem. Soc Trans 1910, 97, 1006) these flowers contain pratol C<sub>15</sub>H<sub>8</sub>O<sub>2</sub>(OH)(OCH<sub>3</sub>), free quercetin, and a glucoside of quercetin C<sub>21</sub>H<sub>20</sub>O<sub>12</sub>,3H<sub>2</sub>O, to which the name incarnatrin is applied. This latter crystallises in yellow prismatic needles, meltingpoint 242°-245°, dissolves in sulphuric acid with formation of a green fluorescent solution, and when hydrolysed yields quercetin and glucose according to the equation

$$C_{21}H_{20}O_{12}+H_{2}O=C_{15}H_{10}O_{2}+C_{6}H_{12}O_{6}$$

Incarnatrin is not identical with the quercimentum of Perkin (Chem. Soc. Trans. 1909, 95, 2181)

In addition to these substances the flowers yield furfuraldehyde, benzoic and salicylic acids, a trace of p-coumanc acid, incarnatyl alcohol  $C_{34}H_{69}OH$ , hentriacontane  $C_{31}H_{64}$ , a phytosterol C27H46O, and palmitic, stearic, oleic, linolenic, and isolinolenic acids

Trifolium repens -The flowers of the white clover, T. repens, according to Perkin and Phipps (Chem. Soc. Trans 1904, 85, 58), owe their tinctorial property to quercetin which is present as glucos:de. A G P.

CLOVES. The cultivation of cloves forms the most important industry of Zanzibar and of the island of Pemba In 1913-14 the total erop was 25,431,735 lbs of which 20,692,771 lbs came from Pemba, where about two-thirds of the total area, or about 35,000 to 40,000 acres, are under clove cultivation The crop is, however, very uncertain, the amount in some years not being above one sixth of the above yield. Large trees may furnish from 60 to 70 lbs of dry cloves, although the average yield is only about 31 to 4 lbs per tree per annum The cloves are picked by hand and dried in the sun The value of cloves depends upon the amount of essential oil present, the quantity of which may vary from about 6 to 20 p c Of this oil about 84 to 90 pc consists of eugenol (V Clove Oil in art Oils, Essential)
COAL v Fuel

COAL GAS v. GAS, COAL

COBALT. Sym Co At wt 58 97.

Cobalt usually occurs combined with arsenic or sulphur, and almost invariably associated with nickel and other metals Free cobalt occurs only in meteorites.

The most plentiful and important ore of cobalt is smalline or tin white cobalt, consisting of arsenide of cobalt, nickel, and iron It occurs abundantly at Schneeberg and many other localities where bismuth is found (v. BISMUTH), ın Bohemia, at Balmoral near Pretoria, near Sudbury in Ontario, very pure at Tunaberg in Sweden, and free from nickel in Colorado. triarsenide, it forms skutterudite CoAs3 Sulpharsenide of cobalt occurs as cobalt glance, grey cobalt, or cobaltine, abundantly at Vena (Sweden) and in Norway; in smaller quantities in Silesia and Cornwall As hydrated arsenate it forms cobalt bloom or erythrite Co3As2O8,Aq, found frequently as a peach-coloured incrustation on other arsenical ores, being produced by their decomposition. As sulphide, cobalt occurs

in small quantities in cobalt pyrites and in linnaeite.

Black earthy cobalt is a hydrated oxide of cobalt and manganese. It is essentially a variety of wad (hydrated manganese oxide), containing sometimes as much as 40 p.c. of cobalt. Surface deposits containing oxides and sulphides of cobalt and of nickel occurat New Caledonia, with veins of chrome ore and nickel and magnesium silicates, it is smelted on the spot and exported as a regulus to England Cobalt occurs in small quantities in many other minerals, in pyrites and in certain non ores, as oxide in black oxide of manganese, as arsenide in mispickel, sometimes to the extent of 10 p c., in selenide of lead, and in cerite, and in some peats and coals.

At the present time the most important source of cobalt, whence almost the whole of the cobalt of commerce is obtained, is the district surrounding Cobalt city, near the shores of Lake Temiskaming in North Ontario very rich silver ores are found, containing frequently 2000 or silver per ton, and sometimes as much as 60 p c These ores also contain a high percentage of cobalt and nickel, chiefly as arsenides, and, after extraction of the silver, are

sent to nickel and cobalt refiners.

Extraction —The methods adopted by manufacturers for the extraction of cobalt from the ores vary with the nature of the ores 
The ores are first concentrated by fusing with a suitable flux, whereby silicious matter and a portion of the iron are removed In the case of the Ontario ores, the speiss consists chiefly of arsenides of cobalt, nickel, iron, and copper The speiss is then calcined, the iron is oxidised to ferric oxide, part of the arsenic is removed, a little less being left than is required to combine with the iron. The mass is then dissolved in hydrochloric acid, and bleaching powder and lime are added, whereby the iron is precipitated as arsenate and hydroxide. The liquid is then treated with sulphuretted hydrogen to precipitate copper and other heavy metals, excess of the gas evaporated off, and the cobalt precipitated as oxide by the addition of bleaching powder The mekel in the solution is subse-

quently precipitated by potash or lime
In the case of New Caledonian ores, consisting chiefly of sulphides, the treatment of the concentrated ore is somewhat different The matt, consisting chiefly of sulphides, is carefully calcined at a moderate temperature, and converted into sulphates. These are dissolved in water, and the non precipitated by careful addition of the requisite quantity of sodium carbonate. The copper and heavy metals are separated by sulphuretted hydrogen, and after boiling off the excess, the cobalt is precipitated by sodium hypochlorite (bleaching powder cannot be used, owing to the sulphates in the solution), and the nickel in the remaining liquid

precipitated by sodium carbonate.

In Herrenschmidt's process for the treatment of New Caledonian ores, these are treated with a solution of ferrous sulphate, when cobalt, nickel, and manganese pass into solution, and iron remains in the residue. Copper is first separated, and then the cobalt, nickel, and manganese precipitated by sodium sulphide. The manganese is extracted by ferric chloride, the nickel and cobalt sulphides roasted to

sulphates, and converted into chlorides by calcium chloride. A part of the solution is treated with lime water, and the suspended precipitate oxidised by chlorine and added to the other portion, whereby the cobalt is precipitated and nickel chloride remains in solution.

Until of late years the only market for cobalt compounds was in the ceramic industry, in which it was employed for whitening the body of china and in the production of blue colour employed in the enamelling trade, the manufacture of blue-coloured signs, the colouring of glass and in the manufacture of pigments and

as driers for paints.

Metallic cobalt is not at present used to any extent in the arts, though its utility is becoming more fully recognised, especially in electroplating and in high-speed steels and in 'stellite, an alloy of cobalt, chromium and tungsten. Commercially it is always prepared by heating the oxide with charcoal, it then contains small quantities of carbon Pure cobalt is prepared by heating cobalt oxalate Co(CO2)2, when the carbon and oxygen pass off as carbonic anhydride, leaving the pure metal as a powder A button of the fused metal may be obtained by fusion of the oxalate alone, or of the oxide mixed with charcoal under a layer of powdered glass containing no lead or other reducible metal, the fusion is preferably performed in a lime crucible at the highest temperature of a furnace Pure cobalt may also be prepared by electro-deposition with platinum electrodes (Winkler, Zeitsch anorg Chem 8, 1) Cobalt is also produced as a grey metallic powder by igniting the oxide at a red heat in a current of hydrogen If too low a temperature has been used, the metal produced is pyrophoric, igniting and becoming reoxidised in contact with the air

Carbon monoxide rapidly reduces cobalt oxide at all temperatures above 600°, the reduction being complete at 900° With chair coal complete reduction takes place at 900°, the final product containing about 2 p c of earbon Reduction with aluminium by the thermit process gives a metal containing about 0.1 p.c. of aluminium or even less (Kalmus, J Ind Eng.

Chem. 1914, 6, 107).

Properties —Cobalt is a brilliant silver-white metal, with a slight bluish cast It is the most magnetic metal after iron, and retains its magnetism at high temperatures; above 1115°

it passes into a non-magnetic variety.

Pure cobalt is harder and more tenacious than iron, its breaking weight being about 50 kilos per sq millimetre Valenciennes states that cobalt, after fusion and annealing, may be rolled into spiral bands like good iron (Compt. rend. 70, 607). Cobalt melts at 1530° (Copaux), 1490° (Henning), 1478° (Kalmus & Harper), and boils at 2415°/30 mm. Its sp gr. is 88, and hardness, 5.5. Its specific heat (20°-100°) is 0.104, specific resistance, 55; and the temperature coefficient of resistance, 0 0055. According to Kalmus & Harper (J Ind. Eng. Chem 1915, 7, 6), pure cobalt prepared by reduction of cobalt oxide has  $D_4^{17}$  8·7918 (unannealed),  $D_4^{14}$  5·8·8105(annealed),  $D_{1}^{\overline{16}5}$  8 9253 (swaged); m p. 1478±1·1°; tensile strength, cast and unannealed, 34,400 lb.

per sq. in.; annealed 36,980 lb. per sq. in; and rolled, more than 100,000 lb. per sq. in.; compressive strength, cast and unannealed, 122,000 lb per sq. in, annealed, 117,200 lb. per sq. in., electrical resistance 89 64 ohms per cm 3, specific heat, from 0° to 890° 0.1058  $+0.0000457t+0.000000066t^2$ . The hardness of cast cobalt is considerably greater than that of either iron or nickel, being approximately 124 0 (load 3500 lb ) The presence of from 0 06 to 03 pc. of carbon in the metal increases the tensile and compressive strength, and the electrical resistance, but has little effect on the hardness. Pure cobalt may be machined in a lathe as readily as pure iron or nickel, but it is somewhat brittle, cobalt containing a small quantity of carbon machines like mild steel, and whilst pure cast cobalt cannot be rolled or swaged without developing cracks, cobalt containing carbon may be rolled or swaged down from cast bars to any extent provided that the metal is worked at a red heat (Chem Soc Abst 1915, ii 99). It is slowly soluble in hydrochloric and sulphuric acids, with evolution of hydrogen, quickly soluble in nitire acid; it is not attacked by hot or cold alkali, but if used as anode in the electrolysis of alkaline electrolytes, passes into a colloidal solution of cobaltous oxide It exhibits also the passive state, similar to that of iron and other metals, but the tendency to do so is slight Cobalt occludes from 60 to 150 times its own volume of hydrogen, the amount varying with the mechanical condition of the metal. The powdered metal combines, when heated, with chlorine, bromine, and rodine. It decomposes steam at a red heat, and, if finely divided, ignites when heated in oxides of nitrogen, and in a current of amimonia gas at  $500^{\circ}$  forms a intiide  $\mathrm{Co_4N_2}$ . When finely divided cobalt is heated to  $150^{\circ}\text{--}200^{\circ}$  in carbon monoxide at 100 atmos pressure, orangered crystals of cobalt carbonyl Co(CO), are formed, melting at 51°, soluble in alcohol or ether, but insoluble in water, and decomposing above 52°, yielding Co(CO)<sub>3</sub> which further decomposes (Mond, Chem. Soc. Trans. 1910, 798) (v. COBALT. CARBONYLS)

Cobalt may be deposited as a fine lustrous coating on metals by the electrolysis of a neutral solution of the double sulphate of cobalt and ammonium, using a current of about 6 amperes per sq foot of cathode surface, and an anode of cobalt (A Gaiffe, Compt rend. 1878) presence of ammonium chloride appears to hinder, whilstammonium sulphate assists, the deposition. The deposit of cobalt is harder, more tenacious, and of greater beauty than that of nickel (Troost).

(For the properties of cobalt containing other

metals, v Alloys of cobalt)

Detection —All ores and compounds of cobalt impart a blue colour to a bead of borax on fusion; if present in too great excess, the bead appears black. Heated on charcoal with reducing flux, all substances containing cobalt produce metallic particles which are attracted by a magnet (distinction from all other bodies except iron and nickel).

Salts of cobalt, when strongly heated with alumina before the blowpipe, give a fine blue colour (Thenard's blue) Sulphuretted hydrogen produces no precipitate in an acid solution, but gives a black precipitate of sulphide in

alkaline solutions, the precipitated sulphide being insoluble in dilute acids. In a solution containing free acetic acid, potassium nitrite produces a yellow precipitate of potassium co-baltimitrite  $K_3\mathrm{Co}(NO_2)_6$ , the whole of the cobalt being thus precipitated on standing in a warm place for some time If a mineral acid is present, it should first be neutralised by potassium hydroxide.

Potassium thiocyanate with cobalt salts gives a red colouration, and, on the addition of alcohol and ether, forms a deep blue ethereal layer.

Cobalt is also precipitated as a red precipitate by the addition of a solution of nitroso-βnaphthol to its alcoholic or acid solutions nickel is not thus precipitated, this forms a useful means of separation of the two metals, the sulphides being dissolved in aqua regia, evaporated to dryness, taken up with a little water and hydrochloric acid, and precipitated as above

Estimation —Ores containing cobalt are usually very complex; they almost invariably contain nickel and many other metals from which the separation is difficult. The following scheme will serve for the estimation of both cobalt and nickel in any mineral, though in the case of certain ores it may be somewhat simplified.

From 2 to 7 grams of the finely ground ore are roasted in a porcelain crucible in a muffle. The residue is boiled with hydrochloric acid containing a little nitric acid until the metallic oxides are dissolved; the solution is diluted, nearly neutralised with ammonia, and boiled with sodium acetate solution, whereby the iron and aluminium are precipitated as acetates containing the greater part of the arsenic (as arsenate of iron) and a little nickel and cobalt The solution is filtered, and the precipitate redissolved in hydrochloric acid and reprecipitated as before, the precipitate being this time free from nickel and The two filtrates are mixed, just neutraised with ammonia, and treated with sulphuretted hydrogen gas; cobalt, nickel, zinc, copper, bismuth, &c, are thus precipitated as sulphides, leaving the manganese and the alkaline and earthy metals in solution. The precipitate is filtered, washed, dried, and roasted, the oxides so produced dissolved in hydrochloric acid, and treated with sulphuretted hydrogen, the copper, bismuth, lead, &c, are thus precipitated and fil-

The solution is boiled till free from sulphuretted hydrogen, and the two metals coprecipitated by the addition of potash to the boiling solution; the precipitate is filtered and washed, nearly dried, and separated as far as possible from the filter. The precipitate is digested in a dish with dilute hydrocyanic acid, then with potash solution, and again with hydrocyanic acid, and warmed until no further solution A minute quantity of residue is left, consisting of paracyanogen and a small quantity of the oxides of nickel and cobalt, it is added to that still remaining on the filter paper, and the whole washed, dried, heated, and weighed.

The solution containing the two metals is boiled to expel the excess of acid, and consists of cobalticyanide of potash and double cyanide of nickel and potash. Precipitated yellow mercuric oxide is added, and the solution boiled for some time. The nickel is thus precipitated, is filtered, washed, dried, heated, and weighed as nickel protoxide NiO. The filtrate is nearly neutralised with nitric acid, and a neutral solution of mercurous nitrate added in excess. cobalt is thus precipitated as mercury cobalticyanide, which is washed, dried, and heated in an open crucible until of constant weight. The product consists of Co<sub>3</sub>O<sub>4</sub>.

From the proportions of cobalt and nickel thus found, the amount of each in the residue on the filter paper, &c., which has been weighed may be calculated, and added to the amounts

directly determined.

In other methods, the copper, &c, are first precipitated by sulphuretted hydrogen, non by ammonia, and the nickel and cobalt precipitated as sulphides. The sulphides are redissolved, and in (1) the total nickel and cobalt estimated by electro-deposition, in (2) the cobalt is first precipitated and the nickel alone determined.

The cobalt in the above solution may also be determined by precipitation as potassium cobaltinitrite, dissolving the precipitate in warm dilute sulphuric acid, evaporating nearly to dryness, and electrolysing after addition of ammonium oxalate and excess of ammonium carbonate

(Copaux, Bull Soc chim iii 29, 301)

The solution containing the sulphides of nickel and cobalt may also be estimated after freeing from excess of acid by boiling with barium carbonate, and adding excess of bromine After 10 minutes the cobalt is completely separated as sesquioxide (Taylor, Proc.

Man Phil Soc 46, 1).

Various volumetric methods have been suggested, but none appears to be perfectly satisfactory

Alloys of Cobalt. Cobalt unites with incandescence on fusion with antimony or arsenic,

forming brittle iron-grey alloys.

With arsenic the freezing-point curve indicates the existence of compounds, Co,As,; Co<sub>2</sub>As, and Co<sub>3</sub>As<sub>2</sub>. Smaltine CoAs<sub>2</sub> forms tin-white octahedra, and skutterudite CoAs<sub>3</sub> occurs as greyish-white octahedra The arsenides are attacked by oxidising agents, and all containing over 38 p.c arsenic are non-magnetic.

Antimon $\hat{y}$  and cobalt are completely miscible in the liquid state. Three classes of alloys exist: magnetic alloys with less than 67 pc. antimony; alloys of from 67 to 80 27 pc, which on heating form CoSb; and alloys above 80 27 pc, which by acids yield a compound, CoSb<sub>2</sub>, as a grey crystalline powder (Ducellicz, Compt rend. 147, 1048)

Cobalt has a marked effect on the electrical properties of copper; 35 pc. gives an alloy with a temperature coefficient of the resistance of only 0 00077. Two series of mixed crystals appear to exist, 0-10 p.c, and 95-100 p.c copper, and alloys within these limits consist of conglomerates of the two sets of mixed crystals Some of the alloys are very ductile. As in many of the cobalt alloys at high temperatures, the cobalt is present as a non-magnetic form  $(\beta)$ , which changes to the magnetic (a) form on cooling. Alloys containing 99 p.c of copper are stated to be magnetic (Sahmen, Zeitsch anorg Chem 57, 1).

Tin and cobalt are perfectly miscible as liquids, and, according to Ducelliez (Compt. partly as cyanide and partly as sesquioxide. It | rend. 148, 562), all alloys are mixtures of CoSn

and Co<sub>3</sub>Sn<sub>2</sub> with one another, or with either of the components. Below 57 6 p.c tin, the alloys are magnetic and brittle; above 66.8 pc tin, the alloys are less brittle and non-magnetic; all are much harder than either component. A compound Co<sub>2</sub>Sn of m p 1151° has also been described

With gold a brittle alloy is produced, that containing 1 part gold and 17 parts cobalt is dark-yellow and very brittle, all the alloys are magnetic. With platinum a fusible alloy is produced, and a silver-white magnetic amalgam

with mercury is used in dentistry.

Alloys with aluminium containing less than 68 5 p.c. cobalt are non-magnetic, the freezingpoint curve indicates a compound CoAl. Cobalt and nickel do not appear to form a compound, the freezing-point curve is almost straight, the composition of the solid being almost identical with that of the liquid in all cases

Addition of iron has no appreciable effect on the melting-point of cobalt, the curve being almost horizontal to 5 pc. cobalt, the crystals having the same composition as the fused mass. Two series of non-magnetic, which transform into four series of magnetic crystals, appear to exist (Tammann, Zeitsch anorg. Chem 45, 205) Steels containing up to 60 pc cobalt show perlitic structure, and the mechanical properties are only slightly affected by the cobalt.

Lead, silver, or bismuth are only partially miscible with cobalt when molten, and no definite

alloys appear to be formed.

With silicon in the electric furnace, steel-grey metallic-looking silicides are formed, Co2S1 and CoSi2. Both are stable and hard, and attacked by hydrofluoric acid and aqua regia. A very hard magnetic bonde is prepared in a similar manner; and by the action of phosphorus on cobalt chloride at a low red heat a hard brittle phosphide, Co, P3, not affected by heat or acids, is produced.

It is possible to combine cobalt or nickel with manganese with the production of valuable alloys. For this purpose commercial manganese or ferro-manganese may be used, the presence of iron not being injurious From 2 to 5 pc. of the manganese is added in small portions to the molten metal, much gas is evolved after each addition, and the metal is poured into moulds when tranquil (v. Hunting-

don, J. Soc Chem. Ind 1882, 258)

It is stated that when in the molten condition cobalt and nickel take up both carbon and oxygen, but evolve the greater part of the latter substance on cooling, leaving a porous metal containing carbon A method has been patented whereby this carbon is said to be eliminated. The porous cubes of metal are immersed in a 4 p.c solution of alkaline manganate or permanganate, dried and melted in a crucible at a high temperature The carbon is said to be thus oxidised and removed, and the metal takes up a small quantity of oxygen As soon as the metal flows easily, a little black flux and charcoal. or a small quantity of aluminium, or the calcium zinc produced in Caron's process, is added, and the oxygen thus removed (Dingl. poly J. 1884, 254-315).

Oxide of cobalt is used for the preparation of the salts of cobalt and of smalt, and by enamellers and porcelain manufacturers for the production of the finest blue glaze and colour on porcelain, glass, and other vitrifiable substances. The presence of  $z_0 \bar{b}_{00}$  of this substance imparts a bluish tinge to clear glass.

The presence of other oxides has an injurious effect on the colour produced by this substance (v Smalt), it is therefore necessary, for the more delicate work, to ensure its complete freedom from such impurities.

When heated strongly with magnesia, it pioduces a pink mass, with alumina, a fine blue (Thenard's blue); and with zinc oxide, a green (Rinmann's green) Zaffre consists of a very impute oxide of cobalt, produced by roasting the ore. It is usually mixed with 2 or 3 parts of fine sand or ground quartz, and is used for the same purpose as smalt for the coarser work.

Cobalt blue, Cobalt ultramarine, Thenard's blue, Azure blue. The composition of this pigment varies considerably, according to the It consists sometimes method of manufacture of the oxides of cobalt and aluminium, at other times it contains phosphate or aisenate of cobalt

It may be prepared by first precipitating a solution of a cobalt salt, usually the nitrate, free from iron and nickel, with potassium or sodium phosphate, or with sodium arsenate gelatinous violet precipitate is thoroughly washed and well mixed with 3-5 volumes of freshly precipitated, well-washed alumina (when cobalt arsenate is employed, a larger proportion of alumina may be added), precipitated from a solution of alum free from iron by the addition of sodium carbonate The mixture is dried until it becomes brittle, and calcined at a cherryred heat for 30 minutes in a well-covered clay crucible When the desired blue colour has been developed, the mass is ground with water and dried In order to prevent the possibility of entrance of reducing gases, which much injure the value of the product, Regnault recommended the addition of a little mercuric oxide This becomes decomposed before the ignition and evolves oxygen, which effectually prevents reduction, while the mercury escapes as vapour (Regnault, Cours Elém de Chimie, 3, 150)

A similar but less fine colour is produced by the simultaneous precipitation of the oxides of cobalt and aluminium by the addition of sodium carbonate to a mixture of cobalt nitrate and alum, the mixed precipitate is washed and

treated as above

According to the process recommended by Binder (Technologiste, 5, 55), the oxide of cobalt is precipitated by the addition of the requisite quantity of ammonia to a solution of pure cobalt chloride The washed precipitate is mixed with alumina as before, dried, and calcined at a red heat in a clay crucible for two hours

According to Hedvall the fusion of cobaltous oxide and alumina is best made in presence of potassium chloride as a flux When the cobalt oxide is in excess the resulting product can be obtained pure by treatment with water to remove the flux and with hydrochloric acid to remove the excess of cobalt oxide At 1100° the blue aluminate  ${\rm CoO\cdot Al_2O_3}$  is formed, at temperatures above 1100° a green aluminate 4CoO,3Al<sub>2</sub>O<sub>3</sub> is produced. McEachern has pointed out that alumina and cobalt oxide alone do not produce a satisfactory colour, but that additions of sodium or magnesium salts greatly improve it. Wagner and Vanino found

that cobalt sulphate could be used instead of phosphate, and that the alumina was most suitably introduced as ammonium-alum To 25 pts. of alum were added 25 to 5 pts cobalt sulphate, 03 pt. magnesium or zine sulphate; the mixture is melted in its water of crystallisation and constantly stirred until dry, when it is heated to the required temperature (Vanino, Chem Zeit 1911, 35, 497).

Thenard's blue is of a fine ultramarine colour; the presence of excess of cobalt imparts a somewhat greenish tinge. It has, however, the disadvantage of appearing violet by gaslight It is one of the most permanent blue pigments, being unaltered by acids or alkalis, and is largely used as an oil and water colour, but works better as water colour. It is non-poisonous, and has the advantage of miscibility with other pigments without alteration.

Correleum, Coline, Bleu céleste, is a corresponding colour containing oxide of tin and usually calcium sulphate—Its general composition is said to be oxide of tin, 49 66; oxide of cobalt, 18 66; calcium sulphate and silica, 31 68, and a pigment of this nature is obtained by precipitating a solution of cobalt intrate with sodium stannate, and heating the precipitate as

above.

It is a fine light-blue pigment of slightly greenish tinge, of greater density than Thenard's blue, and retains its colour in aitificial light

New blue. A blue pigment of shade varying from a pale-greenish blue to a deep-turquoise blue, largely used for enamels, consists of aluminates of cobalt and chromium produced by the action of alum on carbonates and hydroxides of cobalt and chromium

Cobalt green, Rinmann's green. The constitution of this pigment corresponds with that of Thenard's blue, with the substitution of zinc oxide for alumina

It may be prepared by mixing a solution of pure cobalt sulphate (or nitrate) into a paste with zinc oxide, and heating in a muffle furnace for 3 or 4 hours, until the desired colour is obtained. Pigments of a more uniform nature are produced by the precipitation of a mixture of solutions of zinc and cobalt salts, by means of sodium carbonate, phosphate, or arsenate, drying and heating the precipitate. The phosphate is said to give a purer and brighter green than the carbonate

R. Wagner (Technologiste, 18, 409) gives the composition of fine green pigments as varying from 71.68 p.c. zinc oxide and 11.62 p.c. cobalt oxide, to 88.04 p.c zinc oxide and 11.52 p.c. cobalt oxide, the quantity of phosphoric oxide also varying greatly

According to Hedvall (Arkw Kem Min Geol

According to Hedvall (Arkw Kem Min Geol 1913, 5, No. 6, 1), Rinmann's green is not a definite substance, but rather a series of solid solutions of its component oxides in isodimorphic

crystals.

Rinmann's green is a fine permanent pigment; its colour varies through many shades, according to the proportion of zinc present, the darker greens containing less of that substance It is non-poisonous and unacted upon by dilute acids or alkalis, but is attacked by ammonia. It has also the advantage of not affecting or being unaffected by other pigments. Rinmann's green, on account of its high price, and

the number of other permanent greens, finds little use as a pigment. Another permanent green pigment used in porcelain painting is said to be prepared by the calcination of a dried mixture of the following proportions of the freshly precipitated substances; 20 cobalt carbonate, 40 alumina, 20 chromium oxide

Cobalt bronze is a phosphate of cobalt and ammonia, of a violet colour, with a bronze-like

metallic lustre

Cobalt silicates. A cobalt silicate is produced by the addition of sodium silicate (soluble glass) to a cobalt salt—It produces a blue pigment well suited for painting on glass and porcelain

Smalt. Bleu d'azur, Bleu de Saxe This important substance consists of a silicate of cobalt

and potash

In the preparation of smalt in Saxony, smaltine is principally used. In Norway and Sweden the principal ore is cobaltine. The general process is identical for both ores.

The selected one is powdered, freed from the lighter earthy impurities by washing, and placed, in charges of about 3 cwt, in a layer of 5 or 6 inches thick on the bed of a reverberatory furnace or in a muffle. It is then roasted, and the sulphur and arsenic expelled as oxides, the latter being condensed in chambers through which the fumes are passed The roasting must be discontinued when only sufficient arsenic is left to combine with the less oxidisable metals, especially copper and nickel, while the greater part of the cobalt is converted into oxide order to ascertain whether the roasting has proceeded sufficiently, a small portion is occasionally tested and the process stopped when the The material finest tinted glass is produced. is then ready for fusion for 'smalt'

The fluxes used consist of powdered quartz and potassium carbonate. They must be of great purity, as the presence of lime and many other substances impairs the beauty of the colour. The quartz is first heated to redness, plunged into water, and, thus disintegrated, is powdered in a mill. It is then suspended in water and allowed to subside for a short time, and the liquid, which contains most of the iron and other impurities still suspended, is poured off; this operation is repeated until a pure quartz

powder is obtained

The proportions of the ingredients used vary extremely according to the richness of the ore and the colour required. The potassium carbonate used generally amounts to about one-third of the weight of ore and quartz together. White arsenic is usually added to oxidise any ferroussalt, and thus prevent the injurious effect of that substance on the smalt, and to otherwise heighten the colour of the product. The exact amount of each of these substances is found by a test fusion and comparison with a smalt of the required colour.

The smalt furnace is similar to a glass oven; it usually contains eight crucibles. The pots are of very refractory clay, free from lime, and capable of holding \( \frac{3}{4} \) cwt. When charged, they are strongly heated, and the mass fuses in about 8 hours; it is frequently stirred to render it homogeneous, and to break the crust which forms on the surface. At a white heat combination occurs, the quartz and potash react with the production of fusible potassium

COBALT. · 299

silicate, which dissolves the cobalt oxide, forming the blue 'smalt,' while the mixed arsenides of nickel, copper, and iron, with the small quantity of cobalt arsenide which is always present, fuse and form a brittle metalliclooking speiss beneath the smalt. This usually contains about 3 p c. of cobalt and a considerable quantity of nickel, and is used for the preparation of the latter substance and of cobalt oxide. The mass is left at a white heat for some time without stirring, to allow the speiss to settle, and is ladled out with long-handled iron ladles into vessels of cold water It is thus rendered granular, brittle, and easy to pulverise When nearing the bottom of the crucible, the ladle contains both speiss and smalt, the former, being much more fusible, is poured first from the ladle, from beneath the smalt, into a niche in the side of the furnace, whence the dense fumes of arsenic, &c., which it evolves pass up the chimney

The blue glass is next powdered with gramte stampers, or ground between gramte nullstones under water, and the product passed through a series of depositing vats. In the first vat it remains for a short time only, thence it passes to a second for a somewhat longer time, to a third for about 15 minutes, to a fourth and fifth for a corresponding period, and finally to a sixth vat, where it is allowed to deposit completely. The deposit in the first vat is returned to the mill and re-ground, that in the second and third is the best marketable smalt, whilst in the fourth and fifth vats a lighter-coloured deposit, sold under the name eschel, is produced. The deposit in the last vat is not marketable, and is

usually resmelted

The marketable deposits are again washed, deposited, dried, powdered, and sifted. About three-fifths of the glass taken from the pots is thus available. The presence of other oxides than those of cobalt and potash, even in small quantities, exerts a powerful influence on the colour of the small. Baryta produces an indigotinge; sodium, calcium, and magnesium produce a reddish shade; iron, a blackish green, very prejudicial to the brighter-coloured smalts; manganese violet, nickel violet, but less intense, copper, zinc, bismuth, and antimony, dull shades.

The composition of smalt varies considerably; the silica is usually from 56-70 p.c.; potash, 12-22 p.c., and cobalt, 6-16 p.c. The following analyses by Ludwig (J. pr. Chem. 1850,

51, 129) are typical -

•			
	Norwegian deep- coloured smalt	German deep- coloured	German pale coarse
Silica Potash and soda Cobaltous oxide Alumina Ferrous oxide . Arsenious acid . Water and car- bonic acid .	70 86 21·41 6·49 0 43 0 24 trace	66·20 16·31 6·75 8·64 1·36 — 0·92	72 12 20 04 1 95 1 80 1 40 0 08 0 46
		,	

Smalt is a very permanent blue pigment, largely employed for the production of blue colours with vitreous substances and in painting, both for delicate and coarse work. It was formerly much used for colouring starch, paper, &c., but for this and many other purposes it is now almost superseded by the far less permanent artificial ultramarine.

Cobalt yellow consists of potassium cobaltimitrate, prepared by the addition of potassium intrite to a solution of a cobalt salt acidulated with acetic acid. Its composition varies with the proportions used and the strength of the solution. Erdmann and Sadtler state that when precipitated from a solution containing much acetic acid it consists of  $K_3({\rm 'o(NO_2)_6})$  with a variable amount of water

Cobalt yellow is a bright yellow crystalline powder, very free from impurities, unacted upon by cold water or cold acids, and but slowly blackened by sulphuretted hydrogen.

It forms a fine pigment for artistic purposes. When used for painting porcelain it produces the usual blue colour on baking, which, on account of the purity of the substance, is of great beauty

Salts of cobalt. The soluble salts of cobalt are prepared by solution of the oxide, carbonate, or metal in the various acids. They are generally pink and deliquescent, forming pink solutions when dilute, but when dired, or in concentrated solution, they are blue. For this reason, solutions of initiate and other salts of cobalt are used as sympathetic inks, such writing being almost invisible until brought near a fire, when the letters appear blue. They gradually reabsorb moisture and again become invisible. They are largely used for neutralising the yellow colour of porcelain, and giving a pure white body, the porous clay being for this purpose soaked in solution of cobalt salts.

When ammonia is added to a solution of a cobalt salt, oxygen is absorbed, giving rise under various conditions to many complex series of salts containing ammonia and different oxides of cobalt.

Cobaltous hydroxide is obtained by adding an alkali to a solution of a cobalt salt either as a blue or red piccipitate depending upon the proportion of alkali. The red modification is more reactive than the blue compound they are regarded by Hantzsch as 'chromo-isometic' modifications (Zeitsch. anorg. Chem. 1912, 73, 304).

Cobaltous chloride CoCl<sub>2</sub> is prepaied by dissolving the oxide or carbonate in hydrochloric acid, and evaporating the solution According to the temperature at which it crystallises, it may contain 6, 4, or 2 molecules of water of crystallisation, the latter at the higher temperatures.

Cobaltous chloride is soluble in water or alcohol A strong solution, or a weaker solution containing strong hydrochloric acid, is of a blue colour.

Cobaltous sulphate CoSO<sub>4</sub>,7H<sub>2</sub>O is prepared by solution of the oxide or carbonate in dilute sulphuric acid It crystallises in red crystals, isomorphous with magnesium sulphate, soluble in 24 parts of cold water.

Cobaltous nitrate Co(NO<sub>3</sub>)<sub>2</sub>,6H<sub>2</sub>O is a red deliquescent crystalline salt prepared by dissolving the oxide in nitric acid. It is very soluble in water, the solution being used as a

reagent in blowpipe analysis.

Combinations of cobalt, as resinate, cleate or linoleate, tungate, oleo-resinate and resinotungate, are employed as driers. (See Krauss, 8th Int. Cong Appl. Chem. 1912, 12, 127; Journ. Soc Chem Ind 1912, 31, 938.)

Cobalt Carbonyls. Cobalt forms two carbonyls the tetracarbonyl Co<sub>2</sub>(CO)<sub>8</sub>, and the

tricarbonyl Co(CO)3.

Cobalt tetracarbonyl. This compound is prepared as follows. cobalt oxalate is carefully heated until completely converted into the ovide It is then washed free from traces of chlorides which hinder the formation of the carbonyl, and dried at 120°. The oxide is placed in the apparatus described under 'carbonyls,' and reduced to cobalt by means of hydrogen, at a temperature of 300° and a pres sure of 5 atmospheres After the reduction, the hydrogen is displaced by carbon monoxide, the pressure being increased and the temperature lowered For the production of the carbonyl, a minimum pressure of 40 atmospheres and a temperature of 150° is needed, and at pressures up to 250 atmospheres and temperatures between 150° and 250°, varying quantities of the carbonyl are collected in a tube attached to the outlet of the apparatus, and immersed in a freezing mixture.

The tetracarbonyl forms orange-coloured crystals of sp gr 1.73, which melt at 51°, and

decompose above that temperature.

The molecular formula is Co<sub>2</sub>(CO)<sub>8</sub> The compound is insoluble in water, but is soluble in most organic solvents, and in nickel tetracarbonyl It is not attacked by aqueous acids but concentrated acids and the halogens decompose it, forming the corresponding salt, carbon monoxide, and hydrogen

$$Co(CO)_4 + H_2SO_4 = CoSO_4 + 4CO + H_2$$

In contact with air a deep violet-coloured compound is formed, which is most probably a hydrated basic carbonate. The crystals are best preserved in a sealed tube in an atmosphere of hydrogen or carbon monoxide.

When the tetracarbonyl is heated at 60°, one quarter of the carbon monoxide is evolved at a regularly decreasing rate, leaving a compound having the formula Co(CO)<sub>3</sub> in the form of black cobalt and carbon monoxide, no intermediate

product being formed

When the tetracarbonyl is heated at 100°, one quarter of the carbon monoxide is evolved almost instantaneously, the rest coming off very irregularly and comparatively slowly (Mond, Hirtz, and Cowap, Chem Soc Trans. 1910, 97, 798)

COBALT BLOOM. Erythrite, Hydrated co-

balt arsenate (v COBALT).

COBALT BLUE or THENARD'S BLUE v.
COBALT.

COBALT BRONZE v. COBALT.

**COBALT GLANCE.** Sulpharsenide of cobalt (v Cobaltine)

COBALT GREEN OF RINMANN'S GREEN v. COBALT.

COBALT PYRITES v. COBALT.

COBALT SPEISS v. COBALT.

COBALT ULTRAMARINE or COBALT BLUE v. COBALT

COBALT YELLOW v COBALT.
COBALTITE (or Cobalt-glance). Sulpharsenide

of cobalt, CoAsS, containing Co 25–35 p.c. with Fe 2–5 p.c It frequently occurs as well-developed crystals, fine specimens coming from Tunaberg and Håkansboda in Sweden and Skutterud in Norway; these are cubic with parallel hemihedrism and with the same characteristic form as crystals of the isomorphous mineral iron-pyrites. Gianular and compact masses also occur The colour is tin-white with a reddish tinge and a brilliant metallic lustre; the streak is greyish-black Sp gr. 6.2, hardness 5½ A bed of pure cobaltite averaging 0.2 metre in thickness occurs at Dashkessan, govt Tiflis, Caucasus In the Cobalt area of Ontario massive cobaltite and small crystals are present with the more abundant smalter. Under the name sehta the mineral, found at Kheit in Rajputana, is used by the Indian jewellers for producing a blue enamel on gold and silver.

COCAINE AND THE COCA ALKALOIDS. The coca leaves of commerce are derived from Erythroxylon coca (Lam) (Bolivian or Huanuco), E coca var novogranatense (Morris) and Etruxillense (Rusby) (Peruvian or Truxillo coca).

Coca was introduced into Ceylon in 1870, and later into India, but whilst production in Ceylon attained considerable proportions, the Indian supply was never of commercial importance Subsequently, cultivation was taken up in Java where, after considerable initial difficulties, production attained large proportions constituting about 45 p c of the world's supply, the remainder being produced in South America, Cultivation was stopped in Ceylon in 1915 by order of the Government

Coca leaves contain alkaloids of four types:—
1 Cocaines; alkyl acyl derivatives of ecgo-

Cocames; alkyl acyl derivatives of ecgonine.

2 Acvlecgonines, acyl derivatives of ecgonine

3  $\psi$ -tropeines, acyl derivatives of  $\psi$ -tropine

4 Hygrines

The first and second types can be used as commercial sources of cocaine, since on hydrolysis they yield ecgonine which by benzoylation and methylation yields cocaine. The coca leaves derived from South America contain up to 1 p c of ether-soluble alkaloids (mostly cocaine), whilst Java leaves contain up to 1 5 p c, or sometimes more ether-soluble alkaloid of which little or none is cocaine.

Estimation of total alkaloids in coca leaves.-The following process is given in the United States Pharmacopæia, 8th ed. . 10-grms. of coca leaves in No 60 powder are mixed with 50 c.c. of a mixture of chloroform 1 vol, ether 4 vols., in a stoppered Erlenmeyer flask and set aside during 10 mins. Two cc of ammonia solution, diluted with 3 c c. of water, are then added, and the flask shaken at intervals during 1 hour. small glass percolator with the lower end obstructed by a wad of cotton wool is then placed in the neck of a separating funnel containing 6 cc of N-sulphuric acid, diluted with 20 cc. of water, and the contents of the flask poured into the percolator. When all the liquid has passed, the contents of the percolator are packed in firmly by means of a glass rod, and then the flask is rinsed out with 10 c.c. of the chloroformether mixture, followed by several successive washings with 5 c.c. quantities of the same mixture, the percolation being continued with

this liquid until 50 c.c. have been used in all. The separator is now shaken during one minute, the acid liquid drawn off, and the extraction repeated twice, using each time 10 c.c. of acid (6 c.c. N-acid with 20 c.c. of water); the combined acid liquors are made distinctly alkaline with ammonia solution and the alkaloids extracted with successive portions of other (25, 20 and 15 c.c.). The other is allowed to evaporate spontaneously, and the residue dissolved in 4 c.c. of N/10 sulphuric acid, and the excess of acid determined by N/50 potassium hydroxide using cochineal or iodeosin as indicator. This process gives the amount of other-soluble alkaloids in the drug. No satisfactory process is yet available for the estimation of cocaine in the 'total alkaloids'

Estimation of Ecgonine in total alkaloids.— The total alkaloids from 15-grm. of leaves are boiled for 1 hour with 30 times their weight of dilute hydrochloric acid, the mixture is cooled, extracted with other, evaporated to dryness and the residual ecgonine hydrochloride weighed (Greshoff, Pharm Weekb 1907, 44, 961)

Cocaine C<sub>17</sub>H<sub>21</sub>O<sub>4</sub>N occurs to a variable extent in the different varieties of coca leaves

(see above)

Preparation —The coca leaves are exhausted by maceration and agitation with a mixture of aqueous sodium carbonate and petroleum, whereby the cocaine with cinnamyl cocaine, &c, passes into solution in the petroleum The latter is shaken with dilute hydrochloric acid, when cocaine hydrochloride crystallises out and is collected, pressed and dried. This crude salt is purified by solution in water, liberation of the free base by ammonia, and solution of this in alcoholic hydrogen chloride, when the hydrochloride crystallises out. The motherliquors contain the other coca alkaloids Java coca leaves are used, the coca alkaloids other than cocaine alone are obtained, and these and the residual coca alkaloids referred to above are converted into cocaine by heating with boiling hydrochloric acid and pouring into water, when trustlic and cinnamic acids separate and may be filtered off The filtrate is concentrated until ecgonine hydrochloride crystallises out The ecgonine is recovered and then benzovlated and methylated in turn to produce cocaine

The crude cocaine manufactured in Peru and imported to Europe for the preparation of cocaine salts is obtained by extracting the leaves with 0.5 p.c. sulphuic acid, making the filtrate alkaline with sodium carbonate and extracting with petroleum. The latter is again shaken out with dilute sulphuric acid and the crude alkaloid precipitated from this as a white, bulky powder by sodium carbonate. The powder is washed with water and dried by ex-

posure to air for export

Properties — Monoclinic prisms, mp 98°; at higher temperatures it sublimes with decomposition. Lavorotatory [a]p—158°; sightly soluble in cold water, more soluble in hot water, by which it is slowly hydrolysed. Soluble in alcohol, ether, benzene, light or heavy petroleum, or carbon disulphide. The aqueous solution is alkaline to litmus, has a bitter taste and produces a tingling and numbness of the tongue and lips, or, more generally, local anæsthesia of the surface to which[it is applied.

On this account it is largely employed in minor surgical operations, although in recent years it has been replaced to some extent by synthetically prepared anæsthetics, such as benzamine (β-eucaine), stovaine, novocaine, &c alkaloid also causes dilatation of the pupil of the eye, but is not so powerfully mydriatic as some of the solanaceous alkaloids. When administered internally in small doses it acts as a stimulant, and diminishes the feeling of fatigue consequent on muscular exertion Coca leaves have long been known to possess this property, and in places where the plant is abundant, a mixture of lime and coca leaves is chewed by the natives before engaging in severe excition. Larger doses of the alkaloid are poisonous

The ordinary salts of cocaine are crystalline. The hydrochloride, which is the salt used in medicine, forms white micaceous scales, possessing a characteristic lustre, or as white needles, readily soluble in water. The chromate B H<sub>2</sub>(10<sub>4</sub>, H<sub>2</sub>O is thrown down, as a pale-yellow precipitate when potassium chromate is added to an acid solution of the hydrochloride. This salt is very slightly soluble in cold water, in p. 127°. The aurichloride B HAu(1<sub>4</sub> is crystalline and spatingly soluble in water. When aqueous mercuire chloride is mixed with a solution of cocaine hydrochloride a bulky precipitate of the mercurichloride. B HCl Hg(1, 2H<sub>2</sub>O is formed this salt may be crystallised from alcohol, in p. 124°.

d-Cocaine, mp. 46°, was obtained by Liebermann and Giesel (Ber. 1890, 23, 508, 926) from coca leaves, but was probably formed by the racemisation of 1-cocaine by the action of alkalis, it has been synthesized from d-ecgonine dd-cocaine, mp. 80°, was prepared from decgonine by Willstatter and Bode (ibid. 1901, 34, 1457)

Detection —In isolating cocaine, the aqueous solution should not be heated for any length of time, and not at all in the presence of acids For the liberation of the alkaloid or alkalıs from its salts, ammonia, and not caustic alkalis, should be employed Cocaine may be detected by its characteristic action on the tongue (see above) The following reactions are useful for the detection of the alkaloid: One c c of a 3 p e solution of potassium permanganate gives a violet precipitate with 0.01 grm of the hydrochloude dissolved in two drops of water (Chem Zeit 1886, 10, 71; Amer J Pharm 1911, 83, 195-201, 265-268). The hydrochloride heated with alcoholic potash gives off an odour of methyl benzoate When moistened with nitric acid, the mixture evaporated to dryness and alcoholic potassium hydroxide added, a characteristic peppermint odour is produced.

The most important criteria of purity of cocaine hydrochloride are the mp, the permanganate test, and the ammonia test

The mp. of cocame hydrochloride varies with the rate of heating. When inserted into a bath previously heated to 195° and the temperature then slowly raised, it melts at 201°-202°; when 3 drops of N/10 solution of potassium permanganate are added to 0·1 grm. of cocame hydrochloride in 5 c.c. of water containing 3 drops of dilute sulphuric acid a purple colour is produced, which should show no diminution in shade in half an hour. If 0·1 grm. is dissolved

in 100 c.c. of water, 0.25 c.c. solution of ammonia added and the mixture set aside 15 minutes, the side of the beaker being occasionally gently rubbed with a glass rod, a crystalline deposit should be formed leaving the supernatant liquid clear (limit of amorphous alkaloids).

Reactions and Constitution —Cocaine exhibits most of the general reactions of the vegetable alkaloids When boiled with water it is gradually hydrolysed into methyl alcohol and 1-

benzoylecgonine (see below)

 $C_{17}H_{21}O_4N+H_2O=C_{16}H_{19}O_4N+CH_3OH$  (Paul, Pharm J. 1887-1888 [III] 18, 781; Einhorn, Ber. 1888, 21, 47), and the same change gradually occurs if the alkaloid remains in contact with an alkaline liquid With alkalis or mineral acids methyl alcohol and benzoylecgonino are first formed, but the latter undergoes further hydrolysis into benzoic acid and 1-ecgonine (see below)

$$C_{17}H_{21}O_4N + 2H_2O_4$$

=CH<sub>3</sub>OH+C<sub>7</sub>H<sub>6</sub>O<sub>2</sub>+C<sub>9</sub>H<sub>18</sub>O<sub>3</sub>N Natural cocaine is therefore methylbenzoyl-

1-ecgonine

i-Eegonine C<sub>9</sub>H<sub>18</sub>O<sub>3</sub>N This base does not occur as such in coca leaves, but is the ultimate basic product of hydrolysis of cocaine, cinnamyl-cocaine, the truxillines and benzoyleegonine fit crystallises from dry alcohol in monoclinic prisms with 1H<sub>2</sub>O, mp 140°, or 198°-199° (anhydrous); lævorotatory It is readily esterified by alcohols yielding a series of alkyl esters, and by acid chlorides or anhydrides, giving the acyleegonines from which the corresponding 'cocaines' are obtained by alkylation

Ecgonine is now known, principally as the result of the researches of Einhorn, Liebermann, Willstatter and their collaborators (see below), to be closely related to tropine, the ultimate basic hydrolytic product of atropine and hyosoyamine, and may be regarded as tropine-β-carboxylic acid Ecgonine and its most important derivative 1-cocaine may be represented by the following formula:—

Some of the chief papers bearing on the constitution of ecgonine are: Einhorn, Ber 1889, 22, 399, 1890, 23, 1338; 1893, 26, 324; 1894, 27, 2823; Annalen, 1894, 280, 96; Liebermann, Ber 1890, 23, 2518; 1891, 24, 606; Willstatter, *ibid* 1897, 30, 702, 1898, 31, 1546, 2500, 2655, 1901, 34, 519; Annalen 1903, 326, 79.

1-Benzoylecgonine  $C_{16}H_{19}O_4N$  is the first product of the hydrolysis of cocaine; it occurs in coca leaves from which it was isolated almost simultaneously by Skiaup (Monatsh 1885, 6, 556), and by Merck (Ber. 1885, 18, 1594). The base crystallises from water in needles, mp 86° (anhydrous 195°), and is hydrolysed by hydrochloric acid to benzoic acid and ecgonine. On methylation it furnishes cocaine.

1-Cinnamylcocaine  $C_{19}H_{23}O_4N$ . This alkaloid is now known to occur in all varieties of cocalcaves, but is found in largest proportion in the Java leaves, whence it was isolated by Giesel

(Pharm. Zeit. 1889, 34, 516). It forms needles, m.p. 121°, from benzene on addition of light petroleum Lævorotatory. On hydrolysis it furnishes cinnamic acid, methyl alcohol, and lecgonine, and is therefore methylcinnamyl-1-ecgonine. The d-isomeride has been prepared (Einhorn and Deckers, Ber. 1891, 24, 7)

Truxillines  $C_{38}H_{46}O_8N_2$ . Two of these,  $\alpha$ - and  $\beta$ -, occur in coca leaves, the former is methylatruxillyl-1-ecgonine, and the second the corresponding  $\beta$ -truxillyl compound (Hesse, Pharm. Zeit 1887, 32, 407, 668; Liebermann, Ber. 1888, 21, 2342; 1889, 22, 130, 680) They are amorphous, and, according to von Liebreich, are heart poisons, and exert no local anæsthetic action

Tropacocaine (Benzoyl  $\psi$ -Tropeine)  $C_{15}H_{19}ON$  was isolated by Giesel (Ber 1891, 24, 2336) from Java coca leaves. It crystallises from ether, has mp 49°, is optically inactive, distils unchanged. The salts are crystalline, the hydrochloride melting at 283° and the aurichloride at 208°. On hydrolysis tropacocaine furnishes benzoic acid and  $\psi$ -tropine, and it may be prepared synthetically by the benzoylation of  $\psi$ -tropine (Liebermann, ibid 1891, 24, 2336, 2587, 1892, 25, 927), the latter is a physical isomeride of tropine, and has been synthesised by Willstatter (Ber 1896, 29, 936; 1901, 34, 129, 3163), who assigns to it the same structural formula as to tropine, and states that since tropine and pseudotropine are optically inactive the isomerism appears to be of the cis-trans type, a view subsequently confirmed by Barroweliff and Tutin (Chem Soc Trans 1909, 95, 1970). Tropacocaine, like the pseudotropeines as a class, is almost devoid of mydriatic action, but like cocaine is a local anæsthetic, though the effect is more transitory

Hygrines.—These secondary alkaloids were discovered by Lossen in Peruvian coca leaves (Annalen, 1862, 121, 374; 1865, 133, 352), and mvestigated by Liebermann and collaborators (Ber 1889, 22, 675; 1891, 24, 407, 1893, 26, 851; 1895, 28, 578; 1896, 29, 2050; 1897, 30, 1113), who showed that Lossen's supposed homogeneous material contained two liquid alkaloids, one boiling at  $193^{\circ}-195^{\circ}$  and having the formula  $C_8H_{16}ON$ , which makes it isomeric with tropine, and a second having the composition  $C_{14}H_{24}ON_2$ , bp.  $215^{\circ}$ , under 50 mm. pressure The low boiling alkaloid is that now known as hygrine Willstatter and Ettlinger's synthesis of hygrine acid shows that hygrine probably has the following constitution (Annalen, 1903, 326, 91):—

$$\begin{array}{c} \text{Me N} \\ \begin{array}{c} \text{CH}_2 \\ \\ \text{CH(CH}_2 \text{ CO Me) CH}_2 \end{array} \end{array}$$

From 'cusco' leaves Liebermann and Cybulski isolated a third substance of this type, viz. cuscohygrine  $C_{17}H_{24}ON_2$ , b p. 185° under 32 mm. pressure, which forms a crystalline hydrate B,3 $\frac{1}{2}H_2O$ , m p 40°. It probably has the constitution (L a C, Ber. 1895, 28, 578)—

COCAETHYTIN. Trade name for the ethyl ester of benzoylegonine.

COCATANNIC ACID v. TANNINS.

COCCINE v. AZO- COLOURING MATTERS.

α-COCCINIC ACID, COCCININ, v. COCHINEAL.

COCCULIN v. PICROTOXIN. COCCULUS INDICUS or INDIAN BERRY 18

the fruit of the Anamrta paniculata (Colebar), [A Cocculus (Wight and Arnott)] nat ord Menispermaceæ, a tree growing upon the coasts of Malabar, Ceylon, &c The berry is about the size of a large pea and of a dark-grey colour It possesses strong poisonous and narcotic qualities from the fact of its containing about one-fiftieth part of its weight of piciotoxin, and has been employed to increase the bitterness as also the mebriating effect of beer It is, however, exceedingly deleterious to health, and the use of it for this purpose has been prohibited by the Legislature under heavy penalties

Besides picrotoxin, Schmidt and Lowenhardt (Bull Soc chim 14, 817) obtained from the seeds cocculin C19 H26O10, crystallising in white needles, sparingly soluble in hot water, insoluble in cold water, alcohol and ether. It is probably identical with Barth and Kretschy's anamutin C19H21O10 which they regarded as a constituent of piero-

Cocculus indicus berries contain menispermine, paramenispermine, and picrotoxin (v Picrotoxin).

COCCUS CACTI, C. ILICIS, C. LACCAL, v. COCHINEAL.

COCHIN CHINA WAX v WAXES.

COCHINEAL. This important natural dyestuff, which, in its native country, Mexico, was used as a dye and cultivated by artificial means at a remote period of history, was for a long time considered to be of vegetable origin Cochineal, however, consists of the dried body of an insect, the Coccus cacti, which lives upon a species of cactus (the Nopalea coccinellifera (S -Dyck) or Nopal), a plant which is found in the wild condition, but which, for the sake of the insect, is cultivated in gardens which are termed Nopaleries. The collection of the insects takes place before the commencement of the rainy season, and they are then brushed either into straw baskets or into basins of tinned iron A number of insects are left upon each plant, and a new generation is produced, which is again gathered at a suitable period The insects are killed by immersion in boiling water, or are enclosed in a linen bag and placed in an oven; by the latter process the peculiar white down covering the insect is preserved, but in the former case is lost.

In Mexico and Central America two varieties of cochineal are known-the home-grown, or fine cochineal (grana fina), and the wild, or forest cochineal (grana silvestra). The former is more valuable than the latter, and is richer in colouring Since 1830 the cultivation of cochineal was introduced into Spain and the Canary Islands, Algeria, and Java, but the most productive of these newer plantations were those Since the discovery of the coal-tar of Java colours, the consumption of cochineal as a dyestuff has gradually decreased, and at the present time it is only employed in a minor degree. According to Liebermann, cochineal contains about 10 per cent. of colouring matter (Ber. 18, 19).

Carminic acid, the colouring matter of cochineal, was first isolated by Pelletier and Caven-

tou (Ann. Chim Phys (2), 8, 250), and was subsequently examined by Preisser (Annalen, 52, 375; J. Pharm. Chim (3), 5, 191) and Arppe (Annalen, 55, 101); but Warren de la Rue (ibid 64, 1) was the first to isolate this substance in a pure condition, and described it as a purplebrown mass, which, on grinding, yields a bright-red powder, easily soluble in water and alcohol, but not in ether. The mean of his analyses gave 54 13 p.c. of carbon, 4 62 p c of hydrogen, and 41 25 pc of oxygen (by difference), and from these figures he deduced the formula  $C_{14}H_7O_8$  or  $(C_{14}H_7O_8)_2$  Schutzenberger (Ann. Chim Phys (3) 54,

52), on the other hand, was the first chemist who succeeded in obtaining carminic acid in a crystalline condition. He precipitated the colouring matter, from an aqueous cochineal extract, in the form of its lead compound, suspended this in water, and decomposed it with sulphuretted The carminic acid thus liberated dissolved in the water and was recovered from this solution by evaporation at a low temperature. The product, dissolved in alcohol, was treated with ether to precipitate certain impurities, and the liquid partially evaporated, when, on cooling, a crystalline mass was obtained, which Schutzenberger considered to consist of two substances, carminic acid,  $C_9H_8O_8$ , and oxycarminic acid,  $C_9H_8O_8$ , the latter being distinguished by its solubility in ether

Schaller (J 1864, 410), who prepared carminic acid by the same method, assigned to it,

however, the formula CoH,O,

The work of Hlasiwetz and Grabowski (Annalen, 141, 329) indicated that carminic acid was a glucoside which could be decomposed into a sugar and a new colouring matter, carmine red-

$$\begin{array}{c} {\rm C_{17}H_{18}O_{10}} + 2{\rm H_2O} = {\rm C_{11}H_{12}O_7} + {\rm C_6H_{10}O_8} \\ {\rm Carmine\ acid} \end{array}$$

but according to Liebermann (Ber 18, 1969; Will and Leymann, ibid. 18, 318; and Von Miller and Rohde, ibid. 26, 2647), this is incorrect

Coccinin, according to Hlasiwetz and Grabowski, is produced when carminic acid is fused with caustic potash It crystallises from alcohol in straw-yellow needles or leaflets, dissolves in alkalis with a yellow colour, which, by air oxidation, develops first a green, then violet, and, finally, a purple tint. The analyses of this substance were in agreement with the formula C14H12O5.

Ruficoccin—By heating carminic acid with sulphuric acid to 130°-140°C, Liebermann and van Dorp (Annalen, 163, 105) obtained a new colouring matter ruficoccin, C<sub>16</sub>H<sub>10</sub>O<sub>6</sub>, and this consisted of a bright-red powder, sparingly soluble in hot water and ether, with a greenishvellow fluorescence. On distillation with zincdust, it yielded a colourless crystalline hydrocarbon,  $C_{16}H_{12}$ , melting-point 183°-188°C, from which, by oxidation, a quinone melting at 250° C. could be produced.

Furth, somewhat later (Ber. 16, 2169), prepared the same hydrocarbon by the distillation of both cochineal carmine and coccinin with zinc-dust.

Ruficarmine, C<sub>16</sub>H<sub>12</sub>O<sub>6</sub>, can be obtained, according to Liebermann and van Dorp, by heating carminic acid with water in a sealed tube at 200°C. It consists of a carmine red

powder, easily soluble in alcohol

In view of the uncertainty existing as to the percentage composition of carminic acid, Schunck and Marschlewski (Ber 27, 2980) submitted this substance to an elaborate process of purification, and, using in their operations as low a temperature as possible, obtained a product which crystallised from alcohol in red prismatic needles. Their analyses agreed closely with that required by the formula  $C_{11}H_{12}O_6$ , and the percentage composition approximately with the figures given by Warren de la Rue, and also by Schutzenberger

On the other hand, analyses by Miller and Rohde (Ber 30, 1762) pointed to the formula  $C_{12}H_{11}O_7$  or  $(C_{12}H_{11}O_7)_2$ , but according to the more recent work of Liebermann, Horing and Wiedermann (Ber 1900, 33, 149), it now appears that the correct formula for carminic acid is

 $C_{22}H_{22}O_{13}$ The most simple method of purification of Rohde. A solution of the crude colouring matter in five times its weight of water is diluted with four times its volume of acetic acid. filtered liquid, on standing over sulphuric acid, gradually deposits the carminic acid in a crystalline condition

Carminic acid crystallises in red prisms, easily soluble in water and alcohol, with a purple-red colour. It possesses no melting-point, but darkens at 130°, and at 250° becomes quite black.

By the action of alcoholic potassium acetate, carminic acid (Perkin and Wilson, Chem Soc Trans 1903, 83, 139) yields two potassium salts,

Monopotassium carminate C22H21O13K, which is red coloured; and

Dipotassium carminate  $C_{22}H_{20}O_{13}K_2$ , soluble in water with a violet-red colouration

Hexabenzoylcarminic acid C22H16O13(C7H5O)6, obtained by digesting carminic acid with benzoyl chloride (Liebermann, Horing and Wiedermann), is an orange-coloured powder, easily soluble in benzene

Octacetylearminic acid  $\rm C_{22}H_{14}O_{13}(C_2H_3O)_8$ , crystallises in golden-yellow needles, meltingpoint 155°-165°, and is readily prepared by the action of acetic anhydride in presence of zinc chloride or sulphuric acid on carminic acid (Miller and Ronde)

Constitution of Carminic Acid —Nitrococcussic acid was obtained by W de la Rue from carminic acid by the prolonged action of boiling nitric acid It was subsequently studied by v Kostanecki and Niementowski (Ber 18, 250), and was found to be identical with the trinitrocresotime acid, of the following constitution :-

When carminic acid dissolved in 50 pc acetic acid is treated with an excess of bromine, and the solution digested at the boiling heat, two substances, known as  $\alpha$ - and  $\beta$ -bromearmines, are produced (Will and Leymann, Ber. 18, 3180).

ingly soluble substance, crystallises in colourless needles, and melts at  $247^{\circ}-248^{\circ}$ , with decomposition. When oxidised with potassium permanganate in alkaline solution, it gives dibrommethylhydroxyaldehydobenzoic acid-

and dibrommethylhydroxyphthalic anhydride-

$$\begin{array}{c}
\operatorname{Br} & \operatorname{CH_3} \\
\operatorname{OH} & \operatorname{CO} \\
\operatorname{OH} & \operatorname{CO}
\end{array}$$

On treatment with hot caustic soda solution. a-bromearmine yields, in addition to a purplered colouring matter, dibrommethylhydroxyphthalic acid, and bromoform (Miller and Rohde). As a result of this reaction, these authors assigned to a-bromcarmine the constitution of a methylhydroxytetrabromdiketohydrindene-

for Zincke (Ber 20, 3227, 21, 2388) had previously shown that dibromdiketohydrindene itself-

under similar treatment yields both phthalic acid and biomoform

B-Bromcarmine C11H5B13O4 is obtained as a yellow amorphous powder, easily soluble in alcohol (Will and Leymann), and is best purified by means of its potassium salt Crystallised from acetone, it separates in orange needles, melting at 232° (Will and Leymann) or 288° (Miller and Rohde). By the action of bromine in the presence of 50 p c. acetic acid solution, it is converted into a-bromcarmine. As a result of their investigation, Miller and Rohde ascribed to  $\beta$ -bromcarmine the constitution of a methyldihydroxytribrom-a-naphthaquinone, possessing one of the following formulæ

This suggestion was supported by the fact that bromoxynaphthaquinone (1), on treatment with bromine and caustic soda solution, is converted into dibromdiketohydrindene (2) (Zincke, loc. cit)-

Additional support for this view was obtained a-Bromcarmine C<sub>10</sub>H<sub>4</sub>Br<sub>4</sub>O<sub>3</sub>, the more spar- by a study of the behaviour of \$\beta\$-bromcarmine

with zinc-dust in alcoholic solution. Thus the acetyl compound of the reduction product, melting-point 206°, gave, on analysis, numbers agreeing with those of the acetyl derivative of methyldibromdihydroxynaphthahydroquinone, possessing the following formula:

From a consideration of the points above enumerated, Miller and Rohde considered that the constitution of carminic acid could be represented by one or other of the following expressions :-

As, however, such formulæ require C=64 7 p c, H=3.92 p c, figures which are much higher than those given by the analysis of carminic acid itself, these authors suggested the addition of two molecules of water of hydration, as shown below-

A substance of this constitution would require C=55 p c , and H=5 p c Somewhat later, Liebermann and Voswinkel

(Ber 30, 688) studied the oxidation of carminic acid with alkaline potassium permanganate at the ordinary temperature, and in this way

succeeded in producing two important acids.

Cochenillic acid, C<sub>10</sub>H<sub>8</sub>O<sub>7</sub>, crystallises in colourless needles, which melt at 224°-225° with evolution of CO. It is tribasic, and at 260° is converted into hydroxymethylphthalic anhydride (1). When heated with water in a sealed tube at 210° it yields symmetrical cresotinic acid (2)-

and in the same manner at a lower temperature, 170°, gives a-coccinic acid or m-hydroxyuvitic acid (3) The constitution of cochemille acid is therefore as follows:-

VOL II -T.

a-Coccinic acid CoH &Os, the second product of the oxidation, which, as already indicated, can also be prepared from cochemilic acid, proved to be identical with the hydroxyuvitinic acid of Oppenheim and Pfaff (Ber. 7, 929) It consists of colourless needles, melting-point 239°.

Liebermann (ibid 30, 1731), whilst agreeing with the diketohydrindene constitution which had been assigned to α-bromcarmine by Miller and Rohde, considered that \$\beta\$-bromcarmine was an indone rather than a naphthoquinone derivative, and could be better represented as follows .---

It was probable, indeed, that carminic acid itself was a hydrindene or bishydrindene derivative, and the following constitutions were at the time suggested for it-

In a subsequent paper, however, Liebermann and Voswinkel (Ber. 37, 3344) consider that carminic acid is possibly a tetrahydrate of adimethyldihydroxynaphthacenequinonedicarboxylic acid-

$$\begin{array}{c|c} \operatorname{CH_3} & \operatorname{CH_2} & \operatorname{COOH} \\ \operatorname{COOH} & \operatorname{CH} & \operatorname{CH_2} & \operatorname{CH_3} \\ \operatorname{OH} & \operatorname{OH_2} & \operatorname{OH_2} \end{array}$$

and it was observed that the dimethyltetrahydroxynaphthacenequinone-

prepared by these authors not only possessed weak tinctorial property of a cochineal-like character, but in several respects closely resembled carminic acid itself

Rohde and Dorfmuller (Ber. 1910, 33, 1363) further examined & bromo-carmine, and obtained results which support the naphthoquinone constitution assigned to it by Miller and Rohde (loc cit.), but disprove Liebermann's contention that it is a derivative of indone By reduction with zinc-dust and acetic acid and subsequent acetylation,  $\beta$ -bromo-carmine gives the compound  $C_{17}H_{14}O_8Br_2$ —

colourless needles, m.p. 208°C.

Simultaneous hydrolysis and oxidation converts this into the substance

orange prisms, m.p.  $258^{\circ}\mathrm{C}$ , and it thus appears that by the latter treatment an hydroxyl has entered the quinone nucleus. The diacetyl derivative melts at  $233^{\circ}\mathrm{C}$ . When distilled with zinc-dust, this product, and also  $\beta$ -bromo-carmine itself, give naphthalene.

A valuable contribution to the subject was made by Dimroth (Ber. 1909, 42, 1611), who studied the oxidation of carminic acid with potassium permanganate at 0° in presence of sulphuric acid. The solution thus obtained gave nothing to ether, but on heating for three-quarters of an hour at 90°, it evolved carbon dioxide, and ether then extracted carminazarin

Carminazarin crystallises from water in gainet-red needles, decomposing at 240°-250°. It possesses the constitution (1), and is very similar to isonaphthazarin (2)—

Its alkaline solution when treated with a stream of oxygen, is quickly decolourised with formation of 5.6-dicarboxy-4-hydroxy-o-lolylgly-oxylic acid—

A further point of resemblance of caiminazarine to isonaphthazarine is shown by its behaviour with nitric acid in glacial acetic acid, for whereas the latter gives tetraketotetrahydronaphthalene, the former yields the analogous Carminizarinquinone crystallising in colourless prisms—

and which, when heated with water or acetic acid, passes back to carminazarin

The intermediate product formed by the oxidation of carminic acid with permanganate at 0° insoluble in ether, and which, on heating, is transformed into carminazarin, is termed by

Dimroth carminoquinone. The constitutions assigned to this substance (1) and tentatively to carminic acid (2) are given below—

Carminic acid was, therefore, not a symmetrical compound, and the nature of the group  $C_{10}H_{15}O_7$  was not then determined. At the time, Dimroth considered that the coccurum (see above) of Hlasiwetz and Grabowski, and which is prepared by fusing carminic acid with potassium hydrate, had probably the constitution of a tetrahydroxymethylnaphthalenc—

Later, Dimroth (Annalen, 1913, 399, 1) reexamined this product in detail, preparing it by fusing carmine acid with caustic potash at  $170^\circ$ – $200^\circ$ C. To it he gave the formula  $C_{17}H_{14}O_{6}$ , and described the pale yellow crystalline *tetra-acetyl derivative*,  $C_{17}H_{10}O_{6}(CH_{3}CO)_{4}$ , m.p.  $242^\circ$ – $244^\circ$ C.

When coccinin was oxidised by means of air, or oxygen, in alkaline solution (6 p c NaOH), the colour changes above described occurred, and when the pure violet colour had been obtained, acidification with hydrochloric acid yielded a substance coccinone, C<sub>17</sub>H<sub>12</sub>O<sub>7</sub>, which forms dark brown glistening crystals and decomposes at 250°C; it yields a tri-acetyl derivative, C<sub>17</sub>H<sub>19</sub>O<sub>7</sub> (CH<sub>3</sub>CO)<sub>3</sub>, orange-red crystals, m.p. 210°C, and also forms three different barium salts (one of which has a composition analogous to the sodium hydrogen salt of 2.6-dihydroxyles-methyl-a-naphtho-quinone-3:5-dicarboxyles-dickled coccinines in the content of the other hand, alkaline oxidation by means of hydrogen peroxide below 20°C, gives rise to two products, (1) cochenilic acid, and (n) an unexamined acid

Dimroth now formed the opinion that coccinin and coccinone are derivatives of anthranol and anthraquinone respectively, and the position of one of the hydroxy, methyl, and carboxyl groups in coccinone is made clear by the production from it of cochenillic acid. To coccinone Dimroth ascribes the structure—

and to coccinin either the structure-

or

When heated with water at 200°—or dilute sulphuric acid at 170°C.—coccinone loses carbon dioxide, yielding decarboxy-coccinone to which the structure-

is given, it forms red-brown crystals, and dissolves in alkalis to form purple red solutions and in concentrated sulphuric acid with a blue colour which becomes violet on addition of boric acid

Not only coccinin, but carminic acid itself, has been further examined by Dimroth (loc cit.), and as a result he considers that this substance is also a derivative of anthraquinone He has oxidised carminic acid by means of hydrogen peroxide in aqueous caustic soda, using cobalt sulphate as catalyst, and in this way obtained in the first instance carminoquinone, but the reaction proceeded further with the formation, after acidification with 80 pc acetic acid, of a yellow crystalline compound, C<sub>26</sub>H<sub>13</sub>O<sub>16</sub>Na<sub>3</sub>, 5H<sub>2</sub>O, which, when triturated at 0°C. with dilute hydrochloric acid, yielded 2 · 6-dihydroxy-8-methyl-a-naphthoquinone-3 5-dicarboxylic acid. pale yellow, hygroscopic crystals-

trisodium salt. C<sub>13</sub>H<sub>5</sub>O<sub>8</sub>Na<sub>3</sub>,4H<sub>2</sub>O, orange needles

The orientation of this acid has been established by Dimroth by comparison of its colour reactions with those of 2 6-dihydroxy-anaphthoquinone, synthetically prepared by Dimroth and Kerkovius, as also by its conversion into carminazarin by treatment with acid permanganate Moreover, the structure previously assigned by Dimroth to carminazarin, viz 2:3:6-trihydroxy-8-methyl-a-naphthoquinone-

5-carboxylic acid, has been supported by conversion of carminazarin-quinone—the oxidation product of carminazarin—into a diphenazin,  $C_{24}H_{14}O_3N_4$ , by treatment with an alcoholic solution of o-phenylene-diamine. The product

crystallises in yellow needles, and yields an acetyl derivative,  $C_{26}H_{16}O_4N_4$  When 2 · 6-dihydroxy-8-methyl- $\alpha$ -naphthoquinone-3 : 5-dicarboxylic acid—the oxidation product of carminic acid referred to above—is warmed with water, carbon-dioxide is eliminated with the production of 2: 6-dihydroxy-8-methyla-naphthoquinone-5-carboxylic acid: brown-yel-

low needles-

potassium salt, C12H2O6K, lemon-yellow crystals; dipotassium derivative, C12H6O6K2, orange-red crystals.

This compound when brominated in glacial acetic acid at 40° yields a monobrom derivative, yellow needles, m p. 240°-244°.

which on treatment with hydrobiomic acid yields a-bromo-carmin, whilst with bromine in cold methyl alcohol, the product is \$\beta\$-bromocarmin (Will and Leymann), which proves that this body has the structure

assigned to it by Miller and Rohde.

Beyond the above-mentioned decomposition products of carminic acid, Dimioth has obtained a 5 pc yield of hydrocarbons of the anthracene series by distillation with zinc-dust in an atmosphere of hydrogen After · oxidation of the mixture of hydrocarbons he isolated anthraquinone, and possibly a-methyl-anthra-

By treatment of carminic acid with boiling dilute sulphuric acid, Dimioth has also obtained a 10 pc yield of trihydroxy-methylanthraquinone carboxylic acid (C16H10O7), needles, melting-point above 300°, and this acid when heated with water at 230°-240°, passes into trihydroxy-methyl-anthraquinone by loss of carbon dioxide.

Dimroth considered the possibility that the anthraquinone nucleus is produced during the reactions described above, but concluded that this is not the case, and that it is present as such

both in carminic acid and coccinin. Dimroth formulates carminic acid thus—

$$C_{22}H_{22}O_{13} = HO \longrightarrow \begin{pmatrix} CH_3 & O \\ OH \\ OH \\ OH \\ OH \end{pmatrix}$$

On the other hand, C and H. Liebermann (Ber 1914, 47, 1213) bring forward arguments, chiefly the smallness of the yield of anthraquinone or anthracene derivatives obtained by Dimroth, in favour of the view that the anthracene nucleus is formed during the degradation reactions.

These authors have also re-examined the 'ruficoccin' of Liebermann and van Dorp (see above), and conclude that it consists of a mixture of trihydroxy-methyl-anthraquinone carboxylic acid, and trihydroxy-methyl-anthraquinone, which is confirmation of the work of Dimroth. Incidentally they described carmine anhydride,  $C_{22}H_{20}O_{12}$ , prepared by heating carmine acid with thionyl chloride—vivid red powder, resembling carminic acid, though less soluble It is reconverted into the latter by the action of aqueous alkalis.

Technical Preparations of Cochineal -Ammoniacal Cochineal - When a solution of carminic acid in ammonia is allowed to stand for some time, there is formed a new compound, which appears to consist of carminic acid in which one of the hydroxyls has been replaced by an amino group The formula assigned to this substance by Schutzenberger was C. H. NO. This reaction, long known, has been utilised for the production of a new colouring matter termed ammoniacal cochineal, or carminamide, and this comes into commerce either as cakes or in the form of a paste

The first variety is usually prepared by allowing one part of cochineal to stand in a closed vessel for three months with three parts of ammonia. The clear liquid is decanted, treated with about half its weight of gelatinous alumina, evaporated, and when the mass has become thick, it is cut up into cakes and dried

The paste is manufactured in a somewhat similar manner, the ammonia, however, being only allowed to react for eight days. The clear liquid is then evaporated to about one-third its bulk without addition of alumina (Crookes, Dyeing and Calico Printing).

Ammoniacal cochineal dyes much bluer shades than cochineal itself, and it is sometimes employed in conjunction with the latter. Its use is now very limited, but it finds some application in the blueing of bleached cotton

Dyeing Properties of Cochineal—Cochineal has been little employed in cotton dyeing, but was very largely used in silk and wool dyeing It has now been practically replaced by the azo-scarlets.

Two shades of red are obtained upon wool with cochineal, namely, crimson, which is produced by means of aluminium sulphate, and a very fiery scarlet, for which stannous and sometimes stannic chlorides are employed

For crimson, wool is mordanted with aluminium sulphate and tartar, and is then dyed

Dimroth in a separate bath with cochineal. The employment of calcium salts in the dyeing operation is not beneficial. Fairly good shades of crimson can also be obtained by mordanting and dyeing in a single bath with aluminium sulphate, oxalic acid, and cochineal For scarlet, wool can be mordanted with stannous chloride and tartar, and dyed in a separate bath with cochineal. single-bath method, employing stannous chloride, oxalic acid, and cochineal, has, however, been very largely used for this purpose On the other hand, preparations of stannic chloride, known as 'tin spirits,' 'scarlet spirits,' and 'nitrate of tin,' are and have been much employed by division of stannic chloride. ployed by dyers of cochineal scarlet. Though, when used alone, stannic (hloride does not give such brilliant shades as the stannous mordant, a mixture of both is considered to be beneficial. For very yellow shades of scarlet, yellow colouring matters can be added to the cochineal dyebath, and of these 'flavine' has been employed to a considerable extent

> Cochineal red on wool possesses considerable fastness to light, but has the defect that weak alkalis and soap cause it to acquire a duller or more bluish shade

> Wool mordanted with potassium dichromate gives with cochineal a good purple colour, whereas with ferrous sulphate and tartar, purplish, slate, or lilac colours can be produced. These mordants, however, are not employed in practice. A good erimson shade is produced by mordanting silk with alum, and dyeing with an extract of cochineal. In scarlet dyeing, silk is preferably first dyed yellow, then mordanted with 'intromuriate of tin,' and finally dyed in a second bath with the assistance of cream of tartar. Silk can also be dyed in a single bath with cochineal, stannous chloride, and oxalic acid.
>
> A G. P. & A E. E.

COCHINEAL CARMINE v. Lakes. COCHINEAL RED, SCARLET v Azo- col-OURING MATTERS.

cochineal wax and FAT. The wax is obtained by extracting the cochineal with benzene, and consists almost entirely of a substance to which Liebermann gave the name coccern. After the removal of the coccern, the fat is extracted with ether; it consists of myristin and an oil containing fatty acids. The following are the percentages of coccern in various samples of cochineal (extracted whole with benzene):

Silver cochineal . 1.0 p c

Mexican silver cochineal . 1.7 ,,

Teneriffe silver cochineal . 2.0 ,,

Zaccatille . . . 0.5 ,,

Teneriffe black cochineal . 0.7-1.0 ,,

Mexican black cochineal . 1.5 ,,

Granilla . . . 4.2 ,,

By powdering the cochineal before extraction, the percentage is increased by about one-half, except in the case of Granilla, where the granules are exceedingly small. The percentage of myristin is on the average about 15-20, and of fluid fat and fatty acids 4.0-60, so that the average total quantity of wax and fat contained in a silver cochineal is about 12 p.c. After recrystallising from benzene or acetic acid, coccern forms thin glistening plates of meltingpoint 106°. It is sparingly soluble in all cold

COCOA. 309

solvents, and nearly involuble in alcohol and ether. It has the constitution  $C_{30}H_{60}(C_{31}H_{41}O_3)_2$ . On saponification with alcoholic potash, it gives cocceryl alcohol  $C_{30}H_{60}(OH)_2$  and coccerc acid  $C_{31}H_{62}O_3$ . The alcohol forms a white crystalline powder of mp  $101^\circ-104^\circ$  The acid melts at 93°, and is soluble in hot alcohol, benzene, &c; its ethyl ether melts at about 70°. The myristin  $C_3H_5(OC_{14}H_{27}O)_3$  forms colourless crystals of mp  $55^\circ$ , easily soluble in warm alcohol, &c; on saponification it gives glycerol and myristic acid  $C_{14}H_{25}O_2$  of mp.  $54^\circ$ , and b p.  $248^\circ$  at 100 mm (J Soc Chem. Ind 4, 585).

COCHINEALIC ACID v COCHINEAL

COCINIC ACID v. Hendecatoic acids. COCOA. The cocoa tree is an evergreen shrub, indigenous to Mexico and the tropical regions of Central America, whence it was introduced into Europe by the Spaniards. It belongs to the natural order Sterculaceæ, and comprises several species, that most highly valued and cultivated being the Theobroma cacao, so called by Linnæus, from  $\theta\epsilon \delta s$  (god) and  $\beta\rho\hat{\omega}\mu\alpha$  (food), in allusion to the esteem and almost veneration in which the beverage of Čentral America, and 'cacao,' the name by which it was originally known. The seeds were

used as a kind of money even down to the time

of Humboldt's visit to Mexico.

The tree grows to a height of from 12 to 40 feet, the average being about 25 feet, but in plantations it rarely exceeds 18 feet. It flourishes best in a moist and warm climate, in latitudes between 25° north or south of the equator, and at altitudes up to 1000 feet. It is now cultivated more or less extensively in Brazil, Caraccas, Central America, Ecuador, Essiquibo, Grenada, Guayaquil, Peru, Trinidad, Venezuela, and in the West Indies (particularly Jamaica, Dominica, and St Lucia); also in Ceylon, the East Indies, the Philippines, Mauritius, Madagascar, San Thomé, Gold Coast, Cameroons, and to some extent in Australia

The cocoa tree begins to bear fruit when 3 years old, and attains its full productiveness when 7 or 8 years old, but a healthy tree will bear for 40 or 50 years. Fruit is produced all the year round, but harvesting usually takes place only twice a year—from February to April, and again in October and November.

The fruit consists of an elongated pod, in shape between a melon and a cucumber, measuring from 5 to 12 inches in length, and from 2 to 4 inches in diameter, with a thick pericarp varying in colour from yellow or red to purple, according to the variety. There are from 20 to 40 or more seeds in a pod (though the number is smaller in the West Indian varieties) arranged in 5 longitudinal rows, and embedded in a rose-coloured mucilaginous pulp fresh they are usually white (some varieties purple), and have a bitter taste, but after drying and exposure to the air and light they acquire a yellow, red, or brown colour, and become hard, brittle, and less bitter They are ovate, compressed, and about the size and shape of harroot beans or olives.

After removal from the pods, the seeds or beans are either freed at once from adhering pulp and dried in the sun (or occasionally by artificial means), to produce 'unformented'

cocoa,' or they are removed to the 'sweating' or curing house to undergo a process of fermentation before drying Unfermented cocoa is distinguished by its pale skin, which separates with difficulty from the kernel Owing to its bitter taste, it is suitable only for use in cocoa mixtures containing starch and sugar. It is quite unfit for chocolate making, and commands a lower price than fermented cocoa.

To produce fermented cocoa, the beans were formerly buried in the ground, or in trenches covered with plantam leaves, but in order to avoid the consequent adherence of earthy matter, they are now generally buried in casks or boxes, or heated in piles in sheds, and turned over once or twice daily. The temperature rises considerably during fermentation, and the process requires great care and skill, as on it the commercial value of the finished beans chiefly

depends In the fermentation—for which the optimum temperature is 35°-40°-alcoholic fermentation is succeeded on the second or third day by acetic fermentation; during the subsequent drying, oxidation of the astringent constituents The market value of the seeds varies occurs inversely as their tannin content When the seeds contain over 20 pc of moisture butyric fermentation often occurs and causes mouldiness Oxidation is most rapid at 50°-60°, but must not be allowed to proceed too far, or loss of aroma occurs For observations on the fermentation of cocoa, see Brill, J. Soc. Chem Ind 1917, 36, 1245

The objects arrived at in the 'sweating' operations are the loosening of the adherent pulp, the removal or modification of the bitter principle, the development of an agreeable aromatic odour, and the colouration of the beans to a rich mahogany tint. The fermentation process generally lasts from 2 to 7 days. In some parts, as Trinidad and Jamaica, red earth is often employed to assist in the removal of the mucilage from the beans, which are afterwards rubbed clean and dried. Occasionally Venetian red is used to impart a uniform red tint to the finished beans.

As the fermentation process produces considerable change in the composition of the nibs it has recently been replaced by treatment with a 1 p c solution of sodium carbonate for 10 minutes at 45°, whereby the beans are 'stabilised' and sterilised. Cocoa prepared in this way is perfectly stable, and contains no pre-formed sugar. Theobromine is not present in the free state, but is formed together with sugar and tannin by hydrolysis (v infra).

In the drying operation the beans are spread out in trays and systematically turned so as to expose every part of the bean to the light, and, when thoroughly air-dried, are in a suitable

condition for shipment

The preliminary processes of manufacture consist in (a) sifting out sand, dust, and small stones; (b) picking out by hand the empty beans and foreign matter, as larger stones, grass, wood, &c; (c) roasting; and (d) cleaning, by breaking up the beans and removal of the husks or shells. Berhardt (Chem. Zeit. 1889, 32) has found, from the results of actual factory work, that the losses in these operations amount to 2 or 3 pc in sifting; nearly 1 p.c. in picking; about

310 COCOA.

5 p.c. in roasting; and 12 or 13 p.c. in cleaning, making a total loss of over 20 p c.

The roasting process, which requires great care, skill, and experience on the part of the operatives entrusted with it, is conducted in rotating cylinders heated by coke fires or by means of superheated steam in pipes lining the cylinders, to a temperature varying from 500° to 600°F (160° to 315°C.) Uniformity of roasting is facilitated by grading the beans in the same operation to a uniform size. The beans are then cooled, gently crushed in a 'kibbling mill,' to fracture the crisp husk and break down the kernel into its natural angular fragments or 'nibs,' after which the nibs, husks, and hard rod-shaped germs are separated by winnowing and sieving, and the nibs graded into different classes according to size

The object of roasting is to develop to the fullest extent the characteristic aroma and flavour of the cocoa. This is probably due to the development of a volatile principle in the nature of an essential oil, somewhat analogous to the caffeol of coffee (v. infra, Coca-red)

The roasted husks or shells constitute on an average about 12 or 13 p c of the weight of the whole bean, varying from 7 or 8 p.c. in the thin-walled beans of Ceylon and Java to 18 p c in those of Trinidad and the West Indies, which The shells form a have a much thicker shell low-priced product extensively used, after grinding, in the cheaper grades of cocoa and chocolate, and occasionally, under the name of 'cocoa tea,' as a beverage which has the taste and flavour of weak cocoa Cocoa shells have proved to be a useful addition to feeding stuffs, especially for milch cows, and gave good results on the experimental farms of Canada when used as manure. (For analysis of cocoa shells, see table below)

The nibs enter into commerce as such, but although in this form the consumer may be assured of the purity of the article, yet, owing to the large proportion of fat in them (about half their weight) and the difficulty of disintegrating them even by boiling water, they are hardly suitable for popular use as a beverage.

For the manufacture of the various forms of prepared cocoa, the nibs are ground by means of millstones or steel mills so constructed as gradually to reduce the cocoa to the finest and smoothest possible condition If the cocoa is to go into consumption as ground nibs in powder, the stones or rollers are kept cool during the operation, but if, as is usual, the ground cocoa is afterwards to be made up into any of the numerous forms of prepared cocoa or chocolate, the stones or rollers are usually kept heated during the grinding. This causes the fat to melt, and the whole is reduced to a liquid mass, which is run into moulds and cooled so as to form blocks, which are sold as such or kept as stock for subsequent conversion into prepared cocoa or chocolate

The leading types of manufactured cocoa are flake cocoa, rock cocoa, and cocoa powders sold under such names as 'soluble' cocoa, and 'extract' or 'essence' of cocoa. Flake cocoa is prepared simply by grinding the whole beans or the nibs only in steel mills somewhat resembling those used for grinding coffee Rock cocoa is

made in the form of a paste by grinding in a mill with heated stones or rollers, and with the addition of sugar and starch.

The admixture of starch or other gelatinising substance tends to mask the presence of the fat and to render the cocoa more readily miscible with boiling water. These mixtures, sold under such names as 'homeopathic,' 'Iceland moss,' &c., are frequently described as 'soluble cocoa,' but this is a mismomer, since the absence of a sediment in the cup is due, not to solution of the cocoa, but to the formation of an emulsion, which holds the cocoa in a finely divided state suspended in the liquid. In some of these, as 'granulated cocoa,' 'cocoatina,' &c, sugar is added as well as starch.

The large proportion of fat (about 50 pc), being found by many to be difficult of digestion, is reduced in preparations intended for use as a beverage, not only by the addition of sugar and starch (which incidentally reduces the proportion of cocoa present), but also by the actual abstraction of fat to form the so-called 'essence' or 'extract' of cocoa

The partial extraction of the fat is accomplished by submitting the pure cocoa paste enclosed in strong canvas bags to a pressure of 1200-1400 lbs per square inch in hydraulic presses heated to 70°-80°. The cocoa butter flows out through a special orifice, and is afterwards solidified into blocks or slabs ready for use in surgical and pharmaceutical preparations, for soap making, and in the manufacture of chocolate confectionery. The commercial value of cocoa butter just before the war was twice that of the raw bean, and this no doubt was an inducement to its removal, in addition to the alleged purpose of making the cocoa more directible.

digestible
To prepare the so-called 'essence' or 'extract' of cocoa, the cakes of compressed cocoa, now containing only about 25 pc of fat, are turned out in a nearly dry condition, and ground to a fine powder Some cocoa powders are said to be rendered more digestible by treatment with alkalis, which soften the cellulose tissue and act upon the fat so as to produce a more perfect emulsion than that obtained by other 'soluble' cocoas prepared with starch; also there is less tendency to separation of the melted fat in the cocoa bever-Another method is to soften the tissue by heating in closed vessels with or without water to a temperature of 150°. In some preparations, as 'solidified cocoa,' the fat is not removed, whilst as preparation of cocoa practically free from fat is obtained by extracting the roasted and ground beans with ether or benzene ordinary flavouring materials are vanilla and cinnamon, but other spices and artificial vanilla are occasionally used.

The principal constituents of cocoa are fat ('cocoa butter'), alkaloids, albuminoids, starch, tannin, cocoa-red, and mineral matter.

The fat, cocoa butter, or cacao butter, oleum theobromæ, is the principal food constituent of cocoa, and forms from 45-55 p c. of the husked bean It has a pleasant taste and odour, suggestive of chocolate At ordinary temperatures it is a yellowish-white, somewhat brittle solid, melting below the temperature of the body, viz 28°-33°, and solidifying at 21°-23°;

COCOA. 311

sp gr. from 0.95 to 0.97 at 15°, and 0.856 at 100°; rodine number from 33 to 38; saponification value, 193-195; refractometer reading (Zeiss) at 40° is 46-478; and index of refraction, between 1.4565 and 1.4578 at 40°. It is readily soluble in ether, chloroform, turpentine, and boiling alcohol, from which it crystallises out almost completely on cooling Chemically, it is a mixture of the glycerides of stearic, palmitic, oleic, and arachidic acids (Benedikt, Analyse der Fette und Wachsarten). It is sometimes adulterated with such substances as coco-nut oil, cotton seed, and sesamé oils, stearin, tallow, and paraffin way Illipé oil is often used as a substitute for cocoa butter (which it closely resembles in certain respects) in the manufacture of chocolate

ANALYSES OF RAW COCOA NIBS.

			Bell	Bousein- gault	
Moisture Fat		•	5 23 50 44	7 6 49 9	
Starch			4 20	24	
Albuminous matte Soluble . Insoluble .	·		6 <b>30</b> 6 96	} 10 9	
Astringent princip	i le	•	$\begin{array}{c} 6 \ 71 \\ 2 \ 17 \end{array}$	02	
Celluloso .	•	•	6 40	10 6	
Alkaloids Cocoa-red	•	. :	$\begin{array}{c} 0.84 \\ 2.20 \end{array}$	3 3	
Indefinite organic (insoluble)	ma •	itter	5 80	5 3	
Ash (Tartaric acid)	:	:	2 75	4 0 3 4	
			100 00	100 0	

Alkaloids — Theobromine C<sub>5</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>N<sub>4</sub>O<sub>2</sub>, or dimethylxanthine, is the principal alkaloid in cocoa, and is closely related to the alkaloid of tea and coffee, caffeine (trimethylxanthine), which is also present to a small extent in cocoa Theobromine crystallises in white microscopic needles, having a bitter taste, and is present in cocoa to the extent of about 15 pc The published analyses show variations from 0 4 to 3 3 p.c., due to the different methods of analysis employed. One of the best methods is that of Decker, as modified by Welmans and Fromme (Pharm. Zeit 47, 789, and 858; and Apoth Zeit 18, 68) The caffein is usually included with the theobromine, from which, however, it can readily be separated by means of cold benzene, in which caffeine is soluble, but theobromine practically insoluble The amount of caffeine varies from 0 1 to 0 3 p c. (see art. Theorrowine).

According to Kreuz, the proportions of theobromine hithorto shown in published analyses represent only free theobromine. He maintains that part of the alkaloid is combined and is not extracted until the cocoa has been subjected to hydrolysis By this method Kreuz found from 2 8 to 5 1 p c. of theobromine in cocoa beans (Zeitsch. Nahr. Genussm. 1909, 17, 526-7).

Albuminoids.—These have been but little studied, but Stutzer (Zeitsch. angew. Chem

1891, 368) has classified the nitrogenous constituents of cocoa as follows:—

(1) Non-proteids, substances soluble in neutral water solution in presence of coppes hydroxide (theobromine, ammonia, and aminocompounds)

(2) Digestible albumen, insoluble in neutral water solution in presence of copper hydroxide, but soluble when treated successively with gastric juice and alkaline pancreas extract

(3) Insoluble and indigestible nitrogenous substances

According to Stutzer's analyses, the proportion of albuminoids in cocoa powder manufactured without chemicals is between 17 and 18 p.c., of which 10 p.c. is soluble or digestible albumen, the remainder (over 40 p.c. of the whole) consisting of insoluble and indigestible substances. Forster, however, maintains that the proportion of digestible albuminoids reaches as high as 80 p.c., his experiments being made on the human subject, whilst Stutzer's were conducted under laboratory conditions (Hyg. Rundschau, 1900, 314)

Starch, as in oleagmous seeds generally, is not very abundant. It has been variously estimated from 2 to 20 pc of the nib, but many of the results are too high, owing, no doubt, to the conversion of cellulose into sugar when the acid method for the starch conversion is employed. The diastase method is preferable in the presence of cellulose. The average amount of starch is from 4 to 5 pc, or about 8 pc calculated on the fat-free bean. The starch grains are nearly spherical, with a very indistinct nucleus. They have a tendency to unite in small groups of 3 or 4, and rapidly lose the colour imparted by ioding. The average size is about 0.005 mm.

Tannin —An astringent principle resembling tannin exists in the raw bean, but, according to Mitscherlich and Bell, it is rapidly oxidised to 'cocoa-red,' to which the characteristic colour of cocoa is due Bell (Foods, 1 79) states that this astringent principle is different from the tannin of tea and coffee, and that its rapid change to cocoa-red, even during the process of analysis, renders its quantitative estimation difficult. It is precipitated by basic acetate of lead, and can be obtained after removal of the lead by sulphuretted hydrogen. It gives a green precipitate with ferric chloride not unlike caffeic acid under similar treatment.

Cocoa-red is not present in the fresh seeds, which are nearly white, but arises from the oxidation of the natural tannin of the seed. It consequently appears in different quantities in different kinds of cocoa. On hydrolysis it decomposes into glucose, tannin, and a resinous substance. By many the peculiar taste and aroma of cocoa are associated with the formation of cocoa-red, which increases in amount during the process of fermentation, drying, and roasting, simultaneously with a corresponding increase in flavour, thought to be due to the development of a volatile principle or essential oil. Details of methods for its extraction and estimation are given in Zipperer (Die Schokoladen Fabrikationen) and Blyth (Foods).

tionen) and Blyth (Foods).

Mineral matter—The ash in cocoa nibs varies from 2 5 to 4 5 p.c., the average, according to Zipperer, being 3 6 p c. in raw, and 3 9 p.c. in

312 COCOA.

the roasted kernels. The shells contain two or three times this amount. Approximately half the ash (which is rich in potash and phosphates) is soluble in water. The ash is naturally lower than the average when starch or sugar has been added, and higher when fat has been abstracted, or when the cocoa has been adulterated with husks or coloured with pigments The shell is rich in alkaline carbonates and silica as compared with the nib A high alkalinity in the ash of cocoa powders indicates treatment with alkalis as described in the preparation of the so-called 'soluble' cocoas The composition of the ash of cocoa nibs and husks is shown in the table given below.

Whilst cocoa ranks higher than tea or coffee as a nutritious beverage, its value as a substitute for solid food has been greatly exaggerated. In its content of alkaloid and tannin substances, it stands midway between tea and coffee, and, like them, it should be considered a stimulating beverage rather than

a true food.

The tables on p 313 will serve to show the composition of cocoa nibs, shells, the ash, and

several kinds of commercial cocoas

Adulterants —Besides sugar and starches (which have come to be considered as essential ingredients of certain proprietary cocoa preparations), the usual adulterants are foreign fats (particularly coco-nut oil) to replace the abstracted cocoa butter, farmaceous substances, ground cocoa shells, and pigments such as ferric oxide, ferruginous earth, &c For the estimation of cacao shell, see Knapp & McLellan, Analyst, 1919, 2, which contains a bibliography on the subject

COCO-NUT OIL GROUP. Under this heading is grouped a number of vegetable fats which differ from all other vegetable fats by having high saponification values in conjunction with high Reichert-Meissl values The considerable proportion of volatile fatty acids which the fats belonging to this group contain, brings them into relationship with the milk fats of mammals. They differ essentially from the latter, however, in that they do not contain any The high proportion of volatile butyric acid fatty acids contained in the members of these groups differentiates them from other fats which are also characterised by high saponification values, but do not contain volatile acids.

The fats belonging to the coco-nut oil group are the following: (a) Muriti fat, from Mauritia vinifera (Mart); Acrocomia vinifera (Oerst);
(b) mocaya oil, from A sclerocarpa (Mart); Cocos sclerocarpa, C. aculeata (Jacq); Bactris minor (Gart); (c) cohune oil, from Attalea cohune (Mart), (d) maripa fat, from Palma (Attalea) maripa (Aubl), (e) aouara-kernel oil, from Astrocarpum vulgare (Mart); (f) palm-nut oil, from the kernels of Elons gumeensis (Linn; Jacq); (g) coco-nut oil from Cocos nucifera (Linn) and Cocus butyracea (Linn); (h) fat from Cocos acrocomordes.

It may be pointed out that aouara-kernel oil and palm-nut oil differ considerably from the corresponding fats contained in the fruit flesh surrounding the seeds (see Palm oil)

All these fats are derived from trees belonging to the palm family The two most important

latter will be described subsequently (v. PALM KERNEL OIL).

Coco-nut oil is the fat obtained from the kernels of the coco-nut, especially from those of the two species Cocos nucifera (Linn.), and C buturacea (Linn).

Cocos nucifera grows on all coasts and islands of the tropics; in fact, it may be considered a cosmopolitan of the tropical coast-lines in both hemispheres. The coco-nut tree is indigenous to the islands of the Malayan Archipelago, whence the nuts were carried by sea currents towards the east to the Pacific Islands and the coasts of ('entral America, and towards the west to Ceylon and the east coast of Africa.

Being well protected by the 'coir' (fibres), the seeds are able to germinate on all shores (islands and even coral reefs) where they may be

deposited.

The coco-nut tree forms large forests, especially on the coast-line of Ceylon, Java, and the Philippines. On account of its manifold uses, the tree has been cultivated early in the history of mankind, and the fruits were, therefore, gathered, as it were, on a commercial scale, not only in the localities named, but also in the South Sea Islands, Madagascar, Mauritius, Ceylon, Zanzıbar, the east and west coast of Africa, the Antilles, and the coast-lines of tropical South America

Cocos butyracea is indigenous to Brazil, and yields, together with Cocos nucifera (as also with the other cocos species mentioned above), the chief quantities of coco-nut fats which are consumed locally The coco-nut oil which is exported to Europe is mainly derived from

Cocos nucrfera

The coco-nut tree flowers when about 6 or 7 years old, and begins to bear fruit from an age of 8 years up to an age of 60 years. The nuts contain a fleshy endosperm, built up from a The nuts milky juice, rich in oil At the period of maturity of the fruit the milky juice has disappeared almost completely, and the endosperm forms the well-known coco-nut kernel, inside which the remainder of the milky juice, coco-nut milk, The kernel contains from 30 to is enclosed 40 pc of fat, and about 50 pc of moisture

Since the earliest times in the history of mankind, the kernels have been used by the natives of the South Sca Islands, and especially of India, as one of their chief articles of food. The mode of preparing the oil, as practised centuries ago, was of the most primitive kind The kernels were cut into small pieces and exposed in piles to the sun, when the oil would run off spontaneously Improved methods consisted in converting the kernels into pulp, placing the latter in vessels perforated with holes, and exposing them to the sun, when the oil would run through the holes. The oil was then collected and carried in bamboo sticks to the market, a practice which up till a few years ago still obtained in the outlying parts of the Philippines. In order to render the process independent of the heat of the sun, as in the rainy season, the kernels were, at a later stage of evolution laid on hurdles, formed from bamboo staves and coco-nut leaves, and dried over a slow fire, the fuel for which was derived from the shell of the kernels. In India, where fats are coco-nut oil and palm-kernel oil. The the nuts became at an early date an extensively

used food product, the nuts were split into | dried kernels were then triturated and expressed, halves by a blow with a sharp instrument, the coco-nut milk was poured off, and the halves exposed to the sun to dry, whereby the kernels became readily detachable from the shell. The used as cattle food ('coco-nut poonac') This

## COMPOSITION OF COCOA SHELLS (WEIGMANN).

Commercial varieties	Moisture	Nitro- genous sub- stances	Theo- bromine	Fat	Nitrogen- free extract	Cellulose	Ash	Sand	Total nitro- gen
Caraccas Trinidad	12·49 14·64 13 93 14·89	13·18 14·62 16·25 16·18	0 58 0 74 0 78 0 75	2·38 3 45 2 54 2 01	40·30 44 89 42 47 43·32	16·33 15·79 17 04 15 25	9 06 6·19 6 63 8·08	6 26 0 42 0 85 0 27	2·11 2·34 2·60 2·59

# ROASTED BEANS AFTER REMOVAL OF THE HUSK (HEISCH).

	Husk	Fat	Nitro- gen	Albu- minoids	Ash	Ash soluble in water	Ash soluble in HCl	H <sub>3</sub> PO <sub>4</sub> m ash	Moisture	Starch, gum cellulose, &c
Caraccas . Trinidad . Surinam . Guayaquil . Grenada . Bahia . Cuba . Para .	13 8	48 4	1.76	11·14	3.95	2·15	1.80	1.54	4 32	32·19
	15 5	49·4	1.76	11·14	2.80	0·90	1.90	0.93	3 84	32·82
	15 5	54 4	1.76	11·14	2.35	0·80	1.85	1.23	3·76	28·35
	11 · 5	49 8	2.06	13·03	2.50	1·75	1.75	1.87	4 14	30·47
	14 · 6	45 6	1.96	12·40	2.40	0 60	1.80	1.35	3 90	35·70
	9 6	50 3	1.17	7·40	2.60	0 90	1.70	1.26	4 40	35·30
	12 0	45·3	1.37	8·67	2.90	0 95	1.95	1.13	3 72	39·41
	8 5	54 0	2.00	12·66	3.05	1·40	1.65	1.00	3·96	26·33

## ANALYSES OF THE ASH OF COCOA NIBS AND HUSKS (BELL).

			1 Guayaquil nıbs	2 Surinam nibe	3 Grenada nibs	4 Finest Trinidad nibs	5 Finest Trinidad husks
Sand		•	0·15 0 46 0 46 23 35 19·18 3 24 0 10 0 21			0 65 0 83 29 30 18 23 6 51 0 08 0 10	5·12 2 87 0 44 0·94 37·89 13 04 7 30 0 55 0 63
Carbonic anhydride Sulphuric ,, Phosphoric ,,	$(SO_2)$ . $(SO_3)$ . $(P_2O_5)$	:	0 69 2·77 49 39 100·00	3·31 4·29 37·78 100 00	2 92 4·53 39 20 100 00	4 19 3 91 36 20 100 00	10 80 3 25 17 17

## The following table shows the general composition of commercial cocoas: -

#### ANALYSES OF COMMERCIAL COCOAS (BELL).

	Prepared cocoa	Iceland moss cocoa	Rock	Flake cocoa	Cocoa- tina	Choco- latine	Finest Trinidad nibs	Chocolat de Santé	Cocoa extract
Moisture Fat Starch (added) Sugar (cane) Non-fatty cocoa	4.95 24 94 19.19 23 03 27 89	5·47 16 86 24·70 29·23 23·74 100·00	2.58 22.76 17.56 32.20 24.90 100.00	5 49 28·24 none none 66 27	3 52 23 98 none none 72 50	4·40 29 60 none none 66 00	2·60 51·77 none none 45·63	1 44 22 08 2 00 61 ·21 13 27 100 ·00	5 76 29 50 none none 64·74 100 00

process was carried out with special care at Cochin and the coast of Malabar. Hence the reputation of Cochin coco-nut oil for best quality has been maintained to this day.

As large proportions of moisture left in the kernel readily lead to putrefaction, a further step in the preservation of the fat was found in drying the kernels thoroughly, so as to remove the bulk of the water. The absence of moisture rendered the growth of fungi impossible, and thus prevented excessive hydrolysis and subsequent rancidity. The process of producing the dried kernel—copra—developed into a manufacturing process.

Coconut oil became first known in Europe in the 18th century, but was not imported in any notable quantities before the year 1820. At present three qualities of oil are distinguished in commerce (1) Cochin oil; (2) Ceylon oil,

(3) Copra oil.

(1) Cochin oil represents the finest and whitest quality, and, is obtained as stated above, by boiling the kernels with water and

skimming off the oil.

(2) Ceylon oil is prepared in a similar manner in the plantations of Ceylon. This oil is not of such high quality as Cochin oil, owing mainly to the want of care on the part of the natives, as compared with that exercised by the natives of the Malabar coast. Hence Ceylon oil represents, as a rule, a second quality of coco-nut oil, and is characterised by a somewhat higher percentage of free fatty acids than is found in Cochin oil. At present, however, in some plantations in Ceylon, managed by Europeans, the same care is exercised as in Cochin, and some brands of Ceylon oil are now imported which yield little in quality to Cochin oil

(3) Copra oil is the oil prepared in modern factories in Europe, United States, Australia, from copra (i.e. the dried kernels), which are imported in enormous quantities into these countries. The conversion of the kernel into copra must not only be looked upon as a convenient method for reducing the cost of freight, but, as pointed out above, is pre-eminently a cheap process for preserving the coco-nut oil

as far as possible

In the Manila market copra oil is sold as 'Cebu sun-dried,' 'fair marketable Manila,' and 'Laguna,' but the distinction between the

different grades is not well marked

At present great care is bestowed on the preparation of copra, it is used in growing proportions for the production of edible coco-nut oil (see Edible oils), and coco-nut stearin (see CHOCOLATE FATS) In commerce we differentiate 'sun-dued copra' and 'kiln-dried copra' The sun-dried copra was considered the better quality, as it gives a white coco-nut oil, with only a slight yellow tinge, and could be readily converted into an edible oil Kiln-dried copra is obtained by a more rapid process, the kernels being dried over a bamboo grating heated by burning underneath it the shells. In these primitive kilns, the fumes obtain ready access to the copra itself, and not only burn it occasionally, but impart to the copra a peculiar empyreumatic smell and a dark colour Hence copra thus dried yields a yellowish oil which proves refractory in the refining process. At present, however, notable improvements are being

introduced in the kiln-drying process (rotary driers, vacuum driers), so that it may be expected that in the near future kiln-dried copra will take its place next to, or rank equally with, sun-dried copra. Sun-dried copra contains on an average 50 pc of fat; in kiln-dried copra the percentage rises to 63 and even 65 pc. Copra dried by means of hot air contains even as much as 74 p.c.

Its acidity (as oleic acid) should be less than 1 p.c. By reducing the amount of moisture to about 6 p c immediately after opening the nuts the action of moulds is checked and a better

yield of oil of good quality is obtained

The copra is treated in oil mills in a similar manner to that in which oleaginous seeds generally are worked up. The meal is expressed twice at a temperature of 55°-60°. The practical yield varies from 63 to 66 p.c., according the content of oil in the copra. The expressed oil contains a notable quantity of free fatty acid, the amount depending on the quality of the raw material

Copra press-cake is used for fertilisers and feeding-stuffs for cattle, but by separating the oil by appropriate methods a press-cake suitable for human food may be obtained. A meal suitable as a substitute for wheat flour and laid in pastry had the following composition: water, 7.35, oil, 32.14, ash, 4.05, crude fibre, 37.12; and proteins 20.34 p.c. (Parker and Brill, Philippine

J Sci, 1917, A, 12, 87)

In our climate coco-nut oil is, at the ordinary temperature, a solid white fat of bland taste, having the peculiar, though not unpleasant, odour of coco-nut. The odorous substances in coco-nut oil consist chiefly, according to Haller and Lassieur (Compt rend 1910, 151, 697), of methylheptylketone and methylnonylketone These two ketones, together with certain undefined volatile substances, are removed in the process of refining coco-nut oil for edible purposes. In order to obtain coco-nut stearin (chocolate fat), the liquid glycerides are removed by expression in hydraulic presses, when 50-80 pc of the liquid glycerides are expressed Inferior copra oils also are frequently subjected to the same process, and thus a commercial coconut stearin is obtained, which, on account of impurities, can only be employed in the manufacture of composite candles and night-lights The lower qualities of coco-nut oil are used in enormous quantities in the manufacture of soap, whilst the best grades are used for margarine

Coco-nut oil contains large proportions of myristin and, especially, of laurin, small quantities of palmitin, stearin, and triolein, and of

caprin, caprylin, and caproin.

Elsdon (Analyst, 1913, 38, 8), using the method of alcoholysis, found coco-nut oil to contain glycerides of the following fatty acids in approximately the proportions named: caproic, 2, caprylic, 9; capric, 10; lauric, 45; myristic, 20; palmitic, 7, stearic, 5; and oleic acid, 2 p.c. Methods for the detection of coco-nut oil in butter are based on the large proportion of lauric and myristic acids in the fat (Shrewsbury and Knapp, Analyst, 1910, 35, 385; 1912, 38, 3).

The saponification value of coco-nut oil lies between 246 and 260; the Reichert-Meissl value

of the oil is 7-8; its iodine value about 8-10; and its Hehner value 88 to 90 Owing to its peculiar chemical composition, coco-nut oil is not readily saponified by weak caustic leys. It requires for saponification alkaline leys of high strength, which, however, so easily convert it into soap, that it is only necessary to stir the fat and caustic alkali together at 30°-35°, and allow the mixture After a short time saponification takes place, with liberation of heat (Soapmaking by the 'cold process.') Soaps so formed are very hard, and combine with a large amount of water without becoming soft The soaps are further distinguished by their remarkable property of requiring large quantities of salt to throw them out of their aqueous solution. Hence they are used as 'marine soaps'

COCOSITOL v Inositol. CODAMINE v Opium CODEINE v OPIUM

CODEONAL. Trade name for a mixture of 1 part of medinal and 2 parts of codeine-veronal

COD LIVER OIL is obtained from the liver of the cod, Gadus morrhua (and of the dorsch, the young of G. morrhua, formerly considered a separate species, G callarius). The cod appears annually at certain times in enormous shoals on the coasts of the northern seas, chiefly on the coasts of Norway, Scotland, Newfoundland, the east of the United States and of Canada, the west coast of the United States and the coasts of Japan and Siberia The best-known and oldest fishing-grounds are those of Norway, where the fish make their appearance at regular

intervals of the year

In the early years of the manufacture of cod liver oil, the livers were heaped up in barrels and allowed to undergo a process of spontaneous putrefaction, whereby the liver cells burst and the oil exuded The oil so obtained was contaminated with decomposition products of the liver, and possessed that rank odour and nauseous taste with which even nowadays popular prejudice associates medicinal cod liver Of late years the manufacture of cod liver oil has been carried out in a scientific manner, and the best oil is at present obtained by steaming the livers in the freshest state obtainable The livers are washed immediately they are taken out of the body of the fish, and are placed in jacketed vessels in which they are heated to a temperature of 70°-80°, when the cells burst and the oil flows off This oil is known as 'raw medicinal oil,' and is consumed as 'congealing oil' in those countries where the Pharmacopeia The British Pharmapermits congealing oils copœia prescribes, however, a non-congealing oil, which will not 'set' at the freezing-point is prepared by expressing the fresh liver at a temperature not exceeding 85°, cooling the crude medicinal oil below zero, and filtering it at a temperature of about - 5°. The 'stearme' is sold as fish stearm of best quality for soapmaking purposes All unsound livers, and those which have been exposed for some time to the atmosphere, are then worked up for brown oil.' This oil is, of course, a genuine cod liver oil, but is (or should be considered) unfit for consumption, and is mostly used up in leather industries, under the name 'brown cod oil 'or 'cod oil.

light yellow, and the best qualities have a slight fishy odour only, and a by no means unpleasant taste Such oil has a sp gr of about 0 922-0 93 at 15°, and a saponification value of 175-192, and an iodine value of 140-175. The best qualities have an iodine value of about 175, for the longer the livers have been exposed, the lower will be the iodine value, so that this value alone may indicate, preliminarily, a good medicinal oil. The chemical composition of the cod liver oil glycerides is not yet completely Small quantities of palmitin occur in the oil, and are the cause of the separation of 'steame' The older statement that small quantities of lower saturated fatty acids, such as those of acetic, butyric, valeric, and capric acids occur in cod liver oil must be rejected, these statements being due to the fact that the cod liver oil in which those volatile fatty acids were found had already undergone decomposi-The greater number of the liquid fatty acids in cod liver oil have not been identified. Among the unsaturated fatty acids there occurs undoubtedly therapic acid, this can be identified and isolated by means of its octobromide, C<sub>17</sub>H<sub>36</sub>O<sub>2</sub>Br<sub>8</sub>, which does not melt at 200°, and decomposes, with blackening above that temperature

Japanese cod liver oil has a similar composition to that of the European oil Specimens examined by Tsujinioto (Chem Rev Fett Ind. 1913, 20, 70) had sp gr 0.9263-0.9290, saponification value 183 8-188 l; and iodine value 159 1-160 8 The fatty acids yielded 34 4-

36 17 p.c. of therapic octobromide.

The therapeutic value of cod liver oil which was attributed to the organic bases occurring in badly rendered cod liver oil must (in the author's opinion) be ascribed rather to the easy manner in which cod liver oil is digested than to any specific constituent, for the purest cod liver oil practically represents a mixture of pure glycerides with a small amount only of cholesterol and other unsaponifiable substances.

Like all liver oils, cod liver oil may be identified by the colour test which it gives with concentrated sulphuric acid in a solution of carbon tetrachloride or carbon disulphide better the quality of cod liver oil the fainter is the sulphuric acid colour test, this points to the fact that the colour test is due to small quantities of foreign substances which are dissolved by the oil from the tissues of the liver This is borne out by the fact that when completely hydrogenated, cod liver oil no longer gives any colour reaction with sulphuric acid A specimen of such fully hydrogenated oil prepared by Man-nich and Thiele (Ber. deut Pharm. Ges. 1916, 26, 36) melted at 65°, and had iodine value 1.2, and saponification value 186 2

CŒLESTIN v. STRONTIUM

CŒLINE, CŒRULEAN BLUE, CŒRULEUM v. Cobalt; also Pigments

CŒRULEIN, Anthracene green (v Alizarin AND ALLIED COLOURING MATTERS).

CŒRULIGNOL C<sub>3</sub>H<sub>7</sub> C<sub>6</sub>H<sub>3</sub> (OH)OCH<sub>3</sub>. A colourless oil of a burning aromatic taste, and a smell resembling creosote, obtained from beech pine and fir tars. Boils at 240°-241°; sp gr. 1.0564 at 15°. It is somewhat soluble in water, toil or cod oil. freely soluble in alcohol, ether, and acetic acid, The pure medicinal oil is pale yellow or It is coloured red by sulphuric acid, and

dark-brown by potassium hydroxide in contact with the air In alcoholic solution it gives with barium hydroxide a characteristic deep-blue colouration. Ferric chloride in alcoholic solution gives a green, in aqueous solution a deepred, colouration Carulignol is decomposed on treating with hydrochloric acid at 145°, according to the equation

 $C_{10}H_{14}O_2 + HCl = C_9H_{12}O_2 + CH_3Cl$  **CŒRULIGNONE** (Cedriret)  $(CH_3O)_*C_6H_9 \cdot O \cdot O \cdot C_6H_9(CH_3O)_9$ 

A steel-blue crystalline solid obtained by oxidising pyrogalloldimethylether in acetic acid solution with potassium dichromate (Hofmann, Ber 11, 335) It combines with primary amines to give deep-blue dyestuffs (Liebermann and Flatau, *ibid* 1897, 234, Moir, Chem Soc. Proc. 1906, 110, 1907, 307) (v CEDRIRET).

COFFEE. The beverage known as 'coffee' is prepared from the seeds (dried, roasted, and ground) of Caffea arabica (Linn), an evergreen shrub, indigenous to North-East Africa, and The use of coffee has been known from time immemorial in Abyssinia, whence it spread to the neighbouring Mahometan countries (where its use was at various times forbidden as being an 'intoxicating' beverage), and subsequently to all parts of the civilised world lt did not find its way into Europe, however, until the beginning of the 17th century It was introduced into England in 1652, and rapidly spread in popular favour in spite of heavy taxation and attempts to suppress the 'coffee houses,' for political reasons, until its importation reached a maximum of 371 million lbs ın 1847 Since then the consumption has declined, due, no doubt, chiefly to the greatly extended use of tea, and partly to the introduction of coffee substitutes At the present time the consumption of coffee in the United Kingdom is less than 30 million lbs, or less than I lb per head In the principal coffeedrinking countries of the world the consumption of coffee per head of the population is as follows Holland, from 14 to 15 lbs , United States of America, 12 lbs , Scandinavia, nearly 12 lbs ; Germany, 7 lbs , and France, 4 lbs

· The coffee tree belongs to the natural order Cinchonacea, which embraces, according to Lindley and Paxton, two species, C arabica and C paniculata (Aubl.) Other botanists distinguish many other species, as C microcarpa, C umbellata, C acuminata, C subsessilis, C laurina, C. racemosa, C bengalensis, C indica, &c. Some of these are probably only varieties However this may be, four species distinct from each other are now grown. (1) Arabian or Mocha coffee, having short upright branches with a brittle leaf, and seeds usually single in the (2) Jamaica coffee, bearing longer and more phable branches, a tougher leaf, and seeds almost always double in the berries. (3) East India or Bengal coffee, of which the leaf is smaller than the Jamaica variety, and the berry (4) Liberian coffee (C liberica) is very small taller and has larger leaves and fruit than the preceding varieties, but owing to its coarse flavour, it was not until recently of much commercial use It has, however, rapidly attained importance on account of its hardy habit and

power of resistance to disease, and is now largely cultivated by grafting on to its stem the more delicate and highly esteemed Arabian coffee. Trees growing wild in the Cingalese jungle attain a height of 14–18 feet; but under cultivation and subjected to systematic pruning, the plant rarely exceeds 6–9 feet. Within the tropics the coffee tree succeeds best at 1200–3000 feet above the sea-level, but grows at a height of 6000 feet. When the mean temperature is about 70°, it may be grown as far as lat. 36°. It is cultivated in the West Indies, Mexico Central America, Brazil, and Cayenne, in Western Africa, Egypt, Mozambique, and Natal; in Arabia, India, Ceylon, Siam, Sumatra, Java, Celebes, and other parts of the Eastern Archipelago, Fiji, and the islands of the Pacific. At the present time more than three-quarters of the world's supply comes from Brazil

The coffee consumed in the United Kingdom comes principally from Costa Rica and British India as regards the finer qualities, and the

inferior grades from Brazil

The manner of forming plantations varies somewhat in different countries The surface soil should be good and the subsoil not too stiff. Perhaps the best soil is that known in Brazil as 'terra roya,' a topsoil of red clay 3 or 4 feet thick, with a subsoil of gravel Virgin forest land is as a rule found very suitable from wind is most important, and a wet soil is Steep slopes are unsuitable owing to sh The loss due to this cause should be fatal repaired by good manure, well-rotted dung and ground bones being the best The young plants may be reared in nurseries, whence they are transplanted to their permanent position, or they may be gathered out of the old plantations; the age at which this is done varies, as in Brazil where the shrub is usually two years old on its removal, or as in the West Indies where it is planted out at 6 months Fruit is produced in the fourth year, and continues to be gathered for several years

The tree blossoms and bears fruit practically all the year round, and the fruit takes about 4 months to upen. There is thus in some places an almost perpetual crop, but, as a rule, there are two principal harvest seasons in the

vear

The coffee fruit consists of a bilobed berry somewhat resembling a small elongated cherry (by which term it is described commercially) The fleshy pericarp changes colour, as the fruit ripens, from green through yellow and red to a deep crimson When dried, it blackens and becomes hard and brittle, resembling a shell or Beneath the skin of the pericarp is a saccharine and somewhat glutinous pulp containing the seeds, which are usually double, lying face to face, but sometimes occurring singly. Each seed or 'bean' is enclosed in two membranous coverings—an outer, consisting of tough buff-coloured tissue, technically known as parchment,' which becomes loose and readily separates on drying; and an inner film of thin white tissue, closely adherent to the bean, and known as the 'silver skin.'

The principal methods in vogue for gathering and preparing the coffee berries for commerce are as follows—In Arabia, Egypt, and the Antilles, the berry is left on the tree till completely dried,

when it is easily shaken down and the dry outer envelope and the parchment removed by pounding in a mortar, beating with a flail, or simply rubbing in the hands. In many other places the berries are gathered by hand as soon as ripe but not dry, and are spread out on the ground in layers from 4 to 6 inches deep. These are exposed to the sun's heat for 3 or 4 weeks, with frequent turning by means of spades, when the bean becomes easily removable from the fruit by gentle trituration. As this process of drying sometimes imparts to the coffee a disagreeable flavour owing to putrefactive changes in the pulp, recourse is frequently had to artificial heat. The most modern method is to 'pulp the berries by crushing between rollers so as not to injure the beans, which are then subjected to a fermenting process in tanks, with or without the addition of water, after which they are separated from the pulp by 'washing,' and dried, either by the sun or by artificial means. Coffee beans in the form of 'cherry' rarely enter into commerce, but frequently retain the parchment and silver skin, which, however, before roasting, are removed by rollers and separated by winnowing They are then graded into different sizes by special machinery, which also separates the round or 'pea-berries' from the ordinary flat-faced beans. The colour of coffee beans varies from a pale-yellow passing through various shades of green to a slatey grey. The following table will indicate the variations ın sıze:-

				a measu	
	nord	ung ac	grai	ns of wat	er
Fine brown Java				187	
Fine Mysore .		•		198	
Fine Neilgherry		•		203	
Costa Rica .				203	
Good ordinary Gua	teme	ıla		207	
Good La Guayra				210	
Good average Sant	os			213	
Fine long-berry Mo	cha			217	
Good ordinary Jav				223	
Fine Ceylon planta	tion			225	
Good average Rio				236	
Medium plantation	(Cey	zlon)		238	
Manilla .				248	
Ordinary Mocha			. `	270	
West African				313	

Coffee undergoes important changes in the process of roasting, the object of which process is to develop its aroma and to destroy its toughness so that it may easily be reduced to powder Great care is necessary to ensure an equable roasting of all the beans, and to attain the exact point at which torrefaction should cease the one hand, rawness is not removed, the flavour of the bean is not developed, and the coffee is hard to grind, on the other hand, if too much heat is permitted, carbonisation takes place, and a strong empyreumatic flavour is produced, whict renders the coffee infusion nauseous

The roasting process is conducted in cylinders (preferably of strong wire gauze to permit of the escape of steam and other volatile products as fast as they are formed) heated by coke fires or gas. The loss in weight is from 15 to 18 pc, of which about one-half is due to moisture, and the rest chiefly to volatile products derived from the fat, and from the sugar during the conversion of the latter into caramel. There is also some loss of caffeine. At the same time, an empyreumatic oily substance is developed, known as caffeol, which, in spite of its small amount, imparts to roasted coffee much of its characteristic aroma. The most favourable temperature for developing the maximum aroma during roasting is a little above 200°. After roasting, it is most important that the beans should be rapidly cooled, and this is effected by spreading them on a large sieve through the meshes of which they cannot pass, but through which the air is forcibly drawn by means of a powerful exhaust connected with a closed chamber immediately below the sieve. Since the essential oil rapidly escapes after the bean is broken up, coffee should not be ground until required for infusion

Bernheimer (Wien Akad 81, [2] 1032, 1042) found that during the roasting of coffee, water vapour is first given off, then suddenly an evolution of carbonic anhydride, and distillation of volatile substances commences The distillate consists of an aqueous liquid and a solid part. The former contains caffeine, caffeol, acetic acid, quinol, methylamine, and acctone, the latter consists of fatty acids, chiefly palmitic Pyrrol escapes during roasting Of fatty acids, 0.48 p c of the coffee employed was volatilised, of caffeine, 0.18 p.c., and of caffeol, 0 05 р с.

Erdmann (Ber 1902, 35, 1846) attributed the aroma of roasted coffee to traces of acetic acid furfural, valeric acid, an unstable nitrogenous compound and a phenolic body resembling creosote Various samples of roasted coffee yielded from 02 to 025 grm of pyiidine per kilo (Bertrand and Weisweiller, Compt rend

1913, 157, 212)

Bernheimer found that raw coffee beans (50 kg) roasted in a revolving drum yielded a liquid distillate (5 lities) and a solid portion (680 grams) floating thereon The uncondensable vapours contained carbonic anhydride, and, on passing them through dilute hydrochloric acid, a resinous body was deposited, having the ap-

pearance of pyrrol-red.

A Liquid distillate -The crude liquid distillate filtered from solid particles has a yellow colour and the fragrant odour of coffee. On agitating it with ether, and evaporating the ethereal extract over the water-bath, a dark heavy oil was obtained, smelling like coffee, and yielding, on redistillation, first a few drops of liquid having the odour of acetone, afterwards, for the most part, acetic acid and water. The thermometer then 10se quickly to 200°, above which point there passed over an oil smelling like coffee, the temperature at the same time rising to 300°. Above this point there passed over a very small quantity of liquid, which immediately solidified to a mass having the buttery aspect and rancid odour of the higher fatty acids This portion, and the liquid which passed over from 200° to 300°, were neutralised with sodium carbonate, whereupon a thick darkcoloured oil was thrown down, which was separated from the supernatant liquid by means of a tap-funnel, and washed with water rendered very

slightly alkaline by potash.

The alkaline wash waters contained no fatty acids, but, when acidified with sulphuric acid and agitated with ether, they yielded a substance

which, after evaporation of the ether, filled the liquid with a pulp of crystals, and after precipitation crystallised in colourless needles, exhibiting the characters of quinol (m.p. 171°), converted into quinhydrone by the action of ferric chloride.

The oil above mentioned, designated by Bernheimer as caffeol, was dried over calcium chloride and fractionally distilled, the greater part passing over from 195° to 197°. The distillate smells very strongly of coffee, is apparently insoluble in cold water, but imparts to it the characteristic odour, slightly soluble in hot water, very slightly in aqueous potash, very casily in alcohol and ether, the alcoholic solution giving, with ferric chloride, a red colouration, which does not disappear on addition of sodium carbonate. Analysis led to the formula  $C_8H_{10}O_2$ . Caffeol is oxidised by chromic acid mixture, without formation of any definite products. By fusion with potash it yields salicylic acid. The portion of the crude liquid distillate

not taken up by ether, when concentrated over the water-bath, yielded groups of needle-shaped crystals, exhibiting the composition and properties of caffeine  $C_8H_{10}N_4O_2$  (0.28 p.c. of the coffee). The mother liquor was found to contain methylamine and trimethylamine.

B Solid distillate —The black mass floating on the crude liquid distillate was found, by a preliminary examination, to consist of a mixture of the higher fatty acids. By saponification, and treatment of the product by Heintz's method of separation, this mixture yielded, as chief constituent, palmitic acid melting at 60°, and a very small quantity of another acid having a melting-point a few degrees higher than that of lauric acid.

The chemical composition of coffee has been investigated by many chemists with various results. The following table gives analyses by O Levesic (Arch Pharm [3] 8, 294) of some of the principal commercial varieties of coffee:—

	Gummy matter	Caffeine	Fat	Tannic and caffetannic acids	Cellulose	Aslı	Potash	Phos- phoric acid
						-		2.4
Finest Jamaica plan-								
tation .	25.3	1 43	14 76	227	33.8	38	1 87	0.33
Finest green Mocha	22 6	0.64	21.79	23 1	29 9	41	2 13	0.42
Cevlon plantation .	23 8	1 53	14 87	20 9	360	40		0 27
Washed Rio	27.4	1.14	15 95	20 9	32 5	45		0.51
Costa Rica	20 6	1.18	21.12	21.1	33 0	49		0 46
Malabar	25 8	0.88	18 80	20 7	31 9	43		0 60
East Indian .	24 4	1 01	17 00	19 5	36.4			

Commaille (Mon Sci [3] 6, 779) found in undressed Mysore coffee.

Hygroscopic v	vater	(extre	mes		
of 24 determ				6 3 to	157
Fatty matters				$12\ 68$	
Glucose				$2\ 60$	
Dextrin .					
Legumin-casei	n.			1 52	
Albumen .	•			1 04	
Total quantity	of ca	ffeine		0 42 to	131
Ash		•		$3 \; 882$	
Coffee extract					
,, ,,		arm wa		37 20	
Coffee extract	with	alcoho	l of		
60 pc				23 15	

Dragendorff gives the following results of analyses of 25 different samples of coffee from Ceylon, Java, Malabar, Mocha, Mysore, and other places —

*	3		Minimum per cent.	
Caffeine .		2 21	0 64	1 16
Ash		4 87	3 83	441
Phosphoric act	d.	0.72	0.28	0.42

The principal constituents of coffee are alkaloids (caffeine and caffearine), caffetannic acid, cellulose, fat, gum, protein, sugar, water, and mineral matter.

Caffeine  $C_5H(CH_3)_3N_4O_2$ , trimethylxanthine, is identical both chemically and physiologically with theme, the alkaloid of tea, and is closely related to the obromine (dimethylxanthine), the principal alkaloid of cocoa (q.v.). According to

Payen (Ann Chim Phys [3] 16, 108), caffeine is present as a caffetannate, Gorter (Annalen, 1908, 358, 327) states that caffeine occurs in coffee combined with potassium as potassium caffeinechlorogenate  $\mathrm{C_{32}H_{36}O_{12}K_2(C_3H_{10}N_4O_2)_2}$ ,  $\mathrm{2H_{2}O}$  When pure it consists of white silky needles, slightly bitter and odourless, almost insoluble in ether, sparingly soluble in cold water and alcohol, but more soluble in hot water, and readily soluble in chloroform, by means of which it may be extracted from coffee. The crystals begin to sublime below the boiling-point of water

Caffeine is found also in guarana leaves, kola-nut, maté or Paraguay tea and to a small extent in cocoa as well as in coffee and tea, and gives to these their exhilarating qualities. Taken in excess, caffeine is a poison. In coffee it is present to the extent of from 0.6 to 2.2 p.c., the average being 1.1 p.c. Lendrich and Nott bohm (Zeitsch Nahr Genussm. 1909, 18, 299) report nearly 3 p.c. of caffeine in certain African coffees, whilst in Madagascar and other islands off the east coast of Africa a variety of coffee has been found entirely devoid of caffeine Accoffee has recently been placed on the market after being deprived of practically the whole of its caffeine.

Numerous methods have been proposed for the estimation of caffeine That of Lendrich and Nottbohm is detailed in the Analyst, 1909, 214, and gives accurate results. A less tedious method, but sufficiently accurate for ordinary purposes, is that of Fendler and Stuber (Zeitsch. Untersuch. Nahr. Genuss. 1914, 28, 9-20) abstracted in the Analyst, 1914, 394.

Another alkaloid, Caffearine, having narcotic properties, is stated by Paladino (Gazz. chim. ital. xxv. 1895) to be present in coffee in very

small quantity.

Caffetannic acid presents certain of the characteristics of tannin (which, however, is not found in roasted coffee) and to it the bitter taste of coffee is chiefly due. It is very susceptible to change, and its composition has been much disputed. According to Gorter (vide supra), it is a mixture of a new acid discovered by him and called caffalic acid, with chlorogenic acid. A method for its estimation in coffee has been devised by Krug (U.S.A. Dept. of Agric Div. of Chem. Bull. 13, 908) According to Bell, caffetannic acid is present in coffee to the extent of from 3 to 5 p.c., and when distilled with manganese dioxide and sulphuric acid yields quinone.

Cellulose.—In coffee this is of a very hard and horny character, and difficult to define under the microscope, especially after roasting, although the chief characteristics of the cellular tissue are more or less retained. Coffee cellulose is dissolved by an ammoniacal solution of copper oxide, and gives a blue colour with sulphuric acid and iodine. Remnants of the 'silver skin' are always present even after roasting, and its structure, a thin membrane with adherent, thick-walled, spindle-shaped, and hollow cells,

is quite peculiar to coffee.

Fat forms a considerable constituent of coffee, varying from 10 to over 20 p.c. A loss occurs in roasting, due to the separation of volatile fatty acids When pure, coffee fat is white, inodorous, and of the consistency of butter, melting at 37 5°.

Protein, chiefly in the form of insoluble albuminoids, occurs in coffee to the extent of

from 10 to 14 p.c., the soluble albuminoids constituting about 2 p.c. There is but little difference in this respect between raw and roasted coffee.

Sugar.-König and Bell found from 5 to 9 p.c. of sugar in raw coffee, but this practically disappears in roasting, being almost entirely converted into caramel. Herfeldt and Stutzer (Zeitsch. angew. Chem. 1895, 469) state that sugar is not present as such in the raw bean, but only in the form of a glucoside. Bell, however, found that when the saccharine extract of the beans is boiled with a little dilute sulphuric acid, a quantity of glucose is obtained nearly equivalent to the weight of dry extract operated upon, a result inconsistent with the existence of sugar as a glucoside. Bell considers that this sugar is not present as sucrose, since it is not inverted even after prolonged boiling with acetic acid. On the other hand, the with acetic acid chemists of the United States Department of Agriculture have isolated cane sugar in considerable amount as pure, well-defined crystals (v. Bulletin, 13). It is usually accompanied by some invert sugar probably formed during the process of extraction, and a small percentage of dextrin. Starch is rarely found in the coffee bean.

Water—The moisture in raw coffee varies from 8 to 14 pc, the average being about 11 pc. After roasting, it rarely exceeds 3 pc, and frequently falls as low as 0 5 pc.

Mineral matter—The total amount of ash varies from 3.5 to 4.0 p.c. in law, and from 4.0 to 5.0 p.c. in roasted coffee. The composition of the ash is shown in the following table (U.S.A. Dept. of Agric. Div. of Chem. Bull. 13, 904).—

Constituents	 		Mocha	Maracaibo	Java	Rio
Sand			1 44	0 72	0 74	1.34
Silica (SiO <sub>2</sub> ).		.	0 88	0 88	0.91	0 69
Ferric oxide (Fe <sub>2</sub> O <sub>3</sub> )		. [	0 89	0 89	1.16	1 77
Lime (CaO) .		.	7 18	5 06	4 84	4 94
Magnesia (MgO)		.	10 68	11 30	11 35	10 60
Potash (K 2O)		. 1	59 84	61 82	$62\ 08$	63 60
Soda (Na, Ö)			0 48	0 44		0 17
Phosphoric acid (P <sub>2</sub> O <sub>5</sub> )			12.93	13 20	14 09	11 53
Sulphuric acid (SO <sub>2</sub> )			4 43	5 10	4.10	4 88
Chlorine (Cl)		1	1.25	0 59	0.73	0.48
		1	100 00	100 00	100 00	100 00

The principal constituents are potash, magnesia, lime, and phosphoric acid. Ludwig, in his analyses of various parts of the Brazilian coffee tree, shows that the silica increases from the root upwards to the leaves and pericarp of the fruit, after which it declines in the parchment skin, and is nearly absent in the bean (Arch. Pharm [3] 1, 482). Graham, Stenhouse, and Campbell inferred that sodium compounds are entirely absent in coffee (vide table in Bell's Food Adulteration, 46), but although this is not absolutely correct, the amount, as shown in the above table, is very minute.

Adulteration of coffee is very common, and

the number and variety of the adulterants is almost unlimited

Colouring is frequently employed in order to disguise damaged beans and give them the appearance of superior quality, a practice analogous to the 'facing' of tea. The substances most commonly employed are, for the raw beans, chrome yellow, yellow ochre, indigo, Prussian blue, azo- dyes, and soapstone for polishing; and for the roasted beans, burnt umber, oxide of iron, Venetian red, ferruginous earth and clay. The inorganic colouring matters are best sought for in the ash or in the sediment obtained by shaking the beans in cold water; indigo and

Prussian blue by their usual reactions; and other organic colouring matters after extracting them with alcohol.

Glazing is a process resorted to for the alleged purpose of improving the keeping properties of the roasted beans, and the substances used, a mixture of egg albumen with glucose or gum, are said to have the effect of clarifying the infusion. The improved appearance of the beans and increased weight due to the retention of moisture, are probably also important considerations. The latter result is sometimes attained by the addition of sugar before roasting, or by actually steaming the beans and coating them with glycerine, palm oil, or vaseline, to prevent evaporation of moisture Bertarelli (Zeitsch Nahr Genussm. 1900, m 681) describes the adulteration of roasted coffee beans with a solution of borax, which not only gives the beans a glossy appearance, but also adds as much as 12 p c to their weight.

Shellae is occasionally used for glazing, and Nottbohm and Koch draw attention to the fact that shellac is sometimes coloured by arsenic trisulphide Coffee beans which had been glazed with arsenical shellac were found to be strongly contaminated with arsenic (Zeitsch

Nahr. Genussm 1911, 21, 288).

Coffee substitutes comprise a large number of vegetable substances, roasted and ground, and having for their object either the production of a stronger infusion than can be obtained from coffee alone, or the simulation of coffee in order to give weight by the use of thick-walled vegetable tissue of hard or leathery consistency The former, of which chicory is the type, includes acorns, beet refuse, biscuits, bran, cereals (barley, buckwheat, maize, malt, oats, rye, and wheat), caramel (burnt sugar), carrots, dandelion root, figs, lupin seeds, mangolds, parsnips, peas, beans, and other leguminous seeds, turnip, &c, whilst the latter comprise such substances as coffee husks, cassia seeds, date stones, grape stones, locust or carob beans, sawdust, vegetable ivory dust, &c

Imitation coffee -Although adulteration is naturally more prevalent in ground coffee than in the beans, yet the latter have been prepared artificially and on a large scale from the above adulterants (generally without any admixture of coffee), held together by means of gum, molasses, or stiff clay, and skilfully moulded

into the form of coffee beans

The detection of the adulterants above referred to is best effected by means of the microscope By simply digesting with water, a ready means is afforded of differentiating between the pale soft-walled tissues of cereals and roots and the dark-coloured and hard-walled tissue of coffee The former may readily be detected and identified by the presence and characteristics of their starch, coffee (and chicory) being practically free from starch The use of bleaching reagents is of great assistance in the microscopical examination, and reference might usefully be made to the drawings and photographs reproduced in the works of Bell, Leach, Villiers et Collin, and Wiley, and in the Coffee Report of Graham, Stenhouse, and Campbell, referred to below.

The chemical differences between coffee and its adulterants are also strongly marked, as

mixtures usually give high aqueous extract and sugar, and low fat and caffeine. The density is a useful factor in detecting the sophistication of coffee, and, in particular, the artificial moistening of the berries In the Paris Municipal Laboratory, Regnault's volumenometer, as modified by Dupré, is used for this purpose. According to the French chemists, the density of raw coffee varies between 1 041 and 1 368, and that of roasted coffee between 0.500 and 0.635.

By far the most important adulterant of coffee is chicory (qv.), but owing to the public preference for an admixture of chicory with coffee, it is now regarded in the light of an adjunct to rather than an adulterant of coffee, provided its presence in the mixture is disclosed. Chicory has little in common with coffee except a considerable proportion of sugar, which is converted into caramel in the roasting, and thus imparts to the infusion a bitter taste and aroma somewhat resembling that of coffee This and the stronger infusion obtained as well as the comparative cheapness of chicory and similar substances, has, no doubt, led to the present extensive use of chicory and other coffee substitutes

An elaborate research into the nature of coffee and chicory, with a view to the detection of adulteration, was made at the instance of the Commissioners of Inland Revenue in 1852 by some of the leading botanists and chemists of the day, viz. Lindley and Hooker, and Graham, Stenhouse, and Campbell, and the results, chemical and physical, as well as microscopical embodied in their reports are of great practical

value even at the present time

The following table by Graham, Stenhouse, and Campbell, shows the spgr at 60°F. of sulutions made by treating one part of each of the substances named (roasted and finely ground) with 10 parts of water (by weight), raising the mixture to boiling-point, where it is maintained for half a minute, then filtering and cooling :-

Specific Gravities of Solutions at 60°F. One Part of Substance to 10 Parts of Water.

)	ne Part of Sv	ıbslanı	ce to	10	Paris	of	n c	ue
	Spent tan					100	)2	1
	Date stones					100	)2	9
	Lupin seed					100	)5	7
	Acorns					100	07	3
	Peas .					100	07	3
	Mocha coffee	,				100	80	0
	Beans .					100	98.	4
	Neilgherry co	offee				100	8(	4
	Plantation C	eylon	coffe	е		100	980	7
	Java coffee					100	980	7
	Jamaica coff	ee				100	98	7
	Native Ceylo	n coff	f <b>e</b> e			100	9	0
	Costa Rica c	offee				100	$99 \cdot$	5
	Parsnips					10	14	3
	Carrots					10	17:	1
	Black malt					102	21	2
	Turnips					109	21.	4
	Rye malt					109	21.	6
	English chice	orv				109	21.	7
	Dandelion ro					109	21.	9
	Red beet					102	22	1
	Foreign chic	ory				102	22	6
	Guernsey chi	icory				103	23	2
	Mangold wu	rzel				102	23	5
	Roasted figs					102	24 ·	9
	Maize					10	25.	3
	Bread raspin	gs				10	26	3
		0						

It will be seen from the above table that the low sp.gr. of the coffee infusion distinguishes it from the roots and cereals, and when chicory is the only substance admixed with the coffee, its percentage may be calculated with approximate accuracy by a comparison of the density of the infusion of the mixture with the average densities of coffee and chicory given in the

foregoing table.

The same observers lay stress on the relative colouring power of roasted coffee and chicory, the latter being four and a half times as great as the former. This property affords a simple device for detecting the presence of roasted chicory or other substances containing caramel when admixed with coffee. A few grains of the sample are placed on the surface of cold water in a glass, without stirring, when, if roasted chicory or similar substance be present, each particle emits a yellowish-brown coloured cloud, which is rapidly diffused in streaks through the water till the whole acquires a brownish colour, whilst pure coffee gives practically no colour to the water in similar circum-The chicory readily sinks, whilst nearly the whole of the coffee floats.

If examined under the microscope, the sediment of chicory which reaches the bottom of the vessel will be found to exhibit the characteristic loose, large cells, with dotted vessels and branching lactiferous ducts Even to the touch the difference is evident, for, whilst chicory under these conditions will feel soft and smooth, the coffee grains remain hard and

A. Winter Blyth (Foods; their Composition and Analysis, 359) gives the following summary of the influence of chicory on the coffee

mixture :-

(1) It decreases the gum, the latter seldom rising in chicory to more than 15 pc, whilst in coffee it has not been found less than from 21 to 28 pc.

(2) It increases the sugar, roasted coffee having seldom so much as 2 p.c. of sugar; whilst chicory, when roasted, has at least 8 or 9

(3) It decreases the fatty matter, the fat of chicory ranging from 1 to over 2 p.c., that of

coffee being about 14 to over 20 p c. (4) It decreases the tannic and caffeotannic

acids, chicory being destitute of tannin. (5) It decreases the caffeine, chicory pos-

sessing no alkaloid.

(6) It profoundly modifies the constitution of the ash, especially by introducing silica, which is not found to any great extent in coffee

Graham, Stenhouse, and Campbell laid stress upon the presence of silica as a means of detection of the admixture of chicory. analysis, showing the principal differences in the composition of the respective ashes, gives:

	in conee ash	in chicory ash
Silica and sand		10 69 to 35 85
Carbonic acid	14.92	1.78 ,, 3 19
Ferric oxide	0.44 to 0.98	3.13 ,, 5.32
Chlorine .	0.26 ", 1.11	3.28 ,, 4.93

There is a marked difference in the proportion of ash soluble in water, that from coffee Vol II.—T.

amounting to from 75 to 85 p c, and in chicory from 21 to 35 p c.

Negro or Mogdad coffee consists of the seeds of Cassia occidentalis It contains no caffeine nor starch, and its infusion more nearly approaches the character of genuine coffee than any other known coffee substitute

Mussaendar coffee is derived from the seeds of Gaertnera vaginata It contains no

caffeine

French coffee is a description commonly applied to a mixture of coffee with burnt sugar and a large proportion of chicory. As the latter ingredients readily absorb moisture on exposure to the air, such mixtures are usually preserved

Various fraudulent coffee substitutes have been imposed upon the public by specious advertising under fancy names, and as possessing certain desirable qualities not found in coffee. Such are Kunst kaffee, consisting of roasted roots or cereals, and often moulded into the form of coffee beans; Pelotas coffee and Coffee surrogate, made from ground acorns, Date coffee from date stones, or torrified dates mixed with a small proportion of coffee; and many others.
The grosser forms of adulteration above

referred to are much more prevalent in the coffee-drinking countries of the continent of Europe and in the United States of America than in the United Kingdom.

COGNAC v BRANDY.

COHUNE NUT FAT. The nuts of the Cohune Palm (Attalia cohune), indigenous to British Honduras, yield from 65 to 72 p c. of a white solid fat resembling coco-nut oil Sp gr at 100°/15 5° 0 868 to 0 871; acid value 1.2 to 204, saponification value 252 4-256 5; iodine value 11 0 to 13 7; Hehner value 87 7; Reichert-Meissl value 6 8 to 8 3; Polenske value 12.5 and Solidifying point of fatty acids 197°-21 0° (Bull. Imp. Inst. 1913, 11, 226; 1914, 396).

#### COKE MANUFACTURE AND THE RE-COVERY OF BY-PRODUCTS.

Introduction —Previous to the early nineties. practically the whole of British metallurgical coke was made in the old beehive oven, which, as will be seen from what follows, is a semicombustion process, involving the loss of the whole of the gas and by-products, together with With the a portion of the possible coke yield gradual disappearance of the old prejudices against coke made in by-product ovens, the latter and much more economical process began to assert itself, and in 1898, 1.25 million tons of coal were being carbonised in by-product coke ovens, in 1905 the quantity of coal so treated reached 3.31 millions; and in 1916, about 14 60 millions Meanwhile, the number of beehive ovens in operation had decreased from 16,037 in 1910, to 6892 in 1916 According to figures published by the Coal Conservation Committee at the end of 1917, some 8700 byproduct ovens were in operation in Great Britain (of which 8000 were fitted with benzol recovery arrangements) with a total carbonising capacity of 17 5 million tons of coal per annum. In 1917, 166,354 tons of ammonium sulphate were produced in British coke-oven plants. The practice of by-product recovery had become universal in Germany years ago; whilst in the

United States the by-product plants, either building or in operation at the end of 1917, were capable of producing 24 million tons of coke per annum. In view of this rapid adoption of the by-product coking process in all these countries, the following article will be confined to a description of modern coking methods viewed chiefly from a by-product standpoint, with only a passing reference to the wasteful beehive process.

History of the by-product coking process— The earliest reference which we can find of any observation being made on coal as a source of by-products is that contained in the memoirs of Johann Joachim Becher, duting about the end of the 17th century, Becher states that he has found means for treating coal 'so that it no longer smoakes nor stinks,' and in doing so

made a tar equal to the Swedish.

The use of coke in iron smelting became general within 50 years of Becher's discovery, and a French metallurgist, De Gensanne, describes a process working at Sulzbach near Saarbrucken prior to 1768, where coal was coked for iron smelting, tar being recovered as a byproduct, Stahl is said to have been the inventor of this process, and he produced therefrom an 'oil,' bitumen, sal-ammoniac, and lamp-black

In 1781 Archibald, Earl of Dundonald, was granted a patent for 'allowing coals to burn or ignite without flaming, so by their own heat to throw off the tar or oils that they contain Dundonald stated that the only method whereby tars could be obtained prior to his discovery, was by a distillation in closed retorts, where the admission of external air was prevented, and where other coals were required for heating besides the coal contained in the closed vessel It is interesting to note that Dundonald was perfectly aware of distillation in closed retorts, which is the basis of modern by-product coking methods, and it is also evident that his knowledge of closed retorts was 16 years previous to the introduction of coal gas for street and household illumination in 1797

In 1838 Frederick Neville obtained protection for an invention which consisted in heating ordinary gas retorts by the waste heat from beehive coke ovens, the distilled gases from the retorts being passed though condensing and scrubbing appliances William Newton, in 1852, introduced the use of an aspirator or pump for withdrawing the gases from coke ovens through and  ${f rubbers}$ Newton likewise mentions the use of acids for removing the ammonia from the gases. In 1860 W. H Kingston patented a coke oven provided with external furnace, from which the heat evolved by the combustion of an auxiliary supply of coal was conducted round the oven through a system of flues; the evolved gases were passed through a series of pipes and condensing vessels for the recovery of the by-products Kingston's patent is especially noteworthy, since he recognises that the true method for the recovery of by-products from coal lies in heating the coal in closed chambers, and not in chambers to which air is admitted

With regard to Continental practice, in 1850 M M Pauwels and Dubouchet, and in 1862, Pernelot, introduced systems of carbonising

coking coals in long rectangular chambers, heated from below by the regulated combustion of part of the gas in flues specially constructed under the floor, for the purposes of manufacturing metallurgical coke and removing tars, oils, and ammonia The Pernelot system was tried in the north of England in 1870, but the coke produced was too soft for metallurgical purposes

In 1856 Knab erected a coal-distillation plant at Commentry, on the closed-retort system with bottom flues, but made no attempt to recover by-products Later Knab was joined to recover by-products (1866-1873) introduced by Carvès,  $\mathbf{w}$ ho improved methods of heating by side flues in addition to the bottom flues, thus obtaining a more even distribution of temperature and a Carvès likewise quicker rate of carbonisation utilised the gases (from which the by-products had been extracted) in the side-wall flues, in addition to the auxiliary coal as employed by The Knab-Carves ovens at Commentry gave the following results -

Large coke .		70 00
Breeze		1.50
Dust		2 50
Graphite		0.50
Tar		4.00
Ammoniacal liquor		9 00
Gas .		10.58
Loss		1 92
		$100 \ 00$

Further plants on the Carvès system were erected at Bessèges and Terre-Moire, near St. Etienne, between the years 1866 and 1873, and were still further improved, the Carvès ovens working at Besseges in 1880 were built with carbonising chambers only 24 inches wide, to facilitate rapid heat penetration and quick carbonisation In 1881 Mr. H. Simon of Manchester introduced a recuperator, whereby the air necessary for the combustion of the gas in the side-wall flues of the Carves oven was preheated by the waste gases escaping from the flues, and thus the rate of carbonisation was still further increased, the combined work of the two investigators resulting in the wellknown Simon-Carvès oven, which may be held to be the precursor of the modern by-product coking retort

Following on the success of the Simon-Carvès process, other coke-oven patents were filed, viz Semet-Solvay, 1880, Hussener, 1881; Lurmann, 1882; Otto and Brunck, 1883, but although development was rapid on the Continent, little was done in this country before the nineties, owing to the prejudice of iron smelters against by-product coke; that this prejudice has now disappeared will be seen from the rapid adoption of the process, indicated by the figures at the beginning of this article. The by-product coking plants which have generally found acceptance in this country are the Otto-Hilgenstock, Simon-Carvès, Koppers, Semet-Solvay, Simplex, Coppée, and Hussener.

Non-by-product and by-product ovens compared.—Prior to the establishment of the by-product coking process, the type of oven universally employed was that known as the beehive.

This oven (Fig. 1) consists of a fireclay chamber, bechive shape in elevation and circular in plan, provided with a loose brick charging door at ground level, and with an exit flue in its crown for carrying the waste gases to the chimney. The 'coal is charged into the oven by spades, through the charging door, to a height of 2 feet 6 inches to 3 feet, is then ignited, and the charging door bricked up and plastered, a small hole 6 inches square being left for the admission of air. The upper layer of the coal burns and or an. The upper layer of the coar burns and sets up distillation of the under layers, the evolved gases passing upwards and burning in the dome of the oven, with the air drawn in at the door. The dome is thus soon heated up to a high temperature, and by radiation powerfully assists in carbonising the charge of coal below. This operation of distillation and immediate combustion proceeds until the whole of the volatile constituents of the coal have been evolved, which for a 10-ton charge occupies 70 hours. The coke is then quenched with water and withdrawn by rakes

It will be seen that by this process, the whole of the volatile products, that is to say, benzols and ammonia together with the permanent gases, are burned immediately they are evolved within the coking chambers itself; further it is not possible to so regulate the combustion of these gases without burning a portion of the coke in addition; in actual practice, from 15 to 20 pc of the coke is burned to waste the reason, too, that the coke is quenched inside the oven, the temperature of the latter is lowered, so that before the next charge can be coked, considerable expenditure of heat is necessary to raise the oven to coking temperature. The removal of the coke by hand-rakes

is also tedious and expensive With the object of overcoming the main drawbacks of the bechive oven, Coppée introduced a non-by-product retort of rectangular form, from which the coke could be discharged en bloc by a ramming machine and quenched outside the oven, thus expediting the operation In Ćoppée's and greatly increasing the yield oven (Fig. 2) the gases evolved from the coal are drawn into flues in the side walls where burned, and in this way they are

bustion within the coking chamber and loss of coke, as occurs in the beehive oven, is avoided.

The by-product retort (Fig. 3) is built on the same lines as the Coppée, that is to say, it is a rectangular chamber heated by flues in the side walls, and the coke is discharged by a ramming machine; the essential point of difference lies in the fact that in the Coppée the whole of the gases containing the by-products are drawn immediately they are evolved into the side-wall heating flues, where they are burned; in the by-product retort, the gases and products are drawn from the oven through cooling and washing plant for the recovery of the tars, ammonia, and benzol, and the permanent gas only is burned in the flues after the said products have been extracted. In the Coppée, the whole of the permanent gas is burned in the heating flues; in the by-product oven, only one-half to two-thirds of the gas is utilised in the flues, the remainder being available for heating or power purposes.

The following diagrams illustrate the essential differences in the three types of oven




FIG 1 -BEEHIVE.




Fig. 2.—Coppee Fig. 3.—BY-PRODUCT RETORT.

It will be seen, therefore, that in addition to effecting the theoretical coke yield, and a large output, the by-product retort yields valuable by-products and a considerable yield of power or heating gas The average yields (calculated upon 100 parts of dry coal) of coke and by-products obtainable from British coking coals of different origin by the most up-to-date methods. are approximately as follows -

Origin	Coke	Anhyd tar	Crude ben <b>z</b> ol	Ammo- nium sulphate	
Durham	75 0	3 85	0 93	1 16	
South Yorkshire	70 0	3 62	1 37	1 35	
Lancashire	70 0	3 50	1 20	1 25	
South Wales .	80 0	2.62	0 60	0 95	
Scotland	68 0	3 25	0 70	1 00	

The by-product process described in detail -The by-product retort usually measures from 30 to 33 feet in length, from 5 feet 9 inches to 7 feet 6 inches in height, from 17 to 22 inches in width, and is built in batteries of from 25 to 60 ovens, capable of carbonising from 100 to 350

Table showing the Return from carbonising a Lancashire Coal containing 30 p.c. Volatile Matter, i.e a Theoretical Coke Yield of 70 p.c. in the Three Types of Oven.

Beehive	Coppée	By-product retort  Coking chamber heated externally by gas from which by-products have been extracted: quenching outside		
Combustion within coking chamber, quenching inside oven	Coking chamber heated ex- ternally by gas from which by-products not extracted quenching outside oven			
Yield of dry coke = 56 p.c.	70 p c.	oven 70 p.c.		
,, tar —		40 to 45 p c.		
" sulphate of ammonia		1 0 to 1 2 p c.		
" benzol —		25 to 30 gallons of crude benzol testing 62 p.c. at 120° per ton of coal		
Waste heat-		Per con er com		
In most cases not utilised but approximately equal to 0.75 ton of water evapo- rated per ton of coal carbonised	Always utilised · equal to 1 ton of water per ton of coal carbonised	Always utilised. equal to 1 ton of water per ton of coal carbonised Surplus gas if required for other purposes than steam		
A		raising, $eg$ gas engines or furnaces, 3000 to 5000 cubic feet per ton of coal		
Amount of coke produced per oven per week = 7 to 10 tons	24 to 26 tons	24 to 26 tons		

tons of coal per day. Each oven is provided with cast-iion or rolled-steel doors at each end, lifted by a small crane whenever an oven is discharged, and with three holes in the roof through which the coal is charged into the oven from the charging hoppers, a fourth hole is provided, through which the gases are led to an ascension pipe from which they pass to the by-product plant. The oven limings and combustion flues are constructed of silicious firebricks containing from 80 to 84 p.c. of silica, from 13 to 18 pc of alumina, and 2 pc of other constituents (iron oxide, lime, magnesia, and the alkalis) In the United States, however. silica bricks (SiO<sub>2</sub>=94 p c) are extensively employed They are superior in heat conductivity to a more aluminous brick, but are less satisfactory when 'washed' coals (containing say 10 to 15 p c water) are being coked. The design of the heating flues constitutes the main difference between the various types of oven; the Simon-Carvès, Semet-Solvay, Hussener, and Simplex containing horizontal flues, the Otto-Hilgenstock, Otto-Hoffmann and Koppers prefer vertical flues (Figs 4 and 5) In all cases the air required for the combustion of the gas in the heating flues is preheated either by passing it through recuperators in which it travels through a system of flues situated alongside the waste gas flues of the oven, or through regenerators similar to those employed in open-hearth steel furnaces, where the hot waste gases and air are alternately taken over the same surface (Fig 6). Where a large supply of surplus gas is desired from a coking plant, the regenerative method of preheating the air is to be preferred, since it is more effective and less gas is required for heating the flues; the amount of surplus gas available from a Lancashire coal containing 30 p.c. of volatile matter, may be taken at 2000 cubic feet per ton, where recuperative heating is employed,

and 5000 cubic feet per ton where regeneratives are used

The coal is generally charged into the oven through the crown from small hoppers, carrying the coal on rails from the storage bunker, the coal flowing into the oven being levelled by hand-rake or levelling machine In cases where the coal employed is not of a strong coking character, the fuel is stamped into a solid cartridge in a compressing machine and charged into the oven en bloc through the end doorway, The coke is discharged from the oven by a ram, generally electrically driven, often the levelling machine or compressor is built on the same carriage as the ram, and driven by the same The coke is received from the ovens on a sloping bench, where it is quenched with water, and from which it slides into railway trucks, furnace barrows, or a conveyor heating flues of each oven discharge into a common flue, which leads the waste gases in the case of recuperator types to steam boilers; with regenerator-ovens, the gases are led direct to the chimney stack The gases distilled from the coal collect above the level of the fuel and pass forward to the ascension pipe or stand pipe, which delivers the gas into the tar or hydraulic main, which receives the gases from the entire battery. A valve or damper is provided on each ascension pipe for shutting off each oven whilst the latter is being discharged and recharged In the tar or hydraulic main the heavier tars or pitches are deposited, and in the 'dry' types of main, it is necessary to pump the thinner tars made at a later stage of the cooling, through the main to clear the pitch and prevent stoppages.

In the older systems of by-product recovery, which are now being rapidly superseded by more 'direct' methods (q v), the gases are drawn from the main by an exhauster (either

FIG. 5A.—SECTION THROTCH SEMET-SOLVAY OVEN.





FIG. 5.—HORIZONTAL-ELUED COKE OVEN THE SEMET-SOLVAY, NON RECENERATIVE



steam or electrically driven), a Root's blower, or some form of aspirator, through coolers. These coolers are either plain pipes or cylinders offering a large amount of atmospheric cooling surface, or are water-cooled chambers, in these condensers, the bulk of the remaining tar, together with the greater proportion of the ammonia liquor, is precipitated. From the condensers, the gases pass into the exhauster or aspirator, which then delivers them at a pressure into a tar-fog extractor This vessel is employed to remove the last traces of tar

state of division, and which no amount of The tar extractor, of cooling would remove which the Pelouze Andouin and Livesay are types, operates by dividing the gas into numerous thin streams by means of very small gas passages, the agglomeration and removal of the tar particles being effected by the wire drawing and friction thus afforded.

From the tar catcher the gases pass into the ammonia scrubber, where the last of the ammonia not recovered by the condensers is removed by contact with weak ammonia liquor and water. which exist in the gas in an extremely fine | The types of ammonia scrubber generally em-




FIG. 7.—SHOWING ARRANGEMENT OF GAS COOLING, SCRUBBING OF SULPHATE OF AMMONIA PLANT, OLD METHOD.

- A, Oven B, Hydraulic main
- Condensers or coolers
- Tar catcher (Pelouze)
- Exhauster.
- Scrubber
- G, Return main to ovens
- Tar pump to store

- M, Liquor feed to still
- H, Liquor feed to scrubber N, Cooler for sulphate plant R, Lime feed to still.
  - waste gases
- J Tar pump to store

  K, Liquor pump to sulphate O, Superheater
  plant P, Purifiers for sulphate U, Mother liquor well
  plant waste gases V, Spent liquor outlet
  W. Spent liquor se
- - S, Saturator T, Sulphate draining table.


  - Spent liquor outlet Spent liquor settling
- W, Spent

ployed are similar to those employed at gas works, and are either of the tower or rotary In the tower scrubber, the gases pass upwards through a mass of coke or other open material, meeting a downward current of water or ammonia liquor pumped in at the top. In the rotary scrubber, the gas is caused to impinge on revolving brushes or circular wooden screens, which dip into the water or liquor in the lower half of the vessel, and pass upwards with wet surfaces into the upper half of the scrubber, where the gas is circulating, and solution of ammonia is attained. The gas then passes forward to the benzol scrubber, which is constructed on similar lines to that employed for ammonia removal, creosote or anthracene

heavy oil being employed for the solution and recovery of the benzol vapour.

The gas having been freed from its tar, ammonia liquor, and benzol, is then passed forward to the ovens, where it is fed into the heating flues, the surplus being passed forward to boilers, gas engines, or gas holder

The tars and ammonia liquors made in the hydraulic main, condensers, tar catchers, and ammonia scrubber are collected and run into separators or tanks, where the denser tar settles to the bottom, is withdrawn and despatched to the tar stores. The liquors which remain above the tar are run off to the sulphate of ammonia plant, where the liquor is pumped into overhead store tanks, whence it is allowed



to flow by gravity through a preheater (heated by the waste gases from the sulphate plant), and thence into a distilling column, where it flows downwards, meeting in its travel an upward current of steam, which liberates the free or combined ammonia. In the lower half of the still, milk of lime is mixed with the liquor for the liberation of the ammonia from the fixed salts; the whole of the ammonia, together with steam, sulphuretted hydrogen, carbonic acid, and hydrocyanic acid, pass upwards through the still and into a lead-lined vessel called the saturator, containing sulphuric acid. The ammonia combines with the acid to form ammonium sulphate, which, when the liquor in the bath becomes saturated, crystallises out and is removed either by copper scoops, a valve, or by a steam injector. The sulphuretted hydro-gen and other gases pass from the saturator to the preheater, where they serve to heat the liquor passing to the still; thence the gases travel to a condenser where they are cooled, and afterwards to purifiers where the sulphur-etted hydrogen is removed by oxide of iron The waste liquors, which should not contain more than 0 01 p.c ammonia, are run into the sewers or watercourses, and are a frequent source of trouble to the river boards. By the Otto-Hilgenstock method of by-product recovery, to be described later, the production of these waste liquors is entirely obviated.

The recovery of crude benzol from its solution in creosoto or anthracene oil is effected in a distilling column somewhat similar to that employed for the recovery of ammonia from liquor, the heating of the solution of benzol in the absorbent being effected by wet steam assisted by dry steam. The vapours, steam, and benzol pass from the top of the still to a water-cooled coil, where they are together condensed, the condensates being separated by reason of their different densities. The grade of crude benzol thus obtained varies with the class of absorbent oil employed; with creosote, 62 p.c. of the product distrils at 120°, with anthracene oil the benzol is richer, 80 p.c. distilling at 120°. For the rectification of crude benzol, the product is washed successively with strongest oil of vitriol, water, and caustic soda, and redistilled.

Direct Recovery Systems -Two successful attempts have been made to simplify the old system of treatment of coke-oven gases for the recovery of ammonia, viz the inventions of the Koppers and Otto-Hilgenstock Coke Oven Companies. Koppers treats his gases by the old system up to the point of the tar-fog extraction He then reheats his tar-free gas by passing it in pipes surrounded by the hot gases coming from the ovens in an apparatus called a heat exchanger, and then passes the heated gas direct into a sulphate saturator, into which is likewise passed the ammonia from the liquor made in the condensers, which is distilled in an ordinary ammonia still. In short, Koppers abolishes the usual ammonia scrubber and substitutes the sulphate saturator, thus making his sulphate direct on the gas main (Fig 9). By thus using the sulphuric acid in the saturator for scrubbing out the ammonia, Koppers dispenses with the water utilised in the old scrubber, thereby reducing the total amount of liquor produced, and required in the ammonia distillation.

A more extensive simplification is effected the invention of the Otto-Hilgenstock Co of Dahlhausen, who dispense entirely with the old condensing plant and remove the tar from

effecting an economy in the amount of steam | the gases by means of a tar spray at a temperature above the dew-point of the liquors tar-free gases then pass direct into a saturator, where the whole of the ammonia is caught by sulphuric acid. The gases issuing from the saturator still contain their liquors in steam




FIG. 9.—KOPPERS' BY-PRODUCT RECOVERY PROCESS.

- Hydraulic main
- Heat exchanger Condensers
- Exhauster
- F, Tar extractor
- G, Liquor pump to still. H, Feed tank to still
- Liming chamber
- , Ammonia still K, Saturator.
- Acid spray catch-box. Tar store tank
- N, Tar of liquor receiving tank.
- O, Liquoi tank
  P, Pump delivering tan to railway trucks.

form, and the whole passes forward to the ovenflues, and the offensive and troublesome waste liquors are thus got rid of The Otto system, by abolishing the condensing plant, liquor tanks, ammonia stills, lime mixers, and pumps, waste hquor tanks, &c, effects a great economy in ground space. It likewise saves nearly the whole of the steam required to distil the ammonia liquor made by the old condensing

process, and, by passing the whole of the ammonia through sulphuric acid, losses of ammonia, which are hable to occur in the old process of ammonia liquor distillation, are rendered impossible. The ammonium sulphate made by the Otto direct-recovery, tests from 25 to  $25\ 5$  p c of ammona, and contains only the merest traces of tar, viz  $0\ 05\text{--}0\ 1$  p c (Figs 10, 11, and 12).




FIG 10.—OTTO HILGENSTOCK BY-PRODUCT RECOVERY PROCESS.

- E, Exhauster
- A, Tar-spray
  B, Tar-spray receiving tank G, Acid spray catch-box
  C, Tar-spray pump.
  B, Tar-spray feed pump.
  E, Exhauster

  H, Mother liquor return
  pipe.
  E, Exhauster
  - K, Tar spray tank overflow L, Condensing tank


Tar store

M, Pump delivering condensings to saturator N, Pump delivering tar to railway trucks.

### MODERN METHODS FOR THE RECOVERY OF NAPHTHALENE AND BENZOLS.

Recovery of Naphthalene.—The gas from the saturator, which is now freed from both ammonia and tar, but still containing the benzol, moisture and naphthalene, passes through a lead-lined eatch pot (not shown in the diagram, but whose

situation prevents all fear of any acid spray being carried over from the saturator) and on to the naphthalene extractor and spray cooling tower, H (Fig. 13.) This tower is quite devoid of any filling or apparatus, but at the top there are several jets through which cold water is sprayed at high pressure. The action of this mass of



finely-divided cold water beating suddenly on to the hot gas produces a rapid chill, which, together with the concussion brought about by the water, precipitates and washes out practically all the naphthalene, which flows away along with the water into a catch tank beneath this apparatus. The naphthalene is filtered off whilst the water can be either allowed to flow away, or re-cooled and used over again continuously, as circumstances permit This tower also serves the purposes of cooling the gases. It is necessary to extract the naphthalene, otherwise it would pass on to the benzol scrubbers, where it would be dissolved by the benzol wash oil, which would very soon become saturated with naphthalene and rendered useless for the purpose of collecting benzol naphthalene thus extracted is of a pale yellow colour, and is readily sold in this country for 50s a ton, which more than pays the cost of

running this part of the plant.

Recovery of Benzols.—The gas now being cooled but still containing the benzol, passes through the exhauster, K, and a second water cooler, L, and thence to the benzol scrubbers, MM, which vary in number according to the size of the plant (Fig. 13). These scrubbers are from about 50 to 70 feet high, the lower portion forming a reservoir for the circulating benzol wash oil, and the remaining portion being filled with wooden grids. Benzol wash oil is constantly circulated downwards over these grids, the gas having to pass upwards through them By this means the gas is brought into vory intimate contact with the oil, with the resulting absorption of the benzol. After leaving these scrubbers the gas passes either wholly back to the ovens, or in the case of regenerative ovens about one-half of it returns to the ovens, and the other half is used either for firing boilers, town lighting, operating gas engines, or any other purpose


other purpose

A fuller account of the process is given in a paper by Mr. E Bury before the Institution of Mining Engineers in the year 1914, from which the following temperatures of a plant at New Brancepeth Colliery (Co Durham) are

taken:-

Hydraulic main		$rac{\mathrm{Cent}}{225^\circ}$	
Before tar extracts		86°	
After ,, ,,			58°
In saturator .			67°
After ,, .			58°
After exhauster			66°

Distillation and Rectification of Benzols—
The 'wash oil' containing the benzols in solution is sent on to the continuous crude benzol still, N (Fig. 14), where, after being pre-heated by the hot vapours coming away from the still, it passes downwards over a series of trays. Dry steam at about 100 lbs pressure, reduced, however, at the still to 10 lbs. pressure, is delivered into the bottom of the apparatus, and passing upwards bubbles through the wash oil liberating the benzol hydrocarbons which it carries over into the condenser, P. Here the steam and benzols are condensed, and the condensed water and oil separated. The crude benzol then either passes into a storage tank, or (if its rectification is to be carried out) into the 'primary rectifying still,' s. The debenzolised



'wash oil' leaving the crude benzol still is passed through the refrigerator, R, whence, after being cooled down to the atmospheric temperature, it is sent into the 'oil storage tank,' T, and from there it is used over again in the benzol scrubbers

The crude benzol is first of all fractionally distilled in the primary rectifying still, s, 3000-5000 gallons capacity and intermittent in its working, in which it is heated under high pressure by means of steam coils, and, with the aid of a long dephlegmating column, it is separated into—

(1) Crude 90 p c Benzol

(2) ", 90 ", Toluol

(3) ", 90 ", Xylol

(4) ,, 90 ,, Solvent naphtha

Each of these fractions is separately condensed and pumped into a large vessel, v, termed the

agitator, where it is successively treated with (a) pure strong sulphuric acid, and (b) a solution of caustic soda, in order to remove all resinous, basic, or acidic substances. Each of the 'washed' fractions is finally refractionated under reduced pressure in the 'secondary rectifying still,' w, and the redistilled resulting fractions, after condensation, are sent forward into their respective storage tanks, xyz, ready for transport

Other firms (e q the Simon-Carvés and the Simplex Companies) have also adopted efficient 'direct recovery' processes, on much the same lines as the Otto process just described, the chief difference between them being in the precise mechanical arrangement for the removal of tar and tar fog from the hot gas before it enters the saturator where the ammonia is absorbed.<sup>1</sup>

Alcohol from Coke Oven Gas —Attention





Fig 13.—Otto Direct Recovery Process (Benzol Recovers)

should here be drawn to the successful process worked out by E Bury and O. Ollander at the Skinningrove Ironworks for the absorption of the ethylene in coke oven gas by cone H<sub>2</sub>SO<sub>4</sub> (forming ethyl hydrogen sulphate), and its subsequent recovery therefrom as ethyl alcohol The process, which is fully described in the Coal and Iton Trades Review for December, 1919 (q v), consists essentially in (a) absorbing the C2H4 from the dried, debenzolised, and H<sub>2</sub>S-free gas by means of 95 pc H<sub>2</sub>SO<sub>4</sub> in a series of towers, on the counter-current principle, at a temperature between 60° and 80°C.; and subsequently (b) treating the strong acid containing the ethyl hydrogen sulphate with a current of steam in a special form of distilling column, so as to dilute it to between 70 and 75 p.c. strength, thereby also hydrolysing the ethyl hydrogen sulphate and liberating the resultant ethyl alcohol, which then distils over,

and is condensed. The whole process is worked in a continuous cycle, the diluted acid being (c) reconcentrated to 95 p c. strength in a Guilland 'concentration tower,' and then being reused for extracting ethylene from the gas. A notable feature of the process is that the small amount of  $\mathrm{SO}_2$  formed by the reducing action of the gas upon the hot conc.  $\mathrm{H_2SO}_4$  in the absorption towers is subsequently utilised to remove  $\mathrm{H_3S}$  from the gas by the well-known reaction  $2\mathrm{H_2S} + \mathrm{SO}_2 = 2\mathrm{H_2O} + 3\mathrm{S}$ , so that all the S contained as  $\mathrm{H_2S}$  in the gas is removed and recovered from it without the employment of elaborate 'iron-oxide' or other purifiers. From the gas (containing between 1.75 and 2.25 p c. ethylene) evolved in carbonising a typical Durham coking coal in regenerative ovens, with the 'direct'

<sup>1</sup> See a paper on Recent Developments in Ryproduct Coking, by Mr G. Blake Walker. (Proc. Inst. Civil Engineers, 1917.)

COLCHICUM.



system of recovering tars, naphthalene, ammonia, and benzol, it is claimed to be possible to obtain about 15 gallons of 95 p.c. ethyl alcohol per ton of coal at a present-day cost not exceeding one shilling per gallon.

Utilisation of Coke-Oven Gas. The composi-

tion of the surplus gas obtained in working the process depends somewhat on whether it is 'debenzolised' or not, as well as on the character of the coal carbonised, but for gas from a typical Durham coking coal it would be approximately as follows -

Gross calonific value B Th U per cub ft at N<sub>2</sub> 760 mm.

333

(1) Without benzol CO<sub>2</sub> CO  $C_nH_m$   $CH_4$   $H_2$   $N_2$  recovery 2 5 6 5 3 2 28 0 48 0 11 8 (2) 'Debenzo-

 $2\ 5\ 6\ 5\ 2\ 0\ 25\ 0\ 55\ 0\ 9\ 5\ 485$ lised?

With a regenerative oven giving 5000 cubic feet of such surplus 'debenzolised' gas per ton of coal carbonised, the corresponding heat units available for outside purposes would be 2,450,000 (gross) and 2,150,000 (net) B Th Us. respectively This gas may be utilised (a) for generating steam in a Bonecourt surface-combustion boiler with an efficiency of 92 pc: or (b) for generating power in internal combustion engines with an efficiency of from 20 to 25 p c Where coke-ovens are erected in conjunction with blast-furnaces and steel-works, the surplus gas is usually mixed with cleaned blast-furnace gas, and the mixture used both for generating power in gas-engines and also for heating purposes in open-hearth steel-furnaces, soaking pits, &c. In both Germany and America, cokeoven gas (after purification in the usual manner from H<sub>2</sub>S and eyanogen compounds) is extensively used as a public supply for domestic purposes As long ago as 1910, Parliament granted powers for the supply of gas from Loid Ellesmero's Brackley Coke-Works to the urban district of Little Hulton (Lanes), under a guarantee of a minimum illuminating power of 14 candles when tested in a No 2 Metropolitan Argand burner, and a gross calorific value of 550 B Th U per cubic foot at 60°F and 30 mm pressure Its average composition was CO2=2 5, CO = 7.0,  $C_nH_m = 3.8$ ,  $CH_4 = 31.0$ ,  $H_2 = 49.0$  and N<sub>2</sub>=67 pc More recently, the town of Middlesbrough now draws its entire gas supply from a battery of Otto regenerative ovens at the Newport Ironworks Also Leeds takes part of its supply from the Simon-Carvès ovens of the Middleton Estate and Colhery Co, Ltd · whilst at the Birmingham (Saltley) Gasworks a battery of 66 Kopper's regenerative ovens, fired by Mond gas, has been installed for the production of furnace coke and town's gas

Literature — W A Bone, Coal and its Scientific Uses, chap xvi 1918, and G Stanley Cooper, By-Product Coking, 1917; Report of Coal Conservation Committee, 1918, pp 79-81. W A B and E. B.

COLALIN. Syn for cholalic acid

C24H40O5,H2O

COLATANNIN v TANNINS. COLCHICEINE v Colchicum.

COLCHICUM. Meadow saffron. (Colchique, Fr.; Herbstzeitlose, Ger) Colchicum autumnale (Linn); the corms and the seeds are official in the BP. and U.SP., the alkaloid colchicinn only in the U.S.P. The plant is found in meadows and pastures throughout Europe, where its crocus-like flowers, usually of a lilac hue, occur in September and October.

334 COLCHICUM

The only constituent of importance is the alkaloid colchicine  $C_{22}H_{25}O_6N$ , which was regarded as identical with veratrine by Pelletier and Caventou as early as 1820, and was recognised as a separate individual by Geiger and Hesse (Annalen, 1833, 7, 274) Its constitution has in part been clucidated, chiefly by Zeisel (Monatsh 1883, 4, 162, 1886, 7, 557; 1888, 9, 1; Zeisel and Johanny, ibid. 1888, 9, 865; Zeisel and Stockert, ibid. 1913, 34, 1327, 1339), and more recently by Windaus (Chem. Soc Abstr. 1911, 1. 904; 1915, 1. 708) According to these investigations the formula may be resolved into:—

$$\begin{array}{c|c} \textbf{MeO} \\ \textbf{MeO} \\ \hline \\ \textbf{MeO} \\ \end{array} \begin{array}{c} \textbf{C}_{\textbf{10}}\textbf{H}_{\textbf{9}}\textbf{O}(\textbf{OMe)}\textbf{NH} \ \textbf{CO} \ \textbf{CH} \,, \end{array}$$

By boiling with 02 pc sulphuric acid one methoxy group is hydrolysed and there results colchiceine  $C_{21}H_{23}O_0N, 1H_2O$ , crystals, m p 139°, or anhydrous, m p 172°, which has been described, probably erroneously, as a constituent of the plant. Zeisel regards colchiceine as a carboxylic acid; Windaus, on account of the green colouration by ferric chloride, considers By methylation it is transformed it an enol back to colchieme, and to N-methylcolchieme. Concentrated hydrochloric acid hydrolyses colchicine further, removing the N-acetyl group and finally the remaining three methyl groups, yielding colchicinic acid C<sub>16</sub>H<sub>11</sub>NO(OH)<sub>3</sub>OH Windaus obtained, by permanganate oxidation of colchicine, 3 4 5-trimethoxy-o-phthalic and succinic acids, and by potash fusion of colchicinic acid terephthalic and trimellitic (1·2·4-benzenetricarboxylic) acids, which indicates that the complex C<sub>10</sub>H<sub>8</sub> of the above formula contains a (second) benzene ring Iodine and sodium hydroxide remove one carbon atom, oxidising to  $C_{20}H_{22}O_5NI$ , which may be reduced to a substance  $C_{20}H_{23}O_5N$  Colchicine apparently contains no heterocyclic nitrogen According to D R P 279999 of Hoffmann-La Roche and Co tetrahydrocolchicine is formed by reduction with hydrogen and palladium; this base is amorphous and less toxic than the parent alkaloid.

Properties—Pure dry colcheme usually forms a yellowish varnish, softening at 142°, and wholly melting at 147°; but Clewer, Green, and Tutin (Chem Soc Trans 1915, 107, 839) obtained from ethyl acctate soft, pale yellow needles, m.p 155°-157°; [a]<sub>D</sub>-121° (0 88 gram per 100 c.c) in chloroform, -429° in water. Zeisel first crystallised it from chloroform, with two molecules of chloroform of crystallisation; according to Merck (Apoth. Zeit. 1916, 31, 399), there are two such compounds, B,CHCl<sub>1</sub> and 2B,CHCl<sub>3</sub>. The crystalline colchicine of commerce is the latter compound, in which the 14-16 p.c. of chloroform is retained tenaciously, and is only given off slowly at 100°. According to Merck, pure colchicine, dissolved in 3 parts of water, slowly crystallises as the hydrate B<sub>2</sub>,3H<sub>2</sub>O. Colchicine has a very bitter taste. It dissolves slowly but abundantly in water, readily in alcohol, chloroform, and hot benzene, hardly at all in ether. Being an acid amide, it is a very feeble base,

and one of the few alkaloids which can be extracted from acid solution by chloroform. Its salts are generally hydrolysed by water; the only crystalline one appears to be the aurichloride B, HAuCl<sub>4</sub>, mp. 209°. Dilute mineral acids and alkalis colour colchiene solutions intense yellow Nitric acid (d. 14) colours the solid alkaloid a deep violet, changing to yellow, and finally to green (characteristic) Tannic acid, phosphomolybdic acid, potassium triiodide, and bismuth potassium iodide are the most sensitive precipitants; potassium mercuric iodide only precipitates in acid solution.

Extraction, Toxicological Recognition, Assay. —These are all based on the removal from acid solution by chloroform To prepare colchicine, uncrushed seeds are exhausted with 85-90 p c. The extract is evaporated and diluted with much water, filtered from fat, and extracted with chloroform, The chloroform solution is evaporated to the consistency of olive oil and diluted warm with small quantities of alcohol, until the whitish masses which separate at first again dissolve. After cooling to 0° for some time yellow needles of the chloroform compound separate, which may be suspended in very little hot water and decomposed by steam. The aqueous solution is then evaporated in vacuo On a smaller scale, eg in the toxicological recognition, the acid solution is first agitated with petrol and then with chloroform residue, after evaporation of the latter solvent, is dissolved in water, and precipitated with tannic acid. The precipitate is decomposed with lead oxide and extracted with alcohol or chloroform On evaporation the solid product so purified can be tested by the reactions given above; 1 mg will give the nitric acid test (see also Allen's Commercial Organic Analysis,

1913, vol vn pp 4-12)
The assay of colchicum seeds and the estimation of colchicine is based on weighing the residue from a chloroform extract of an acid solution, previously washed with petrol (Farr and Wright, Pharm. J. 1891, [iii.] 21, 957; for the assay of corms, v. Lyons, ibid. 1909. [iv.] 28, 270, v also the mere complicated process of the U.S.P.)

The alkaloidal content of the seeds and corms varies greatly, mostly from 0 2 to 0.8 p c. in the seeds, from 0.1 to 0.5 pc in the corms. The Brussels Conference agreed that the seeds The alkaloid occurs to a only should be used slight extent also in the flowers. It is present in other species of the same genus, and also in the large tubers of Gloriosa superba, L. (NO. Lihaceæ), common in India and Ceylon (Clewer, Green, and Tutin, lc, found 03 p.c. of dry weight) Perhaps this plant would afford a suitable commercial source of colchicine. Colchicum is a specific in acute gout, and an ingredient of many advertised gout remedies. It is gastro-intestinal irritant, and may cause vomiting and purging. Its degradation and substitution products are much less active or without action (Fuhner, Arch. exp Path. Pharm 1913, 72, 228). G. B. Pharm 1913, 72, 228).

**COLCOTHAR.** The reddish-brown residue of ferric oxide formerly obtained in the manufacture of fuming oil of vitriol from iron sulphate.

COLEMANITE. A hydrated calcium borate Ca<sub>2</sub>B<sub>6</sub>O<sub>11</sub>,5H<sub>2</sub>O, containing 50 9 p.c. boron

trioxide It is found in California at Death Valley in Inyo Co., and the Calico district in San Bernardino Co; and although not discovered until 1882 and 1883, is now of some importance as a material for the preparation of borax and boracic acid Cavities in the massive, white material are lined with beautiful, water-clear crystals with a profusion of brilliant facets; these are monoclinic, with a perfect, pearly cleavage parallel to the plane of symmetry They are considerably harder (H 4-4½) than borax, sp gr 242. The mineral forms, together with other borates and borosilicates, a bed 7 to 10 feet in thickness in sandstones and clays

The compact minerals, priceite from Oregon and pandermite from Asia Minor, have very nearly the same composition (Ca<sub>5</sub>B<sub>12</sub>O<sub>23</sub>,9H<sub>2</sub>O), and they have been regarded as impure, massive varieties of colemante. E. S. Larsen (1917) has, however, by an examination of the optical characters established the identity of priceite and pandermite, and proved them to be triclinic and distinct from colemanite. These also are mined as borates v Boron. L. J. S.

COLLIDINE v. Bone oil

**COLLIN.** A preparation made by heating a solution of hide powder or gelatin with caustic soda and neutralising with acetic acid. Used in the analysis of tannin and tanning materials (Parker and Payne, J. Soc Chem. Ind. 1904, 648; Wood and Trotman, *ibid.* 1904, 1071)

**COLLODION** v PHOTOGRAPHY **COLLODION COTTON** v. EXPLOSIVES.

COLLOIDS. In 1861 Graham pointed out that substances which crystallise readily from water are characterised by a high power of diffusion and by the ability to pass through animal or vegetable membranes, substances, on the other hand, which cannot easily be obtained in the crystallised condition are characterised by a low power of diffusion and by mability to pass through animal or vegetable membranes. Graham termed substances of the first class, such as sucrose or sodium chloride, crystalloids; and those of the second class, such as starch, gum, or albumin, colloids The separation of a colloid from a crystalloid is effected by dialysis, a process based on the inability of a colloid to pass through parchment or other membrane

At the present time, the term 'colloid' is used in a sense different from that in which Graham employed it, and is to be interpreted as referring, not so much to a particular class of substances, as to a condition which very many, if not all, chemical elements or compounds may be made to assume From the modern standpoint 'colloidal solutions' are to be regarded as forming a special group of the physical systems built up from two substances, one of which is present in relatively small quantity. Such systems may be described as 'dispersed systems,' the substance present in excess being termed the 'dispersion medium,' and the other the 'dispersed phase.' The main groups of dispersed systems are: (1) mechanical suspensions; (2) colloidal solutions; (3) molecular solutions, the feature which characterises each group being the degree of dispersion or division of the dispersed phase From this point of view a colloid solution is a

dispersed system in which the degree of dispersion is intermediate between that of a mechanical suspension and that of a molecular solution.

Now the smallest suspended particle which can be observed directly under the microscope is of the order 0  $1\mu$ , and this size of particle may be taken as roughly marking the boundary between mechanical suspensions and colloidal On the other hand, the size of an average molecule is 0 0001 µ-0 001 µ, and although there is no haid-and-fast line of demarcation to be drawn between the three groups of dispersed systems, the dispersed phase of a colloidal solution may conveniently be regarded as consisting of particles of the dimension  $0.1\mu$ -0  $001\mu$ This conception of the continuity of dispersed systems, from mechanical suspensions at one extreme to molecular solutions at the other, finds its justification in the graded character of the physical properties of the three groups, as evidenced, for example, by observations on diffusion and filtration, and by osmotic, optical, and electrical experiments (see below). It should further be observed that the colloid condition is independent, not only of the chemical character of dispersion medium and dispersed phase, but also of their states of aggregation whilst in many colloidal solutions, such as those of platinum, arsenious sulphide or ferric hydroxide, the dispersion medium is a liquid, and the dispersed phase is a solid, there are other cases, as, for example, an aqueous solution of gelatin, in which the dispersed phase also is a liquid. The terms 'suspension colloid' and 'emulsion colloid' are commonly used to differentiate the Again, there appear to be many cases in which both the dispersion medium and the dispersed phase are solids, instances of this kind being furnished by ruby glass, some coloured minerals, and probably also alloys in certain conditions

Osmotic properties of colloidal solutions — The osmotic pressure of a colloidal solution is extremely low According to Duclaux, for instance, the osmotic pressure of an 886 pc ferric hydroxide solution is only 22.6 cm. of water, which is about equal to the pressure given by a solution of sucrose containing onethirtieth of a gram in 100 grams of water. It has, indeed, been suggested that the osmotic pressures which have been observed for colloidal solutions are due, not to the colloids themselves, but to small quantities of electrolytes associated with the colloids While it is certainly true that the removal of the last traces of electrolytes from a colloidal solution is no easy matter, and that electrolytes are extremely active material from the osmotic point of view, still the most recent investigations indicate that colloidal solutions do possess a definite, if small, osmotic pressure In this connection the epoch-making work of Perrin is of great significance, for his determination of the Avogadro number, based on a study of the distribution of suspended gamboge particles in a vertical column, really involves the extension of the gas laws to mechanical suspensions. This extension is justified by the experimental results (see Perrin's book, Atoms) and it is therefore to be expected that the osmotic pressure of a colloidal solution will be related in the usual way to concentration and temperature.

It appears, however, that the osmotic pressure of a colloidal solution may not be in every case completely defined, as is that of a crystalloidal solution, by concentration and temperature. Thus Moore and Roaf have found (Biochem J 1906, 2, 34) that the osmotic pressure of a given gelatin solution at a given temperature depends, to some extent, on its previous history, while Lillie has found (Amer. J. Physiol 1907, 20, 127, see also Bayliss, Proc Roy Soc B, 1907, 81, 269) that the osmotic activity exhibited by the colloid itself is modified by the presence of electrolytes in a remarkable degree. The osmotic pressure of a colloidal solution, it is clear, will be determined not only by the total solute concentration and by the temperature, but also by the actual degree of dispersion

The depression of the freezing-point and the elevation of the boiling-point of water by a dissolved substance are quantitatively related to the osmotic pressure of the solution, and, like the latter, may be made the basis for a determination of the molecular weight of the solute freezing-point and boiling-point of a colloidal solution are, however, very slightly different from those of water itself, and the experimental error of the determination of the molecular weight of the colloid by this method is relatively In view, further, of the fact that the osmotic activity of a colloid depends on other factors than those of concentration and temperature, it is obvious that the values for the molecular weight of a colloid deduced from its effect on the freezing-point or boiling-point of water have a very limited significance values have reference only to the particular condition of the colloid at the time of the determination

In connection with the interesting question of the osmotic activity of colloids, attention should be directed to the investigations of McBain and others on soap solutions (Chem Soc Trans 1919, 115, 1279)

Colloids in an electric field -In 1892 Linder and Picton observed (Chem Soc Trans 1892. 61, 148) that when two wires, connected with the terminals of a battery, are placed in a colloidal solution of arsenious sulphide, this substance is attracted by the positive pole and is gradually transported thither Ferric hydroxide in colloidal solution is, on the other hand, attracted by the negative pole It appears, therefore, that the particles of colloidal arsenious sulphide carry a negative charge, whilst those of colloidal ferric hydroxide carry a positive charge Other colloids are similarly found to carry a definite charge, thus chromium and aluminium hydroxides, methylene blue, Bismarck brown and hæmoglobin, are electropositive; tannin, caramel, starch, platinum, gold, and indigo are electronegative Very significant in this conelectronegative nection is Hardy's observation (J Physiol 1899, 24, 288), that protein is electronegative in an alkaline medium, but electropositive in an acid medium. A reversal of the electric charge on a colloid has been observed in other cases also (see Burton, Phil. Mag. 1906, 12, 472). The (see Burton, Phil. Mag. 1906, 12, 472). behaviour of substances in colloidal solution, when exposed to the action of an electric force is very similar to the behaviour of suspensions in the same circumstances, for when wires connected with the terminals of a battery are immersed in suspensions of quartz powder, gum mastic, or shellac, the suspended particles move towards the positive pole.

Colloids and the ultramicroscope.—The view that there must be a close relationship between colloidal solutions, on the one hand, and mechanical suspensions on the other, is strongly supported by various other lines of evidence. Thus the majority of colloidal solutions exhibit the Tyndall phenomenon, i.e. the opalescence observed when a powerful beam of light is passed through a fluid medium containing definite suspended particles. Further, the ultramicroscope, devised by Zsigmondy and Siedentopf, makes it possible to detect the individual particles in colloidal solutions, even when the most powerful microscope fails to reveal any trace of heterogeneity.

There are, however, cases on record in which colloidal solutions, even when examined with the ultramicroscope, have been found free from distinct particles. On the other hand, evidence has been adduced (Lobry de Bruyn, Rec. trav. chim 1904, 23, 155, 218) showing that solutions of a heavy crystalloid, such as sucrose, are non-homogeneous. Hence it is evidently impossible to draw a hard-and-fast line of division between colloidal and crystalloidal solutions, so far at least as the size of the particles of the dispersed phase is concerned.

The particles which are detected in a colloidal solution with the help of the ultramicroscope are seen to be in a state of constant motion—a phenomenon essentially the same as the Brownian movement exhibited by gamboge or fat particles suspended in water. The finer the particles the more rapid is their movement, and a study of the relationship between the size of the particle and its rate of movement has led to a calculation of the rate at which a particle of molecular size would move. The value so calculated is in good agreement with the value deduced from the kinetic theory, and this agreement is regarded as amounting to a definite proof of the kinetic nature of heat.

Substances in colloidal solution pass through the finest filter paper, but if the close relationship between mechanical suspensions colloidal solutions is borne in mind, it will be seen that the question whether filtration can be effected in the latter case as in the former reduces itself to the question whether filters with sufficiently small pores can be procured. The most successful attempts that have been made to solve this problem depend on the impregnation of a suitable supporting medium with a colloid such as gelatin or collodion. Filter paper, for instance, may be soaked in gelatin solution and then hardened with formaldehyde, or impregnated with a solution of collodion in glacial acetic acid and then dipped in water As the concentration of the impregnating gelatin solution is increased, the permeability of the filter to colloids diminishes, so that a series of graded filters can be prepared which may be employed to sort out a number of colloidal solutions, according to the size of particles they contain (see Bechhold, Zeitsch. physikal. Chem. 1907, 60, 257). It is noteworthy that these 'ultra-filters,' as they are called, may be

employed to separate oil-water emulsions (see Hatschek, J. Soc. Chem. Ind. 1910, 29, 125).

Another method of separating a colloid from its solution is to precipitate the colloid and then filter off in the ordinary way. It is found that in relation to precipitating or coagulating agents, solutions of suspension colloids behave differently from solutions of emulsion colloids. The former give non-viscous solutions which are coagulated by small quantities of electrolytes, these colloids are known also as 'non-reversible,' for when once they have been separated they do not again pass into solution by mere contact with water; their solutions can be obtained only by indirect methods. Emulsion colloids, on the other hand, give viscous solutions which are not readily coagulated by electrolytes, they are also termed 'reversible' colloids, because, even if separated from solution by any means, they will again pass into solution on mere contact with water.

The coagulation of suspension colloids, such as arsenious sulphide or ferric hydroxide, is brought about by extremely small quantities of electrolytes. When experiments are made to determine the minimum concentration of various electrolytes required to precipitate colloidal arsenious sulphide, it is found that this minimum concentration is of the same order for all salts containing cations of equal valency higher the valency of the cation, the smaller is the concentration of the electrolyte required to bring about coagulation In relation to colloidal ferric hydroxide, on the other hand, the coagulating power of a salt is mainly determined by the valency of the negative ion; the valency of the cation is relatively unimportant The contrast in this respect between colloidal arsemous sulphide and colloidal ferric hydroxide is very interesting in view of the fact that the colloid particles of arsenious sulphide are negatively charged, while those of ferric hydroxide are positively charged. The full significance of this was first emphasised by Hardy (Zeitsch physikal. Chem. 1900, 33, 385), who formulated the rule that the ion of an electrolyte which determines the coagulation of a colloidal solution is the one which has a charge opposite in sign to that on the colloid particles. The validity of this rule has been extensively confirmed.

The 'coagulum' or 'hydrogel' obtained when a solution of a suspension colloid is precipitated by a salt contains some of the acidic or basic part of the salt. Thus the coagulum obtained when colloidal arsemous sulphide is precipitated with barium chloride, is found to contain barium, and in proportion as the coagulum retains barium, the filtrate becomes acid. The barium cannot be removed by continued washing with water, but may be replaced by an equivalent quantity of another metal by digesting the coagulum with a colution of the chloride of this metal

If, as suggested by the facts just mentioned, the coagulation of a suspension colloid by a salt is essentially a neutralisation of electric charges, it may be expected that colloids of opposite electrical sign will precipitate each other. In harmony with this, it is found that while no coagulation occurs when colloidal solutions ('hydrosols') of the same electrical sign are mixed, precipitation immediately follows the

addition of the solution of a positive colloid to the solution of a negative colloid, unless one of the colloids is in very large excess. For a certain proportion of the colloids, the precipitation of both is complete, but as the quantities deviate from this optimal ratio, precipitation is increasingly incomplete. The precipitation of egg albumin by solutions of various complex acids—e g molybdic, tungstic, and tannic acids—furnishes an example of the mutual coagulation of two colloids. Metaphosphoric acid, too, forms a pseudo-solution or hydrosol which precipitates albumin, while the crystalloidal orthophosphoric acid has no such effect.

The separation of emulsion colloids, such as serum protein, from their solutions, may also be effected by the addition of electrolytes, but the process is essentially distinct from the corresponding precipitation of suspension colloids; it is not electrical in character When a number of alkalı salts are arranged according to their efficiency in precipitating emulsion colloids, the order is practically the same as when they are airanged according to their efficiency in salting out an organic compound from water or according to their power of lowering the solubility of gases in water. The precipitation of emulsion or reversible colloids by neutral alkalı salts appears, therefore, to be closely allied to the process of 'salting out,' and this result of itself shows that solutions of reversible colloids approximate more closely to true solutions than do solutions of suspension colloids

Not only are emulsion colloids themselves very slightly sensitive to electrolytes, but in presence of an emulsion colloid the precipitation of a suspension colloid by salts is more or less inhibited. This is not due to an increase in the viscosity of the medium and consequent increase of resistance to sedimentation, for the protective effect is exhibited by quantities of the reversible colloid insufficient to cause any appreciable change in viscosity. It is probable that the emulsion colloid forms a thin envelope round each particle of the suspension colloid and so prevents the aggregation and subsequent

flocculation of the particles.

Adsorption - Reference has already been made to a difference of electrical potential as determining in some cases the relation between a colloid and its environment There is, however, another factor which must be taken into account in the consideration of this question. and that is the relatively enormous surface of contact between the colloid and the surrounding medium. The evidence sketched in the foregoing paragraphs shows that a colloidal solution is really a two-phase system, consisting of a dispersion medium in which are distinct suspended particles of extremely small size Now, it is obvious that as a given quantity of matter is divided up more and more finely, its surface area increases regularly. The surface area, for instance, of a single sphere with a diameter of 1 mm is 0 0314 sq cm.; if the matter composing this sphere were broken up into a number of smaller spheres, each 0 0001 mm in diameter, the total surface area of the spheres would be 314 sq. cm. It is evident, therefore, that in a colloidal solution with its fine suspended particles, there must be relatively a very large surface of contact between the colloid and its medium.

Vol.  $\Pi - T$ .

On the basis of the two-phase character of colloidal solutions and the large surface development of the colloid phase, the relation of colloids to each other and to various solid and dissolved substances may be regarded as being essentially of the nature of 'adsorption.' This term is used in reference to the observation that when a substance is introduced into a two-phase system, and provision is made for its distribution throughout the system, it is frequently found concentrated at the surface of one of the phases. The nature of this phenomenon will perhaps be best appreciated by contrasting it with cases in which the substance is distributed uniformly in each phase of the two-phase system. When iodine is shaken up with two immiscible liquids, such as water and carbon tetrachloride, a state of equilibrium is soon reached at which the concentration of the iodine is uniform throughout the aqueous phase, uniform also (but different) throughout the carbon tetrachloride phase If  $c_i$  is the concentration of iodine in the aqueous layer. and  $c_2$  its concentration in the carbon tetra-chloride layer, then  $c_2/c_1$ , the 'partition coefficient, is found to be a constant, independent of the absolute values of  $c_1$  and  $c_2$ . This is a rule applicable to the distribution of any substance between two immiscible liquids, provided the molecular condition of the substance is the same in each liquid If, on the other hand, the molecular weight of the substance in the first liquid is n times its molecular weight in the second liquid, then  $c_2^{n}/c_1$  is a constant, independent of the absolute values of  $c_1$  and  $c_2$ versely, the distribution of a substance between two liquids at various concentrations has been found to be such that  $c_1^{n}/c_1$  is a constant, the conclusion may be drawn that the molecular weight of the substance in the first liquid is ntimes its molecular weight in the second liquid

The distribution of a substance in a twophase system which consists, not of two immiscible liquids, but of a solid and a liquid, is generally quite different in character. for instance, acetic acid solutions of various concentrations are shaken up with earbon, a state of equilibrium is rapidly attained, and it is found that some of the acetic acid has been taken up by the carbon If  $c_s$  and  $c_l$  represent the equilibrium concentrations of the acetic acid in the solid and the liquid phase respectively, then the experiments show that  $c_s^{\ 2\ 35}/c_l$  is a constant According to the foregoing paragraph, this would mean, on the assumption of a uniform distribution of the acetic acid throughout the carbon, that the molecular weight of acetic acid in the carbon must be less than half what it is This conclusion must obviously be rejected, and the assumption of a uniform distribution of the acetic acid throughout the carbon is incorrect. The acetic acid must be concentrated at the surface of the carbon, a view supported by the rapidity with which equilibrium is established The acetic acid is said to be 'adsorbed' by the carbon, and the phenomenon is termed 'adsorption.'

The distribution of a substance in a twophase system where one of the phases is a solid with largely developed surface, is generally found

to consist in an adsorption equilibrium. This is indicated, qualitatively, by the concentration of the substance in the solid phase increasing more slowly than its concentration in the liquid phase, and quantitatively by the validity of the formula  $c_8^n/c_l = \text{constant}$ . In this way it has been shown, to quote one or two examples, that the taking up of gases by charcoal, the removal of arsenious acid from solution by freshly precipitated ferric hydroxide, and the dyeing of silk with pieric acid and numerous other colouring matters, are all cases of adsorption. The complexes formed, although not due to chemical combination in the strict sense, and although lacking the characteristics associated with a definite chemical substance, are frequently referred to as 'adsorption compounds' As van Bemmelen especially has shown, colloids in the form of hydrogels have a very marked power of adsorbing other colloids, crystalloids and water, and evidence is accumulating in favour of the view that the behaviour of colloids in solution towards other substances is also to be interpreted in terms of adsorption.

The question may be asked: Is there any physical reason for the concentration of a substance at the surface of one of the phases in a two-phase system, such as a colloidal solution undoubtedly is? The answer is that if surface tension considerations only are taken into account, then it follows on thermodynamical grounds that the most stable relation between the two phases of a colloidal solution is the one characterised by a minimum surface tension Hence if the substance present in the solution lowers the surface tension of the solvent, that substance will accumulate in the surface layer of the solution That such spontaneous accumulations may occur in the surface layers of solutions has been shown by Ramsden (Proc Roy Soc A, 1903, 72, 156), who observed the formation of solid or highly viscous coatings on the free surfaces of protein and many other colloidal solutions

Bearing of colloid chemistry on technical problems.—The  $\imath \delta lc$  played by colloids in the operations of applied chemistry is of great importance In the manufacture of starch, cellulose, glue, leather, caoutchouc, artificial silk, paper, and photographic plates, colloids are involved, and a knowledge of the general behaviour of colloids throws much light on these and other industrial operations Many dyes give colloidal solutions, and dyeing, in a large number of cases at any rate, depends on the adsorption of the colouring matter by the fibre The relation between hide and tannin appears to be of the same description, and the investigation of the tanning process from this point of view shows the definite bearing which colloid and adsorption phenomena have on this opera-The problem of sewage disposal also is closely related to colloid chemistry, and it has been found practicable to retain the colloids of the sewage in the form of adsorption compounds. The tints of coloured glasses are frequently due to the presence of metals in the colloidal state. This is notably the case with gold ruby glass, which owes its colour to the presence of ultramicroscopic particles of the metal. Colloid chemistry has an important bearing, too, on

agricultural chemistry. Many of the properties of soils—for example, permeability, absorptive power, and moisture content-depend more on the size of the constituent particles than on their chemical composition The retention by the soil particles of materials of nutritive value for plants is closely related to the phenomenon of adsorption already discussed, and it has, in fact, been shown that the removal of dissolved substances by soil from solutions of different concentrations takes place in harmony with the adsorption formula. Colloids are of importance further in connection with the pottery trade Highly plastic clays behave like colloids, and the treatment of clay with a colloid, such as tannin, notably increases the plasticity of the material. On the view that the plasticity of a clay is determined mainly by its colloid content, and that the latter is chiefly responsible for the power of the clay to adsorb colouring matters, attempts have been made to utilise the adsorptive power of clays, say, for malachite green, as a measure of relative plasticity

Colloid clay has been found to be capable of use as a catalyst in process of oxidation and

hydrogenation

Of great interest are recent attempts to interpret the properties of many alloys in terms of colloid chemistry. The view is taken that the technical and physico-chemical properties of an alloy are largely dependent on the state of division of the constituent particles, and that in many cases the size of these particles is such as to bring them within the colloid range. The relation between tempering carbon and hardening carbon, is on these lines described as a 'dispersion allotropy' (For development of this point of view, see Benedicks, Journ Iron and Steel Inst. 1905, 352., Koll. Zeitschrift, 1910, 7, 290.)

Reference has been made above to the protective action of emulsion colloids, and this phenomenon is of frequent occurrence presence of such colloids, generally and vaguely referred to as 'organic matter,' may altogether prevent the flocculation and precipitation of insoluble compounds In presence of gelatin, for example, silver nitrate gives with sodium chloride only an opalescence which is not altered by filtration This is a fact of the greatest importance in connection with photographic emulsions (losely related to this phenomenon is the use of glues and gums as retarders in the setting of plaster of Paris, and the inclusion of such materials in mixtures which have for their object the prevention of scale in boilers Detailed references to the bearing of colloid chemistry on technical problems will be found m a monograph, Die Bedeutung der Kolloide für die Technik (Dresden, 1909), by K Arndt; in a paper on Colloid Chemistry and some of its Technical Aspects, by J. Alexander (J. Soc Chem Ind 1909, 28, 280), in W. W. Taylor's book on The Chemistry of Colloids, 1915; and in Wo. Ostwald's volume, Theoretical and Applied Chemistry, 1917. J. C. P.

collophanite. A colloidal hydrated phosphate and carbonate of calcium forming an important constituent of phosphorites and rockphosphates. It is amorphous and optically isotropic, and forms concretionary masses resembling opal or gum in appearance, with a

vitreous to resinous lustre and a conchoidal fracture. The material may be opaque or translucent to nearly transparent, and the colour ranges through colourless, white, yellow, brown, and reddish. An earthy variety is known as monte. Sp gr 2 69-2 77 (of monte 2 1), haidness 4½, refractive index 1 569-1612 A Lacioix (1910) gives the formula as:—

$$\begin{array}{l} (a[({\rm Ga_3(PO_4)_2}] + y({\rm GaCO_3} + z{\rm H_2O}) \\ x[({\rm GaF})_2({\rm Ga_8(PO_4)_6}] + y({\rm GaCO_3} + z{\rm H_2O}) \end{array}$$

which includes also the variety called fluocollophanite W T Schaller (1911) gives the formula as  $9 \text{CaO} 3P_2 O_5 \text{CaO} 2O_2 H_2 O_7 + nH_2 O$  The immeral is found in three classes of deposits; in guano deposits, having been first recognised in the guano of Sombreio, West Indies, in pockets of phosphorites, and in the phosphatic nodules and coprolites of sedimentary rocks tanging in age from the Eocene to the Siluian. Material of the last of these modes of occurrence frequently contains fluorine in addition to carbonate and phosphate, and is distinguished as fluorollophanite L J S

COLLOTYPE v. PHOTOGRAPHY.

COLOCYNTH. Bitter apple Colocynthidis Pulpa, BP, Colocynthis, USP. (Coloquinte, Fr; Koloquinte, Ger)

Fr; Koloquinte, Ger)
The dried pulp of the fruit, freed from seeds, of Citrullus Colocynthis (Schrader), a chimbing plant of the gourd family, inhabiting the warm dry districts of the Old World—India, Persia, Arabia, Syria, and the African and European shores of the Mediterianean. It is cultivated in Spania and Cyprus. (Cf. Bentl. a. Tum. 114.)

m Spain and Cyprus (Cf Bentl a Tim 114)
The BP, and USP demand that not more than 2 pc, of fixed oil be extracted by petrolcum spirit (r. Dowzard, Phaim J. 1903, 71, 400)
The B.P. demands at least 9 pc. of ash, the USP not more than 15 pc.

Chattaway and Mooi (Analyst, 1903, 28, 205) consider 8 pc a more reasonable minimum, Umney (Pharm J. 1903, 71, 879) found the ash to vary from 72-135 pc, and thinks a microscopical examination and the estimation of fixed oil better ways of detecting impurities. The seeds contain the oil, and only 2-3 pc. of ash, as they constitute 75 pc of the peeled fruit they raise, when mixed with pulp, the amount of fixed oil, and lower the ash content. The two chief varieties imported are Turkey and Spanish colocynth, of which the former is the whiter, the richer in pulp, and the more esteemed

Colocynth was well known to Greek, Roman, and Arabian medicine, and in the form of solid extract enters into many of the purgative pills of modern pharmacy.

Power and Moore, who gave an account of the earlier literature (Chem Soc Trans 1910, 97, 99), have cast doubt on the existence of the colorynthin and colorynthin of previous investigators. The former substance was described as an extremely bitter glucoside and drastic purgative, soluble in water and alcohol, but not in ether, the latter substance as insoluble in water, but soluble in ether Power and Moore examined the pulp of the fruit separately from the seeds, and the following substances were isolated · citrullol, a dihydric alcohol C<sub>22</sub>H<sub>\*</sub>, O<sub>2</sub>(OH)<sub>2</sub>, m.p. 285°-290°, yielding a

diacetyl derivative, melting at 167°; an amorphous alkaloidal substance possessing an extremely bitter taste and representing one of the purgative principles of the fruit; a-elaterin (see ELATERIUM); a small amount of hentriacontane C<sub>31</sub>H<sub>64</sub>, m.p 68°; a phytosterol C<sub>27</sub>H<sub>46</sub>O, and a mixture of fatty acids. The seeds, which represent 755 p.c. of the peeled fruit, were found to contain traces of an alkaloidal principle, a small amount of an enzyme hydrolysing \(\beta\)-glucosides, and fatty oil proportion of glucosidal substance in the fruit is found to be very small. The purgative action depends on at least two compounds, one of which is alkaloidal. After removal of the alkaloidal compound, the fruit still possesses marked purgative properties, but the substance to which these are due has not yet been obtained in a definite form

It has been sometimes required to search for colocynth in beer (v Dragendorff, Chem Zentr 1881, 285, 299). G. B.

COLOGNE YELLOW v Chromium.

COLOPHON v OLEO-RESINS.

COLOPHONY v RESINS

COLOPTENE v RESINS

CHROMOMETERS. COLORIMETERS. **CHROMATOMETERS** Under this heading are included those instruments (colorimeters) which are used for determining the relative intensity of colour in solutions, and those (chromatometers, tintometers) which are used to obtain a numerical record of the hue and intensity of a colour.

Colorimetric measurements depend on obtaining equal intensity of colour in two columns of liquid, one of which contains a known amount of the substance to be estimated This can be done either (a) by varying the concentration of one liquid, keeping the lengths of columns equal, or (l) by altering the length of one column,

keeping the concentration the same

The first method is used in Nessler's test, in Eggertz' method of estimating 'carbon in steel, and in the simple colorimeter of the US Geological Survey (Bulletin 422) It is ap-Tĥe plicable to all colorimetric measurements second principle is adopted in most cases because it allows the repeated use of the standard solution and facilitates rapid work In certain cases, however, eq. in comparing manganese solutions, this method fails, as varying the length of column changes the hue as well as the intensity of the colour.

The standard used may be of the same composition as the liquid under examination, or may be made of any suitable materials, but its colour must be identical in quality with that of the liquid tested, and its value per unit volume in terms of the substance estimated must be accurately known Thus the permanent standards used in the estimation of combined carbon in steel are generally made of mixed solutions of metallic salts, and in the estimation of ammonia in water a solution coloured with caramel and a little aniline red has been used (Leeds, Chem. News, 37, 229)

Equality of tint having been obtained, the lengths of the columns are noted, and the comparison is repeated, using, preferably, a different length of column of the standard.

The relative strength of the solutions varies inversely as the lengths of the columns, so that if 1 cc. of the standard solution correspond with x parts by weight of the substance estimated, the weight of that substance in 1 c c of the test solution will be

length of column of standard solution  $\times x$ length of column of test solution

If the strength of the standard differs by more than 10 pc. from that of the test liquid, it is advisable to dilute one or other before comparison

For the examination of precipitates, a solution of considerable suspensory power is prepared by dissolving 100 grams of gelatine, 100 grams glacial acetic acid, and 1 gram of salicylic acid (to prevent decomposition of the gelatine by

keeping) in 1 litre of distilled water.

The solution is clarified with white of egg and filtered hot For estimation of sulphuric acid, this solution is charged with barium chloride, and is added to a weak standard solution of sulphuric acid or a sulphate and to the solution to be tested, the two turbid liquids being then compared as above By the use of other reagents, lime, vinc, and a large number of other substances may be estimated.

A useful colorimeter for general purposes, devised by Mills, is shown in Fig. 1. It con-

sists of a glass jar closed above by a cap of which the length is sufficient to cover the top of the liquid The cap is perforated in middle and carries a short tube in which slides a glass rod bent as shown, and supporting, exactly at right angles to the axis of the jar, a flat circular opaque white In the bottom of the jar are laid, one above another, a red and a green glass disc which form a black background to the white disc When examining solutions of certain colours, a background of a colour other than black may be advantageous, and the white disc itself may be covered by a coloured plate The white disc is lowered to the bottom of the par by moving its rod, and the jar is filled to exactly 100 divisions with a standard liquid.

The white disc is raised until, on looking vertically down through the liquid at the disc, the depth of tint appears suitable. A second apparatus is then filled to 100





Fig 1.

divisions with the liquid to be tested and its disc is adjusted until the colours of the columns of liquid in the two jars appear of equal intensity.

By placing on the white disc a small, black, hemispherical button, this instrument may be used for determining the strength of turbid solutions, such as milk, or for ascertaining the amount of a precipitate suspended in a liquid. Milk is diluted before examination with 99 parts of water, and is compared with a similarly diluted milk of known quality, the disc with the button being adjusted in each tube as above until just invisible.

The slight alteration in the level of the liquid caused by the varying length of the sliding tube contained in the liquid when adjusting the height of the white disc is found by experiment, and a correction must be introduced into the calculation for variations of more than two or three divisions.

A very useful colorimeter, especially designed for the estimation of carbon in steel, but of course available for other purposes, has been introduced by Stead (Proc Iron and Steel Inst 1883, No 1, 213) A similar, but more convenient apparatus, devised by Ridsdale (J. Soc. Chem. Ind. 1886, 586) is shown in Fig. 2. Three glass tubes, each fitted below with a small,




opaque, white cylinder, are placed beneath a movable mirror, in a stand which may be turned to any position upon its weighted base. The central tube is removable and is closed below; the outer tubes are drawn out and pass through indiarubber corks into ruby glass bottles, containing standard solutions of different strengths. Each bottle has a tube with a stopeock through

which the liquid may be introduced or removed, and each is connected by a flexible tube with a syringe.

The liquid under examination is introduced into the central tube, and by working the syringes the standard solutions are raised in their tubes until the intensity of colour in each tube, as seen by looking into the mirror which is adjusted to reflect the light passing upwards through the tubes, is identical. Two distinct comparisons are thus simultaneously made, and, by noting the height of each column by means of the sliding pointer, the strength of the tested liquid may be ascertained from the formula already given. A very important advantage of this apparatus is that every part except the central tube is cleansed and recharged without disconnection.

Ridsdale (J Soc Chem Ind 1888, 70) has devised an apparatus for the examination of liquids of deeper tint than can be used in the foregoing apparatus. The central upper tube (Fig 3) is sealed below, the upper side tubes are perforated at the bottom. Each tube contains a loose white opaque plug, and each is held at



any height by a movable jaw pressed against it by an indiarubber band

The centre tube is charged with, say, 10 divisions of the standard solution, and the test liquid is run into one of the side tubes, whence it passes at once into the lower tube. When this is nearly filled, it is gradually raised to bring a portion of its contents into the upper

tube, until, on looking down vertically, the colours in the upper side tube and in the standard tube are of equal intensity. The calculation is made as already described. The lower tube beneath the standard tube must be raised with that of the side tube in order to equalise the effect of the double thickness of glass. When the solutions to be tested differ widely in tint, the standard tube may be perforated so that the length of its column may be readily adjusted.

Stokes' apparatus (Chem Soc Proc 1887, 135) has two tubes placed vertically in a stand with a reflector below. One tube is connected with a reservoir of the standard solution, by raising or lowering which the level of liquid in

the tube can be altered

A similar arrangement is used in Le Docte's colorimeter (Eng. Pat. 20667, 1898)

The Kennicott-Sargent colorimeter (Chem Engineer, 1907, 213) differs from the above only in altering the level of the standard liquid by moving a glass plunger in a fixed cylindrical reservoir

Another arrangement, in which the standard liquid is forced into a graduated tube from a reservoir, is described by Gunsberg (Dingl poly J. 228, 457).

The Nesslerising tubes adopted by Hehner (Chem News, 33, 185) and recommended by Milne (J. Soc. Chem Ind 1887, 33) are two cylinders of equal and perfectly uniform diameter, holding about 100 c c Each is graduated from below, and at about the 30 c c division has

a glass stopcock.

The two cylinders are placed side by side on white paper, and to one of them is added 100 c c of the liquid to be tested, followed by 2 c c of To the other tube a suitable Nessler's solution and known amount of standard solution of ammonium chloride is added, and the volume is made up to 100 cc. with distilled water, after which 2 c c of Nessler's solution is added two liquids are looked at vertically, and the darker is allowed to slowly run off into a clean beaker, until their colours are equal The comparison may be repeated after filling up from the beaker The mode of calculation is the same as in former cases

A modified form of Eggertz tube has been described by Hogg (J. Soc Chem Ind 1895, 1022)

In a colorimeter devised by Leeds (Chem News, 37, 229) comparison tubes are arranged in a rack, light being reflected through them by an inclined upper mirror, and again reflected to the eye of the observer from a mirror similarly placed beneath them. The standard liquid is contained in a bottle prism which is moved beneath one of the comparison tubes containing distilled water only until the colour observed is identical with that in the tube containing the test liquid. The strength of the test liquid is calculated from the thickness of liquid in that section of the prism which is interposed.

Gallenkamp's colorimeter (Chem Zeit Rep 1891, 324) consists of two vertical troughs enclosed in a metal frame. That which contains the standard solution is wedge-shaped, while that for the test liquid is rectangular. An observation apparatus, moving vertically over a scale of 100 parts, has two slits at the same level, one

opposite each trough, the light from which is reflected and brought into one eyepiece. The observation consists in finding the thickness of the stratum of standard solution in the prism, which shows the same depth of colour as the test liquid.

White's apparatus differs from the above in having two wedge-shaped tanks, both movable with reference to two fixed slits through which


the depth of colour is observed

A colorimeter, described by Autenrieth and Koenigsberger (Chem Zentr 1910, 1 2032) and recommended especially for the determination of hæmatine in blood, depends on comparison of the test liquid with a stable solution of similar hue contained in a wedge-shaped reservoir which is movable vertically beside the trough containing the test liquid. The movement of the wedge is measured on a millimetre scale, and the reading translated directly into percentages by reference to a plotted curve The two colours are observed through a Helmholtz 'Doppelplatte,' which obviates a dividing space in the field of view By substituting a suitable wedge, the instrument can also be used for the estimation of titanium, iron, chromium, iodine, ammonia, nitric acid, &c

Schreiner's colorimeter (J. Amer. Chem. Soc. 1905, 1192) consists essentially of two parts.

wide flat-ended tube, B, about 26 cm × 3 cm, graduated in millimetres, and the flat-ended immersion tube A, about 26 cm × 2 cm, which slides inside the wider tube


Two such pans of tubes are mounted on stand (Fig. 4), the inner tubes being fixed to the top the camera οf while the outer tubes are held in brass clips which them in position, but permit their being moved up and down. The length of the column of liquid enclosed between the ends of the outer and inner tubes can thus be adjusted at will Light,




reflected from a white surface D, passes through the two columns of liquid, and is then reflected at a mirror E to the observer's eyes at F. The axes of the tubes are inclined to one another so as to bring the images close together Standard glass slides may be inserted at G in place of one of the wide tubes, the immersion tube being left in position.

Patterson has described (J Soc Chem Ind 1890, 36) a more complex apparatus employing the same principle. In this case the adjustment of the concentric tubes is made by rack and pinion, and total reflection prisms are employed to bring the two images into one eyepiece. By substituting a direct-vision spectroscope for the eyepiece, the apparatus becomes a spectrum absorptimeter.

The simple and effective colorimeter devised by Steiger (J. Amer Chem Soc. 1908, 215) consists of two wooden boxes painted dead-black inside, one of which (Fig. 5A) is mounted on a



stand and has a rectangular opening c in the base, while the second box (Fig. 5B) fits into the



first, and has a rectangular opening, E, which coincides with c

In the base of this box are two grooves in which slide two rectangular glass cells  $\kappa$ ,  $\kappa$ , about 15 cm long  $\times$  25 cm wide  $\times$  5 cm deep.

Fixed vertically over the opening E are two glass rods F, F, held in place by the springs R, each carrying at the lower end a small plane mirror, G, fixed at an angle of 45°. Light coming through the ground-glass A is reflected by a mirror, B, through the aperture E into the tank, where it is again reflected at the mirror G and passes through a column of liquid, the length of which can be altered by moving the tank backwards and forwards. When equality of colour is obtained, the lengths of the columns are read

off on millimetre scales engraved on the base of each tank. The reading is taken where the lower edge of the mirror touches the bottom of the tank


Otto's ozonometer has two parallel tubes for gases or liquids, and employs wedges of coloured glass moving over a graduated scale as a means of measuring the depth of tint (Eng. Pat. 17524, 1892).

H Kruss suggests (Zeitsch anorg Chem 5, 325) the application to colorimeters of the Lummer-Brodhun double prism, which gives a field of view in which there is no visible line of separation between the two colours under observation. This condition gives maximum accuracy in matching the colours (see also Chem Soc Abstr 1894, ii 158)

Andrieux (Compt rend 103, 281) describes an apparatus in which the *quality* and *intensity* of a colour are determined by comparison with a ray of polarised light

An apparatus has been described by Meisling (Zeitsch anal Chem 1904, 137), in which the hue is matched by the colour of a quartz plate in polarised light, and the intensity is measured by varying the length of the column of liquid

An ingenious instrument, introduced by Mr. J. W Lovibond, for the examination of liquids, powders, fabrics, &c, and known as the tintometer, is shown, partly in section, in Fig. 6. It



consists of a box, closed at one end by a ground glass plate, e, and separated into two divisions b, c. The box is carried by a stand, adjustable to any angle A square glass cell, of known width, along the line q'  $q^2$ , containing the solution to be examined, is placed in one division, and glass plates h of similar colour (or, if necessary, of different colours, producing the proper tint in combination), and of known intensity, are placed in grooves in the other division, until, on looking through both tubes simultaneously through a single eyepiece, d, shaded to prevent access of side light to the eye, the appearance through the glass plates is identical with that through the solution.

The standard plates are specially prepared of various colours, and each is distinguished by a number indicating its colour, and another number showing the intensity of colour, i.e. the number of plates of that colour of unit intensity required to produce an intensity equal to that of the plate. The unit chosen by the inventor for each colour is an arbitrary one, but its actual value can be ascertained at any time by comparison with a standard solution of similar colour.

In some cases the glass plate e is wanting, and an adjustable plane glass mirror is used to reflect the light to the apparatus (Eng Pat 12867, 1886)

When the colour of a fabric is to be estimated, the fabric is stretched on a block placed in the division c, and a white fabric of similar texture, previously compared with a plate of pure plaster

The stanof Paris, is placed in the division b dard plates are then inserted as above until the appearances through the eveniece are identical. Powders, sugars, &c, pressed in a small tray placed in the division c, may be similarly examined, a tray containing pure plaster of Paris being placed in the division b (J Soc Chem. Ind 1888, 234)

For very delicate observations, the intensity of the light used may be found and noted, before commencing, by closing the division c and placing in the divisions b as many standard plates of neutral tint as serve to obstruct all the

light

The substitution of a movable wedge of coloured glass for the standard plates makes the apparatus available for ordinary colorimetric determinations († Soc Chem Ind 1888, 234). An improved form of Lovibond's apparatus is binocular, and has a movable diaphragm makes it applicable to tests of colour vision (ibid 1891, 489) (see also Lovibond, ibid. 1909,

Procter has described (ibid. 1895, 123) a modified form of tintometer in which, by a system of mirrors, the colours to be compared are brought close together in the field of view,

thus giving greater accuracy Sczepanik (Eng. Pat. 22995, 1907) has patented an ingenious apparatus with which a numerical record of any colour can be obtained Three strata of liquid of the three primary colours are superimposed. The thickness of each stratum can be measured and altered independently of the others, and thus any hue or intensity of colour can be matched

Gardner and Dufton have introduced the 'Dalite' lamp, which gives a light of the quality of daylight, and may be used for tintometric work (J Soc Chem Ind 1904, 598) This lamp is also of use in performing titrations by

artificial light.

A useful device, introduced by Dupré (Analyst, 5, 123), may be mentioned under this heading In the volumetric estimation of silver with potassium chromate as an indicator, especially when the solution is weak, it is somewhat difficult, on account of the original yellow colour of the solution, to observe the first tinge of red produced by the formation of the silver Dupré interposes, between the eye chromate and the test liquid, a flat glass cell (of which the faces are about half an inch apart) containing a solution of potassium chromate of such strength as will neutralise the colour of the test liquid, which thus appears perfectly colourless, and in which the faintest appearance of red is readily visible The cell filled with other liquid may be similarly adopted in many other titrations

Kong (Proc Amer Phil Soc [18] 7, 29, 184, 208, 211) has devised a method for estimating metals by the intensity of the colour imparted by them to the borax bead He dissolves a known weight of the oxide in a known weight of borax, grinds the bead until its faces are flat, coats it with Canada balsam to render its rough faces transparent, and places it in an apparatus where it is viewed by a lens of small power, while, by means of a rack, a glass prism whose colour is complementary to that under examination, is moved before it until such a thickness is interposed that the colour of the bead is extinguished.

A photochemical process for estimating differences in the intensity of a colour has been described by E. J. Mills and Buchanan (J Soc. Chem Ind 1888, 309) It has been applied to the comparison of dyed goods. A number of discs of the tissues are photographed on one negative from which a number of positives on paper sensitised with silver, are copied are treated as usual, soaked for a considerable time in sodium thiosulphate to remove the whole of the unaltered silver salt, well washed, and dried. The disc photographs of each tissue are cut from a number of positives and each set is burned separately The silver in the residue is dissolved and estimated volumetrically relative amount of silver found in the different sets of discs directly represents the relative intensity of the colours.

The following colorimetric process for the estimation of minute quantities of sulphur in iron, mineral waters, black ash, slag, &c, is due to Wiborgh (Chem Zeit 10, 92, Chem News,

1886, 54, 158).

The divided metal, ore, &c, or the concentrated solution, is introduced into a small flask provided with an indiarubber cork, through which passes a stoppered funnel for introduction of acid, and a narrow-mouthed funnel ground at A disc of white calico, which has been previously saturated with a solution of 5 grams crystallised cadmium nitrate (or, preferably, cadmium acetate) in 100 c c of water and dried in a horizontal position on a cloth, is fixed between two indiarubber rings of the same interior diameters as the standard discs hereafter described, and the rings are firmly clamped over the funnel mouth Dilute sulphuric acid is poured into the flask and heat is applied from 30 to 45 minutes the whole of the sulphur which is capable of conversion into sulphuretted hydrogen has passed off in that form and has been wholly absorbed by the cadmium salt on The disc is removed and compared the calico with a set of standard discs, similarly prepared from iron containing a known amount of sul-The standard discs should not be exposed too freely to light Arsenic, carbon, silicon, and copper do not appear to affect the determination. This method is so delicate that a disc will show a tinge of the yellow colour due to the cadmium sulphide when so small a quantity as the 1000th of a milligram of sulphur is present per square centimetre

COLOUR AND CHEMICAL CONSTITUTION. The object of this article is to give as concise a description as possible of the various rules which have been established, and the theories which have been proposed, with regard to the relation between colour and chemical constitution, particularly to the colour of dyes, with special reference to those rules which have proved of value in the preparation of dyes and which have served as guides for modifying the

colour in a group of dyes

Outline -The first Historical synthetic dye (mauve) was produced by Sir W. H. Perkin in 1856. This branch of synthetic chemistry developed with great rapidity, and in the next twenty years the first members of many important groups of dyes had been prepared, eg. fuchsine (1859), rosaniline blue (1860), and alkylated triphenylmethane dyes (1863–1866);

Bismarck brown (1864), chrysoidine (1876), and naphthol orange (1876) in the azo- group, Martius yellow (1864) in the nitro- group, Magdala red (1868) in the azine series; alizarin (1869) of the anthraquinone group; fluorescein and cosin (1874), representing the phthaleins, and Lauth's violet (1876), foreshadowing the thiazines. By this time sufficient material had been accumulated to allow the formulation of the well-known Witt's Rule (1876), according to which substances attained the potentiality of colour when their molecules contained certain atomic arrangements or groups, which Witt termed chromophores The colour and dyeing properties were made manifest by the introduction of additional groups termed auxochromes

Within the next three years additional triphenylmethane and azo- dyes were produced, deeper colour (red) being attained in the latter group by the use of naphthalene components in place of the benzene derivatives used in the earlier members, as eg in the Ponceaus (1878), and also by the introduction of an additional chromophore, as in Biebrich scailet (1878) consideration of the results now obtained, especially in the triphenylmethane and azogroups, led to the enunciation of Nietzki's Rule (1879), which states that the colour of a dyestuff can be deepened by adding groups so as to increase its molecular weight, and that the deepening of colour produced is more or less proportional to the increase of molecular weight (It should be noted that the term depth is used by colour-chemists in a very definite and limited way In ordinary language the term depth of colour has at least two meanings Blue and red are said to be deeper colours than yellow, and also one yellow is said to be deeper than another yellow if it is stronger or more intense. ie less admixed with white Colour-chemists restrict the use of the term, and employ it for the first purpose only. Blue is deeper than red or yellow, and red is deeper than yellow. The ordinary colours arranged according to depth are green, blue, violet, red, orange, yellow. i e. the complements to the ordinary colours of the spectrum taken in order from the red to the violet end This rule served for many years as a most valuable guide in extending the range of colours in many groups of dyes, especially in the azo- group, in which the deepest colours were now soon obtained (blue-black. 1882, Glaser) It may be said that this rule is still a valuable guide to the colour chemist

In 1885 Liebermann and v. Kostanecki published the results of their investigations of the colour and dyeing properties of the hydroxyderivatives of anthraquinone Although these authors did not commit themselves to any very definite statement, yet their publication drew the attention of colour-chemists to the ease with which colour could be deepened by the multiplication of auxochrome groups in the molecule, provided these groups were suitably This general principle may be said arranged to have been of equal value with Nietzki's Rule, and for many years the development of dye manufacture was almost entirely controlled by these two rules. Very interesting and ingenious methods were developed by Bohn and Schmidt in 1888 and 1890 for the multiplication of hydroxyl groups in the anthra-

quinone molecule with the result that a complete range of colours was soon available in this group of mordant dyes. The continued application of the same idea led somewhat later to the preparation of anthragumone dyes containing both amino- and hydroxyl- groups, such as alizarin saphirol (1898), which combines the advantages of deep colour and fastness with brightness and cleainess of tone. Liebermann and Kostanecki's Rule led at once to the careful study and production of dihydroxy-, amino-, hydroxy-, and diamino-naphthalene derivatives and their employment in the manufacture of azo- dyes This general principle led to the production between the years 1887 and 1897 of an enormous number of azo- dyes of all colours

In 1888 H E Armstrong put forward the best-known theory which has ever been formulated as to the cause of colour in organic com-This is known as the Quinonoid Theory. Its author pointed out that the chemical formulæ of the well-known dyestuffs either contained the quinonoid grouping or could be slightly modified or rearranged, in accordance with accepted ideas on tautomerism, so as to contain this grouping More recent work has shown that the original quinonoid theory cannot stand Modifications have been put forward by Baeyer (1907) and H v Liebig (1908), neither of which,

however, is entirely satisfactory

Of recent years it has come to be generally recognised that visible colour is only a special case of selective absorption. That some substances are coloured whilst others, nearly related, are colourless, must be regarded as more or less accidental, and due to a physiological cause rather than to any essential physical difference between them, viz to the fact that the human eye can only detect selective absorption within a limited range of the spectrum The problem of colour has thus resolved itself into two parts (1) the cause of selective absorption, (2) the cause of the absorption lying within the limits of the visible spectrum

As to the cause of selective absorption various theories have been put forward in iecent years. We may mention Baly and Desch's theory, according to which absorption bands are due to dynamic isomerism (1904-1905), Stewart and Baly's theory of isorropesis as the cause of selective absorption (1906), Watson and Meek's theory that absorption in the visible spectrum and its neighbourhood is due to the reversal during tautomeric change of a conjugate chain of alternate double and single bonds (1915), the theory that absorption bands in the visible part of the spectrum are due to the vibiation of elections, and Baly's theory (1914) that absorption bands in the visible and ultra-violet parts of the spectrum are due to overtones or summation or difference tones derived from fundamental vibrations causing infra-red absorption bands (infra-red absorption bands have been attributed by most investigators to the vibrations of definite chemical radicals, Abney and Festing\_(1881), Julius (1888), Coblentz (1904–1905), and Weniger (1910)) In 1897 Pauer showed that under certain circumstances ordinary broad absorption bands are replaced by a large number of fine bands, recalling the lines in the emission spectra of substances examined in flames or under the influence of the electric discharge. Hartley (1908) came to the conclusion that the ordinary broad absorption bands are formed by the fusion of these narrow bands. Various observers have shown the intimate connection between such fine absorption bands and the lines in the emission spectra of the same substances, e.g. Wullner (1863), Mitscherlich (1864), Salet (1872), Eder and Valenta (1899). And, finally, in certain cases (hydrogen and helium) the lines in the emission spectra have been deduced with great accuracy on the hypothesis that they are due to rotational vibration of electrons within the atoms.

The fact that so many different theories have been formulated makes it obvious that no final conclusion has been reached as to the cause of absorption bands in general

As to the causes which bring the absorption bands into the visible spectrum and influence their positions within the visible spectrum, many investigators showed that Nietzki's rule was only a first approximation to the facts, and many exceptions to this rule were collected by Schutze (1892) Hewitt (1907) suggested that the chief oscillation frequency was less, and consequently the colour was deeper, the longer the conjugate chain of alternate double and single bonds in the molecule. The present writer pointed out in 1914 that those dyes of which the molecular formula can only be written in a quinonoid form have all a deep colour, whilst if the formula can be written either in a quinonoid or a non-quinonoid form, even great molecular complexity may fail to produce any great depth of colour Watson and Meek (1915) showed that in several series the period of the absorption band is proportional to the length of the conjugate chain, which may be assumed to reveise during tautomeric change

No explanation has yet been offered of the great effect on depth of colour which is so frequently produced by a multiplication of auxochromes. The relative position of the auxochromes has a great effect on the colour Certain regularities in this respect have been pointed out by Kauffmann and Franck (1906), and Meek and Watson (1916)

It cannot be said that the theories which have been formulated since 1885 have proved of much service for the preparation of dyes In the older group of dyes a complete range of colours had already been attained by assiduous experiment, guided by Nietzki's and Liebermann and Kostanecki's rules It may perhaps be claimed that the quinonoid theory had some influence in the development of the azopyrazolone dyes. The present writer was able to apply one of the rules which he formulated to the production of deep-coloured dyes from the yellow flavones, and the work in this direction was perhaps of service for the recognition of the molecular structure of the colouring matters of flowers (anthocyanins) As to the new groups of valuable synthetic dyes which have been developed recently, the sulphide dyes, and the anthraquinone vat-dyes, the constitution of the former has been too little understood to allow of anything but empirical investigation, whilst in the latter group (the anthraquinone vat-dyes), our theories seem at

present quite unable to explain the wealth of colour phenomena exhibited by the dyes already synthesised

A short account is appended of some of the more important theories and rules above mentioned

Wut's Rule—This investigator pointed out (Ber 1876, 9, 522) that two things are required to give an organic substance colour and dyeing properties. The molecule must possess a certain grouping of atoms to give it the potentiality for colour, and it must also possess a salt-forming radical to bring out the colour and dyeing properties. The atomic groupings giving the potentiality of colour he termed chromophores, and molecules containing them were called chromogens. He specified the following chromophores and gave illustrations of chromogens containing them—

It will be noticed that some of the chromogens are colourless, eg nitrobenzene and phenolphthalein, whilst others have a feeble colour, eg azobenzene and anthraquinone; but none of the chromogens has any dyeing property. The colour is developed or strengthened by the introduction of sait-forming ladicals into the chromogens. These ladicals ne therefore termed auxochromes. Hydroxyl, aminoand substituted amino- groups, eg. NHCH<sub>3</sub>, N(CH<sub>3</sub>), &c, are the principal auxochromes. Not all salt-forming groups can act as auxochromes, eg. sulphonic and carboxyl groups are without this function. The following table shows how the chromogens already mentioned are converted into dyes by the introduction of auxochromes.—

$$\begin{array}{c} \operatorname{Chromogen} & \operatorname{Dye} \\ \operatorname{C}_6\operatorname{H}_5\operatorname{NO}_2 & \operatorname{C}_6\operatorname{H}_4(\operatorname{NO}_2)\operatorname{OH} \\ \operatorname{Nitrophenol} & \operatorname{C}_6\operatorname{H}_5(\operatorname{NO}_2)\operatorname{NH}_2 \\ \operatorname{C}_6\operatorname{H}_5-\operatorname{N}=\operatorname{N}-\operatorname{C}_6\operatorname{H}_5 & \operatorname{C}_6\operatorname{H}_6-\operatorname{N}=\operatorname{N}-\operatorname{C}_6\operatorname{H}_4\operatorname{OH} \\ \operatorname{Benzene-azo-phenol} & \operatorname{C}_6\operatorname{H}_5-\operatorname{N}=\operatorname{N}-\operatorname{C}_6\operatorname{H}_4\operatorname{OH} \\ \operatorname{Benzene-azo-phenol} & \operatorname{CO\cdot\operatorname{C}_6\operatorname{H}_4}\operatorname{NH}_2 \\ \operatorname{Anline vellow} & \operatorname{CO\cdot\operatorname{C}_6\operatorname{H}_3}\operatorname{OH} \\ \operatorname{C}_6\operatorname{H}_4 & \operatorname{O} & \operatorname{C}_6\operatorname{H}_4 & \operatorname{O} \\ \operatorname{CO\cdot\operatorname{C}_6\operatorname{H}_4} & \operatorname{CO\cdot\operatorname{C}_6\operatorname{H}_3}\operatorname{OH} \\ \operatorname{Fluoiescein} \end{array}$$

THE QUINONOID AND MODIFIED QUINONOID THEORIES.

In 1888 (Proc. Chem. Soc. 1889, 27-33) H. E. Armstrong pointed out that the chemical formulæ of the well-known dyestuffs either contained the quinonoid grouping or could be slightly modified or rearranged, in accordance with accepted ideas on tautomerism, so as to contain this grouping. The quinonoid grouping may be defined as a benzene nucleus to which

other atoms or groups are attached by double bonds in the para- or ortho- positions.

Armstrong was probably much impressed by the recently discovered fact that benzene azo-β-naphthol was identical with the phenyl hydrazone of  $\beta$ -naphthogumone, so that there was as much justification for the formula

as for the more usual one based on the formation of the substance from  $\beta$ -naphthol and a benzenediazonium salt, viz

$$HO$$

$$C_6H_5N=N$$

and if it were allowable to assume tautomerism to a quinonoid form in this case, it might be allowable in other cases, e.g. in the case of fluorescein

whilst in the case of p-rosaniline it was quite consistent with experimental fact to suppose that on the conversion of the base into salts there was elimination of a molecule of water and formation of a substance with a quinonoid structure

$$\begin{array}{cccc} \operatorname{HO-C} & \overset{\operatorname{C}_6H_4\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}}{\overset{\operatorname{HCl}}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{\operatorname{C}_6H_4-\operatorname{NH}_2}{\overset{C}_6H_4-\operatorname{NH}_2}{\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6H_4-\overset{C}_6$$

In many cases the ordinary formula already contained the quinonoid structure

The theory has been shown to be untenable On the one hand, substances have been prepared which have a quinonoid structure, but are colourless, and on the other hand, coloured substances are known to which a quinonoid structure cannot be assigned.

Colourless Substances possessing a Quinonoid Structure—Immoquinone (I), diimmoquinone (II), quinonedimethylimine (colourless in crystalline form, yellow in solution) (III), diphenoquinone (exists in two modifications, one of which is colourless) (IV), and fuchsonimine (V); also the dimethyl ether (VI), the diacyl

derivatives (VII), and the anhydride (VIII) of phenanthraquinone dioxime

Coloured Substances to which a Quinonoid Structure cannot be assigned -Nitroquinol dimethyl ether, the sulphate of p-trichlortriphenyl carbinol, the fulvenes, eg dimethylfulvene, and some fatty compounds, such as diacetyl, tertiary nitroso- compounds, e.g. tert-nitrosobutane, and dinitroparaffins.

Baeyer's Modification of the Quinonoid Theory—It has already been mentioned that fuchsonimine is colourless. The introduction of a para- amino group into one of the phenyl groups causes the development of colour

This and similar facts led A v Baeyer to suggest that the cause of colour is an oscillation of the quinonoid condition between two or more benzene nuclei (Jour Chem Soc Abs 1907, 1 504) If we call the three benzene nuclei in the above compound A, B, and C, it will be seen that the oscillation of a hydrogen atom will allow A and B alternately to become quinonoid:

$$\begin{array}{c|c}
NH = & A \\
NH_2 & B
\end{array}$$

$$\begin{array}{c|c}
NH_2 & A \\
\hline
NH = & B
\end{array}$$

$$\begin{array}{c|c}
C & C
\end{array}$$

Baeyer suggested that this oscillation was the cause of the colour.

Many strongly-coloured substances are known in which such an oscillation is not possible And necessarily all the arguments against the original quinonoid theory apply with still greater

H v Liebig's Theory that — A—A—A—is the Essential Organic Chromophore—The fact that the fulvenes are coloured, although they do not contain the quinonoid structure, led H. v Liebig to suggest that the essential cause of colour is the arrangement of linkages which is possessed both by quinonoid compounds and

the fulvenes, viz ——A—A——A—— (Annalen, 1908, 360, 128) This theory explains the colour of nitroquinol dimethyl other, which contains this arrangement of linkages —

The theory will not explain the colour of the tertiary fatty nitroso-compounds, and, of course, the existence of colourless quinonoid compounds is as much an argument against this modification as against the original quinonoid theory.

as against the original quinonoid theory
NIETZKI'S RULE The colour of a dyestuff
can be deepened by adding groups so as to increase
its molecular weight, and the deepening of colour

produced is, more or less, proportional to the increase of its molecular weight (Verhandl des Vereins zum Beforderung des Gewerbefleisses, 1879, 58, 231). This rule was probably formulated by Nietzki to summarise well-known facts in connection with the triphenylmethane group of dyes, but apparently was not based on any systematic investigation. The table on p. 349 shows that the rule applies to a considerable extent to other groups besides the triphenylmethane series.

The rule was subjected to a careful examination by Schutze (Zeitsch physikal Chem 1892, 9, 109 et seq), who pointed out many exceptions. He showed that in some cases, e.g on the addition of a nitro-group to indigo, the shade even becomes lighter instead of deeper, and he introduced the terms bathychrome and hypsochrome, the former being applied to those groups which produce a deepening of colour, and the latter to those having the contrary effect. Another example of a hypsochrome is the acetyl group All dyestuffs containing hydroxyl- or aminogroups become much lighter in colour when completely acetylated.

The following table is of interest, as showing that in some cases the molecular weight may be made very large without a deep colour being attained

Moleculai tormula	Mol wt	Shade of dyemg	
$('_{6}H_{5}-N=N-('_{6}H_{4}OH))$	198	yellow	
$C_{6}H_{5}-N=N-C_{6}H_{4}-N=N-C_{6}H_{2COOH}^{OH}$	346	brownish yellow	
${\rm C_6H_5-N\!=\!N\!-\!C_6H_4\!-\!N\!=\!N\!-\!C_6H_4\!-\!N\!=\!N\!-\!C_6H_{2\rm COOH}}$	450	brownish yellow	
$('_{6}H_{5}-N=N-('_{6}H_{3}('_{6}H_{9}))$	274	orange	
$C_6H_4$ — $N=N-C_6H_3$ $COOH$ $C_6H_4$ — $N=N-C_6H_3$ $OH$ $COOH$	482	yellow	
$({}^{\circ}_{6}H_{4}-N=N-({}^{\circ}_{6}H_{3COOH}^{3COOH}$ $({}^{\circ}_{6}H_{4}-N=N-({}^{\circ}_{6}H_{3COOH}^{3COOH}$ $({}^{\circ}_{6}H_{4}-N=N-({}^{\circ}_{6}H_{3COOH}^{3COOH}$	752	yellow	

HEWITT'S RULE This investigator's researches on the colour of azo-compounds led him to the conclusion that the chief oscillation frequency is less, and consequently the colour is deeper, the longer the conjugate chain,  $\iota$  e the chain of alternate double and single bonds, in the molecule (Chem. Soc. Trans. 1907, 91, 1251)

He noticed that in the case of many azodyes the introduction of a mitro-group into the benzene nucleus not containing auxochromes produced very pronounced deepening of colour in alkaline solution, eg p-nitrobenzeneazophenol and p-nitrobenzeneazonaphthol have much deeper colours (crimson and violet) than benzeneazonaphenol and benzeneazonaphthol (orange and crimson respectively) He brought forward evidence in favour of the assumption that in alkaline solution these nitro-derivatives

have an iso- nitroso structure, and in this form the nitro- derivatives have a much longer conjugate chain than the parent substances, eg —

$$N = \begin{cases} N - N = \\ N - N = \end{cases} = 0$$

compared with:

$$\langle$$
 NH-N= $\langle$  =0

Sircar (Chem Soc Trans 1916, 109, 757, et seq) showed that in very similar azo-compounds the length of the conjugate chain is not proportional to the depth of colour, e.q. 4-nitronaphthalene-1-azophenol only gives a red colour in alkaline solution, whilst p-nitrobenzene-azo- $\alpha$ -naphthol dissolves in the same solvent with a blue colour, whilst the two substances

	OLOUR AND CHEMICAL CONST	ITOTION	349	
Name	Name Molecular formula		Shade of dyeing	
Pararosaniline	$C_{0}^{+}H_{4}NH_{2}$ $C_{0}^{+}H_{4}NH_{2}$ $C_{0}H_{4}=NH_{2}CI$	281.5	Magenta	
Methyl Violet R .	$\begin{array}{c} C_6H_4NHCH_3\\ C - C_6H_4NHCH_3\\ C_6H_4 = NHCH_3CI \end{array}$	323 5	Reddish violet	
Methyl Violet 3B .	$\begin{array}{c} C_{0}H_{4} \text{ N(CH}_{3})_{2} \\ C_{0}H_{4} \text{ N(CH}_{3})_{2} \\ C_{0}H_{4} = \text{N(CH}_{3})_{2}\text{Cl} \end{array}$	365 5	Violet	
Rosaniline Blue .	$\begin{array}{c} C_{6}H_{4}NHC_{6}H_{5} \\ C = C_{6}H_{4}NHC_{6}H_{5} \\ C_{6}H_{4} = NHC_{6}H_{5} Cl \end{array}$	509 5	Blue	
Iodine Green .	$\begin{array}{c} C_{6}H_{4} \text{ N(CH}_{3})_{2} \\ C_{6}H_{4}\text{ N(CH}_{3})_{2} + CH_{3}I \\ C_{6}H_{4} = \text{N(CH}_{3})_{2}I \end{array}$	599	Green	
Benzene-azo-phenol .	$C_6H_5$ —N=N— $C_8H_4OH$	- 198	Yellow	
Benzene-azo-naphthol.	$C_6H_5$ — $N=N$ — $C_{10}H_6OH(\beta)$	248	Red	
Biebrich Scarlet	$\begin{array}{c} SO_3H C_6H_4-N=N-C_6H_3(SO_3H) \\ -N=N-C_{10}H_6 OH(\beta) \end{array}$	512	Red	
Naphthol Black	$(SO_3H)_2C_{10}H_5 - N = N - C_{10}H_6 - OH$ $N = N - C_{10}H_4 - (SO_3H)_2$	772	Black	
	$N = N - C_{10}H_4 (SO_3H)_2$			
Alizarın .	$C_{\theta}H_{4}$ $CO$ $C_{\theta}H_{2}(OH)_{2}$	240	Red (on alum)	
Alızarın Bordeaux .	(OH) <sub>2</sub> C <sub>6</sub> H <sub>2</sub> CO C <sub>6</sub> H <sub>2</sub> (OH) <sub>2</sub>	272	Bordeaux (on alum)	
Alizarın Cyanine .	$(OH)_2C_6H_2$ $CO$ $C_6H(OH)_3$	288	Violet (on alum)	
Fluorescoin	но о он	332	Yellow	
Eosin	C <sub>6</sub> H <sub>4</sub> COOO Br HOOODH	648	Red	
	Br COO			
Gallein	HO COO OH OH	364	Violet to blue (according to mordant.)	
Indigotin Ciba Blue BB (Tetra-bromindigotin)	C <sub>16</sub> H <sub>10</sub> N <sub>2</sub> O <sub>2</sub> C <sub>16</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> Br <sub>4</sub>	262 578	Indigo blue Greenish blue	

have conjugate chains of the same length He came to the conclusion that in azo- dyes the depth of colour is proportional to the length of the conjugate chain reckoned from the auxo-

chrome to the azo- linkage

WATSON'S RULE —The present writer pointed out (Proc Chem. Soc 1913, 29, 348. Chem Soc. Trans 1914, 105, 759) that dyes which are quinonoid in all possible tautomeric forms have a deep colour, no matter how simple their structure or how small their molecular weight.

'In the following table there are placed opposite each other in the two columns dyes of very similar constitution which yet show remarkable differences in depth of colour -

Hexahydroxybenzophenone (Dyes yellow on alum and chrome )

$$C_6H_2(OH)_3$$
 CO  $C_6H_2(OH)_3$ 

Leather yellow (Dyes yellow on leather )

 $C_6H_4$  NH 2

 $C_6H_4$  NH 2

4 4'-Diaminoazobenzene (Yellow) NH2 C6H4 N N C6H4 NH2

Quercetin (Yellow on atum yellowish brown on chrome)

Myncetin (Dyes similar shades to queicetin)

Anthracene blue W R (Dyes violet on alum, blue on chrome )

O: 
$$C_6H(OH)_2$$
  $C_6H(OH)_3$ 

Magenta (Dyes bluish red ) C<sub>6</sub>H<sub>4</sub> NH<sub>2</sub> CH2N.C6H4 CC6H4NH2 Indamme (Blue)

NH · C<sub>6</sub>H<sub>4</sub> N C<sub>6</sub>H<sub>4</sub> NH<sub>2</sub>

Gallacetein (Dyes violet on chrome )  $C_6H_2(OH)_2$   $C_6H_2(OH)_2$  CH

'The dyes in the right-hand column can only be represented by formulæ containing a quinonoid arrangement. Whatever tautomeric change they may be supposed to undergo their formulæ still contain the quinonoid arrange-

Those on the left hand differ in that they ment can be represented by formulæ not containing this arrangement They can, of course,t be supposed to tautomerise into forms containing this quinonoid airangement, but in all cases it is possible to represent them also in a non-quinonoid form.' 'A survey of all the better known dye-stuffs fully bears out this theory. The triphenylmethane dyes, the pyronines, benzeins, and fluorones, the indophenols, indamines, oxazines, thiazines, azines (safianines, &c), all satisfy this requirement and exhibit deep colour, in many cases with little molecular complexity. Hæmatein and brazilein conform to this rule. They have almost the same molecular weight as the flavones, and there is considerable resemblance in their structure, but they conform to this rule, whilst the flavones do not, and consequently their colour is much The alizarin series, and also naphthazarin, illustrate the rule On the other hand, the monoketonic dyes, ey the benzophenone and xanthone derivatives, the flavones and flavonols, ellagic acid, and the azo-dyes can be represented by non-quinonoid formulæ, and many members of these groups, despite very considerable molecular complexity, have no great depth of colour

WATSON'S AND MEEK'S MODIFICATION OF HEWITT'S RULE—In passing from one tautomeric form to another there is a reversal of the conjugate chain, so that double and single bands change places all along the conjugate In that group of substances which can only tautomerise from one quinonoid form to another the conjugate chain reversed is generally longer than in the other group of substances which can tautomerise from a quinonoid to a non-quinonoid form An appreciation of this fact led to the formulation of a further modification of Hewitt's rule, viz. that the colour is deeper the longer the conjugate chain reversed during tautomenic change (Watson and Meek, Chem Soc Trans 1915, 107, 1567) This rule was tested by measuring the wave-lengths of the maxima of the absorption bands of seven pairs of substances which were selected as being very similar in structure, but having conjugate chains of different lengths reversed during tautomerism. It was found that in each of these seven pairs the period of the vibration was proportional to the length of the conjugate chain reversed (see table, p 352)

The reversal of the chain is shown in the following structural formulæ for 4 · 4'-tetramethyldiaminobenzhydrol (Michlei's hydrol) and 4 4'-tetramethyldiaminobenzophenone (Michler's ketone)

$$-(\operatorname{CH}_3)_2 \operatorname{N} = \operatorname{CH} - \operatorname{N}(\operatorname{CH}_3)_2 = \operatorname{CH} - \operatorname{CH} - \operatorname{N}(\operatorname{CH}_3)_2 = \operatorname{CH} - \operatorname{CH} -$$

Name	Molecular formula	Shade of dyeing
Alızarın .	$C_6H_4$ $CO$ $C_6H_2(OH)_2$	Red (on alum)
Alızarın Bordeaux	(OH) <sub>2</sub> C' <sub>6</sub> H <sub>2</sub> CO (C <sub>6</sub> H <sub>2</sub> (OH) <sub>2</sub>	Bordeaux (on alum)
Alizarın ('yanıne	$(\mathrm{OH})_2\mathrm{C_6H_2} < \frac{\mathrm{CO}}{\mathrm{CO}} \mathrm{C_6H}(\mathrm{OH})_3$	Violet (on alum)
Fluorescem  Gallem .	$\begin{array}{c} HO & O \\ C_6H_4 \\ CO \\ O \\ HO & OH \\ C_6H_4 \\ CO \\ O \end{array}$	Yellow  Violet to blue (according to mordant)
And Andre Yellow . Chrysoidine .	. $({}^{'}_{6}H_{4}-N=N-C_{6}H_{4}NH_{2})$ . $({}^{'}_{6}H_{4}-N=N-C_{6}H_{3}(NH_{2}))$	Yellow Orange
Azooiseilline .  Congo Violet . Congo Red .	. (' <sub>6</sub> H <sub>4</sub> —N=N—C <sub>10</sub> H <sub>6</sub> (SO <sub>3</sub> H)OH (' <sub>6</sub> H <sub>4</sub> —N=N—C <sub>10</sub> H <sub>6</sub> (SO <sub>3</sub> H)OH . ditto (isomeric) . (' <sub>6</sub> H <sub>4</sub> —N=N—C <sub>10</sub> H <sub>6</sub> (SO <sub>3</sub> H)NH <sub>2</sub>	Brownish-red Violet Red
Naphthocyanino	$\begin{array}{c} ({}^{\prime}_{6}H_{4}-N=N-({}^{\prime}_{10}H_{6}(SO_{3}H)NH_{2})\\ ({}^{\prime}_{6}H_{4}-N=N-({}^{\prime}_{10}H_{4}(SO_{3}H)(OH)_{2})\\ ({}^{\prime}_{6}H_{4}-N=N-({}^{\prime}_{10}H_{4}(SO_{3}H)(OH)_{2}) \end{array}$	Blue
Diamine Black, B	$\begin{array}{c} \cdot & C_{6}H_{4}-N=N-C_{10}H_{4}(SO_{3}H)(OH)(NH_{2}) \\ \cdot & C_{6}H_{4}-N=N-C_{10}H_{4}(SO_{3}H)(OH)NH_{2} \\ \cdot & \text{ditto (isomeric)} \end{array}$	Black Blue
Brasileïn .	. НО СН <sub>2</sub> С(ОН)	Red
Hæmatein .	OH   O	Blue

	Length of conjugate chain reversed during tautomerism	Wave-length of absorption band
4 4'-Tetramethyldiaminobenzhydrol 4 4'-Tetramethyldiaminobenzophenone	$\frac{5}{3}$ ratio=1 $\dot{6}$	6100 ratio=1 65
4 · 4'-Diaminobenzhydrol 4 · 4'-Diaminobenzophenone	$\frac{5}{3}$ ratio=1 $\delta$	$\frac{5640}{3414}$ ratio=1 65
Quercetin reduction product Quercetin	$\frac{5}{3}$ ratio=1 $\dot{6}$	$\frac{6100}{3690}$ ratio=1 65
Morin reduction product  Morin	$\frac{5}{3}$ ratio=1 $\dot{6}$	$\frac{5860}{3800(?)}$ ratio=1.54(?)
Apigenin reduction product Apigenin	$\frac{5}{3}$ ratio=1 $\dot{6}$	$\frac{5700}{3410}$ ratio=1.67
Pyronine G 3 6 Tetramethyldiaminoxanthone	$\frac{4}{3}$ ratio=1 3	$\frac{5480}{3861}$ ratio=1.41
3 Hydroxyfluorone 3 6 Dihydroxyanthone	$\frac{4}{3}$ ratio=1.3	$\frac{5049}{3810}$ ratio=1 32

# EFFECT OF MULTIPLICATION OF AUXOCHROMES ON DEPTH OF COLOUR

Additional auxochromes frequently deepen the colour to a very great extent. This may be regarded as an example of Nietzki's rule, but the effect produced is generally much greater than we should expect from the very moderate weight of the additional auxochromes, eq from the first table given in the paragraph on Nietzki's rule, it will be observed that two hydroxyl groups added to fluorescein (producing gallein) deepen the colour much more than four bromine atoms (producing cosin), although their weight is much less. The table on page 351 illustrates the great effect of additional auxochromes

The relative positions of the auxochromes has a very great effect on the colour, eq there are two isomeric hexahydroxyanthraquinones, anthracene blue WR and rufigallol, one of which is blue and the other brown

From a study of the polyhydroxyanthraqumones, Meek and Watson (Chem Soc Trans. 1916, 109, 557, et seq) suggested the following rules as to the influence on the colour of the relative positions of the auxochromes —

(a) Two hydroxyl groups in one benzene nucleus in the o- or p-position with respect to one another are necessary to produce a deep colour

(b) The colour is still further deepened if both benzene nuclei contain pairs of hydroxyl groups in the o- or p-position to one another

(c) Three hydroxyl groups in the 1 2 4-position in one benzene nucleus produce a deeper colour than a pair of hydroxyl groups in the o- or p-position.

(d) Three hydroxyl groups in the 1:2·3-position in one benzene nucleus produce a brown colour.

And these rules are apparently applicable to all groups of dyes, eq

(a) The hydroxy- derivatives of azobenzene dissolve in potash with the following colours:—

p-Hydroxyazobenzene
o-Hydroxyazobenzene
Benzeneazoresoreinol
Benzeneazoetechol . Carmine
Benzeneazoquinol . Blue
Benzeneazopyrogallol . Brick-red

(b) Fluorescein, quinolphthalein, and catecholphthalein dissolve in potash with a yellow colour, hydroxyquinolphthalein gives a cheiryred, gallein a blue, and phloroglucinolphthalein an orange under the same conditions

(c) The same rules are seen to apply to the pyranol dyes

E R W.L.

COLUMBIA BLACK, -BLUE, -GREEN v.

Azo- colouring matters

COLUMBIA YELLOW v Primuline and its derivatives.

COLUMBIC ACID, COLUMBAMINE, and COLUMBIN v CALUMPA

COLUMBITE. A columbate (mobate) of usually containing some manganese ' manganocolumbite') and tantalum, the general formula being (Fe,Mn)(Cb,Ta)<sub>2</sub>O<sub>6</sub> With an increase in the amount of tantalum, there is a transition to the isomorphous species, tantalite The orthorhombic crystals are dark brown or black and opaque, and range in sp gr. from 5 3 (columbite) to 7 3 (tantalite) It was in this mineral that C Hatchett discovered, in 1802, the element columbium, so named because the specimen he examined came from Columbite is usually found as single crystals embedded in pegmatite or granite; for example, at Haddam and Middletown in Connecticut, Chesterfield in Massachusetts, brilliant crystals in granite at Standish in Maine, large masses up to 2000 lbs in granite veins in the Black Hills of South Dakota, and at At lyigtut several other American localities in Greenland it has been found in the cryolite deposit Large crystals are found in the felspar quarries near Moss in Norway. Other localities are Rabenstein in Bavaria, Bushman Land in South Africa, Madagascar, &c.

COLUMBIUM (Niobium). Sym. Cb. At. wt 93.1 (Smith and van Haagen). Although this metal is still commonly termed niobium by the mineralogist and the metallurgist, the name columbium, which was given to it in 1802 by Hatchett, who first isolated it in the form of a new 'earth' (Cb2O5) from a mineral found in Connecticut, has been officially adopted by the chemist in English-speaking countries, and generally throughout the world, except perhaps in Germany.

Columbium is a member of a group of comparatively rare metals, including tantalum, tungsten, molybdenum, titanium, and uranium, commonly occurring together in minerals found in the oldest Plutonic rocks and especially in and around intrusions which traverse pegmatites and other extremely coarse-grained granito rocks. Monazite, and other minerals containing metals of the cerum group, zircons, cryolite, and fluorspar, and minerals containing lithium, cassium, and rubidium, are often associated with these minerals, and are probably intimately connected with their

Although an extremely rare metal, columbium exists as the pentoxide in a considerable number of minerals, in some of which—such as pyrochlore and columbite-it may be regarded as the essential constituent, while, in others, although practically always present, its proportion varies from a mere trace up to an amount which justifies its being considered as the principal oxide.

Columbium occurs in minerals only as the pentoxide Cb<sub>2</sub>O<sub>5</sub>, in combination with a base, the columbium in the few minerals (such as wohlerite) which contain silica, being present as a silico-columbate, and never as a silicate of

columbium

A list of thirty or forty minerals in which columbium is commonly found in association with tantalum might be given, but the principal will be given in the article TANTALUM, with which metal columbium invariably occurs in From the technical point of view, the principal columbium minerals are those which consist essentially of columbates of iron, man-ganese, and lime, zirconia and metals of the cerium group being also usually present in, or at any rate intimately associated with, such minerals as contain lime.

The mineral richest in columbium is pyrochlore, a columbate of lime containing titanium, cerium, and thorium, together with other alkaline earths and alkali bases and some fluorine The most important mineral, however, is the columbate of iron and manganese known as columbite or niobite, in which the Cb2O5 may even exceed 78 pc, a condition nearly reached by certain specimens occurring in the cryolite of Arsukfiord (Greenland).

The columbium is always partly replaced by tantalum, the proportion of which may vary from a trace through all gradations, until the mineral merges into tantalite which is isomorphous with columbite, and which occurs in much larger quantity and furnishes the bulk of the tantalum of commerce and, as a by-product, of the columbium. These two minerals are found in alluvial deposits with cassiterite and wolfram, and in isolated masses left by the denudation of

intrusions in pegmatites, &c, at Greenbushes and elsewhere in Western Australia, in Colorado (Cañon City), Dakota (Black Hills), North Carolina and other districts of the United States. and in Sweden, Norway, Greenland, Madagascar, Rhodesia, and many other districts. They may be considered to possess the general formula (FeMn)O(CbTa)<sub>2</sub>O<sub>5</sub>, the relative proportions of ferrous and manganous oxides varying within extremely wide limits

New deposits are constantly being discovered, and columbium, like many others of the rare metals for which a demand now exists, will doubtless be found to occur in much larger

quantities than is generally believed

Amounts, far in excess of present requirements, are obtainable as by-products from the preparation of the rare earths employed for incandescent mantles, from tin and wolfram ores, and from the heavy waste material obtained

in the dressing of cryolite

Metallic columbium may be prepared by passing a mixture of the vapour of the pentachloride and hydrogen through a red-hot tube (Roscoe), by electiolysis of the oxyfluoride; in a less pure condition, by heating the pentoxide with carbon in the electric furnace (Moissan), by the alumino-thermic process, and by the method of von Bolton, introduced with a view to the manufacture of metal filament lamps, before tantalum came into use. In this process, the pentoxide is mixed with paraffin and formed into threads which are heated in vacuδ to a temperature of over 1900° by passage of an electric current

By the alumino-thermic reaction, von Bolton has obtained columbium containing about 3 p c of aluminium which can be driven off with other impurities by prolonged heating in vacuo in the electric furnace The purest columbium hitherto prepared in bulk is said to have been thus obtained Like tantalum, it can also be obtained by reduction of the fluoride K2('bF7 with potassium, as a black powder from which the residual hydrogen may be expelled by shaping it into small cylinders and melting in vacuo in an electric furnace

Although the most important use of tantalum is for manufacture of electric lamps, in which case it requires to be separated from columbium, the amount actually consumed for the purpose is extremely small. The standard tantalum lamp, introduced in 1904, contained only one twenty-thousandth of a pound of tantalum, and those now manufactured contain much

The method employed for the separation of columbium from tantalum in the preparation of the latter element for electric lamp filaments probably depends essentially upon a fractional crystallisation of the mixed double fluorides of potassium with tantalum, columbium, and

Metallic columbium melts at 1950°, and has a sp.gr of 127, and an electrical resistance of 0.1870 ohm for a wire 1 metre long by 1 mm. diameter It is less malleable or ductile than tantalum, but the rolled metal has a hardness greater than that of wrought iron, and may be welded at a red heat.

Although the powdered metal oxidises rapidly when heated in air, the rolled or cast

COLUMBIUM.

metal is but little affected, as a protective coating of oxide rapidly forms. It is practi cally unaffected by an acid except hydrofluoric acid, or by solution of the alkalis, but is rapidly attacked by fused alkalis or alkaline nitrates

It combines with carbon when heated, and at dull redness rapidly absorbs hydrogen or nitrogen and becomes brittle, the hydride and nitride respectively being apparently formed.

It reacts violently when heated with sulphur or selenium, and forms the volatile penta chloride when heated to redness in chlorine

It does not form an amalgam with mercury, but alloys with many other metals and in all

354

proportions with iron
The compounds of columbium have not yet been applied commercially, but the pentoxide, generally known as columbic acid, and the fluorides, especially the double fluoride with potassium 2KF CbF, are of interest in connection with the preparation of the pure com-

pounds or metal from nunerals.

For methods of analysing columbites and tantalites, and of estimating columbium in presence of tantalum, see G Chesneau (Compt rend 1909, 149, 1132, J Soc Chem. Ind 1910, 29, 93), J Moir (J Chem. Met and Min Soc S Africa, 1916, 16, 189, J Soc Chem Ind 1916, 35, 637), Giles (Chem News, 1909, 99, 1), Simpson (Chem News, 1909, 99, 243), Metzger and Taylor (Zeitsch anoig Chem. 1909, 62, 383), Weiss and Landecker (Zeitsch anorg Chem. 1909, 64, 65), cf Foote and Langley (Amer J Sci. 1910, [iv] 30, 401), cf. Hauser and Lowite (Zeitsch angew. Chem. 1912, 25, 100), Meimberg (Zeitsch angew Chem 1913, 26, 83), Meimberg and Wintzer (Zeitsch angew Chem 1913, 26, 157), Levig (Analyst, 1915, 40, 204)

For a method of estimating columbium and tantalum based on the difference between the densities of the oxides of columbium (4.552) and tantalum (8.716), see Foote and Langley (Amer J Sci 1910, [iv ] 30, 393), see also Ruff and Schiller (Zeitsch anorg. Chem 1911, 72, GT. H 329).

## Compounds of Columbium.

Columbium dioxide CbO or Cb2O2, formed by heating potassium columbium oxyfluoride with sodium, is a white powder, which burns when heated in the air, forming the pentoxide In chlorine it ignites, forming the oxychloride CbOCl<sub>3</sub>. Is soluble in hydrochloric acid with evolution of hydrogen.

Columbium tetroxide CbO, or Cb<sub>2</sub>O<sub>4</sub>, a black powder, formed by heating the pentoxide in hydrogen, or with magnesium. Burns when

heated in the air to redness

Columbium pentoxide  ${\rm Cb}_2{\rm O}_5$ , a white infusible powder of sp gr 4.53-4.568 (4.8 Balke and

Smith).

Columbic acid HCbO<sub>3</sub>, a white powder formed by decomposing the oxychloride or pentachloride with water. Slightly soluble in hot hydrochloric acid. Yields the pentoxide when strongly heated. Potassium hexacolumbate K<sub>8</sub>Cb<sub>6</sub>O<sub>19</sub>,6H<sub>2</sub>O, formed by fusing the pentoxide with potassium carbonate. Large monoclinic crystals, efflorescing on exposure to dry air. Other potassium columbates are K<sub>6</sub>Cb<sub>4</sub>O<sub>13</sub>,13H<sub>2</sub>O; 2K<sub>2</sub>Cb<sub>4</sub>O<sub>11</sub>,11H<sub>2</sub>O; K<sub>4</sub>Cb<sub>2</sub>O<sub>7</sub>,11H<sub>2</sub>O Sodium columbate 7Na<sub>2</sub>O,6Cb<sub>2</sub>O<sub>5</sub>,32H<sub>2</sub>O (Bedford, J. Amer. Chem. Soc. 1904, 26, 1235); Sodium metacolumbate Na<sub>2</sub>O,Cb<sub>2</sub>O<sub>5</sub>,7H<sub>2</sub>O (Balke and Smith, J. Amer Chem Soc. 1908, 30, 1637) For the composition of other columbates, see Balke and Smith (lc)

Percolumbic acid HCbO4, an amorphous vellow powder, obtained by treating columbic acid with hydrogen peroxide Is decomposed on warming with sulphuric acid Potassium percolumbate or pyrocolumbate K<sub>4</sub>Cb<sub>2</sub>O<sub>11</sub>,3H<sub>2</sub>O, formed by mixing hydrogen peroxide with a solution of potassium columbate and adding Decomposed by water Yields hydroalcohol gen perovide and ozone with concentrated For other percolumbates, see sulphuric acid Balke and Smith (l c)

For the hydrosol's of columbic acid see Hauser and Lewike (Zeitsch angew Chem

1912, 25, 100)

For the electrolytic reduction of columbic acid and its salts, see Ott (Zeitsch-Elektrochem

1912, 18, 349)

When columbic acid is dissolved in concentrated sulphuric acid, the solution diluted, and then reduced with zine, an indigo-blue solution is obtained which reduces a solution of copper sulphate to metallic copper, the solution is thus more strongly reducing than one containing tervalent titanium.

Columbium pentafluoride  ${\rm CbF_5}$ , colourless, strongly refractive monoclinic prisms,  ${\rm D^{18}\,3\,293}$ ; 75.5° (corr), bp 217°-220°/760 mm (corr ) Obtained by the action of fluorine on Very hygroscopic; readily soluble columbium in water Solution precipitates columbic acid on addition of ammonia or sodium hydroxide (Ruff and Schiller, Ber 1908, 42, 492)

Columbium oxyfluoride CbOF, forms double salts with other metallic fluorides, of which the alkaline salts have been investigated by

Marignac.

Columbium trichloride CbCl3, a non-volatile, non-deliquescent crystalline substance resembling iodine, forming long dichroic needles Oxidised on exposure to air, and yielding columbium oxychloride CbOCl<sub>3</sub>, and carbon monoxide when heated in carbon dioxide (Roscoe)

Columbium pentachloride CbCl<sub>s</sub>, vellow needles, obtained by heating columbium pentoxide mixed with charcoal in a stream of chlorine Also formed by heating the pentoxide with sulphur monochloride S2Cl2, or, together with the oxychloride, by heating the pentoxide with carbon tetrachloride; mp 194°; bp. 240 5°; sp gr 2·73-2·77. Gives a normal vapour density. Soluble in carbon tetrachloride, chloroform, and alcohol. Its solution in hydrochloric acid gelatinises, and on addition of water, forms columbic acid. On reduction with sodium amalgam gives chlorocolumbrum chloride (Cb<sub>6</sub>Cl<sub>12</sub>)Cl<sub>2</sub>,7H<sub>2</sub>O, black, shining crystals, forming an olive-green powder, insoluble in cold water, but soluble in hot water to an olive-green solution Only two of the chlorine atoms are ionic. When treated with the equivalent amount of sodium hydroxide, chlorocolum-brum hydroxide (Cb<sub>6</sub>Cl<sub>12</sub>)(OH)<sub>2</sub>,8H<sub>2</sub>O is formed as a black microcrystalline precipitate, and from this the bromide  $(Cb_6Cl_{12})Br_2,7H_2O$  is obtained. The hydroxide is soluble in excess of sodium hydroxide, and from the solution excess of concentrated hydrochloric acid precipitates a brown powder  $C_6H_{14},9H_2O$  (Harned, J. Amer Chem. Soc. 1913, 35, 1078)

Reduction of a solution of columbium pentachloride at a platinum cathode gives a blue solution, which is of a colloidal character; with gold choude it gives a magnificent purple liquid, from which, on boiling, a purple precipitate deposits. This precipitate is similar to purple of Cassius; it colours glass a violet-red (Styhler, Ber 1914, 47, 841).

Columbium oxybromide CbOBr<sub>3</sub> and pentabromide CbBr<sub>5</sub> also exist, and an iodide has been formed (Bair, J. Amer Chem. Soc. 1908,

30, 1668)

**Columbium sulphides** CbS<sub>2</sub> and Cb<sub>2</sub>S<sub>5</sub>, corresponding to the oxides, have been prepared (Biltz and Gonder, Ber 1907, 40, 49, 63).

Columbium oxysulphide CbOS<sub>3</sub>, a black crystalline powder formed by heating the pentoxide in the vapour of carbon disulphide mixed with carbon dioxide

Columbium nitride  $\mathrm{Cb_2N_5}$ . A yellow powder formed by heating columbium in nitrogen (Moissan, Compt 1 end 1901, 133, 20, Muthman, Weiss and Reidelbauch, Annalen, 1907, 225, 529

The spectrum of columbium contains thirty-one more or less complete groups of five lines showing wave-number differences equal to 720 25, 1027 02, 355 19, and 303 29 respectively (cf. Siogbahn and Freman, Ann. Physik 1916, [iv.] 49, 611)

For are spectrum of columbium, see Hildebrand (J. Amer. Chem. Soc. 1908, 30,

1672)

For the magnetic resolution of the spectrum lines of columbium, see Jack (Proc. Roy. Irish Acad. 1912, A, 30, 42)

COLZA OIL v RAPE OIL

COMENAMINIC ACID v Bone oil.
COMMON CAMPHOR v CAMPHORS.
CONCRETE v. CEMENT.

CONDENSERS. A condenser is an apparatus in which the vapour of a liquid may be cooled down to a temperature below that at which it is converted into the liquid state under the prevailing pressure The condensation of the vapour is usually effected by bringing it into contact with a cold surface, to which it imparts its heat, suitable arrangement being made for conveying away this heat from the cooling surface. Although air cooling is sometimes depended upon for this latter purpose, the arrangement usually adopted consists in causing a stream of cold liquid (water in the majority of cases) to circulate over the exterior of the con-densing chamber The following points should be considered in the designing of an efficient condenser :-

(a) The cooling surface must be made of material upon which the vapour has no chemical action, and should be capable of withstanding repeated variations of temperature without fracture

(b) The area of cooling surface should be as great as possible, without making the apparatus unduly large and heavy.

(c) The material should preferably have a

high specific heat, so that its rise in temperature may be kept within reasonable limits, and it should have a high thermal conductivity, in order that the heat it absorbs may be rapidly carried away

(d) The design should be such as to admit of

the condensor being readily cleansed.

It is not possible to obtain materials entirely satisfactory in each of the above respects, and so in practice the materials employed are those which combine cheapness with reasonable satisfaction in the foregoing particulars. For laboratory apparatus, the material commonly used is glass, metals (block tin, copper, &c) are less often employed, except for constructing fairly large pieces of apparatus, of iceent years, silica has found a limited use.

Condensers are used in two different kinds of operations. (1) for condensing vapours in distillation processes, in which case a still is attached to the upper end and a suitable receiver to the lower end; and (2) as 'reflux' apparatus in cases where it is necessary to boil a liquid for some time without sensible loss through evaporation, the vessel containing the boiling liquid is then attached to the lower end of the condenser and the condensed vapours flow back into the vessel. A number of condensers described below are specially designed for this

ourpose

The simplest type of condenser is the familiar form devised by Weigel in 1771, but usually known as Liebig's condenser It consists simply of a straight cylindrical glass tube, in which the vapour is condensed. The upper end is widened out slightly to facilitate connection with the still, and its lower end is cut off obliquely, a cylindrical glass jacket is slipped over the greater portion of the length of the tube, and through it, by appropriate entrance and exit tubes, cold water is circulated in an upward direction, so that the vapours passing down the inner condenser tube encounter a surface the temperature of which continuously falls in the direction of the receiver This type of condenser admits of considerable variation and modification in the precise details of construc-

tion, thus, with a view to exposing a large cooling surface, the portion of the condenser within the jacket is sometimes made of approximately rectangular section, thus

, being broadened considerably in one direction; in other condensers the central portion carries a series of bulbs blown on it Cooling surface is obtained in some condensers by replacing the central portion by a closely wound glass spiral, as in Fig. 1, the water-jacket being sealed on to the condenser proper. The modified Weigel condenser, shown in Fig. 2 (Chem. Zeit. 1902, 26, 633), is arranged for use with the two

limbs a and c of the inner U-tube





Fig. 1.

on the same level to prevent accumulation of liquid at b. The inner tube is fused into the outer tube at d.

Glass condensers with water-jackets sealed on are restricted in their use to the condensation of the vapours of liquids of fairly low boilingpoint (below 100°), since otherwise the glass joints are liable to crack For liquids of moderately high boiling-point (140°-200°), it




usually suffices to employ merely the inner tube

of a Weigel (Liebig) condenser (an condenser) In the reflux condensers illustrated in Figs 3 and 4 (J. Soc Chem. Ind. 1897, 16, 979) the relative positions of the vapour and cooling stream of water are reversed, the water entering as shown by the arrows (at a in Fig. 3) large area of exposed surface is obtained in these condensers, which, it will be observed, are of the 'double surface' type, since, in addition to the inner water-cooled surface, condensation also occurs on the outer surface, which exposes an even larger area to the cooling action of the an

Several vertical condensers have been devised by Friedrichs in which one of the two surfaces between which the vapours condense is constructed spirally (Zeitsch angew Chem 1910, 23, 2425)

A number of extremely useful, compact condensers of the 'double-surface' type have been described. The vapours are condensed in the annular space between two surfaces, both of




which are water-cooled, the design being such that one stream of water flows in succession over each The efficiency of this cooling process is so great that these condensers can be made quite short; they are readily fixed in a vertical position and occupy little bench space.

In the Burgess condenser (J Soc Chem Ind. 1905, 24, 1190), the tubes a and c (Fig. 5) are the cooling jackets, and b is the condensing tube

two concentric tubes joined together at the top; condensation is effected in the interior chamber thus formed. Water is conveyed into the inner tube by the tube a; overflowing, it passes round the exterior of the outer tube and flows away by the side tube b The vapour is led in through the inlet tube c

Another double-surface condenser of simple construction is illustrated in the article Analysis (Vol 1)

The condensers illustrated in Figs. 7 and 8




have the advantage of exposing four cooling surfaces to the vapour, and may be readily cleansed when necessary by removing the inner portions with the stopper (Chem. Zentr. 1908, ii 277)

An inquiry into the relative efficiency of laboratory reflux condensers made by Dover and Marden (J Ind Eng Chem 1916, 8, 834) showed that where a long condenser can be conveniently used, the Weigel is preferable, whilst the Friedrichs, or some modification of it, seems best where a short condenser must be used.

The condensers so far described are usually constructed of glass, but occasionally metals are employed. The use of aluminium for the inner tube of a Weigel condenser has been advocated by Norton (J Amer Chem Soc 1897, 19, 153) Many agricultural chemists, who have to deal with numerous ammonia distillations, prefer block-tin condensers; they are constructed like inverted U-tubes, one arm (the condenser arm) being much longer than the other and carrying a water-jacket if required, while the short arm is attached to the distilling A convenient method of constructing a number of such condensers in series to be cooled by one stream of water is described in J. Amer. Chem Soc 1906, 28, 999

A useful metallic condensor is the 'ball' type, used on the reflux principle in numerous experiments involving extractions with volatile solvents. Soxhlet's pattern is shown in Fig. 10. while the construction of a double-surface condenser of this pattern is sufficiently indicated in proper. Cribb's condenser (Fig 6) consists of Fig. 9. Storch's metallic reflux condenser is depicted and described in J. Soc. Chem. Ind. 1897, 16, 979.

The spiral (worm) type of condenser is frequently constructed of metal (copper) and enclosed in a metal tank which can be filled with cold water: these condensers find considerable use in small plants for preparing dis-



tilled water For this latter purpose, a novel form of condenser is described in Eng. Pat. 6916, 1905, which combines efficiency with extreme simplicity of construction The condenser, which is placed vertically above the evaporating chamber, consists of two metallic truncated cones, the lower being open at the top, while the upper is closed by a dome-shaped The outer surface of the upper cone is cooled by a continuous stream of cold water, and condensation is effected upon its inner surface. The condensed vapour collects in the annular space between the two cones, and the water is continuously run off into large earthenware jars Special apparatus, devised for preparing pure distilled water (so-called 'conductivity water') on a laboratory scale are described by Bousfield (Chem. Soc. Trans 1905, 87, 740); Golding (J. Soc. Chem. Ind 1906, 25, 678), and by Haitley, Campbell, and Poole (Chem Soc Trans 1908, 93, 428).

Condensers of various designs are described m J pr. Chem. [2] 34, 427; [2] 49, 44, Compt rend. 118, 478; Ber 1892, 24, 3950; 1895, 28, 2388, J Soc. Chem Ind 1899, 18, 707; Chem Zeit. 1902, 26, 325, 1904, 28, 598, 686, 1905, 29, 809, 1910, 34, 116; Bull Soc chim. 1904, 31, 1116; Chem. Zentr. 1906, n. 993; and reflux condensers in Chem. Zeit. 1896, 20, 462, 1904, 28, 598; Bull Soc chim 1901, 25, 476, 1908, 3, 855; Zeitsch anal Chem. 1901, 40, 769, Chem. Zentr 1906, n. 993, Chem News, 1907, 95, 52; and J Soc Chem. Ind 1908, 27, 962

CONDURANGO. The dried bark of Marsdenia condurango (Nichols) (Gonolobus Condurango triana, Reichenbach; family Asclepradaceæ). It is obtained from a half-climbing The bark comes or prostrate shrub of Ecuador into commerce in quills or quilled pieces, about 10 cm. long, and rarely exceeding 2 cm in diameter; externally brown grey, roughened, with scale like warts; internally, greyish or pale brownish-white, striate; fracture short, granular or slightly fibrous in inner layer. Taste, somewhat bitter; odour of fresh bark, aromatic.

Kubler (Arch. Pharm. 1908, 246, 620) states that the principal constituents of the bark are dextrose, other usual components of plants, the glucoside condurangin and a polyhydric alcohol

condurated. Condurangin C40H60O16 contains two methoxyl groups, is soluble in water, alcohol, and chloroform, insoluble in other. Dilute sulphuric acid hydrolyses it to dextrose and an amorphous powder, containing several substances; alcoholic potash yields some cinnamic acid. Conduitol  $C_6H_{10}O_4$  crystallises from hot alcohol in colourless prisms, and yields a tribenzoyl derivative, m p 142°-143°. According to Carrara (Gazz chim. ital 21, 1 204, 22, 1 236), commercial condurangin can be separated into two parts, one insoluble and the other soluble in water The former is a white powder,  $C_{20}H_{32}O_6$ , mp  $60^\circ$ - $61^\circ$ , and the latter a yellow substance,  $C_{18}H_{24}O_7$ , mp  $134^\circ$ . Hager (Handbook, 1900) has further investigated these substances and named them a- and b-condurangin; a-condurangin gives a green colour with Froehde's reagent (cone sulphuiic acid and molybdic acid), b-condurangin does not rara (l.c) also isolated a yellow powder,  $C_{30}H_{50}O_2$ , which he named conduransterm, mp 52 Fluckiger has isolated a small quantity of an alkaloid having a strychume-like action. Fulbas alkaloid having a strychnine-like action (Pharm. J 70, 417) treats the chloroform extract of the crude glucoside with a mixture of equal volumes of sulphuric acid, hydrochloric acid, and alcohol (Lafon's reaction) On warming, a green colour is produced, changing to bluish-green on the addition of a trace of ferric chloride Certain other glucosides give this reaction, but they are not likely to be present in the bark

Condurango has been employed for many years in South America as an alterative in the treatment of syphilis. It is used as a stomachic, and lessens gastuc catarrh and relieves the vomiting and nausea which are usually associated with caremoma of the stomach, it is also used to lessen the pain consequent upon gastric ailments, Condurango produces in large amounts incoordination of the muscles and convulsions, which may be followed by para-Dose, I fluid drachm of the fluid extract lvsis of condurango, or 4 fluid drachms of the wine of

condurango

There are many other varieties of condurango and marodenias The Guayaquil condurango has several peculiarities in the wood of the bank, and has not so many medullary rays Mexican condurangos contain a strong collenchyma next to the bark, while nearer the pith is a layer of bast fibres containing large crystals of calcium oxalate (Chem Zeit 11, 1601; 12,

CONDY'S FLUID v MANGANESE. CONESSINE or WRIGHTINE  $C_{24}H_{40}N_{2}$ exists in the bark and seeds (0 6 p c ) of Wrightia antidysenterica and in Holarihena africana (A DC), H congolensis (Stapf), and H anti dysenterica (Wall) Crystals, mp 122° (Polstorff, Ber 1886, 19, 78, 1682; Wainecke, ibid 1886, 19, 60, Arch Pharm 1888, [iii] 26, 248, 281; Pyman, Chem Soc Trans 1919, 163; Giemsa and Halberkann, Arch Pharm. 1918, 256, 201, Ulriei, Arch. Pharm. 1918, 256, 57). Soluble in acetone, from which it separates in large colourless plates, efflorescing in air; m p. 125° (coιr) [α]-1 90°

CONGO BLUE, BROWNS, CORINTH, FAST BLUE, ORANGE, REDS, RUBINE, CONGO -SKY BLUE v. Azo- colouring matters.

CONHYDRINE (Conydrine), Coniceine. See under Coniine, below.

CONIFERIN v GLUCOSIDES.

### CONIINE AND THE ALKALOIDS OF HEM-LOCK.

Conline  $C_sH_{17}N$  is one of several poisonous, volatile alkaloids found in hemlock, Continumaculatum (Linn) (0.2–0.9 p.c. in the not quite tipe seeds). Hemlock (Grande ciquë, Fr; Schierling, Ger) is a tall biennial of the temperate regions of Europe, Asia, and America, and was the essential ingredient in the potion administered to condemned criminals by the Greeks Besides contine, hemlock contains N-methyl-decontine, N-methyl-1 contine,  $\gamma$ -contictine, conhy-

drine, and pseudoconhydrine

Preparation — Comme is most readily separated from the crushed, unripe hemlock seeds by digesting them with water made alkaline with sodium carbonate, to decompose the conine malate, and distilling the mixture in a current of superheated steam. The distillate contains the comme, together with much ammonia derived from the ammonium salts present in the seeds The liquid is neutralised with hydrochloric acid and evaporated to dryness The dry residue is extracted with absolute alcohol, when most of the ammonium chloride remains undissolved, whilst the alkaloidal hydrochlorides go into solution If ether is added to the alcoholic solution a further quantity of ammonium chloride is separated. The residue left by the evaporation of the alcoholic solution is dissolved in cold water, ether added, and the comme liberated by the addition of potash. The ethereal layer is decanted, washed, dried with potassium carbonate and distilled at a low temperature, when the comme remains as an only liquid. It may be purified from the accompanying conhydrine by fractional distillation, the comme passes over first So prepared the comine contains y-coniceine from which it may be freed by conversion into the hydrochlorides and extracting these when dry with acctone, in which the γ-coniceine salt is soluble and the comme salt insoluble the separation of the subsidiary alkaloids by benzoylation, see von Braun (Ber 1905, 38, 3108)

Properties — ('olourless, alkaline liquid, having a remarkable, penetrating odour and a burning taste, b p  $166^{\circ}-167^{\circ}$ , m p  $-2^{\circ}$ , rel. den 0.8626 at  $0^{\circ}$ , dextrorotatory [a]<sub>p</sub>=  $+13.79^{\circ}$ 

Comme is slightly soluble in cold water, but less so in hot water, so that a clear, cold solution becomes turbid when warmed It dissolves about one-quarter of its own volume of water at the ordinary temperature. When boiled with water the alkaloid distils with the steam Conine mixes with alcohol in all proportions, and is also readily soluble in other, but only sparingly so in chloroform or carbon disulphide Exposed in the air it darkens and slowly resim-When administered internally confine acts as a powerful poison. It is a monacidic base The ordinary salts are crystalline and soluble in water or alcohol. The hydrochloride B HCl forms rhombic crystals and melts at 220° The aurichloride B HAuCl<sub>4</sub>, m p. 77°, forms minute yellow needles.

Reactions. -- Conline is a strong base, which

precipitates many metallic hydroxides from solutions of their salts.

It is a secondary amine. When oxidised with nitric acid it yields normal butyric acid among other products. When permanganate is used a-pyridinecarboxylic acid (picolinic acid) is obtained. On distillation with zinc-dust confine is decomposed into hydrogen and a-n-propylpyridine.

Ladenburg has inferred from these and other reactions that comine is a-propylpiperidine and has confirmed this by effecting the synthesis of the alkaloid by hydrogenating (with sodium

ın alcoholic solution) α-allylpyridine

 $C_5H_4N(C_3H_5)+8H=C_5H_{10}N(C_3H_7)$  The liquid thus produced possesses all the chemical and physical properties of the natural alkaloid except one, it is optically mactive (Ber. 1886, 19, 2579, 1906, 29, 2486). Ladenburg has also succeeded in resolving the mactive comine into dextrorotatory and lævorotatory forms and has shown the dextro compound to be identical with the natural alkaloid. This

was the first synthesis of a natural alkaloid. The toxicological recognition is somewhat difficult, in particular there are no reactions distinguishing it sharply from incotine. Auric and platinic chlorides do not precipitate a 1 p c aqueous connine solution, but do precipitate nectine at much greater dilution. A trace of conline evaporated in a desiccator with a drop of hydrochloric acid readily yields a crystalline hydrochloride, nicotine does not. Differences in odour, density, and solubility in water may also be used.

N-Methyl-d-conine  $C_8H_{16}(CH_3)N$  is stated to occur in hemlock (v-Planta and Kekulé, Annalen, 1854, 89, 150), and remains with  $\gamma$ -conceine when this is separated from comine by fractional crystallisation of the acid tartrates—It has been synthesised by methylating comine with potassium methyl sulphate (Wolffenstein, Ber. 1894, 27, 2611, cf. Passon, ibid—1891, 24, 1678). It is a colourless, only liquid, b p—173°—174°, closely resembling comine

N-Methyl-l-conline was obtained by Ahrens (*ibid* 1902, 35, 1330) from residues accumulated

in the preparation of conine.

Coniceines. This name has been applied to a series of isomeric bases, C<sub>8</sub>H<sub>15</sub>N, obtained in various ways from the hemlock alkaloids. Only one of them γ-coniceine (2-propyl-Δ'-tetrahydropyridine) occurs in hemlock and was isolated from crude comine obtained therefrom by Wolffenstein (Ber 1895, 28, 302); occasionally as much as 70 pc of the crude alkaloid was γ-coniceine. This base was first prepared by Hofmann by the action of alkalis on bromoconine. It strongly resembles connine in odour and appearance, boils at 171°−172° under 746 mm. pressure, yields crystalline salts, and furnishes connine on reduction. It has the constitution

$$\begin{array}{cccc} \operatorname{CH}_2 & \operatorname{CH}_2 \\ \operatorname{CH}_2 & \operatorname{CH}_2 & \operatorname{CH}_2 \\ \operatorname{CH}_2 & \operatorname{CH}_2 & \operatorname{CH}_2 & \operatorname{CH}_2 \\ \operatorname{NH} & \operatorname{NH} \\ \operatorname{\gamma-Coniceine} & \operatorname{Conine} \end{array}$$

Cf Hofmann, Ber. 1885, 18, 16, 27, 112; Lellmann, Annalen, 1890, 259, 127; Loffler and

colleagues, Ber. 1909, 42, 94, 107, 116, 124, 948, 3420; Gabriel, *ibid*. 1909, 42, 4059.

Conhydrine C<sub>2</sub>H<sub>17</sub>ON, also found in hemlock, may be separated from contine with which it occurs either by cooling the concentrated ethereal solution of the two alkaloids, when the conhydrine crystallises out; or it may be obtained by distilling the crude contine, when it remains behind in the retort, if the temperature has not exceeded 180°.

Conhydrine is a crystalline, basic substance, mp. 121°, bp. 226°,  $[\alpha]_{\rm D}+10^\circ$ . On heating with phosphorus pentoxide, or with concentrated hydrochloric acid, it furnishes a mixture of  $\alpha$ - and  $\beta$ -conicenes (see above). According to Willstatter it is 2: $\beta$ -hydroxy-propylpiperidine (Ber. 1901, 34, 3166; cf, however, Loffler and Tschunke, *ibid.* 1909, 42, 920)

42, 929).

Pseudoconhydrine  $C_8H_{17}ON$ , which is also found in hemlock, closely resembles conhydrine. It is a crystalline, strongly basic substance, in p.  $105^\circ-106^\circ$ ; b.p.  $236^\circ$ . It is a secondary base and forms crystalline salts (Ladenburg and Adam, Ber 1891, 24, 1671). Engler and Baur regard it as  $2 \cdot a$ -hydroxypropylpiperidine (*ibid* 1894, 27, 1777), and Löffler has suggested that it is 4-hydroxy-2-propylpiperidine (*ibid* 1909, 42, 116) G B

CONIMA RESIN, CONIMENE V OLEO-RESINS.
CONQUINAMINE V CINCHONA ALKALOIDS
CONSTANTAN V NICKEL.

CONVALLAMARIN, CONVALLARIA v. Digitalis.

CONVALLARIN  $\mathrm{C_{34}H_{62}O_{11}}$  (Walz),  $\mathrm{C_{25}H_{40}O_{10}}$  (Lindner). A glucoside contained in Convallaria majalis (hly of the valley). On heating with dilute sulphuric acid it is hydrolysed to a hexose and convallarelin ( $\mathrm{C_{19}H_{28}O_4}$ ?), a crystalline substance which forms a stable monohydrate. It contains a benzene ring, but is free from ethylenic linkings, ketonic, and methoxyl groups, and contains two hydroxyl radicals; the two remaining oxygen atoms appear to constitute part of a carboxy-ester group (Lindner, Monatsh 1915, 36, 257). (†f Digitalis.

CONVICINE v VICINE.

CONVOLVULIN v GLUCOSIDES.

COOMASSIE NAVY BLUE, -WOOL BLACKS, v. Azo- Colouring matters

COPAIBA BALSAM v OLEO-RESINS

COPAIDA OIL v OILS, ESSENTIAL COPAIVENE, COPAIVI, COPAIVIC ACID v.

OLEO-RESINS.

COPAL v. RESINS.

COPAL VARNISH v VARNISH

COPALIN v. RESINS.

COPPER. Sym. Cu At. wt 63.57. Melting-point in air, 1062°; in reducing atmosphere, 1084°. Specific gravity: Roberts Austen (Int Study of Met. 1910, 15) gives 8.217 as the sp.gr. of the molten metal. Copper, deposited by electrolysis from an acid solution of copper sulphate and not subjected to any mechanical treatment, was found to have sp.gr. of 8.9544 at 15.5° in air (Watson, J. Soc Chem Ind. 2, 153)

Commercial copper, prepared in the usual way, is never pure, and the examination of the fractured surface of a cast ingot of copper shows a minutely vesicular structure. On this account the density of such metal is less than

the same sample of metal after it has been subjected to a compressing mechanical treatment, and varies from 8.2 to 8.6

Colour.—Copper has a characteristic red colour which, in a fractured surface of worked metal, may be described as salmon-red.

Lustre —A smooth cut or polished surface shows a bright metallic lustre which, however, soon tarnishes.

Crystalline system —Copper crystallises in the cubical system

Ores of copper. The following minerals are those from which copper is extracted on a commercial scale: native copper, cuprite, melaconite, malachite, azurite, chrysocolla, atacamite, chalcocite, bornite, chalcopyrite, fahl ore

The importance of the several species in the above list as ores of copper varies very much with the locality. Thus, while native copper is the chief ore in the Lake Superior district of North America, and atacamite in one or two localities in Chile, they would in other localities, and generally, be looked upon as rather rate, especially so in the case of the last species mentioned So also in the case of malachite, this species, as an ore of copper, is seldom met with except in small quantities in the upper and oxidised portions of some copper lodes, yet in the case of the once celebrated Burra Burra mine in Australia, and at Nischne-Tagilsk in Russia, it formed the bulk of the ore produced, and is also the chief ore in the Tanganyika district, Africa

Generally, in a copper-mining district, all the oxidised species mentioned above occur in varying proportions, but as depth is attained, and almost invariably at or about the level at which water is reached, the copper-bearing minerals change and the oxidised species give place to the richer sulphide species, and these finally to chalcopyrite intimately mixed with varying proportions of pyrite The intermingling of the two species last mentioned is frequently so intimate that it is not easy to distinguish the component minerals or to separate them. Chalcopyrite thus diluted down, as it were, with one or more of the mineralspyrite, pyrotite, &c —forms the ore from which the greater proportion of the world's copper supply is derived and the bulk of the material treated at the smelting works generally. Such ore frequently contains appreciable quantities of gold and silver The silver is frequently carried by intermixed fahl ore, but also, in many instances, by the pyritic mixture in which distinct silver-bearing minerals are not observable.

As by far the larger proportion of ore presented for metallurgical treatment is of the nature indicated above, viz. chalcopyrite mixed with varying quantities of the pyrite family of minerals, it will best serve the purpose of this article if the methods adopted for the extraction of copper from this material are first considered. The ores treated generally contain only small percentages of copper, frequently not more than 1 p c

The elimination of the several elements in the ore and the final production of the copper in the metallic state, are effected by two distinct series of chemical operations. One of these—the most important, because of its very general application—treats the ore by smelting, and

causes the necessary reactions to take place while the material is in a molten condition. The other-of limited application-treats the material in the wet condition, and causes the necessary reactions to take place in solutions

These two distinctly different methods of procedure may be distinguished as (a) dry

methods, and (b) wet methods

Dry methods. The object which the smelter keeps in view in the first stages of the processes is to concentrate the copper contained in the ore in as small a quantity of material as may be economical. For this purpose, the product known as 'matte' or 'regulus' is universally selected. This consists of cuprous sulphide Cu2S and non sulphide FeS in varying proportions, together with sulphides of certain The whole of the other metallic elements copper contained in the ore thus passes into the matte, while the silica combines with iron oxide and earthy bases, if present, and forms the The slag separates easily from the matte, owing to its lower specific gravity, and is thrown away Having thus concentrated the copper into regulus, or matte, the next stage of the process concerns itself with the decomposition of this matte in such manner as to eliminate the iron and sulphui and, as far as possible, other metallic elements which may be present and to yield the copper in the metallic state

The final process consists in refining the copper produced as above, or, in other words, removing, as far as practicable, the small quantities of such elements as may still be present in the product of the preceding operation which would injuriously affect the copper for

manufacturing purposes

The treatment of the ore involves, therefore, three distinct operations

I Smelting the one with the production of matte and slag.

II. Treatment of the matte for the production of crude copper

III Refining

The methods and the plant used for carrying out each of the three operations vary considerably, and the choice of any one depends upon local conditions, the nature of the ore as regards physical character, and the nature and quantity of the foreign elements present will not be possible within the limits of this article to study these considerations and the bearing they may have on the methods to be adopted, all that the space will allow of is a description of the methods as actually carried out under the most generally prevailing conditions

I Smelting the ore with the production of matte and slag -The object of this process being the concentration of the copper in the matte, the chief considerations kept in view are the production of a matte as rich in copper, and a slag as poor in copper, as is practically and economically possible

The grade of the matte to be produced having been decided upon, the smelter regulates this by suitably proportioning the sulphur in the ore to be treated so that there shall be sufficient to form Cu2S with all the copper, and a further proportion to form FeS with the iron present in such quantity as will cause the matte to contain the desired percentage of copper. From these

considerations it will be seen that the factor which regulates the grade of the matte is the quantity of sulphur present in the ore mixture smelted. With the class of ore under consideration, it invariably happens that the quantity of sulphur present is considerably in excess of requirements, and the production of the most suitable grade of matte necessitates the removal of this excess, which may be effected before the actual smelting of the ore is commenced. Under certain favourable conditions, however, it may be removed during the smelting process (see Pyritic smelting). This preliminary operation is designated 'calcining' or 'roasting.'

The process consists in subjecting the ore to

a dull-red heat under oxidising conditions, so that the sulphur is caused to combine with oxygen derived from an admitted into contact with the ore with the production of sulphur dioxide, while the non is also oxidised at the

The method of carrying out this calcining process depends upon whether the ore is to be

(a) In the rough—say in pieces from 3 inch cube upwards, or

(b) As fines or 'concentrates,' i.e. in which the particles are finely divided.

Under (a). The one may be calcined by forming heaps of it in the open air and staiting the combustion by means of a small proportion of wood, the combustion being maintained by the sulphur present in the ore.

The ore may also be calcined in stalls more or less open These consist of series of chambers in rows built back to back. The chambers are each bounded by three walls, viz back and two side walls, but are open in front and have no 100fs. Small flues, opening out into the chambers, are built in the walls separating them, and these flues connect with a larger flue in the central division wall between the two rows, and these again connect with a chimney stack

The ore is charged into these open chambers on to a layer of wood, when filled up, the wood is fired and combustion extends to the ore itself in the same way as in the case of the heaps The gases produced in the operation are to a large extent led away by the flues in the walls above referred to The annoyance caused by the escape of these gases at or near the ground level of the works—as is the case in open heap roasting-is in this way to a large extent avoided

The ore may be calcined in kilns - These are similar to the stalls, but are generally built of much larger size, are closed in on all four sides, and are roofed over They are provided with suitable openings in the roof through which the ore is introduced, and a doorway in the front wall (bricked up when the kiln is in operation) through which the calcined ore is removed. An outlet flue, connected to the roof or back wall, allows the gases to be conveyed to a chimney.


The ore may be calcined in 'burners.'-These are constructed and worked on very much the same lines as the 'burners' used for burning 'pyrites' at sulphuric acid works. In this plant, a number of closed cells or 'burners' are built in rows, back to back. Each cell is roofed in and connects at the back with a

central flue common to all the cells, by which the gases are led to the acid chambers. In the roof of each cell is an opening through which the ore can be charged from a hopper placed above. In this class of plant, the ore does not rest on the floor as it does in each of the cases previously considered, but upon a series of square bars suitably spaced. Below these bars is an enclosed space with suitable door through which the calcined ore is removed and by means of which the air supply is regulated

In operating this plant, the cell having been brought up to a good ied heat by the combustion of wood or other suitable fuel, ore in small quantities is gradually added as combustion proceeds, until the burner is full. At suitable intervals, a small quantity of the calcined ore is removed from the bottom of the

burner and a corresponding fresh supply of ore is added through the opening in the roof. In this way, the operation of the burner is rendered continuous, and no fresh extraneous fuel is required after the first starting of the fire

In all the methods of treatment described under (a), it is essential that proper care be taken to prevent the temperature reaching anything like melting-point. If the one melts of sinters together, calemation is delayed owing to the reduced surface on which oxidation can take place. Further, in the case of such sintered or melted material, difficulties are created in working the process. In the case of heap roasting and also in the stalls and kilns, the formation of large masses of sintered or melted one introduces undesirable inregularities in the air current through the mass of one under treatment, and



Figs. 1, 2, and 3.

causes trouble when the ore is removed. In the burners, the formation of sintered masses of ore must be avoided, as such masses would have to be broken up in the burner with suitable tools before it could be made to pass out between the grate bars.

Under (b) The ore may be calcined in hand calciners or mechanical calciners—The former are not much used at the present time, as the necessary labour is effectually replaced by mechanical appliances. The hand calciner, which was universally used some years ago, consisted of a long flat bedded reverberatory furnace, with the usual fire box at one end and the outlet flue for the exit of the gases at the opposite end (Figs. 1, 2, and 3). The ore was dropped from a hopper through an opening in the roof on to the bed at the flue end of the furnace, where it was spread out by means of


suitable tools worked by men at both sides of the furnace, operating through doors in the side walls

The conditions in the furnace were intensely oxidising, and it was necessary to stir the ore frequently to prevent it sintening. In the process of stirring, the ore was turned over and moved towards the fire end, thus making room for a fresh charge from the feed hopper. This process of stirring and moving the ore forward towards the fire end was continued until the ore reached close up to the fire box, at which point it was discharged through openings in the bed of the furnace into vaults, or chambers, constructed below, from which—when sufficiently cool—it was removed. The labour required on these furnaces was very heavy and the output small, and, except in unimportant installations, they have given way to furnaces in which the

ore is stirred, moved forward, and discharged by mechanical means operated by power. It will be impossible within the limits of this article to describe the many furnaces of this type which have been introduced, and it will suffice to describe one or two which will serve as examples of the class of plant employed

The McDougall furnace (Fig. 4) —This is very largely used, and consists of a series of flat

circular superimposed firebrick hearths, enclosed in a brick-lined cylinder of steel plate An iron shaft passes centrally up the cylinder through openings in the several hearths, and is



(From Gowland's Metallurgy Charles Griffin & Co , Lt1)

caused to revolve by suitable gearing attached either to its upper or lower end. This shaft may be made hollow and cooled by air or water circulation.

One or more iron arms are attached to this shaft above each bed The arms extend from the shaft to the periphery of the circular beds, and have attached to them a series of iron plates or ploughs set at a slight angle to the arm. The ore is fed from a hopper on to the top bed near the centre, and the angle of the ploughs is such that, as the arms revolve, they not only turn the ore over, but gradually move it from the centre to the edge of the bed. At one point in the edge of the bed is an opening through which the ore

falls on to the second bed. On this bed the ploughs are so set on the arms that the ore travels in the reverse direction and is drawn to the centre of the bed. Here it discharges on to the third bed, and so on until it escapes at the bottom of the furnace. The openings through which the ore falls from one bed to the other also serve as flues through which the gases pass upwards through the furnace, finally escaping through a flue above the top bed.

Attached to the lowest bed is a fire box by which the furnace is heated up in order to start it, but, after being once started, little or no fuel is required, as the oxidation of the ore provides sufficient heat throughout the apparatus

The straight-hearth mechanical furnace—This type of furnace passes under various distinctive names, according to the several methods adopted for operating the ploughs, and slight modifications in the construction furnace consists of a flat bed about 100 feet At one end is the long covered with an arch inlet for the ore, and at the opposite end is the outlet. At the ore inlet end is the flue for carrying off the gases, and at the opposite end, but to one side, is a fire box with the inlet to the furnace constructed in the roof of the latter similar fire boxes—one or two in number—are placed at intervals along the length of the A channel is constructed in the side walls of the furnace and in these a rail is laid. A suitable carriage with wheels iesting on the iails stretches from one side of the furnace to the other, and the bearings pass through slots in the two side walls and are carried by the wheels above referred to To this carriage, suitable ploughs are attached in such manner that, in the course of its travel from one end of the furnace to the other, the ploughs turn the ore over and also move it forward. The carriage is moved by a pair of chains or ropes travelling in the channels formed in the side walls, and at the two ends of the furnace these chains or ropes are carried round wheels sufficiently large in diameter to lead them back under the furnace, thus making them continuous. One pair of these wheels is driven by suitable gearing, thus causing the pair of ropes or chains-and with them the carriage -to travel the whole length of the bed of the furnace, returning through the tunnel under the bed, where rails are laid for supporting the carriage Two or more such carriages are attached to the ropes, or chains, at equal distances apart. The ore is fed mechanically on to the bed at intervals just as a carriage is about to enter, and the ore is spread, turned over, and moved forward gradually from one end of the furnace to the other. Each time a carriage emerges from the furnace, it draws with it a quantity of calcined ore The carriage and its attached ploughs are cooled during the passage through the tunnel The ends of the furnace are closed by light iron doors which hang down vertically and are automatically opened by the passage of the carriage and close themselves as soon as the carriage has passed.

There is yet another method of removing the excess sulphur which is sometimes used, known as 'pot roasting,' or 'blast roasting.' This method of calcining is advantageously applied to concentrates and fines, and, while it eliminates the sulphur, the heat attained is at the same


time sufficiently intense towards the end of the process to cause the ore to partially melt or sinter together, forming a mass which requires to be broken up before it can be treated in the smelting furnace. The process is carried out in a similar manner to that used in the treatment of lead ores, but is not very extensively used in the metallurgy of copper.

After the sulphur in the ore mixture has been reduced to the requisite amount, the ore is ready to be smelted. For this operation one of two distinctly different types of furnace are used,

these are:

(1) Blast furnace. (2) Reverberatory furnace

The choice between these two methods of carrying on the smelting operation depends on a



variety of circumstances, all of which must have proper consideration, but the following are the chief points, viz: If the ore is to be smelted in the state of fines or concentrates, it is unsuitable for blast-furnace treatment, while it is eminently suitable for the reverberatory furnace. On the other hand, rough ore is unsuitable for reverberatory-furnace smelting

Blast-furnace smelting—The furnace used may be built of brick, but is more generally, nowadays, constructed with water-cooled walls of steel plate (Fig 5), each section of which is termed a jacket. In the smaller furnaces the horizontal section is circular, but in the ordinary sized and larger furnaces the section is always rectangular. As the working of this type of furnace is—as its name implies—dependent upon an air blast projected into the interior,

the width of the furnace is limited by the strength of the blast which is required to penetiate the charge from the sides to the centre of the furnace, while the length of the furnace is only limited by the capacity required. The increase in length is obtained by clamping together two or more of the jackets on each side, thus extending the longer axis of the furnace.

The jackets stand on a thick cast-iron plate carried on non supports. In some cases these jackets are only used in the lower part of the furnace, all the upper part being constructed of brick, but the brickwork may, in certain cases, be advantageously replaced with a second tier of jackets, reaching up to the feed floor the feed floor the furnace is continued in the form of a covered-in brick chamber with lifting doors at each side and an outlet in the roof of the chamber, connecting, by means of a suitable iron flue, to the general flue system of the works Through the doors the charge, consisting of a proper mixture of ore—either raw and calemed, or partially calemed ore—is introduced. To this one mixture there is also added such flux, either limestone or silicious material, and the necessary amount of coke as is required to form a suitable smelting mixture As previously indicated, the object of this is to produce a matte of the desired grade and a slag which shall be as free from copper as possible and which shall be of such fusibility as to flow readily from the furnace On the lower or working floor, the furnace is provided, either at the ends or sides, with one or more spouts through which the molten products of the smelting operation Except in very large furnaces, only one such spout is used at a time, and if more than one is fitted, it is with the object of having one in reserve in case of a stoppage to the one in use.

The side jackets of the furnace are provided with circular openings (tuyères) passing from the outer to the inner surface. When the jackets are set up, these tuyères form a horizontal line of openings on each side of the furnace. The blast is brought down from the air main by light iron piping to the tuyères, and through these it passes to the interior of the furnace.

The blast, at a pressure of 10-42 oz. per square inch, is generated by a suitable blowing machine, which must be of the positive, and not the fan, type. A plentiful supply of water is circulated through all the jackets, and the overflow pipes from these, carrying away the hot water, discharge into sinks or launders.

The products of the smelting operation performed in this furnace flow out through the spout above described into the settler In the case of small furnaces this is a square cast-iron vessel open at the top and mounted on wheels to allow of easy replacement It is furnished with a cast-iron spout at its upper edge through which the slag overflows, and is provided with a tap hole near the bottom from which the matte is tapped from time to time. In the case of larger furnaces for this form of settler, there is substituted a circular tank, formed of steel plates lined with refractory material, placed permanently in position under the spout of the furnace. This form of settler may be 10 or more feet in diameter, and, like the smaller

pattern mentioned above, is provided with a slag spout near the top and a tap hole near the bottom.


In the settler, a thorough separation of the slag and matte takes place. the matte, having the higher specific gravity, falling to the bottom of the vessel and the lighter slag rising to the top, where it overflows through the spout into suitable pots provided for its removal to the slag dunip. From time to time the matte is tapped off through the tap hole at the bottom of the settler The working of this type of furnace proceeds continuously, fresh ore charges being dumped in at the top at short intervals, while a constant flow of slag and matte passes out through the spout

Reverberatory-furnace smelting -The furnace used is essentially different both in construction and method of working from the one previously described. In the blast furnace, the fuel—coke -is burnt in contact with the materials to be smelted, and the intense heat required for the operation is obtained by uiging the combustion of the fuel by means of an air blast. In reverberatory furnace the material to smelted occupies one compartment of the furnace, while the fuel-coal or wood--is buint The draught required for the comin another bustion of the fuel is quite moderate, and is obtained by a chimney of suitable height and area. The use of pulverised coal has recently

been introduced in reverberatory furnace practice, with gratifying results, from a fuel economy point of view The working bed of the furnace in plan is a parallelogram (Fig. 6) with rounded corners At one end, and separated from it by a low partition, is the fire box, in which the fuel is consumed. At the opposite end is the outlet flue connecting with the chimney A brick aich extends from the back wall of the fire box to the flue This arches the furnace across from side wall to side wall, and is constructed with a fall from the fire-box end to the flue end of the furnace. The partition, or bridge, as it is termed, to which reference is made above as separating the working bed from the fire box, reaches to such a height above the bed that an opening is left between its top surface and the roof of the furnace. Through this opening above the bridge, the flame from the burning fuel passes over the bed on its way to the outlet flue. The side walls of the furnace are pierced with a number of openings used in working the furnace for spreading the ore charge, drawing off the slag, &c, and one opening is cut down to the lowest point of the bed for tapping the matte

The furnace walls and roof are constructed of highly refractory bricks

with an outside casing of



with steel girders placed vertically and tied together from one side to the other by non tie rods. The working bottom is variously constructed. It may be formed of highly refractory bricks, or of silicious sand or crushed quartz

The charging of this furnace takes place through one or more openings in the roof from hoppers supported above. Assuming that the furnace is heated up to the required temperature, a charge of ore mixture is dropped from the hoppers above through the openings in the roof. This is then spread over the bed by means of suitable tools operated by men working through the side and front doors. When this operation is completed, the doors are closed and the fire attended to so as to raise the temperature of the furnace as rapidly as possible to full heat. The charge is occasionally stirred, if necessary, to remove any agglomerated masses from the bottom of the furnace, and

finally, a molten bath is obtained composed of two distinct layers—the matte forming the bottom layer, and the slag floating upon it. When this stage is reached, another charge may be dropped and treated in the same way; and this may be repeated from time to time until the furnace is full of molten slag and matte A side door and end door are opened and the slag allowed to run off, and towards the end, this is drawn off by means of tools handled by the furnace-men until nearly all the slag has been removed. The tap hole is then carefully opened, and this being at the bottom of the furnace, the matte flows through it and is led by a launder into a ladle or into moulds placed When nearly all the on the floor to receive it matte has been run off, the tap hole is again closed and made secure, fresh charges are dropped in, and the operations repeated

The slage may be received in ladles mounted

on wheels, and in these may be removed to the dump, run into a stream of water and thus granulated and removed; or it may be run into sand moulds formed on the floor of the furnace house, and, when cool, broken up and removed to the dump.

The object attained in this operation in the reverberatory furnace is the same as with the blast furnace, viz. the production of the proper grade matte and a slag of easy fusibility

reasonably free from copper.

Pyritic smelting.—This method of working secures in one operation what, in the preceding descriptions, is attained in two processes carried on in separate appliances It has been shown that in order to produce a proper grade of matte from cupriferous iron pyrite, a certain proportion of sulphur must be oxidised to sulphur dioxide, and that, accompanying this action, a portion of the iron sulphide must also be oxidised. It has also been shown that, in ordinary working, this picliminary treatment and the actual smelting of the ore are carried out as two distinct operations. It has also been indicated that in the smelting operation proper, the high temperature necessary is obtained by the combustion of carbonaceous fuel. In pyritic smelting, the two processes referred to above are carried out as one operation in one furnace, and the high temperature necessary is derived from the rapid oxidation of the sulphur and iron (chiefly the latter) contained in the ore While this method of working is of the highest importance from an economical point of view, it is of somewhat limited application, owing to the fact that comparatively few mines produce ore which—as regards its physical condition and chemical constitution—is suitable for this method of treatment

The process is carried out in blast furnaces, and for its successful operation demands a larger volume of air per unit of time than the ordinary smelting process and an ore mixture in which the pyritic material is present in fairly large pieces. The silica must be, for the most part, free silica (quartz), and not combined with earthy bases as silicates. Under these circumstances the ore may be satisfactorily smelted, producing the required matte and slag with the consumption of but 1 to 3 p.c. of coke instead of 10 to 12 p.c. used in the ordinary method of smelting. A small amount of coke is found to be desirable in order to keep the charge open

II Treatment of the matte for the production of crude copper—It has already been pointed out that matte consists of cuprous sulphide Cu,S, with varying proportions of iron sulphide FeS, and the carrying out of the stage which is now to be considered involves the climination of the sulphur and iron, while the copper is left in the metallic state. The process is carried out by different methods, the chief of which will now be considered. In all of these the operation is performed in two stages, although, in some cases, these follow one immediately after the other and in the same apparatus, while in others the stages are performed as two separate operations and in two or more separate furnaces. The first stage in each case consists in the removal of the FeS with the production of Cu<sub>2</sub>S, and the second stage treats the Cu<sub>2</sub>S with the production of metallic copper. The carrying out of the

first stage depends upon the fact that, when matte is exposed to oxidising conditions, the iron sulphide is the first of its constituents to be decomposed, the sulphir being oxidised to sulphir dioxide. The resulting iron oxide is then caused to combine with silica, forming a slag which can be removed from the residual cuprous sulphide.

The second stage in the process depends upon one or more of the reactions represented

by the following equations .-

the first being the most important as representing the change which takes place in the largest proportion of the material. The reactions indicated are effected by bringing the several compounds together in a molten condition. The oxidised compounds may be added to the molten Cu<sub>2</sub>S from outside sources or may be, and generally are, produced by the oxidation of a portion of the cuprous sulphide itself.

The reactions outlined above are, at the present day, almost always brought about by what is termed the 'converter process,' also named 'Bessemeising,' the former name derived from the designation of the vessel in which the operation is performed, viz. converter, and the latter from the name of the distinguished inventor, Sir Henry Bessemer, who had many years previously applied a similar method and plant to the production of steel from pig iron

The converter consists essentially of a vessel formed of steel or iron plate, with a thick lining of either highly silicious rock to which a small quantity of clay has been added, sufficient only to give the necessary coherence, or of basic material The vessel is provided with a suitable aperture serving the purpose of a charging opening through which the matte to be operated upon is poured in a molten condition, also as an outlet through which the products of the operation are poured, as well as being an escape for the gases produced in the process. The vessel is supported on horizontal trunnions or on fuction rollers, and is capable of being iotated by electric or by hydraulic power (Fig. 7). The form of the vessel is more or less cylindrical with the axis of the cylinder placed vertically or horizontally. In the case of the former, a shallow circular enclosed space, termed the air box, is attached on the outside at the bottom of the vessel In the latter form, this air box takes the shape of a long rectangular box attached horizontally to the outside of the converter. From these boxes holes are led through the plates forming the casing of the converter, and are continued through the lining to the Air, at a pressure of from 8 to 12 lbs per square inch, is supplied to the air boxes as required, and the supply is controlled by suitable valves. Charging of the converter with matte is effected by means of a launder direct from the furnace, or settler, or by tapping these first into a steel ladle handled by an overhead crane which conveys it to the converter and pours it

the FeS with the production of Cu<sub>2</sub>S, and the second stage treats the Cu<sub>2</sub>S with the production of metallic copper. The carrying out of the fire, is placed upon the stand and connected up

to the air main. A charge of molten matte is then poured in, the converter having been tilted into a suitable position for receiving it The air being turned on, the converter is then tilted over so that the tuyères are brought beneath the surface of the matte, and the air,




Fig. 7.

forced through being molten sulphides, causes a rapid oxidation of the constituents of the iron sulphide The sulphur is oxidised in this way to sulphur dioxide and the iron to oxide, which is immediately converted to silicate by combination with the silica of the lining

he oxidation proceeds, the temperature of the harge increases and the operation is continued ntil practically the whole of the iron sulphide as been decomposed, and the contents of the onverter consist of cupious sulphide and slag. he converter is now turned over and the slag in off into pots or moulds, and when this is one, it is again turned into the blowing position, hen the second stage of the oxidation is comienced. The passage of the air through the iolten cuprous sulphide brings about the xidation of a part of that material to cuprous xide, which immediately reacts upon unaltered iprous sulphide with the production of an juivalent quantity of metallic copper, in scordance with the reactions already given his action continues until all the cuprous sulhide has been decomposed and the contents of ie converter consist of metallic copper; any nall quantity of slag which may be on the irface is removed by turning down the vessel id skimming off. The metallic copper is ien poured by tilting the converter. As the ags produced in the operation contain a certain nount of copper they are resmelted in the rnace in which the matte was produced in the st place As soon as the copper has been oured, the converter is ready to receive another large, and the operations are repeated. When veral charges, generally 5 to 10, have been put rough (the number depending upon the nature the material of which the lining is formed and the care exercised in ramming it into place), the lining becomes too thin for further use; the converter is therefore removed to the relining department by means of the crane, the old lining is taken out and a fresh one put in, when,

Owing to the frequency of this relining necessary in the acid converters, basic lined converters have been introduced, magnesite bricks being used for this purpose The silica which is necessary to flux off the oxide of iron formed by the oxidation of the iron sulphide in the matte, is charged into the converter in the form of diled silicious ore. Very large basic lined converters are now being used and the method of working varies somewhat from that used in the case of acid lined converters. The copper from a previous charge having been poured, 30 to 40 tons of matte may be poured into the converter followed by about 3 tons of silicious ore which may itself be cupriferous. The blast is then put on and the converter slowly turned until the tuyères are below the level of the matte For the first few minutes a blast pressure of only 5 lbs is used, and after the matte and silica have thoroughly intermingled, the blast pressure is increased to about 12 lbs. per square inch. When the silica has been fluxed by the iron oxide, the converter is turned down and the slag poured off A further charge of 6 to 7 tons of matte and sufficient silicious ore are now added and the blow is resumed removal of slag and additions of matte and ore are continued until sufficient white metal or cuprous sulphide has accumulated in the converter to be blown to metal in the usual way

Another method for the production of metallic copper from the matte is conducted entirely in reverberatory furnaces, and is some-times employed where the output does not justify the more expensive converter plant previously described. In this method working, the first stage, viz. the removal of the iron sulphide of the matte by oxidation of its constituent elements, is effected by smelting the material with a proper proportion of calcined matte together with silica sufficient to form a silicate of iron slag with the iron oxide; or the matte may be smelted with oxidised copper ore, silica being also present. In either case, the result is that the iron sulphide is decomposed, the sulphur passing from the furnace as sulphur dioxide, and the iron forming, with the added silica, a slag, which is removed. The contents of the furnace at this stage consist chiefly of cuprous sulphide, Cu.S. Currents of heated air are directed over the surface of the molten bath. with the resulting formation of cuprous oxide and the concurrent reaction between this and the unchanged cuprous sulphide results in the production of metallic copper, in accordance with the equations previously set out. When this reaction has extended to the whole contents of the furnace, the heat is raised and the metallic copper is tapped out into moulds

III Refining—The method of treatment to be adopted in this operation depends upon whether (a) the copper is practically free from gold and silver and from deleterious elements; (b) the copper contains appreciable quantities of gold and silver, or contains undue quantities of

deleterious elements.

If the metallic copper produced by the operations previously described under I and II comes under the description (a) above, it can be refined by furnace treatment only; if it comes under (b) it must be refined electrolytically in order to recover the gold and silver or to obtain it free from the impurities.

Under (a), the furnace employed is of the reverberatory type, and the removal of the impurities is effected by slowly melting the pigs of copper under highly oxidising conditions, and maintaining the bath of molten copper under this condition until the impurities are oxidised and removed, either by volatilisation or by passing as oxides into the copper silicate slag formed. The oxidation of the impurities is facilitated by the fact that metallic copper has the property of dissolving a small proportion of cuprous oxide, which compound is brought into contact with the elements to be oxidised under the very favourable condition of fluidity.

The oxidising condition of the bath of molten copper is maintained until the impurities are removed, as far as practicable, and the copper contains a maximum quantity of dissolved cuprous oxide. In this condition the metal is said to be 'dry'. After removal of the slag, the next stage consists in reducing the greater portion of the dissolved cuprous oxide, which is effected by depressing poles of green wood below the surface of the bath of metal, when considerable agitation takes place and the cuprous oxide is subjected to the reducing action of the gases given off by the pole

The process is continued until only a small quantity of cuprous oxide remains, and the copper attains the physical condition known as tough pitch, when it is ladled out into moulds

of any required shape.

Under (b), the copper is refined electrolytically. This process depends for its operation upon the fact that if an electric current is caused to pass through a solution of copper sulphate, the copper in the solution is deposited at the cathode, while the acid is set free at the anode If the anode is composed of metallic copper, then the acid set free on its surface immediately dissolves an equivalent portion of copper which passes into solution, and is, in turn, deposited at the cathode, and the process thus becomes continuous so long as the current passes and there is copper at the anode to be dissolved. Assuming that the solution of copper sulphate is properly made up and maintained, and the voltage and current are suitably adjusted, then only copper will be deposited at the cathode, while the gold and silver and all the impurities contained in the copper anodes will either remain as insoluble slime, collected at the bottom of the vessel in which the operation is conducted, or will pass into the electrolyte. Gold, silver, and lead pass entirely into the insoluble slime; other elements, such as iron and nickel, pass wholly into the solution, and others again are to be found partly in the slime and partly in the electrolyte. cathode surface is, to start with, a very thin sheet of electro-deposited copper, and when-as the result of the operation-sufficient copper has accumulated upon its surfaces, it is removed from the bath and is ready for melting and casting into any required shape. In practice, a number of anode plates are hung in a lead-

lined wooden tank, and are suitably connected to the positive lead. Alternating with these anode plates are hung the cathode sheets, connected in parallel with the negative lead large number of tanks are connected in series, and the current is passed through the whole set of tanks. The electrolyte is caused to circulate through the whole series, and, after it has been in use for some time, a certain proportion must be removed regularly for purification. A quantity of anode copper remains undissolved, and when the tank is to be cleaned, this is taken out for remelting into anodes, the solution is run off, and the slime is removed. This latter is washed and dried, any large particles of copper removed by sieving, and the fine slime is then treated for the recovery of any silver or gold which it may contain

Copper flue dust. A considerable amount of copper may be carried forward in the waste gases from the furnaces and precautions are now taken to recover it. In modern well arranged plants, capacious flues and dust chambers are provided between the furnaces and the chimney for the deposition of the fine particles of ore carried forward, and for the condensation of copper fume, which is frequently arsenical. The Cottiell piocess of electrostatic precipitation of dust is sometimes used. The dust obtained is collected from time to time and retreated in reverberatory furnaces or is briquetted and smelted in blast furnaces.

Wet methods. Until recently these had a very limited application in the extraction of copper, but within the last few years have made remarkable strides, and extensive plants are now in operation in which new methods or modifications of the older methods are being used. The wet extraction of copper consists of leaching the ore either in its natural condition on after roasting and in a suitable state of subdivision, with a suitable solvent, in vats fitted with false bottoms and precipitating the copper from the solution thus obtained by suitable means.

Wet methods are specially suitable for the treatment of low grade oxidised ore and for oxidised tailings, but have also been successfully applied to the treatment of material containing the copper as sulphide. The most widely used method consists in leaching the ore with dilute sulphuric acid and recovering the copper from the solution by electrolysis or by precipitation on metallic iron. This method can be applied directly to oxidised ores, but in the case of sulphide ores a preliminary roast, in order to render the copper soluble in the acid, is frequently necessary.

At Rio Tinto the massive cupriferous pyrites is treated without roasting, large heaps of the mineral being allowed to oxidise slowly under the action of air and moisture, and the copper sulphate thus formed is washed out and precipitated on metallic iron. In some cases in which sulphuric acid leaching is followed by electrolytic deposition of the copper, the regeneration of sulphuric acid in the precipitation plant yields a solvent suitable for further leaching operations.

Another method used consists in roasting the cupriferous sulphide material with salt. Soluble copper salts are thus formed which may be recovered by solution and precipitation of the

copper on iron. A modification of this process is also used whereby silver is recovered with the

Other processes are sometimes used in which sulphurous acid, ferric sulphate, ferrous chloride and ammonia are used as solvents  $C \cup B$ 

#### COMPOUNDS OF COPPER.

Five oxides of copper are said to be Oxides. known: the quadrantovide Cu4O, the trientoxide Cu<sub>3</sub>O, the sub- or hemi-oxide Cu<sub>2</sub>O, the monoxide Other oxides CuO, and the peroxide CuO<sub>2</sub> have been described, but their existence is not proved

Copper quadrantoxide is stated to be formed by adding a solution of copper sulphate to a cooled dilute solution of stannous chloride in caustic potash (Rose, Pogg Ann. 1861, 120, 1) It is an olive-green powder, which rapidly absorbs oxygen from the air. According to Moser (Zeitsch anorg Chem 1909, 64, 200), this product is really a mixture of cuprous oxide and copper.

Copper trientoxide is obtained as a hard yellowish mass by heating cupile oxide at 1500°-2000°. It is unaffected by any acid except hydrofluoric (Bailey and Hopkins, Chem

Soc Trans 1890, 269)

Copper suboxide, or hemioxide, red oxide of copper, cuprous oxide  $\mathrm{Cu}_2\mathrm{O}$ , occurs native as cuprite or red copper ore, and as chalcotrichite

It may be prepared (1) by heating finely divided copper in an below a red heat, (2) by gently heating a mixture of 5 parts cuprous chloride and 3 parts sodium carbonate in a covered crucible, and separating the oxide by lixiviation; (3) by reduction of an alkaline solution of a copper salt by sugar or certain other organic bodies, as in Barreswil's (Fehling's) solution, (4) by heating in a covered crucible a mixture of 5 parts copper monoxide and 4 parts copper filings, (5) by the electrolysis of copper sulphate solutions under certain conditions (Miller, J. Phys. Chem. 1909, 256, Gillett, *ibid.* 1909, 332); (6) by heating cupric oxide in sulphur dioxide and treating the product with water, when cupric sulphate is dissolved and the  $3CuO + SO_2 = Cu_2O$ cuprous oxide remains +CuSO<sub>4</sub> (Hammick, Chem Soc. Trans 1917, 111, 384)

This oxide is decomposed by most acids into a cupric salt and copper; hydrochloric acid, however, converts it into cuprous chloride

Cuprous oxide fuses at a bright-red heat dissolves in metallic copper and renders it brittle or 'dry' when present in quantity much as 4 6 p.c. has been found by Abel in very dry copper, and, according to Watson (J. Soc. Chem Ind 1883, 154), from 1 to 2½ p c is present in ordinarily good copper

It is largely used in the manufacture of ruby glass, and for the production of a red glaze on pottery (Louth and Dutailly, Mon Ceram et Verr 19, 237), and, together with the black oxide, forms one of the so-called copper paints

used for painting ships' bottoms

A yellow form of the oxide is produced by the action of alkali on cuprous chloride, or better by the reduction of a cupric salt by means of hydroxylamine in presence of alkali. It can also be prepared electrolytically, using an alkali sulphate as the electrolyte and an anode of pure

copper. It may also be obtained by the reduction of Barreswil's (Felling's) solution in presence of a deficit of tartrate. The yellow precipitate thus formed is not a definite hydrate, but is a colloidal form of the oxide with an indefinite amount of water (Groger, Chem Zeit. Rep 1902, 197). It quickly changes to an orange or brick-red colour and can then be dried unchanged. The dried product contains from 2-3 pc of adsorbed water. At a low red heat it loses this water and changes into the red crystalline form. The yellow cuprous oxide is to be regarded as a primary metastable form, the red crystalline form being the stable one. It is a powerful reducing agent, especially in ammoniacal solution. This solution, which is colourless, immediately becomes blue from the formation of the monoxide on exposure to the atmosphere, and thus forms a delicate test for oxygen. The hydrated oxide dissolves in a concentrated solution of magnesium chloride

This oxide is produced in Hoepfner's process for extracting copper from ores, &c , with cupric chloride solution. Cuprous chloride is formed and treated with alkali to obtain cuprous oxide (J Soc Chem Ind. 1893, 932, Eng Pat 18900,

1892).

Copper monoxide, cupric oxide, black oxide of copper, 'copper oxide' ('uO, occurs as metaconite' or black copper, especially at Lake

It may be prepared by heating the nitrate or carbonate to dull redness or the sulphate to intense redness. The decomposition of the sulphate is the basis of Adcock's process for the preparation of the oxide direct from the natural carbonate or sulphate (Eng. Pat. 11678, 1900) The copper oxide used for organic analysis, &c., may be prepared by moistening copper scale (the mixture of suboxide and black oxide obtained while working copper sheet, &c) with nitric acid, and igniting

An electrolytic method of preparing the oxide has been described by Miller (J physikal. Chem. 1909, 256), and by Luckow (Zeitsch.

Elektrochem 1897, 482).

Copper oxide is a black powder which agglomerates when heated, and fuses at the melting-point of copper, forming cuprous oxide, or Cu<sub>3</sub>O, according to the temperature. It can be made to crystallise in regular tetrahedra is slightly hygroscopic, especially when in powder When heated with organic substances or certain gases, it is reduced to metal. The metal produced by reduction in hydrogen always contains that gas, but if the oxide be reduced in formic acid vapour, the metal is quite free from hydrogen and is suitable for organic analysis (Weyl, Ber. 15, 1138).

It dissolves in acids with formation of cupric salts. In oils, &c, it is soluble to some extent, it is for this reason that copper cooking

vessels require such constant cleansing.

Copper oxide is used in ultimate organic analysis; to make green and blue glass and glazes, and to some extent as a pigment. It has also been used as one electrode of a galvanic cell (v Lelande and Chaperon, Bull Soc chim. [2] 49, 173).

Cupric hydroxide CuO H2O may be prepared by precipitating a solution of a copper salt by addition of alkali and washing rapidly.

When heated with water, it becomes black and amorphous.

A colloidal form of the oxide has been obtained by Paal and Leuze (Ber 1906, 1545), which gives blue-violet solutions, and when dry retains its solubility indefinitely.

The hydroxide is used as a blue pigment by paper stainers, but becomes green on exposure. It corresponds closely with certain of the highly basic carbonates sold as verditers According to Dumas, it is best prepared as follows: 6 parts of copper sulphate are dissolved in water and mixed with a solution of 3 parts of The clear liquid is decanted calcium chloride from the precipitated calcium sulphate and is mixed with 1½ parts of lime made into a cream with water. The greenish precipitate is collected, washed, and mixed with one-fourth its weight of slaked lime and as much pearl ash, and to render the colour more permanent, onefourth its weight of ammonium chloride and one-half its weight of copper sulphate are usually added.

The hydrated oxide dissolves in ammonia, the saturated solution in strong ammonia being known as 'Schweitzer's reagent,' or curpanimonium. This solution has the property of gradually dissolving cellulose with formation of a viscid solution which is used for the preparation of the 'Willesden papers,' and for the production of 'artificial silk' on the large scale. This process consists in injecting the coppercellulose solution into strong solutions of alkah or acid. In either case, coagulation takes place with formation of a fibre of cellulose corresponding in thickness to the diameter of the jet

An aqueous solution of sodium hydroxide of sp gr 1 345-1 370, or of potassium hydroxide of sp gr 1 453-1 498, will dissolve 0 78 grm of cupric hydroxide in 100 c c giving a bright blue solution which yields no precipitate when boiled If a more dilute alkali solution be used the solution is not stable when further diluted and subsequently boiled (Justen-Mueller, Compt rend 1918, 167, 779)

Cuprammonium solutions are prepared on the large scale by the joint action of oxygen and ammonia on metallic copper at temperatures near 0° (Schaefer, U.S. Pat. 884298, 1908, Bronnert, Frémery, and Urban, Eng. Pat. 1763, 1900. Leccour, Fr. Pat. 374277, 1906).

Bellot has patented a continuous process of manufacture, using a series of closed vessels (Fr Pat 335207, 1903).

The solution should be purified by dialysis, as the presence of crystalline salt favours its decomposition (Leccui, Fr. Pat. 362986, 1906). Addition of 1–2 p.c. of polyhydric alcohols, carbohydrates, gum, &c., to cuprammonium solutions increases their stability and prevents the deposition of cupile hydroxide (Friedrich, Eng. Pat. 4104, 1909), and the presence of such substances facilitates manufacture of the solution, making it possible to employ copper concentrations up to 5 p.c. without cooling (Fr. Pat. 399911, 1909).

Chattaway has shown (Proc Roy Soc. 1908, 80, A, 88) that by the reduction of a cuprammonium solution with aromatic hydrazines, copper mirrors can be deposited on glass

Dawson states that a solution of cupric hydroxide in ammonia contains copper as the Vol. II.—T.

base Cu(NH<sub>3</sub>)<sub>4</sub> (OH)<sub>2</sub> (Zeitsch physikal Chem 69, 110) (For a general discussion of the constitution of cuprammonium salts, see Horn and Taylor (Amer. Chem. J. 1904, 253), and Horn (tbil 1907, 475))

Copper dioxide, copper peroxide CuO,, is obtained by the action of hydrogen peroxide on cupire hydroxide in neutral solution at 0° (Kruss, Ber 1884, 17, 2593, Moser, Zeitsch anorg Chem. 1907, 54, 121) The crystalline precipitate is washed with water, alcohol, and ether at 0°, and dried in vacuo 1t has a yellow-ish-brown colour, and decomposes at 180°, forming cupire oxide

ish-brown coron, conforming cupric oxide

Salts Two series of copper salts are known, cuprous and cupric. The former are mostly insoluble in water, and are of but little commercial importance. They act as powerful reducing agents and readily pass into the cupric condition. Cupric salts are mostly soluble, and are of wide application.

Cuprous chloride Cu<sub>2</sub>Cl<sub>2</sub> may be prepared, amongst other methods, by boiling cupric chloride with hydrochloric acid and copper turnings. On addition of water it is precipitated in white crystals

Zuloaga has suggested (Rev Chim Ind. 1900, 107) that the cuprous chloride solution used in the extraction of silver from its ores should be prepared by mixing solutions of cupric sulphite and of cupric sulphate and sodium chloride

Cuprous chloude melts below a red heat into a yellow transparent mass—The dry crystals become yellowish on exposure to light, if moist, they acquire a dirty-violet tinge—Plates of copper coated with this chloude by the action of chlorine will take photographic impressions, but cannot be made to withstand exposure to light (v Carlemann, J pr Chem 63, 475; and Priwoznik, Dingl-poly J 221, 38)

Cuprous chloride combines with ammonia to form three compounds

Cu<sub>2</sub>Cl<sub>2</sub> NH<sub>3</sub>, Cu<sub>2</sub>Cl<sub>2</sub> 3NH<sub>3</sub>, Cu<sub>2</sub>Cl<sub>2</sub> 6NH<sub>3</sub> Cuprous bromide and rodide form compounds analogous to the last two (Lloyd, J. Phys. Chem.

The hydrochloric acid solution, when exposed to air, absorbs oxygen and acquires a brown colour, subsequently depositing a pale bluishgreen insoluble copper oxychloride

An identical substance used as a pigment, and known as Brunswick green, is piepaied by boiling a solution of copper sulphate with a small quantity of bleaching powder solution; or by moistening copper turnings with free exposure to air. It occurs in nature as alacamite

The use of the oxychloride in preference to the sulphate for destroying mould on vines, &c., is recommended 200-250 grams per litre of water are employed (Deletrez, J. Soc. Chem. Ind. 1909, 438).

When a solution of cuprous chloride in potassium chloride, acidified with hydrochloric acid, is treated with acetylene, a yellow precipitate of Cu<sub>2</sub>Cl<sub>2</sub> C<sub>2</sub>H<sub>2</sub> is formed Other compounds containing copper oxide or potassium chloride can be obtained by varying the conditions (Chavastelon, Compt. rend. 132, 1489).

Acetylene reacts with solutions of cuprous

2в

chloride in ammonia to form a blood-red precipitate of cuprous accipiede, which, on dehydration, has the formula Cu<sub>2</sub>C<sub>2</sub>, and is highly explosive (Scheiber, Ber. 1908, 3816). Its decomposition by acids has been used as a means of preparing pure acetylene A colloidal solution of copper acetylide has been prepared by Kuspert (Zeitsch anorg. Chem 34, 453) (v. ACETYLENE)

Cuprous bromide Cu2Br, is obtained by the union of its elements or by boiling a solution of cupric bromide with copper, and forms a

brown crystalline mass

Cuprous iodide is the only known iodide of copper, and is obtained as a white precipitate by adding potassium iodide to a solution of copper sulphate containing ferrous sulphate or sulphurous acid It has a sp gr 5 653 at 15° (Spring, Rec trav chim 1901, 79), melts without decomposition at 628°, and is practically insoluble in water.

Cuprous sulphite Cu<sub>2</sub>SO<sub>3</sub>,H<sub>2</sub>O is obtained in microscopic nacreous plates by passing sulphur dioxide into a hot solution of cuprous accetate in acetic acid It forms double salts with the alkalı sulphites Cupro-cupric sulphite Cu<sub>2</sub>SO<sub>3</sub> Cu<sub>2</sub>SO<sub>3</sub>,2H<sub>2</sub>O is a red microcrystalline powder.

Cuprous sulphate Cu<sub>2</sub>SO<sub>4</sub> can be prepared by the action of dry dimethyl sulphate on dry powdered cuprous oxide (Recoura, Compt rend 148, 1909, 1105) A compound of cuprous sulphate and carbon monoxide Cu<sub>2</sub>SO<sub>4</sub> 2CO, H<sub>2</sub>O has been obtained by Joannis (ibid 1903, 615), by reducing cuprammonium sulphate hydroxylamine. Péchard has made ammoniocuprous sulphate Cu<sub>2</sub>SO<sub>4</sub> 4NH<sub>3</sub> (ibid 1903, 504) Cuprous sulphide Cu<sub>2</sub>S occurs as chalcone or

copper glance It may be prepared by heating copper to redness in sulphur vapour, or by heating a mixture of copper and sulphur—It melts at  $1130^{\circ}$   $D_{4^{\circ}}^{25^{\circ}}$  5 785—It is dimorphous, and has an inversion temperature at 91°

Cupric chloride CuCl2 may be prepared as a fused, liver-coloured, anhydrous mass by heating copper in excess of chlorine In solution it may be prepared by dissolving the oxide in hydrochlone acid - It crystallises in grass-green prisms or needles containing 2 molecules of They become pale-blue when dried in The anhydrous salt is obtained on gradual addition of a large excess of cone sulphuric acid to a solution of the chloride (Viard, Compt. 1end 1902, 135, 168)

Cupric chloride is very deliquescent, and is soluble in alcohol When a small quantity of potash is added to the aqueous solution, a paleblue basic oxychloride 2CuO CuCl<sub>2,4</sub>H<sub>2</sub>O is precipitated It turns to a black anhydrous powder when heated, but again becomes green with absorption of 3 molecules of water when moistened. It is an intermediate product in the manufacture of verditer (v. Copper carbonates)

Cupric chloride is used in calico printing, in the manufacture of methyl violet, and for the oxidation of Cutch colours

Green has shown that the chloride is a very effective disinfectant (Zeitsch. für Hygiene),

1893, 495).

When cupric chloride solution is mixed with excess of potassium thiocyanate solution, cupric throcyanate Cu(CNS)2 is produced. This is slowly acted on by water and is ultimately reduced to cuprous thiocyanate CuCNS with formation of carbon dioxide, ammonia, carbamide, hydrogen cyanide, throcyanic and sulphune acids (Philip and Bramley, Chem Soc. Trans 1916, 109, 597

Cupric bromide CuBr, is obtained by dissolving cupric oxide in hydrobromic acid and evaporating in vacuô It is dark-coloured, very deliquescent, and, when heated, decomposes into cuprous bromide and bromine It is used as an intensifier in photography, and for that purpose is made by dissolving in water a mixture of anhydrous copper sulphate and potassium bromide (Akt Ges f. Andinfabr D R. P. 201168, 1907)

Cupric sulphate, Blue vitrol CuSO<sub>4</sub>,5H<sub>2</sub>O, occurs naturally at Copaquire, Chile, the rock containing over 12 pc of the hydrated salt (Walker, Eng and Min J 1903, 710) This salt may be prepared by dissolving copper or

its oxide in sulphuric acid

It is prepared on the large scale from metallic copper in the following way. Old copper plates from the sheathing of ships, &c, are heated to strong redness in a reverberatory furnace, and an excess of sulphur is thrown in and the doors are closed When combination is complete, the doors are reopened and the heat is raised to dull redness to oxidise the sulphide into sulphate mass, while hot, is thrown into dilute sulphuric acid, allowed to settle, the solution decanted, concentrated, and crystallised The sulphate so produced is of considerable purity copper and copper glance or other sulphurous copper ores are similarly treated Pauffin and Couvert have patented a process for the direct manufacture of the sulphate from the metal (Fr Pat. 399413, 1909) U.S. Pat. 788862, 1905) (See also Walker,

When much iron is present, as in most copper ores and especially in copper pyrites, it would be impossible to separate the copper and ferrous sulphates by crystallisation, because, although copper sulphate crystallises normally with 5 molecules of water, it forms, in presence of ferrous sulphate, crystals isomorphous with that salt and containing 7 molecules of water By carefully adjusting the temperature at which the sulphide is roasted, however, the iron salt may be converted into the oxide while the copper sulphate remains almost unaffected. The iron may be removed by heating the solution of the sulphates to 180° in a copper boiler under pressure Ferrous sulphate crystallises out and is filtered off (Gin, Fr. Pat 328800, 1903) Tho iron may be oxidised and thus separated from the copper by boiling the solution with a little nitic acid. It may also be precipitated as oxide by boiling with copper oxide, or by boiling with lead peroxide with subsequent addition of a little banum carbonate.

Crude copper or the ore may be roasted in air, and then treated with sulphur dioxide to form the sulphate (Gin, Zeitsch. angew Chem.

1903, 566; Eng Pat 5230, 1903)
Darier has patented a process in which cupric chloride is formed by the action of chlorine and water on the metal, and is then decomposed by sulphuric acid, with formation of the sulphate and regeneration of hydrochloric acid (Fr. Pat. 350421, 1904).

When the sulphate is required for purposes

where the presence of iron is not injurious, such liquors, or the mother liquors from which the purer salt has been separated, may be at once crystallised out, with the formation of crystals containing both iron and copper. Much of the 'agricultural copper sulphate' is of this class. The 'Sulzburg vitriol,' prepaied at Buxweiler, also contains both iron and copper. According to Lefort (Compt rend. 26, 185), it consists of CuSO<sub>4</sub> 3FeSO<sub>4</sub>,28H<sub>2</sub>O 'Cupram vitriol,' prepaied at Chessy from zinciferous copper ores, forms blue rhombic prisms, which contain according to Lefort, CuSO<sub>4</sub> 3ZnSO<sub>4</sub>,28H<sub>2</sub>O.

The copper may be separated from the ferrous mother liquors by placing them in vats containing iron plates upon which the copper is

slowly deposited.

Apparatus for the solution of copper oxide in sulphuric acid upon a large scale has been described by Coste (Fr Pat 392617, 1908) If copper be known to contain silver or gold, as is the case with that from the Harz, it is treated with sulphuric acid diluted with its own volume of water, the copper being thus dissolved while the silver and gold are unattacked

Argentiferous copper ones are roasted in a reverberatory furnace, and added in small charges to sulphuric and and digested until the solution contains but little free acid. The solution is then decanted from the precipitated lead and gold, and is concentrated and run into leadlined tanks containing plates of copper, upon which all the silver and part of the antimony and arsenic are deposited, while the greater proportion of the bismuth is precipitated as a basic sulphate, and the iron is reduced to the ferrous condition The liquor is then crystallised, the mother liquors being used for treating a fresh quantity of ore Copper sulphate is obtained of great purity and in considerable quantity in the refining of silver by precipitating it upon plates of copper from its solution as sulphate (v Silver)

An electrolytic process for the production of the sulphate consists in using a solution of sodium sulphate and copper electrodes, a current of carbon dioxide being passed through the liquid. Copper is dissolved from the anode as sulphate, while sodium carbonate is produced at the cathode. These react, regenerating sodium sulphate and precipitating copper carbonate, which is collected and dissolved in sulphuric acid (Kroupa, J. Soc. Chem. 1nd. 1906, 78)

Palas and Cotta have also described electrolytic processes for the production of the salt (Eng. Pat. 9806, 1899, and 17485, 1900)

(Eng Pat 9806, 1899, and 17485, 1900)
The electrolytic production of the sulphate from 'coment copper waters' has been described by Rambaldini (Chem. Zentr. 1909, 1–1675).

It is stated that copper sulphate is sometimes adulterated with ferrous sulphate and Prussian blue to the extent of 90 pc (J. Soc Chem. Ind. 1900. 84)

Copper sulphate crystallises in large transparent blue, doubly oblique rhombic prisms of sp gr 2 28, of the formula CuSO<sub>4</sub>,5H<sub>2</sub>O. According to Poggiale, their solubility in 100 parts of water is:

10°C.  $20^{\circ}$ 30° 40° 50° 36 59 42 31 48 81 56 9 65 83 80° 90° 70° 100° 203.32 156.44 94 60 118.0

When heated, it loses 2 molecules of water at 30°, 2 more at 93°–98°, and at 232° becomes anhydrous, white, and hygroscopic, and combines with water with considerable evolution of heat. At 341° the anhydrous salt begins to decompose (Wanjukow, Chem. Zentr. 1909, if 1124). The sp.gr. of the anhydrous salt is 3606 (Thorpe and Watts). It combines with water in alcohol and other organic liquids with production of a blue colour, and may be used for ascertaining the presence of water in those substances, but the test is not delicate. The anhydrous salt is also soluble in anhydrous methyl alcohol, and produces a blush-green solution (Klepl, J. pr. Chem. 1882, 25, 520)

Copper sulphate absorbs hydrogen chloride with evolution of heat, forming cupit chloride and liberating sulphuric acid—It is therefore used to remove hydrogen chloride from such gases as chlorine or carbon monoxide—A similar

reaction occurs in solution.

When mixed in solution with molecular proportions of potassium and other sulphates, it produces well-defined, crystalline double sulphates. Several basic sulphates are known

The sulphate is the most important of the copper salts. It is largely used in calico printing, principally in connection with potassium dichromate or iron mordants, in black dyeing with logwood, directly or indirectly for the preparation of most pigments containing copper, and to prevent rot in timber. In agriculture the crude sulphate containing ferrous sulphate is used to a considerable extent to prevent smut in seeds. The seeds are soaked for some hours in a weak solution of the sulphate, and are sown within 24 hours.

A 10-20 pc solution of copper sulphate, preferably mixed with lime, is applied to vines with great benefit. A small quantity is found in the vines and must, but the tannin and sulphur introduced before fermentation appear to remove it in most cases.

Kaisten has stated (v. J. Soc. Chem. Ind. 1896, 367) that attacks of illness resembling dysentery have been caused by a wine which contained copper in such quantity as to give a visible deposit on steel in 12 hours. Its presence was due to spraying the vines with a copper salt.

An effective wash for vines is prepared by boiling 25 parts of sodium carbonate and 25 parts of resin in 100 parts of water, and then diluting and adding copper sulphate (J. Soc. Chem. Ind. 1899, 155)

Copper sulphate is used in preparing copper ferror yande, a solution of which, in potassium citrate solution, is used as a toning-bath for photographs (Ferguson's Copper Toning, ibid. 1900, 465),

Copper sulphate is also used in dyeing to make diazo-colours permanent to light. The dyed fabric may be treated with the salt (Graebling, ibid. 1905, 797), or it may be added to the diazotising bath (Grossmann, Farber-Zeit 9, 328)

Cupro-tetrammonium thiosulphate  ${\rm CuS_2O_3}$   ${\rm (NH_3)_4}$ , has been proposed as an ingredient of detonator compositions, which although less sensitive are alleged to possess advantages over mercury fulminate compositions, as being cheaper, less unhealthy, simpler, and less dangerous to manufacture. The thiosulphate is

made by adding excess of strong ammonia solution to a hot saturated solution of copper sulphate, and thereafter adding an equivalent amount of sodium thiosulphate, dissolved in The salt separates out as the soluhot water tion cools; it is washed with a weak solution of ammonia, and dried at a temperature not exceeding 50° The violet-blue crystals are slightly soluble in water, not hygroscopic, insensitive to light, and very stable positions suitable for various purposes are made by mixing with potassium chlorate, powdered glass, antimony trisulphide and lead thiosulphate (Herz, Zeitsch ges Schiess u Spieng-stoffw 1912, 7, 284; J. Soc Chem Ind 1912, 31, 903)

Potassium copper nitrite K<sub>3</sub>Cu(NO<sub>2</sub>)<sub>5</sub> 18 obtained by mixing cold saturated solutions of copper sulphate and potassium nitrite, adding methyl alcohol, filtering, and adding ethyl acetate to the clear, green filtrate when long crystals of the salt separate Rubidium copper nitrite of similar composition may be obtained In like manner (Kurtenacker, Zeitsch anorg Chem 1913, 82, 204). Basic copper nutrites  $3\mathrm{Cu}(\mathrm{OH})_2\,\mathrm{Cu}(\mathrm{NO}_2)_2$  and  $2\mathrm{Cu}(\mathrm{OH})_2\,\mathrm{Cu}(\mathrm{NO}_2)_2$ 

are also known

Cupric natrate Cu(NO<sub>3</sub>)<sub>2</sub> may be prepared by dissolving copper or the oxide or carbonate in nitric acid. The solution at temperatures above 20° deposits dark-blue crystals containing 3 molecules of water, which melt at 1145° and boil at 170° with evolution of nitric acid and formation of a green basic salt. This basic salt formation of a green basic salt may also be prepared by boiling the solution of the nitrate with copper or copper hydrate, or with a little alkalı (v. also Copper carbonates for an application of the basic mitrate)

The anhydrous salt may be obtained by the action of nitric anhydride on the hydrated salt It is a white, deliquescent powder, decomposing at 155°-160° (Guntz and Martin, Bull Soc chim

1909, 5, 1004)

Copper nitrate is used to a slight extent in dyeing and calico printing, and a solution, prepared by mixing solutions of copper sulphate and calcium nitrate, is used in viticulture (Gouthière et Cie and P Ducancel, Fr Pat. 395806, 1908)

Cupric sulphide CuS occurs as covelite or ındıgo copper in blue flexible plates, in Vesuvian lava, in Chile, and in the copper-mines of Mansfeld It may be prepared by heating cuprous sulphide with sulphur at a temperature not above the boiling-point of sulphur, by digesting cuprous sulphide with cold strong nitric acid, or by precipitating a copper solution with sulphu-

retted hydrogen  $D_4^{25}$  4 68. On the large scale it is obtained as a paste by precipitating a solution of the sulphate with sodium sulphide. This paste is used in calico printing for the development of anılıne black

The preparation and use of copper sulphide for thermo-couples has been described (Eng Pat 11300, 1905).

Rieder has used (Zeitsch. Elektrochem 1902, 370) an electrolytic method for the pre-

paration of this compound

A brownish-red persulphide of copper has been obtained by Bodroux by adding a solution of calcium pentasulphide to a solution of copper acetate at 0° (Compt. rend. 130, 1397).

Copper acetate v. Metallic acetates, ACETIC ACID.

Copper arsenite v. Scheele's green. ait. ARSENIC.

Cupric phosphate is of no technical interest. A bulliant green colour containing a basic phosphate, prepared by heating a mixture of copper acetate, sodium phosphate, and ammonia, has, however, been used in calico printing (Bourcart, Mon Sci [3] 4).

Phosphorus combines in several proportions with copper, and produces true alloys, which in

many cases are of great value.

Heyn and Bauer state that the phosphorus exists in the alloy as Cu<sub>3</sub>P and Cu<sub>5</sub>P<sub>2</sub> (J Soc. Chem. Ind. 1906, 1047) The latter compound has been obtained by Rubénovitch by the action of phosphine on copper sulphate (Compt rend 127, 270)

Copper silicide (Cuprosilicon) CuaSi is obtained when copper and silicon are heated together in the electric furnace (Vigouroux, Compt 1end 1896, 122, 318, Lebeau, sbid. 1906, 142, 154) It is now manufactured in Germany and America by this process (Steinhart, Eng and Min J 1899, 67, 710), and is used for purifying and hardening copper, and for adding to tin- zinc-, and aluminiumbronzes to increase their tensile strength and hardness

Cuprosilicon can be prepared pure by heating together 17 parts silicon and 9 parts copper in hydrogen, and removing the excess of silicon by solution of sodium carbonate. It is silverwhite, with a metallic lustre, and is haid and The sp gr is 758, and it is attacked by chlorine and aqua regia (Vigouroux, Compt. 1end 1906, 142, 87, see also Philips, Métallurgie, 1907, 4, 587, and Lebeau, ibid 141, 828)

Copper carbonates. The normal salt has not been prepared, but several basic carbonates are known, and used as blue and green pigments Basic copper carbonate occurs native as malachite Cu(HO)<sub>2</sub> CuCO<sub>3</sub>, and as azurite or chessylite Cu(HO)<sub>3</sub> 2CuCO<sub>3</sub> When ground, these minerals

form a fine but expensive pigment

An anhydrous basic carbonate is precipitated on addition of a solution of sodium carbonate to a hot solution of copper sulphate or nitrate. (For a complete description of all the known basic carbonates of copper, see Pickering, Chem Soc Proc 1909, 188 Cf Dunnieliff and Lal, Chem Soc. Trans. 1918, 113, 718) Commercial basic carbonates of copper differ considerably in composition: the amount of carbon dioxide present is below that required for malachite or azurite. A definite basic salt is known of the composition 2CuCO<sub>3</sub>,5Cu(OH)<sub>2</sub>.

Under the names Verditer green, Verliter blue, Bremen green, and Bremen blue, this salt is largely prepared for paper staining, &c According to Habich (Technologiste, 17, 413), the following processes are in use. A hundred parts of old copper sheathing from which impurities, oxide, &c, have been removed by digestion in dilute sulphuric acid, is cut into pieces and mixed with a paste formed by moistening a mixture of 99 parts potassium sulphate and 100 parts common salt, and the mixture is stirred occasionally for some months to ensure thorough exposure to the atmosphere The action appears to consist in the production of cupric chloride,

which, in presence of the metal, becomes converted into a basic chloride, and finally, by the action of the oxygen, into an oxychloride. the oxidation be incomplete, the subchloride is not completely oxidised, and, as it, when subsequently decomposed by alkali, gives an orangered precipitate of suboxide, the pigment would be lowered in value. The oxychloride may also be similarly prepared from a mixture of 100 parts copper, 60 parts common salt, and 30 parts strong sulphuric acid diluted with three times its volume of water; or by treatment, with free atmospheric exposure, of copper with a solution of cupric chloride in hydrochloric acid.

To 100 kilos. of this thick magma of oxy-

chloride, which has been washed by decantation, a concentrated solution containing 7 kilos copper sulphate and 40 kilos concentrated soda solution (32°-36°Bé) is added. The mixture is vigorously stirred and poured at once into 150 kilos of soda solution (20°Bé.). The precipitate is washed, passed through a fine hair sieve, and dried spontaneously or at a temperature below 25°. If hot air be used in the drying, it must be quite free from acid or sulphin gases produce the finest colour, desiccation must be perfect

Habich recommends the following process for making a verditer which has a very different composition from the above A solution of copper oxide in nitric acid is treated with potassum carbonate, which is added in portions until most, but not all, the copper has been preci-The green precipitate is washed and introduced into a solution of copper nitrate whereby it is converted into a heavy green insoluble basic nitrate. This is digested in a solution of potassium zineate (formed by dissolving zinc in caustic potash solution) A dark-blue pigment, of great 'body' but little weight, is thus produced It contains zinc and copper, apparently combined as a double oxide or as 'copper zincate,' and a small quantity of basic

copper nitrate

Verditer may be prepared by agitating a solution of copper nitrate with chalk. The washed pasty precipitate is mixed with 8-10 p c. of freshly burnt lime with production of a fine The precipitate without velvety blue pigment

addition of lime is green

According to Gentele, the following process adonted at Bremen, Cassel, &c; 225 lbs is adopted at Bremen, Cassel, &c; 225 lbs sea salt and 222 lbs blue vitriol, both free from iron, are mixed dry and ground between millstones with water to a thick paste paste is mixed in strata with 225 lbs of clean copperplate cut into pieces about 1 inch square, in wooden chests made without nails, and the mixture is digested for about 3 months with the maximum exposure to air, the whole being well mixed and exposed about once a When the action is complete, the undissolved metal is removed and the precipitate is washed with a minimum of water, filtered and thrown into a tub To 180 lbs of the magma 12 lbs of hydrochloric acid (13°Bé) is added, and the mixture is stirred and left for 24 or 36 hours. Six volumes of this solution is mixed with 15 volumes of caustic alkali (19°Bé) followed after an interval by 6 volumes of water. The mass is well mixed and left for 36-48 hours, and is then well washed by decantation, filtered,

exposed for some time in a moist condition to the air, and finally dried (v. also Copper monoxide'

COPPER-GLANCE or Chalcocite, Native cuprous sulphide Cu<sub>2</sub>S, crystallising in the orthorhombic system. Excellent twinned crystals were formerly found in abundance in the neighbourhood of Redruth in Cornwall (hence the name redruthite) and Bristol in Connecticut, usually, however, the mineral is found as compact masses. It is iron-black with metallic fustre, but, on exposure to light, it soon becomes dull with a powdery black coating It is soft (H 2½) and sectile, and can be readily cut with a knife; sp gr 5.7 It occurs in veins of copper ore, and is often of importance as an ore, especially when it contains small amounts of Isomorphous with copper-glance is the mineral stromeyerite (Ag, Cu)2S, which has been found in considerable quantities in the Altai Mountains in Siberia

COPPER PAINT v COPPER

COPPER PYRITES or Chalcopyrite. A sulphide of copper and iron CuFeS<sub>2</sub>, containing when pure 34.5 p.c. of copper, and the most important of the ores of copper It has a characteristic brassy-yellow colour with metallic lustre, and on the surfaces often displays a brilliant iridescent tarnish (peacock-copper-ore). It often occurs together with, and intimately intermixed with, iron pyrites, from which it can be distinguished by its deeper colour and lower degree of hardness, copper pyrites (H  $3\frac{1}{2}$ -4) is readily scratched with a knife, giving a greenishblack powder, whilst iron pyrites (H. 6-6½) is scratched only with difficulty or not at all It occurs in mineral-veins and in bedded deposits, and by its alteration gives rise to many secondary copper minerals Crystals are tetragonal with inclined hemihedrism, but only rarely are they distinctly developed, sp gr 4 2 L. J. S.

COPRA and COPRA OIL v. COCO-NUT

COPROLITES. Concretionary mixtures of calcium phosphate and calcium carbonate, derived from the excrements of certain extinct fishes and reptiles Formerly much used in the manufacture of superphosphate of lime (v FERTILIZERS)

COPROSMA. The bark, and more especially the roots, of the C grandiflora, C linariifolia, and C areolata, which belong to the natural family of the Rubiaca, and are widely distributed in New Zealand, possess tinctorial property (Aston, New Zealand, J Sci Tech 1918, 1, 3), and being related to madder probably contain either alizarin itself or an allied colouring matter. Thus an alcoholic extract of the C grandiflora is coloured purple by alkali, and becomes yellow on addition of acid. From the acid solution, by means of ether, an orangeyellow crystalline substance can be isolated in small amount

COPYING or INDELIBLE INK PENCILS. Unlike iron-gall copying-inks, which will only yield good copies for a relatively short time after writing, solutions of aniline dyestuffs produce writing from which copies may be taken at any subsequent period. In copyingink pencils the aniline dyestuff, usually methyl violet, is incorporated with graphite, kaolin clay or other basic material. The preparation of pigments for this purpose was claimed by Petit (Eng. Pat. 4090 of 1874), and their composition was first described by Viedt (Dingl. poly J 1875, 216, 96) At the present time these pencils are seldom employed for copying purposes, and then chief use is as a substitute for ink, since they produce a writing which cannot be erased so readily as the marks of a lead pencil Hence they are frequently described as indelible pencils, although the description is only relatively correct Lehner (Ink Manufacture, 1902, p 125) gives the composition of four kinds of these pencils made by Faber in Bayaria aniline dyestuff ranged from 255 to 50 pe, graphite from 24 5 to 37 5 pc, and kaolin clay from 12 5 to 50 p c

Specimens of these pencils examined by Mitchell (Analyst, 1917, 42, 1) had the following composition .-

Origin	Mois- ture	Residue insol, in water or alcohol	D <b>y</b> est uff	Loss on ignition (gra- phite, &c )	Ash (kaolin, &c)
	p e	рe	рe	p e	D C
American	$2\ 35$	76 45	1 21 20	27.09	47 36
,,	7 48	51 53	40 99	25.48	31 92
,,	2 30	60 89	36 81	56 84	4 05
,,	3 75	63 95	32 30	13 90	50 05
Austrian	3 56	47 14	49 94	44 14	3 00
British	2.72	65 48	31.80	62 08	3 40
,,	4.53	73 05	22 42	69 17	3 88
German	3 72	64 18	31 80	11.86	52 86
,,	6.33	59.83	33 84	7 61	52 22
,,	3 83	57 59	38 58	40 01	17 58
	I				

Pigments used in copying-ink pencils may be classified into four main groups-viz (1) those composed of graphite and dyestuff only, (2) those in which the dyestuff has been incorporated with approximately equal proportions of kaolin and graphite, (3) those in which the kaolin is largely in excess; and (4) those in which the graphite predominates The dyestuff is usually methyl violet, but copying-ink pencils in which red or blue aniline dyes are used are also sold

The copying properties depend not only on the proportion of the dyestuff, but also on the influence of the basic material. Pencils in which a large proportion of alumina is present yield only faint copies, even when the pigment contains a large amount of the dyestuff Although methyl violet is usually regarded as a very fugitive dyestuff, experiments have shown that under certain conditions the writing done with these pencils is more permanent than writing done with ordinary ink

C A M CORAL is the calcareous substance secreted by many varieties of five kinds of organisms, viz. (1) Actinozoa, the principal constituents of coral reefs, (2) Hydrozoa, the large common corals known as millepores; (3) Bryozoa, delicate corals in various forms, resembling mosses, branches, thin filamentary deposits, or

lowly organisms, mostly of very minute size; (5) Nullipores, which, although closely resembling millepores, are veritable algae or sea-weeds; they may be readily distinguished from the millepores by the absence of perforations

Coral is simply a secretion corresponding to the bone of vertebrates or the shell of mollusca, and is composed almost entirely of calcium carbonate derived from sea-water and from the food of the coral-making polyps

Ordinary coral is somewhat harder than . limestone or marble, and, when tapped with the hammer, gives out a ringing sound clearer than is clicited from ordinary calcium carbonate. The sp gr of 15 specimens examined by Silliman, averaged 2 523

### Composition of Coral

S P Sharples (Amer J Sci 31, 168) found

	Calclum carbonate	Calcium phos- phate	Water and organic matter
Oculina arbuscula,			
N Carolina Manicina areolata,	95 37	0 84	3 79
Florida	96 54	0 50	2 96
Florida Siderasti wa radians.	97 73	0 53	1 64
Florida . Madrepora cervicor -	97 30	0 28	2 42
nis, Florida Madrepora palmata,	98 07	0 32	1 93
Florida	97 19	0 78	2 81

Small and variable quantities of magnesium carbonate, calcium sulphate, calcium fluoride, alkaline chloride, silica, and oxide of iron are found in corals

The red coral, Corallium subrum, varies in colour from crimson to rose, the latter being the rarest and, therefore, the most prized It is gathered from the rocky bottom and shores of the Mediterianean, and it occurs most abundantly at a depth of 25-50 feet, although, according to Dana, it may be found as deep as 1000 feet Forchhammer found 2.1 pc magnesia in Corallium rubrum, and 6 36 in Isis According to Trommsdorff, the red pigment of coral is a resin soluble in oil of turpentine, and, after removal from the coral by this solvent, is easily soluble in alcohol and ether, but insoluble in caustic potash. In a red madrepore, Vauquelin found a red colouring matter, which was turned violet by alkalis. Moseley has described in some corals and allied organisms a red colouring matter (Polyporphyrin) which is insoluble in water, glycerol, alcohol, ether, and in strong solutions of ammonia and potash It is soluble in moderately strong sulphuric, nitric, or hydrochloric acids. For the formation and distribution of coral and coralreefs, &c v Corals and Coral Islands, J. D Dana: more solid masses (in the Palæozoic limestone formations this species of coral abounds); (4) Edinb 1880, 505 See also Forchhammer, Foraminifera, chambered shells belonging to J Pharm. Chim 49, 52; J 1847, 1291, 1849,

CORK. 375

813; 1851, 865; 1852, 958-961; Sharples, *lc*; Moseley, Quart. Journ. Microscop. Sci. xvii. 1877; also Max Bauer, Edelsteinkunde, 2nd ed. 1909, Eng trans. Precious Stones, I.

Spencer, 1904.

It is now known that the materials constituting coral-reefs sometimes undergo remarkable chemical changes, and that the calcium carbonate is largely replaced by magnesium carbonate or calcium phosphate. In 1846 J D Dana and B. Silliman showed that the rock of an upraised coral-reef in the Pacific contained as much as 38 pc. of magnesium carbonate. The deep boring made in the Atoll of Funafuti in 1897-8, yielded a series of cores, of which a chemical examination was made by E W Skeats and Hart Smith in the laboratories of the Royal College of Science It was found that down to the depth of 637 feet dolomitization had only gone on to a small extent, the proportion of magnesium carbonate never exceeding 16 pc, but the lower portions of the core were very nich in the magnesium salt, containing from 30 to 43 pc, the rock thus becoming almost a true dolomite A large collection of coral limestones obtained by Alex Agassiz, from reefs in the Pacific, was subsequently examined by Skeats, who showed that they contained varying amounts of magnesium carbonate up to nearly 42 pc Specimens from Christmas Island in the Indian Ocean gave similar varying results, percentages of magnesium carbonate up to over 43 p.c. being found A coral limestone obtained by boring at Key West, Florida, yielded to G Stieger, in the laboratories of the U.S. Geological Survey, over 14 p c of magnesium carbonate

Coral reefs that have been covered by deposits of guano are found to have their carbonates largely replaced by phosphates Thus the rocks of the coral reef known as Christmas Island in the Indian Ocean have been largely worked for the deposits of calcium phosphate, and Teall has shown that Clipperton Atoll in the North Pacific contains rocks with as much as

38½ p.c. of phosphotic acid (For changes in the chemical composition of coral rock, see Dana and Silliman, Amer J Sci 2, 1 1846, 189; Judd, Chemical Examination of Materials from Funafuti in 'the Atoll of Funafuti,' Phil Trans 1904, 362-389, Skeats, The Chemical Composition of Limestones from Upraised Coral Islands, Bull Mus Comp Zool Harvard, vlu 1903, 53-126, Teall, Quart Journ Geol Soc liv 1898, 230-232)

CORALLIN, CORALLINPHTHALIN, PHE-NOLCORALLIN v AURIN

CORDIALS and LIQUEURS are spirituous compounds flavoured with essences obtained from fruits, herbs, spices, or other substances, usually sweetened with sugar or glycerol, and often strongly coloured They are manu factured and used to a much greater extent in France and other European countries than in the United Kingdom, and although there is no essential difference between them, the term 'cordials' is by many restricted to British, and 'liqueurs' to the foreign preparations.

The spirit in the best qualities of liqueur is derived from the grape, but grain and other spirit is frequently employed, especially in cordials; but, whatever the origin of the spirit, it should be highly rectified and neutral in character, so as not to impair the flavour of the finished product. The alcoholic strength of cordials and liqueurs varies greatly, but, on the average, they contain from 30 to 40 p.c. of alcohol, or 50 to 70 p c. of proof spirit. The sugar must be pure, and is added in the form of a

clear syrup.

The principal flavouring ingredients employed are (a) those used solely on account of their essential oils, as aniseed, bitter almonds, caraway, juniper berries, mint, lemon-peel, orange blossom, &c; (b) those valued for their bitter principles as well as their essential oils, as cinnamon, cloves, curaçoa apples (a kind of orange), gentian, ginger, orange-peel, sweet calamus, wormwood, vanilla, &c, and (c) fruits, chiefly cheries, pineapples, raspberries, and strawberries. The flavourings are obtained either by digesting the flavouring ingredients (bruised or cut up) with alcohol and distilling the mixture to obtain an essence, or by filtering the alcoholic extract to form a tincture, or by simply expressing and clearing the juice of fruits or digesting the latter for a more or less prolonged period in spirit

Artificial essences simulating the above flavourings are frequently employed, but their use in beverages is questionable, since it seems probable that their physiological action differs from that of the natural flavouring substances

which they are intended to imitate.

The colouring matters most commonly used are as follows "Red, amline red, brazilwood, cochineal, cudbear, and sandalwood, yellow, saffron and turmeric, blue, indigo, chlorophyll (from nettles, parsley, spinach, &c) and mixtures of blue and yellow colourings, violet, and brown, a solution of buint sugar or caramel (v Liqueurs)

CORDIERITE v IOLITE

CORDITE v Explosives.

CORDYLITE (or Barium-parisite). Carbonate and fluoride of cerium-metals and barium (CeF)<sub>2</sub>Ba(CO<sub>3</sub>)<sub>3</sub>, containing Ce<sub>2</sub>O<sub>3</sub>, 23 72, and (La,Di),O<sub>4</sub>, 25 67 p.c. It occurs as small, pale wax-yellow, club-shaped hexagonal crystals in syenite-pegmatite at Narsarsuk near Julianehaab in south Greenland, sp gr 431 It is isomorphous with, and closely iclated to, parisite.

L. J S

J. C

**COREINE** v Oxazine colouring matters. CORIANDER FRUIT, Cornandra fructus,
The dried ripe fruit of Cornandrum satirum (Linn)

CORIANDER, OIL OF, v OILS, ESSENTIAL. CORIARIA MYRTIFOLIA v SUMACH

**CORINDITE.** A refractory and abiasive substance obtained by heating a mixture of bauxite and anthracite in a cupola furnace Aluminium carbide is first formed and is ultimately transformed into a vesicular fused mass of crystalline alumina of high melting point, low thermal expansibility, and great hardness, suitable for the manufacture of fire bricks for rotary cement kilns, gas and heavy oil ovens, coke ovens, &c.

CORIOFLAVINE v ACRIDINE DYESTUFFS. CORIOPHOSPHINES v ACRIDINE DYESTUFFS. CORK. Cork is derived from the bark of two evergreen species of oak tree, the cork oak, Quercus suber (Linn) and Q occidentalis (Gay) The former yields the best cork, and grows mostly in those countries bordering on the Mediterranean up to the 45° lat The latter occurs on the Atlantic seaboard of Portugal, Spain, and

France up to 48° lat

The cork tree is from 30 to 60 feet high, and 3-4 feet thick. It grows slowly and reaches the age of 150-200 years. The tree has the peculiarity of throwing off its outer bark when it reaches a certain age, and then reproducing it. It is this that makes the cork industry possible. The chief countries in which cork is produced are:

1 Spain Particularly the province of Catalonia Here the best cork is grown

Seville is the chief market.

2 Portugal Centres Lisbon and Oporto

3 Algeria Here the cork trees form great forests, those belonging to the State were some years ago fully 600,000 acres in extent

- 4 Southern France and the Pyrenean district Large forests of Q suber exist in the more mountainous parts, but in the low-lying 'Landes' the tree is Q occidentalis, which grows as far north as Belle Isle
- 5 The Adratic coast. Some is produced in the Austrian provinces, but it is of inferior quality.

6 Sicily, Sardinia, and Corsica also yield cork of good quality.

For an account of the cork industry in Sardinia, see J Chem Soc Ind 1919, 38, 352 R. From U.S. Com. Rep. July 25, 1919

Speaking generally, the cork tree seems to thrive best on dry granitic and sandy soils, and not to do well in damp or limestone soils

The complete bark of the cork tree consists

of two layers

(1) The dry dead bark, comprising the epidermis and the cork proper, with the meso-

(2) The green living skin which clothes the wood immediately under the dry back. This may be looked upon as the active cork-producing layer, and contains bast fibres and the endodermis

The epidermis on one-year-old twigs is green and smooth, but in the second year between it and the green under-skin irregular white waxy masses are formed, these represent the beginning of the cork formation and increase from year to year, changing gradually into cork proper. At the end of the third year the epidermis dries up and clacks

Barking the trees. This is the most important operation in the management of a cork forest, because on the care with which it is carried out depends the life of the trees and the

regularity of the slabs produced.

In barking, the workman takes a special tool and makes a cut round the stem above and below the portion of bark to be removed. A vertical cut or cuts are made so that the bark may be stripped off the tree, in which operation the shaft of the tool is used as a lever. Care must be taken not to bruise the green underskin, the real producer of the cork

The first barking produces so-called 'virgin cork,' which is of little value, as it has not elasticity enough for use as bottle corks. It was formerly used for the production of Spanish

black, but of late years it has been employed in the manufacture of ground cork.

During the first year of growth the tree produces from 3 to 5 mm. of cork, but afterwards less, the mean for the first 9 years being about 2 1-3 mm., giving a total thickness of 20½ mm. After 9 or 10 years, when barking usually occurs, the growth is slower, but where thicker cork is required, the period is of course longer. The first stripping usually takes place on the 15-20 year-old tree. The best cork is yielded by trees of from 50 to 100 years old. Those growing in the mountainous districts yield a smaller quantity but better quality of cork.

After the cork bark has been removed from the tree, it is stacked and weighted with heavy stones to flatten it, and when dry it is scraped with an iron tool to remove the epidermis. The cork is then immersed for 5 or 6 minutes in a tank of boiling water, when it swells, and, according to the quality, retains this swelling more or less permanently. The water also removes tannin, vegetable acids, and their salts. In Spain the cork is occasionally passed through a fire, whereby it is superficially blackened, acquiring, it is supposed, by this treatment especially good qualities or 'nerve'

After steeping, the plates are placed in heaps, weighted, and by hydraulic presses compressed

into bales and bound with hoop iron

Chemistry of cork. ('ork extracted with hot water under pressure, yields gallic acid, colouring matters, iron and magnesium salts. By extraction with alcohol it yields 'cerin,' or cork wax  $C_{32}H_{20}O$ . Siewert examined the alcoholic extract and isolated from it 'decaciylic acid'  $C_{10}H_{18}O_2$ .

The composition of cork has been further studied by Kugler (Arch Pharm 1884, 217), Gilson (*ibid* 1890, 690), and Hy Ingle (J Soc Chem Ind 1904, 1197)

Suberine, according to Kugler, is a real fat, which cannot be entirely extracted by solvents, because they do not penetrate the material

sufficiently

Gilson saponified cork from Q suber with 3 p.c. alcoholic potash, and isolated from the filtrate 44 p.c. of fatty acids, consisting of Phellonic acid  $C_{2\nu}H_{3\nu}O_{2\nu}$ , mp. 95°-96°, a monobasic acid giving a violet colour with zine chloude-nodine solution; yield 8 p.c. Suberinic acid  $C_{1\gamma}H_{3\rho}O_{3\nu}$ , a semi-fluid sticky substance; 36 p.c. Phlononic acid  $C_{12}H_{4\rho}O_{3\nu}$  fine white needles, m.p. 120°-121°; in small amount

Gilson considers suberine not to be a true fat, but rather a mixture of complex esters and possibly condensation products of different acids.

Treated with iodine solutions cork does not give the cellulose reaction, and there is some dispute as to whether cellulose or a pentosan is the carbohydrate present in cork. Gilson, from microscopic examination, considers that cellulose is present.

Cork also contains certain aromatic compounds, tannin and the aromatic aldehyde of wood, hadromal, besides coniferin and vanillin.

Treated with strong nitric acid, cork produces suberic acid (CH<sub>2</sub>)<sub>8</sub>(COOH)<sub>2</sub>, oxalic acid, and other products. Strong sulphuric acid charsit, but it is little affected by hydrochloric acid. Chlorine, bromine, and iodine cause cork to shrink, the two former first bleaching it—Ignited

it burns, swells up, and forms a charcoal called Spanish black

Physical properties—The sp gr of cork is variously given from 0 12 (Rollmann) to 0.24 (Muschenbroek), on the average 0 16

The uses of cork depend on its compressibility, elasticity, imperviousness to liquids, its lightness, its stability to weathering, &c It is used in making stoppers, mats, soles, linings, springs for heavy waggons, as anti-vibratory and sound-deadening materials

Waste cork is granulated by suitable machinery, sifted, and then ground to powder. The finest dust is mainly used in the manufacture of linoleum. Slightly coarse cork is used for the manufacture of cork carpet, but in both cases the binding material is prepared linseed oil.

Still coarser cork, of the size of peas, mixed with a proportion of fine, is employed in the manufacture of mats, insulating materials for water and steam pipes and cold storage chambers, packing, artificial corks for bottles, &c

In order to bind the particles of cork together are employed (1) collodion, used for bath mats and stoppers. The granulated cork of various grades in suitable proportions is mixed with the solution and the pasty mass pressed into perforated moulds, these are then heated in a suitable chamber provided with a condensing appliance to recover the solvents, ether and alcohol. 'Suberite' belongs to this class The mat is removed from the mould and 'buffed' (sand-papered) on a buffing machine similar to that used by cabinet makers. Cork stoppers are made in a similar manner, and also insulating slabs and tubes for covering steam and water pipes.

(2) Gluten, casem, and glue mixed with only materials and resins and dissolved in suitable solvents, after mixing with the cork (rendered more or less insoluble by means of formalin, tannin, dichromates, lime, &c), have also been used and patented for the production of cork slabs, &c, from granulated cork. The cork of various degrees of fineness is mixed with the cement, subjected to powerful hydraulic pressure in moulds, and in some cases heated up to 200° to harden the albuminous matters of the cement.

(3) Mixtures of cork with silicate of soda and barytes or other pigment have been used for steam-pipe coverings

Slabs of cork have also been made by soaking cork in soap solutions, then drying, treating with milk of lime, again drying and pressing the resultant mass into heated moulds

Another use for cork waste, more or less finely ground, is in the production of floor. In this operation ground cork is mixed with a cement, such as Portland cement, or more often a mixture of cork and calcined magnesia is mixed with a solution of magnesium chloride and the pasty mass spread on the floor and allowed to harden, other salts, such as lead acetate, are sometimes added

Cork dust is also used as a packing for the hollow spaces between the walls of cold-storage chambers, round the pipes of ice plants, &c, no binding materials being employed Mixed with pitch, granulated cork is used as the filling between the outer and inner soles of boots,

especially American. It forms a warm, water-proof and resilient layer

Granulated cork of various degrees of fineness is also mixed with the paint used to coat iron surfaces where there is danger of the condensation of moisture, such as in the cabins of steamers, in breweries, factories, laundries, and the like

Hy I

CORNISH CLAY v CLAY

CORNISH STONE v CLAY and CHINA STONE

CORN OIL v MAIZE OIL.

CORNUTINE v ERGOT

CORONITE. An explosive consisting of nitroglycerine, 38-40 parts, nitro-cotton, 1-1½ parts, ammonium nitrate, 26-28 parts, potassium nitrate, 3-5 parts, aluminium stearate, 11-14 parts, rye flour, 8-11 parts; wood meal, 2-4 parts, liquid paraffin, 2-4 parts (v Explosives)

COROZO. Vegetable ivory The seeds of

Phytelephas macrocarpa

CORROSION OF METALS. By the term 'coriosion'is generally understood the disintegration of a metal through oxidation. Usually a skin of oxide is formed which more or less protects the underlying metal from attack Copper, zinc, lead and aluminium are cases in point. Iron,1 on the other hand, is exceptional, in that, when oxidised by exposure to ordinary atmospheric influences, a pulverulent, hygioscopic mass of oxide, known familiarly as fust, is produced, which not only fails to protect the underlying metal, but actually accelerates further corrosion For this reason, and also because non is of such enormous commercial importance, the various problems associated with the corrosion of non have received wide consideration and been made the subject of numerous researches; only in very recent years have systematic attempts been made to investigate the corresivity of the other commercial metals

Corrosion of iron. It is impossible here to review or discuss the voluminous literature which has appeared on the subject during recent years, and for particulars, including bibliography, the reader is referred to Friend's Corrosion of Iron and Steel (Longmans and Co., 1911), and to The Corrosion and Preservation of Iron and Steel, by Cushman and Gardner (McGraw-Hill Book Co., New York, 1910)—It will suffice here to give a short account of the questions at issue

Conditions essential to the corrosion of non—Before passing on to consider the various theories that have been offered to account for the mechanism of corrosion, it is important to have a clear conception of a few of the fundamental facts. These may be briefly summarised as follows—

- I Liquid water, alone, both at ordinary temperatures and at the boiling-point, is without any appreciable action upon iron. In the course of time the metal frequently becomes very slightly tarmshed, but no trace of rust-formation can be detected
- 2 Water vapour is without visible action upon iron at ordinary temperatures upwards to about 300°C. At 350°C, and somewhat above, tarnishing occurs owing to a superficial layer of ferrous oxide being produced. At still higher
- 1 The word 'iron' is, both here and in the sequel, used in its broadest sense to cover wrought iron, steel, and cast iron

temperatures a protective layer of ferroso-ferric oxide is produced, which, however, is not rust.

3. Dry air or oxygen has no visible action upon iron at ordinary temperatures. It is not until the metal is raised to about 220°C, that a pale yellow tint is acquired, which, as the temperature rises, becomes successively straw-These 'temcoloured, purple, and finally blue pering' colours, as they are termed, are due to the formation of thin layers of oxide, and are consequently not produced in vacuó (Roberts, Trans Inst Mech Eng 1881, p 710), but are produced in perfectly dry air, such as that obtained by prolonged desiccation over phosphorus pentoxide (Friend, J. Iron Steel Inst. 1909, n 172).

4. A mixture of water vapour and air is without action on iron at ordinary temperatures. This was demonstrated by Dunstan and his coworkers (Trans. Chem. Soc. 1905, 87, 1548), who further showed that if, under the above conditions, liquid water is allowed to condense on the surface of the metal corrosion proceeds rapidly

From the foregoing it is evident that the rusting of iron is not an example of direct aerial oxidation, but that the presence of both liquid

water and air or oxygen is essential

Theories of rusting -- Now although all ordinary nons readily corrode when exposed to the combined action of distilled or natural waters and air, it is of considerable academic importance to inquire whether or not a pure specimen of iron will corrode when exposed to perfectly pure air and water. In 1888 Crum Brown (J Iron Steel Inst 1888, n 129) summarised buefly the most important facts then recorded on the corrosion of non and supported, though he did not originate, the then current idea that the action is primarily the result of acid attack—the acid, in most cases of aerial oxidation, being carbonic According to this theory the metal is first attacked by dissolved carbon dioxide (carbonic acid) giving ferrous carbonate, or bicarbonate, and hydrogen ferrous salt is then oxidised by atmospheric oxygen, and the basic ferric carbonate potentially formed is, in presence of water, converted into hydrated ferme oxide (rust). As the carbon dioxide is thus wholly eliminated at the end of the cycle, it may be supposed to renew its attack on the iron

This theory received vigorous support from Moody, who proved by a very ingenious experiment (Proc Chem Soc 1907, 23, 84) that rust is formed by the action of oxygen upon iron in solution and not by its action upon the metal In other words, the rusting of iron is preceded by its solution Moody further satisfied himself, as the result of a careful series of experiments (Trans Chem. Soc 1906, 89, 720), that water and oxygen alone are quite insufficient to induce rusting In experiments where iron had remained bright for many weeks in contact with pure air and water, the introduction of a minute quantity of carbon dioxide at once determined rusting. The experiments at once determined rusting The experiments of Friend (J Iron and Steel Inst 1908, ii 16; Proc Chem. Soc. 1910, 26, 179) led to the same conclusion.

On the other hand, Whitney, in 1903 (J. Amer Chem Soc 25, 394), applying modern electrochemical theory to the process, expounded the

diametrically opposite view that water and air alone suffice to establish rusting According to modern views, the action of an acid on a metal is ascribed to an exchange of electric charges between the hydrogen ions of the acid and the electrically neutral atoms of the metal, the hydrogen being deposited on the metal in the molecular state, and the metal atoms passing into the solution as metal ions. Inasmuch as pure water is regarded as being to some small extent ionised into hydrogen ions and hydroxyl ions, it is to that extent, electrochemically speaking, an acid, and when iron is immersed in it, there should be a deposition of hydrogen on the iron and an equivalent formation of iron ions The water round the iron now contains ferrous ions and hydroxyl ions, or in other words, is a dilute solution of ferrous hydroxide. If oxygen has access to the ferrous hydroxide solution, hydrated ferric oxide will be formed and deposited as rust

Such in outline is Whitney's theory, apart from the detailed application of Nernst's theory of electrolytic solution pressure, which he also introduces. The explanation just given for the action of pure water on pure and physically homogeneous iron, involves the deposition of hydrogen as a film of gas on the immersed portion of the metal As this film would prevent further contact between the metal and the water, it is supposed that it slowly dissolves in the water and escapes by diffusion, or is oxidised by the dissolved oxygen, so enabling the action to continue This theory received considerable support from W H Walker and his co-workers (J Amer. Chem Soc 1907, 29, 1251), who concluded that they had 'substantiated the work of Whitney, and shown that non does dissolve in pure water in the absence of carbon dioxide and oxygen

Dunstan, Jowett, and Goulding (Trans. Chem Soc 1905, 87, 1548).

It may be added, in passing, that the names of Cushman (The Corrosion of Iron, Bulletin No 30, U.S. Dept of Ague 1907), Tilden (Trans Chem Soc 1908, 93, 1358), and of Heyn and Bauer (Mitt Konigl Materialprufungsamt, 1908, 26), must be added to those who believe that iron will corrode in the presence

of pure water and oxygen

Nearly all the experiments just referred to were carried out with iron of ordinary com-mercial purity. This circumstance made it possible for supporters of the 'acid theory' to ascribe the rusting in absence of carbonic acid, to the presence of acidic substances generated from impurities in the iron itself Lambert and Thomson (Trans. Chem Soc 1910, 97, 2426) found that carefully purified iron remained unchanged when exposed for several months to purified air and water Any specimen of commercial iron, or of iron prepared with fewer precautions than the immune sample, rusted quite readily in the same circumstances. From their experiments, the authors concluded that whilst pure non will not rust in pure air and pure water, iron will do so if it contains only a small amount of impurity, even if the impurity is neither acid in itself nor likely to give rise to acid.

From the foregoing it is evident that opinions widely differ as to the chemistry of corrosion. Academically, therefore, the problem still awaits solution In actual commercial practice, however, since neither water, air nor iron itself, is ever quite pure, it may be safely postulated that corrosion will result whenever these three are

brought into contact.

The Mechanism of Corrosion —Aitchison (J. Iron and Steel Inst, 1916, I., 77) has outlined an attractive theory according to which, in the case of a perfectly annealed specimen of pure iron, a difference of potential exists between the crystals of ferrite and the amorphous, intercrystalline material, thereby leading to corrosion. Any potential difference between the crystal grains themselves, due to strain, in a less perfectly annealed specimen will add to this effect, the total corrosion being the sum of the two actions. In the case of iron containing carbide the main differences of potential will lie between the carbide and ferrite, the latter functioning anodically and oxidising away. The case of steels containing alloying metals is interesting. Potential difference exists between the carbide and solid solution, the latter acting as anode If the alloying element is associated entirely with the carbides, as in the case of small quantities of W, V, and Mo, they do not decrease corrosion Ni, on the other hand, enters the solid solution, reducing its solution pressure and hence the potential difference between it and the carbide, thus hindering corrogion

The ferroxyl indicator The study of the susceptibility of non to rust, and especially of the localisation of susceptibility, has been aided considerably by a device due to Cushman (lc and J. Iron and Steel Inst 1909, 1–33) and Walker (J Amei Chem Soc 1907, 29, 1257) The iron to be studied is fixed in a jelly made of agar-agar, in which small quantities of phenol-phthalein and potassium ferricyanide have been incorporated. Wherever the iron is being attacked, blue ferrous ferricyanide (Turnbull's blue) appears, whilst the second electrode in the circuit declares itself by the pink colour developed from the phenol-phthalein. The pink colour indicates the presence of hydroxyl ions

· The chemical nature of rust -Rust is a pulverulent, hygroscopic mass, ranging in colour from brown to red, and consisting of iron oxidised more or less completely to hydrated ferric oxide As a rule, the browner the rust the greater is the extent of hydration ferrous oxide is generally present also, particularly if the original metal has not been completely oxidised away. This ferrous oxide has a blackish appearance and may be found in the innermost layers close to the metal (see analyses by Moody, Trans. Chem Soc. 1906, 89, 726, Tilden, ibid. 1908, 93, 1358; Gaines, Chem. News, 1910, 101, 205; Steel, J Soc. Chem Ind. 1910, 29, 1141, Rust also contains free water in addition to that chemically combined with it, and it readily absorbs carbon dioxide from the air These two properties serve to render it an active stimulator of corrosion, for the carbon dioxide dissolves in the moisture yielding carbonic acid This attacks a portion of the iron converting it into ferrous carbonate, which in turn is oxidised by the air to rust. The liberated carbon dioxide is now free to attack further portions of iron in a sımılar manner.

Graphitisation is a form of corrosion to which cast iron is particularly liable, the mass becom-

ing so soft as to be readily cut with a knife The metallic iron is more or less completely oxidised, mainly to ferrous oxide Frequently portions of this latter have been washed or dissolved away, the remainder collecting within the pores of the now isolated mass of graphite. Examples are afforded by Berzelius (Traité de Chimie, 1831, vol 111 ), Rennie (Min. Proc Inst Civil Eng 1845, 4, 323), Draper (Chem. News, 1887, 56, 251), Le Naour (J. Iron Steel Inst. 1898, 1 526), Holgate (Gas World, 1913, 58, 479), and Of these Berzelius drew attention to the cast-iron cannon balls raised at Karlskrona from a ship that had been sunk fifty years previously They had been converted into a soft, porous mass which spontaneously heated upon exposure to air Rennie mentions the case of cast-iron guns reclaimed from the sea in 1822, off Holyhead, after an immersion of about 100 years. When brought out of the water they were quite soft, but upon exposure to the air, they hardened to such an extent that they were used to fire salutes when King George IV passed through Holyhead on his way to Dublin. It was observed that these old guns gave louder reports than any others As a rule, this kind of corrosion does not take place with the purer forms of commercial iron, although a few cases of similar softening are on iccord where wrought iron has been subjected to prolonged immersion in sea water (Lidy, Engineering News, 1897, 39, The percentage of carbon in such cases is, of course, much less than with cast iron, but is frequently much higher than the normal in consequence of the solution and removal of some of the iron

Influence of dissolved substances on rusting. The effect of dissolved salts in the water has received great attention in both the laboratory

and industrial study of rusting

As a general rule, a dissolved salt might be expected to stimulate corrosion by increasing the conductivity of the water If the substance gives an acid solution this action will be all the more marked On the other hand, concentrated salt solutions do not dissolve oxygen to the same extent as pure water, so that if the metal is completely immersed, retardation of rusting might be expected on this account The above conclusions are borne out by the experimental work which has been done. Acids and ammonium salts stimulate corrosion very considerably; sodium chloride stimulates in dilute solutions and retards in concentrated solutions. Generally speaking, as the concentration is increased the corrosivity increases to a maximum (critical concentration), and then, unless interrupted by saturation of the solution, falls practically to zero (limiting concentration). The limiting concentration in certain cases is found to have an extremely low value, eg in alkaline solutions and in chromates and dichromates. Chromates and dichromates, and other oxidising agents are known to render iron passive, and Dunstan and Hill (lc) have shown that alkalis also produce passivity, so that the effects may all be due to this cause In any case there is no doubt that above a small concentration, many substances, such as potassium and sodium carbonates, permanganates, iodates, chromates, and dichromates, borax, &c, will prevent rusting entirely. This property of

the chromates is beginning to find industrial application in the painting of iron work, and the protection of boilers. When more than one electrolyte is present in the water the effects are more complicated and it is found that the concentration of the inhibitor must usually be considerably greater than when it is present alone, if corrosion is to be prevented. An interesting special case is the behaviour of potassium dichromate in the presence of other salts. In such cases the dichromate being an acid salt reacts with the other salt to produce free acid, which destroys the passivity and allows rusting unless the dichromate is in large excess. For practical application as a rust preventive in such cases, potassium chromate which produces no acid is equally effective (Friend and Brown, J. Iron and Steel Inst. 1911, I. 125)

The temperature of a solution exerts an important influence upon its relative corresivity A 3 pc solution of sodium chloride at 10°C, is considerably more corrosive than tap water at the same temperature, but as the temperature rises the relative corresivity falls, until at 21°C the solution is appreciably less corrosive than tap water. The behaviour of sea water, which contains approximately 3 p c of sodium chloride, is of practical interest to the engineer it is considerably more corrosive than fresh water, at 13°C the two waters act similarly; whilst at all higher temperatures sea water is less corrosive than tap water (Friend and Brown, Trans Chem Soc 1911, 99, 1302) In the western part of the tropical Pacific Ocean a temperature of 32°C. is frequently attained; whilst in the Red Sea and Persian Gulf temperatures of 34 4°C and 35.5°C. respectively have been recorded. Such waters, therefore, are much less corrosive than river waters at the same temperature. In the Aictic Ocean, on the other hand, the temperature often hes below 0°C, and such waters are much more corrosive than fresh waters Between these two extremes all grades of temperature occur with corresponding fluctuations in the corrosivities

It is a matter of common knowledge that alkalis, if present in sufficient quantity in solution, will completely inhibit corrosion The merest traces of sodium or potassium hydroxide will suffice to do this provided the solution is protected from neutralisation by the carbon dioxide of the air Upon exposure to the ordinary atmosphere, however, carbonates are produced, and these tend to stimulate corrosion at ordinary temperature unless the concentration of the alkali hydroxide is sufficiently great to yield at least the limiting concentration of alkali carbonate, namely, a 0.25 p.c. solution. An interesting case arises when iron is immersed in alkaline solutions containing dissolved inorganic salts For example, iron will remain untarnished for an indefinite period in a 1 p c solution of potassium hydroxide Upon addition of potassium chloride however, corrosion is readily induced. It is possible, however, to increase the concentration of the alkali to such an extent that corrosion is entirely inhibited, no matter what concentration of chloride be added. The minimum amount of alkali required to inhibit corrosion varies with the nature of the added salt, and rises with the concentration of the same

until saturation of the last-named is reached. It is important to note that when iron corrodes in alkaline solution, the rust invariably takes the form of pitting, being localised at various points on the surface of the metal. When scraped away its interior is found to be dark, consisting largely of ferrous oxide, and the metal is observed to be eaten out or corroded into a small pit The original cause of this pitting is probably the presence of traces of superficial impurity; nevertheless, even the purest forms of commercial iron, such as Kahlbaum's electrolytic metal, readily pit in this manner. This is a point of considerable importance to boiler engineers, masmuch as alkaline feed waters are hable to produce serious trouble. It is not difficult to imagine that a large iron boiler or tank might very well lose several pounds in weight through a uniform superficial corrosion, and not be seriously injured thereby loss of only half an ounce, if localised in the form of a deep pit passing entirely through the metal, might be disastrous. The remedy in such cases appears to lie in raising the concentration of the alkalı until the inhibiting point is reached

Various factors influencing corrosion -1t has already been indicated that the concentration of dissolved oxygen at the surface of the metal has a very important influence on the progress of rusting Practical experience confirms this in many ways Pipe lines conveying water have been known to fail mainly on account of the extreme aeration of the water. Bridge structures always rust most readily near the water level, particularly if the water is in slight motion (see Cobb, J. Iron Steel Inst. 1911, I 170). Iron work which is deeply immersed in water or deeply buried, rusts much more slowly than that which is nearer to the atmosphere The oxygen used up at the metal-water surface, is gradually replaced by diffusion from the airwater surface, and anything which prevents or decreases the rate of diffusion, such as reduction in the size of air-water surface, or covering it with a layer of material impermeable to oxygen, will delay rusting The rate of flow of the water past the metal might be expected to have an influence in this respect, and Heyn and Bauer (Mitt Konigl Material-prufungsamt, 1910, 28, 93) have actually found that slight motion increases the rate of corrosion for both cast iron and steel, especially the former. With an increased rate of flow, however, the corrosion decreased again. At high temperatures, it is naturally found that the tendency to rust is greatly increased although, under many conditions, this may be compensated for by the diminution in the supply of oxygen caused by the lower solubility

Influence of physical condution—It is an experimental fact that two samples of iron may possess precisely the same chemical composition and yet exhibit a marked difference in their powers of resisting corrosive influences. This is attributable to variation in the physical properties of the metals. A series of interesting experiments was carried out by Andrews (Min. Proc. Inst Civil Eng. 1894, 118, 356) on this point, with the object of determining the influence of tensile, torsional and flexional stresses, respectively, upon corrosion. In so far as the first of these, namely, the tensile stress, was

concerned, it was found that the strained portion was cathodic to the unstrained when both were connected in a cell containing sodium chloride solution; in other words, the strained portion resisted corrosion more successfully than the unstrained On the other hand, Hambuechen Bull Univ. Wisconsin, Eng Ser. 8, 1900), after a similar series of experiments, was led to an opposite conclusion. This apparent discrepancy is explained by the work of Richards and Behi (Pub Carnegie Inst, Washington, 1906), which makes it clear that any difference of potential existing between strained and unstrained portions of metal is very small and is greatly influenced in direction by local encumstances Burgess (Trans Amer Electrochem Soc 1908, 13, 17) found that non deformed by stretching beyond the elastic limit corrodes more rapidly than unstrained portions, and this is supported by the work of Heyn and Bauer (J Iron Steel Inst 1909, I 109). As regards the influence of torsional and flexional stress, Andrews found the unstrained portions to be more susceptible to corrosion than the strained Here again the extent of the strain and the local conditions will each exert due influence upon the result

In 1911 Hanemann drew attention to some interesting effects produced by heat treatment upon the resistance of steel to acid attack (Stahl und Eisen, 1911, 31, 1365). He showed that each steel, according to its carbon content, appears to have an optimum quenching temperature, at which a maximum tensile strength is obtained. His results further indicated the existence of a close connection between the tensile strength and solubility of the steel, the steel with the highest tensile strength exhibiting the greatest solubility in acid. From the foregoing it is evident that the physical condition of the metal exerts an important influence upon its corrodibility.

Influence of chemical composition—The chemical composition of non or steel has an important bearing upon its resistance to corrosion. This is most marked in the case of east iron and steel, since these have a wider range of composition than wrought non. The most usual effects of alloying elements may be summarised as follows (Friend, Paper read before Staffs Iron and Steel Inst. Feb. 21, 1914).

1 Some elements, such as sulphur, exist in steels in the form of relatively oxidisable compounds. In mangamferous and cupriferous steels the sulphur is present mainly as the sulphides of copper and manganese. These readily undergo oxidation to sulphureous acids which stimulate corrosion.

2 Not a few elements combine with iron forming alloys or compounds highly resistant to corrosion. Such appears to be the case with phosphorus, nickel, chromium, and silicon.

3 The presence of one element may tend to alter the chemical or physical condition of a second element present in the iron or steel. For example, silicon tends to throw the carbon out of combination with the iron, and effect its separation in the form of graphite. This is a well-known reaction in blast furnace practice, and the consequence is that the corrodibility of the metal is not only influenced by the presence of the silicon per se, but also by the altered condition of the carbon.

4. Alloying elements tend to produce segregation or unequal distribution of the components of the metal during solidification. Serious differences of potential are thereby initiated and the liability to corrode is proportionately enhanced.

In addition to lack of homogeneity in the metal itself it must be remembered that contact with a less electro-positive metal, the presence of small quantities of mill scale (Fe<sub>3</sub>O<sub>4</sub>), or even of rust itself, will cause the iron to become the soluble electrode in a galvanie circuit and thereby hasten its corrosion It is therefore always advisable to avoid making contacts with other metals, or even with other kinds of iron, at points where electrolytic action might take place The fact that non in constant use does not corrode so fast as when at rest has been ascribed to the fact that any oxides formed are quickly removed and do not remain to exercise a galvanic influence

Influence of carbon and manganese—Two of the most important alloying elements in steel are carbon and manganese, and the influence of these upon the corrodibility of the metal has been made the subject of considerable study Hadfield and Friend (J. Iron Steel Inst. 1916, I. 48) summarised their results as follows. I The addition of carbon from 0.03 to 1.63 p.c. to pure iron containing less than 0.2 p.c. of manganese, results in a steadily increasing rate of corrosion both in tap water and in sea water: an initial fall in corrosion with a rapid rise to a maximum with 1.05 p.c. carbon in alternate wet and dry, and a rapid rise in corrosion to a maximum with 0.8 p.c. carbon when exposed to the action of dilute sulphuric acid.

2 The addition of 0.7 p.c. of manganese to the above steels results in a slightly increased corrosion in tap water, sea water, and in alternate wet and dry tests, with a carbon content of up to about 04 to 06 pc. With higher carbon contents the manganese generally affords a slight protection, which is very decided in the wet and dry tests. On the other hand, the presence of the manganese enormously increases the solubility of the steels of all carbon contents m dilute sulphuric acid. This is a point of great commercial importance since in ordinary steels the manganese content is very hable to fluctuate between 0.2 and 0.7 p.c. and it is very evident that the latter are unsuited for acid atmospheres or liquors For example, in large towns considerable quantities of sulphur acids are found in the atmosphere, owing to the combustion of bituminous coal, which, containing a small percentage of non pyrites, leads to the formation of sulphur dioxide, and this dissolving in rain rapidly oxidises to sulphuric acid, giving the rain water an acid reaction. It has been shown that in the industrial quarters of a large manufacturing town this acidity will sometimes represent as much as 90 lbs of sulphuric acid per acre per annum, whilst in the residential quarters it will probably not be more than about 25 lbs The sulphune acid, however dilute, coming in contact with iron-work, accelerates the rusting, and by its corrosive action renders more difficult the protection of the metal

For such districts steels of very low manganese content would appear to be most suitable.

3 By increasing the manganese content to

2 pc. upwards the corrodibility of the steels is greatly decreased in neutral media, although in dilute sulphuric acid the solubility is greatly enhanced.

Numerous other researches on the influence of carbon, nickel, chromium, copper, &c, have been carried out by Bieuil, Chappell, Altchison, Buch Diegel, and others, but for accounts of these the leader is referred to the J Iron Steel Inst.

Relative corrodibilities of wrought iron and steel—The question as to which of the two metals, wrought non or steel, is the more resistant to corrosion is one which has attracted much attention The subject has been discussed by several investigators (see Trans Faraday Soc. 1916, 11, part 2), who point out that the mean result of examining a large number of samples of wrought non and of steel after exposure to corroding influences indicates that the two metals are practically equally resistant. Nevertheless wide divergences exist between the different irons and steels in individual cases This is not necessarily due to any irregularities in the metals themselves, but rather is it attributable to the fact that no one metal can be expected to offer an equal resistance to all kinds of corrosive influences The problem thus becomes one of determining which variety of iron or of steel is the most suitable for any particular purpose. It is conceivable that certain makes of wrought iron will prove most useful in certain circumstances, whilst under other conditions it will prove better to employ Field tests, which are urgently needed to throw light upon these obscure problems, are in progress.

Corrosion of cast iron —Comparatively little research work has been published on the corrosion of east non In 1883 Gruner found that cleaned cast non was less attacked by an and moisture than wrought iron or steel (Compt rend 1883, 96, 195), but was more readily attacked by dilute acids. This latter observation was confirmed in 1915 by Friend and Marshall (J Iron Steel Inst 1915, I 353), working with a mild steel and grey east iron In neutral corroding media there was little to choose between the two metals. The same investigators found (ibid 1913, 1 382) also that a variation in the percentage of silicon between the limits of 1 2 and 2 3 p c exerts no influence per se upon the corrodibility of the The fact that in many cases cast iron objects appear to have a longer life than corresponding ones in wrought non or steel is attributable to the protection afforded by the outer skin formed on the cast iron during casting This may also account for the greater resistance offered by cast-iron pipes to the destructive action of stray electric currents (Brown, Electrical Engineer, New York, 1898, 26, 441)

Acceleration tests—In order to determine what particular variety of iron or steel will resist corrosion most effectively many attempts have been made to employ rapid chemical tests in the laboratory. One of the most usual of these consists in immersing different samples of the metal in dilute sulphuric acid for definite intervals of time and noting the losses in weight Those specimens that dissolve most readily are regarded as most readily corrodible. It cannot be too strongly emphasized, however, that such

tests are of no general value whatever. The only reliable method consists in experimenting with the samples of metal under precisely similar conditions to those to which the metal will be exposed in practice. This, it is true, is a slow process, but the results are sure.

Prevention of rust.—Where the use to which iron is to be put does not admit of any protective coating, it is obviously necessary to consider all the above questions, with a view to providing conditions which shall keep corrosion as slow as possible. The steel plates of a steam boiler, for instance, are subjected to very stringent conditions which often cause extensive pitting, so that special attention has to be given both to the metal of which the boiler is contained and to the feed water An example is described by Huntley (J Soc. Chem. Ind. 1909, 28, 339) of remarkable corrosion in a stand-by borier which could not be prevented by adding alkalis to the water Blisters were found near the water level of the boiler, filled with a liquid which contained ferrous sulphate and sulphuric acid The production of the sulphuric acid was found to be due to the presence of manganese sulphide in the steel. The blister of rust had retained the sulphuric acid, and oxygen passed in to oxidise the sulphide more rapidly than sodium hydroxide to neutralise the resulting acid. In this case, a remedy was effected by the addition of sodium arsenite, which presumably acted as a reducing agent combining with the dissolved oxygen There are various other devices for the removal of oxygen, such as passing the feed water over scrap iron in a closed box, or the addition of an alkaline solution of tannin, which in the boiler forms strongly reducing pyrogallate. The use of dichromates or chromates has already been 1eferred to, although the practice has not yet been widely tested on the large scale Other attempts to protect non have been made by placing pieces of zinc in contact with the metal so that the iron would be protected at the expense of the zine. This practice is not very economical, and usually the effect of the zinc is confined to its immediate locality. In certain circumstances, however, the method is of value. A method of preventing or retaining the corrosion of boilers consists in applying an external current so that the boiler shell is the cathode, an immersed piece of wrought iron the anode, and the boiler-water the electrolyte of 1 to 2 amperes at 4 to 8 volts' pressure is said to show a satisfactory reduction in the extent of the attack on the plates (Harker and Macnamara, J Soc Chem Ind 1910, 22, 1286).

Where conditions permit it, and especially in the case of structural iron work, it is of course a matter of economy to give to exposed iron work a protective coating. The first essential is to provide a coating which shall be as impervious as possible to air and water. Further considerations arise, particularly with regard to the course of events at points where the coating has given way. For small articles a film of vaseline, black lead, or similar material which can be readily renewed is of great practical value. A coating of hot tar or pitch (Angus Smith) has also found wide application. Wire and articles of sheet metal may be protected by means of a coating of a less corrodible metal,

usually zinc or tin. Tin plate finds its widest application in the manufacture of vessels to contain foodstuffs, and other articles for which a long life is not desired. It is less suitable for severe conditions, since at any point where the iron has become exposed a couple is produced in which the more electro-positive iron acts as the soluble electrode and suffers rapid corrosion It is therefore of importance that the tin should be as free from pinholes as possible, and the ferroxyl indicator applied in a thin layer to the surface of the metal is a very valuable indication of the extent to which they exist pinhole is indicated by the development of a blue spot which shows that iron is passing into solution In galvanised or zinc-coated iron the state of affairs is reversed The zinc protects the non first by its own resistance to corrosion, and then when the coating is broken, by the fact that the non is protected at the expense of the more electro-positive zinc. There are several methods now in use for providing the zinc coating in the most suitable form In the hot dip method the iron, after passing through a 'fluxing solution,' is dipped into molten zine The fluxing solution which has for its object the cleaning of the iron surface usually contains hydrochloric acid, and the traces of this which are carried over often have an undesirable effect in producing corrosion of the finished product. In addition to the care necessary to avoid this effect, it is very important to control the bath so that a sufficiently uniform and thick layer of zine shall be produced on the iron Other methods of depositing the zinc are by electrolysis and by exposure to the vapour of zine (Sherardising), and it is claimed that both these methods produce a more resistant coating than the hot dip process. The uniformity of a zine coating can be tested by dipping the metal in a hot concentrated solution of caustic soda (Walker, Proc Amer Soc, Testing Materials, 1909, 8, 430) Pure zinc, under such conditions, is practically unaffected, but in contact with iron, as it would be where there were cracks or pinholes in the coating, a current of hydrogen is produced from the non Tested in this way, hot galvamsed non was found by Walker to be relatively free from imperfections, but wet or electrolytically galvanised was frequently porous, indicating the necessity for careful supervision of the rate of deposition of the zinc so as to obtain the most adherent coating number of other protective devices, mostly patented, depend upon the production from the non itself of a resistant surface layer best known of these is the Baiff-Bower method of producing a fine closely adhering layer of magnetic oxide, by the action of steam on the heated metal. In this way a very uniform and durable coating can be obtained on articles which are not too large When the metal does become exposed, however, the oxide forms with the metal a couple which results in stimulated In the Tatlock process, corrosion of the non the magnetic oxide is obtained by heating in a bath of fused sodium or potassium nitrate Coslett process, which is largely used in bicycle manufacture, depends on the production of an insoluble layer of phosphate by the regulated action of phosphoric acid. These and similar methods of protection are all fairly efficient for for this reason, other things being equal, the

the protection of smaller articles or sheets of iron under not too stringent conditions. where, as is often the case with structural and massive nonwork, the water may contain much soluble foreign matter (often of an acid nature), the progress of corrosion must be carefully watched and checked This involves the use of a protective coat which shall be renewable in situ and easily applied For this reason, a considerable amount of attention is now being paid to the protection of non work by paint.

Paints usually consist of an intimate mixture of pigment and liquid vehicle. As regards the latter it is essential that it should excel in chemical permanence, imperviousness towards air and moisture, electric insulation, chemical mertness towards non, resistance towards physical damage, elasticity, and finally in its drying, hardening or setting power The vehicle most commonly in use is linseed oil, which, upon exposure to an, absorbs oxygen, being transformed into a tough, clastic solid called lmoxyn This oxidation proceeds only slowly in the case of the raw oil as obtained by pressure from Imseed, and in order to facilitate the setting, a substance termed a drier is usually added, which, by some means not thoroughly understood, assists the oil to combine with the atmospheric oxygen. To the same end linseed oil is frequently heated, either in the absence of air or whilst a current of air is blown through, a duer may or may not be added in this case, according to circumstances

The absorption of oxygen by the oil is at first accompanied by a slight expansion, amounting in the case of pure raw ('alcutta oil to about 3 pc at 15°C (Friend, Trans Chem. Soc 1917, 111, 162) Hence when thick layers of paint are exposed to the air the outer layer becomes oxidised most rapidly and by expanding causes the familiar crinkled appearance For this reason the painter prefers to apply his paint in thin layers, so that, upon diving, a smooth and uniform coat results

If the oxidation of the oil stopped at the formation of the solid linexyn the paint film would be very permanent Unfortunately such is not the case. Further oxidation takes place, although very slowly, water, carbon dioxide, and other products being evolved, and accompamed by a simultaneous contraction and embrittling of the film In course of time the contraction becomes so great that the film cracks and peels away. Such old films are a source of danger to nonwork, masmuch as they not only fail to protect the metal, but actually stimulate corrosion (Walker and Lewis, J. Ind. Eng. Chem. 1909, I. 754). Hence when cracking of the paint film has once begun it is desirable to scrape off the old paint and apply an entirely fresh coating. All the drying oils are open to this objection, and attempts have been made to utilise other liquid vehicles, such as nitrated cellulose in amyl acetate, &c

The heating of linseed oil, referred to above, induces a certain amount of polymerisation, reducing its permeability to moisture and generally increasing its protective power when used in paint for iron work. Light greatly accelerates the setting of the oil, but unfortunately it also accelerates its subsequent decomposition;

most permanent paints are those containing black or red pigments, since these absorb the shorter rays of light and prevent them from hastening the destructive oxidation of the linoxyn. As regards the pigments, they should be in a fine state of division in order to admit of thorough incorporation with the oil They should also be insoluble in water, and chemically both permanent and mert. The last-named property is largely dependent upon the solubility and in the case of certain pigments, such as the chromes, a very slight solubility in moisture is not altogether a disadvantage, since the dissolved chromate tends to inhibit corrosion. as already mentioned. This was found experimentally to be the case in the Atlantic City tests carried out under the ægis of Committee U of the American Society for Testing Materials. These tests were begun in 1908 and for details the reader is referred to Cushman and Gardner. Corrosion and Preservation of Iron and Steel (New York, 1910), and Gardner, Paint Technology and Paints (New York, 1911), suffice it to say, however, that paint mixtures containing chromates yielded promising results

It is dangerous, however, to use pigments that are markedly soluble in water, since when these have dissolved out of the paint film by repeated rainfalls, pores are left through which moisture and air readily pass to the metallic surface beneath, hading to serious corrosion.

The chief functions of a pigment consist in (1) offering a support to the Imoxyn, thereby affording a harder and tougher film, proportionately more resistant to mechanical injury, (2) reducing the permeability of the paint film to air and moisture, since the pigmentary particles are themselves impervious to these (There are other functions, but for further details reference may be made to Friend, Iron and Steel Institute Carnegie Memoir, 1918) In the case of acrial iron work, a thick coat of paint protects the underlying metal from attack more efficiently than a thin coat, provided the thick coat is not so heavily laid on as to cause crinkling or running. The best results are obtained by multiple coats, two thin coats being more efficient than one thick one of equal weight The non structures should be painted whilst the scale is still on after loosely adherent flakes and rust have been scraped off. In this case the paint will last rather longer than if applied to the pickled or sandblasted surface and the labour of removing the scale is saved. Rust need not be so carefully removed prior to painting as was formally believed to be necessary

A useful paint for structural non work not exposed to sea-water consists of oil with red lead, Indian red and lamp black, the three pigments being in approximately equal proportions by weight

Almost equally good as the red or white lead paints for protection is a bituminous paint made from pitch, the distillation of which has not been carried too far. This must be absolutely free from free carbon and tai acids or ammonium salts, and if applied hot to the structure to be protected, gives very good results, the colour being the main drawback.

There are many circumstances in which Pitting is, of course, an impossible phenomenon such a bituminous paint is preferable to an oil in the case of a liquid. This leaves us with the paint, as, for instance, in gas works, urinals, or following commercial metals to consider, namely

slaughter houses, where there is much ammonia in the air, which has an extraordinarily destructive effect upon oil paints

Corrosion of non in concrete -In view of the widespread use of concrete reinforced with iron or steel the question of the corrosion of the metal in these circumstances is fraught with unusual importance. Owing to the fact that iron, upon rusting, undergoes considerable expansion (Bauermann, J. Iron Steel Inst. 1888, II 135), it follows that should corrosion of the metal occur when it is embedded in concrete, a cracking of the latter becomes inevitable, more air and water enter, so that corrosion proceeds with an increased rapidity and the structure becomes proportionately weakened If the concrete is well made and the whole is undisturbed by stray electric currents it would appear that ferro-concrete structures may last almost indefinitely. The ingredients of the concrete should not be too coarse, otherwise thorough mixing and good contact will hardly be possible. The mixture must be well punned into position in order to eliminate voids in so far as is possible A sufficient thickness of concrete should be applied, since, if it is too thin, it may crack mechanically and thus admit an and water to the metal Substances likely to contain acids or acid producing constituents should be avoided, such as, for example, coke breeze and slags, which frequently contain dangerous sulphur compounds (see Report of Science Standing Committee of Concrete Inst Mar 9, 1911 Friend, Trans Concrete Inst 1917-18, vol 1x)

Concrete may be advantageously coated with some waterproofing material to render it still more impervious, provided such proofing is entirely free from acid or acid producing substances. This course is specially to be recommended in the case of reinforced concrete structures exposed to the action of sea-water, since the salt exerts a deteriorating effect upon the concrete. For this reason Creighton points out that if beach gravel is used in making the concrete it should first be very thoroughly washed to remove the salt, and furthermore salt should not be added to the concrete during building in cold weather to prevent its freezing (J. Franklin Inst., Nov. 1917.)

(J. Franklin Inst., Nov. 1917) Corrosion of Non-Ferrous Metals --Tho corrosion of non-ferrous metals has not been made the subject of so much study as that of non and steel, nevertheless a considerable amount of useful information has been accumulated The rare or precious metals, such as platinum, midium, and gold, are so resistant to oxidation or corrosion at ordinary temperatures that they may be said, for all practical purposes, not to cornode at all Nickel and cobalt are also very resistant to corrosion, although less so than the foregoing metals. Silver tainishes, particularly in contact with sulphur or sulphureous substances, and is readily pitted or corroded by chlorides such as common salt Mercury or quicksilver likewise tarnishes upon exposure to air, but the layer of oxide tends to protect the underlying metal from attack, if the liquid is kept stationary Pitting is, of course, an impossible phenomenon in the case of a liquid. This leaves us with the aluminium, copper, lead, tin, and zinc, together with their alloys. Such metals as rhodium, ruthenium, &c, hardly concern us here, being rather of the nature of chemical curiosities than otherwise

Aluminium —The study of the corrosion of aluminium has acquired a considerably enhanced importance of late years owing to the ever-increasing use of this metal for air-craft, fittings of sea-going vessels, and for domestic culinary operations

Aluminium tarnishes upon exposure to air, and the layer of oxide, adhering closely to the metal, affords good protection in temperate climes against further corrosion. In hot, tropical countries, where moisture is plentiful, the aluminium corrodes more rapidly, and the coating of oxide, although apparently continuous, may be minutely porous. Aluminium dinking vessels acquire a foul taste in such climates, probably owing to the absorptive properties of this porous oxide layer (Desch, Trans. Faraday Soc. 1916, 11, Part 2).

Aluminum is oxidised by prolonged contact with hot distilled water in the presence of air, the insoluble hydroxide being formed. Scala (Atti R. Accad. Linea, 1913, 22, (1), 43, 95) concludes that this is mainly due to the presence of iron, which passes into pseudo solution in colloidal form as forrous hydroxide and catalytically accelerates the oxidation of the aluminum.

The greater the purity of the metal, and the less its variation in homogeneity, the greater is the resistance of the metal towards corroding influences. Consequently annealed samples are more resistant than those which have not been so treated (Bailey, J. Inst. Metals, 1913, 9, 79). Organic acids, particularly when hot and in the presence of dissolved chlorides, act upon aluminium (Watson Smith, J. Soc Chem Ind 1904, 23, 475), whilst boiling with dilute sodium carbonate solution causes the formation of soluble sodium aluminate, from which the hydrated oxide of aluminium is precipitated on passage of carbon dioxide Both pure and crude alcohol attack aluminium fairly readily, and are rendered turbid thereby (Karpinski, Z Spiritus-industrie, 1912, 35, 660) Hydrogen peroxide in dilute solution readily effects the oxidation of the metal (Droste, Chem Zeit 1913, 37, 1317), as do also dilute solutions of ammonia, sodium hydroxide, and sodium carbonate respectively, in the cold (Hale and Foster, J Soc Chem Ind 1915, 34, 464)

Copper and its alloys —Copper tarnishes upon exposure to moist air at ordinary temperatures, and the skin of oxide thus formed tends to protect the underlying metal from attack Water alone does not appreciably affect copper, and the employment of this metal for kettles for boiling water for culmary purposes is quite Copper differs from iron in that it does not decompose steam, even at red heat cupric oxide heated in steam is reduced to cuprous oxide. Nitric acid readily attacks copper, and the metal oxidises in contact with alkaline solutions. Dilute hydrochloric acid attacks copper in the cold more seriously than nitric or sulphuric acid under similar conditions (Hale and Foster). When embedded in damp concrete copper remains in excellent condition (Gaines, J. Ind. Eng Chem 1913, 5, 766) Sulphur tarmshes copper, and organic acids gradually convert it into basic salts (see Verdigris, &c). Solutions of electrolytes such as sea-water, &c, also attack the metal. [See report of Corrosion Committee J. Inst. Metals, 1919, 21, 87]

The corrosion of an alloy is usually a more complicated process than that of a single homogeneous element, since the different constituents in an alloy do not behave alike towards corroding media. Owing to the corrodibility of non and steel various alloys of copper are used commercially on a large scale for special purposes, as, for example, in the manufacture of condenser tubes, millions of which are in use at any moment by the Admiralty. Consequently a study of the conditions under which corrosion takes place assumes a position of corresponding importance. During the last few years increased attention has been given to this subject, notably by the Institute of Metals, and for full details of the work carried out under their auspices, the reader is referred to Bengough's Reports to the Corrosion Committee, as published in the Journal of the Institute

Much so-called corrosion of metal-work really consists of mechanical abrasion. For example, furnace gases carrying with them fine particles of solid material on passing through metal tubes tends to wear them away. This action is accentuated by the presence of chlorides or sulphur compounds, since these yield with the metal compounds less hard in character, and which are in consequence more easily abraded, or they may even give rise to volatile salts which readily disappear as vapour. This is a common source of trouble with copper and its alloys.

Brass is an alloy of copper and zine, and the resistance offered by it to solutions of electrolytes has been studied by many investigators. The action of chlorides in particular has received attention, notably by Desch and White (J Inst. Metals, 1913, 10, 305, 1914, 11, 235; Desch, J Soc Chem Ind 1915, 34, 258), whose results may be briefly summarised as follows: Corrosion of brass takes place mainly by dezincification, for although both copper and zinc are removed in solution, the proportion of the latter metal in solution and in the flocculent precipitate is much higher than in the alloy Consequently a layer of metal is left, in which the percentage of zinc is much less than in the original alloy, and which has an open, spongy texture that readily yields to oxidation under the influence of air or dissolved oxygen This explains the presence of the layer of cuprous oxide frequently observed on the corroded surface of brass tubes, and which is thus probably not of direct, but of secondary origin. The removal of zinc grains proceeds at first along the boundaries of crystal grains, and in such brasses as exhibit twinning (a-brasses) along the dividing planes between twin crystals.

The presence of iron in solid solution in brass accelerates the corrosion, but small quantities of lead up to 1 pc apparently do no serious harm (Bruhl, J. Inst Metals, 1911, 6, 302) With 2 pc lead a protective layer of basic salts is formed, which tends to retard further action (Desch).

An interesting example of corrosion is afforded by a condenser tube taken from a vessel on the Manchester Ship Canal, and described by Desch The composition of the original metal was Cu, 72 72, Zn, 27 04, Fe, 0 15, and Pb, 0 09 Along the bottom of the tube the metal had been almost completely dezincified, and a transverse section showed that practically only spongy copper remained somewhat higher level the spongy copper extended only partly through the metal, but the boundary between the copper and unaltered brass was perfectly clear

Tin, as is well known, greatly enhances the resistance of biass to attack by sea-water, and for this reason bronzes are in considerable demand for marine work The standard Admiralty alloy for condenser tubes contains Cu, 70, Zn, 29, Sn, 10 The protective influence of the tin appears to be largely mechanical, in that it forms in neutral or faintly alkaline solutions of electrolytes a tough adherent layer of basic salt, which can only be

detached with difficulty

The behaviour of bronzes towards acid solutions has been studied by Giolitti and Ceccarelli (Gazz chim ital 1909, 39, n. 557), who find that their susceptibility to acid attack is intimately associated with microscopical

appearance

When embedded in damp concrete (Gaines, J. Ind Eng Chem 1913, 5, 766) both monel metal and manganese bronze appear to resist corrosion exceedingly well Addition of phosphorus to bronzes increases their resistance to corrosion, and phosphor bronzes, which contain up to 175 pc of phosphorus, have proved very valuable in marine work

Alloys of copper and aluminium, such as those containing up to 10 p.c. of the latter metal, have also proved very resistant to seawater and dilute alkalies, a property that is further enhanced by the addition of nickel (Read and Greave, J Inst Metals, 1914, 11, 169), although, curiously enough, the nickel, if it has any action at all, tends to increase the corrodibility of the alloys in tap water, dilute

mineral acids, and vinegar

Lead—(1) Pure Lead The corrosion of lead of exceptional purity has been investigated by Lambert and Cullis (Trans Chem Soc 1915, 107, 210) The metal, having been prepared in a high state of purity by the method of Stas (Bull Acad roy Belg 1860, 10, 295), was finally distilled in vacuo in quartz tubes at a temperature approximating to 1200° C. Its behaviour towards air, water, and a mixture of oxygen and water was then investigated observed that, if the metal was exposed to air only a short time after distillation, its surface was quickly tarnished in the cold, indicative of slight superficial oxidation If the metal, however, was kept in vacuo for several months, subsequent exposure to air did not cause any appreciable decrease of lustre for several days

Other samples of freshly distilled lead underwent rapid corrosion upon exposure to pure water and oxygen. In the case of totally submerged specimens, white crystals of lead hydroxide were formed, together with a brown deposit, probably consisting of hydrated plumbous oxide Pb<sub>2</sub>O,2H<sub>2</sub>O. When the metal was only partially immersed in the water a bright yellow oxide was formed within half an hour of the admission of oxygen The colour gradually turned to a dull red, the change requiring many months for completion, being accelerated by

exposure to sunlight.

A sample of lead kept in contact with water for 12 months in vacuó presented an undiminished lustre, and no evidence was obtained in other experiments indicative of the slightest solubility of the metal under such conditions Admission of pure oxygen led to no visible corrosion or tainishing for more than a week, and even after six months oxidation had proceeded to such a slight extent that the layer of oxide was suffiattenuated to display interference ciently colours

From the foregoing it is evident that pure, distilled lead behaves somewhat differently towards air or aerated water, according to its 'age,' the specimens that have been most recently distilled proving more susceptible to corrosion than those that have had time to age, although the actual mechanism of the corrosion appears to be the same in either case probably this is due to a difference in the physical character of the metal, and it is in accordance with the behaviour of other substances to suppose that the most unstable form or forms will predominate in the freshly distilled metal, thus giving it a heterogeneous character which gradually settles down to a homogeneous condition of maximum stability

(2) Commercial Lead Although commercial lead is a remarkably pure metal for technical and trade purposes, it cannot be compared with the distilled product of Lambert and It is important, therefore, to study its behaviour towards such corroding agents as are commonly met with in practice This is particularly necessary, since the subject is intimately associated with the national health owing to the extensive employment of leaden pipes for the conveyance of potable waters to private dwellings

The results obtained by early investigators are somewhat conflicting in many of their details (see Summary by Heap, J Soc Chem Ind 1913, 32, 771, 811, 847), but the following facts may be regarded as well established Although Lambert found that pure water exerts no appreciable solvent action upon pure distilled lead in finite time, it is generally agreed that distilled water, freed from air by boiling, readily dissolves commercial lead, and even refined lead. Clowes (Proc Chem. Soc. 1902, 18, 46) found that a sample of very pure lead dissolved to the extent of 0.3 gram per million parts of water, and according to Traube-Mengarini and Scala (Mem R Accad Lincei, 1911, 8, 576), the dissolved metal is suspended in the water in colloidal form Admission of oxygen converts the colloidal metal into colloidal hydrated oxide, the particles of which slowly aggregate to give a crystalline deposit, the liquid becoming turbid Oxygenated waters are thus very corrosive towards lead

The question now arises as to the influence exerted by atmospheric carbon dioxide upon the foregoing reactions Water containing dissolved carbon dioxide has but little action upon lead in the absence of oxygen, and the presence of carbon dioxide in oxygenated waters reduces

their corrosive action by effecting the precipita tion of an insoluble carbonate of lead which protects the metal from further attack. for example, Graham (J. Soc Chem. Ind. 1851, 4, 400) found that pyrophoric lead, which is simply the metal in a state of exceedingly fine subdivision, rapidly renders aerated waters poisonous if carbon dioxide is not present, in consequence of the formation and solution of lead hydroxide, which is soluble to the extent of 1 part of PbO in 10,000 parts of water (Pleissner, Chem Zentr 1907, 11, 1055) The presence of 3 p.c of carbon dioxide, however, prevents the water from becoming poisonous owing to the formation of relatively insoluble lead carbonate, 1 part of which requires 4 million parts of water to effect its complete solution Lead is readily attacked by dilute nitric acid in the cold, much less rapidly by hydrochloric acid, and still less by sulphuric acid under like conditions (Hale and Foster, J Soc Chem Ind 1915, 34, 464). Yorke, in 1834, had already observed that

dissolved salts exert an important influence upon the solubility of lead in waters the most powerful inhibitors are phosphates, such as those of the alkalı metals, small quantities of which almost entirely prevent the solution of lead This is due to the formation of a protective layer of lead phosphate which dissolves in water to the extent of only I part in 7 million parts of water Carbonates and bicarbonates are also very effective, and it has proved a useful procedure to pass certain moorland waters through beds of chalk or limestone to reduce their corrosivity. This does not, in all cases, entirely prevent the solution of lead, since a mere trace of carbonate is not sufficient for the purpose, 4 parts of CaCO<sub>3</sub> or MgCO3 per 100,000 of water being required as a minimum Nevertheless, the procedure is highly beneficial (Thresh, The Examination of Waters and Water Supplies, Churchill, 1913; Liver-eege and Knapp, British Association, Birmingham, Section B, 1913). Birmingham water, which comes chiefly from Wales, is treated at its source with powdered chalk for the same purpose.

The foregoing offers an explanation of the fact that waters which are temporarily hard do not attack lead, in other words, the lead does not pass into solution continuously, since a coat of insoluble carbonate is formed on its surface Permanently hard waters, on the other hand, frequently attack lead rather vigorously Sodium silicate diminishes the corrosivity of water. Dissolved nitrates, particularly ammonium nitrate, attack lead vigorously; but sulphates, chlorides, and acctates, as a rule, much less so Calcium oxide and alkalıs generally have a marked action, and damp concrete may produce serious corrosion (Gaines, J. Ind Eng. Chem 1913, 5, 766). Pure rain water and melted snow resemble distilled water in their activity, but rain waters collected in localities where macadam roads are in common use are laden with calcareous dust, which reduces their activity (Heap). Hence the first rain water from such a district, after a spell of fine weather, may have a comparatively slight action on lead, whilst that which falls after the air has been washed may be expected to act more

vigorously. On the other hand, rain water collected in the neighbourhood of manufacturing cities, is charged with intrates, haloid salts, and sulphur compounds resulting from the combustion of coal, and may or may not be very corrosive according to circumstances

The influence of organic matter upon the solubility of lead in water is obscure It is generally believed that waters derived from moorland and peaty sources are particularly corrosive, and this is usually attributed to the presence of peat acids or of carbon dioxide (Thresh) Experiments carried out by Heap, on the other hand, suggest that such waters, though variable in activity, are considerably less active than pure rain or distilled water. Indeed, both distilled and rain water exhibited m his tests less action on lead after contact with peat, even although their acidity was in some cases increased thereby Water containing certain forms of organic matter yield, after prolonged contact with lead, a flocculent precipitate containing the metal and organic material. This tends to deposit on the lead surface and to form a loosely attached protective layer (Heap)

Rise in temperature in general increases the solubility of lead in water

When sheet lead is immersed in a slightly acid solution of lead intrate or acetate it gradually disintegrates, becoming loose in structure, and ultimately is converted into a spongy crystalline mass, owing to the metal having dissolved and separated out again (see Heller, Zeitsch physikal Chem 1915, 89, 761, Cohen and Heldermann, bid 733, Creighton, J. Amer. Chem. Soc. 1915, 37, 2064). This phenomenon is characteristic both of the purest assay foil and of commercial lead (Desch), and possibly indicates the existence of stable and metastable allotropes (Cohen), the latter of which dissolves and recrystallises as the former. Such a hypothesis is, at any rate, in harmony with the observations of Lambert, to which reference has already been made.

Lead antiques in museums frequently undergo a gradual disintegration, being converted into a white powder consisting essentially of lead carbonate. In such cases chlorides are always present in the powder (Matignon, Compt. 1end 1912, 154, 1619)—Lead that has been immersed in sea-water and dried so that crystals of salt remain attached to the metal undergoes progressive destruction, for which the following cyclo of reactions is suggested

$$\begin{array}{c} \text{NaCl+Pb+O+CO}_2 \xrightarrow{>>} \text{Na_2CO}_3 + \text{PbCl}_2 \\ \xrightarrow{>>} \text{PbCO}_3 + 2 \text{NaCl} \end{array}$$

The sodium chloride thus acts as a catalyser, and is able to effect, in the presence of air and carbon dioxide, the disintegration of indefinite quantities of lead

Tin —Metallic tin remains bright and untarnished for long periods, both in dry and in moist air, at ordinary temperatures. In a piece of tin from an aboriginal cemetery in Florida, Keller (J. Amer. Chem. Soc. 1917, 39, 2354) noticed a number of cavitics lined with small shining crystals, both in the form of thin plates and accular in shape. These consisted of basic chloride SnCl<sub>2</sub> SnO. Water is piactically without action upon tin both in the absence

of, and in the presence of, air Tin is also very resistant to the action of dilute sulphune and vegetable acids, and is thus exceedingly useful for protecting iron or steel from corrosion in the manufacture of articles required for culinary purposes. Caustic soda attacks it, but sodium carbonate, and particularly ammonia, are much less corrosive (Hale and Foster). When alloyed with copper or its alloys it greatly enhances their resistance to corrosion (see Bronze, above), and it exerts a powerful protective influence in other alloys, such as type metal (Meyer and Schuster, Zeitsch angew Chem 1914, 27, 12)

Zinc.—Ordinary commercial zinc readily oxidises in air, and is rapidly attacked by acids Minute quantities of aisenic, antimony, copper, and tin, particularly the first two elements,

accelerate the corrosion of the metal

Zinc is largely used in the manufacture of galvanised iron, and waters passing through galvanised iron pipes or stored in galvanised iron tanks usually give a reaction for zinc. As a rule, however, this is not in sufficient quantities to be toxic.

Boiling water attacks zine (Watson Smith, J Soc Chem Ind 1904, 23, 475), the hydrated oxide of the metal separating out—It is attacked by boiling solutions of sodium chloride, as also by hot dilute solutions of alkalis, such as sodium carbonate, zine hydroxide being produced—The hydroxide does not form a produced—The hydroxide does not form a produced—The hydroxide does not form a produced taken In these cases, but, being readily detached from the metal, exposes fresh surfaces to attack.—Dilute solutions of sodium hydroxide or ammonia corrode zine in the cold, but sodium carbonate acts less vigorously (Hale and Foster See also Report of Corrosion Committee J Inst Metals 1919, 21, 59)—J N F

CORROSION, PROTECTION OF METALS FROM. Of the metals in common use iron (including steel) is not only the most corrodible, but because of its extensive and varied applications under widely diverse conditions, its protection from corrosion constitutes by far the most important case for consideration. With the other common structural metals and their alloys corrosion has in some special cases to be provided against, as in the corrosion of condenser tubes, but, in general, copper, tin, lead, and zine seldom need this attention.

In order to protect the surfaces of iron and steel structures methods based on several

distinct principles are employed

1 Exclusion of the agents causing corrosion air, moisture, or water in the case of submerged structures

This may be accomplished by .-

(a) Coating with an impervious oil film, or asphalt, pitch, &c

(b) The use of paints, varnishes, &c.

(c) Coating with another metal not easily attacked by the corresponding agents

2 By forming a non-corrodible skin consisting of some compound of the metal on its surface, such as the magnetic oxide (Bower-Barff process), phosphide, or nitride

3 By the use of a substance which greatly

reduces ('inhibits') corrosion.

4. Placing the iron in an electro-negative at the same temper condition to another metal, so that the former the use of linseed oil is protected at the expense of the latter, or making the surface to be protected the cathode tough elastic coating.

(negative) and furnishing a continuous current from an external source

It will be seen that there is a wide choice of methods, but practical considerations limit their application. In the case of large structures, ships, bridges, tanks, &c., the large surface, the fixed character of such structures, and the necessity for renewal of the protecting coat, determine the use of some form of paint or vainish paint. Deposition of another metal is applicable to a variety of conditions of use, ranging from the oldinary galvanising for tanks, roofing, &c., to electro-deposition of nickel, as in the case of bicycle parts. For small articles generally, processes of inhibiting, oiling, and formation of a surface compound are alternatives.

Before any system of protection can be properly applied it is essential that the metal surface shall be properly cleaned from dirt, mineral oil, rust (the ordinary ferric oxide  $Fe_2O_3$ ), and mill scale (the black magnetic oxide  $Fe_3O_4$ ). Plates from the rolling mills generally show patches of this oxide. Although it is an excellent protection against rusting when thinly and evenly covering the metal, patches of mill scale may set up active corrosion locally by galvanic action. Further, patches of scale are generally covered with fine cracks through which corrosive agents may pass, and the patch may scale off, carrying any protective composition with it, so that it is essential that all scale shall be removed before coating

Rust and scale may be removed by mechanical means, scraping, who brushing, and sand blasting, the latter being very efficient. Commonly they are removed by pickling the plates in sulphuric acid, 25 p.c. In this acid red rust dissolves, and mill scale is thrown off by penetration of the acid, hydrogen being generated and forcing the scale from the plate. Arsenic in the acid is regarded by many as injurious. The plates should then be washed with water jets under fair pressure (if the plates are immersed in still water a 'gummy' film forms), and then steeped in hot milk of lime to neutralise any acid remaining

Temporary protection from rusting is commonly done by smearing small articles with vaseline or heavy oil Several processes of similar nature, but giving more lasting protection, have been patented, the principle of which is immersion of the metal at temperatures up to a red heat in various oils, and withdrawing while still hot, also spraying of metal hot from the rolls with oil. A layer of slaked lime pasted on is said to be more efficient than smearing with oil, and one patent has been granted for a paste of slaked lime made with a volatile solvent, like naphtha

Asphaltic and coal-tar pitch compositions have great value as preservatives of iron and steel. For many purposes the black colour is prohibitive, but for buried pipes, &c, no better preservative can be employed. In 1848 Dr Angus Smith took out his well-known patent covering the use of pitch. The pipes were preferably first heated, dipped in linseed oil, then heated to 300° F, and dipped in hot pitch at the same temperature. He also mentions the use of linseed oil as an addition to the pitch to keep it of the right consistency, and give a tough elastic coating.

Undoubtedly the finest material for this method of protection is natural asphaltum or bitumen, to which enough heavy mineral oil has been added to give a tough and elastic Pitch from petroleum ('maltha') has much the same characters. The Sabin process consists in dipping in a hot coating mixture like the above, draining, and then heating for two hours in a vertical oven at 400° F Steel water mains up to 5 feet diameter, and in sections weighing 5 tons, have been treated by this process in America, and the United States Navy adopted the process for copper fire mains and flush mains, which are rapidly acted upon by salt water. It has also been employed for the small steel tubing for electric conduits which require an elastic and tough coating.

Heavy coal tar is commonly employed for hot dipping of cast-iron pipes. In practice the process amounts to distillation of the more volatile portions of the tar from the hot pipe, leaving a pitch coating, so that success is largely dependent on the correct dipping temperature. The soft pitch obtainable from tar produced in the low temperature distillation of coal will undoubtedly prove one of the best commercial products in this country. Tar acids and ammonia compounds are both undesirable constituents of tar for this purpose, as both tend to

increase corrosion

The anti-corrosive or protective paints are widely applicable. They may be classified as — 1. Linseed oil paints, containing some direr, and usually also a proportion of varinish

2 True varnish paints (enamels) consisting mainly of resins and oil

3. Varnishes made from a natural resin with a volatile solvent. Each of the above will carry pigment, of which a large proportion is usually iron oxide (Fe $_2$ O $_3$ )

4 Asphalts, or coal-tar pitch, with a certain amount of boiled linseed oil, dissolved in a volatile solvent, such as coal-tar naphtha or petroleum

naphtha

In general, whilst the linseed oil paints are cheaper and easy to apply, they are not so satisfactory as the varnish paints, in which the natural gum or resin reduces the porosity of the film. This has been particularly noticed in the case of coatings for plates submerged in water. Also the more clastic the kind of varnish the better the result Class 3 constitutes an important class of protective compositions specially suited to ship use, where the slow drying ordinary type of varnish can seldom be employed owing to lack of time for drying. The volatile spirit rapidly evaporates, leaving the resin and oil binding the pigment.

Cushman and others, from the results on steel plates in water carrying various pigments in suspension, obtained a classification of these ingredients of paints into 'inhibitors,' 'indeterminates,' and 'stimulators' (Proc. Amer. Soc. for Testing Materials, Vols III to VIII., 1903–1908.) The result is, in part, dependent on the purity of the pigment, but, broadly, zinc pigments, chrome green, Dutch process white lead, and some Prussian blues fall in the 'inhibitor' class; lamp black, other carbon blacks, precipitated barium sulphate (blanc fixe), ochre and bright red oxide, and some Prussian blues are 'stimulators.' Such a classification would

appear to have little value if the pigment particles are thoroughly 'sealed' in the paint medium, but if through porosity, or absorption of water, electrical contact between the pigment particles and the iron is established, the activity of the pigments is probably of the order given, the results certainly suggest an advantage of a first coating with one of the inhibitor class

The most perfect protection with paints is afforded by enamels (varnish paints or 'japans') which have been stoved on to the metal, a good example is seen in cycle frames, &c. The stoving treatment causes the resin and oil to flow into a perfect film, free from porosity, and to set with a fine hard surface. The one objection is the hability to chip with a blow. For under-water work stoved metal is said to show little advantage, and Sabin states that unbaked varnish paints are better.

Coating with another metal less hable to corrosion includes zinc galvansing and tin plating, both of which are carried out by hot dipping the thoroughly cleaned plate in the molten metal. For more special work electrodeposition of copper or nickel (usually on a first coating of copper), and of other metals is occasionally employed. Electro-deposition of zinc is also largely employed for certain classes of malk with a former for the second.

of work, such as frames, &c, for ships
These deposits afford a fair protection, but
even with the best class of nickel plating, as
applied to cycle parts, rusting under wet conditions is found to occur. Should the metal protecting film become broken, so exposing the
iron beneath, in most cases this exposite leads
to more rapid rusting, as the iron is electropositive to the protecting metal, and galvanic
action at its expense is set up. Zinc coatings
are, however, free from this defect, and as long
as a good surface of zinc is still in contact with
the iron, it will be the former metal which is
mainly attacked. Tin plate offers a complete
contrast in this respect.

A special process for applying zine is the sheardising process of Cowper Coles. The cleaned articles are packed in metal vessels with zine dust, a mixture of finely divided zine and zine oxide, and heated for several hours at a temperature below the melting-point of zine. This metal slowly penetiates the iron, and to a far greater depth than with hot galvanising; a hard alloy is formed which is practically non-coirodible, and is capable of being polished.

Processes dependent upon the formation of a protective skin date from the old Bower-Barft process, where the heated non was treated with steam, so that a film of the black oxide (Fe<sub>3</sub>O<sub>4</sub>) was formed. The good surface of 'Russian iron' is due to a similar film, and as long as the film remains intact, protection is excellent, but rusting is accelerated once the film is broken A more recent process is that of Coslett, in which the iron is immersed in a dilute solution of phosphoric acid Later the inclusion of a subsequent dipping in chromic acid was patented by Coslett. Richard and Adams have patented a process in which the article is immersed for from 2 to 20 minutes in a solution made by dissolving 12 lbs of manganous carbonate in boiling phosphoric acid (60 fluid ounces of B P. phosphoric acid to 87 fluid ounces of water).

Another process aims at the production of a

film of nitride of non, the article being heated in a suitable gaseous mixture, such as ammonia, or treating the highly heated metal in a bath of salts which will yield gases capable of forming the nitride.

Protection by 'inhibiting' the corrosive action almost invariably involves dipping in dilute chromic acid

One of the most successful methods of preventing coriosion, especially in the case of boilers and condensers, is that of Elliott Cumberland The galvanic action leading to corrosion is overcome by inserting insulated iron anodes. these being connected to the positive pole of a small dynamo; current always passes from the anode to the part to be protected, which forms the cathode A difference of potential of 6-10 volts is usually employed. In the case of brass tube condensers I ampere per 500 square feet of surface is found to ensure complete pro-In the case of boilers it is found that the hydrogen set free at the cathode plates or tubes is very effective in preventing the formation of a hard scale

Corrosion of Boilers—A serious form of corrosion is to be found in the rusting and pitting of the plates and tubes of high-pressure boilers. Before the adoption of mineral lubricants, it was supposed that the decomposition of the lubricating oils by the superheated steam in the cylinders caused the corrosion owing to the fatty acids liberated, but the same effect has been found with hydrocarbon oils, so that the pitting is evidently not due to this cause

There are several causes which lead to the corrosion found in boilers, amongst which are ordinary corrosion, set up by the carbon dioxide and oxygen which occur naturally in all water, and which act on the metal when the boiler is out of use. This form of corrosion can be prevented by filling the boiler quite full with boiled water rendered slightly alkaline with a little lime and closing air-tight whilst at rest The most frequent cause of corrosion in land boilers is dissolved salts, or free acids and alkalis in the feed water, which become concentrated solutions by evaporation Acid waters, notably those collected in peaty districts, are particularly injurious Nitrates and nitrites, chloride and nitrate of magnesia are salts most hable to cause corrosion. Complete analyses of the saline constituents of the water can alone determine the requisite treatment required to prevent corrosive action. Secondly, galvanic action, induced by the difference in potential between the rust formed by ordinary corrosion and the iron, and also by metallic impurities in the iron and specks of foreign metals deposited upon it This action can be prevented to a great extent by the use of zinc blocks inside the boiler and in metallic connection with the iron of the plates, or by the Cumberland system already described A third cause of corrosion is to be found in the solvent action of distilled water upon the metal of the boiler, the action getting less with increase in the quantity of salts in solution in the water. The fourth cause arises from high local temperatures, often not far below a dull red heat, and then iron decomposes water, forming black magnetic oxide of iron and liberating hydrogen.

 $4H_2O + 3Fe = Fe_3O_4 + 4H_2$ 

Overheating is most likely to be due to deposits or incrustations, and in such cases is checked, or stopped, by the sole use of condensed water, or a properly softened natural water, for the feed JS.S.B.

CORROSION AND FOULING OF STEEL AND IRON SHIPS. AND ITS PREVENTION. In the days of wooden ships the hull troubles that had to be guarded against were the attacks of woodboring animals and the loss of speed due to the growth of weed and shell at and below the water line, but with the launch of the first sea-going vessel built of iron, in 1822, comparatively simple problems became of great complexity, and at the present time the protection of our Navy and mercantile matine from corrosion and fouling is a question of the greatest importance, affecting, as it does, both the life and speed of the vessels

Corrosion.—The troubles due to corrosion are by no means limited to the exterior plates of the ship, but also affect the interior of the

structure, boilers, and condensers

The corrosion of non and steel is brought about by the actions which ultimately result in the conversion of the metal into oxides, and these being accelerated by galvanic actions, which it is almost impossible to climinate in a huge metallic structure like the hull of a modern ship in the presence of sea water, makes prevention far more difficult than in the case of an iron building or bridge

In sea water the process of corrosion is accelerated by the salts contained therein, as these aid the galvanic action set up in the metal by various means, such as impurities in the iron, the presence of mill scale on the plates, high temperature, or even the different stresses imparted to portions of the same plate, whilst in certain waters, such as the mouths of rivers on the west coast of Africa, or in harbours where the water contains much sewage, the sulphates are decomposed by various vegetable and animal organisms with evolution of sulphuretted hydrogen, which greatly increases the corrosion of the metal

The examination of the condition of the bottom plates of vessels that have been at sea for long periods, shows, when the protective paints and compositions have been removed, that three distinct forms of rusting may be distinguished

1 Rust patches, generally near the edges of the plates, consisting of small areas of fine rust, probably caused by moisture on the plate when the protective paint was on. These patches are never thick, and cause but little damage to the plates if all rust be removed before a new coating of paint is put on.

2 Rust nodules are a more localised form of corrosion which has attacked the metal where particles of mill scale have been left on the surface of the new plate, or where small particles of slag or foreign matter have been rolled into the plate during manufacture. These give slight pitting of the plate, but seldom to a serious extent.

3 Rust cones are an accentuated form of rust nodules, and are due to the most local form of galvanic action, caused by the presence of a speck of deposited copper, lead, or other foreign metal, or even a small particle of rust

or mill scale left on the surface of the iron, and covered by the compositions used as protectives and antifoulers. As soon as the sea water penetrates these, galvanic action is set up, water is decomposed, rust formed, and the escaping hydrogen pushes up the composition, forming a blister, the hydrogen escapes and water leaks in, the action becoming more and more rapid, and the blister gradually filling with the product of the action—rust. The blister bursts, but the cone of rust has by this time set fairly hard, and continues to grow from the base, the layers of rust being easily distinguished in a well-formed cone, and when the rust cone is detached, the pitting of the metal at the base of the cone is, as a rule, found to be of considerable depth.

Certain portions of a ship's hull are exposed to special risk of corrosion from galvanic action, owing to the bearings of the sciew shaft and valves for the escape of waste water, bilge, &c, being made of alloys containing copper. In order to prevent this, zinc protectors, consisting of heavy masses of the metal, are placed round all openings where copper or copper alloys are present, and the zinc, being more electro-positive than the iron, places the non in the electronegative condition, so that whilst the non escapes injury the zine rapidly dissolves Cumberland process (vide supra) has been successfully employed in such cases, insulated non anodes being fixed to the hull close to the part to which it is necessary to afford protection

The general procedure for protecting the exterior of the bottoms of non and steel ships from corrosion is to coat them with so-called anti-corrosive or protective paints and varnishes In order that this should be successful, much depends upon the state of the hull at the time

at which the protective is put on

The bottom plates of the newly built vessel are thickly coated, as a rule, with mill scale or magnetic oxide of iron Fe<sub>1</sub>O<sub>4</sub>, which is formed on the surface of the plate during rolling, and which, if it remains perfectly intact, would be an excellent protective, as it forms, indeed, the basis of the Bower-Barff process for the protection of iron, but during the shearing and riveting of the plates this scale gets broken off and exists only in patches, which, if left upon the plate, would excite rapid local galvanic action, owing to the fact that this scale is strongly electronegative to the iron, so that it would need only the exciting influence of the sea water to start the action

In naval vessels, the plates are pickled in dilute sulphuric acid, as already described, until all the scale is removed. In the mercantile marine it is an ordinary practice to launch a vessel without any paint on the hull, leaving the active, but fairly even, justing which ensues to remove the scale, the ship being scrubbed down and then protective compositions painted on when it is docked

This method has the advantage of cheapness, and also of giving a very slightly roughened surface, to which the protective composition adheres firmly, and answers well with tramp steamers and other craft where speed is not an object of paramount importance; but with service vessels and liners, slight as the roughening is, it undoubtedly has an influence on the speed.

The hull of the vessel, being perfectly free from mill scale, and as free as it can possibly be got from the ordinary oxide, is then painted with the protective, but for this to prove effective, the hull must be absolutely dry, a condition which it is exceedingly difficult to obtain, as the big mass of iron is generally a few degrees in temperature below that of the moisture-laden air, and under these conditions a trace of moisture condenses on the surface, which is spoken of as 'sweating' of the metal, and this prevents the proper adhesion of the paint to the metallic surface, and also starts rusting under the paint

Oil paints are the oldest form of protectives in use, and, when the plates are painted when hot and dry, amongst the best and most lasting, but the drying of such paints, being dependent upon the slow oxidation of the boiled oil by au into a resm, requires that a considerable time must clapse between the painting of the ship and the coating becoming hard enough to allow of the anti-fouling composition being painted on over it, and as a rule so short a time is allowed for docking that this is a great drawback

The old paints for this class of work used to consist of red or white lead mixed with boiled linseed oil, but it was shown by M. Jouvin that with the perishing of the vehicle the lead compounds were converted into chlorides and oxychlorides by the sea water, and these were then reduced by the action of the iron of the hull with formation of crystals of lead, which set up galvanic action and led to rapid corresion.

Spirit variash protectives came into use when it became necessary to cut down the time expended in docking, and they owe their rapid drying powers to the fact that, instead of waiting for oxidation to form a resinous mass, as in the drying of boiled oil, a resin is dissolved in a highly volatile solvent, being mixed with boiled linseed oil to give the necessary elasticity, and when this is painted on the hull of the vessel, the rapid evaporation of the solvent leaves the resin and oxidised oil binding the body colour together

Shellac, rosin, kauri, and many other gums and resins are used, whilst the solvents may consist of light hydrocarbons, such as the various grades of solvent naphtha, alcohol, and even fusel oil, but in the Navy very low flash-point solvents have now been discarded owing to the

danger in storage

The drawbacks to the last two classes of protectives are that the solvents employed for dissolving resins and gums are mostly affected by water, with the result that if the plates are not dry, the resins are apt to be deposited in a pulverulent condition, whilst the cold produced by the rapid evaporation of the solvent tends to increase the amount of moisture deposited, which assists in the deterioration of the coating Alcohol variishes have, however, the great advantage that they are not so affected by moisture on the metal, or deposited moisture, since the spirit will absorb a moderate amount of water.

These varnishes also, even when dry, are not impervious to gases and liquids. Under a microscope the coating is seen to be full of minute pores, through which the sea water finds its

way to the iron beneath, and, when once corrosion has been started, the protective coating is gradually forced away from the surface of the iron, exposing more metal to be acted upon The best results are secured by employing several coats of the varnish, and applying it only in weather favourable to the operation, plenty of time being given for the thorough hardening of the material before water comes in contact with it

In most of the various paints red oxide of iron is used to give body and colour to the mixture, and, as it is the same colour as the rust that may form under it, it is very popular with the composition manufacturer, and, providing it be free from acid and soluble sulphates, is unobjectionable, but perhaps a better material would be the mixture of finely divided zinc and zinc oxide, such as is obtained in the Belgian process of distilling zinc, as, even after the varnish had perished, the zinc would afford a further term of protection. After extensive trials the U.S. Naval Authorities adopted an anti-corrosive paint of the following composition gum shellac, 7.9 lbs., turpentine, 0.6 gallon, pinc oil, 0.6 gallon, alcohol, 7½ gallons; dry metallic zinc, 9½ lbs., white zinc oxide, 28½ lbs., the above making 10 gallons.

The protection of the interior surface of the hold of a ship presents some special complexities, as it is exposed to the wear and tear of taking in and discharging cargo and one has to take into consideration such sources of trouble as the action of bilge water, the high temperature, and the presence of foreign bodies such as coal-dust and leakages from cargo There are three classes of protectives used for

the interior portion of a ship

Cements,

Bituminous coatings;

Paints

Although coment is very largely used, yet there are certain drawbacks attached to it which render it by no means the ideal material. It has a certain amount of porosity, so that any corrosive liquid penetrates it and acts on the iron beneath; its rigidity causes it to come away from the metal on any strain being thrown on the latter, variations in temperature having the same effect, and where the coating has parted from the iron, corrosion will start and remain unseen until perhaps too late

Bituminous coatings applied hot are far preferable to cement, but are rather more

troublesome to put on

Fouling of ships and its prevention. The question of the fouling of ships' bottoms is one that has existed from the carliest days, and was of even more importance formerly than it is at the present time, owing to the increase in dock accommodation allowing the more frequent cleaning and painting of the ships; but even now in the case of war the loss in speed during blockading operations, owing to the fouling of the vessels, might prove an important factor in a struggle for supremacy

Fouling consists of various marine growths, both animal and vegetable, which attach themselves to the bottom of the vessel, and by increasing the skin friction whilst the vessel is

in motion, seriously reduce its speed

The nature of these growths needs to be

studied in properly equipped marine biological laboratories, as but little is at present known of the life-history of many of the organisms which are found on a ship's bottom. The main factors, however, are that it is only during the germ or spore period of their existence that they can attach themselves to the bottom of the vessel, and that although the full-grown results are found when the vessel is docked, these growths have taken place in situ

The open ocean and the northern seas are comparatively free from these organisms, but directly inland seas or tropical waters are reached they increase to an enormous extent in variety and numbers, and some harbours, such as the port of Sourabaya, are noted for the fouling influence of their waters. Moreover, it is found that the period of the year exercises a very important influence upon the fouling nature of the water, the spring and early summer being the chief time for these growths to find a lodgment on the bottom of a vessel.

The more common growths are serpulæ, generally called coralline, barnacles, mussels, membranæ, and oysters, usually classified together under the name of 'shell,' whilst at or near the water line are found luxuriant growths of sea grass (Enteromorpha compressa and Zostera marina), and towards the bottom ribbon and other weeds, like the tufted weed or bugula, flourish

Fouling commences with the deposition of a slight slime on the side of the vessel, which is apparently of an albuminous nature, and contains the germs and spores of various marine growths, and is so slight as to escape notice

The first visible sign of fouling on a ship is the appearance of green grass at the water line, and this, in bad waters, will begin to show after a few weeks, whilst the growth will spread downwards from the surface for 5 or 6 feet, but rarely extends below 10 feet, as the growth needs plenty of light and warmth.

It is the most widely distributed form of fouling, and appears in nearly all waters, and, once formed, the grass protects the germs of other growths, and so helps on the general fouling.

In northern waters fouling seldom goes beyond the 'grass,' but in warm climates, especially if long in port, 'shell' soon begins to appear, unless a really good anti-fouling composition is employed for coating the vessel's bottom.

Ships trading to or stationed near the mouths of rivers often show curious results in the kind and extent of the fouling, the marine growths being often killed by the fiesh water, but such vessels foul again very rapidly when once more going to sea, owing to slight deposits of mud and calcium carbonate from the river water.

With wooden ships the necessity of protection against the wood-boring teredo worm as well as fouling led to the adoption of copper sheathing, and when the metal used was of a fairly pure character, it was found to be thoroughly satisfactory, and not only to give complete protection to the vessel, but also to enable it to remain at sea for a very considerable period. The cost, however, was high, and attempts were made to prevent the excessive waste of the copper by using zinc protectors, as advocated by Sir Humphry Davy, but it was soon found

that with the prolongation of the life of the copper plates, fouling again became serious, whilst when later rolled sheets of alloys contaming zinc, such as Muntz metal, were tried, the same trouble again appeared, and it has been clearly proved that the power a copper plate exercises in preventing fouling is inversely proportional to its power of resisting the action of the sea water.

The action of the copper in anti-fouling is that, in the first place, the salts in solution in the sea water act upon the copper with the formation of chlorides and oxychlorides, the former poisonous to all forms of marine life in the germ state, whilst the latter formed a coherent pellicle to which the growths attached themselves and which was torn off during the passage of the vessel through the water by the skin friction, thus enabling it to shed any form of growth that might have attached itself to the plate by a process of scaling

It was found, however, that if the copper were not of the purest, coralline growths very soon attached themselves so firmly through the surface of the skin of copper compounds that it was difficult to remove, and gave points of

adherence for fresh vegetable growths

With the introduction of steel and iron ships, as was natural, attempts were made to utilise the same form of protection from fouling which had proved so satisfactory with wooden ships, and from time to time during the past efforts have been directed to sheathing non ships with copper in the same way soon found, however, that this sheathing set up such severe galvanic action with the hull of the vessel as to be impracticable, and although these attempts were abandoned at a very early period in the mercantile marine, the naval services of several countries have tried its use for service vessels, the generally adopted method being to sheathe the iron hull with wood and to attach to this the copper in such a way as to be insulated from the electro-positive iron, but if water penetrated the insulation was broken The alternative of saturating div wood down with paraffin wax has been suggested

With increase in dock accommodation, the tendency has been more and more towards the use of anti-fouling paints or compositions, which could be painted on over the protective coating of the vessel, and a number of compositions have been placed on the market with

varying success for this purpose

When anti-fouling paints were first manufactured, the only fact at the service of the maker was the experience which had been gleaned from copper sheathing, and on the assumption that the action had been due to the poisonous character of the chloride and oxychloride of copper produced on the plates by the salts in the sea water, compositions were made containing large quantities of corrosive metallic poisons, which, held in a varnish of a more or less lasting character, gave the piotection from fouling for a few months

As has already been pointed out, the action of the copper sheathing was due to two causes: first, the poisonous action of the copper salts on the germ life; and, secondly, the stripping of the pellicle by skin friction when any quantity of growth had formed on its surface, an action

termed 'exfoliation' How important this latter action was may be judged from the fact that when the vessel remained at anchoi in a harbour, the sheathing would last from 10 to 12 years, whilst when on active service the same sheathing would last for only 4 or 5.

Finding that poisons alone gave very unsatisfactory results, the manufacturers of anti-fouling compositions began to make perishable varnishes and grease paints, which should, by the deterioration and wasting of their surface, imitate as nearly as possible the exfoliation of the copper sheathing, and such anti-foulers gave under certain circumstances much more successful results. The idea was that the ship, having been coated with a protective paint or composition to preserve the non, a second or more coatings was put on over the protective, consisting of varnish paints which should slowly yield soluble poisons to the sea water, and in doing so would lead to the disintegration of the varnish surface, which, being washed off, exposed fresh quantities of the poisons to the water

Such compositions proved very satisfactory in the mercantile marine, and all the time that a vessel was actively engaged in going from port to port gave excellent results, but directly these same compositions, which had gained a reputation in the merchant service, were tried on Service vessels, they were found to be as unsatisfactory as those which before had been in use. The reason for this failure is Take the case of a liner trading between New York and England: she is run at a high speed between these two ports for the main portion of her life, remaining in the basin for as short a period as possible in order to utilise to the full her money-earning capacity, for such work it is easy to arrange an antifouling composition, which, with the aid of the skin friction of the vessel, would cause just sufficient wasting of the coating to give the best results. When, however, this is applied to a Service vessel, she probably goes out from dock into the basin and iemains there stationary for perhaps months, and then goes to sea, where she is kept manœuvring at various rates of speed for a considerable period During the period she is in the basin, the composition undergoes practically no wasting, and a thin film of carbonate of lime and mud deposits on the surface, which forms an excellent foothold for any germs and spores that may be in the water, and which, on going to sea, have often advanced too far for the growth to be readily got 11d of

The bad results so obtained led to a composition being put on which wasted with sufficient rapidity to prevent any such deposition in harbour, with the result that when the vessel went to sea, the deterioration being accelerated to an enormous extent by skin friction, the composition was practically all exhausted in a very short time

The most efficient anti-fouling compositions all contain copper or mercury as the poison In some cases soluble salts of these metals are employed with a fairly strong and lasting varnish; in others oxides or insoluble salts are used with varnishes which perish more rapidly on their surfaces and allow the action of the

sea water to convert the compounds into soluble salts.

It is evident that with compositions containing such metallic compounds every attention must be paid to the protection of the hull, as otherwise the formation of soluble salts of copper and mercury might result in the deposi-tion on the iron of these metals by reduction, in which case rapid galvanic action would be set up, and would lead to senous pitting of the plates of the vessel

The compositions of the present day may be

divided into two classes

1 Grease paints containing a metallic poison,

2 Quick-drying varnish paints, also charged with the same poisonous constituents The former class has met with a considerable degree of success, but they have also several very serious drawbacks. As regards the prevention of fouling, they give results not very far removed from the best varnish paints, and in certain waters, such as the Mediterranean, have proved themselves to be excellent antitoulers, but they undoubtedly increase the skin friction of the vessel, and so tend to diminish speed, whilst a second trouble is that, although a varnish paint may often be scrubbed and touched up where fouling has been but slight, with a grease paint the whole has to be removed before recoating, and they are much more costly than the varnish paints, with the result that in the merchant service they are used only to a very limited extent Another drawback to their use is that it is impossible to properly scrape and recoat the vessel during the short period of docking which would suffice for a varnish paint

In the manufacture of varnish paints it is necessary to mix with the solution of gum or resm in a volatile solvent a sufficient proportion of linseed oil to give elasticity to the vehicle, and these paints have been in use in England since about 1870, the greatest advance in them being due to the introduction of the hot process in their manufacture, which enables a more intunate mixture to be made with greater adhesive power and a smoother surface

The rate at which disintegration of these paints takes place is governed largely by the kind of gum used in the varnish, and by the amount of mineral matter introduced as colouring and poisonous material, the solubility of these having a considerable influence on the

rate of disintegration

With a tough and tenacious vehicle, which by itself would undergo but little disintegration under the action of sea water, it is quite possible to use soluble mercury or copper salts, which are liberated only very slowly during the gradual destruction of the varnish On the other hand, if a rosin varnish be employed, insoluble bodies, like the oxides of copper and mercury, the subchloride of mercury, or Scheele's green (copper arsenite), may be used, the oxides and other salts, when exposed to the sea water, being very slowly converted into soluble chlorides.

Many compositions contain arsenic and arsenic compounds, such as sodium arsenite, as these had been found effective as weed killers under ordinary conditions, and it was expected that in a composition they would prevent the

growth of grass at the water line, which, as it caught the eye, gave the impression of the ship being very foul Experience showed, however, that the arsenic had but little effect, and mercury and copper still remain pre-eminently the metallic poisons of the greatest value in anti-foulers. H Williams, Naval Constructor, U.S. Navy, has given the following composition for a paint adopted with great success for naval vessels—grain alcohol, 6 gallon, gum shellac, 13† lbs, turpentine, I gallon, pine-tar oil, I gallon, white zinc oxide, 13‡ lbs, Indian red, 13‡ lbs.; red oxide of mercury, 4‡ lbs. It was found that the cost of this paint was only one-third of the price charged for the proprietary paints

In some anti-fouling varnish paints the poisonous metals themselves have been carried in a very fine state of division, some paints containing very finely divided copper, piecipi-tated from the solution of a copper salt by non or zinc, whilst others contain metallic mercury in a comminuted condition, obtained by adding stannous chloride to the solution of a mercurous salt, these paints, however, show but little advantage over those containing salts of the

One of the most difficult portions of the ship to deal with is what is known as the 'boot top, that is, the portion between the heavy and light load line, as when a ship is discharging cargo the anti-fouling composition over this portion is exposed to the sun and rapidly perishes and peels off, whilst when the ship is again loaded, it is once more immersed, and the alternate action of sunshine and air and sea water perishes the paint at the very spot where it is most likely to be abraded by contact with lighters and barges, the result being that it is often necessary to touch up this portion of the vessel with paint during the voyage, and as good anti-fouling paints are rarely carried for this purpose, this portion of the ship is often painted with ordinary paint, and hence is liable to more damage than other parts of the vessel

**CORROSIVE SUBLIMATE** (Mercuric chloride) v Mercury

CORUBIN v ABRASIVES

**CORUNDUM.** (Coundon, Fr., Kound, Ger., from the Sanskit name korund.) Crystallised alumina ( $Al_2O_3$ ). The crystals belong to the thombohedral system, and are isomorphous with hematite (Fe $_2$ O $_3$ ) They vary considerably in habit, having the form of hexagonal prisms, pyramids, or tables. In some crystals the combination of six faces of the primary rhombo hedron with two faces of the basal plane pro duces a form strikingly similar to the regular octahedron There is no true cleavage, but often the crystals and crystalline masses show a pseudo-cleavage or parting parallel to the basal plane or to one or more faces of the primary thombohedron, due to lamellar twinning on these planes. On these surfaces the lustre is sometimes pearly, but otherwise it is usually vitreous, or inclining to adamantine in character The fracture is uneven to conchoidal; and the sharp jagged edges of the irregular grains give effect to the cutting power of the material when used as an abrading agent. Analyses show the presence of 95-99 p.c. of alumina, with small amounts of ferric oxide, magnesia, silica, water, &c, and sometimes traces of chromium. Sp.gr. 39–41. The crystals are optically unaxial and negative, refractive indices for sodium light  $\omega=17690$ ,  $\epsilon=17598$ . The dichroism is marked. Next to diamond, it is the hardest of minerals, being placed as No 9 in the scale of hardness, on this account it was

formerly known as adamantine spar

A distinction is made between the transparent 'precious corundum' and the dull, opaque 'common corundum,' while a third still less pure variety is known as emery (q v) This division also corresponds with the economic uses of the mineral, 'precious corundum' being used for gem-stones, and 'common colundum' and emery for abrasive purposes The pure, well-crystallised material is perfectly colourless and transparent with a high degree of brilliancy Traces of morganic impurities, however, produce a wide range of colours, which are more vivid in the stones of gemquality, namely red (ruby), orange and yellow ('oriental topaz'), green ('oriental emerald'), blue and indigo (sappline), and violet ('oriental Parti-coloured stones are not amethyst') The colour of common corundum uncommon is usually grey or brown

Corundum is not a widely distributed mineral, but it is found in abundance at certain localities It occurs in igneous and metamorphic rocks, such as granite, syenite, gneiss, peridotite, and serpentine, and in crystalline limestones Material of gem-quality is usually collected as water-worn pebbles in gravels derived from Material for gem-stones and for these rocks. abrasive purposes is not, as a rule, mined in the same localities. Taking first the localities for the gem-varieties, the following may be noted as a few of the most important. The best noted as a few of the most important rubies are obtained from the famous mines near Mogok in Upper Burma, where they occur in a white crystalline marble near its contact with gness, they are mined in this rock and also in its products of decomposition Darkcoloured tubies are obtained with sapphire from alluvial deposits in the provinces of Chantabun and Krat in Stam Stones of good quality have also been mined together with common corundum in Macon Co, North Carohna The best sapphires come from the gem-gravels of Ceylon, other localities of importance are Siam and Kashmir. Dark-blue and greenish-blue sapphires occur abundantly in gravels near Anakie in Queensland, while paler stones of various shades ('fancy stones') are plentiful in the neighbourhood of Helena in Montana

Common corundum as used for abiasive purposes (apart from emery, qv) was formerly mainly mined in the United States; later the Canadian deposits came into prominence; while now the latter are being ousted by the artificial products carborundum and alundum. India has also since remote times produced a certain amount which has been largely used by the native lapidaries. In the United States the corundum deposits extend along the Appalachian Range from Alabama to Massachusetts, reaching their greatest development in North Carolina and Georgia. Here they occur in peridotites near their contact with gneiss. In Canada a belt of corundum-bearing syemites

(nephelme and related alkalme syemites), associated with the Laurentian gnesses, extends for 100 miles across Renfrew, Hastings, and Halburton counties in Ontario, the principal mining district being near Craigmont in Renfrew Co Large rough crystals of corundum are abundant in gness in Madagascar and at Steinkopf in Namaqualand, South Africa More recently, important deposits have been discovered and worked in the Zoutpansberg and Pietersburg

districts in northern Transvaal

The corundum-bearing rock as mined is crushed between hardened steel rolls and the heavier corundum separated in water by means of jigs and concentrating tables. It is sifted into different grades of fineness and used (1) as loose grams, (2) mixed with glue or other soft cement as a coating on paper or cloth; and (3) as corundum-wheels for grinding (a) The last named are of three classes (a) The 'vitrified wheel,' in which the corundum is mixed with clay and baked at a high temperature, with the result that the giains of corundum are set in a matrix of porcelain (b) The 'chemical wheel, in which sodium silicate is the binding material (c) The 'cement wheel' with shellac, rubber, linseed-oil, &c, or magnesium oxychloride Another use for corundum, depending on its hardness, is for the pivot supports for watches (watch-jewels) and other delicate instruments, the poorer qualities of gem-material being used for this purpose Although corundum is the richest ore of aluminium, there are difficulties in its use for the extraction of the metal—It has, however, been used directly in the preparation of aluminium-copper and aluminium-iron alloys.

The artificial preparation of crystallised corundum was first effected by M A Gaudin, m 1837, by decomposing potash alum with charcoal It has since then been produced by a variety of methods, and is now manufactured on a large scale both for use as gem-stones and for abrasive purposes Beautiful crystals of ruby were prepared by E Frémy and Feil, in 1877, by heating a mixture of alumina and redlead with potassium dichromate in a porous fireclay crucible, a fusible aluminate of lead was thereby formed, which was decomposed by the silica of the crucible with the separation of crystallised alumina The crystals so obtained, though of some size, good colour, and perfectly transparent, were unfortunately too thin for cutting as gems A far greater measure of success was achieved by A Verneul in 1902, who has produced fine gems of all colours (v Gems, Artificial) Interesting experiments from a petiological point of view were conducted by J Moiozewicz in 1898, in which he obtained crystallised colundum in silicate magmas supersaturated with alumina Artificial corundum, used as an abrasive under the trade name of alundum (aloxite, adamite), is now manufactured on a large scale at Niagara Falls by fusing bauxite in an electric furnace 'Corubin' is also an artificial colundum, formed as a

by-product in the Goldschmidt thermite process

\*References\* — A E Barlow, Corundum, its
Occurrence, Distribution, Exploitation, and
Uses, Canada, Dept Mines, Geol. Survey, Mem
57, Ottawa, 1915, 378 pp (with special reference
to the Canadian deposits); T. H. Holland,

396 CORUNDUM.

Corundum, Manual of the Geology of India, 2nd edit, Calcutta, 1898 (with special reference to the Indian deposits), G. P. Merrill, The Nonmetallie Minerals, 2nd edit, New York, 1910; J. H. Pratt, Bull. U.S. Geol. Survey, 1906, No. 269; J. H. Pratt and J. V. Lewis, North Carolina Geol. Survey, 1905, 1; P. A. Wagner, South. African Journ. Industries, 1918, 1, 776, Trans. Geol. Soc. S. Africa, 1919, xxi. 37, M. Bauer, Edelsteinkunde, 2nd edit., Leipzig, 1909, and English translation by L. J. Spencer, London, 1904, Sapphire-Mining Industry of Anakie, Queensland, Bull. Imp. Inst., London, 1916, xiv. 253.

CORYBULINE, CORYCAVINE v. Corydaline

CORYDALINE AND THE CORYDALIS ALKALOIDS. These alkaloids occur in the loots of the Corydalis tuberosa (DC) (syn C cava, C bulloosa, Bullocapnus cavus), belonging to the Fumariacoæ, and also in some other species of this genus (C ambigua, C vernyi, C nobilis) Corydaline was discovered by Wackenroder, in the year 1826, and forty years later Wicke (Annalen, 1866, 137, 274) described the alkaloid more fully, assigning to it the formula C<sub>18</sub>H<sub>19</sub>O<sub>4</sub>N. An investigation by Additional (Inaug Dissert, Dorpat, 1890) indicated that C cava contained three or four different alkaloids, and the formula C<sub>20</sub>H<sub>27</sub>O<sub>4</sub>N was attributed to the one which is now regarded as corydaline

This base was more completely examined by Dobbie and Lauder (Chem Soc, Trans 1892, 61, 244, 605), and by Freund and Josephi (Annalen, 1893, 277, 1), who respectively ascribed to it the formula ('21H2904N and C22H2704N, the latter being probably the more correct formula. About the same time Freund and Josephi isolated three other alkaloids from ('cava roots, namely bulbocapnine, conycavine, and corybulbine, whilst Dobbie and Lauder described a fifth, conytuberine (l.e. 1893, 63, 485).

Several other corydalis alkaloids have since been isolated, the chief of which are dehydrocorydaline, isocorybulbine, corycavanine, corycavidine, and coryline Gadamei, Ziegenbein, and Wagner have classified the principal alkaloids into the following three groups —

		_	*
(1)	(Corydaline . Dehydrocorydaline Corybulbine .	:	$C_{22}H_{27}O_4N$ $C_{22}H_{23}O_4N$ $C_{21}H_{25}O_4N$
	(150Corybulbine	:	$\mathrm{C_{21}H_{25}O_{4}N}$
(2)	{Corycavine {Corycavamine	:	$C_{23}H_{23}O_{6}N$ $C_{21}H_{21}O_{5}N$
	(Corycavidine . (Bulbocapnine		$C_{22}H_{25}O_5N$ $C_{19}H_{19}O_4N$
(3)	Corytuberine Corydine	:	C <sub>19</sub> H <sub>21</sub> O <sub>4</sub> N C <sub>20</sub> H <sub>23</sub> O <sub>4</sub> N
	(isoCorydine .		CaHaOAN

The alkaloids in the first group are of weakly basic character, those in the third are strong bases, and the members of the second group are intermediate in this respect. All are oxidised by rodine, the first group yielding berbennelike compounds, except in the case of dehydrocorydaline, which is, itself, already such a compound. The third group all contain phenolic hydroxyl, and are therefore dissolved by alkalis

Protopine and two other alkaloids, one of which has the formula C<sub>20</sub>H<sub>17</sub>O<sub>4</sub>N, and is isomeric with berberine, have been found in

Chinese corydalis tubers (C. ambigua), as well as the first three members of Group I (Makoshi, Arch. Pharm. 1908, 246, 381). Eve more alkaloids, including glaucine, have been obtained from parts of C. tuberosa by Haars and by Gadamer (Arch. Pharm. 1905, 243, 154; and 1911, 249, 224).

To separate the chief corydalis alkaloids Gadamer, Ziegenbein, and Wagner extract the ground root with alcohol, distil off the solvent, dilute and acidify the residue with acetic acid. After filtering, the acid filtrate is purified by extraction with ether, then made ammoniacal, and the alkaloids extracted by means of ether. From the ethereal solution, after most of the solvent has been distilled off, a mixture of corydaline, bulbocapnine, corycavine, and corybulbine is deposited; these are separated, in the order given, by successive extractions with boiling alcohol More corydaline is obtained from the mother liquor on further concentration The residue evaporated to dryness is converted into hydrobromides and fractionally precipitated with ammonia, this treatment giving more corydaline, and also corybulbine, isocorybulbine, corycavamine, corycavine, corydine, and bulbocapnine, in the order named Corytuberine remains in the ammoniacal solution, and is obtained by extraction with chloroform after evaporating the solution to a syrupy consistency. To purify the crude corydaline it is converted into the hydrochloride and precipitated, together with some corycavine, from an aqueous solution by addition of alkali; any bulbocapnine present remains in solution, and can be recovered by passing carbon dioxide into the alkaline liquid. The corydaline and corycavine may then be separated by repeated crystallisation from absolute alcohol, corydaline, being the more soluble of the two, accumulates in the later fractions

Corydalis roots have been used in pharmacy, and are credited with antiperiodic properties. The alkaloids have a narcotic effect, but are not employed in medicine. The members of the three groups above-mentioned are closely related in physiological action as well as chemical constitution According to Peters (Arch expt. Path Pharm 1904, 51, 130) and Gadamer (Arch Pharm 1905, 243, 147), the alkaloids of the corydaline group produce paralysis of the spinal cord, and those of the corycavine group stimulate the motor centres, whilst the members of the bulbocapnine group give rise to increased All except corytuberine induce reflex action in flogs narcosis like that due to morphine, and have a prejudicial effect upon the action of the heart It has been suggested that bulbocapnine may be useful as a narcotic in veterinary practice, in the case of animals on which morphine has a stimulating effect.

Group 1. Corydaline  $C_{22}H_{27}O_4N$  crystallises from alcohol in hexagonal prisms, mp 134 5°,  $[\alpha]_D^{20}+317^\circ$  in chloroform or  $+310^\circ$  in alcohol. It dissolves easily in ether, chloroform, carbon disulphide, benzene, or warm alcohol, but is sparingly soluble in cold alcohol, and insoluble in water or alkalis. Corydaline and its solutions are quickly coloured yellow by the action of light or heat. It gives well-crystallised salts The aurichloride (B·HCl)<sub>2</sub>AuCl<sub>3</sub> crystallises from

dilute alcoholic hydrochloric acid as orange-coloured needles, mp 207°, the platinichloride (B·HCl)<sub>2</sub>PtCl<sub>4</sub> forms brown crystals, mp 227°; the hydrochloride B HCl,2H<sub>2</sub>O, columnar crystals, mp 206°–207°; and the ethyl sulphate B·C<sub>2</sub>H<sub>5</sub>HSO<sub>4</sub>H<sub>2</sub>O, large colourless prisms, mp 152 5°.

On exidation with iodine in alcohol corydaline yields dehydrocorydaline, which occurs with it in Corydalis tuberosa (Schmidt, Arch Pharm 1908, 246, 575) Four methoxyl groups are present in corydaline, and the base contains a tertiary nitrogen atom, since it reacts with methyl iodide to form corydaline methiodide B CH<sub>3</sub>L. Oxidised with permanganate, it gives eventually hemipine and meta-hemipine acids, showing the presence of two benzene nuclei in the molecule At the same time, another exidation product, corydaldine C<sub>11</sub>H<sub>13</sub>O<sub>3</sub>N, is formed; and from the constitution of this the presence of an isoquinoline nucleus in the original alkaloid is inferred

On oxidation of corydaline with dilute nitric acid instead of permanganate, dehydrocorydaline is first produced. One of the benzene nuclei is next destroyed, and a yellow dibasic acid, corydic acid C19H17O6N+1H2O, is formed When corydic acid is in turn oxidised with permanganate, it yields first an insoluble, colourless, tribasic acid  $C_{17}H_{15}O_8N$ , coryddic acid, and eventually metahemipinic acid, and a methyl pyridine tricarboxylic acid, which latter on further oxidation is converted into pyridine-2 3 4 6-tetracarboxylic C<sub>2</sub>H<sub>5</sub>O<sub>8</sub>N. The position of the methyl group in the tricarboxylic acid is thus indicated, and from this and the foregoing results the following constitutional formulæ for corydaline dehydrocorydaline are deduced (Dobbie and Lauder, Chem Soc Trans 1892, 61, 605, 1894, 65, 57; 1897, 71, 657; 1899, 75, 670, and 1902, 81, 145) -

This formulation indicates the close relationship shown by these alkaloids with tetrahydroberberine and berberine respectively. The former differs from corydaline only in having a dioxymethylene group substituted for the two methoxyl groups in ring 4, and a hydrogen atom for the methyl group in ring 2. Between the absorption spectra of corydaline and tetrahydroberberine there is also a very close relation; in fact, the only difference is that the general absorption shown by tetrahydroberberine is slightly greater than that of corydaline (Chem. Soc. Trans. 1903, 83, 598)

By reduction of dehydrocorydaline two optically inactive corydalines are produced, r-corydaline, mp 135°, and r-mesocorydaline, mp. 158°-159°, the two compounds being stereoisomeric. The former is obtained when an aqueous solution of dehydrocorydaline hydro-

chloride is treated with platinised zinc and dilute sulphuric acid at the temperature of the waterbath. Its sulphunic acid can readily be resolved by means of brucine into the corresponding l- and d-salts. r-Mesocorydaline is obtained together with its isomeride under milder conditions of reduction, by keeping an alcoholic solution of dehydrocorydaline hydrochloride with zine-dust and sulphuric acid at the ordinary temperature until it is decolourised. The separation of the two bases is easily effected by crystallisation from ether, the r-mesocorydaline alone separating (Gadamer and Klee, Arch Pharm 1916, 254, 295). By means of the camphorsulphonate it can be resolved into active components, but neither the l-form nor the d-form is identical with natural corydaline.

Dehydrocorydalme C<sub>22</sub>H<sub>25</sub>O<sub>4</sub>N occurs in the roots of C tuberosa, C ambigua, C solida, and C verim. From corydalme it is produced by the action of mild oxidising agents, such, as dilute intrie acid. It is a yellow crystalline powder, m.p. 112°-113° (deomp.), and gives crystalline salts, the aurichloride B HAuCl<sub>4</sub> forms red-brown needles, m.p. 219°. It gives a crystalline oxime, m.p. 165°, and, like berberine, unites with one molecule of chloroform, yielding a colourless crystalline compound, m.p. 154°. Haars ascribes the formula C<sub>22</sub>H<sub>25</sub>O<sub>5</sub>N to dehydrocorydaline, and regards the alkaloid as having a keto-structure when in the free state, but as being a quaternary base when in solution (Arch. Pharm. 1905, 243, 165)

Corybulbine was first separated by Freund and Josephi (lc) from commercial corydalme. and more fully investigated by Dobbie and Lauder, who assigned to 1t the formula C21 H25 O4N (Chem Soc Trans 1894, 65, 25) It is soluble in boiling alcohol, from which it separates, on cooling, as a fine crystalline powder, mp 238°-240° It is practically insoluble in water, sparingly soluble in ether and in methyl alcohol, readily in carbon disulphide, chloroform, and hot benzene The solution in chloroform is dextrorotatory,  $[a]_{D}^{20} + 303 \, 3^{\circ}$ . Corybulbine gives well-crystallised salts, the hydrochloride B HCl forms yellowish prisms, m p  $245^{\circ}-250^{\circ}$ oxidation with iodine in alcohol corybulbine gives dehydrocorybulbine C<sub>21</sub>H<sub>21</sub>O<sub>4</sub>N, m p 210°–211°, and on treatment with hydriodic acid furnishes 3 molecules of methyl iodide methylation with methyl iodide in the presence of potassium hydroxide, it yields corydaline, so that it differs from the latter only in containing an -OH group in place of an -OMe group, probably in the isoquinoline nucleus (Dobbie, Lauder, and Paliatseas, Chem Soc Trans 1901, 79, 87)

iso-Corybulbine C<sub>21</sub>H<sub>25</sub>O<sub>4</sub>N, first obtained by Gadamer, Ziegenbein, and Wagner, and subsequently examined by Bruns (Arch. Pharm 1902, 240, 19), crystallises from alcohol in voluminous leaflets, mp 179°-180°, [a]<sub>D</sub>+299 8° in chloroform, and on oxidation with iodine in alcohol yields dehydroisocorybulbine

## C21H21O4N

The latter resembles dehydrocorydaline and dehydrocorybulbine, and its hydriodide reduced with zine and sulphuric acid gives 1-180corybulbine, m.p. 165°-167°. Three methoxyl groups

are present in isocorybulbine (cf. Bruns, idem 1903, 241, 634).

Group 2. Covycavine C<sub>23</sub>H<sub>23</sub>O<sub>8</sub>N was first obtained by Freund and Josephi (Annalen, 1893, 277, 15). It is insoluble in water and in alkalis, sparingly soluble in cold, but more readily in hot absolute alcohol, from which it crystallises in rhombie tablets, mp 218°-219°; the alkaloid is optically mactive. The hydrochloride B HCl,H<sub>2</sub>O forms broad needles, mp 219°, the hydrodide B HI,H<sub>2</sub>O has mp 236°, and the aunichloride B HAuCl<sub>4</sub> has mp 178°-179° (decomp.) With methyl iodide it reacts as a tertiary base, the methiodide produced forming rhombic tablets, mp 218°. Corycavine contains a dioxymethylene group, but no—OH or —OMe group (Ziegenbein, Arch. Pharm. 1896, 234, 523, and Gaebel, idem. 1910, 248, 207)

Conyavamine  $C_{21}H_{21}O_5N$ , like corycavine, contains no methoxyl group, but probably a dioxymethylene group. It is a crystalline base, forming rhombic columns, in p. 149°, and has  $[a]_D^{20}+166°6°$  in chloroform; but on heating at 180°, or by warming with acetic anhydride, it may be converted into an optically inactive modification, in p. 213°–214°, resembling cryptopine (Gadamer, Ziegenbein, and Wagner, Arch Pharm. 1902, 240, 81)

Conjeaviding  $C_{22}H_{25}O_5N$  has been described by Gadamer (idem, 1911, 249, 30). The mixed amorphous alkaloids of C cava on conversion into thiocyanates can be separated into two groups, one giving crystalline thiocyanates, soluble with difficulty in alcohol, the other readily soluble, and amorphous From the second group corycavidine was isolated It is crystalline, and has m p.  $212^{\circ}-213^{\circ}$ ,  $[a]_{D}^{20}+2031^{\circ}$  in chloroform It appears to be corycavamine, in which a dioxymethylene group is replaced by two methyoxyl groups Lake corycavamine, it is converted on heating into the optically mactive form, mp 193°-195° after recrystallisation, the change occurs at 209° Corycavidine crystallises from hot chloroform with ICHCl3 on addition of alcohol It forms a crystalline hydrochloride and nitrate The aurichloride BHAuCl, is a red powder, which sinters at 85° and melts at 170° (decomp).

obtained by Freund and Josephi from the roots of Corydalis cava (Annalen, 1893, 277, 10) It separates from hot absolute alcohol in rhombic needles, mp 199°, [a]<sub>0</sub>+237·1° in chloroform, and yields crystalline salts. The hydrochloride B HCl forms needles, mp about 270°; the hydrodromide B HIs has mp 250°-270°; the hydrodromide B HI, mp 230°-270°, and the platinichloride, mp 200°-230° (decomp). The methicalide B CH<sub>3</sub>I forms shining needles, mp 257°.

Bulbocapnine is insoluble in water, but dissolves in alkalis, and is precipitated from its solutions in these by addition of carbon dioxide or ammonium chloride. Its four oxygen atoms are accounted for by the presence of a methoxyl, a hydroxyl, and a dioxymethylene group (Gadamer and Kuntze, Arch Pharm 1911, 249, 503, 598). Further, 'exhaustive methylation' gives rise to 3:4-dimethoxy-5:6-dioxymethylene-8-vinyl-phenanthrene, which brings

bulbocapnine into close relationship with apomorphine. The same authors have investigated the other members of this group, and assign the following formulæ to the four alkaloids concerned (ibid 641) —

Corntuberine C19H21O4N,5H2O was obtained by Dobbie and Lauder from commercial corydaline by extraction with hot water (Chem Soc Trans 1893, 63, 485) It crystallises from alcohol in fine, silky needles, mp 240°.  $[\alpha]_{\mathbf{D}}^{20}$  +282 6° in alcohol The alkaloid is nearly insoluble in cold water, ether, or chloroform, slightly soluble in benzene, and dissolves readily in hot water or in solutions of alkalis crystalline salts, including a methiodide methoxyl and two hydroxyl groups are present On methylation with methyl sulphate or diazomethane a mixture of two isometic monomethyl ethers is obtained, these correspond with corydine and isocorydine The dimethyl ether of conytuberine is glaucine ('orytuberine in alkaline solution is readily oxidised by air, giving a dark green liquid Its physiological action is very similar to that of anomorphine; and as with bulbocapnine and colydine, the characteristic emetic effect disappears when the two hydroxyl groups are methylated (Gadamer, Arch Pharm 1902, 240, 81, and 1911, 249, 503, 641)

Corydine C<sub>20</sub>H<sub>24</sub>O<sub>4</sub>N was first described by Merck, and later by Gadamer and his collaborators (idem 1902, 240, 81, 1911, 249, 669). It crystallises from alcohol with ½EtOH, mp 124°-125°, or 149° dry, [a]<sup>20</sup><sub>D</sub>+204 3° in chloroform. Three methoxyl groups and one hydroxyl group are present in the alkaloid. Oxidised with rodine, it yields the hydrodide of dehydrocorydine (1<sub>20</sub>H<sub>19</sub>O<sub>4</sub>N HI, which, on reduction, gives i-corydine, mp 165°-167°. From this l corydine is obtained by recrystallisation of the l-bartrate. As shown above, corydine is a monomethyl ether of corytuberine.

isoCorydine  $C_{20}H_{23}O_4N$  is obtained, as already shown, by methylating corytuberine. It crystallises in tablets, mp 185°, and has  $\lceil \alpha \rceil_D^{20} + 195 \ 3^\circ$  in chloroform. Its methiodide has m.p. 213°-214°, and  $\lceil \alpha \rceil_D^{20} + 143 \cdot 3^\circ$ .

CORYFIN. Trade name for ethyl glycollic

COTTON. 399

menthyl ester C<sub>10</sub>H<sub>19</sub>O CO CH<sub>2</sub> O C<sub>2</sub>H<sub>5</sub> used in | acetyl derivative of methody-hydroxy-phenylthe treatment of nasal catarrh

COSMOS REDS v Azo- colouring matters. COTARNINE v OPIUM

COTO BARK. True coto bark is obtained from Bolivia, the Amazons (Brazil), and Vene-It has been considered identical with Winter's bark [from Drimys Winters (Forst)], but its source is really unknown (Jobst and Hesse, Annalen, 282, 191) It comes into the market in long strips about 8 cm broad and 12 mm. thick, and is tough and haid. The colour is cinnamon brown, and a cross section shows numerous yellow spots It has an aromatic odour, and contains cotoin, dicotoin, hydrocotoin and methyl-hydrocotom, resins, tannin, The supply of the true coto bark is limited, so inferior banks, chiefly para-coto bank, are sold These contain little or no cotoin, but instead protocotoin and methyl-protocotoin

Cotoin or Benzoyl-phloroglucinol-mono-methyl ether C<sub>14</sub>H<sub>12</sub>O<sub>4</sub>, is found, together with various

related substances, in true coto back, but not in para coto bark, and has been the subject of much investigation by Chamician and Silber (Ber 24, 299 and 2977, 25, 1119, 26, 777, 27, 409 and 841; 28, 1549), Jobst (N Repert Pharm 1876, xxv 23, Ber 11, 1031), Jobst and Hesse (Ber 10, 249, Annalen, 199, 17), and Hesse (Ber 26, 2790, 27, 1182, Annalen, 282, 191). As a result its constitution has been made clear and a number of its derivatives described.

Cotom can be isolated from true coto bark by extracting the powdered bark with cold ether, distilling off the ether from the extract, and mixing the residue, whilst still hot, with light petroleum, whereupon a resinous-oily mass separates, from which the solution of cotoin can be decanted and the product obtained from it in the form of large yellow crystals. A further quantity can be obtained from the resinous mass mentioned above by boiling it with limewater, and treating the solution obtained with hydrochloric acid, when the cotom is precipitated Cotoin may be recrystallised from alcohol, or hot water, when it separates in yellow prisms, melting-point 130°-131°C. It is difficultly soluble in cold water, readily soluble in hot, is fairly soluble in alcohol, ether, and chloroform, but sparingly soluble in light petroleum or benzene It dissolves in alkalis forming yellow solutions from which it is reprecipitated on acidification As decomposition products of cotoïn, phloroglucinol as also benzoic acid have been obtained

Cotoin has been used pharmaceutically in cases of diarrhora and of phthisis

The diacetul derivative C<sub>18</sub>H<sub>18</sub>O<sub>6</sub>, produced by the action of acetic anhydride, or acetic anhydride and sodium acetate, on cotoin, consists of large prisms, melting-point 94° C. sodium acetate is used in the acetylation, the diacetyl derivative is accompanied by the

commann acctate, melting-point 142°C -

Benzoul cotom, obtained by the action of benzoyl chloride on cotoin, crystallises in prisms, melting-point 110°-112°C

Dibenzoyl-cotom consists of needles, meltingpoint 134°-135°C

Dimethyl-cotoin, obtained by the action of methyl iodide and alkali on cotoin, crystallises from alcohol, and has melting-point 138° C It yields a mono-acetyl derivative, colourless needles, melting-point 150° C, hence the formula

has been proposed for it by Ciamician and

Dibromcotoin C14H10O4B12, which crystallises in large colourless needles, melting-point 116° C, is prepared by the action of bromme on cotom in chloroform solution

Cotom oxime C14H13NO4 crystallises in plates, soluble in alcohol, difficultly soluble in water, and has been prepared by the action of hydroxylamine hydrochloride on a cold aqueous solution of cotom in sodium carbonate

Mono-nitroso-cotom consists of dark red leaflets, or orange-yellow needles (from acetic acid), melting-point 153°-154° C (both forms) The needle variety possibly contains acetic acid of crystalisation (Pollak, Monatsh 1901, 22, 996)

A dilute solution of cotoin in sodium carbonate is capable of coupling with diazonium compounds, and in this way Perkin and Martin (Chem Soc Trans. 1897, 1149) have prepared the following derivatives

Coloin-azo-benzene C14H11O4 C6H5N2 crystallises from acetic acid in orange-yellow needles, molting-point  $183^{\circ}-184^{\circ}$  C , it yields a diacetyl derivative,  $C_{14}H_{9}O_{4}$  ( $C_{2}H_{3}O)_{2}$   $C_{6}H_{5}N_{2}$ , when treated with acetic anhydride and sodium acetate, searlet needles, melting-point  $155^{\circ}$ - $156^{\circ}$ C, on treatment with hot alkalis the acetyl compound is hydrolysed with formation of the free azo compound

Cotom-azo-o-toluene, similarly prepared from diazotised o-toluidine and cotom, orange-yellow needles, melting-point 203°-204°C.

Cotom-azo-p-toluene consists of orange-yellow needles, melting-point 207°-208°C

Hydro-cotoin, which occurs together with cotoin, is a dimethyl ether of trihydroxy-benzoylbenzene, C<sub>6</sub>H<sub>5</sub> CO C<sub>6</sub>H<sub>2</sub>(OH)(OMe)

A number of poly-hydroxy benzophenone derivatives, including products found in coto bark and related to cotoin, have been prepared by W H Perkin and Robinson (Proc Chem Soc, 1906, 305). A G P.

COTTON consists of the long hairs which cover the seeds of various species of Gossymum, a genus of the nat ord Malvacew The botanical classification of cotton is in a confused state, and this is doubtless mainly due to the cultivation of the plant for extensive periods, the large 400 COTTON.

interchange of seed which has taken place between the different countries of production, and the consequent gradual modification of the distinctive characters of the original species.

American cottons.—The ordinary Upland cotton is fairly generally considered to be derived from Gossypium hirsutum (Linn) Some authorities, however, are of opinion that it is derived from G. herbaceum (Linn), whilst others regard these two species as identical Sea Island cotton has usually been regarded as G burbadense, but is considered by Watt (Wild and Cultivated Cotton Plants of the World) to be a hybrid form

Egyptian cottons—The Egyptian cottons probably cannot be referred to any particular species, but are mostly hybrid forms of G burbadense with other species. Most commonly, Egyptian cottons are regarded as G. barbadense, but Watt places them under G. peruvanum (Cav.), which some authorities consider as identical with the former

Induar cottons—With reference to these varieties, G. A. Gammie, of the College of Science, Poona, writes, in his monograph on The Indian Cottons. 'It is clearly evident that we have at the most only one true species of cotton in India, Gossiphum obtust folium, with its two sub-species, G. arboneum and G. herbaceum. All other forms should be treated as derivatives of these.' Other botianists, however, regard the Indian cottons as derived from various distinct species, including G. herbaceum, G. neglectum, G. Wightmanum, and G. arboneum.

Brazilian and Peruvian cottons—Brazilian cotton is generally considered to be the product of G brasiliense (Macf.), and rough Peruvian cotton that of G peruvianum (Cav.)

The cotton plant grows as a shrub or small It is of perennial habit, but, in cultivation, is usually treated as an annual certain districts of Brazil and Peru, however, arborescent forms are grown, which are allowed to remain in the ground for several years the flower-bud first opens, the petals are generally white or pale yellow, but they gradually become darker and redder until the third or fourth day, and then fall to the ground young fruit remains attached to the flower-stalk and enveloped in the calyx, it gradually increases in size, and is known as the 'boll.' When the boll is ripe, it dehisces by from three to five valves, exposing the cotton, which, now that the pressure on it is released, rapidly expands and forms a large fluffy mass As soon as the boll has opened completely, and is quite dry, the cotton is ready to be gathered. The seeds, with the cotton fibre adhering to them, form the product known as 'seed cotton'; this is collected from the ripe bolls, and the cotton (or 'lint'), after being torn from the seeds by the process of 'ginning,' constitutes the law Seed cotton usually concotton of commerce sists of about one-third of its weight of cotton fibre and two-thirds of seeds.

The cotton fibre is composed of a single hollow cell, and, while unripo, has the form of a long narrow tube. As the fibre matures, it becomes more or less flattened, and presents the appearance of a narrow, somewhat opaque ribbon or band, with slightly thickened, rounded edges. The ripe fibre possesses a peculiar characteristic twist. This twist appears to be

an acquired character; it is not present in wild cotton, and is increased by care in cultivation. The number of twists in a given length is very variable, and, as a rule, is, roughly speaking, inversely proportional to the diameter of the fibre, the twist being most numerous in the finest variety (Sea Island) Unripe cotton consists of thin, transparent fibres with little or no twist. Such fibres are known technically as 'dead cotton,' and, being very weak and brittle, they break up in the process of manufacture, thus increasing the amount of waste, and also tending to weaken the yarn or fabric into the composition of which they enter; they also possess the defect of not being able to take dyes satisfactorily.

According to the length of the individual fibres, cottons are classed in commerce as longmedium-, or short-stapled. The long-stapled cottons are 11 inches or more in length, the medium-stapled are about 1 inch long, and the short-stapled are less than I inch The principal long-stapled varieties are Sea Island cotton. Egyptian varieties, and the improved American Upland forms, which have been evolved from the ordinary Upland kinds by continuous selection and careful cultivation The Brazilian and Peruvian cottons vary from about 1 to 11 inches, and are, therefore, either medium- or long-stapled The chief medium-stapled cottons are the ordinary American Upland kinds, and the short-stapled class is represented by the Indian varieties

Sea Island cotton is grown in South Carolina, Georgia, Florida, and the West Indies The fibre is very long, fine, silky, and of a lustrous creamy tint, and this cotton is therefore the most valuable variety. It is used in commerce for spinning the finest and best yarns. Considerable care is required in the cultivation of this cotton, and the yield per acre is usually lower than that of other kinds.

American Upland cottons are produced in enormous quantities in the United States, and are the principal kinds used in the Lancashire spinning industry. They are of longer staple and better quality than the Indian varieties, and are employed for the manufacture of yarns of medium quality.

Egyptian cottons exist in several varieties, most of which yield a fibre of a brownish tint. These cottons are long, strong, and fine, and more lustrous than American Upland, but inferior to Sea Island cotton. Egyptian cotton is preferable to American Upland for the manufacture of goods requiring a soft, smooth finish and high lustre.

The Indian cottons are mostly short and coarse, and are consequently less valuable than American Upland kinds. The chief markets for these varieties are Japan and Germany, comparatively little being imported into the United Kingdom.

Brazilian cotton is harsh and wiry, of medium strength and of good length. One of the varieties grown in North Brazil is a tree cotton, which lives for 8 years or more. The ordinary annual cottons are more largely cultivated and are more productive than the arborescent form, but the fibre of the latter has the longer staple.

Peruvian cotton is chiefly of two kinds,

known respectively as 'rough' and 'smooth.' The former is the product of a cotton tree which is allowed to remain in the ground for from 4 to 6 years; the fibre is well known for its peculiarly harsh and crinkly character, which renders it especially useful to spinners for mixing with wool. 'Smooth' Peruvian is very similar to American Upland cotton, and is used for the same purposes.

The following table gives the approximate length and average diameter of the fibres of the principal commercial varieties. The diathe principal commercial varieties meters of the fibres of any one variety show great variation among themselves, but the figures given under this head are the average of a large number of determinations by various observers, and can, at least, be regarded as

relatively accurate .-

Variety	Usual length	Average diametei
Sea Island American Upland Egyptian Indian Brazilian Peruvian (rough) , (smooth)	 niches 1 6-2 2 0 8-1 1 1 3-1 6 0 6-1 0 1 1-1 4 1 2-1 5 0 9-1 1	mch 0 00064 0 00077 0 00068 0 00084 0 00080 0 00078

Cotton consists of about 83 7 p c of cellulose (qv.), together with 7 p c. of water, 0 6 p c of wax and oil, 1 5 p c. of protein, and about 1 65 pc of mineral matter (ash) (Bull 33, U.S. Dept Agric) The actual composition of The actual composition of

cotton varies with its origin

The wax is present as a thin layer on the surface of the fibre, and renders it incapable of absorbing water readily This substance, to which Schunck (Manchester, Mem Lit. Phil Soc. [III ] 4, 95) has given the name 'cotton wax, resembles other vegetable waxes, such as cerosin prepared from the leaves of the sugar-cane, and the waxes from the leaves of the carnauba palm [Copernicia cerifera (Mait)], and the Laphia palm [Raphia pedunculata (Beauv ) It is insoluble in water, but soluble in alcohol or ether; it melts at 86°, and solidifies again at 80°-82° Puro cotton wax is insoluble in solutions of caustic alkalı, ıt contains C 80 4, H 14 5, O 5 1 crated with this wax is a fatty acid, which was regarded by Schunck as margaric acid, but is more probably a mixture of steame and palmitic acids; it melts at 55.5°, and resolidifies at 50 5°.

The oil in the cotton fibre appears to be identical with that contained in the seed the preparation of the so-called absorbent cotton wool, the wax and oil are removed by boiling the fibre with dilute caustic alkali under pressure, the material is subsequently washed, treated with bleaching powder and hydrochloric acid, again washed, and dried.

The nitrogenous constituents of cotton consist of the remains of the cell protoplasm and

allied substances.

The mineral substances contained in the fibre are chiefly phosphates, chlorides, and sulphates of potassium, sodium, magnesium, and calcium, together with small quantities of iron

and traces of aluminium. If the amount of ash is much more than 2-3 pc, the excess is probably due to the presence of sand and other mechanical impurities.

When cotton is immersed in a strong solution of caustic soda, it undergoes certain remarkable changes, the chief of which is the production of a silky lustre. This phenomenon was discovered by John Mercer in 1844, and has led to results of great commercial importance. The process is now carried out on an extensive scale under the name of 'mercensation' (q v), and is sometimes applied to the yarn and sometimes to the woven The material is either kept in a state fabric of tension during the operation, or is allowed to shrink in the alkaline liquid, and afterwards stretched to its original length before being The process not only produces an washed exceedingly high lustre on the cotton, but also increases its power of absorbing dyes. The structure of the cotton fibre undergoes alteration, the flattened, twisted tube, characteristic of ordinary cotton, swells out, and is thus converted into a hollow, cylindrical form. This change is accompanied by a decrease in length and also by an increase in tensile strength, mercerised cotton being in some cases as much as 50 pc stronger than ordinary cotton Egyptian cotton generally mercerises better than ordinary American cotton

When cotton is treated with a mixture of nitiie and sulphuric acids, it is converted into nitro-cellulose or gun-cotton (v. Explosives). On dissolving gun-cotton in ether, or a mixture of ether and alcohol, the product known as collodion is obtained (v Photography) Nitiocellulose also enters into the composition of

celluloid (q v)

Among the more important works on this subject may be mentioned The Cotton Plant, Bull No 33, Office of Expt Stat US Dept. of Ague, Bowman's Structure of the Cotton Fibre; and Oppel's Die Baumwolle

COTTON BROWNS, SCARLETS, YELLOWS, v Azo- colouring matters, Primuline.

COTTON FLOWERS. Among the various portions of the cotton plant which have been industrially employed must be included the flowers which constitute one of the numerous Indian dyestuffs According to Watt (Dietionary of the Economic Products of India) they are thus used in the Manipur district Gossypetin was first isolated in small quantity from the flowers of the ordinary Indian cotton plant, G herbaccum (Perkin, Chem Soc Trans 1899, 75, 826), and has been more completely studied at a later period (ibid 1913, 103, 650) For its preparation a concentrated alcoholic extract of the flowers is treated with hot water and the mixture digested when boiling with addition of hydrochlone acid for three hours. After removal of tar by filtration the hot liquid on cooling deposits a brownish-yellow powder, which contains a mixture of quercetin and gossypetin These colouring matters are separated by a fractional crystallisation of their mixed acetyl derivatives from acetic anhydride, acetylgossypetin being in these circumstances the least soluble. The acetyl compound is finally hydrolysed by sulphuric acid in the presence of acetic acid in the ordinary manner.

Gossypetin C18H10O8 forms yellow needles,

melting-point 311°-313°, closely resembles quercetin in appearance, and is readily soluble in alcohol, but only very sparingly soluble in water. Concentrated alkaline solutions dissolve it, forming orange-red liquids which on agitation and dilution with water, become green, then blue, and finally assume a dull brown tint. Alcoholic lead acctate gives a deep red precipitate in the cold, passing into dull brown at the boiling-point, and with alcoholic ferric chloride a dull olivegreen liquid is obtained.

Gossypetin like quercetin forms compounds with mineral acids which crystallise in orangered needles, and with alcoholic potassium acetate the monopotassium derivative  $C_{15}H_{9}O_{8}K$  is produced. The hexa-acetyl derivative

C15H4O8(C2H3O)6

forms colourless needles, melting at 228°–230° By fusion with alkali or by dissolving the colouring matter in cold 50 p c aqueous potassium hydroxide and allowing the solution to remain exposed to air for twenty-four hours, with shaking, protocatchiae acid is obtained A phenolic substance also produced by these methods of hydrolysis has not yet been identified

When methylated with excess of methylic rodide and methyl alcoholic potash gossypctim hexanethyl ether C<sub>15</sub>H<sub>4</sub>O<sub>2</sub>(OMe)<sub>6</sub>, colourless needles, melting-point 170°–172°, is produced, and this by hydrolysis with alcoholic potash gives pretocatechia acid dimethyl ether and gossyptiot tetramethyl ether C<sub>12</sub>H<sub>16</sub>O<sub>6</sub>, needles, melting-point 115°–116° By ethylation in a similar manner, gossypctin hexacthyl ether C<sub>15</sub>H<sub>4</sub>O<sub>2</sub>(OEt)<sub>6</sub>, melting-point 144°–146°, can be prepared, and with alcoholic potash yields duthylprotocatechiae acid and gossyptiol tetracthyl ether C<sub>16</sub>H<sub>24</sub>O<sub>6</sub>, melting-point 110°–111° The latter forms an oxime C<sub>16</sub>H<sub>25</sub>O<sub>6</sub>N, melting-point 127°–129°, and when oxidised with permanganate gives gossypctonic acid, yellow needles, melting-point 154°–155° The constitutions assigned both to this latter (1) and to gossyptiol tetraethyl ether (2) are given below—

(1) HO C<sub>6</sub>H(OEt)<sub>3</sub>CO CO<sub>2</sub>H (2) HO C<sub>6</sub>H(OEt)<sub>3</sub>CO CH<sub>2</sub> OEt

and from these that of gossypetin naturally follows —

When an alkaline solution of gossypetin is exposed to air the blue liquid on acidification yields a reddish-brown precipitate which on gentle warming becomes crystalline. Again, if benzoquinone is added to a solution of gossypetin in cold absolute alcohol a similar reaction takes place. On keeping for a few minutes crystals commence to separate, and by then heating to 50° a semi-solid mass is obtained.

Gossyptone C<sub>15</sub>H<sub>8</sub>O<sub>8</sub>, the name assigned to this substance, consists of microscopic needles of a dull red colour, which are difficultly soluble in the usual solvents 1t dissolves in dilute alkalis with a pure blue colouration and its solution in concentrated sulphuric acid is dull brown. Sodium hydrogen sulphite solution reconverts it into gossypetim, Gossyptone possesses strong

dyeing properties, and gives the following shades on mordanted woollen cloth .—

Chromium. Aluminium. Tin
Dull brown. Orange-brown. Orange-red
Iron.
Deep olive.

These, it is interesting to note, are identical with those given in these circumstances by gossypetin itself, and it is accordingly evident that during the dyeing operation oxidation of the latter to gossypitone takes place Until a definite knowledge of the tetrahydroxybenzene nucleus in gossypetin has been obtained the position of the hydroxyl groups in this portion of the molecule can only be conjectured. Existing as it does side by side with quercetin it seems natural to consider that gossypetin is a Again, should gossypitone hydroxyquercetin be a p-quinone, the constitution of gossypetin will be the same as that which Neirenstein and Wheldale have suggested (Ber 1911, 48 3487) for the flavonol (1) which they obtained from quercetone, an oxidation product of quercetin, but the descriptions of the two compounds are not in agreement <sup>1</sup> On the assumption of the necessity for a quinol nucleus in gossypetin a second constitution (2) is however possible -

Again, a third expression (3), representing gossypetin as an hydroxyquercetin, is available, though in this case gossypitone must be regarded as an ortho-rather than a para-quinone:

The glucosides present in cotton flowers have been examined by Perkin (Chem Soc Trans. 1909, 95, 2181), who in the first place employed the Egyptian variety. The Egyptian cottons appear, according to the researches of Messrs Fletcher and Balls, to be derived from natural crosses of brown Peruvian cottons with the Sea Island variety.

A concentrated alcoholic extract of the flowers deposits on cooling a semi-crystalline powder, and this consists mainly of the potassium salts of two glucosides, quereimeritrin and gossypitrin, whereas a third glucoside iso-quercitin is present in the filtrate. A rough separation of the latter from this mixture may also be effected by means of aqueous lead acetate, for in these circumstances the precipitation of isoquercetin does

This compound, more recently synthesised by Neirenstein (Chem. Soc. Trans 1915, 107, 872), is described as melting at 854°-355°, and its hexamethyl ether at 146°-147°.

not occur, at least to any extent The isolation of quereimeritiin and especially of gossypitrin in a pure condition from these flowers is however of too tedious a character to be described here, and for details of the processes involved the original paper must be consulted

Quercimeritrin C<sub>21</sub>H<sub>20</sub>O<sub>12</sub>, 3H<sub>2</sub>O, melting-point 247°-249°, consists of small, glistening, bright yellow plates, insoluble in cold and fairly readily soluble in boiling water. Its alkaline solutions possess a deep yellow tint, with aqueous lead acetate it gives a bright red precipitate, and with ferric chloride an olivegreen colouration Octa-acetylquercimeritrin, needles,  $C_{21}H_{12}O_{10}(C_2H_3O)_8$ , melting-point 214°-216°, is sparingly soluble in alcohol, whereas monopotassium quercimeritiin, a yellow powder, can be obtained by means of alcoholic potassium acetate By hydrolysis with dilute sulphuric acid quereimentiin yields quercetin and glucose according to the following equation -

$$C_{21}H_{20}O_{12} + H_2O = C_{15}H_{10}O_7 + C_6H_{12}O_6$$

and is thus analogous to quercitrin which in this manner is conveited into quercetin and rham-

On wool mordanted with aluminium, tin, chromium, and iron, quereimeritim gives the following shades -

Aluminium. TuChromium Orange-yellow. Bright orange Reddish-brown. Iron Olive-brown

And these results are interesting, because with the exception of the iron mordanted pattern, which is of a rather browner character, the colours thus produced closely resemble those which are given by quercetin itself when dyed in a similar manner. They are widely different from those given by rutin and quercitiin, and mainly as a result of this property there can be little doubt that querementrin is to be represented by one of the two following formulæ .--

Querementin is also present in the flowers of the Prunus emarginata? (Finnemore, Pharm J. 1910, (iv), 31, 604).

Gossypitrin C21H20O13, only isolated with considerable difficulty is more readily prepared from the G neglectum (see below). When crystallised from dilute acetic acid it contains 2H<sub>2</sub>O of crystallisation, and is readily dissolved by hot water, but on boiling this solution yellow needles of the anhydrous substance, melting-point 240°-242°, separate When anhydrous it does not absorb water of crystallisation from moist air, but on boiling with acetone the hairlike needles are broken up into much finer needles, which when dried at 100° possess 1 molecule of water of crystallisation. Such a preparation melts at 200°-202°, and when dried

at 160° on standing in air now absorbs water. Preparations possessing this lower melting-point can be obtained in other ways, and it is accordingly evident that gossypitiin exists in two interchangeable modifications

Gossypitrin dissolves in alkalis with a pale yellow colouration, and gives with lead acetate a deep red precipitate It is almost insoluble in boiling acctone, a property which readily distinguishes it from quereimeritrin, which is fairly soluble in the latter solvent

Acetyl qossyptrin  $C_{21}H_{11}O_{14}(C_2H_3O)_9$ , colourless needles, melting-point  $226^\circ$ – $228^\circ$ , is almost insoluble in alcohol

When hydrolysed with dilute sulphune acid gossypitim yields gossypetin and dextrose according to the equation-

$$C_{21}H_{20}O_{13}+H_{2}O=C_{15}H_{10}O_{8}+C_{6}H_{12}O_{6}$$

The shades given by this glucoside on moidanted wool are as follows .-

Chromium Aluminium  $T'\iota n$ Reddish-Dull Bright Dark brown yellow orange olive; brown.

Gossypitrin reacts, like gossypetin itself, with benzoquinone, and yields in this way a quinone to which the name Gossypitrone C21H18O13, has been assigned This consists of maroon-coloured needles, which possess no definite melting-point, although fusion of the product occurs about 255°-259° By the action of warm dilute sulphurous acid solution it is reconverted into gossypitin, and the same change appears to occur in the dyeing process, for the shades produced are identical with those yielded by this latter glucoside—It is considered probable that the sugar group of gossypitim is attached to its tetrahydroxybenzene nucleus, though until the exact nature of this has been decided, its position is necessarily uncertain

Isoquercitrin  $C_{21}H_{20}O_{12}$ ,  $2H_2O$ , crystallises from dilute alcohol in pale yellow needles, melting at 217°-219° It is sparingly soluble in water, and dissolves in alkaline solutions with a deep yellow tint, but its most interesting property is the fact that with aqueous lead acctate solution it gives a bright yellow precipitate entirely distinct from the deep red deposit which is produced in this manner from

the isomeric quereimeritrin

Again, though more readily susceptible to hydrolysis than the latter glucoside, it yields identical products —

$$C_{21}H_{20}O_{12}+H_{2}O=C_{15}H_{10}O_{7}+C_{6}H_{12}O_{6}$$

Dyeing experiments with isoquercitiin give hades entirely distinct from those given by quereimeritrin, and these, although slightly paler, resemble those yielded by quercitiin.

Aluminium. Chromium Tin. Iron. Golden-Lemon-Brownish-Brownishyellow yellow. yellow. olive.

The properties of this substance indicate that its sugar group is not attached as in querementrin to the phloroglucinol nucleus of quercetin Indeed, it is probably constituted similarly to quercitrin, which, however, contains a rhamnose in the place of the glucose residue Three formulæ are possible for iso-quercitrin, which may be briefly expressed by the statement that the position of the sugar residue in respect to the quercetin group is at | one or other of the points in the following which are marked with an asterisk :-

The aqueous extract of the Egyptian cotton flowers employed in this investigation gave by hydrolysis with acid 186 p.c. of crude colouring matter, and in this approximately 10 pc of gossypetin was present. Dyeing experiments with the flowers in the usual way gave the following shades .-

Chromium Aluminium  $T_{2n}$ Iron. Reddish-Olive-Green-Orangebrown brown. brown. yellow

and these though duller were somewhat similar in character to those given by quereimeritrin In comparison with the shades similarly produced from other natural dyes, they most nearly resemble those of the so-called 'Patent Bark,' a preparation of quercitron bark in which quercetin and no quercitrin is present.

Among the types of cotton flowers there are (a) red, (b) pink, (c) yellow, and (d) white flowered plants. In the offspring of a cross between (a) and (c) there occurs in the second and subsequent generations red and yellow plants which breed pure, whereas in the off-spring of a cross between (a) and (d) all four forms occur which breed pure As a supplementary investigation to that of the Egyptian flowers the petals derived from such pure plants occurring among the offspring of one or other of these crosses have been examined (Perkin Chem. Soc. Trans 1916, 109, 145) (cp. Leake, Proc. Roy Soc 1911, (B), 83, 147)

The types were as follows red flowered, G. arboreum (Linn); pink, G sanguineum (Harsk), yellow and white, two varieties of G neglectum (Tod), usually now treated as one species but originally described as G neglectum and GAs a result it has been found that the red flowers of G arboreum contain isoquercitrin, quereimeritrin and gossypitrin in this case being absent, whereas in the yellow flowers of G neglectum, gossypitrin and isoquercitrin were present, and querementrin appeared to be absent On the other hand, the white flowers of G neglectum and the pink flowers of G sanguineum gave but traces of colouring matter too small for complete identification, though the respective products obtained resembled in their properties apigenin and quercetin An examınation of the ordinary Indian cotton flower, G. herbaceum, available only in small amount, gave the same results as the G neglectum

Gossypetin is also present in the flowers of the Hibiscus sabdariffa or 'red sorrel' of the West Indies, a small shrub which is widely cultivated throughout the hotter parts of India and Ceylon (Perkin, Chem. Soc Trans 1909, 95, 1855) The stems yield the 'Rozelle hemp' of commerce, and this is obtained by tetting the twigs as soon as the plant is in flower. The yellow flowers are just capable of dyeing yellow, but are not used at all in India for this purpose; on the other hand, the red calyces are employed | ing,' which may also occur during the voyage);

for dyeing in an obscure degree in two remote parts of the country (Burkill, Agricultural Ledger, Calcutta, 1908, No. 2, 13).

COTTON SEED OIL is obtained on a large scale from the seeds of the several species of the cotton tree, Gossypium. In the United States, Gossypium hirsutum (Linn), 'Upland cotton,' covered with 'linters,' is extensively cultivated, whereas the plant grown in the Sea Islands and The Indian in Egypt is G barbadense (Linn) variety of cotton seed is generally assumed to be G herbaceum and G arboreum, whilst the South American cotton plants are frequently classified as a separate species, G acuminatum or peruvianum Formerly the seed not required for planting was mostly wasted, as the oil contained therein could not be refined Cotton seed oil was first expressed in this country in the year 1847. Since then the industry has assumed such enormous dimensions that the annual import of cotton seed to this country from Egypt, the East Indies, and South America amounts to more than 600,000 tons, in addition very con siderable quantities of cotton seed oil are imported from the United States, which produce almost 200,000,000 gallons of oil per annum, of which approximately 50,000,000 gallons are exported to Europe

The seed contains, on the average .-

			1	er cen
Oıl .				200
Water				85
Proteins				20 0
Carbohy	drates			32 0
Crude fit	ore			150
Ash				45

The proportion of oil in the seed varies considerably with the origin of the seed. Thus Egyptian seed contains up to 24 pc of oil, East Indian seed on the average only 18 pc, American 'Uplands' and Jamaica seed 23-24 pc; and Brazilian and Peruvian no more than 20-21 p c of oil The seeds consist on an average of 60 p c of kernel and 40 p c of husk.

A distinction must be made between seed which must be decorticated before crushing, owing to the fine 'linter,' which cannot be removed readily, and seed which is crushed in the undecorticated state Undecorticated seeds are chiefly crushed in England, France, and Germany, from Egyptian, Bombay, and also Smyrna seeds, whereas the vast majority of the American 'Upland seed' must be decorticated before being crushed In some European establishments Bombay seed is decorticated before crushing.

The kernels ('meats') are permeated with cells containing a deep-brown colouring matter; hence the clude cotton seed oil running from the hydraulic presses has a ruby-red to almost black colour The depth of the colour of crude cotton oil depends, in the first instance, on the state of freshness of the seed The American seed, which is, as a rule, crushed shortly after having been harvested, yields a crude oil of a much paler colour than that of crude oil obtained ın England, France, and Germany from Egyptıan seed The palest Egyptian crude oil is obtained When the from fresh seed arriving in October. seed is stored it undergoes deterioration ('heathence the crude oil obtained from Egyptian seed in July and August has a very dark colour.

In the American trade three grades of crude cotton seed oil are recognised, viz. 'prime,' 'choice,' and 'off' qualities The crude oil is refined by treatment with a dilute caustic soda solution, at a temperature not exceeding 120° F, whereby the free fatty acids are neutralised; the soap so formed falls to the bottom and carries with it the brown colouring matter, so that the supernatant oil is of a bright-yellow colour. The oil is then washed free from soap, or, in the case of a technical oil, is treated with bleaching powder, to yield the bright-yellow oil of commerce. The first qualities of cotton seed oil which are not bleached with chemicals are used as edible oils, and are consumed in enormous quantities either as such, as a cooking oil, for tinning sardines, table oil, &c. or in the manufacture of lard substitutes, under such names as salad oil, sweet nut oil, butter oil The last-named product is chiefly used in the manufacture of margarine and compound lard

Official methods for analysing cotton seed oil, and tests for determining its behaviour when refined or bleached, have been published by the Interstate Cotton Crushers' Association, U.S.A. (Oil, Paint and Drug Rep., 1913, June 30; 1914,

Cotton seed oil is, at the ordinary temperature, a pale-yellow oil of characteristic, but not unpleasant, taste. At about 12° 'stearine' deposits This 'stearine' would interfere with the use of cotton seed oil as a table oil or salad oil; therefore the 'stearine' is removed on a large scale by cooling the oil to a low temperature, and filtering off the deposit. The filtered oil is known in commerce as 'winter oil' ('demargarinated cotton seed oil'). According to the requirements of the market, it will not 'cloud' at 40° F. or 32° F, or even at 28° F. The sp gr. of cotton seed oil is 0 923–0 925 at 60° F. The saponification value varies from 190 to 193, and its iodine value from 105 to 115.

The solid fatty acids in cotton seed oil consist chiefly of palmitic acid, small quantities of arachidic acid seem to be present, but stearic acid appears to be absent. The liquid fatty acids would seem to consist of oleic and linolic acids only.

Cotton seed oil eminently typifics a semi-drying oil. In the Livache test (see Oils and Fats) it absorbs 5 9 p.c. of oxygen in 24 hours. By blowing air through cotton seed oil at a temperature of about 90°-100°, oxygen is absorbed, with the formation of a small amount of oxidised acids, the acetyl value of the oil rising at the same time considerably. This reaction is used on a large scale in the preparation of 'Blown Cotton Seed Oil' (see Oils and Fats), which is used in considerable quantities is a lubricating oil in admixture with mineral pils, to produce the 'Marine Lubricating Oils' of commerce.

Cotton seed oil hardened by hydrogenation s used as a solid fat in the manufacture of margarine A typical commercial hardened oil melts at about 38°, and has a saponification value of about 196 and iodine value of about 70.

A reliable test for the identification of cotton eed oil, and for the detection of it in other oils ind fats, down to even 1 p.c. (under favour-

able circumstances), is the Halphen colour test, which is carried out in the following manner: 1-3 cc. of the oil is dissolved in an equal volume of amyl alcohol; to this is added 1-3 ce of carbon disulphide, holding in solution 1 pc. of sulphur flowers The test-tube is immersed in boiling water, or, better, in boiling brine, the carbon disulphide being allowed to evaporate off In the course of 5-20 minutes, a deep-red colouration appears, the intensity of which stands in direct proportion to the amount of cotton seed oil in the sample. The reagent which produces the colouration appears to be an impurity in the amyl alcohol, and the test may be rendered more delicate by substituting pyridine for amyl alcohol. In this way as little as 0 25 pc of cotton seed oil may be detected (Gastaldi, Giorn. Farm Chim. 1912, 61, 289) Although the test cannot be used as a quantitative one, yet it must be considered, if positive, as one of the best means of recognising the presence of cotton seed oil qualitatively. A negative Halphen test is, however, no conclusive proof of the absence of cotton seed oil in a mixture, as the colouring matter is destroyed by heating cotton seed oil to 250°, or by keeping it at 200° for some prolonged time, or by 'blowing' it. Hence 'blown cotton seed oils' do not show the Halphen colour reaction.

**COTTON SEED, COLOURING MATTER OF.** Cotton seeds contain, in addition to cotton-seed oil, a phenolic substance gossypol, which remains dissolved in the oil when this is expressed

By extracting the oil with caustic soda and neutralising the solution thus obtained a voluminous precipitate separates which consists of the colouring matter admixed with fatty acids, and the products of its own oxidation. For the isolation of gossypol Marchlewski (J. pr. Chem 1899, 60, 84) employed a partially purified preparation of this character from which the main bulk of the fatty acids had been eliminated. By extraction with ether, repeated crystallisation from glacial acetic acid, and from a mixture of alcohol and 50, c. acetic acid, it was isolated in the form of small yellow prisms.

As the result of analyses of this substance dried in vacuo over sulphuric acid, Marchlewski suggested two formulæ,  $C_{13}H_{14}O_4$  and  $C_{32}H_{34}O_{10}$ , and of these the first is considered preferable. Preparations dried at higher temperatures appeared to suffer alteration as they thus became darker, and had then a lower melting-point.

Thus the gossypol when quickly heated melted , but when previously dried at 125°-130° at 179°-180°. It is easily soluble in the usual organic solvents, and dissolves in sulphuric acid with a cherry-red colouration, a reaction by which its presence in cotton-seed oil can be detected In alkalis, gossypol dissolves with a yellow colouration which rapidly changes to yiolet, and then slowly fades. The ready violet, and then slowly fades susceptibility of alkaline gossypol solutions to air oxidation accounts largely for the impure nature of the crude colouring matter isolated by such reagents from the oil, and it appears that if care is taken in the process, it is possible to obtain a much cleaner material in the first instance. An alcoholic solution of gossypol gives with ferric chloride a dark-green colouration, and with the neutral and basic acetates of lead a deep-yellow precipitate. The analysis of the

lead salt corresponded to C<sub>13</sub>H<sub>12</sub>O<sub>4</sub>Pb. According to Marchlewski, gossypol yields amorphous acetyl and benzoyl derivatives and appears to possess two hydroxyl groups. It contains no methoxy groups, and is not a glucoside dyes iron mordanted material a grey shade, but the product obtained by a careful oxidation of its alkaline solution possesses, it is stated, tinctorial properties of greater importance. Gossypol can be employed on cotton fabrics as a mordant for the basic colouring matters

F E Carruth (Amer Chem J 1918, xi 4, 647) isolates gossypol from the decorticated cotton-seed kernels which have been crushed by rollers before they are 'cooked' in the oil mill From these a considerable amount of oil is removed by percolation with ligroin, and the residue is then extracted with other and the extract evaporated The product is now treated with one-half to one-third its volume of glacial acetic acid, and on keeping for some days, or more quickly by gentle warming, crystals of gossypol acetate separate purification it is dissolved in ether, glacial acetic acid is added, and the solution evaporated until crystallisation commences

To obtain gossypol itself an ethereal solution of the acetate is treated with water, and the ether distilled off. The gossypol thus remains suspended in the water as crystalline crusts, and may be recrystallised from alcohol

A second interesting method consists in adding aniline to an ethereal extract of cotton seed. The solution after warming on the waterbath gradually deposits on standing an orangeyellow microrystalline precipitate, which apparently consists of an aniline salt of gossypol,  $C_{30}H_{29}O_{9}$ ,  $2C_{6}H_{5}NH_{2}$  or  $2C_{30}H_{29}O_{9}$ ,  $5C_{6}H_{5}NH_{2}$ The yield is practically quantitative, and the process may be used to estimate gossypol in extracts of cotton-seed products

To regenerate gossypol from the aniline compound, it is dissolved in hot alcoholic potassium hydroxide, and the aniline removed by steam distillation From the alkaline liquid the gossypol is precipitated on acidification, and may be purified by conversion into the acetate according to the method given above

Gossypol acetate, a molecular compound of gossypol and acetic acid is evidently the substance described by Marchlewski as the free colouring matter It is a comparatively stable substance, and when heated does not evolve acetic acid readily below 180° Its formula determined by alkali titration is

$$C_{30}H_{28}O_9$$
,  $C_2H_4O_2$  or  $C_{30}H_{30}O_9$ ,  $C_2H_4O_2$ 

Hence gossypol itself is  $C_{30}H_{28}O_9$  or  $C_{30}H_{30}O_9$ , and this result is in harmony with other analytical

figures given in Carruth's paper

Gossipol acetonate, C<sub>30</sub>H<sub>28</sub>O<sub>9</sub> (H<sub>3</sub> CO CH<sub>3</sub>, is obtained as glistening crystals when gossypol is dissolved in acctone, and the solution concentrated Like the acetate it is a stable compound, and when heated at 140°-150° it alters but little in weight

Acetyl and benzovl derivatives of gossypol could not be obtained in a crystalline condition, but as a result of their examination the evidence was sufficient to indicate that four or five hydroxyl groups are present in gossypol. When By the interaction of coumarane and benzoyl gossypol is heated to 186°-190° it fuses with the chloride in the presence of aluminium chloride

evolution of vapour, and then solidifies to a black mass On treating the residue with ether a sparingly soluble yellow crystalline compound B. gossypol separates, which is much less toxic than gossypol and melts at about 246°-248°. Again, by fusion with alkali, a second new crystalline substance C. gossypol can be obtained, and this, which is soluble in alkalis with a blue colour is probably closely related to the purple oxidation product of gossypol referred to above

('otton-seed meal which has been submitted to the 'cooking process' loses much of its toxic property, and contains then practically no gossypol By extracting this product with aniline, the aniline compound of D gossypol which crystallises in orange-yellow prisms can be isolated

Alcoholic potash converts this compound into D quesypol itself, yellow crystals, which

darken and soften at about 256°.

From the cooked meal 1 2 p c of the anilmo compound can be obtained These products have not as yet been closely examined.

COUCH GRASS or Triticum, Agropyrum BP. The dried rhizome of Agropyron repens (Beauv) Is frequently adulterated with Cynodon Daetylon or dog grass A common weed in pasture and arable land Used in pharmacy Therapeutic arable land Used in pharmacy value obscure

COUMARANE GROUP Leuco-benzoyl commarane-

according to v Kostanecki and his co-workers is probably the mother substance of the catechin present in Gambier catechu, though the correctness of this view has not yet been fully established Coumarane or hydro-coumarone, the anhydride of o-hydroxyphenyl ethyl mner alcohol--

$$\mathrm{C_6H_4} \underbrace{\mathrm{CH_2\,CH_2\cdot OH}}_{\mathrm{CH_2\,CH_2\cdot OH}}$$

was first prepared by Alexander (Bei 1892, 25, 2409) by reducing coumarone in alcoholic solution with sodium—

$$\begin{array}{ccc} C_{0}H_{4} & O \\ CH & CH_{2} & CH_{2} \\ \hline Coumarone & Coumarane \end{array}$$

It forms a colourless oil, boiling-point 188°-189°, and gives in sulphuric acid solution by addition of ferrie chloride a violet colouration Stoermer and Gohl (Ber 1903, 36, 2873) have obtained coumarane and its homologues by the action of sodium on o-bromo-phenol brom-ethyl esters according to the following equation -

4 benzoyl-coumarane, colourless crystals, meltingpoint 44°-

is produced (v. Kostanecki, Lampe and Marschalk, Ber 1907, 46, 3660), and when reduced in alcoholic solution, the leuco-benzoyl-coumarane above referred to is obtained

COUMARANDIONE, COUMARANONE v.

KETONES.

COUMARANONE GROUP. Though colouring matters derived from coumaranone (ketocumaran) have not as yet been isolated from natural sources, there is some probability of

their presence in certain flowers, and they are not only historically connected with the earlier attempts to synthesise the flavones, but have in certain instances been successfully converted into flavonols.

Coumaranone (1) is obtained (Friedlander and Neudorfer, Ber 1897, 30, 1077) by boiling acetyl o-hydroxyacetophenone bromide (2) with an aqueous suspension of chalk, or by the action of dehydrating agents on phenoxyacetic acid (3)—

(2) 
$$\begin{array}{c} -0 - \text{COCH}_3 \\ -\text{CO CH}_2\text{Br} \end{array} \rightarrow (1) \begin{array}{c} 0 \\ \text{CO} \end{array}$$

$$\begin{array}{c} \text{CH}_2 \\ \text{HOOC} \end{array}$$

(Friedlander, Ber 1899, 32, 1867), and consists of colourless needles, melting-point 97° condenses with benzaldehyde in the presence of hydrochlone acid with formation of benzylidenecoumaranone-

$$C_{\rm CO}$$
 C=CH  $C_6H_5$ 

which was at first considered by Friedlander and Neudorfer (Ber 1897, 30, 1077) to consist of flavone On substituting protocatechnic aldehyde for benzaldehyde these authors obtained the dihydroxy compound-

and this is a strong dyestuff which gives on aluminium mordanted fabrics an orange, and with iron and chromium mordants brown-coloured shades.

The first colouring matter of this group was, however, prepared in 1896 by Friedlander and Rudt (Ber. 29, 878) from chlorogallacetophenone and benzaldehyde-

$$\begin{array}{c} \text{OH} & \text{OH} \\ \text{OH} & \text{OH} \\ -\text{CO} - \text{CH}_2 \text{CI} \end{array} \Rightarrow \begin{array}{c} \text{OH} & \text{O} \\ \text{CO} \end{array} \begin{array}{c} \text{CH}_2 \end{array}$$

$$\rightarrow \begin{array}{c} \text{OH} & \text{O} \\ \text{OO} & \text{C.CH} \\ \end{array}$$

and from the analogous resacctophenone derivative other hydroxy compounds of this group were obtained later. These are of an orange or orange-red colour, as a rule form oxonium salts, and those possessing tinctorial property dye alum mordant a much more orange shade than the flavone colouring matters. There is indeed a close resemblance in this latter respect between these compounds and the colouring matters derived from chalkone-

$$\bigcirc -\text{CO}: \text{CH-CH-C}_6\text{H}_5$$

(Perkin, Chem Soc Trans 1904, 85, 1465), for butem (a tetrahydroxy chalkone obtained from the Bulea frondown) and the 3 4 3' 4' tetrahydroxybenzylidene coumaranone of Friedlander and Rudt give almost identical shades The benzylidene coumaranones dissolve in sulphuric acid with the production of a red or redviolet colouration, and can thus again be distinguished from flavones which in this manner give either colourless or pale yellow solutions

In 1896 Kesselkaul and v Kostanecki (Ber 29, 1886) suggested that judging by its colour reactions the colouring matters of Friedlander and Rudt were not, as these authors suggested, flavone but rather coumaranone derivatives, and the true constitution of these substances became evident on the synthesis of flavone itself by Feuerstein and v. Kostanecki (l.c.) That benzylidenecoumaranones can be converted into flavanol derivatives has been shown by Auwers and Muller (Ber 1908, 41, 4203), and this transformation is discussed under Flavonol.

Coumaranone is also interesting as the source of ox-indigo-

the oxygen analogue of indigotin. This crystallises in yellow prisms, but does not possess any tinctorial value

Analogous colouring matters can be obtained from diketohydrindene (indandione), this also possessing the reactive methylene group, with aldehydes. Thus by heating this compound with protocatechnic aldehyde at 110°-120° v Kostanecki prepared the dihydroxybenzylidene denvative-

which consists of brownish-yellow needles, melting-point 257°, soluble in alkaline solutions with a red-violet colour and gives yellow shades on aluminium mordanted fabrics

From indoxyl also in a similar way Perkin and Thomas (Chem Soc Trans 1909, 95, 798) prepared dihydroxybenzaldehydeindogenide-

crystallising in orange-red needles, and producing with aluminium mordanted fabrics dull scarlet shades

Chalkone or benzylideneacetophenone is readily prepared by the interaction of acetophenone and benzaldehyde (Claisen, Ber 20, 257)—

$$\begin{array}{c} \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}} & \mathrm{CO} & \mathrm{CH}_{3} + \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}} & \mathrm{COH} \\ \longrightarrow & \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}} & \mathrm{CO} & \mathrm{CH}_{2}\mathrm{CHOH} \cdot \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}} \\ \longrightarrow & \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}} & \mathrm{CO} & \mathrm{CH} & \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}} + \mathrm{H}_{2}\mathrm{O} \end{array}$$

It consists of yellow rhombic prisms, melting-point 57°-58°.

When submitted to hydrolysis it is reconverted into the aldehyde and ketone, a reaction which is the reversal of that given above—

which is the reversal of that given above—
$$C_6H_5 CO CH CH C_6H_5 \\ \rightarrow C_6H_5 CO CH_3 + C_6H_5 CH(OH)_2 \\ \rightarrow C_6H_5 CO CH_3 + C_6H_5 CH(OH)_2$$

$$C_6H_5 CO CH_3 + C_6H_5 CH(OH)_2$$

By the employment of hydroxyacetophenones and hydroxybenzaldehydes, in general previously methylated, numerous chalkone derivatives have been obtained by v. Kostanecki and his co-workers. Only one natural colouring matter of this group, known as butein, has at present been isolated, and the synthetically prepared compounds have been more specially interesting owing to their employment for the synthesis of the flavonol group of colouring matters.

of the flavonol group of colouring matters

Ryan and O'Neill by the aid of diacetoresorcinol have obtained dichalkone compounds.

Thus diacetoresorcinol dimethyl ether and
benzaldehyde form dibenzylidendiacetoresorcinol dimethyl ether, and this by the action of
aluminum chloride is demethylated with production of the free hydroxy compound—

It was obtained as yellow crystals melting at  $196^{\circ}$ – $198^{\circ}$ 

Interesting is the fact shown by these authors that by a variation in its method of preparation four distinct varieties of this dichalkone can be produced. Whereas three of these designated as  $\alpha$ ,  $\beta$  and  $\delta$  are stereo-isomers, the fourth or  $\gamma$  variety is structurally related to the other three

When 2-hydroxychalkone is digested at the boiling temperature with alcoholic sulphuric acid, flatanone (dihydroflavone) is produced according to the following scheme:—

and this is the method generally adopted for the synthesis of flavanone derivatives. On the other hand, though as a rule a chalkone only is produced by the interaction of aldehyde and

hydroxyketone, when gallacetophenone dimethyl ether (Woker, v Kostanecki, and Tambor, Ber 1903, 36, 4235) or quinacetophenone monomethyl ether (v. Kostanecki and Lampe, ibid 1904, 37, 773) are condensed with benzaldehyde in the presence of sodium hydroxide the respective flavanones (1 and 2) are thus directly obtained—

On hydrolysis the flavanones yield chalkones, a reversal of the equation given above. The reaction in these cases is not undirectional, and the conversion of flavanone into chalkone, or chalkone into flavanone, is never complete.

Chalkones which contain hydroxyls in the ortho position to one another, in the position 3′ 4′ are powerful colouring matters, whereas flavanones possessing similar hydroxyls are not dyestuffs, as indeed their constitution indicates. During the dyeing operation, employing mordanted woollen material hydrolysis occurs, with subsequent chalkone production, and the shades produced by them are in reality those derived from the latter. Flavanone crystallises in small colourless needles, melting at 75°-76°.

Biomflavanone (1) by elimination of hydrobromic acid readily yields flavone (2)—

and, again, flavanones may be converted into flavonols.

A G P.

COUMARIN 
$$C_0H_6O_2=C_6H_4$$

$$CH \cdot CH$$

$$O = CO$$

$$(1, 2)$$

$$CH \cdot CH$$

$$O = CO$$

was obtained by Vogel (Gilb Ann 64, 161), in 1820, from the tonka bean (the seed of Dipteryx (Coumarouma) odorata, Willd), or from the melilot (Melilous officinalis, Lam), and from the woodruff (Asperula odorata, Linn), but was mistaken by him for benzoic acid Guibort, however, showed that the crystalline odoriferous principle of the tonka bean was distinct from benzoic acid, and named it coumarin. It occurs in Peristrophe angustifolia (Nees), Achylis triphylla (DC), and in many other plants, and is a normal constituent of lavender oil. It is present in plants as a glucoside, which is decomposed by a ferment present in the plant. This ferment is destroyed by alcohol or boiling water (Molish, J Pharm. Chim. 1902, 279).

It was first obtained synthetically by Perkin by heating the sodium compound of salicylaldehyde with acetic anhydride (Chem. Soc. Trans. 1868, 53); or more simply by boiling together salicylaldehyde, acetic anhydride, and dry sodium acetate (*ibid* 1887, 388; see also Tiemann and Herzfeld, Ber 1877, 284) Acetyl coumaric acid is first formed:

$$\begin{array}{c} \text{C}_{\bullet}\text{H}_{4} \swarrow \text{CHO} + \text{CH}_{3} \text{CO} \searrow \text{O} \\ = \text{C}_{\bullet}\text{H}_{4} \swarrow \text{OCO CH}_{3} + \text{H}_{2}\text{O} \end{array}$$

which then parts with acctic acid, yielding coumarin.

$$\begin{array}{c} \mathrm{C_6H_4} \diagdown \mathrm{CH} \cdot \mathrm{CH} \cdot \mathrm{COOH} \\ \mathrm{OCO\ CH_3} \\ = \mathrm{C_6H_4} \diagdown \mathrm{CH} \cdot \mathrm{CH} \\ \mathrm{O-CO} \\ \end{array} + \mathrm{CH_3 \cdot COOH} \\ \end{array}$$

By treating salicylaldehyde (1 molecule) with excess of acetic anhydride (4 molecules) for 6 hours at 180°, coumarin is formed (Reychler, Bull Soc chim. 1897, 17, 517)

Coumarin is also formed in small quantity by heating a mixture of phenol and malic acid with sulphuric acid

$$\begin{array}{c} \text{CH}_2\text{COOH} \\ \text{C}_6\text{H}_5\text{ OH} + \Big|_{\text{CH(OH) COOH}} \\ = \text{C}_6\text{H}_4 \Big|_{\text{O}-\text{CO}} + \text{CO} + 3\text{H}_2\text{O} \\ \end{array}$$

(Von Pechmann, Ber 1883, 2119, 1884, 929, 1649)

It has also been obtained from o-chlorocinnamic acid which is reduced electrolytically to o-chloro-hydrocinnamic acid and then heated with aqueous sodium hydroxide to 250°, when mellotic acid is produced Thionylchloride converts mellotic acid into mellotic anhydride, from which coumarin results by oxidation with bromine vapour at 270°–280° (Moyer, Beer and Lasch, Monatsh 1913, 34, 1665)

Substituted commarins may be obtained by condensing phenols with acetoacetic ester or with malic acid in the presence of concentrated sulphunic acid

'The condensation of acetoacetic ester or malie acid with substituted phenols (containing alkyl, hydroxyl or dialkyl-amino groups) gives good yields if the groups are present in the positions as shown—

The condensation is, however, prevented by the presence of nitro, carboxyl or carbethoxy groups (Clayton, Chem. Soc. Trans., 1908, 2018). It is also very considerably affected by the concentration of the sulphuric acid used

Coumarm forms rhombic crystals melting at 70°, boils at 290°-290 5°; has a pleasant, aromatic odour; is scarcely soluble in cold water, readily soluble in boiling water, very soluble in alcohol and ether. Crude coumarin may be purified by repeated boiling with light petroleum (Claassen, Pharm J 1897, 161).

It dissolves in caustic alkalis, and is reprecipitated on addition of acids If, however, it is heated for some time with concentrated caustic alkali, or better, with sodium ethoxide, a solution is formed from which acids precipitate

This compound can be reconverted into coumain by heating it with acetic anhydride; but when heated alone it is decomposed, yielding phenol and other substances. Reduction of coumarin with zinc in alkaline solution or sodium amalgam in aqueous solution yields mellotic acid as the chief product, with sodium amalgam and alcohol hydrocoumaric acid is formed (Williamson, J. 1876, 587)

On account of the differences shown between coumarin and thiocoumarin, it has been suggested that the two compounds should be represented by the annexed formulæ:

(Clayton, Chem Soc Trans 1908, 524).

The thiocommarin is produced by heating coumarin with phosphorus pentasulphide, and in contradistinction to coumarin is coloured. It also reacts with phenylhydrazine, yielding coumarin phenylhydrazine (with elimination of hydrogen sulphide), which cannot be obtained directly from coumarin itself. For an account of the synthetic benzotetronic acids (hydroxycoumarin carboxylic acids), see Anschutz, Annalen, 367, 174

In Germany, woodruff (Ger Waldmeister) is steeped in white wine, to which it imparts the pleasant aroma of coumain (Maiwein, Maitank) Instead of woodruff, an alcoholic solution of coumain is employed for the same purpose under the name of Maiweinessenz. In larger doses coumain is stated to have a narcotic action

Coumarin has the agreeable odour of asperula, and is used in perfumery for the preparation of asperula essence. It is also frequently used to adulterate extract of vanilla (q, v) (v) Lactones).

COUMARONE C<sub>8</sub>H<sub>4</sub> CH CH. An oil ob-

tained from the coal-tar fraction boiling between 165° and 175° After the pyridine bases and phenols have been removed by acid and alkali. it is obtained as the dibromide on addition of bromine. The dibromide is reduced to coumarone by alcoholic potash and sodium amalgam (Kraemer and Spilker, Ber 1890, 78) It can also be obtained from this source by adding pieric acid to the mixture of oils, whereby the picrate of coumarone crystallises out. This is separated, decomposed by hot water or dilute alkali, and the coumarone obtained by steam The pieric acid is recovered and distillation used for another operation (Ch. Fab. Act. ges. Hamburg, Eng. Pat. 1422, D. R. P. 53793; Ber. 1891, Ref. 233). Coumarone has been synthesised by heating o-hydroxy-ω-chloreinnamene OH C<sub>6</sub>H<sub>4</sub>·CH: CHCl with potash, hydrogen chloride being eliminated (Komppa, Ber 1893, 2968). It has also been prepared by heating phenoxyacetaldehyde with zinc chloride in

glacial acetic acid (Stoermer, ibid. 1897, 1700).

Coumarone boils at 174 0° (corr.) (Perkin,
Chem. Soc. Trans. 1896, 1249). It is unacted

upon by potash or by hydrochloric acid. Sulphuric acid reacts with coumarone in benzene solution, producing a soluble resinous material paracoumarone ( $C_8H_6O)_4$ , which is used to preserve wood and paper. Other polymerides are known. Coumarone is of interest as, when passed through a red-hot tube with benzene, phenanthrene is produced, and with naphthalene, chrysene is produced (Kraemer and Spilker, Ber 1890, 84). The methyl coumarones are present in the coal-tar fraction of b p 185°–195° (Stoermer and Boes, ibid 1900, 3013)

coumarone resins are prepared by treating solvent naphtha with sulphuric acid. They differ from the pitches by their leady solubility in acctone, and from the condensation products of phenols with formaldehyde by their solubility in petroleum spirit and by not yielding phenol when heated with soda-lime at 260°. They differ from the natural resins in being optically inactive, in having a lower melting-point, saponification value and iodine value. They are classified for commercial purposes according to their colour and hardness, those softening above 50° are termed 'very hard,' between 40° and 50° 'hard,' and between 30° and 40° 'medium hard'. For methods of their examination, see Maicusson, Chem. Zeit. 1919, 43, 109–122. Analyst, 1919, 44, 203

122, Analyst, 1919, 44, 203 If coumarone resins are distilled under ordinary or under reduced pressure they yield yellow oils between 150° and 240°, consisting mainly of coumatone, or partially depolymensed paracoumatone, to gether with indene hydrindene and small quantities of phenol By treatment with condensing agents and catalysts these oils again yield resins varying in character with the nature of the catalyst Coumarone oils are not autoxidisable like drying oils, but when mixed with 4-6 pc of driers, such as the resinates of lead or manganese, they yield dry films. The distinctive oxidation process of coumarone oils is to be attributed to indene (E. Stein, Zeitsch angew Chem 1919, 32, 246)

COUPIER'S BLUE v INDULINES.

COVELLITE. Native cupric sulphide, CuS, crystallising in the hexagonal system—Crystals are tare and have the form of thin six-sided plates. Fine large crystals have been found at Butte in Montana—The mineral has a characteristic indigo-blue colour, hence the name indigo-copper (German, Kupperindig)—It occurs in most copper-mines as a blue, earthy coating on other sulphide ores of copper, and is sometimes found in sufficient quantities to be of importance as an ore, eg at Butte in Montana, and Copiapo in Chile—Covellite—containing platinum (as the mineral speriylite, PtAs<sub>2</sub>) is mined in the Medicine Bow Mountains in Wyoming

LJS

COW TREE WAX v WAXES

COXPYRIN. Syn. for acetyl salicylic acid CRANBERRY, the fruit of Vaccinium Oxycoccos (Linn)

American analyses give as the average composition of the berries

 Water
 Protein
 Fat
 Carbohydrates
 Ash

 88 5
 0.5
 0.7
 10.1
 0.2

 Konig
 gives
 Other

 Water
 Protein
 Free acid
 Sugar
 carbohydrates
 Ash

 89 6
 0.12
 2.3
 1.5
 6.3
 0.15

The juice of cranberries was examined by Mach and Portele in 1888 (Landw. Versuchs. Stat. 1889, 38, 69), who found a litre of the fresh juice to contain (in grammes)

Invert Acidity (as sugar malic acid) Benzoic Tannic acid acid acid Ash Nitrogen 87.1 19.0 0.75 2.2 3.0 0.11

The sp gr. of the juice averaged 1 0591.

The juice will not ferment with yeast, owing, it is said, to the presence of benzoic acid, but this may be due to the poverty of the juice in introgenous matter, for Otto (Bied Zentr. 1899, 28, 284) found that bilberry juice would only ferment vigorously with yeast after the addition of asparagine (about 0 6 gram per litre) or of ammonium chloride (about 0 3 gram per litre) three.

Rising (Bied Zentr 1915, 44, 163) found in cranberries, benzoic, isovaleric, malic and citric acids, fermentable sugars and a glucoside—vaccinin  $\rm C_6H_{11}O_5$  Bz. The amount of sugar varied from 5 7 to 10 4 pc, and that of benzoic acid from 0 108 to 0 455 pc. The keeping qualities of the fruit are dependent upon the amount of benzoic acid present

The pigment of the cranberry consists of an anthocyanin—a compound of galactose with a cyanidin, the chloride of which has the composition  $C_{15}H_{11}O_6CI$  (Will-statter, Sitzungber K Acad Wiss Berlin, 1914, 12, 402).

Classen (ibid 16, 70) isolated a bitter sub-

Claassen (ibid 16, 70) isolated a bitter substance from cranberries, to which he gave the name oiycoccin, and which he found to give reactions similar to those of arbutin  $C_{12}H_{16}O_{7}$ , occurring in the leaves of the red bearberry, Arctostaphylos Uva-ursi (Spreng)

The ash of cranberry junce contains about 3 pc of phosphorus pentoxide and 47 pc of potash

H I

CREAM OF TARTAR v TARTARIC ACID. CREATINE, Methylglycocyamine, Methylguaninoacetic acid H2N C(NH)NMe CH2 CO2H,H2O discovered by Chevreul (Annalen, 1835, 4, 293) m muscular flesh (χρέας=flesh), occurs in human flesh, in the flesh and blood of many warm and cold blooded animals and in urine (Liebig, Annalen, 1847, 62, 257; Gregory, thid 1848, 64, 105; Schlossberger, thid 1848, 66, 80, Voit, Z. Biol 1868, 4, 77; Price, Chem. Soc Trans 1851, 3, Marcet, thid 1864, 2, 406; Folin and Denis, J Biol Chem 1914, 17, 488, Rose, ibid 1911, 10, 265); the amount varies with the species and the organ; the flesh of fowl yields 3 21 pc; of pigeon, 0 825 pc, of ox, 0 697 pc.; of ox heart, 1 375 pc, of herring, 1 324 p c., and of brill, 0 614 p c (Gregory, l c Beker, Zeitsch physiol. Chem 1913, 87, 28; according to Myers and Fine (J Biol. Chem 1913, 14, 9-12) it is constant for a given animal Okuda, 8th Int Cong App. Chem. 1912, 18, 275.

According to Folm (Amer J. Physiol 1905, 13, 45, 66, 117), van Hoogenhuyze and Verploegl (Zeitsch physiol ('hem 1908, 57, 161), Mellanby (J. Physiol 1908, 36, 447), muscular creatine has its origin in the liver. This organ is continuously forming creatinine from substances carried to it by the blood from other organs, and in the developing muscle. This is changed to creatine, until the muscle is saturated with creatine, when the excess of creatinine is excreted by the kidneys. Cf., however, Folin

CREATINE. 411

and Denis (ibid. 493-496 and 500-501) who suggest that creatine is part of living vertebrate protoplasm and the creatine of muscles a postmortem product, and that creatining elimination is a measure of the total normal tissue metabol-18m and independent of any special function of the liver; see also Myers and Fine (abid 1913, 15, 283-285 and 301-304), Inooye (Zeitsch physiol Chem. 1912, 81, 71-79), Ricser (abid 1913, 86, 415-453, and 1914, 90, 221-235), Myers and Fine (abid 1913-14, 16, 169, and abid 1915, 21, 377), Rose and Dimmitt (J Biol. Chem. 1916, 26, 345); Baumann and Hines (L Biol. Chem. 1917, 21, 540). The acceptance of the literature o (J Biol Chem 1917, 31, 549). The creatine is presumably combined with the muscular proto plasm, for though a readily dialysable substance, it does not pass into the surrounding lymph or blood in vitro, nor is it readily obtainable by diffusion from muscular flesh (Urano, Beiti Chem physiol Path 1907, 9, 104) Cf, however, Folin and Denis (J Biol Chem 1914, 17, 499); Leo and Howe (Pro Amer Soc Biol Chem 1912-13, xlin For the origin of urinary creatine, see Benedict and Osterberg (J. Biol Chem 1914, 18, 195), Myers and Fine (ibid. 1913, 15, 301)

In order to extract creatine, the flesh is heated with its own weight of water at 60°, and the juice expressed. The flesh extract is then heated to coagulate the protein, and filtered, the filtrate is treated with basic lead acctate until no further precipitation occurs, and again filtered after removing the excess of lead from the filtrate by sulphinetted hydrogen, when it is evaporated until creatine separates on cooling; this is washed with alcohol and recrystallised from water (Neubauer, Zeitsch anal Chem 1863, 2, 26; 1867, 6, 33). Creatine may be prepared by the decomposition of creatinine zine chloride by means of calcium hydroxide (Benedict, J. Biol. Chem. 1914, 18, 186).

Creatine is prepared synthetically by heating sarcosine with evanamide at  $100^{\circ}$ 

$$CN NH_2+NHMe CH_2 CO_2H$$
  
=  $NH C(NH_2) NMe CH_2 CO_2H$ 

(Volhard, Z. Chem. N. F. 1869, 5, 318, Strecker, J. 1868, 686), or, together with creatinine, by fusing sarcosine with guanidine carbonate (Paulmann, Beilstein Suppl. 1–657)

Creatine forms colourless transparent monoclinic prisms, containing 1H<sub>2</sub>O, which it loses at 100°; it dissolves in 74 4 parts of water at 18°, or in 9810 parts of cold absolute alcohol, and is insoluble in ether (Liebig, Annalen, 1847, 62, 257). Heat of combustion at constant pressure is 4240 calones (Emery, Benedict, Amer J. Physiol 1911, 28–307). The presence of urea, creatinine, or certain salts, raises the solubility of creatine in water and alcohol (Neubauer, lc). The aqueous solution is neutral, and has a bitter, harsh taste.

Creatine is decomposed by boiling baryta water into sarcosine, urea, and methylhydantoin NH  $C(NH_2)$  NMe  $CH_2$   $CO_2H + H_2O$ 

$$= CO(NH_2)_2 + NHMe CH_2 CO_2H$$

$$CO(NH_2)_2 + NHMe CH_2 CO_2H \rightarrow CO$$

$$NMe CH_2$$

$$NH \cdot CO$$

(Neubauer, Annalen, 1866, 137, 294); or by mercuric oxide and water into oxalic acid and

methylguanidine, or by heating with soda-limeinto ammonia and methylamine. Treated with mercum; acetate in presence of sodium carbonate it yields a-methylguanidinoglyoxylic acid NH2 C(NH) NMe CO ('O<sub>2</sub>H,2H<sub>2</sub>O; glistening flakes, m p 203°-204° (Baumann and Ingvaldsen, J. Biol Chem 1918, 35, 277) When potassium hydroxide is added to an aqueous solution of creatine and silver nitrate until the white precipitate first formed just redissolves, the liquid solidifies to a transparent gelatinous mass immediately reduced on heating, or in the course of a few hours at the ordinary temperature addition of potassium hydroxide to a solution of creatino and mercuric chloride precipitates a white crystalline compound (C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>N<sub>3</sub>),Hg,HgO (Engel, Compt. rend. 1874, 78, 1707, 1875, 80, Creatine is converted into creatinine by heating with dilute mineral acids for some hours (Dessaignes, J 1857, 544), or at 117° for 15 minutes (Benedict and Myers, Amer J Physiol 1907, 18, 362), or by boiling down to dryness the solution of creatine to which has been added an equal volume of hydrochloric acid (Benedict, J. Biol. Chem 1914, 18, 192), or by heating with water only in scaled tubes at 100° for 2-3 days (Neubauer, Zeitsch anal Chem 1863, 2, 33) Smoro-dincev (J. Russ. Phys. Chem. Ges. 1915, 47, 1275-9) obtained a double salt of silver nitrate and creatinine, C<sub>4</sub>H<sub>7</sub>ON<sub>3</sub> AgNO<sub>3</sub> (decomposes 188°-191°) on treating pure creatine with silver nitrate in a slightly acid solution

Creatine forms soluble normal salts with the mineral acids (Dessagnes, Annalen, 92, 409), and readily soluble compounds with zine chloride  $C_4H_9O_2N_3$ , $Z_nC_1$ , and cadmium chloride  $C_4H_9O_2N_3$ , $CdC_1$ , $2H_2O$  (Neubauer, *ibid* 137, 300)

Creatine is not usually estimated directly, it is converted into creatinine (q|r), and estimated as such; for the estimation of creatine in urine, r Benedict (J. Biol. Chem. 1914, 18, 192) and Walpole (J. Physiol. 1911, 42, 301), who makes use of the fact that a pale red colour is given by alkaline solutions of creatine and not by creatinine when a trace of diacetyl is added. For its estimation in blood r Greatine wald (Proc. Soc. exp. Biol. Med. 1917, 14, 115). a. Guaninopropionic acid, alacreatine.

## NH2 ('(NH)NH ('HMe'CO2H

from alanine and cyanamide (Baumann, Annalen. 1873, 167, 83), an isomeride of creatine forms, sparingly soluble prisms

The homologues of creatine are prepared by the action of cyanamide on the corresponding amino-acid in the presence of ammonia, e g α-aminocapiocyamine

from leucine and cyanamide (Duviller, Compt. rend 1886, 103, 211, 1887, 104, 1290). According to Duviller (lc) methyl- or ethylamino- acids yield with cyanamide the corresponding creatinine. The creatine is formed only in the case of methylglycocine and β-methylaminopropionic acid, the latter yielding methylaminopropionic acid, the latter yielding met

from a-methylaminopropionic acid and cyanamide, but prepared  $\beta$ -methylguaninopropionic acid NH<sub>2</sub>·C(NH)NMe·[CH<sub>2</sub>]<sub>2</sub>CO<sub>2</sub>H,H<sub>2</sub>O, mp 201°–202°, from cyanamide and  $\beta$ -methylaminopropionic acid, and  $\gamma$ -methylguaninobutyric acid NH<sub>2</sub> C(NH)NMe [CH<sub>2</sub>]<sub>3</sub>CO<sub>2</sub>H, mp 307°, from cyanamide and  $\gamma$ -methylaminobutyric acid.

Formaldehyde and creatine yield the compound C<sub>6</sub>H<sub>11</sub>O<sub>3</sub>N<sub>2</sub>,2H<sub>2</sub>O, decomposing at 250°, and forming a dibenzoyl derivative C<sub>6</sub>H<sub>9</sub>O<sub>3</sub>N<sub>3</sub>Br<sub>2</sub>, mp 265°-266° (Jaffé, Ber 1902, 35, 2896)

Phthalyldicreatine

C<sub>6</sub>H<sub>4</sub>[CO NH·C(NH) NMe CH<sub>2</sub> CO<sub>2</sub>H],

has m.p 212° (Urano, Beit. Chem Physiol Path 1907, 9, 183). Diacetylcreatine

NH C(NHAc)NMe CH, CO, Ac

has m p 165° (Erlenmeyer, Annalen, 1895, 284, 50) M A W.

CREATININE, Methylglycocyamidine

discovered by Liebig in human urine (Annalen 1847, 62, 268), the amount excreted being 26-30 mg per kilo weight of body per diem (Koch, Amer J. Physiol 1905, 15, 1), is also a constant constituent of the unne of horses, cows, dogs, pigs, and rabbits, it is found in muscle (Shaffer and Reinoso, Proc Soc Biol Chem 1910, 7, xxx, Meyers and Fine, J Biol Chem 1915, 21, 383) and occurs in small quantitative of the control o tities in the flesh of fish, about 0.2 p.c. in the case of cod (Poulsson, Chem Zentr 1904, 11 30), for other estimations, see Okuda (8th Int Cong. App Chem. 1912, 18, 275); and traces (0 000098 p.c., Cooper Colls, J. Physiol. 1896, 20, 107) are found in the blood. According to Folin and Denis (J. Biol. Chem. 1914, 17, 487) normal human blood contains 1 mg creatinine and 6-8 mg of creatinine and creatine per 100 grs. blood In domestic animals it is about the same; in birds 0.1 mg creatinine but 11 mgs creatinine and creatine It is probably a normal constituent of all soils (Shorey, J. Amer. Chem. Soc 1912, 34, 99-107, Sullivan, J Amer Chem Soc 1911, 33, 2035-42) and vegetable matter (Oshima and Arizumi, J Coll Agric Tohoku Imp Univ 1914, 6, 17-25)

Under normal conditions, the excretion of creatinine is constant from day to day and hour to hour (Shaffer, Amer J Physiol 1908, 23, 1), it is not affected by diet or exercise (Folin, ibid 1905, 13, 45, 66, 117. Pekelhating, van Hoogenhuyze Verploegh Proc k Akad Wetensch, Amsterdam, 1905, 8, 363. Klercker, Beit Chem Physiol Path 1906, 8, 59) It is now generally regarded as one of the end products of endogenous nitrogenous katabolism, the liver continuously forming creatinine from substances called to it by the blood, cf, however, Folin and Denis (J. Biol. Chem. 1914, 17, 500-501), part of this creatinine furnishes the muscular creatine by the change it undergoes in the developing muscle, v CREATINE; the rest is excreted by the kidneys (Folin, lc, Mellanby, J Physiol 1908, 36, 447; Lefmann, Zeitsch physiol Chem 1908, 57, 476) In cases of fever or starvation, the amount of urinary creatinine is increased, the material wanted for contraction being drawn from the muscular proteids (Pekelharing, van Hoogenhuyze Verploegh, *l.c.*; Leathes, J. Physiol. 1907, 35, 205). In cancer of the liver, the urinary creatinine is diminished and the creatine is increased (Mellanby, *loc.* 

Creatinine is extracted from urine by evaporating to one-third the original volume, decanting from the salts that crystallise out, precipitating with lead acetate and filtering, removing the lead from the filtrate and precipitating the creatinine as the sparingly soluble double salt with mercuric chloride. This is separated, decomposed by sulphuretted hydrogen, and the creatinine crystallised from the filtrate as the hydrochloride (Maly, Annalen, 1871, 159, 279), or by precipitating it as creatinine potassium picrate with subsequent decomposition of the double salt (Folin, J. Biol. Chem. 1914, 17, 463–467, Benedict, ibid. 1914, 18, 184–190)

For extraction from the body tissues and fluids, see Costantino (Chem Zentr 1915, 11, 287, from Arch farm sper 1915, 19, 254-258) Creatinine bears the same relation to creatine as hydantoin to hydantoic acid

and is prepared by boiling creatine with dilute mineral acids for some hours (Liebig, Annalen, 1847, 62, 628) or by heating at 117° for 15 minutes under pressure (Benedict and Myers, Amer. J. Physiol 1907, 18, 362), or by boiling down to dryness with an equal volume of hydrochloric acid (Benedict, J. Biol. Chem. 1914, 18, 192)

('reatinine can be prepared synthetically by heating guanidine carbonate with sarcosine at  $140^{\circ}-160^{\circ}$  (Horbaczewski, J Pharm Chim 1885, 86) A quantitative yield is obtained by heating N-methylglycocyamine with an organic acid (D R P 281051, from J Soc Chem. Ind. 1915, 34, 637)

By heating creatinine with silver nitrate and baryta in excess on the water-bath for about an hour a nearly quantitative yield of methyl guanidine can be obtained (Ewins, Bio-Chem J. 1916, 10, 104). By oxidation with mercuric acetate it yields α-methyl guanidinoglyoxylic acid (Baumann and Ingvaldsen, J. Biol Chem. 1918, 35, 277)

Creatinine forms anhydrous monoclinic prisms soluble in 11.5 parts of water or 100 parts of absolute alcohol at 16° (Liebig, *Le*), and separates on slow evaporation of dilute solutions in square plates containing 2H<sub>2</sub>O (Salkowski, Zeitsch physikal Chem 1880, 4, 133). It is a weak base, displacing ammonia from its salts, and forming soluble salts with the mineral acids (Liebig, *le*), the picrate C<sub>4</sub>H<sub>7</sub>ON<sub>3</sub>·C<sub>5</sub>H<sub>3</sub>O<sub>7</sub>N<sub>3</sub>, m.p. 212°-213°, crystallises in sparingly soluble yellow needles; the acid picrate

C4H,ON3.2C4H3O7N3

has m p  $161^{\circ}$ - $166^{\circ}$  (Mayerhoffer, Wien. Klin. Woch. 1909, 22 [3]); the tartrate

 $(C_4H_7ON_3)_2\cdot C_4H_6O_6$ 

decomposes at  $207^{\circ}$ – $209^{\circ}$ ; the oxalate

 $(C_4H_7ON_3)_2 C_2H_2O_4$ 

is sparingly soluble in alcohol (Poulsson, Chem. Zentr. 1904, 11 30) Creatinine forms characteristic double salts with platinic chloride (C<sub>4</sub>H<sub>7</sub>ON<sub>3</sub>,HCl)<sub>2</sub> PtCl<sub>4</sub>, orange-red plates, m p 220°-225°, with auric chloride

C4H7ON3 HCl AuCl3

yellow crystals, m p.  $170^{\circ}-174^{\circ}$ , with potassium picrate  $\mathrm{C_4H_7ON_3}$   $\mathrm{C_6H_3O_7N_3}$   $\mathrm{KC_6H_2O_7N_3}$ , lemonyellow prisms, 100 parts of water dissolve 0 1806 part of the salt at  $19^{\circ}-20^{\circ}$ , with zinc chloride ( $\mathrm{C_4H_7ON_3}$ ), ZnCl<sub>2</sub>, almost insoluble in alcohol (Heintz, J. 1847, 48, 883), with mercuric chloride 4( $\mathrm{C_4H_5HgON_3}$  HCl) 3HgCl<sub>2</sub>,2H<sub>2</sub>O(Johnson, Proc. Roy Soc 1886, 43, 493)

Creatinine is converted into creatine by the action of cold dilute animonia or calcium hydroxide solutions, or on treatment with iodine (Reichardt, Pharm Zeit 1911, 56, 922); boiling baryta solution converts it into ammonia and methyl hydantoin, with alkaline oxidising agents it yields methylguamidine and oxahe acid, but is comparatively stable towards acid permanganate solutions (Jolles, Bei 1902, 35, 160)

The presence of creatmine in dilute solution or in urine can be detected by (1) Maschke's test (Zeitsch anal Chem 1878, 134), which consists in the formation of a spaningly soluble double compound of creatinine and cuprous oxide, when the suspected solution is heated at 50°-60° with Fehling's solution, as the cuprous oxide is formed by the oxidation of part of the creatinine, it is advisable to add a little glucose to the mixture, the test is sufficiently delicate to allow of the detection of 10000 part of creatinine (2) Weyl's test (Ber 1878, 11, 2175) consists in the production of a ruby-ied colouration, changing to yellow when, to a solution of creatinine, a few diops of 10 p.c. sodium nitroprusside and the same quantity of 10 p c sodium hydroxide or carbonate solution are added acidifying the solution with acetic acid and boiling, the colour becomes emerald green, changing to blue, due to the formation of Prussian blue (Salkowski, Colasanti, Gazz chim ital 1887, 17, 129). Guareschi (Bei 1888, 21, Ref. 372) has shown that this reaction is given by other compounds than creatinine containing the —CO CH,— complex (3) Jaffe's test (Zeitsch physiol Chem 10, 399) is the production of a deep-red colour, when pieric acid and an excess of sodium hydroxide solution are added to the hquid containing the creatinine Chapman (Chem News, 1909, 100, 175) has shown that the colour is due to the sodium salts of aminodinitrophenol (pieramic acid), and diaminonitrophenol formed by the reduction of the pieric acid by the creatinine; if the creatinine is present in excess, reduction to the colourless triaminophenol occurs.

The older methods for the estimation of creatinine were based upon the isolation of the base in the form of its sparingly soluble double salt with mercuric or zinc chloride, and either weighing the precipitate or estimating the metal

or the mtrogen in the compound (Neubauer, Annalen, 119, 35, Salkowski, Zeitsch physiol. Chem. 1886, 10, 113; Grocco, Chem. Zentr. 1887, 17, Kolisch, *ibid* 1895, 1814, Ladd and Bottenfield, Amer. Chem. J. 1898, 20, 869) Edleson (Chem. Zentr. 1909, 1108) recommends precipitating the sulphate in the presence of alcohol and ether, dissolving it in water, and tirrating the solution with standard barrum hydroxide solution, using phenol phthalein as indicator.

The method generally adopted for estimating creatinine in urine is Folin's colorimetric method, based on Jaffé's reaction (Zeitsch physiol Chem 1904, 41, 223) N/2 potassium dichromate solution is placed to a height of 8 mm in one tube of a Duboseq colonmeter; 100 cc of the urine is placed in a 500 c c flask with 10 c c of a 12 pc solution of pieric acid and 5 cc of a 10 pc solution of sodium hydroxide, left for a few minutes and made up to the mark. If x is the length of this solution in the second tube required to give the same colour effect as the 8 mm, of the standard dichromate solution in the first tube, then  $8.1 \times 10/x = mg$  of creatinine in the 10 cc of the urine If a is less than 5, only 5 c c of urine should be used, if a is greater than 13, then 20 cc of urme should be used If creatine is also present in the urine, the amount may be estimated by first determining the creatinine as above, then converting the creatine in another 10 cc of the urine to creatinine by heating it with 5 c c of N/2 HCl on the water-bath for 3 hours or for 15 minutes at 117° under pressure (Benedict and Myers, Amer J Phys 1907, 18, 362), or by boiling down to dryness with an equal volume of hydrochloric acid (Benedict, J. Biol. Chem. 1914, 18, 191-194), The differand estimating the total creatinine ence between the two results represents the amount of creatmene equivalent to the creatine originally present (For precautions to be observed regarding temperature, time, and dilution, in applying this method for the estimation of creatinine, compare Chapman, Chem News, 1909, 100, 175) Folim and Morris (J Biol Chem 1914, 17, 469-473) recommend the use of a standard solution of creatinine instead of a dichiomate solution; see also Folin and Doisy (ibid 1917, 28, 349-356), Gettler (ibid 29, 47) The presence of dextrose, acetone and acetoacetic acid interferes with the final colour reaction and various suggestions have been made for overcoming the difficulty (Graham and Poulton, Proc Roy Soc 1914, B, 87, 212, Greenwald, J. Biol Chem 1913, 14, 87, Rose, J Biol Chem. 1912, 12, 73–80) Morris, J Biol Chem 1915, 21, 201-208, recommends the preliminary precipitation of creatinine as the double potassium picrate See Bauman and Ingvaldsen (ibid 1916, 25, 196-200) for the quantitative precipitation of creatinine with potassium pierate

An isomeride of creatinine, alacreatinine,



was prepared in 1873 (Ber 6, 1371), and other isomerides have been described by Korndorfer, Arch. Pharm. 1904, 242, 620; Schenck,

ibid 1910, 248, 376, and 1911, 249, 463; Schmidt, ibid. 1910, 248, 568; Johnson and Nicolet (J. Amer Chem Soc 1915, 37, 11, 2417) obtained two methyl glycocyamidines somenic with creatinine by the action of methylamine on the ethyl ester of benzoylpscudo-othylthiohydantore acid

Two isomenic nitrosocreatinines were obtained by Dessaignes (Annalen, 1856, 97, 341) and Marcker (ibid 1865, 133, 305) by passing the nitrous fumes from the action of intric acid on assenious oxide into an acid solution of creatinine. The less soluble a-nitrosocreatinine  $C_4H_8O_2N_4$  decomposes at 2103, forming a colourless liquid that quickly solidines to a brown mass from which a new base  $C_7H_{12}O_2N_{10}$  (?) can be extracted, it forms a crystalline nitrate, hydrochloride, and platinichloride, and yields the bromoderivative  $C_4H_7O_2N_4$  melts at 1953 to a brown liquid and decomposes at 2203

According to Kramm (Chem Zenti 1898, 1 37), the yellow colour formed by the interaction of sodium introprusside and sodium hydroxide on creatinine (cf. Woyl, Ber. 1878, 11, 2175), is due to the formation of a nitrosocialization of a nitrosocialization of the yellow alkaline solution and shaking vigorously. According to Schmidt and Hennig (Arch. Pharm. 1912, 250, 345) this is an oxime, they find that on treatment with sodium nitrite in intricacid solution oxime, in p. 193~194°, and "creatinine oxime, which begins to discolour at 250° but does not melt."

The following acyl- and alkyl- derivatives of creatinine have been prepared Benzoylcreatinine C<sub>11</sub>H<sub>11</sub>O<sub>2</sub>N<sub>3</sub>, pale yellow needles, mp 187° (Urano, Beiti Chem Physiol Path 1907, 9, 183), methyl, dimethyl and trimethyl creatinine and salts (Korndorfer, Arch Pharm 1904, 242, 641, Kunze, Arch Pharm 1910, 248, 578–93).

Methyl creatinine HN C | gives an aurichloride, mp. 170°-171°, and dimethyl creatinine gives an aurichloride, mp. 128°-129°

creatinine gives an aurichloride, in p. 128°-129°, and a platinichloride, m.p. 177°-179° According to Kunze the constitution of dimethyl-NMe CH<sub>2</sub>

creatinine is MeN.C. NMe CH<sub>2</sub>; it reacts as a NMe—CO NH—CO

tertiary base; ethylcrcatinine C NEt NMe—CH<sub>2</sub> (Neubauer, Annalen, 119, 50), see Henzerling (Arch Pharm 1910, 248, 594, the platinichloride crystallises in monoclinic plates, mp 197°-211°, ethylcrcatinine hydriodide, mp 217°-219°, diethylcrcatinine platinichloride 201°-202°);

methylamino-a-butyrocyamidine

(Duvillier, Compt. 1end 1882, 95, 456, 96, 1583, 97, 1486, 100, 916, 103, 211, 104, 1290).—For Xantho-, Cruso-, and Amphi-creatinine v Xanthocreatinine. M A W

**CREOLIN.** A disinfectant consisting of an emulsion of coal tar cresols with soap

CREOSOFORM. A condensation product of formaldehyde and creosote Used as an internal antiseptic

**CREOSOL**  $\hat{v}$  CREOSOTE, WOOD TAR

**CREOSOTAL** (Creosote carbonate) is a mixture of carbonates of the phenols present in word creosote, of which the chief are guaracol and creosol

It is produced by heating creosote with a benzene solution of carbonyl chloride (phosgene) under pressure (Heyden, Fig. Pat. 19074, 1890), but is generally prepared by passing phosgene gas through a solution of beechwood creosote in caustic soda solution. The carbonate separates as an oil, and is washed with dilute caustic soda and finally with water

Crosotal is a viscid yellowish liquid, insoluble in water, but soluble in alcohol, chloroform, benzene, &c (Aufrecht, Pharm Zeit 1908, 53, 480) When it is boiled with caustic potash, the odour of crosote is evolved. It is quite odourless, tasteless, and non-initiant in its action upon the mucous membrane, and is therefore much used as a substitute for crosote in the treatment of bronchial affections and pneumonia, especially in children (Weber, Pharm J 68, 115, Toff, ibid 73, 414) The dose for adults is about 60 grains daily, and it is usually given in an emulsion (Haefelin, Pharm Zeit 49, 141)

CREOSOTE or KREOSOTE (from  $\kappa\rho\epsilon\omega s$ , contracted gentive of  $\kappa\rho\epsilon\alpha s$ , meat, and  $\sigma\omega\zeta\dot{\alpha}$ , I save or preserve). Wood tar, and coal tar, when subjected to distillation, yield fractions technically known as 'creosote,' both of which, the latter more especially, find extensive and important uses in the industrial arts

Reichenbach, in the year 1832, applied the term 'kieosote' to a substance derived from wood tar, and which has been found to possess powerful antiseptic properties, shortly afterwards, Runge discovered carbolic acid in coal tar, and the two products were, for a considerable time, regarded as identical, similar compounds, however, have since been procured from other sources, and in order to prevent confusion, it has been suggested to restrict the use of the term 'creosote' to the use in a generic sense as referring to all the phenols and phenoloid bodies derived indifferently from coal, wood, coke-oven, blast-furnace, and shale-oil tars. The enormous extension of the use of creosote oil from coal tar for the preservation of timber has resulted in the terms 'creosote' and 'creosote oil' being used commercially as meaning the heavier distillates from coal tar and coke oventar, the oils from blast-furnace tar being distinguished as 'blast-furnace creosote'

Wood-tar creosote. The tar derived from the distillation of beechwood, when subjected to further distillation, yields certain fractions that are heavier than water, and which, after agitation with solution of caustic soda, and separation from insoluble oils, are boiled with free access of air in order to oxidise various impurities present, the alkaline solution thus obtained is decomposed with dilute sulphuric acid, the crude creosote which separates is again submitted to a similar alkaline and acid treatment, and the product finally distilled, reserving the traction passing over between 200° and 220°

This represents the ordinary 'wood-tar crosote' of commerce It is a complex mixture of phenoloid compounds, the proportions of which are materially influenced by the method originally followed in distilling the beechwood

from which it was obtained.

Wood-tar crossote, when freshly prepared, is a colourless, transparent liquid of an only consistency, and which retains its fluidity at a very low temperature, its sp gr varies from 1 037 to 1 087; it boils at 205°-225°, and dissolves in about 200 parts of water; its odour is strong and penetrating, resembling that of wood smoke, or rather of smoked meat, it refracts light powerfully and burns with a very sooty flame. Wood-tar creosote is powerfully antiseptic, but its power of coagulating albumen has been denied, nevertheless, it preserves animal substances as effectively as carbolic It is less caustic than carbolic acid, and is not poisonous Wood-tar creosote may be regarded as consisting of a mixture of compounds included in several homologous series, chiefly of acid methylic ethers of catechol, and the presence of the following bodies has been established

CS	stablished		
		ormula	Boiling point
1	Monohydric phenols Phenol, carbolic acid C <sub>8</sub> H <sub>5</sub> OH		182°
	Cresols, cresylic acid $\{C_6H_4(CH_4)\}$	(3)OH	203°
	$X$ ylenols or phlorols $\{C_6H_3(CH_3)\}$		220°
2	Ethyl phenol (1-2) $C_6H_4(C_2)$	H <sub>5</sub> )OH nols	203°
	Gualacol or methyl- C <sub>6</sub> H <sub>4</sub> (O	$_{ m H}^{ m CH_3} \}$	200°
	Creosol or methyl- CaH3(CI	${ m P}_{ m OH}^{ m OH}$	219°
	Dimethyl homocate-	I)(OCH <sub>3</sub> ) <sub>2</sub>	214°-128
	Homocreosol or di- methyl guaiacol C <sub>6</sub> H <sub>2</sub> (CF	$I_3)_2\left\{ egin{matrix} \mathrm{OCH}_3 \ \mathrm{OH} \end{smallmatrix}  ight\}$	230°
	Ethyl guaracol(1-3-4)C <sub>6</sub> H <sub>3</sub> (C <sub>2</sub>		
3	Corulignol or pro- pyl guaracol C <sub>6</sub> H <sub>3</sub> (C <sub>3</sub> : Methyl ethers of trihydric phe	H), {OH }	· 241°
J	100	'H <sub>3</sub> )	
	Dimethyl-pyrogallol $C_6H_3$ Of	$\mathbf{H}^{\mathbf{H}_{\mathbf{J}}}$	253°
	Dimethyl - methyl- C6H2(CF	$I_3$ $\left\{ \begin{array}{l} OCH_3 \\ OCH_3 \end{array} \right\}$	265°
	Dimethyl - propyl- pyrogallol (pica- mar)	'UL'	285°
	$\left. \begin{array}{c} Methyl-propyl-pyro-\\ gallol \end{array} \right\} C_{\bf 6} H_{\bf 2} (C_{\bf 3})$	$_{\mathrm{H_7}}$ $\left\{ _{\mathrm{OH}}^{\mathrm{OCH_8}} \right\}$	290°

Of these compounds, phenol is present in have been deprived o extremely small quantity, paracreosol in larger, guaiacol it contained.

and phlorol in notable proportion, but the two principal constituents of creosote are guaiacol and creosol, although not equally present in all varieties, guaiacol, according to Allen, predominates in Rhenish creosote, whilst with Morson's creosote from 'Stockholm tar,' creosol is the chief constituent Homocreosol and cœrulignol are present in small quantity only, but, according to Gratzel, the latter body possesses such energetic and astringent qualities that a single drop placed on the tongue causes bleeding, purified creosote should therefore be absolutely free from this compound, which may be detected by the blue colouration produced by baruum hydroxide

The less valuable portions of wood-tar crossote were found by A W Hofmann to contain the methyl ethers of pyrogallol and its homelogues, these bodies are of interest from the remarkable colouring matters (originally discovered by Runge) which may be derived from them by oxidation, thus, if the sodium derivative of dimethyl-pyrogallol be mixed with the sodium derivative of dimethyl-methyl-pyrogallol and excess of sodium hydroxide and heated in the air, a substance called cupitlonic actal is formed, according to the following equation —

 $(C_6H_{10}O_3)_2 + C_9H_{12}O_2 + 2O_2 = C_{25}H_{26}O_9 + 3H_2O_2$ 

Eupittonic acid has the constitution of a hexamethoxyl aurin  $C_{19}H_8(O(H_3)_8O_3)$  Reichenbach's 'pittical,' called by Wichelhaus 'eupittone,' was a salt of cupittonic acid Wood-tai crosote has been met with adulteriated to a considerable extent with cuide carbolic and cresplic acids, and from the fact that the reactions of these substances closely resemble those of wood crosote, the distinction between the two, especially when in admixture, is attended with some difficulty. Genuine beechwood tai is soluble in solutions of caustic alkalis, and forms a crystalline compound with potassium hydroxide, but not with sodium hydroxide, on the other hand, it is practically insoluble in strong ammonia

An alcoholic solution should give no colouration whatever (blue or red) with baryta water, such colouring indicates the presence of conulignal and other impurities

The composition of beechwood and oak crossotes, freed from hydrocarbons, is given by Béhal and Choay (Compt rend 1894, 119, 106),

as follows ---

			Beech	Beech	Oak
	-				
Distillation peratur		m-	200-220°	200-210°	200-210°
Specific g			1 085	1 085	1 068
Monophei			39 0	39 0	55 0
Guaiacol			19 7	26 5	140
Creosols a	nd hor	110-			
logues			400	32 1	310
Loss			13	24	

Owing to the demand for gualacol for pharmaceutical purposes, wood-tar creosote, as it occurs in commerce, is frequently found to have been deprived of part or the whole of the gualacol it contained.

Methods for the determination of gualacol in wood crossote are given by Béhal and Choay (Compt rend. 1893, 116, 197), M Adrian (Nouv. Rem. 13, 97); L. F Kebler (Amer J. Pharm. 1899, 409)

Pharm. 1899, 409)

Wood-tar crossote is sharply distinguished from the coal-tar acids by its insolubility in absolute glycerol, it is also distinguished from the coal-tar acids by its reaction with an ethereal solution of nitrocellulose, shaken with half its volume of collodion, BP, Calvoit's No 5 carbolic acid coagulates the gun-cotton Morson's crossote does not precipitate the introcellulose from collodion, but mixes perfectly with the ethereal solution; when a mixture of equal volumes of Morson's crossote and Calvort's No. 5 acid is treated with the collodion, the precipitation is very marked.

Creosote from coal tar and other sources. pharmaceutical circles the term 'creosote' is still retained for products of the carbonisation of wood, but, generally speaking, it is recognised in industry as applying to a mixture, consisting principally of hydrocarbons distilled from coal tar, the latter being either of gasworks origin, or from the carbonisation of coal in coke ovens At the time when anthracene was a valuable commercial product, the fraction of coal tar containing this material was usually collected separately, the crude anthracene being removed by filtration, and the oil separated therefrom being placed on the market under the names 'anthracene oil,' 'green oil,' or 'heavy oil' In more recent times many tar distillers have found the recovery of naphthalene unremunerative, and this is especially the case where the crude tar is of such a nature that an appreciable proportion of solid paraffinoid hydrocarbons appears The whole of the anthracene fraction has therefore frequently found its way to the Other distillates of coal tar crossote storage have also been passed into the creosote tanks at various works, including the 'sharp oil' separated during the recovery of crude naphthalene, and the 'middle oils' or 'carbolic oils,' after treatment for the removal of 'tar acids' (crude There is no very serious objection to this development, especially in view of the fact that inclusion of the anthracene fraction results in a mixture containing a large proportion Creosote of this of high-boiling constituents nature is employed in the United States for railway work, and has been exported in recent times to that country in very large quantities by means of 'tankers.' But exception must be taken to the practice that has arisen in some cases whereby the creosote storage has largely assumed the character of a refuse pit for the disposal of various oils, and even tars, that are only marketable with difficulty, such as blastfurnace tar, blast-furnace creosote, water-gas It is probably most satisfactory to tar, &c regard creosote as a fraction obtained during the distillation of coal tar, either of gasworks or coke-oven origin, collected between the light oil fraction and the anthracene fraction; it may or may not include the middle oils, or those portions of them remaining after partial or complete removal of tar acids or naphthalene.

Creosote from coal tar (gasworks). This consists, for the most part, of a mixture of liquid aromatic hydrocarbons with naphthalene.

The naphthalene may be present in such small quantities that the oils retain it entirely in solution, or there may be so much of it, say up to 40 p.c, or more, that the creosote exists as a solid mass at 15°C. Phenolic and basic constituents are also present, together with small amounts of paraffinoid hydrocarbons. The creosote produced from tar that is made in London and the southern districts of England, where Durham and other north country coals are largely carbonised, contains considerably more naphthalene, as a rule, than that obtained in more northerly districts, where Midland coals are retorted On the other hand, the creosote obtained from Midland tar is far richer in tar acids than that made in London. So characteristic are these features that various creosotes are regularly known in the trade as 'Midland creosote,' 'ordinary London creosote,' &c Scotch creosote will be dealt with later. The carbonisation of coal in gasworks has, during the last few years, developed in several directions. It was formerly the custom to work with 'light charges' of coal in the retorts, with the result that a relatively large free-space existed between the surface of the coal and the top of the retort But about 1911 the practice was widely adopted of employing heavier charges, wholeby the 'free space' was greatly reduced The result of this change of procedure was that the volatile products of carbonisation experienced a much more rapid journey to the ascension pipes of the retort, and consequently the changes they underwent, due to thermal decomposition in the free space, were lessened. The effect of this change on the quality of the tar was very noticeable. The quality of the creosote seems to be very similar to that resulting from 'light charge' carbonisation, but there is more of it in the tar The proportion of naphthalene in the crossote is lower, while the proportion of tar-acids is somewhat higher. The average yield of creosote from coal tar is roughly about 20 pc of the latter by volume

Creosote being a mixture of many hydrocarbons, &c, it is not possible to give any definite figures for its physical and other constants, and the more so as it is highly improbable that any two makes are identical. When a sample of good quality is distilled in the laboratory, the main bulk of it usually passes over between 200° and 300° C, the bulb of the thermometer being immersed in the vapour If the anthracene fraction be present there will be a considerable residue at the latter temperature The calorific value is about 16,050 BTh U The evaporating power (in actual practice, 'from and at 212° F.') is 12 0 lbs. The flash-point usually lies about 170°-180° F (closed test). The refractive index of the higher fractions should be above 1.6 (Sage). Determination of the specific heat between 30° and 100° C. gave the figure 0 398. The coefficient of expansion per °F. is about 0.000455. The specific gravity usually lies between 1 030 and 1 080. Creosote generally contains between 0 5 and 0.9 p.c. of sulphur.

Although these figures naturally differ with the various makes of creosote, and can therefore only be regarded as approximate, they, nevertheless, are frequently found useful in industry, and several serve as evidence of admixture of adulterants of a paraffinoid nature, such as have already been mentioned in this article.

Creosote finds its largest application in the treatment of wood, especially railway sleepers, piles, and telegraph poles. Its employment as fuel is being very rapidly developed, chiefly in burners which serve to atomise the liquid, either by direct application of pressure, or by means of steam or air under pressure. Numerous patents have been granted for such burners, and for further information on the subject, consultation of such works as Brame's Fuel may be recommended. The employment of croosote as a fuel for Diesel engines, with or without pilot ignition, is extending, and the 'Diesel Engine Users Association' has recently issued a specification for the quality of oil considered to be best suited for that particular purpose Croosoto oil is (when freed from naphthalene) used in the preparation of disinfectants and sheep-dips. It is also employed for generating artificial light in specially designed lamps such as the 'Lucigen.'

Lamp-black is manufactured by partial combustion of creosote in a limited air supply Recent experiments seem to indicate the possible

value of crossote as a soil steriliser

The best known methods of creosoting timber are perhaps those of Bethel, Boulton, Ruping, Curtis Isaaes, Lawry, Rutgers In some of these processes the croosote is applied under pressure, occasionally after first placing the timber under a vacuum, and at other times the oil is employed at a temperature sufficiently high to boil the sap out of the wood in vacuo.

Many specifications for creosote considered to be best suited for this work have been published; probably the best known is that of Dr. Tidy. The advantages resulting from the creosoting of timber are summarised by Tidy as follows—

Physical —Choking of pores, and thus rendering the material non-absorbent of water.

Brochemical —Inhibition of germ life.

Chemical —Coagulation of albumen by the tar acids

Probably most of the creosote sold for creosoting at the present time is subject to the American 'Maintenance of Way' specification. The most important feature is the insistence on a large proportion of high-boiling oils. It is estimated that in 1913, 35,000,000 gallons of creosote were exported to the United States from the United Kingdom. At various times emphasis has been given to one or more of the following properties of creosote for treatment of timber:

1. Maximum naphthalene content.

2. Minimum naphthalene content.

3 Tar acids.

4. Proportion of high-boiling constituents.

It is considered that the life of railway sleepers is increased five-fold by creosoting. Experiments by J. M. Weiss (USA.) (see J. Soc. Chem. Ind. 1911, 1348), show that coal tar creosote is more efficient as a preservative than is that from water-gas tar, in the proportion of about 6 to 1. Creosote is used for the treatment of wooden paving blocks, for which purpose it is valuable as a preservative and also as a means of rendering the blocks impervious to

water, the absorption of which would give rise to expansion of the wood.

The following are among the tests commonly applied to creosote —

1 Distillation.

2. Liquidity of residue left after distillation.

3 Water.

4. Solid matter (naphthalene, &c.) deposited at  $60^{\circ}$  F

5. Loss on evaporation under specified conditions

- 6 'Absorbent spot'—a drop of the oil being placed on blotting-paper, and examination made for particles of 'free carbon' indicative of adulteration by crude tars
- 7 Taracids
- 8. Specific gravity.
- 9 Colour.
- 10 Fluorescence
- 11 Refractive index—serving to indicate presence or absence of paraffinoid hydrocarbons, or tars containing them.
- 12 Flash point
- 13. Matter insoluble in benzol.

14. Sulphonation

(Cf Sage, J Soc Chem. Ind 1911, 588.)

Creosote from coke-oven tar is very similar

to that from gasworks tar. Oils from low-temperature processes of carbonisation. A great deal of work has been carried out in investigating the products of these systems Some of the processes are The volatile worked under partial vacuum. products of carbonisation are not subjected to the secondary thermal decompositions that are incidental to high-temperature work in horizontal retorts, with the result that the proportion of aromatic hydrocarbons appearing in the oils is very much smaller, the specific gravity of those fractions corresponding to creosote being The content of tar acids is lower than 1 000 very much higher, though there is but little actual phenol From the highest fractions solid paraffinoid hydrocarbons are frequently deposited. Work by Pictet and Bouvier, on the carbonisation of coal at low temperatures and in partial vacuum, points to the conclusion that the primary condensible products are similar to the constituents of certain petrolcums, consisting to a considerable extent of terpenes and naphthenes No solid aromatic hydrocarbons are present. On subjecting these lowtemperature tars to heat by passage through a tube or over red-hot coke, benzene, naphthalene, anthracene, &c, are formed, the resulting oils being very similar to ordinary coal tar creosote. According to Wheeler, 'vacuum' tar is largely a mixture of ethylenic, naphthenic and paraffin hydrocarbons, and homologues of naphthalene, together with phenoloid compounds, chiefly cresols and xylenols. Aromatic solid hydrocarbons are generally absent D. Trevor Jones concludes that the ordinary products of coal tar result from the action of heat on a tar which is formed in the first instance, and which is similar to that obtained under low-temperature conditions. The naphthenes, paraffins and unsaturated "hydrocarbons are decomposed to produce olefines, which are at their maximum at 550° C. The olefines vanish at 750° with the simultaneous appearance of naphthalene and of large quantities of hydrogen. He further concludes that acetylene plays an insignificant part in the formation of coal tar products. Low temperature carbonisation on the large scale yields a very oily tar amounting to 25 or more gallons per ton of coal. This tar is rich in those oils corresponding in distilling range to the creosote of normal tar. In the carbonisation of cannel coal by such processes as much as 50 gallons of tar is obtained.

Creosote from vertical retort tar is characterised by its low content of naphthalene, its high content of tar acids (chiefly higher homologues of phenol), and the large proportion of paraffinoid bodies present, the latter being sufficient to bring the specific gravity below

1.000.

West introduces steam during the process of vertical carbonisation, using continuous vertical retorts. The yield of tar is increased very considerably and reaches 20 gallons per ton. The tar is of low specific gravity and gives a large 'creosote' fraction. The tars evolved during the carbonisation of coal in vertical retorts are not subject to such high temperatures as in the case of horizontal retorts. Consequently these tars resemble, to a certain extent, the low-temperature products already described. Similar remarks apply to the tars from blast furnaces and from power-gas producers, where coal is employed as fuel

Blast furnace creosote. This is sometimes known as 'phenoloid' It resembles vertical retort creosote in many ways. The specific gravity of the hydrocarbons is low, the proportion of tar acids is high, reaching 20-35 p.c., though there is little phenol. The phenoloids from blast-furnace tar, extracted by Watson Smith, contained only 1.33 p.c. of real phenol, boiling at 182°, whereas the tar acids from Lancashire tars yielded 65 p.c. of crystallisable

carbolic acid

The fractions passing over between 210° and 230° contain phlorol and creosol, and those following at a higher temperature, viz 360°, furnish compounds allied to the cupittonic acid obtained from wood tar

This creosote has a very strong and characteristic odour. Blast furnace crossote, and sometimes even the crude tar from which it is distilled, is often mixed with coal tar creosote, and several of the tests generally applied to the latter are employed for the purpose of detecting

this undesirable adulterant

Blast-furnace creosote oil is now produced in large quantities in Scotland. It is employed to a certain extent for creosoting timber, in producing the 'Lucigen' light, and as a liquid fuel. It furnishes from 20 to 35 p c of phenoloid substances soluble in caustic soda, as against 4 to 7 p.c in London coal-tar creosote, and about 16 p.c for Midland and country make. Allen and Angus have secured by patent their use as an antiseptic under the name of 'neosote,' a word signifying 'new preserver' (Eng. Pat. 11689, August 29, 1887) Experiments have shown that the substance compares favourably with carbolic acid, whilst its caustic properties, when applied to the skin, are much less marked. When freshly prepared, the fluid is almost colourless, and bears a great resemblance to wood-tar creosote.

Creosote from producer tar. -In cases where

bituminous coal is gasified a tar is formed, is creosote from which is similar in several respe to the blast-furnace oil. Tar acids and paffinoids are present in large quantity.

Creosote from water-gas tar is characteristy the absence of tar acids. It is usually r in paraffins, of low specific gravity, and m contain anthracene and naphthalene.

Scotch creosote is characterised by its laicontent of paraffinoid hydrocarbons and con

quent low specific gravity.

Employment of creosote oils for lighting Creosote oil has long been employed as a sou of light in the construction of harbours and o door work, where smoke was of little con quence or inconvenience; but when it is wish to burn this oil to the greatest advantage a with the least possible amount of smoke a soot, it becomes necessary to employ lamps constructed that a jet of air or steam, more less superheated, may be introduced in such manner as to ensure more perfect combustic Hartmann and Lucke (D R P 9195, Aug 1879) constructed a lamp specially intended the consumption of creosote oils, in which the is automatically supplied to a dish-shar receptacle, over which is placed a funnel-shar cylinder In the centre of the dish or recepta receiving the oil from its attached reservoir, placed a conical tube, reaching a little abo the surface of the oil, and through which a jet superheated or dry steam may be driven. using the lamp, the dish or receptacle refer to is allowed to fill with oil, it is then ignited its surface, the funnel-shaped cylinder plac over it, and steam cautiously admitted throu the conical tube It will readily be seen th the arrangement closely resembles the 'He path blowpipe,' with the difference that t steam is forced through a body of co bustible fluid, issuing just above its surface, place of air forced through a body or envelo of combustible gas, and issuing at a point position that may be called its surface

The oil thus burns without wick or chimne and affords a light of 180 standard candles, we an hourly consumption of a little over 2 lbs.

creosote.

An apparatus for a similar purpose has be patented by Lyle and Hannay, under the nat of the 'Lucigen' It consists of an oil tank reservoir, fitted with a special burner at the t of a tube, which may be extended to any lengt into this reservoir compressed air is introduc by an indiarubber pipe, so attached as to ma the 'Lucigen' portable. On the compress air being admitted into the oil tank or receptac it forces the oil up through an internal tube, ar escaping at the same time with the oil in t burner, produces a spray which is set on fire. light of about 2000 candle-power is given by t 'Lucigen.' The flame is large, the light w diffused, and the eyes are not dazzled by There is little or no smoke from the flame, a the lamp burns equally well under heavy ra or spray; it requires no lantern, and has parts which can be damaged by rough usa, In the 'Doty' and 'Wells' lights, pressure brought to bear on the surface of the oil or cre sote by means of an air pump, whereby the oil forced through a coil surrounding the lower po tion of the flame, and becomes distilled into ge CRESOL. 419

Creosote oil has also been employed for the injecting into the brick kilns during the burning purpose of illuminating and heating, by direct conversion into permanent gas. A ton of creosote is capable of yielding 13,300 cubic feet of gas, with an illuminating power of 14 candles, a yield of 29,300 cubic feet may be obtained, but of 81 candle-power only; the chief drawback to this method of utilisation is the high temperature necessary to decompose the oils, which results in the production of low candle-power

Employment of creosote as fuel —Creosote oil is extensively used as a fuel, both for marine and stationary purposes and especially for furnaces for heating iron rods for the manufacture of bolts and rivets. From theoretical considerations, it would appear to be eminently adapted to the purpose, I lb. of coal-tar creosote, when effectively burnt, being capable of evaporatmg 11 24 lbs of water, a result which is considerably in excess of that obtained from coal or coke

Various methods have been devised for burning the oil to the greatest advantage These may be briefly summarised under the · following heads :-

1 Injection into the furnace by means of compressed air with 'atomising' apparatus.

2 Injection into the furnace by means of steam with 'atomising' apparatus.

3 Feeding into the furnace by simple gravitation, alone or in combination with coke.

G F Dexter (Journ Gas Lighting, June 29, 1886, 1242-1244), in a table showing the relative calorific values of certain coal-tar distillates, refers to creosote, or heavy oils, as under .

Average formula C<sub>14</sub>H<sub>16</sub>, bp 240° to 270°

and upwards.

Proportionate weight of constituents:

Carbon			0 93913
$\mathbf{Hydrogen}$	•		0 06087

Calorific value:

Carbon units		497
Hydrogen units		210

Employment of creosote oil as an antiseptic and for the purposes of a cattle wash and sheep

dip (v. DISINFECTANTS).

Creosote oil is also now largely used to absorb benzene from coke-oven gas. The gas, after the extraction of the ammonia, is passed up towers or scrubbers down which a shower of creosote The creosote oil takes up from 3 to 4 p c of benzene, and this is regained by distillation with steam, the creosote being used over and over again until, by the accumulation of dirt and tar, it becomes too viscid, when it is generally run into the tar well. This fact should not be lost sight of when estimating the proportion of creosote in coke-oven tars.

The heavier portions of creosote oil are also used to absorb naphthalene from coal gas; the gas is passed through washers containing the oil. As the oil tends to abstract some of the illuminants from the gas, it is best to add to the oil before use a small proportion of benzene sufficient to saturate the oil so that no more is absorbed from the gas.

The heavier oils are also used as a rough lubricant for brick-making machinery, and for of the bricks to impart a superficial blue colour to them. The creosote acts as a reducing agent on the iron compounds in the clay, and the red colour is changed to blue

A preparation of heavy creosote or anthracene oil with lime is used as a cart or axle

Bibliography - Lunge's Coal Tar and Ammonia; Allen's Commercial Organic Analysis: Warne's Coal Tar Distillation. W H. C.

CREOSOTAL. Syn for Creosol carbonate.

CRESATIN. Acetic ester of m-cresol CRESEGOL. Trade name for the mercury-

potassium salt of cresol sulphonic acid CRESOL C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)OH. The three isomeric cresols exist in coal tar (Williamson and Fairlie, Annalen, 92, 319, Ihle, J pr. Chem [2] 14, 442, Tiemann and Schotten, Ber. 11, 767, 783) in about the following proportions. orthocresol,  $35~\mathrm{pc}$ ; metacresol,  $40~\mathrm{pc}$ , and paracresol, 25p.c (Schulze, *ibid*. 20, 410).

The cresols can be separated from the oily,

non-crystallisable mother liquors obtained in the purification of crude phenol by fractionation and crystallisation, by dissolving them in soda lye, freeing the solution from naphthalene or other hydrocarbons by a current of steam, and then fractionally piecipitating with sul-phuric acid. The first fractions, owing to the greater acidity of phenol, contain chiefly cresols, and are further purified by distillation (Muller,

Zeitsch Chem. 1865, 270)

Various processes have been patented for obtaining meta- and para-cresol from crude cresol Raschig (Eng Pat 18334; D. R. P. 112545; J. Soc Chem. Ind. 1900, 37), by repeated fractionation of the crude cresol, obtains a mixture containing 60 p c of meta- and 40 p c This mixture is then sulphonated of para-cresol by means of concentrated sulphuric acid, in which m-cresolsulphonic acid is easily soluble, but the p-compound only with great difficulty. The mixture is allowed to stand for a week, and the crystals of the p-sulphonic acid are removed, or a soluble sodium salt, such as the sulphate, may be added in quantity sufficient to form the sodium salt of the p-cresolsulphonic acid. The latter is separated from the mother liquor, which consists of m-cresolsulphonic acid and a little sodium sulphate The two cresols are obtained from their respective sulphonic acids by hydrolysis with superheated steam. In another process (Raschig, Eng Pat 25269, J Soc Chem. Ind 1900, 1099) the mixture of the two sulphonic acids is subjected to the action of superheated steam at 120°-130°, the mcresolsulphonic acid alone being hydrolysed under these conditions The m-cresol distils off, and on raising the temperature to 140°-160°, the p-sulphonic acid is hydrolysed and p-cresol is obtained. See also Raschig, D. R. PP. 144903, 146752.

According to the Chem Fabr. Ladenburg (D. R P 148703; Chem Soc Abstr. 1904, i. 312) separation of m- and p-cresols, by partial sulphonation, is incomplete The process recommended is to heat crude cresol with sodium hydrogen sulphate at 100°-110°, dissolve the product in water, when p-cresol separates out as an oily layer, the m-cresol alone being sulphonated. Superheated steam, at 180°, is 420 CRESOL

passed in to obtain *m*-cresol, after the *p*-cresol has been separated.

Another process of the same firm (Eng. Pat. 1881; Fr Pat 339880; D R P. 152652; J. Soc Chem Ind 1904, 759) consists in saturating the crude cresol with milk of lime, when the calcium salt of m-cresol, which is much less soluble than that of the p-cresol, separates out. This is filtered off, decomposed with acid, and relatively pure m-cresol is obtained. The crystals of the calcium salt, before decomposition with acid, may be washed with benzene to free them from the last traces of mother liquor Reihm (D. R. P. 53307; Frdl ii 9) separates m-cresol from p-cresol by means of their barium salts.

Rutgers (Fr Pat. 317512; D. R P. 137584 and 141421; J Soc Chem. Ind. 1902, 1410) treats the commercial mixture with anhydrous oxalic acid or an anhydrous acid oxalate By this means, the oxalic ester of p-cresol alone is formed and separates out. This ester is separated and decomposed by water into p-cresol and oxalic acid; the m-cresol is obtained from the mother liquors

Terrisse (D R P 281054) sulphonates the mixture with acid of 60° Bé, at diminished pressure, neutralises with sodium carbonate and drives off the p-crosol in a current of steam. The residual sodium salts are decomposed by sulphuric acid and the m-sulphonic acid then heated with superheated steam, see also Hoffmann Laroche and Co, Fr Pat 454534

Synthetical preparation of oitho- and paracresol -Pure ortho- and para-cresol can be obtained by fusing the potassium salts of the corresponding toluenesulphonic acids with alkali (Engelhardt and Latschinow, Zeitsch Chem 1869, 620), or by diazotising the corresponding For this purpose the toluidine (15 parts) is dissolved in sulphuric acid of sp gr 18 (15 parts) diluted with 500 parts of water, treated with an aqueous solution of potassium nitrite (12 parts), and the product gradually heated with steam. The resulting cresol is then distilled with steam, the distillate treated with soda lye, then with sulphuric acid, and the liberated cresol extracted with ether and finally distilled in a current of carbon dioxide (Tiemann and Schrotter; Ihle (loc cit ))

Orthocresol can also be obtained from carvacrol by treatment with phosphorus pentoxide (Kekulé, Ber 7, 1006). It is crystalline, sp gr 1 0512 at 15 5°, melts at 30° (T. and S; Ihle; Béhal and Choay, Compt rend 118, 1211); boils at 188 5° (B and C), 188° (T. and S) at 191° (corr) (Perkin, Chem Soc Trans 1896, 1182; Fox and Barker 191 8°), is soluble to the extent of 2 5 vols in 100 vols of water, and on fusion with caustic potash is converted into salicylic acid When intrated, it yields dimitroorthocresol (mp 86°-87°) (Nolting and Salis, Ber. 14, 987). Refractive index at 40° ( $\mu_D$ )

1.53719.

Metacresol is prepared by heating thymol (100 parts) with half its weight of phosphorus pentoxide for 10-12 hours until propylene ceases to be evolved, adding the product to 115-120 parts of fused alkali, and further heating the fused mixture for 5-10 minutes to decompose the cresyl phosphate formed. The fused mass is then dissolved in water, treated with ether to

remove any undecomposed cresyl phosphate and other impurities, acidified with hydrochloric acid, and the separated metacresol extracted with ether, and finally distilled in a current of carbon discussed (Figure 1). The second of the second

dioxide (E and L.; T and S).

Metacresol is also prepared from naphthalene 1 3 3'-, 1 3 6-, or 1 3 8- trisulphonic acids, from analogous hydroxy- or amino-naphthalene monoor di-sulphonic acids or from trihydroxynaphthalene, eg 1 3 3'- naphthalene trisulphonic acid together with half its weight of caustic soda and three-quarters of water, is heated for 5 hours at 280°. The product is poured into water, acidified, boiled to expel sulphurous acid, and filtered m-cresol separates out on cooling (Kalle & Co Eng Pat 16559; D R P. 81484; J Soc Chem Ind 1895, 1800; Hoffmann La Roche and Co D. R. P. 247272) It is ordinarily a liquid of sp gr 1 0387 at 15 5°, but can be obtained in crystals, melting at 2 4° (Fox and Barker); it boils at 202° (corr) (P), and is soluble to the extent of 0.53 vol in 100 vols, of water Its aqueous solution is coloured blue by ferric chloude When fused with caustic potash, it is converted into metahydroxybenzoic acid, and on nitiation it yields a trinitro- derivative (m p  $105^{\circ}-106^{\circ}$ ) (Nolting and Salis, Ber 14, 987, 15, 1858) Refractive index at  $40^{\circ}$  ( $\mu_{\rm D}$ )

Paracresol crystallises in prisms, melts at 36° (Barth, Annalen, 154, 358 [B. and C]), 36°-37° (T. and S), and boils at 198° (I.), at 198°-199° (T. and S, B and C), at 202° (corr) (P); and is soluble to the extent of 1 8 vols in 100 vols. of water—Its aqueous solution gives a blue colour with ferric chloride—When fused with caustic potash, it is converted into parahydroxybenzoic acid, and on nitration it yields a dinitro-derivative (mp 85°) (Richter, Annalen, 230, 313;

Nolting and Salis, lc)

The crude cresol, as obtained from coal tar, is only slightly soluble in water, but its solubility in an aqueous solution of potassium cresol, renders its employment as an antiseptic possible. This solution is named solveol (Lacroix-Hunkiabéyendian, J. Pharm. Chim. 28, 34). The crude cresol is also frequently mixed with soaps, and is used in that form for disinfecting purposes (Dammann, D. R. P. 52129; Frdl. i. 538, Holmers, D. R. P. 76133 and 80260, Frdl. iv. 1119 and 1120; Raschig, D. R. P. 87275; Frdl. iv. 1121; Engler and Dieckhoff, Arch. Pharm. 230, 561; 232, 351).

Oxidation with peroxides at a high temperature converts the cresols into the corresponding hydroxybenzoic acids (Friedlander and Low-Beer, D R. P. 170230, J. Soc. Chem. Ind. 1907.

434).

A mixture of nitro- derivatives, obtained by sulphonating coal-tar cresol and nitrating the sodium salt of the resulting cresolsulphonic acid with nitre acid of sp gr 1·2, has been employed, under the names 'Victoria yellow,' 'saffron surrogate,' as a dye for wool and silk; and, although poisonous (compare Weyl, Ber. 20, 2835; 21, 512), has been used as a colouring matter for butter, cheese, &c. By heating crude cresol with sulphur and alkali sulphide, or with alkalipolysulphides, or with a mixture of sulphur and alkalis, at a high temperature, brown sulphur compounds are obtained which are used for dyeing cotton (Farb. vorm.

CROCOITE. 421

F. Bayer & Co., D. R. P. 102897; Chem. Zentr. 1899, n. 352

CRESOLIN. A disinfectant composed of crude phenol and rosin soap

CRESOLINE YELLOW v. Azo- colouring

CRESORCINOL v. PHENOL AND ITS HOMO-

CRESOSTERIL. Acid-o-oxalic ester of mcrosol

OH CO'H

CRESOTIC ACID CH3 A homologue of salicylic acid and obtained from o-cresol by similar methods to those employed in preparing salicylic acid from phenol; m p 163°-164°.

CRESYLIC ACID. The non-crystallisable, higher boiling portion of the phenols obtained in the purification of phenol (q v.) is, after redistillation, known commercially as cresylic acid or liquid carbolic acid. It is a colourless, only, refractive liquid, does not solidify at -- 80°, has a sp gr 1 044, and consists of 40 p c metacresol, 35 p c orthocresol, and 25 p c paracresol (Schulze, Ber. 1887, 410). It is much less soluble than carbolic acid in water, ammonia, glycerol, or caustic soda, but more readily so in petroleum spirit It boils at a higher temperature (about 185°-203°), gives a mixture of m-tribromo and o- and p-dibromo derivatives with bromine. It is even more readily acted upon by concentrated sulphuric and nitric acids than carbolic acid (Lunge, Coal Tar and Ammonia, ed 1900, 182, v also Allen, J Soc Chem Ind 1890, 141). Its antiseptic properties are more marked than those of carbolic acid or of either Cf CRESOL. of the three cresols

For a method of estimating phenol in com-mercial cresylic acid, see Fox and Barker (J. Soc.

Chem Ind 1917, 842, 1918, 265 T CRESYL BLUE v. OXAZINE COLOURING MAT-TERS.

CRETAFORM. Oxymethyl cresol tannin. CRISTOBALITE. A native form of silica (SiO<sub>2</sub>) crystallising in the cubic system (pseudocubic) It was first recognised by G. vom Rath in 1886 as regular octahedra up to 2 mm. across, associated with tridymite, in cavities in andesite from the Cerro de San Cristóbal, near Pachuca in Mexico It has since been found in volcanic rocks in the Rhenish district, Central France, and California, and in certain meteorites. The milk-white, translucent crystals are twinned and exhibit optical anomalies, these disappear suddenly at a temperature of 175°, the crystals being then truly cubic; on cooling, the birefringence reappears Sp gr. 234, hardness 6-7, refractive index 149, birefringence 0 0005.

Quartz when heated passes into \$\beta\$-tridymite at 870°, and this in turn passes into  $\beta$ -cristobalite (or metacristobalite, the cubic, optically isotropic modification) at 1470°. On cooling, the latter passes into the birefringent a-cristobalite at 180°-270°. The refractive index of this artificial cristobalite is about 1 484, being slightly higher than that of tridymite (1.477), m.p 1710°. Cubic crystals of silica have also been obtained artificially by heating an aqueous solution of colloidal silica with hydrofluoboric acid at 200° under a pressure of 26 atmospheres. Cristobalite has also been detected in silica bricks. L. J. S.

CRITH. The weight of a litre of hydrogen at 0° and 760 mm pressure—viz 0 0896 gram. CROCEIN, BRILLIANT, ORANGE, SCAR-

LETS, v Azo- colouring matters.

CROCEINE ACID, 2-Naphthol-8-sulphonic acid Used in making croceine scarlet and other dyes.

**CROCIDOLITE.** (Krokydolith, Ger) mineral of the soda-amphibole group, consisting of sodium and non (ferric and ferrous) silicate, NaFe'''(SiO<sub>3</sub>)<sub>2</sub> Fe"SiO<sub>3</sub>. It is known only in a finely fibrous form, its name, in fact, being from κροκύς, κροκύδος, wool. In the trade it is known as blue asbestos, and it is worked for the same purposes as the other varieties of asbestos It is, however, more readily fusible (to a black magnetic glass) than are tremoliteasbestos and scrpentine-asbestos The mineral has been long known from the Asbestos Mountains near the Orange River in Griqualand West, South Africa, where it occurs abundantly as veins, 1 or 2 inches in thickness, in jasperschists The closely aggregated parallel fibres are arranged perpendicularly or nearly so to the walls of the veins, and the material is extracted as slabs It is now extensively mined for commercial purposes (H. F. Olds, Trans. Inst. Mining and Metall. 1899, vii. 122, G. E. B. Frood, The Cape Asbestos Industry, Ann Rep. Govt. Mining. Engineer, Dept. of Mines, S Africa, 1915, and S. African Mining Journal, 1916).

Being an alkali silicate with ferrous iron, crocidolite is especially liable to decomposition when exposed to weathering, sodium is removed, and the iron oxidised and hydrated to form limonite, while the silica is set free. There then results a ferruginous quartz possessing the finely fibrous structure of the original mineral, or, in other words, a pseudomorph of quartz and limonite after crocidolite. This material is coloured a nich golden yellow, and displays a silky lustre, and being at the same time very hard, it is well adapted for ornamental purposes When cut and polished with a convex surface, it displays a cat's-eye effect, and it is extensively used for making knobs, umbrella handles, beads, Such material is known as tiger-eye, though, unfortunately, in the trade the name crocidolite is very frequently misapplied to it. The name pseudo-crocudolite would be more appropriate. In some stones there has been silicification without oxidation, and the indigoblue colour of the original crocidolite has been preserved · material of this kind is known as hawk's-euc The stones have sometimes been coloured artificially by the same processes as adopted in the staining of agate (q v)

L J. S. CROCOITE or Crocoisite. Lead chromate, PbCrO<sub>4</sub>, crystallising in the monoclinic system. The name was originally given in the French form crocoise by F. S Beudant in 1832, from κρόκος, saffron, this was later changed to croconsite, and afterwards to the more correct form crocoite. This mineral is the only chromate of any importance found in nature, and in it the element chromium was discovered. It is found usually as crystals which are bright hyacinthred in colour with a brilliant lustre, the streak is orange-yellow. On exposure to light the crystals soon deteriorate in translucency and

brilliancy Sp gr 60, H 2½-3 Crocoite is a mineral of secondary origin in veins of lead ore and is often associated with gold. It is found at Beresovsk in the Urals (where it was discovered in 1766), Congonhas do Campo near Ouro Preto in Brazil, Luzon in the Philippines, Umtali in Rhodesia, and as very fine crystals at Dundas in Tasmania The mineral has been used as a pigment, being identical in composition with the artificial product chrome-yellow

### CRONSTEDTITE v THURINGITE

CROOKESITE. A selenide of copper, thallium (Tl 16-18 p c ), and silver forming compact masses with lead grey colour and metallic lustre Sp gr 69, H 21-3 It occurs embedded in calcite in the Skrikerum mine, Smaland, Sweden Although a rare mineral it is of interest in being I tions

one of the few minerals containing thallium in

considerable quantity. L. J. S. CROPS. The composition of crops may be regarded from two points of view: (1) the materials they contain, which render them valuable either as food or as sources of some substance of commercial value, (2) the materials they remove from the soil, which must eventually be replaced if the fertility of the land is to be maintained

In Table I, the composition of some of the principal crops grown in the United Kingdom is set out, cereals being excluded because they are dealt with separately The composition of each of these materials shows considerable variations, and the analyses given are either typical or the means of a small number of determina-

TABLE I -- PROXIMATE COMPOSITION OF CROPS.

	Water	Crude fat	Protein crude	Carbo- hydrates	Crude fibre	Ash
Leguminous .	1			1		
Field beans	. 13 49	1 68	25:31	48 33	8 06	3 13
,, ,, haulm , ,	. 18 40	1 10	8 10	31 00	36 00	5 40
Field peas	. 13 92	1 89	$23 \ 13$	52 68	5 68	2 68
", haulm	13 60	1 60	9 00	33 70	35 50	6 60
Clover hay	. 16 50	2 90	13 50	37 10	24 00	6 00
Luceine hay	16 50	2 60	14 20	29 20	29 50	8 00
Vetches, green	. + 82 50	0 50	3 20	7 20	5 10	1 50
Root crops .	.   02 00	0.00	0 = 0		0.0	
Potatoes .	74 98	0.15	2 08	21 01	0 69	1 09
Mangolds	87 50	0 14	1 34	8 90	0 98	1 14
Sugar beet .	75 00	0 10	1 30	21 40	1 50	0.70
Swede turmps .	87 80	0 21	1.54	8 22	1 32	0 91
White turnips	. 90 78	0 22	1 18	5 89	1 13	0.80
Fosterton hybrid turnips 1	91 30		0 79	6 47	0 94	0.50
Aberdeen yellow 2 .	91 09	i _ 1	1 14	6 08	1 03	0 66
Carrots	86 79	0 30	1 23	9 17	1 49	1 02
Meadow hav	14 30	2 50	9 70	41 40	26 30	6 20
Seeds, hay	. 14 30	2 70	10 20	36 10	30 20	6 50
Mustard, green	85 10	0 40	$\frac{10}{2} \frac{20}{90}$	7 30	2 90	140

[4, Leguminous crops. The leguminous seeds are distinguished by their high protein content. The proteins of the bean and pea and similar seeds have been studied by Ritthausen (Eiweisskorper der Getreidearten, Bonn, 1872) and Osborne (J. Amer Chem. Soc. 1898, 20, 410, Amer. J. Physiol. (1908, 23, 180, &c.) The most abundant of these is the so-called legumin (q v), which, according to Osborne, is a globulin con-Lentils contain a conglutin taining lysine which has been hydrolysed, also some edestin

The leguminous seeds grown in this country contain but a small proportion of oil, which has not received any special examination, the soya bean, however, yields oil up to 20 p c, and is extensively imported from Manchuria and pressed for oil (v. Soya BEAN).

The only carbohydrate in beans, peas, &c, that has received any attention, is the starch, which is abundant and the granules of which possess characteristic shapes The ash of leguminous plants is distinguished by a large proportion of lime, as may be seen from analyses 5-10 in Table II.

Root crops. In these materials, the ether extract contains little, if any, true fat, the mtrogen-containing substances are also very far from being all protein, usually one-half or more of the nitiogen present is combined in various non-protein forms

Potatoes -The characteristic carbohydrate is starch, of which there may be over 20 pc; but sugars (both sucrose and reducing sugars) may be present up to 1 p c, and dextrins or amylans are also present in about the same amount Amongst the nitrogen compounds there are both insoluble and soluble proteins, part of the latter being coagulable by heat or acids, and a large range of amino- bodies, among which tyrosin The non-protein nitrogen has been identified may amount to more than half the total nitro-The darkening of the juice of a fresh potato is due to the black substance formed by the oxidation of the tyrosine under the action of an enzyme, also present in the cells darkening of boiled potatoes is due to some other cause (Ashby, J. Agric Sci. 1905, i. 348). About two-thirds of the ash of potatoes consist of phosphoric acid and potash.

Mangolds.—The characteristic carbohydrate

Aitken, Trans. H & Ag. Soc. 1884, 16.
 Wilson, ibid 1886, 18.

CROPS. 423

present in the mangold is sucrose, which constitutes about 8 p.c of the weight of the root, the total dry matter amounting to about 13 p c. About 0.25 p c of reducing sugar is also present. The variation in the sugar content of mangolds has been examined by Wood (J. Agric. Sci. 1905, 1, 176), who gives the following table.—

Variety	Yield per acre—tons	Dry matter p c	Sugar p c.	Nitrogen p c
White-fleshed globe . , , intermediate Yellow-fleshed tankard . , , , globe . Long red	29 9	10·7	63	0 165
	27 4	12 0	71	0 168
	24 6	13·1	80	0 186
	25 0	13·4	82	0 191
	29 9	13 1	79	0 157

Of the nitrogen in mangolds, only 30-50 p c. is in the protein form, 20-40 p c. is combined as amido or amino-acid and is liberated on hydrolysis as ammonia, while 10 p c. may be in the nitric form

On storage, the mangold loses sugar by respiration, and some of the sucrose becomes changed into reducing sugar, a process which is much accelerated as soon as the temperature rises in the spring (see Miller, J. Roy. Agiic. Soc. 1900, 61, 1902, 63)

1900, 61, 1902, 63)

Sugar beet—In the sugar beet, which has been derived from the same species of Beta as

the mangold, the characteristic carbohydrate is likewise sucrose, which may be present to the extent of 20 p.c. It is always accompanied by a certain proportion of reducing sugars (glucose and maltose), which, in well-grown roots, should not exceed 1 p. c. of the sucrose.

The variations in the amount and nature of the sugars in sugar beet have been subjected to very detailed investigation, for accounts of which the numerous technical works dealing with the beet-sugar industry, should be consulted V Sugar.

The nitiogenous constituents resemble those of the mangold (see Lippmann, Ber 1896, 29, 2645)

Swedes and turnips—In these closely allied roots, the carbohydrates consist of sugars (chiefly reducing sugars, though few data exist) and pectic bodies yielding a mucilage on boiling with water In Wilson's analyses, already quoted, the sugars averaged 4.72 p.c. nitrogen compounds are also largely non-protein, only from 30 to 50 p c. consists of protein. It is generally considered that swedes and turnips grown in the north and especially in the east of Scotland, possess a higher feeding value than the roots grown in the south, analytical data to confirm this opinion are, however, lacking, though it is certain that much larger crops are usually grown in the cooler climate of the north, where the crop can be sown earlier with success.

Ash of crops. In Table II are collected various analyses of the ash of the crops above mentioned

TABLE II -COMPOSITION OF THE ASH OF FARM CROPS

										No. 25			
	-	Ash in dry matter	N m dry matter	Fe <sub>2</sub> O <sub>3</sub>	CaO	MgO	K <sub>2</sub> O	Na <sub>2</sub> O	P2O5	SO <sub>3</sub>	CI	(O <sub>2</sub>	SiO <sub>2</sub>
2 3 4 5 6 7. 8 9 10 11. 12.	Potatoes Mangolds Swedes Grass Clover Beans, corn straw Peas, corn straw Lucerne Flax, seed stem	3·99 6·70 5·58 7·21 8·15 3·27 5·80 2·53 5·87 8·71 3·82 3·70	0 88 1 31 2 · 31 1 65 2 92 4 32 0 94 3 87	0 35 0 44 0·61 0·49 1 03 0 48 1 54 0 07 1 03 2 68 1 54 6·62	1 53 3 58 12 12 12 48 37 37 5 60 28 39 5 74 40 34 24 74 10 96 25 16	3 48 2 13 3 02 3 83 5 10 6 61 3 12 8 97 8 30 6 27 11 47 4 99	35 90 39 17 29 76 13 10 40 58 8 07 44 31 11 78 41 91 27 85 25 56	0 06 19 34 5 24 4 09 0 58 2 12 14 57 0 39 9 90 1 22 4 15 10 16	12 86 9 34 9 83 7 04 3 75 30 69 2 31 29 30 8 26 7 71 41 27 8 94	5 87 3 25 13 49 6 17 1 82 6 35 3 30 10 42 6 76 5 63 2 01 4 02	7 27 3 87 2 48 8 78 2 26 2 43 2 49 0 45 3 84 8 05 0 63 6 64	11 37 21 43 12 43 10 31 34 96 5 28 33 68	0 58 0 97 0 72 18 32 0 54 0 41 3 08 0 37 10 66 3 16 1 47 9 40
13	Hops, cones			1 80	16.77	5 66	28 35	1 99	13 73	5 01	4 35		22 32
14	" leaves			0 64	40.17	5 32	13 65	3 79	7 82	3 49	4 02		22 21
15	", stems	4 70		1 07	29 59	8 34	26.18	5 06	11 23	3 41	8 87	-	8 29
16.	Tobacco .	16 27		2 93	42 34	11 22	17 48	4 26	3.22	4 28	4 87		10.11
ļ									<u> </u>		<u> </u>		

Analyses 8-16 are means taken from Wolff's Aschen-analysen, and have been recalculated to exclude the carbon dioxide present in the ash, as obtained after burning Analyses 1-7 are selected from the Rothamsted records.

Materials removed from the land by average crops. Table III, compiled by Warington from Rothamsted data, shows the amount of the three important constituents—nitrogen, phosphoric acid, and potash, which would be contained in an average crop of the magnitude indicated, and therefore have been removed

from the soil by the crop, except in the case of the nitrogen contained in leguminous crops

A comparison of these figures with the amount of fertilising material usually provided for the crops in question, or the effects of particular manures in field experiments, serves to show that the data contained in Table III. afford no guidance in deciding the appropriate manure for each crop.

The earliest theory of plant manuring, that of Liebig, supposed that the requirements of the plant can best be satisfied by applying to

424 CROPS

TABLE III.—AMOUNT OF FERFILISING CONSTITUENTS PER AGRE REMOVED BY AN AVERAGE CROP

	Weight of crop, dry	Nitrogen	Potash	Phosphoric acid
Wheat, 30 bushels, straw 28 cwts.  Barley, 40 ,, ,, 22 ,, . Oats, 45 ,, ,, 25 ,, . Maize, 30 ,, ,, 19 ,, . Meadow hay, 1½ tons .  Red clover hay, 2 tons .  Beans, 30 bushels, straw 20 cwts Turnips, 17 tons, leaf 5 tons .  Swedes, 14 ,, ,, 2 , . Mangolds, 22 ,, ,, 8 ,, .	10s 4183 3827 3978 3377 2822 3763 3461 4657 4055 7568	1bs 50 49 52 43 49 98 1 107 1 110 98	1bs 28 8 35 7 46 1 36 3 50 9 83 4 67·1 148 8 79·7 300 7	1 bs 21 1 20 7 19 4 18 0 12 3 24 9 29 1 33:1 21:7 552 9
Potatoes, 6 ,, tubers	3360	46	76 5	21.5

the land exactly the constituents taken away by a good crop of that particular kind; but the results of the Rothamsted experiments, confirmed by general farming experience, very soon showed that this view was incorrect, because it failed to take into account the enormous amount of plant food contained in the soil, more or less of which would reach the plant according to the method of cultivation and the plant's habit of Moreover, in practical farming the growth supply of manure is largely determined by the place the crop occupies in the rotation and the treatment which has been given to the previous crops. To take an example, wheat and barley withdraw almost the same fertilising materials from the soil, yet all farming experience goes to show that the yield of wheat is chiefly determined by the supply of nitrogen If this element is present in sufficient quantity the wheat plant with its long period of growth is usually able to satisfy its requirements for phosphoric acid and potash from the soil barley, nitrogen is still a most important element in nutrition, but phosphoric acid also exerts a striking beneficial action and is a necessary element in the fertiliser Neither for wheat nor barley need potash be supplied except on very light soils A still more noteworthy example is The table provided by the swede turnip crop shows that an average crop will take away from the soil about 100 lbs of nitrogen, 20 lbs of phosphoric acid and 80 lbs. of potash, yet the usual fertilisers applied to swedes consist almost wholly of phosphatic compounds, with a comparatively small amount of nitrogen and no potash. The majority of the swedes grown in this country receive from 50 to 100 lbs. of phosphoric acid (i.e. about 5 cwt. of superphosphate or basic slag), and the small quantity of nitrogen that would be provided by a light dressing of farmyard manuse or half a hundredweight of sulphate of ammonia. The fertiliser requirements of a crop are in fact conditioned by its specific habit and by the soil (q v) rather than by its composition Certain broad characters may be distinguished. Cereal crops possess considerable powers of obtaining the phosphoric acid and potash reserves in the soil provided they are well supplied with active forms of nitrogen. Leguminous crops rarely require a nitrogenous fertiliser, but respond to mineral fertilisers and particularly to potash, the soil also should be neutral and well supplied with lime. Cruciferous crops are particularly dependent upon an abundant supply of phosphoric acid. Crops elaborating a considerable amount of carbohydrate, e.g. mangolds and potatoes, require potash. In general it may be said that mitrogen promotes leafy development, phosphoric acid maturity, and potash carbohydrate formation and continued growth.

A. D. H.

CROTON OIL is obtained from the seeds of Croton Tiglium (Linn), a tree belonging to the Euphorbiacew, indigenous to the Malabar coast, and cultivated in Southern Asia and China. The seeds contain 53-56 p c of oil

The oil has an amber-yellow, orange, or brown colour, according to its age, it has a nauseous odour, a burning taste, and acts as a very powerful purgative. It blisters the skin and mucous membrane

The chemical composition of croton oil differs so widely from that of all other oils, that its recognition, by means of the 'quantitative reactions' (see Oils and Fats), is easy It contains a considerable amount of volatile acids, hence its Reichert-Meissl value is about 13, its iodine value is 102–106 Saponification value, 210–215, and specific gravity, 0 940 to 0 960. Croton oil is strongly dextro-rotatory (+14 5° to 16.4°).

The variations in the recorded values are mainly attributable to different methods of extraction. Croton oil contains the glycerides of tighe acid, and of butyric, caproic, lauric, myristic and palmitic acids, and probably also olice acid. The vesicating principle is a resinous substance, C<sub>11</sub>H<sub>18</sub>O<sub>4</sub> (Dunstan and Boole, Proc. Roy Soc. 1895, 58, 238).

CROWBERRY (Empetrum nigrum). The

CROWBERRY (Empetrum nigrum). The leaves of the crowberry contain a wax composed of ceryl cerotate, benzoic acid, urson, tannin, dextrose and probably rutin. The anhydrous urson C<sub>10</sub>H<sub>16</sub>O is identical with that obtained from the leaves of Uva Ursi. Empetrum nigrum contains no alkaloids, andromedextrins or glucosides decomposed by emulsion (Van Itallie, Pharm Weekblad, 1918, 55, 709)

CROWN GLASS v. GLASS.

crucibles. A perfect crucible should withstand the highest temperatures without fusion, should bear sudden changes of temperature without fracture, and should be unacted upon by the materials which are to be heated in it. A semi-automatic apparatus for moulding crucibles has been patented by Sims (Eng. Pat. 445, 1888).

1 Wholly or in part derived from the atmosphere,

Clay crucibles are prepared from selected clay; of the Stourbridge clay worked, only about 5 p.c. is available for this purpose. Its plasticity is much increased by exposure for some months in a moist condition to the air. During this process of 'weathering' it becomes disintegrated, and the removal of foreign matter is thereby facilitated. The presence of non pyrites is especially injurious; furnace gases convert it into the oxide which acts as a flux and forms cavities in the crucible Lime is also hurtful, except when in very small quantities and uniformly distributed. Potash and soda are invariably present in clay, and when in small quantity increase the coherence of the crucible without appreciably impairing its power of withstanding the action of heat.

In order to reduce the tendency to crack, and to increase its refractory character, the clay is usually mixed with sand, ground flints, burnt clay (old broken crucibles), graphite or coke These, which are either less fusible than the clay or altogether infusible, form a skeleton which upholds and prevents deformation of the crucibles at temperatures at which the clay would tend to fuse The most refractory crucibles contain the largest proportion of silica.

The clay, which should be of the greatest possible fineness, is well beaten to remove air bubbles and mixed with the proper quantity of the refractory material. According to Berthier (Traité des Essais, 1, 66), when silica is used, it should not be too fine, as the tendency to combine with the clay and thus increase the fusibility would thereby be increased. Crueibles of fine grain are also more liable to fracture.

Small crucibles are usually baked in a kiln, but the large casting pots of Stourbridge clay are not baked until immediately before use When the temperature of the baking is high, the crucible is more liable to fracture on reheating

All clay crucibles are more or less acted upon by litharge, but metals and most ores may be heated in them without danger. Berthier recommends a lining of such material as silica, alumina, or magnesia previously moistened. A lining of chalk renders a crucible less permeable to litharge.

Queneau recommends lining clay crucibles, before drying, with some substance which will resist the chemical reactions for which they are designed and then compressing strongly, drying, and firing (Fr Pat 354319, J. Soc. Chem. Ind. 1905, 1071).

Ciucibles may be rendered non-porous by treating with ferious sulphate solution and firing (Mitchell, U.S. Pat. 894845, J. Soc. Chem. Ind 1908, 886) or by the addition to the clay of chromic oxide, chromites or chrome iron ore, which also renders them much more refractory (Bach, Eng. Pat. 24041, 1903, Klein, U.S. Pat. 787770, Seigle, Fr. Pat. 383593; and Placet, Eng. Pat. 28728, 1896)

The fracture of ciucibles by expansion of

The fracture of ciucibles by expansion of solidifying metal can be prevented by liming with finable material (Colby, U.S. Pat. 830208, J. Soc Chem. Ind. 1906, 934)

Goldschmidt recommends that crueibles for the 'Thermit' process should be lined with a mixture of magnesia or carbotundum and soluble glass (Fi Pat 354597, J Soc Chem Ind 1905, 1114)

Very large crucibles may be built up from cylinders or annular rings of refractory material, provided arrangements are made for tapping the contents without moving the crucible (Reynolds, Eng. Pat. 1004, 1901, Rousseau, Eng. Pat. 7451, 1902)

Stourbridge clay crucibles are made from a mixture of raw ground clay with about one-half its weight of powdered, sifted, old glass pots, from which all adherent glass has been removed, the mixture being made in a disc mill, and kneaded to the proper consistency. Percy gives the composition of the Stourbridge clay used for the best crucibles as silica, 79.25, alumina, 13.57, inon oxide, 1.38, lime, 0.08, magnesia, 0.42, water, 5.11

The following analyses show the composition of certain varieties of cureibles. Nos. 1-7 are by Berthier, No. 8 by Dick, Nos. 9-12 by Brand, and Nos. 13-15 by Mène.—

	Silica	Alu- nuna	Iron oxide	Mag- nesia	Lime	Potash	Sul- phur	Carbon	Water
1. French, made by Beaufay at Paris 2. French, made by Deyeux	64 6	34 4	10	-			<b>Managara</b>	_	
at Savignies	72 3	19 5	3 9		-				1.8
3 Hessian	70 9	24 8	38	trace					
4 English steel .	71 0	23 0	40					I	
5 St. Etienne steel .	$65\ 2$	25 0	72	trace	-			-	
6 Glass pots, Nemours	67.4	32 0	08	trace	· —				-
7 ,, ,, Bohemia .	68 0	29 0	2 2	0 50	-				
8. Cornish, made by Juleff	$72\ 39$	25 32	1 07	trace	0.38	1 14			
9. Steel crucible; clay .	53 92	40 57				-	0.230		
10. " " coke clay	42.78	34.71			-		0.490	18 60	
11. " " graphite.	24 63	27.89					0.270	40 43	
12 Magnesia	4 80	249		92.62			0.099		
13 English blacklead	51 4	$22\ 0$	35		02			20 0	18
14. Ditto	45 1	16 65	0 95					34 5	25
15. Ditto	50 0	20 0	15	•	05			25 5	30

426 CRUCIBLES.

fracture on sudden change of temperature, but will withstand a very high degree of heat

French crucibles are made from a mixture of 1 part raw Ardennes clay and 2 parts of the same which has been burnt. They are refractory and of fine texture

Cornish crucibles are largely used for copper assaying. They are round, of coarse grain, and of a mottled greyish-white colour. They are kiln-buint, and withstand sudden changes of temperature, but cannot be heated to whiteness Cornish crucibles are now made by the Morgan Crucible Company at Battersea.

London crucibles are of reddish-brown colour and close texture—They will not stand sudden change of temperature, but withstand the action

of litharge fauly well

Silica crueibles may be made of powdered silica, mixed with a highly absorbent substance, e g kieselguhr, saturated with water (Sankey, Eng Pat 3353, 1906, J Soc Chem. Ind 1907, 96) Sand, mixed with a small proportion of cement and water, moulded, dried, and heated to the fusing-point of the sand, forms a very refractory material (Reppey, U S Pat 905295, J Soc Chem Ind 1909, 24) A mixture of sand with a little lime may be used (Berglund, Fr Pat. 366830, J Soc Chem Ind 1906, 1081).

White fluxing pots are made by the Morgan Company from a highly refractory freelay from Rouen (Truchles contaming carbon are now largely used For very high temperatures, crucibles cut from a block of gas carbon or similar material may be employed Clay crucibles may also be filled with a moist mixture of charcoal with starch, treacle, gumwater, or oil, or, for large crucibles, gas tai. The mass is forced with considerable pressure into the crucible and is then cut out, leaving a coating of about \( \frac{1}{6} \) inch in thickness, of the coating is put into the crucible and smoothed with a glass rod. The crucible is then filled with powdered charcoal and heated to redness

Graphite crucibles. For the fusion of metals, plumbago or coke crucibles are of great value, but they are not largely used for assaying. At the Battersea works, the following process of manufacture is employed hand-picked Ceylon plumbago is ground, sifted, digested with dilute sulphuric acid to remove the greater part of the iron, and mixed and kneaded to a stiff paste, with about an equal weight of fine fireclay. The mixture is passed through a pug-mill, and is cut into blocks When required for use, the blocks are again passed through the mill, and made by hand on a wheel or moulded by machinery. They are then dried and baked in saggars These ciucibles may be protected from oxidation by lining them with clay (Woolford, Eng. Pat. 24479, 1898, J. Soc. Chem. Ind. 1900, 52, McDonald, U.S. Pat 828954; J. Soc Chem Ind. 1906, 911)

They are of fine grain, and will withstand the highest temperatures and repeated sudden changes. The graphite is slowly oxidised by metallic oxides or by the furnace gases. To minimise this, the crucible may be coated by dipping in a paste of clay and solution of borax. Any iron which is present in the graphite is liable to become introduced into the melt, and the carbon itself combines with iron or steel (v. Brand, Chem. Zentr. 15, 407; and Dingl. poly. J. 256, 227)

Tar, rosin, &c, may be used to bind the clay and graphite in manufacturing these crucibles (Putz, D R P. 181979; J. Soc. Chem. Ind. 1907, 1030) Digby recommends (Eng. Pat. 3661, 1893) the following composition.—

 Plumbago, 17-mesh granules
 17 parts

 Plumbago, 80 ,
 ,
 2
 ,

 Russian clay
 .
 .
 5
 ,

 China-stone
 .
 .
 5
 ,

 Gas carbon
 .
 .
 2
 ,

The last three ingredients are ground and mixed with water and then the plumbago is added and the crucibles formed, dried at 65° and baked

Salamander crucibles, made by the Morgan Company, contain a much smaller proportion of carbon They withstand sudden changes of temperature perfectly

Lime crucibles are cut from blocks of well-buint (preferably slightly hydraulic) lime.

Magnesia was introduced by Caron as a refractory basic material for making crucibles or for liming reverberatory furnaces for melting and dephosphorising steel Magnesia contracts considerably when heated, it is therefore necessary, before making it into bricks or crucibles, to expose it to a higher temperature than that to which it will be heated. The strongly calcined magnesia is made into a paste with 15-30 pc of raw (slightly calcined) magnesia, and 10-15 p c of water, and is moulded, dried, and baked at a red heat. The highly heated magnesia would not cohere without admixture with the law material Magnesia crucibles are not injured by exposure to the atmosphere Gaudin makes crucibles from a mixture of 1 part magnesia and 1 part bauxite They may also be formed from a mixture of calcined magnesite with from 2 p c to 12 p c of a boron compound (according to the amount of calcium present), moistened, moulded, dried and fired at a high temperature (Rawson and Littlefield, Eng. Pat. 16004, 1900, J. Soc. Chem. Ind. 1901, 992, and *ibid.* 1902, 52). The product is said to resist molten lithage. Porous crucibles for metal refining, are made from a similar mixture with the addition of sawdust (Dobell, Eng Pat 10715, 1897).

Alumina crucibles withstand sudden changes of temperature, and are not affected by the materials fused, even sodium being without action According to Deville's process (Ann. Chim Phys [3] 46, 195), alumina precipitated at low temperature from ammonia alum solution is mixed with an equal weight of powdered marble, and is exposed to the highest temperature of a wind furnace. The mixture so produced, which may be considered as consisting of calcium aluminate, is mixed with its own weight of moist, precipitated alumina, and is made into a crucible, dried and baked (see also Caspar, US. Pat 886111; J Soc Chem Ind. 1908. When the crucible is required to withstand higher temperatures, the amount of the calcium aluminate may be reduced, but the proportion of lime should not fall below 10 to 20 p c.

Bauxite has been used to replace alumina. It is extremely refractory (v. Bauxite, art. Aluminium).

Carborundum crucibles are infusible and

CRYOLITE. 427

resistant to chemical action and have great solidity. They may be made from a mixture of commercial carborundum with sufficient clay to render it plastic (Velter, Fr. Pat 376179; J. Soc Chem. Ind. 1907, 1010; Engels, Eng. Pat 14567, 1901)

A mixture of powdered silica rock or ganister with carborundum or amorphous silicon carbide, may be used. It is mixed with water-glass solution or glue, &c, compressed to shape and heated to the temperature of the electric furnace (Imray and Carborundum Co Eng Pat 19493, 1902; 3308, 1903) (see also Carborundum)

(See further, Kerl, Handb Thonvaaren Ind. 1879, 528-534; and Percy, Metallurgy (Fuel, &c), 1875, 110-141)

CRUMPSALL YELLOW v Azo- colouring

MATTERS

CRURIN. Trade name for quinoline bismuth thiocyanate, used as an antigonoirhoic CRYOGENIN. m - Benzaminsemicarbazide

an antipyretic.

CRYÖLITE.—(Kryolith, Ger) A native fluoride of aluminium and sodium, found almost exclusively in Greenland, whence it has been sometimes called 'Greenland spar.' The name cryolite, or 'ice-spar' (Eisstein) refers to the general resemblance of the nuneral to ice (κρύσς) and to its ready fusibility in a candle-flame The mineral was discovered by the Danes in 1794, and described in the following year by Schumacher Nearly all the cryolite of commerce is obtained from a single locality at Ivigtut (or Evigtok), on Arsuk-fjord in southwest Greenland This locality was visited by Gresecké, and later by J W Tayler (Quart Jour Geol Soc 12, 140) The cryolite occurs as a large deposit in gness, and contains (especially near its junction with the enclosing rock), a number of metallic minerals, such as galena, pyntes, and chalybite (v. P. Johnstrup, N. Jahrb f. Mm. 1886, 1, Ref. 28, R. Baldauf, Zeitsch. piakt Geol. 1910, xviii 432; O. B. Bøggild, Mineralogia Groelandica, Kjøbenhavn, 1905, O. B. Bøggild, Meddelelser om Grønland, 1912, 1. 107, and Zeitsch. Kryst. Mm. 1913, h. 591. C. P. Bannard, The Combine Min , 1913, h 591 C P Bernard, The Cryolite Mine at Ivigtut, Greenland, Mining Magazine, 1916, xiv 202)

Cryolite is usually found in cleavable translucent masses, of snow-white colour, with a peculiar moist-looking vitreous lustre which renders the mineral easy of recognition. H 25; sp gr. 29-3 Crystals are rare, and their relations have been the subject of much discussion, they are cube-like in habit, and belong to the monoclinic system. There are three perfect cleavages approximately at right angles to one another. Although cryolite is usually white, it is sometimes coloured brown or reddish, and is said to pass downwards, in the Greenland deposit, into a black mass. The black variety loses its colour when heated. Before the blow-pipe cryolite fuses to a white enamel

The composition of cryolite may be represented by the formula 3NaFAIF<sub>3</sub> or Na<sub>3</sub>AIF<sub>6</sub>. This corresponds to 12 8 p.c. Al, 32 8 Na, and 54 4 F. An analysis of the Greenland cryolite

by Chodnev yielded 13 23 p c Al, and 32 71 Na, with 0 83 of  $\rm Mn_2O_3$  and MgO (Verh Petersb. Min Ges. 1845–6, 209). Durnev found in the cryolite of Miask in the Urals, 13 41 p c. Al, 32 31 Na, 0 55 Fe $_2\rm O_3$  and Mn $_2\rm O_3$ , and 0 35 CaO (Pogg Ann. 83, 587) Deville found in the Greenland cryolite 0 018 p c of vanadic acid and a small proportion of phosphoric acid An analysis of cryolite from near Pike's Peak, Colorado, by Cross and Hillebrand yielded Al 12 9, Na 32 4, F 53 55, Fe $_2\rm O_3$ 0 4, Ca $_2\rm O_3$ 0 (Amer. J Sci 1883, xxvi 271)

Cryolite was formerly a rare mineral, but about 1855 it was introduced into Europe commercially, and at once found numerous applications in the arts About 10,000 tons are exported annually from Greenland. By heating finely powdered cryolite with sodium or with magnesium, J Percy and A Dick obtained metallic aluminium (Phil Mag 10, 1855, 364) Deville found that fused cryolite is decomposed by the passage of an electric current, with elimination of free aluminium. At the present time, aluminium is prepared by the electrolysis of bauxite, but the addition of some cryolite is necessary to start the process and to keep it in continuous operation (v. Aluminium)

A wide range of applications was suggested when Julius Thomsen showed in 1850 that cryolite is readily decomposed by heating with calcium carbonate, and that sodium aluminate may be thus obtained

 $Al_2F_6 6NaF + 6CaCO_3$   $= 3Na_2O \cdot Al_2O_3 + 6NaF_2 + 6CO_2$ 

If heated with quicklime, a similar decomposition is effected, of course without evolution of CO<sub>2</sub> The sodium aluminate has been used in the manufacture of alum, and in the preparation of caustic soda for soap making By lixiviation, the aluminate is easily dissolved out, and by the passage of CO2 through the solution, Na2CO3 is obtained, while alumina separates in a gela-The decomposition of cryolite tinous form may also be effected by boiling the finely pulverised mineral with milk of lime By adding an excess of powdered cryolite to the resulting solution the aluminate is decomposed, the alumina being precipitated, while sodium fluoride remains in solution.

$$3Na_2O Al_2O_3 + Al_2F_6 6NaF = 2Al_2O_3 + 12NaF$$

Cryolite can be readily decomposed by the action of sulphuric acid, with formation of sodium sulphate, which can be easily converted into carbonate, but the process, after trial at Copenhagen, has been abandoned in consequence of the inconvenience arising from the evolution of hydrofluoric acid.

 $Al_2F_6 6NaF + 6H_2SO_4 = Al_2(SO_4)_3 + 3Na_2SO_4 + 12HF$ 

The manufacture of soda from cryolite is carried on at Oersund in Denmark, and at Natrona, near Pittsburg in Pennyslvania. The mineral is also employed in the preparation of enamels on iron and for the manufacture of an opaline glass, somewhat resembling porcelain. This glass may be formed by fusing together 100 parts of silica, 35 or 36 of cryolite, and 13 or 14 of oxide of zinc. C. Weinreb has investigated the cause of the milkiness of cryolite glass, and believes that it is due to the presence

428 CRYOLITE.

of aluminium fluoride, which on the cooling of the fused mass, separates in a lighty communited condition, and being distributed through the glass renders it more or less opaque (Dingl poly J 256, 361).

L J S.

cryolithionite. Fluoride of lithium, sodium, and aluminium, Li<sub>3</sub>Na<sub>3</sub>Al<sub>2</sub>F<sub>12</sub>, remarkable in containing a higher percentage of lithium (5 35 p c) than any other mineral. It occurs as large, colourless, rhombic-dodecahedra embedded in the cryolite deposit at Ivigtut in Greenland. Sp gr. 2 77; H. 2½–3; n<sub>Na</sub> 1 3395; in p. 710°. The mineral dissolves in 1350 parts of water at 18°, and from this solution crystallises as cubes.

CRYOSCOPY. So long ago as 1788 Blagden proved that the extent to which the freezing point of an aqueous solution lies below that of water—the depression of the freezing point, as it is called—is proportional to the concentration of the dissolved substance. Nearly a century later, the researches of de Coppet and Raoult showed that when equimolecular proportions of different substances are dissolved in equal quantities of a given solvent, the solutions so obtained have, in general, the same freezing point. It is on these facts that the use of the cryoscopic method of determining the molecular weights of dissolved substances is based.

In comparing the effects of different substances on the freezing point of a given solvent, it is obviously convenient, in view of de Coppet's and Raoult's results, to refer these effects to 1 gram molecule of solute and to some definite quantity of the solvent: in practice, 100 grams of the latter are taken as the standard amount The depression of the freezing point for such a concentrated solution as one containing 1 gram. molecule of solute in 100 grams of solvent cannot be determined directly: it is calculated from the depression observed for a dilute solution, on the basis of the proportionality between depression and concentration An example may be taken to illustrate this point tion containing 0 609 gram of ethyl alcohol in 100 grams of water freezes 0 243° below the freezing point of water; from these data it follows, if proportionality between depression and concentration is assumed, that the depression for a solution containing 1 gram molecule of alcohol in 100 grams of water would  $\frac{0.243 \times 46}{0.609} = 18.4^{\circ}$ Again, a solution 4 276 grams of sucrose in 100 grams of water, freezes 0 237° below the freezing point of water; from these data it follows, as above, that the depression for a solution containing 1 gram molecule of sucrose in 100 grams of water would be  $0.237 \times 342 = 19.0^{\circ}$ . If the depressions pro-4 276 duced by I gram molecule of other non-electrolytes in 100 grams of water are similarly calculated from experimental data, values are obtained which are very close to the figures already quoted. The depression, therefore, for a solution containing I gram molecule of non-electrolyte in 100 grams of water is a constant, independent of the particular solute; the average value of the constant for water, deduced in the empirical way just described, is 18 5°.

For each solvent, a characteristic constant

(k) can be similarly deduced from experimental data. The evaluation of this constant—the 'molecular depression of the freezing point'—is effected, as illustrated above, by the formula  $k = \frac{M}{g} \Delta t$ , where  $\Delta t$  is the depression of freezing point observed for a solution containing g grams of solute in 100 grams of solvent, and M is the molecular weight of the solute

It has been shown by van't Hoff that, on thermodynamical grounds, the molecular depression of the freezing point for any solvent is quantitatively related to the temperature of freezing and to the latent heat of fusion. The relationship in question is  $k = \frac{0.0198T^2}{\omega}$ , where T is the freezing point of the solvent on the absolute scale, and  $\omega$  is the latent heat of fusion per gram of the solvent. The following table shows how far the values of the molecular depression, calculated for various solvents by van't Hoff's formula, are in agreement with those deduced empirically.—

		k	$\boldsymbol{k}$
Solvent		Empirical	('alculated
Water .		$18.5^{\circ}$	$18.6^{\circ}$
Formic acid		$28^{\circ}$	$27.5^{\circ}$
Acetic acid		39°	38 5°
Benzene .		50°	52°
Naphthalene		68 5°	69°
p-Toluidine		53°	51°
Phenol .		75°	74°

The formula  $k = \frac{0.0108 T^2}{\omega}$  is based on the assumption that, when a solution freezes, it is the pure solid solvent which crystallises out. It is not surprising, therefore, that in cases where the crystals separating from the solution contain some of the dissolved substance, eg solutions of thiophene in benzene, the empirical value for the molecular depression, deduced in the usual way, differs considerably from the normal value. In such cases the depression observed is smaller than it would be if the pure solid solvent separated out

When the value of k for any solvent has been ascertained, either empirically by experiments with normal substances of known molecular weight, or indirectly by van't Hoff's formula, it may be utilised in determining the molecular weights of new substances. For this purpose, the formula  $k = \frac{M}{g} \Delta t$  is employed, but now k is taken as known, and M is the quantity to be evaluated. In view of this, the equation may be written  $M = \frac{k}{\Delta t}$ .

It is evident that a determination of the molecular weight of a dissolved substance by the cryoscopic method depends, not on the absolute freezing temperature of the solution, but on the difference between the freezing point of the solvent and that of the solution. This difference can be determined with greater accuracy than can the freezing temperatures themselves. A thermometer with a very open scale, covering only 5°-6°, is employed, and it is convenient, as suggested by Beckmann, to have a reservoir attached to the upper end of the thermometer tube, so that the quantity of mercury in the bulb and stem may be varied according to requirements. This device permits

the use of one and the same thermometer at temperatures, and therefore various solvents. In Beckmann's thermometer, which is the one usually employed in cryoscopic investigations, the scale degree is about 4 cm in length and is divided in hundredths: with the aid of a lens, preferably adapted for sliding up and down the stem, it is possible to estimate 0.001°. The form of the reservoir attached to the upper end of the ordinary Beckmann thermometer is shown in the accompanying sketch. When it is desired to adjust the thermometer for experiments with any given solvent, the bulb is immersed in water of the same temperature as the freezing point of If the top of the mercury thread the solvent




Fig. 1.

should reach a steady position on the upper half of the scale, no further adjustment is needed Should the mercury, after sufficient time has been allowed, extend beyond the scale and into the reservoir, it is necessary to detach a portion This is effected by putting the bulb in water 2°-3° higher than the required temperature, and then gently tapping the upper end of the thermometer until the mercury hanging at the top of the reservoir is separated from the thread. On cooling down again to the freezing temperature of the solvent, the top of the thread should now come to rest somewhere on the upper half of the scale If the quantity of mercury in the bulb and stem is too small, so that when the thermometer is immersed in water of the same temperature as the

freezing point of the solvent, the thread does not reach the scale or appears only on the lowest part of it, the bulb is warmed until the mercury appears at the top of the reservoir. By a throwing movement, some of the mercury at the bottom of the reservoir is jerked up to the top, where it joins the end of the thread. The bulb is then immersed in water 2°-3° above the temperature required, and the final adjustment is carried out in the way already described.

The cryoscopic apparatus commonly employed for the determination of the molecular weight of dissolved substances is that known as Beckmann's. The accuracy of this method, as ordinarily applied, is not great, the error being of the order ±5 p.c. For most laboratory purposes, however, this is sufficient, and it is only when extreme accuracy is required, as in the investigation of very dilute aqueous solutions, that special forms of apparatus have to be employed.

The essential parts of Beckmann's apparatus are seen in the subjoined diagram. A stout glass jar, holding about 1½ litres, is provided with a metal top, at the side of which is a hole for a stirrer, and in the middle of which is another hole capable of holding a large stout test-tube, 3-4 cm. in diameter. The lid and the tube are held in position by springs, as will be evident from the diagram. By means of a cork, there is fitted in the tube another, narrower one, to

hold the liquid under investigation. This tube should preferably have a flat bottom, while the part of it which projects beyond the outer jacket is provided with an auxiliary side tube.

and is itself widened out at the upper extremity The cork with which this inner tube-the freezing tube, as it may be called -is fitted carries the thermometer, and allows also free passage to a stirrer The latter is best made of stout platinum wire, but silver or nickel may be used, provided that the possibility of chemical action is ex-The thermomecluded ter must be fitted in an accurately central position, so that the stirrer may be moved up and down without grating

When a determination of molecular weight is to be made, the freezing tube is charged with a weighed quantity of the solvent, usually 10-20 grams, and the jar is filled with water or a mixture of ice and salt kept at a temperature




Fig. 2.

4°-5° below the freezing point of the solvent under examination. The freezing tube, with thermometer and stirrer, is immersed directly in the jar, until the temperature has fallen almost to the freezing point. It is then set in the jacket tube and the contents are stirred regularly, while the mercury in the thermometer falls slowly The fall continues until the solvent is more or less supercooled. In the majority of cases, freezing sets in spontaneously before the supercooling exceeds 0.5°, and the liberation of heat results in a rise of the mercury in the thermometer up to a steady maximum position. This is the freezing point of the solvent—the temperature at which liquid and solid solvent are in equilibrium with each other It is advisable, when the thread is rising towards its steady position, to tap the thermometer before taking a reading, and so avoid the possibility of the mercury sticking in the capillary. Should the supercooling exceed 0.5°, crystallisation ought to be initiated, either by stirring very vigorously for a few moments or by introducing a tiny crystal of the solid solvent. Whenever freezing has begun, the solvent must then be stirred with uniform regularity until the constant temperature has been reached.

The freezing point of the solvent having been determined two or three times in the manner just described, a weighed quantity of the solute, enough to give a depression of not less than 0.2°, is introduced, and the freezing point of the solution is then ascertained in the same way as that of the solvent. Subsequent additions of solute are made and the resulting depressions determined as before. One addition, it is true, yields sufficient data for the calculation of the molecular weight of the solute, but it is desirable

CRYOSCOPY. 430

to ascertain also whether the concentration of the solute has any definite and notable influence on the value obtained for its molecular weight It is advisable, further, to redetermine the freezing point of the solvent at the end of such a series of experiments, and so obtain a guarantee that the zero of the thermometer has not changed appreciably during the course of the

Various suggestions have been made from time to time with the object of rendering the simple Beckmann apparatus a more accurate instrument Thus it has been proposed to stir the contents of the freezing tube mechanically instead of by hand The access of atmospheric moisture to a hygroscopic solvent may be prevented by passing a current of dry air into the freezing tube during an experiment, or by using as stirrer an iron ring which is enveloped in platinum foil and is operated electromagne-In this latter case, obviously, communication between the contents of the freezing tube and the outside air can be completely cut off. The same end is attained by employing an ordinary stirrer with a mercury valve. (For a detailed description of these modifications and additions, see Beckmann, Zeitsch physikal Chem. 1891, 7, 323; 1896, 21, 239; 1897, 22, 616, 1903, 44, 169; also Ostwald and Luther's Physiko-chemische Messungen

For the determination of the freezing point of liquids which are obtainable only in small quantities-physiological fluids, for example-Beckmann's apparatus, in its ordinary form, is too large Modifications of this apparatus, suitable for the investigation of blood, urine, &c, have been suggested by various authors (Guye and Bogdan, J Chim. Phys 1903, 1, 379, Burian and Drucker, Zentr Physiol 1910, 23, 772) In these instruments, only 1-2 cub cm. of liquid is required, and it is claimed that the accuracy is of the same order as that attainable with the ordinary Beckmann apparatus this connection, mention ought to be made also of Eykman's apparatus (Zeitsch physikal Chem. 1888, 2, 964, 1889, 4, 497) This con-sists of a small flask, holding about 10 cub cm, into the neck of which is ground a thermometer, graduated in tenths or hundredths of a degree There is no separate jacketing vessel, but the procedure is similar to that followed in the case

As regards the preparation of the cooling bath required in the cryoscopic investigation of aqueous solutions, the usual mixture of ice and common salt solution may be replaced in some cases by a mixture of ice and a solid salt, which gives a steady eutectic temperature suitable for this purpose, with the eutectic temperatures obtained when they are mixed with ice, are alum,  $-0.47^{\circ}$ ; sodium sulphate.  $-0.7^{\circ}$ ; potassium dichromate,  $-1.0^{\circ}$ , potassium sulphate, -1 5°; copper sulphate, --2 0°; otassium nitrate,  $-3.0^{\circ}$ ; zinc sulphate,  $-5.0^{\circ}$ ; barium chloride,  $-7.0^{\circ}$ . Another kind potassium nitrate, of cooling bath was used by Raoult, who surrounded the jacketed freezing tube with ether, and passed through the latter a current of dry air (Compt. rend. 1896, 123, 476; Zeitsch.

of the Beckmann apparatus

any point between room temperature and  $-15^{\circ}$ .

When it is desired to find the molecular weight of a new substance, the question of the most suitable solvent arises. The choice is frequently limited by the solubility of the substance, it may be appreciably soluble in only one or two of the more common solvents. Apart from this, however, there are other considerations which must be taken into account. Extensive investigation has proved that with certain combinations of solvent and solute, the cryoscopic method gives abnormal values for the molecular weight of the dissolved substance. Pre-emmently is this the case when the solvent is water and the solute is an acid, a base, or a salt, in short, an electrolyte; under these circumstances, the value found for the molecular weight of the solute is abnormally low. On the other hand, the influence of substances containing the hydroxyl group (alcohols, acids, phenols, oximes) on the freezing point of hydrocarbons (benzene) and allied indifferent solvents, is such as to lead to abnormally high values for the molecular weight The cryoscopic investigation of solutions of acetic acid in benzene indicates that the formula of the acid is more correctly represented by (CH3 COOH), than by CH, COOH Again, while the molecular weight of alcohol in benzene is nearly normal in dilute solution, the value increases rapidly as the concentration of the alcohol rises The combinations of solvent and solute which lead to abnormally high values of the molecular weight have been studied in detail by various workers (Auwers, Zeitsch physikal Chem 1893, 12, 689, 1894, 15, 33; 1895, 18, 595; 1896, 21, 337; 1897, 23, 449, 1899, 30, 300, 1903, 42, 513, 629, Biltz, *ibid* 1899, 29, 249; Robertson, Chem Soc Trans. 1903, 83, 1425; 1904, 85, 1427; 1005, 27, 1874, 1006, 20, 247; 1447 1905, 87, 1574; 1906, 89, 567). 1617; should be noted that in cases also where the dissolved substance crystallises out along with the solvent, the depression of the freezing point is abnormally small, and the value of the molecular weight calculated therefrom is correspondingly high.

For the cryoscopic determination of the molecular weights of dissolved substances a method of moderate accuracy is sufficient, and an experiment may be carried out with a comparatively small quantity of liquid The case is different with the problem of finding the exact depression of the freezing point for very dilute aqueous solutions—a problem to which much attention has been devoted because of its bearing on the theory of solution It was early evident that the cryoscopic investigation of very dilute solutions involves special difficulties, for the discrepancies between the results obtained by different workers were very considerable Thus the molecular depression for water, based on experiments with sucrose in which the actually observed depressions amounted only to about 0 05°, was found by Arrhenius to be 20 2°, by Raoult, 20·7°; by Jones, 21·8°; and by Loomis, 18·1°, instead of the theoretical value. 18.6°. Subsequent investigations (Nernst and Abegg, Zeitsch. physikal. Chem 1894, 15, 681; physikal. Chem. 1898, 27, 617). By careful Abegg, ibid. 1896, 20, 207; Wildermann, Phil. regulation of the rate at which the air is forced Mag. 1895, 40, 119; Proc Roy. Soc 1896, 251; through the ether, the bath can be adjusted to Raoult, Zeitsch. physikal. Chem. 1898, 27, 617)

have made it clear what conditions must be fulfilled if the freezing point depression for a dilute aqueous solution is to be determined with certainty to 0 001°. It will be sufficient here to indicate these conditions briefly.

As regards the thermometer, it is essential that a telescope be used for reading the position of the mercury, and that the bulb of the instrument be kept at about 0° during the whole of a series of experiments The influence of alterations in the atmospheric pressure cannot be neglected, for it appears that a difference of 1 mm mercury has about the same effect on a Beckmann thermometer as a temperature change of 0 0002° On similar grounds, the thermometer must be immersed always to the same depth in solvent and solutions It should be mentioned here that platinum resistance thermometers have been used in accurate cryoscopic work instead of the Beckmann instruments (see Chrustchoff, Compt rend 1900, 131, 883, 1901, 133, 955, Bedford, Proc Roy. Soc. A, 1910, 83, 454) The difference between the freezing points of solvent and solution has also been determined thermoelectrically (Haus-Tath, Ann Physik. 1902, [iv] 9, 522; Jahn Zeitsch physikal Chem. 1907, 59, 31, Flugel, *ibid.* 1912, 79, 577; Adams, J. Amer. Chem Soc. 1915, 37, 481).

The temperature of the cooling bath is a matter which has to be carefully considered, if the freezing point depression for a dilute solution is to be correctly determined. Suppose that in an experiment with the ordinary Beckmann apparatus the aqueous solution in the inner tube cools without freezing, it is obvious that the lowest temperature to which the solution can fall must be higher than that of the cooling bath, because of the heat conducted from the air to the solution through the thermometer. stirrer, and tube, and because of the heat generated by the stirring This lowest temperature, which the contents of the freezing tube tend to assume when no freezing takes place, is known as the convergence temperature difference between the convergence temperature and the temperature of the cooling bath varies, of course, with each apparatus, it decreases as the facility for heat exchange between the solution and the external air is lessened, and as the rate of stirring is diminished Suppose now that the solution under examination has become supercooled, and that it has reached the convergence temperature. The introduction of a crystal of the solid solvent at this stage initiates the freezing process; heat is liberated and the temperature of the solution rises. Inasmuch, however, as the solution is no longer at the convergence temperature, there is an influence at work tending to lower the temperature of the solution, so that the latter never reaches its true freezing point the maximum position reached by the mercury is only what may be called the apparent freezing point of the solution. Similarly, when the freezing point of the solution lies below the convergence temperature, an apparent freezing point is observed which is somewhat higher than the true freezing point. In both cases, the apparent freezing point lies between the true freezing point and the convergence temperature. In view of these considerations, it is necessary, in cryoscopic work

of the most accurate kind, either to adjust the cooling bath so that the convergence temperature and the freezing point are practically the same, or to apply a correction to the apparent freezing point (see Raoult, lc; Nernst and Abegg Lc)

Abegg, lc)

The freezing point of a solution is the temperature at which the solution and the solid solvent are in equilibrium, and in order that this equilibrium, and therefore the freezing point, may be well defined, it is essential that a considerable quantity of solid should be in contact with the solution. In the case of aqueous solutions, this has generally been effected by supercooling to the extent of 05° or 10°, and then introducing a crystal of ice If this method is adopted, then a correction must be made for the charge of concentration resulting from the separation of ice Some workers, however, determine the freezing point by adding the cooled solution to a fairly large quantity of finely divided ice, and then observing the steady temperature shown by a thermometer immersed in the mixture (see Walker and Robertson, Proc Roy Soc Edin 1903, 24, 363, Richards, J Amer Chem Soc 1903, 25, 291, Jahn, Zeitsch. physikal Chem 1904, 50, 129; Roth, *ibid*. 1912, 79, 601, Adams, J Amer Chem. Soc 1915, 37, 481, compare also Prytz, Ann Physik 1902, [iv] 7, 882) In this method of procedure, equilibrium between the two phases is very rapidly established and the concentration of the liquid is ascertained by an analysis made subsequently to the determination of the freezing point

It has already been pointed out that the freezing point depression is proportional to the number of dissolved units. The cryoscopic number of dissolved units. The cryoscopic method may therefore be employed, not only in the determination of the molecular weights of dissolved substances, but also in the study of the question whether in a particular case a change in the number of dissolved units takes place. It is found, for instance, that the freezing points of potassium iodide solutions are not appreciably affected by the addition of iodine, which shows that the iodine must combine with the units already present in the solution (see Le Blanc and Noves, Zeitsch physikal Chem 1890, 6, 385, Paterno and Peratoner, Gazz chim ital 1890, The cryoscopic method may similarly be employed in following the course of a reaction between dissolved substances which involves a change in the number of units, as for instance, the inversion of sucrose (see Kahlenberg, Davis, and Fowler, J Amer Chem Soc 1899, 21, 1). J. C. P.

CRYPTOPINE v OPIUM.
CRYSOBERYL or CHRISOBERYL v GLUCI-

CRYSTAL PONCEAU v. Azo- colouring matters

CRYSTAL VARNISH or PAPER VARNISH v. VARNISH.

CRYSTAL VIOLET v TRIPHENYLMETHANE COLOURING MATTERS

CRYSTALLIN v PROTEINS.

NUM

CRYSTALLISATION. Crystals are often produced during the concentration of liquors in evaporating pans. Hand or mechanical scrapers are used for removing these crystals continuously. Such crystals are generally too

impure for sale and are redissolved in sufficient hot water to produce a solution, which, after clarification by subsidence or filtration, will yield, in the crystallising pan, bright and marketable crystals. The temperature and concentration of the solution, to yield the best result, have to be determined for every substance mother liquor is run off and the crystals transferred to drainers or centrifugal machines to be dried before packing In some instances, several repetitions of this treatment are made, and the crystals are washed with water or liquor after each treatment. The crystallising pans are made of materials which give the best economic result, either they resist the mother liquors or, if attacked, do not spoil the product by discolouration or too much impurity iron tanks about 2 feet deep with bottoms sloping to the outlet and rounded corners, are Wooden tanks unlined or lined with common sheet lead, zinc, tin, or other metal, are used Some manufacturers suspend roughened strips of metal in the pans, to which the crystals adhere For rapid production, crystallisation by repose is now giving place in many industries to crystallisation in motion crystallising vessels are provided with circulating blades or with compressed air jets, so that the crystals as they form are prevented from adhering to the sides of the vessel, but remain suspended in the liquor, which, on complete cooling, flows directly to the centrifugal machines for the separation of the mother liquor These crystals are purer and more convenient to handle than those obtained by repose methods In the case of deliquescent substances and also in those instances where the mother liquor is sufficiently pure to make it unnecessary to purify by crystallisation, the liquor is concentrated in film evaporators so that only sufficient water remains in it to form the crystal hauor produced is run into drums, and forms on cooling, a solid block of crystals In some works, the liquor is run into a jacketed worm conveyor so that the crystals which form, leave the apparatus in a loose state

The capacity of crystallising plant can be determined from a knowledge of the physical data involved When crystallising by repose methods the rate of evaporation of the solution is given very closely by the utility formula:

$$\mathbf{W} = \left(\frac{p_{e} - p_{d}}{50}\right)^{1.2}$$

where W is the weight of water in kilograms evaporated per hour per square metre of surface of the liquid,  $p_e$  is the vapour pressure of the solution in mm of mercury at the temperature under consideration, and  $p_d$  is the pressure of the aqueous vapour in the atmosphere of the factory. If a draught with a velocity exceeding (say) 1 metre per second is produced in the crystallising room, the figure given by the above formula will be low, and should be multiplied by a factor which is approximately the square root of the velocity in metres per second

If the mother liquor is run into the crystallising vessels hot, then the time of cooling and evaporation can be determined by a graphical integration of the rates of cooling from the surface of the vessel and cooling from evapora-

tion from the surface of the liquid at a series of temperatures.

Many solutions will not produce bright crystals at temperatures above 20°C.; in such cases, propeller fans are most useful in keeping down the temperature and increasing the output.

Substances which form hard comented masses on the cool surfaces of the crystallising vessels may be most satisfactorily treated in vacuum evaporators, where the crystals form on the surface and then drop to the bottom of the vessel, unless a surrer be provided to keep them in suspension. This method is most useful when crystallising from volatile solvents

Crystals of uniform size are also most efficiently produced in this way, since the rate of evaporation and the temperature can be perfectly regulated. A mother liquor of suitable strength is produced in the pan and allowed to cool to saturation temperature. The solution is then 'sown' with a suitable quantity of finely powdered substance, and the heating coils and vacuum pump set to work at such a rate as to maintain the temperature and produce the required amount of evaporation, the liquid being stirred continuously. As soon as the crystals are known to have reached the required size, the mass is discharged through strainers and the mother liquor returned to form a portion of the next charge.

CUBA BLACK v A70- COLOURING MATTERS CUBEBS. Cubebæ fructus, BP The dried full-grown unripe fruits of Piper Cubeba (Linn fil)

CUBEBS CAMPHOR v CAMPHORS.
CUBEBS, OIL OF, v OILS, ESSENTIAL.
CUCUMBER. The fruit (really a berry) of
Cucumis sativus (Linn) Konig gives as the
average composition

Henze (Zeitsch Nahr Genussm 1903, 6, 529) observed that large encumbers contain a higher proportion of sugar than small ones, but less introgenous matter. He gives as the mean composition

The ash of cucumbers contains:

H. I.

CUDBEAR v. Archil.
CULLEN EARTH or COLOGNE EARTH v.
Pigments.

CULLET v. GLASS.

CULM v Anthracite, art FUEL.

CUMENES. Cumols. The name cumene, originally applied to isopropylbenzene

 $C_0H_0\cdot CH(CH_3)_2$ 

to indicate its connection with cumic acid, from which it was obtained by distillation with lime, is frequently employed as a generic term to denote any of the isomeric benzene hydrocarbons of the formula C<sub>9</sub>H<sub>12</sub>. Theory predicts

CUMENES. 433

the existence of eight of these, all of which have been found in coal-tar oil.

Propylbenzenes.

1. Normal propylbenzene (n. cumene)

C<sub>6</sub>H<sub>5</sub>·CH<sub>2</sub>·CH<sub>2</sub>·CH<sub>3</sub>

Present in coal tar oil in fraction boiling at 153°-157° (Schultz and Perl, Ber 1909, 3602) By the action of sodium on a mixture of bromobenzene and propyl bromide in ethereal solution (Fittig, Schaffer and Konig, Annalen, 149, 324) By the action of aluminium chloride on trimethylene bromide, propylbenzene and diphenylpropane (sym) are formed, the latter of which, on further treatment, yields propylbenzene and cumene (Bodroux, Compt rend 132, 155). From benzene, aluminium chloride and propyl bromide, when kept at 0° (Heise, Ber 1891, 768; Genvresse, Bull Soc chim 9, [3] 219). From benzyl chloride, C<sub>6</sub>H<sub>5</sub> CH<sub>2</sub>Cl and zinc-ethyl (Paternò and Spiza, Ber 10, 294). Liquid boiling at 159° (corr.), sp. gr. 0.867 at 15°[15° (Perkin, Chem. Soc Trans. 1900, 274) Yields benzoic acid on oxidation with chromic acid.

Cumene, together with toluene, may be obtained from cymene by demethylation by the action of aluminium chloride in presence of benzene.

 $\mathrm{C_3H_7\,C_6H_4\,CH_3}{+}\mathrm{C_6H_6}$ 

 $^{6} = C_{6}H_{5} CH_{3} + C_{3}H_{7} C_{6}H_{5}$ 

(Bædlker and Halse, Bull Soc. chim. 1916, 19, 444)

2 İsopropylbenzene (Cumene)

 $C_6H_5\cdot CH(CH_3)_2$ 

Present in American petroleum (Maberry and Dunn, Amer Chem J 18, 215) By distilling cumic acid  $C_6H_4[\mathrm{CH}_{(2H_3)_2}]\mathrm{CO}_2H(1,4)$  with lime or baryta (Gerhardt and Cahours, Ann. Chim Phys [3] 1, 87) From benzal chloride  $C_6H_5$  CHCl, and zinc-methyl (Liebermann, Ber 13, 46) Phenyl magnesium bromide reacts with acetone to give methylethenylbenzene, which, on hydrogenation, yields cumene (Tiffeneau, Compt rend 134, 845) From benzance and either isopropyl bromide or normal propyl bromide in presence of aluminium bromide (Gustavson, Ber 11, 1251) Liquid boiling at 152 5°–153° (cort.), sp. gi. 0.8727, 79°/4° (Perkin, Chem Soc Trans 1900, 275) Yields benzoic acid on oxidation with chromic acid

Methylethylbenzenes (Ethyltoluenes) The three methylethylbenzenes are obtained by the action of sodium on a mixture of ethyl bromide or iodide with the corresponding bromotoluenes (ortho-, meta-, or para-)

1. o-Methylethylbenzene

 $C_6H_4(CH_3)(C_2H_5)(1, 2)$ 

Liquid boiling at  $158^{\circ}$ – $159^{\circ}$ , sp gr 0.8731 at  $16^{\circ}$ . Dilute intric acid oxidises it to o-toluic acid  $C_8H_4(CH_3)CO_2H(1,2)$ 

2. m-Methylethylbenzene

 $C_6H_4(CH_3)(C_2H_5)(1, 3)$ 

Liquid boiling at  $158^{\circ}-159^{\circ}$ ; sp gr. 0.869 at  $20^{\circ}$ . Chromic acid oxidises it to isophthalic acid  $C_6H_4(CO_2H)_3(1,3)$ .

3. p-Methylethylbenzene

 $C_6H_4(CH_8)(C_9H_8)(1, 4)$ 

Liquid boiling at 161° 162°; sp.gr. at 0.8652 at C. H<sub>3</sub>(CH<sub>3</sub>)<sub>2</sub>CO<sub>2</sub>H—xylic acid and p-xylic acid. Vol. II.—T.

21°. Yields on oxidation p-toluic and terephthalic acids.

Trimethylbenzenes.

1. Consecutive trimethylbenzene (Hemillitine, Hemimellithene)  $C_0H_3(CH_3)_3(1,\,2,\,3)$  By heating the calcium salt of a-isoduric acid

with lime Liquid boiling at 175° (Jacobsen, Ber 15, 1857, 19, 2517)

2. Unsymmetrical trimethylbenzene (pseudocumene) C<sub>6</sub>H<sub>3</sub>(CH<sub>3</sub>)<sub>3</sub>(1, 2, 4), occurs together with mesitylene in coal-tar (Beilstein and Koglei, Annalen, 137, 317), and is present in many petroleums (Poni, Chem Zentr 1906, 1 459) By the action of sodium on a mixture of either bromo-paraxylene or bromometaxylene with methyl lodide. In order to isolate it from coal tar, Aimstrong recommends (Chem News, 38, 5; Ber. 11, 1697) that the mixture of pseudocumene and mesitylene (v infia) boiling between 160° and 170°, obtained from coal tar by fractional distillation, should be converted by treatment with an equal volume of concen trated sulphuric acid into sulphonic acids. The mixture of sulphonic acids, which separates as an oily layer on carefully adding water to the sulphuric acid solution, is heated with concentrated hydrochloric acid in a sealed tube for 1 hour at 100°. In this way, the mesitylene sulphonic acid is decomposed, regenerating mesitylene, whilst the pseudocumenesulphonic acid is not attacked After separating the mesitylene the pseudocumenesulphonic acid is precipitated from the hydrochloric acid solution by sulphuric acid, purified by recrystallisation from dilute sulphuric acid and finally decomposed by heating with hydrochloric acid at 140°-150°, when it breaks up into pseudocumene and sulphune acid Jacobsen (Ber 9, 256) treats the mixture of sulphonic acids with barium or calcium carbonate, which removes the excess of sulphuric acid and converts the sulphonic acids into barium or calcium salts, transforms these by double decomposition with sodium carbonate into sodium salts, heats the latter with phosphorus pentachloride, with which they yield the sulphonic chlorides C<sub>2</sub>H<sub>11</sub> SO<sub>2</sub>Cl, converts the chlorides into the sulphonamides

by the action of ammonia, and separates the two isomeric sulphonamides by recrystallisation from alcohol, in which mesitylene sulphonamide is readily, pseudocumene sulphonamide only sparingly, soluble. The sulphonamides are then hydrolysed by heating with concentrated hydrochloric acid at 175°.

Another method proposed by Jacobsen (Annalen, 184, 199) is to separate the two sulphonic acids by fractional crystallisation; pseudocumensulphonic acid is soluble with difficulty in sulphuric acid and crystallises out. On heating to its melting-point (110°) and passing in steam at a temperature of 250°, pure pseudocumene is obtained (see also Schultz and Herzfeld, Ber 1909, 3602). Pseudocumene boils at 168 2°, and has sp.gr. 0 877 at 18° (Schultz and Herzfeld, lc) On oxidation with intric acid, it yields two isomeric acids of the formula  $C_6H_3(CH_3)_2CO_2H$ —xylic acid and p-xylic acid

434 CUMENES

-and a small quantity of the isomeric methylisophthalic and methylterephthalic acids

$$C_6H_3(CH_3)(CO_2H)_2$$

(Bentley and Perkin, Chem Soc Trans. 1897,

3. Symmetrical trimethylbenzene (Misityline) C<sub>6</sub>H<sub>3</sub>(CH<sub>3</sub>)<sub>3</sub>(1, 3, 5) In coal-tar oil, in fraction boiling at 157°-162° (Fittig and Wackenroder, Annalen, 151, 292, Schultz and Perl, Ber 1909, 3602), and in American petroleum (Maberry and Dunn, Amer Chem J. 18, 215). By distilling acetone with sulphuric acid

$$3CO(CH_3)_2 = C_6H_3(CH_3)_3 + 3H_2O$$

(Kuster and Stollberg, Annalen, 278, 207, Meyer and Molz, Ber 1896, 2831), or by dissolving allylene CH<sub>3</sub>C CH in sulphuric acid and distilling with water 3CH<sub>3</sub>C CH=C<sub>6</sub>H<sub>3</sub>(CH<sub>3</sub>)<sub>3</sub> (Fittig and Schrohe, Ber 8, 17) (On the isolation of mesitylene from coal tai and the method of separating it from pseudocumene, v preceding paragraph.) Liquid boiling at 164 5°; sp gi 0 8768 at 4°/4° (Perkin, Chem Soc. Trans. 1896, 1193) Dilute nitric acid oxidises it to mesityleme acid C<sub>6</sub>H<sub>3</sub>(CH<sub>3</sub>)<sub>2</sub>CO<sub>2</sub>H and uvitic acid  $\mathrm{C}_{6}\mathrm{H}_{3}(\mathrm{CH}_{3})(\mathrm{CO}_{2}\hat{\mathrm{H}})_{2}$ ; with potassium permanganate it yields uvitic acid and trimesic acid  $C_6H_3(CO_1H)_{13}(1,3,5)$  CUMIDINES  $C_9H_{13}N$  Of the sixteen iso-

meric cumidines which have hitherto been obtained, pseudo- or  $\psi$ -cumidine and mesidine are perhaps the most important. These bases are derivatives of pseudocumene and mesitylene respectively, the first of which is theoretically capable of yielding three isometic \( \psi\)-cumidines, and the latter one cumidine only

Pseudocumidine, sym - \psi - cumidine

 $(CH_3)_3C_6H_2$   $NH_2[CH_3$   $CH_3$   $CH_3$   $NH_2=1$  2 4 5] can be obtained by nitrating pseudocumene with fuming nitric acid in the cold, and reducing the nitro- compound (m p 71°) with tin and hydro-chloric acid (Schaper, Zeitsch chem [2] 23, 1) The base is prepared technically by digesting xylidine hydrochloride with methyl alcohol in an autoclave at 280°-300°, converting the crude product into the nitrate, separating the spaningly soluble nitrates by means of a centrifugal machine, washing the product with water and The mixture of finally treating with an alkali bases consisting of cumidines and xylidines is then fractionated, and the portion distilling between 225°-245° is allowed to crystallise, and finally freed from non-crystallisable bases by pressure (Aktiengesellschaft fur Anilin-fabrikation, D. R. P. 22265). The cumidine so The cumidine so obtained contains mesidine and other impurities (Hofmann, Ber 17, 1912; Engel, ibid 18, 2229), but consists chiefly of Schaper's pseudocumidine (cf Hotmann, ibid. 15, 2895) which can be prepared in the pure state by digesting o-4-xylidine [NH<sub>2</sub>.CH<sub>3</sub>.CH<sub>3</sub>=1 3 4] or paraxylidine with methyl alcohol under similar conditions (Nolting and Forel, ibid. 18, 2680)

Properties — Pseudocumidine crystallises in large, lustrous prisms, melts at 63° (Hofmann), at 68° (Auwers, Ber. 18, 2661), and boils at 234°. It is insoluble in water, but readily soluble in alcohol. The nitrate is characterised by being very sparingly soluble in water. Its acetyl derivative crystallises in compact white needles, melts at 161° (Edler, ibid. 18, 629; Nölting and Baumann), at 164° (Auwers) and boils above Pseudocumidine is used as a source of azo- dyes (D. R. PP 3229, 1878; 22265, 1882) as-ψ-cumidine [CH<sub>3</sub> CH<sub>3</sub> CH<sub>3</sub> NH<sub>2</sub>=1.2.4:6] has been prepared by Edler (l.c.). It melts

at 36°.

 $\psi$ -cumidine [CH<sub>3</sub> CH<sub>3</sub>.CH<sub>3</sub>:NH<sub>2</sub>=1.2:4.3] is obtained by digesting 0-3-xylidine hydro-chloride [NH<sub>2</sub> CH<sub>3</sub> CH<sub>3</sub>=123] with methyl alcohol in an autoclave at 300° (Nolting and Forel; cf Mayer, Ber 20, 972) It boils at 240° (Nolting and Forel), 236° and does not solidify at -15° (Mayer). Its acetyl derivative melts at 186°.

Mesidine  $[CH_3 CH_3 CH_3 NH_2=1.3.5 2]$  is obtained by treating mesitylene with nitric acid (sp gr 138) and reducing the nitro-derivative with tin and hydrochloric acid (Fittig and Storer, Annalen, 147, 3) It can also be prepared by digesting  $\alpha$ -metaxylidine [NH<sub>2</sub> CH<sub>3</sub> CH<sub>3</sub>=1 2 4] (Hofmann, Ber 5, 715, 8, 61) or  $\beta$ -metaxylidine [NH<sub>2</sub> CH<sub>3</sub> CH<sub>3</sub>=1 2 6] hydrochloride (Nolting and Forel) with methyl alcohol at 300°

Properties—Mesidine is a colourless liquid, which boils at 229°, and has a sp gr 0 9663. The acetyl derivative melts at 216° and crystal-

lises in prisms

Isocumidine [CH<sub>3</sub> CH<sub>3</sub> CH<sub>3</sub> NH<sub>2</sub>=1 2 3 5], prepared by digesting the hydrochloride of γ-metaxylidine

$$[NH_3.CH_3.CH_3=1.3.5]$$

with methyl alcohol, at  $300^\circ$  (Nolting and Forel; Limpach, Ber 1888, 644), melts at  $67^\circ$ – $68^\circ$ , boils at  $245^\circ$ , and yields an acetyl derivative melting at  $163^\circ$ – $164^\circ$ 

Para-aminopropylbenzene, formed by heating aniline, propyl alcohol, and zinc chloride at 280° for 8 hours (Louis, Ber. 16, 105, Francksen, ibid. 17, 1221), boils at  $224^{\circ}-226^{\circ}$ . The acetyl derivative melts at  $87^{\circ}$ .

Orthoaminoisopropylbenzene, - o - cumidine, formed by heating aminocumic acid with baryta (Filetti, Gazz chim. ital. 13, 379; cf Constam and Goldschmidt, Ber 1888, 1158), boils at 213 5°-214 5° at 732 mm The acetyl derivative melts at  $72^{\circ}$ 

Para-aminousopropylbenzene, cumidine, formed by heating isopropyl alcohol, aniline, and zine chloride at 260°-280° (Louis, Ber 16, 111, cf. Nicholson, Annalen, 65, 58), boils at 216°-218°. The acetyl derivative melts at 102°-102 5°. By oxidation with sodium chlorate and potassium ferrocyanide in hydrochloric acid solution, cumidine yields a dyestuff of the aniline black series (Kirptischnikow, Chem. Zentr 1906, 1. By treatment with cyanogen chloride in pyridine solution, condensation is effected and a dyestuff is produced (Konig, D R. P. 155782; Chem Zentr 1904, n 1557).

Ethylorthotoluidine, prepared by treating orthotoluidine in ethyl alcohol with zinc chloride at 270° (Benz, Ber 15, 1650), boils at 229°-230°. The acetyl derivative melts at 105°-105 5°, and boils at 313°-315°.

CUMIDINE RED v. AZO- COLOURING MAT-TERS

CUMIN, OIL OF, v. Cils, Essential. CUPFERRON v. HYDROXYLAMINE. CUPRATION. Copper albuminate.
CUPREINE v. CINCHONA ALKALOIDS.
CUPRENE (C<sub>7</sub>H<sub>6</sub>)<sub>2</sub>. A yellow substance CURARA 435

obtained by passing acetylene over reduced copper heated above 180°, and treating the brown solid so formed repeatedly with acetylene until no further change occurs.

CUPRI-ADEPTOL. Copper m-phenol sulphonate.

CUPRISIN. Trade name for a preparation of cupric cyanide and potassium cyanide, used for the treatment of trachoma.

CUPRITE. Native cuprous oxide, Cu2O, crystallising in the cubic system Brilliant, transparent crystals of a ruby-red colour (hence the name ruby-copper-ore) are not uncommon; spgr 6.0 In the variety called chilcotrichite, the crystals are capillary in form, being enormously elongated in the direction of one of the cubic edges, and are loosely matted together in plush-like masses. Another variety, known as tile-ore (Ger Ziegelerz), is earthy or compact and of a brick-red or brownish colour, it consists of an intimate mixture of cuprite and limonite, and has resulted from the alteration of copper-pyrites Cuprite is sometimes of importance as an ore in the upper oxidised portions of copper veins

CUPROCITROL. Trade name for copper citrate

CUPROL. A compound of nucleinic acid and copper

CUPRONICKEL v NICKEL.

CUPROSILICON, Cuprous silicide (v. Copper) CURARA. Curare, Urari, Woorani, Woorah,

Curara is the airow poison of those tribes of South American Indians which inhabit the districts lying between the Amazon and the It was first brought to Europe by Sir Walter Raleigh, in 1595, and from that time to this its origin and preparation have been much discussed by travellers and botanists. A summary of the earlier observations of La Condamine, A v Humboldt and Bonpland, Waterton, Schomburgk, Castelnau, Weddell, Thirion, Jobert, Crévaux, and others is given by Planchon (Pharm J [3] 11, 469, 491, 529, 589, 693, and It is shown that curara is an aqueous extract of a mixture of many plants, one of which is always a species of Strychnos, and gives to it the chief toxic power, whilst the remainder contribute those other qualities which render the extract suitable for the tipping of airow The method of preparation adopted by one tribe of Indians is described by Schomburgk The product differs some-(*ibid* [3] 10, 646) what with the locality in which it is prepared. distinguishes four varieties. curara of the Orinoco, of the Upper Amazon, of French Guiana, and of British Guiana chief active ingredient in the preparation of the first is a Strychnos plant, closely allied to Schomburgk's S toxifera (J Pharm Chim [5] 5, 20), of the second, S Castelner (Weddell), of the third, S Crevauxiana (Baill.), and of the fourth, S. toxifera (Schomb).

As seen in Europe, curara is a brown to black, nearly solid, or even brittle, extract. It has a bitter taste, and in its poisonous action it is related, though opposed, to strychnine. In medicine, it has been occasionally employed in the treatment of tetanus and hydrophobia, but it is not included in any pharmacopœia. Great care is necessary in its administration, and the solution should first be tested on animals.

For the earlier chemical investigations of curara, see Fluckiger (Arch Pharm 1890, 228, 78). Our present knowledge is mainly due to Boehm (Chem Zentr 1895, u. 1084, 1897, u. 1078; Arch. Pharm 1898, 235, 660), who showed that the alkaloids of curara can be divided into two classes: (1) compounds similar to curine, partly crystalline and partly amorphous, soluble in water with difficulty and precipitated from solution by ammonia, in which the characteristic toxic effect of curara is shown very slightly or not at all, (2) amorphous yellowish-red substances easily soluble in water as curarine, not precipitated by ammonia, and showing the most marked toxic action. Of most specimens of curara, about 70-90 p.c. is soluble in tepid water; sometimes, however, scarcely more than half is soluble Para curara, which is imported in bamboo tubes and is the variety now usually met with in commerce, contains about 11-14 pc water, 12 pc ash, 12-15 pc. curine  $C_{18}H_{19}O_3N$ , which is isolated by extracting the raw material with water, precipitating by aqueous ammonia and expectations. tracting with alcohol. It melts at 212°, and is obtained in crystals from benzene or alcohol with one molecule of the solvent or from methyl alcohol without alcohol of crystallisation Tubocurarine (paracurarine) C19H21O4N was obtained as a reddish-yellow substance which forms about 9-11 pc of raw curara. It is not identical with the previously known curarine, isolated from calabash curara The latter kind, which was sent over from the Orinoco via Venezuela in small gourds, used to be the common variety, but is now seldom met with, it is much superior to tube curare, and seems to be derived from Strychnos toxifera Earlier investigations, as a rule, refer to this The active principle, curarine C<sub>19</sub>H<sub>26</sub>ON<sub>2</sub> (?), consists of haid glistening garnet-red laminæ, decomposing and giving off a smell of trimethylamine when heated to 150°. A minute quantity of a second alkaloid was found to be present

A third variety of curara, from the Upper Amazon, occurs in the market in small jars of unburnt clay. This differs widely in respect to active principles from the other varieties. The following substances have been extracted from 1t protocurine C<sub>20</sub>H<sub>23</sub>O<sub>3</sub>N, colourless needles, which turn brown at 160° and melt, with decomposition, at 306°, protocuridine C<sub>19</sub>H<sub>21</sub>O<sub>3</sub>N, colourless prismatic crystals, m p 274°-276°; and protocurarine C<sub>19</sub>H<sub>23</sub>O<sub>2</sub>N (?), an amorphous red powder, more poisonous than the other curarines.

The mert, or slightly active, curines appear to be tertiary bases, the curarines are quaternary ammonium bases, and in this connection it is noteworthy that, according to Crum Brown and Fraser (Proc Roy Soc Edinburgh, 1869, 560), nearly all ammonium bases paralyse motor nerve endings and have a curare-like effect. In the case of curine and tubo-curarine, at least, there seems to be a close chemical relationship: both appear to be derivatives of methoxylated quinoline Curine methiodide produces the typical curare effect and resembles tubocurarine. The colour reactions of curarine indicate a relationship to strychnine. Pure curarine is extremely active; 100-100 mg. injected will paralyse a frog; given by the 436 CURARA.

mouth it is mactive. It paralyses the motor nerve endings, but, even in large doses, leaves other tissues unaffected. Hence it is much employed in physiological research (cf. e.g. Edmunds and Roth, Amer J. Physiol. 1908, 23, 28, 46).

cURCAS OIL, purging nut oil, physic nut oil (Oleum internal), is obtained from the seeds of Jatropha curcus (Linn), (purging nut'), by hot expression. The seeds consist of 66 p.c. of kernels and 34 p.c. of husks. From the former there is obtained by extraction with ether 52 p.c. of oil of a pale colour, becoming yellow, with a reddish tint, on exposure to the an Characteristic of the oil is its unpleasant odour, which assists in distinguishing curcas oil from other oils. Most prominent are the strong purgative properties of the oil, which are much more pronounced than those of castor oil. It may be distinguished from castor oil by its sparing solubility in alcohol, and ready solubility in petroleum spirit. The oil belongs to the semi-drying oils, its liquid fatty acids consist of about equal proportions of class and broken.

bark contains about four alkaloids, a little ethereal oil, a bitter principle, and a glucoside. Of total alkaloids there is 1 8 p c free and 0 6 p.c. in a combined state. By far the most abundant alkaloids are cusparine and galipine, discovered by Korner and Bohringer (Gazz chim. ital. 1883, 13, 363). They are removed from the ethereal percolate of the bark by shaking with 20 p c acetic acid, which leaves small quantities of other bases behind. The separation of cusparine from galipine is effected by fractional crystallisation of the free bases, from ligion, but principally by conversion into the oxalates, cuspaine oxalate is very little soluble in water, while galipine oxalate remains in solution (Troger and Kroseberg, Arch. Pharm. 1912, 250, 494).

Galipine  $C_{20}H_{21}O_3N$  forms colourless needles, mp 115°, when pure the salts are colourless. Oxidation with permanganate yields veratife and methoxyquinoline carboxylic acids, whence Troger and Kroseberg have suggested the following constitution:

Angostura Bitters. It is believed that Angostura Bark does not form a constituent. See article on Liqueuis and Cordials in Vol.  $\beta$ 

oligandra (Mull ) and J. multifida (Linn )

J L
CURCUMEINE v Azo- colouring matters
CURCUMIN v Turmeric.

CURCUMINES v Azo- COLOURING MATTERS CURCUPHENINE v Azo-COLOURING MATTERS CURINE v CURARA

CUSCOHYGRINE v Cocaine and the Coca

CUSPARIA BARK. Angostura bark, Carony bark. (Ecoice d'Angusture, Fr., Angostura-Rinde, Ger) The baik of the Cuspana febrifuga (Hiemb) or Galipea cuspana (St Illaire), a tree inhabiting the mountain districts drained by the Orinoco (cf Bentl a Trim Confusion has sometimes occurred between cusparia and nux vomica bark. The two barks may, however, be readily distinguished by treatment with a drop of nitric acid, which either deepens the colour of the cusparia or gives it a bluish shade, while nux vomica, similarly treated, exhibits a blood-ied colour Several other barks have, from time to time, been substituted for true angostura (For these, and the mode of detecting them, v Pereira (Mat Med 3rd ed 2, 1915); Maisch (Amer. J Pharm 1874, 50), Cazeneuve (Rép de Pharm 1874, 261); Oberlin and Schlagdenhauffen (Pharm Zeit 1878, 853)) For the bark of Esenbeckia febrifuga, which enters commerce under the name of angostura bark, see Hartwich and Gamper (Arch. Pharm. 238, 568). Cusparia

red, Frohde's reagent a deep-blue, colouration On melting its salts with organic acids, it pyrocusparine C18H15O3N, mp. changes to 255°, which base gives well-crystallised salts Like galipine, cuspărine seems to be a quinoline derivative (Troger and Beck) It is a tertiary base, and contains one methoxyl group. As protocatechnic acid is obtained by potash fusion, it is conceivable that it contains a methylenedioxy group instead of the two methoxy groups of galipine, which would be in accordance with the empirical formula Cuspareine C18H19O2N, according to Troger and Runne (Arch Pharm 1911, 249, 174), is extracted by light petroleum from the amorphous bases left after crystallisation of cusparine and galipine It melts at 56°, distils almost without decomposition, contains two methoxy groups, and yields quinoline on zinc-dust distillation Troger and Runne also describe galipoidine C<sub>19</sub>H<sub>15</sub>O<sub>4</sub>N, m.p 233°, and Troger and Beck (1c) yet another alkaloid C<sub>16</sub>H<sub>13</sub>O<sub>2</sub>M (?), m p 186°. Cusparidine and galipidine of Beckurts and Nehring are, according to Troger and Kroseberg, merely mixtures of cusparine and galipine. Other papers on angostura alkaloids are by Troger and Muller (Arch Pharm. 1910, 248, 1, 1914, 252, 459), by Beckurts and Nehring (Arch Pharm. 1891, 229, 591), and by Beckurts and Frenchs (Arch. Pharm 1905, 243, 470).

The essential oil (from Cusparia trifoliata, Engl.) is described in Schimmel and Co.'s Semi-Annual Report, April, 1913. The yield was 1.03 p.c. of the bark;  $D_{5}^{15}$  0.9285;  $\alpha_{p}^{20}$  -7.32°;

 $n_{\rm p}^{20}$  1 50744; colour, pale brown; acid number, 18; ester number, 5.5; acetyl number, 357, the oil is not completely soluble even in 9 volumes of 90 pc alcohol.

Angosturin C<sub>9</sub>H<sub>12</sub>O<sub>5</sub>, m p 58°, is a crystal-line bitter principle, readily soluble in water and in alcohol, but not in ether (Beckurts and

Nehring)

Angostura bark is a constituent of 'Angostura bitters,' and is used in medicine as a tonic in derangements of the alimentary canal, but it is not included in the BP nor in the USP

CUSSO, KOUSSO, v Kosin, art. Resins. CUSYLOL. Trade name for soluble copper citrate

CUTCH. Considerable quantities of cutch are now prepared in Burma. It is obtained from the Acacia Catechu, the trees being felled while green, the bark taken off, and the timber chopped up and boiled in large cauldrons. The resultant liquid is drained off, and solidifies In the better qualities of cutch as it cools only the heart-wood of the tree is utilised is brought to the market in several forms, the three principal ones being · (1) tablets—small rectangular blocks weighing from 1 to 2 lbs; (2) blocks-more or less square blocks weighing from 28 to 56 lbs; (3) baskets—a soft cutch of a thick consistency. Tablet cutch is the best quality, and basket cutch the lowest and basket cutch usually contain more impurities than tablet cutch On arrival in Rangoon the cutch is packed in wooden cases (usually containing 1 ewt net), and is then ready for export

Cutch trees are found throughout the whole of Burma, but the principal producing districts are Prome, Thayetmo, Myingyan, Minbu, Pegu,

Yamethin, and Pyinmana

Cutch is used largely as a dye, and for tanning, and as a preservative, especially for preserving fishing nets and sails See also CATECHU

CYANACETIC ACID v Malonic acid

CYANELLAGIC ACID Cyanellagic acid  $\rm C_{14}H_6O_{10}$  is prepared by heating cliagic or flavellagic acid with 100 p c sulphuric acid at  $185^{\circ}-230^{\circ}$  (Perkin, Chem. Soc. Proc. 22, 114), or with arsenic and sulphuric acids at 130° crystallises from pyridine in small yellow prismatic needles, and is soluble in strong alkaline solutions with a green-yellow tint, changing to deep blue on dilution. The hexa-acetyl derivative  $C_{14}O_{10}(C_2H_3O_6)$ , forms colourless needles, melts at 330°–332°, and the hexabenzoyl compound, colourless needles, m p 343°-345° Cyanellagic acid is most probably dihydroxyellagic acid, and in this case possesses the following constitution .

It dyes mordanted fabrics more readily than ellagic acid A Ğ. P.

CYANIDES. Compounds of the radical cyanogen UN The history of the cyanid

industry begins with the discovery of Prussian blue by Diesbach in 1704 (H. Kopp, Geschichte der Chemie, 4, 369) While attempting to make Florentine lake by precipitating a mixture of cochineal infusion, alum, and ferrous sulphate with an alkali, he obtained a blue colour The alkali used was potash, over which Dippel had distilled his animal oil, and Dippel recognised in this the source of the blue colouring matter. The new blue very quickly came into use in place of ultramarine, and in 1710 it was sold at one-tenth the price of the latter. The method of manufacture was a secret until 1724, when Woodward showed that the blue could be prepared by calcining blood with potassium carbonate and then treating the aqueous extract 1 with ferrous sulphate and alum, the greenish precipitate so obtained yielding Prussian blue when treated with hydrochloric acid.

This, in its essential features, was the method of manufacture which supplied the world's requirements of cyanogen compounds

for nearly 150 years

Macquer, in 1752, showed that alum is not required, and that Prussian blue, when boiled with an alkali, yields ferric hydroxide and a soluble salt from which the blue can be re-generated. This salt, potassium ferrocyanide, was first obtained in the crystalline form by Sage in 1772

The brilliant researches of Scheele, 1782-3 (Chemical Essays, London, 1786), Berthollet, 1790 (Ann Chim Phys 1790, 1, 30), and of Gay Lussac, 1815 (*ibid* 1815, 95, 156), in the course of which the relationships of the ferrocyanides, hydrocyanic acid, cyanogen, and many of their derivatives were discovered, laid a firm scientific foundation for the industry

Prussian blue was manufactured in Great Britain in 1770 by Bramwell & Co of Newcastle, it sold then at 2 guineas per lb; the price had fallen in 1815 to 10s 6d, in 1820 to 2s 6d, and in 1850 to 1s 9d Crystallised potassium ferrocyanide was first known in commerce about 1825, the price being 5, per lb, although a solution had been sold previously (Graham, Reports of Juries, Exhib of 1851, 1, 95)

The conversion of potassium ferrocyanide into potassium cyanide by fusion with potassium carbonate, was first described by F and E. Rodgers in 1834 (Phil Mag. 1834 [3] 4, 91), but the simple cyanides were not used in any considerable quantity until the process of electro-plating with gold and silver, from a cyanide solution, was discovered by J. R. and H. Elking-

ton in 1840 (Eng Pat 8447, of 1840)

An interesting application of potassium ferricyanide was discovered by Sir J Herschel in 1842 (Phil Trans. 1842) Paper, treated with a solution containing ferric ammonium citrate and potassium ferricyanide, becomes blue when exposed to light, owing to the reduction of the ferric salt and consequent formation of Prussian blue Paper sensitised in this way is used for the reproduction of engineering drawings.

A very remarkable attempt was made by Possoz and Boissière in the year 1843, to manufacture cyanides from atmospheric nitrogen;

1 Liximum sangumis or blood-lye, whence the German name of potassium ferrocyanide, blutlaugen-

this is described under the heading Potassium inch.

The year 1860 witnessed three important developments in the cyanide industry. Gautier-Bouchard (A W Hofmann, Reports by the Juries, Inter Exhib of 1862, 59-63) began to produce prussiate from the spent oxide of the Paris gas works A Gélis (Eng Pat 1816, of 1860) synthesised it from carbon disulphide and ammonia; and Margueritte and de Sourdeval (Eng Pat 1171, of 1860) attacked the problem of the fixation of atmospheric nitrogen by means of baruim oxide and carbon.

The existence of cyanogen compounds in coal gas was known in 1843, and possibly still earlier. E Jacquemyns (Annalen, 1843, 46, 236), whose attention was drawn to them by H. Rose, says that he found none in gas lime, but obtained 0.75 gram of Prussian blue per litre from ammonia liquor The purification of coal gas by means of iron oxide was introduced in 1847. Owing to the repeated use of the same quantity of oxide, the cyanogen compounds accumulate in it in a relatively concentrated form, and it was therefore naturally the first gas works product to be used in their manufacture Prussian blue, made from spent oxide, was exhibited by Gauthier-Bouchard at the London Exhibition of 1862

Both the process of Gélis and that of Margueritte and de Sourdeval were soon abandoned by their inventors, the recovery of cyanides from spent oxide, on the other hand, developed fairly rapidly, and, according to Dupré (Journ fur Gasbeleuchtung, 1884, 885), there were, in 1884, 10 works making prussiate by this means.

The competition of the gas works ferrocyamide, the increasing cost of nitrogenous organic matters (which were being more and more used as fertilisers), and the diminished use of Prussian blue due to the competition of blue colouring matters derived from coal tar, led to the disappearance of the old process in Europe about this time

In the United States of America, it appears to have held its own somewhat longer, the returns of the U.S. Census for 1900 (Bulletin No. 210, 31, 1902) show 18 works engaged in the manufacture of cyanides with a joint annual production of 2752 tons of yellow prussiate, almost entirely made from hoofs, horns, and scrap leather, only 1200 tons of spent oxide having been used

Between 1875 and 1885, the sulphocyanides, especially the tin and aluminium salts, appear to have been used in calico printing instead of the acctates (Bertelsmann, Die Technologie der Cyanverbindungen, 303), and Gélis' synthesis was taken up and improved by Gunzburg and Teherinae (Eng. Pat. 1148, of 1878, 1359, of 1879, 1261, of 1881), but without financial success.

In 1887, J S MacArthur, R. W and W Forrest (Eng Pat 14174, Oct 19, 1887; 10223, of 1888) patented their process for the extraction of gold and silver from their ores by means of a dilute solution of a cyanide. The process was carried out by allowing a solution containing 0.25 p c or less of potassium cyanide to percolate through the ore, which was usually ground to pass a sieve with 40 or 60 holes to the linear

inch. The gold is dissolved in the form of potassium aurocyanide, and is recovered from the solution by passing it through a mass of zinc shavings. The process was first put into practice at Karangahake, New Zealand, in 1889, and near Johannesburg, Transvaal, in 1890. The success of these installations led to a very rapid adoption of the process in all the gold-producing countries of the world.

Prior to the introduction of this process, the cyanide industry was quite a small one. The world's consumption of potassium cyanide (produced exclusively from ferrocyanide by the Rodgers process) (F. Roessler, V. Internationaler Kongress für Angewandte (Themie, Berlin, 1903, 1, 638) did not exceed 50–70 tons per annum (G. Beilby, J. Soc. Chem. Ind. 1898, 134) and that of yellow prussiate was probably under rather than over 5000 tons (worth together about 325,000l).

Ten years later, just before the Transvaal War, the European production of potassium eyanide was estimated by Beilby (V Inter Kongiess Angew Chem Berlin, 1903, I, 630) as 5500 tons, adding the American production of about 1000 tons (U S Census Bulletin, No 210, 31, 1902), we arrive at a total production of 6500 tons

The increased quantities of potassium cyanide required by the gold-mines were at first supplied by fusing potassium ferrocyanide with potassium carbonate, and very soon afterwards by E. Erlenmeyer's process (Ber. 1876, 1840), in which dry prussiate and metallic sodium are used: K<sub>4</sub>Fe(CN)<sub>6</sub>+2Na=2Na(N+4KCN+Fe Synthetic processes producing cyanide direct without the intervention of prussiate, were also proposed in great numbers. The approximate numbers of patents connected with cyanide manufacture applied for were 1889–1898, 104, 1899–1908, 129, 1909–1917, 106. The principal processes which have been worked with some measure of success, are—

Process	Where worked	Began pro-
Siepermann	Stassfurt, Germany.	1892
Beilby.	Glasgow, Scotland	1892
British Cyanides	Oldbury, England	1895
Raschen.	Runcorn, England.	1898
Readman.	Leven, Scotland.	1899
Castner.	Germany, Scotland, USA	1899-1900
Bueb ('Schlempe').	Dessau, Germany.	1902

These, and others, are described fully below

The recovery of cyanides from the products of distillation of coal was also improved by the introduction of special cyanide scrubbers during the decade 1890-1900, whereas the iron oxide purifiers had yielded but little more than one-half of the cyanogen in a useful form, the newer processes give an almost quantitative recovery.

About the year 1900 the fruit trees of California and the Southern States of America were threatened with extermination by a parasitic scale. The investigations of the Experiment Stations of the US Department of Agriculture showed that this could be killed without injury to the trees by fumigation with hydrocyanic acid. The method, which was introduced

about 1903, is now very widely used, both in the United States of America and in other countries. It has also been extended in other directions, such as killing insects in hot-houses, and freeing railway carriages and houses from vermin (in South Africa). Details will be found under the heading Hydrocyanic acid Statistics collected by George P. Gray show that the consumption of sodium cyanide for fumigating citrus trees during 1916 in the State of California alone exceeded 1300 tons (J Ind. Eng Chem. 10, 301, 1918).

Another application which may be mentioned is in case-hardening iron; immersion in fused cyanide produces a very hard skin on mild steel, and both cyanides and ferrocyanides have long been used for this purpose in ordinary workshop practice. According to J E Bucher (J Ind Eng Chem 9, 236, 1917), the reversible reaction  $2\text{NaCN} \rightleftharpoons 2\text{Na} + 2\text{C} + \text{N}_2$  takes place in presence of iron; if pure iron is heated with sodium cyanide the latter is decomposed and the carbon combines with the iron, if, on the other hand, carbonised iron is heated with sodium and introgen, the carbon is removed from it, and cyanide formed.

Potassium cyanide, as used in electro-plating, was a material having the appearance of unglazed porcelain, and containing from 30 to 70 p c. of KCN, together with large quantities of carbonate and cyanate The material supplied to the gold-mines in the early days of the gold-extracting process was a black mass containing some 75 p c of KCN, and all the iron and carbon liberated by the decomposition of the ferrocyanide This was replaced by the pure white crystalline product of the Erlenmeyer process, which contained about 2 molecules of KCN to 1 molecule of NaCN, the cyanogen content being about 40 pc, equivalent to 98-100 pc of KCN. The earlier produce of the Beilby process was in the form of cakes, which were coloured black by a small percentage of charcoal Later, by filtration of the molten cyanide, the charcoal was removed, and the product was white or pale grey, and free from insoluble imputities Shortly after the introduction of the Castner process, in 1900, solid cakes of pure white crystalline sodium cyanide, containing 97 p c. to 98 p c of NaCN (equivalent to 129-130 p c of KCN), became an article of commerce, and this is at present the form in which the greater part of the cyanide employed in gold-extracting is used Sodium cyanide, obtained by the evaporation of aqueous solutions, as in the Readman and 'Schlempe' processes, is compressed into briquettes, which contain about 90 pc of NaCN.

Potassium ferrocyanide has also been replaced to a very large extent by the sodium salt which is quite easily prepared by the more recent methods of extracting the hydrocyanic acid from coal gas by special cyanogenwashers

The rapid rate of increase in the world's consumption of cyanide which followed the introduction of the gold-extracting process has not been maintained. Since 1909 the annual value of the world's gold output has been almost stationary at a little more than 90,000,000, and improvements in the process of extraction have everywhere tended to diminish the quantity

of cyanide needed to produce this. On the Rand, where this tendency seems to have almost reached a limit, the average consumption of cyanide is 0 4 lb. KCN per ton of ore treated (H A. White, J. Chem Met and Min Soc S Africa, 1915, 16, 24). Increased use in the treatment of silver ores (much delayed, however, by political troubles in Mexico) and in agriculture have kept the world's consumption from 1910 to 1920 approximately constant at, in round figures, 20,000 tons per annum. In 1912 and 1913 Germany exported 6700 tons per annum (The Mineral Industry, 1917, 25, 882) The deficit caused by the cessation of these exports on the outbreak of war in 1914 has been covered by increased production in Great Britain; a temporary shortage in the United States was soon overtaken by the American makers

Manufacture of Potassium and Sodium Ferrocyanides—(1) From nitrogenous organic

matter

The method of manufacture from nitrogenous organic matter is now only of historical interest

The sources of nitrogen were horn, dried blood, hair, waste wool, and feathers (15–17 p.c. N), woollen rags (10–16 p c N), pigs' bristles (9–10 p c N), and old leather (45–7 p c N). These materials were sometimes charred before use, about 80 p c. of the nitrogen being driven off in the form of ammonia, hydrocyanic acid, and a complicated mixture of nitriles and organic bases

The process was carried out at first in pearshaped, cast-iron muffles, which were replaced later by reverberatory furnaces, the beds of which were formed of cast-iron pans, 5 feet long, 3 feet wide, 4 inches deep, and about 2 inches A charge of 2 cwt of potash (usually made up of 2 parts of the 'blue salt' recovered in a later stage of the process, and 1 part of fresh potash) was put into the pan and fused with 18-20 lbs of non turnings; 2 cwt of the dry nitrogenous materials were then stiried in gradually (25 to 3 cwt of charred material could be added without making the mass too The temperature was then raised in order to complete the reaction, and the melt ladled out into iron moulds After solidification, it was broken up and digested for 24 hours with water at 50°-60°, and finally extracted completely with boiling water The solution obtained contained potassium ferrocyanide, sulphocyanide, carbonate and sulphide 1t was evaporated to sp gr 1 28, and allowed to crystallise Further evaporation yielded a second crop of impure crystals, the mother liquor of which was evaporated to dryness. leaving a mixture of potassium carbonate and sulphide, known as 'blue salt' which was returned to the fusion pan

The black insoluble residue left from the extraction of the 'metal,' as the product of the fusion was called, contained considerable quantities of potassium, mainly in the form of potassium ferrous sulphide and of double potassium, calcium, and aluminium silicates.

potassium, calcium, and aluminium silicates.

The quantity of 'metal' produced was very little greater than the weight of potash used, and 4 charges were put through in 24 hours, but practice varied considerably: C Karmrodt (Wagner's Jahresber. 1857, 3, 139) speaks of

6 charges of 500 lbs, each as the normal output of a pan in 24 hours

The yield of ferrocyanide was seldom more than 20 pc of the quantity theoretically obtainable from the introgen in the materials used when they were not previously charred. Using charred materials, as much as 57 pc of the remaining nitrogen was converted into ferrocyanide, corresponding with a yield of about 12 pc on the nitrogen originally present. The quantity and composition of the black

residues varied greatly with the kind of organic matter used The main constituents were carbon (4-9 pc), ferrous sulphide, iron, and inorganic substances derived from the ash of the animal matters and adhering dirt, the principal being silica, lime, alumina, and phosphoric acid The quantity of potassium combined with these substances in an insoluble form was considerable According to Karmrodt (Wagner's Jahresber Recording to Karmiott (wagners salresser 1857, 3, 139), 100 parts of 'metal' made (1) from horn gave 18 7 pc of dry residue, containing 12 2 pc of  $K_2O$ , (2) from rags, 28 3 pc of residues, with 16 7 pc  $K_2O$ ; (3) from leather, 35.1 p c of residue, with 10.2 p c K<sub>2</sub>O quantities represent about  $4\frac{1}{2}$ , 9, and 7 pc. of the potassium contained in the original charge, and since only some 10 pc of this is converted into ferrocyanide at each operation (R Brunquell, Wagner's Jahresber 2, 102, 1856), the loss of potash in the residues appears to have been from 45 to 90 p c of that usefully employed Potash was also carried away with The iron pans survived several the flue gases hundred operations

The theory of the process was first given by Liebig (Annalen, 1841, 38, 20). Scheele (Chemical Essays, London, 1786, reprinted 1901) had already shown that lixivium sanguinis is essentially a solution of potassium cyanide and that the addition of a ferrous compound is necessary to convert it into prussiate Liebig confirmed this by extracting the 'metal' with alcohol, which dissolves out potassium cyanide, leaving a residue from which no ferrocyanide can be obtained. He attributed the formation of the evanide partly to the action of ammonia (formed by destructive distillation of the organic matter) on carbon, C+NH<sub>3</sub>=HCN+H<sub>2</sub>, and partly to the action of potassium on the nitrogenous residue of carbon It may be remarked that the temperature of the fusion could not have exceeded 1000°, since cast iron melts about 1100°, and the pans withstood some hundreds of The temperature, too, was lowered by the addition of the organic matter, occasionally sufficiently to cause the charge to solidify, so that the reaction between ammonia and carbon, which requires a temperature of about 1000°, could not have occurred to any great extent. The direct action of ammonia on potassium carbonate would, however, take place, although, the ammonia being evolved on the surface of the mass, the conditions were not favour-As to the second of Liebig's able to it reactions, the writer has found that potassium is not formed in appreciable quantity from a mixture of charcoal and potassium-carbonate below about 1200°, certainly not at 1000°, so that the greater part of the cyanide must have been formed by the action of potassium carbonate itself on the unknown carbon-nitrogen com-

pounds contained in the charred organic matter. That cyanide may be formed in this way is indicated by the fact that carbazol, when heated to redness with caustic potash, yields considerable quantities of it (G Kraemer, Eng Pat. 16529, of 1894). The rôle of sulphur in the process was explained by Liebig thus: potassium sulphate, which is always present in the potash, is reduced by carbon to sulphide, carbonate, and polysulphide The latter dissolves iron, forming a double sulphide of iron and potassium which, when the product is lixiviated, dissolves in the potassium cyanide solution thus:

If insufficient iron is present in the fusion, the polysulphide is not entirely decomposed, and potassium sulphocyanide is also formed R Hoffmann (Annalen, 1860, 113, 81) investigated the behaviour of the sulphur compounds in the process in some detail, and found that the sulphur of potassium sulphide is rapidly and completely removed by iron alone in the fusion process, and that the same is true of potassium sulphiate in presence of carbon. When organic matters are added to the sulphurfree melt, however, sulphocyanide is produced in proportion to the quantity added Hoffmann, therefore, attributes its formation to the organic sulphur

The behaviour of potassium sulphocyanide in the fusion is not clearly understood. Hoffmann found that it is completely decomposed by fusion with excess of potassium carbonate, yielding cyanate and sulphide, 1 but if the mixture is again heated with carbon, the sulphocyanide is regenerated. It appears, therefore, that, under the conditions of the process, sulphocyanide is not decomposed. The reaction between iron and sulphocyanide is described below.

A part of the sulphide of iron formed is present in the form of a potassium ferrous sulphide, FeS xK $_2$ S, which dissolves in hot water to a dark-green, colloidal solution which is coagulated by cooling or by adding excess of potassium carbonate. Another part is probably present in the form of the crystalline potassium ferric sulphide K $_2$ S Fe $_2$ S $_3$ , obtained by K. Preis (J. pr. Chem. 1869, 107, 10), by fusing together finely divided iron, potassium carbonate, and sulphur. He found that the crystalline sulphide was formed at very high temperatures, and the amorphous one, yielding the green solution, at lower temperatures.

The reaction between potassium cyanide and ferric sulphide

13KCN + Fe<sub>2</sub>S<sub>3</sub> = 2K<sub>4</sub>Fe(CN)<sub>6</sub> + KSCN + 2K<sub>2</sub>S

would give a ratio of cyanogen in the form of sulphocyanide to cyanogen in the form of ferrocyanide of  $1\cdot 12$ , whilst Hoffmann, in 10 experiments, found ratios varying from  $1\cdot 5$  to  $1\cdot 18$ 

The improvements in the process which were proposed aimed at saving the ammonia evolved in the early stages by distilling the organic matters separately and passing the ammonia

 $^{\circ}$  I can confirm the accuracy of this observation Equal molecules of potassum carbonate and sulphocyanide were fused together in a closed aluminium crucible at 500°-580° for  $1_{1}^{\downarrow}$  hours, a little more than half of the materials had reacted in accordance with the equation KSCN+K<sub>2</sub>CO<sub>3</sub>=KCNO+K<sub>2</sub>S+CO<sub>2</sub>.—T E.

through a red-hot mixture of potash and charcoal (Brunquell, Wagner's Jahresber 1856, 2, 102), or a similar mixture with iron added (Karmrodt, ibid 1857, 3, 139), or over red-hot charcoal forming hydrocyanic acid which was to be absorbed, together with the unchanged ammonia, in a solution of ferrous sulphate (Brunquell) These proposals, unfruitful at the time, are interesting as foreshadowing processes which came into successful operation nearly half a century later

The belief, which prevailed until about the end of the nineteenth century, that cyanogen compounds are more easily produced from potash than from soda, is not entirely in accordance with the older experimental evidence The truth appears to be that equally good results can be obtained with sodium carbonate under suitable conditions, but the conditions are not so easily

L Possoz (Compt rend 1858, 47, 209) obtained, under manufacturing conditions, 5 parts of sodium prussiate, instead of 25 parts of the potassium salt, from 100 parts of horn Hoffmann (Annalen, 1860, 113, 81), using a very high temperature and sodium carbonate, got the equivalent of 11 parts of potassium ferrocyanide from 100 parts of woollen rags, against 10 to 14 parts with potash, and S Tanatar (Dingl poly J 1880, 237, 234) obtained the results tabulated below, as the means of 3 to 5 experiments:-

Blood charcoal	K <sub>2</sub> CO <sub>3</sub>	Na <sub>2</sub> CO <sub>3</sub>	NaCl	CaCO <sub>3</sub>	Prussiate obtained
10	$25^{\circ}$				2 15
10		25			0.2
10		5	25		2 17
10		5	25	3	23

Manufacture of Potassium and Sodium Ferrocyanides -(2) From the products of distillation of coal The coal used in the manufacture of illuminating gas may be taken to contain from 1 to 19 pc of nitrogen (Drehschmidt, Journ fur Gasbeleuchtung, 1904, 677; McLeod, J Soc Chem Ind 1907, 26, 137) When the coal is distilled, a part of this nitrogen passes into the gas in the form of ammonia, hydrocyanic acid and free nitrogen, a small part is found in the tar, and the remainder in the coke The distribution of the nitrogen between these products varies considerably, being influenced by the nature of the coal, the amount of moisture in it, the temperature of distillation, and the kind of retort used The following table indicates the magnitude of the quantities involved when the coal is distilled dry:

Ne	Wind of soil	P c of the total N of the coal in					
	Kind of coal	Coke	NH <sub>3</sub>	HCN	Tar	Gas	
1 2 3 4	Durham . Westphalia — Scottish .	48·7 50 48 58·3	14 5 12-14 15 17·1	1 6 1 8 2 1 5	15 - 39	35 2 30 0 35 19 5	

 Foster, J Gas Lighting, 1882, 1081 2. Knublauch, J. Soc Chem Ind 1896, 15, 106.

3. Chief Inspector of Alkalı Works, 43rd

Report, 1906, 58

4. J McLeod, J. Soc Chem Ind 1907, 26, 137; and Alkalı Works, 44th Report, 1907, 215

The effect of temperature on the distribution of the nitrogen has been investigated by O Simmersbach (J. Soc Chem Ind 1915, 34, 895), using a Silesian coal containing 1 396 p c of N The following table contains some of his results

Temp		P c of	the tota	ıl nıtrogen	
	as NH3	as HCN	ın tar	as N <sub>2</sub> in gas	ın coke
600	7 81	0 25	2 12	18 13	71 69
$\frac{800}{1000}$	21 28 23 15	$\begin{array}{c} 0.87 \\ 1.23 \end{array}$	3 47 4 11	10 73 21 53	63 65 49 98
1200	22 84	1 42	4 21	45 10	26 43

Even the ammonia liquor from low temperature carbonisation retorts working at 427°-455° contains traces of cyanogen compounds (Alkali Works, 45th Report, 1908, 52)

Hydrocyanic acid is converted into ammonia by water vapour at the temperature of distillation used in gas works and coke ovens

$$HCN + H_2O = NH_3 + CO$$

(Carpenter and Linder, J Soc Chem Ind 1905, 24, 63) The gas from coke ovens distilling coal with some 12 p c of moisture therefore contains about two-thirds of the quantity of hydrocyanic acid found in the gas made from almost dry coal in gas works

The gas leaving the retorts contains the whole of its cyanogen in the form of hydrocyanic acid (Alkali Works, 43id Rep. 1906, 33), the fate of this depends on the process of purification adopted In common practice about one-third of it is condensed along with the ammonia in the virgin liquor and ammonia scrubbers in English works (Alkalı Works, 43rd Rep 1906, 61), but less in many German works, where less water is used in the scrubbers (Bertelsmann, Technologie der Cyanverbin-dungen, 171) Thegreater part of this is converted into sulphocyanide, and escapes finally in the effluent from the ammonia stills From 50 to 70 p c is caught in the iron oxide purifiers, the remainder passing away with the gas, where it is a source of damage to meters and gas-holders A more complete recovery is obtainable when one of the cyanide washers described below is placed before the ammonia scrubbers Up to 1908 attempts to recover cyanogen compounds from coke-oven gases had been unsuccessful (Alkalı Works, 44th Rep 1907, 40)

The total quantity of cyanogen compounds formed during the destructive distillation of coal is very considerable. In 1912 the world's production of ammonium sulphate was 1,144,000 tons. In Great Britain over 70 p c of the production came from gas works and coke ovens in which roughly one molecule of hydrocyanic acid accompanies ten molecules of ammonia Assuming the same conditions in other countries, it appears probable that hydrocyanic acid equivalent to 80,000 tons of KCN is formed annually; only a small part of this is actually

Formation of HCN from Ammonia and Carbon.

-The earliest synthesis of hydrocyanic acid was effected by Clouet (Ann. de Chim. 1791, 11, 30), by passing dry ammonia over wood charcoal heated to bright redness in a porcelain tube. This reaction has been studied quantitatively by E Bergmann (Journ. fur Gasbel 1896, 39, 117), D Lance (Compt rend 1897, 124, 819), G A. Voerkelius (Dissertation, Hannover, 1909), and Carpenter and Linder (Alkalı Works, 45th Rep. 1908, 26; 46th Rep. 1909, 21) The formation of hydrocyanic acid begins at about 700°, and the best yields, calculated on the ammonia used, are obtained at temperatures between 1000° and 1100° The reaction NH<sub>3</sub>+C  $\rightleftharpoons$  HCN+H<sub>2</sub> is reversible, and should lead to an equilibrium between ammonia, hydrocyanic acid, and hydrogen, the equilibrium, however, cannot be realised, because both ammonia and hydrocyanic acid are unstable at the temperatures required The final result depends, therefore, on the rates of decomposition of ammonia and hydrocyanic acid and on the rate of reaction between ammonia and charcoal Even at 1000°, the rate of decomposition of hydrocyanic acid in presence of charcoal is small, but in presence of other substances it may be large Voerkelius, for example, passed a mixture of hydrocyanic acid and hydrogen through an unglazed porcelain tube at 1000°, and found that over 70 p.c of it was decomposed, but after a layer of carbon had been deposited on the tube the decomposi-tion fell to 19 p c. The rate of decomposition of the ammonia is even more variable; for example, at 720°, 28 pc of the ammonia survived 6 minutes' contact with fresh finely ground wood charcoal, whilst 82 pc unchanged when the charcoal was in the form of lumps which had been already used rate of decomposition of the ammonia is diminished by diluting it with an indifferent gas

In the 66 experiments recorded by Voerkelius, the ammonia (diluted with hydrogen) was in contact with the charcoal for from 16 to 360 seconds, within these limits, the ratio between the numbers of molecules of ammonia and of hydrocyanic acid in the product is roughly independent of the time of reaction, but the quantity of ammonia decomposed into its elements increases with the time of contact The following table is calculated from the figures The figures show the given by Voerkelius influence of the quality of the charcoal on the

result :-

		Mols NH <sub>3</sub>	
Tempera-	Pc NH <sub>3</sub>	Mols HCN in product	
ture	in gas used	in product	
720° charcoal A	50	67-134 n	nean 9 80
"В	50	29 - 45	,, 37 7
sugar char	coal 50	71	,, 71
1020° charcoal C	13-15	0 64-0 92	,, 0.76
" D	13-15	0 90-1 80	,, 140
" C	43-47	0 50-0 90	,, 061
1120° ,, D	13 4	0 44-0 48	,, 0 46

Nernst's thermodynamic theorem (Applications of Thermodynamics to Chemistry, London) 1907, and Theoretische Chemie, 5th and 6th ed has been applied to this reaction by v. Wartenberg (Zeitsch anorg. Chem 1907, 52, 299). Since the reaction NH<sub>3</sub>+C=HCN+H<sub>2</sub> absorbs 39,500 cals per molecule, the equilibrium must

lead to the formation of more hydrocyanic aci at higher temperatures, the calculation indicate that the ratio of the partial pressures (in atmo spheres) of the gases in equilibrium should hav the following values -

	$p_{ m NH_3}$
Temperature	$\overline{p_{ ext{HCN}}  imes p_{ ext{Ho}}}$
500°	632
800°	0 27
1100°	0 003
1400°	0 0002

In the experiments tabulated above, the fine mixture of gases usually contained nearly 95 p of hydrogen, the partial pressure of which we therefore almost 1 atmosphere The molecule ratio of ammonia to hydrocyanic acid is, there fore, almost the same as the ratio of the partie pressures. A comparison of the experiments values with those calculated shows that equil brium is never even approximately attained They agree in indicating 700° as the lower temperature at which appreciable quantities c hydrocyanic acid can be formed

Carpenter and Linder's independent invest gation yielded results in close agreement wit those obtained by Voerkelius It seems fairl certain that the greater part of the hydrocyani acid found in the products of destructive dis tillation of coal is formed by the reaction which has just been discussed. The presence of sma quantities of HCN in the products formed a temperatures below 600° indicates, however that it may be, in part, a primary product c

the decomposition Spent oxide. The hydrated oxide of iro first used in gas purification was made by mixin ferrous sulphate and slaked lime, sawdust bein added to increase the porosity (Laming, Eng Pat 11944, of 1847), natural iron ores were intro duced by Howitz in 1870 Sulphuretted hydro gen is absorbed, yielding a mixture of ferrou and ferric sulphides which is regenerated b exposure to air and moisture, free sulphur an ferrous and ferric hydroxide being produced Ferrocyanide is formed from the ferrou hydroxide and ammonium cyanide, and the converted into Prussian blue by interaction wit the ferric iron Sulphocyanides are also forme in very variable amount, doubtless by the inter action of hydrocyanic acid and ammoniui polysulphide. Since ammonia, sulphurette hydrogen, and free sulphur are always preser in the purifiers, it is obvious that polysulphide of ammonium can always be formed; it appear to be generally allowed, however, that sulphe cyanide is only produced in large amount whe the mixture becomes warm, as, for example when the regeneration proceeds rapidly in large mass of material The spent oxide finall contains 30 to 40 pc of sulphur (or more soluble ammonium salts and Prussian blue, i addition to tar and iron oxides Good matern contains 11 to 12 p c of Prussian blue, calculate on the air-dried substance, whilst ammonium

As already mentioned, prussiates were fire produced from spent oxide by Gautier-Bouchar about 1860 Ammonium salts were first re moved by washing with water; the residue wa then mixed intimately with sufficient lime t

sulphocyanide varies from traces to 10 p c.

convert the Prussian blue into calcium ferrocyanide and the mixture submitted to systematic lixiviation with cold water. This operation must be done in the cold, because otherwise calcium polysulphide is produced from the lime and free sulphur, and this converts the ferrocyanide into sulphocyanide. The calcium ferrocyanide was converted into the potassium salt by means of potassium carbonate, and the sulphur in the extracted residue burned to sulphur dioxide and converted into sulphuric acid (A. W. Hofmann, Reports by the Juries, Inter Exhib. of 1862, 59-63). This process contains the essential features of all those which have since been worked successfully

T. L G. Bell (Eng. Pat 257, of 1868) proposed to extract the sulphur with carbon disulphide, and, after removing soluble ammonium salts with water, to boil the residue with lime In order to obtain a pure product, the solution was then acidified and precipitated by the addition of ferric chloride, the Prussian blue being then converted into potassium ferro-The preliminary extraction of the cyanide sulphur has occasionally been carried out, but it is difficult to separate the sulphur from the tar which is extracted along with it. In spite of the great difficulty of filtering and washing Prussian blue, this method of purification appears to have been fairly generally used up to the year 1883, when Kunheim and Zimmer-mann (Eng Pat. 3342, of 1883) described two new methods.

In the first, the ammonia is only partially removed from the spent oxide by the preliminary washing, enough being left behind to produce the double salt Ca(NH<sub>4</sub>)<sub>2</sub>Fe(CN)<sub>6</sub>. The extraction with line then yields an ammoniacal solution of calcium ferrocyanide, which, when neutralised with hydrochloric acid and heated to 80°, deposits the very sparingly soluble calcium ammonium ferrocyanide, which is comparatively easily filtered and washed

By the second method, the solution of calcium ferrocyanide, obtained by following Gautier-Bouchard's original procedure, is treated with a solution of potassium chloride which precipitates the double salt  $\text{CaK}_*\text{Fe}(\text{CN})_6$  in minute crystals, which are very readily washed and are then converted into potassium ferrocyanide by boiling with potassium carbonate By this means one-half of the potassium carbonate is replaced by the cheaper chloride, and a very pure product is obtained (Franz Bossner, Die Verwerthung der Ausgebrauchten Gasreinigungsmassen, 1902). From the calcium ammonium salt, sodium ferrocyanide is readily obtained.

During the decade 1885 to 1895, the spent oxide of the gas works was the principal source of prussiates; its importance, however, has been diminished by the introduction of the more efficient wet methods of recovering the hydrocyanic acid.

Cyanide Washers —The first practicable proposal of this kind was made by O Knublauch (Eng Pat. 15164, of 1887). Coal gas was to be scrubbed with a solution of the oxide, hydroxide, carbonate, or sulphide of ammonium, or of an alkali or alkaline-earth metal containing the hydroxide or carbonate of iron, zinc, or manganese in suspension. The process did not succeed

at the time, probably for commercial reasons. The very similar process of Foulis (Eng. Pat. 9474, of 1892; Foulis and Holmes, Eng Pat. 15168, of 1895) has since been introduced at several gas works with satisfactory results. In this process, the reagent required is prepared by precipitating a ferrous salt with sodium carbonate The washed precipitate (a partially oxidised mixture of ferrous carbonate and hydroxide) is suspended in a solution of sodium carbonate and the gas passed through it in a rotary scrubber Foulis wished to obtain the whole of the ferrocyanide in the form of a solution of the sodium salt, and therefore placed the cyanogen washer after the ammonia scrubbers because, in presence of ammonia, an insoluble ammonium ferrous ferrocyanide is formed which is only decomposed by sodium hydroxide, but not by the carbonate At the Hague gas works, however, this order was reversed in order to catch the hydrocyanic acid which is otherwise condensed along with the ammonia (Rutten, Journ of Gas Lighting, 1902, 80, 879, Johnsen and Rutten, Journ für Gasbeleuchtung, 1903, 716) The saturated liquor from the cyanide washer contains Prussian blue and potassium and ammonium ferrous ferrocyanides in suspension and potassium ferrocyanide and ammonia in solution (potassium carbonate is used instead of the sodium salt) This mixture is filter-pressed and the solution distilled to separate the ammonia and then evaporated in vacuo and crystallised. The press cake is boiled with caustic potash and the ammonia collected, the residual iron oxide returning to the cyanide washer, and the solution of ferroeyanide going to the evaporating plant

In another set of proposals, the use of an alkalı other than the ammonia of the gas is dispensed with, the crude coal gas being simply washed with a solution of ferrous sulphate Since ammonia and sulphuretted hydrogen are present in much larger quantities than hydrocyanic acid, the ferrous sulphate is soon converted into a suspension of ferrous sulphide in a solution of ammonium sulphate, and the subsequent changes may be regarded as due to the action of a solution of ammonium cyanide on ferrous sulphide If a large excess of the latter is present, the main product is ferrous ferrocyanide Fe<sub>2</sub>[Fe(('N)<sub>6</sub>]; as the ferrous sulphide is used up, ammonium ferrous ferrocyanide (NH<sub>4</sub>)<sub>2</sub>Fe[Fe(CN)<sub>6</sub>] is formed in increasing quantities, and finally, when but little FeS remains, ammonium ferrocyanide is found in solution (Feld, Journ. fur Gasbeleuchtung, 1904,

The earlier patentees of processes based on these reactions attempted to obtain the ferrocyanide in a soluble form by using a dilute solution of a ferrous salt, and so having a large excess of ammonia present (Rowland, Eng. Pat. 22347, of 1891; Lewis and Cripps, Eng. Pat. 20883, of 1896; Schroder, Eng. Pat. 19849, of 1898, Teichmann, Eng. Pat. 12485, of 1899), but since insoluble compounds are always formed to some extent, it appears to be easier to obtain all the fefrocyanide (or the greater part of it) in the insoluble form by using a saturated solution of ferrous sulphate, and so having a large excess of iron present as first proposed by Bueb (Eng. Pat. 9075, of 1898). Bueb's process

has been adopted by a considerable number of gas works It is said to absorb 98 pc. of the hydrocyanic acid in the gas, yielding a mud containing 6-7 pc of ammonia and cyanogen compounds, equivalent to 18-20 pc of yellow prussiate of potash. A small part of the ferrocyanide is in solution. This is precipitated by boiling the mixture with ferrous sulphate, the precipitate is then filtered off and treated with lime, the calcium ferrocyanide solution obtained being worked up in the way already described when dealing with spent oxide

As already mentioned, part of the hydrocyanic acid in coal gas is always converted into sulphocyanide owing to liberation of free sulphur by the action of atmospheric oxygen on sulphuretted hydrogen. By suitable means the whole of the hydrocyanic acid may be fixed in the form of sulphocyanide, which is then converted into ferrocyanide by one of the processes

described in the next section

In 1877 W Marriott (Eng. Pat. 3311, of 1877) proposed to scrub gas with a solution of sodium polysulphides, and in 1888 Campbell and Boyd (Eng. Pat. 10186, of 1888) proposed the use of manganese dioxide suspended in water This liberates sulphur from a part of the sulphuretted hydrogen in the gas, and this free sulphur combines with ammonium sulphide to form polysulphid, which converts the cyanide into sulphocyanide The final step, which led to a successful and economical process, was made by H Wood Smith (H. Wood Smith, Gidden, Salamon, and Albright, Eng Pat 13658, of 1901) by simply scrubbing the gas through a suspension of sulphur, whereby the hydrocyanic acid is removed quantitatively, and a solution containing as much as 200 grams of ammonium sulphocyanide per litre may be obtained. This process is in use at several gas works, and yields practically the whole of the sulphocyanides made at present

Manufacture of Sodium and Potassium Ferrocyanides—(3) From Sulphocyanides A Gélis (Eng Pat 1816, of 1860) was the first to manufacture potassium ferrocyanide by heating dry potassium sulphocyanide with iron, made by reducing the oxide, at a dull-red heat, in a closed vessel He states that too high a temperature leads to considerable loss fusion process yields a mixture of potassium cyanide and ferrous sulphide which gives potassium ferrocyanide, when treated water in the same way as the 'metal' of the old process (A W Hofmann, Reports by the Junes, Inter Exhib of 1862, 59-63) Sir G Beilby has informed the writer that he made considerable quantities of potassium ferrocyanide by this process in the years 1873-1876 Potassium sulphocyanide was heated with iron turnings in air-tight iron retorts and, after cooling sufficiently, the product was extracted with water in the retort itself, without coming in contact with In this way the only serious difficulty of the process, which is due to the extreme proneness to oxidation of the fusion product, is avoided and satisfactory yields obtained.

The process was taken up again in 1878 by Gunzburg and Tchermac (Eng. Pat 1148, of 1878, 1359, of 1879, 1261, of 1881) They dried the sulphocyanide by fusing it at 300°, and then mixed it intimately with the oxide-free

iron by grinding them together. The mixture was heated in iron boxes with closely fitting covers at 445°, the vapour of boiling sulphur being used to obtain a constant temperature. After working for two or three years, the process was given up, but it was revived for the third time in 1894 by the United Alkali Co, Ltd. (Hetherington and Muspratt, Eng Pat. 5830, of 1894), and by the British Cyanides Co, Ltd. (Crowther, Rossiter, Hood and Albright, Eng. Pat 8305, of 1894) The patent specifications of Hetherington and of Crowther add but little to the information given above In both cases, some 20 to 30 p c excess of iron is used, and the temperature is almost the same as that recommended by Gunzburg and Tchermac Crowther says that mere fusion is insufficient to dry the sulphocyanide; a current of an indifferent gas must be passed through it for some time Conroy (J Soc Chem. Ind 1896, 15, 12) was able, in a small scale experiment, to convert 76.8 p.c of the sulphocyanide used into ferrocyanide (leaving 18.2 p.c unchanged) by means of Hetherington's method The process has been worked by the British Cyanides Co at Oldbury since 1894 (Brussels Exhib 1910. Catalogue of British Chem Section, 155)

The conversion of sulphocyanides into ferrocyanides is also possible in aqueous solution Sternberg (D R P 32892, of 1882), found that by heating a solution of potassium sulphocyanide with double the theoretical quantities of iron filings and ferrous hydroxide under pressure at 110°–120°, about 80 pc of the material could be converted into ferrocyanide

in accordance with the equation

 $6KCNS + 6Fe + Fe(OH)_2 = 6FeS + K_4Fe(CN)_6 + 2KOH$ G R Bower (Eng Pat 361, of 1896) succeeded in removing the sulphur from ferrous sulphocyanide by means of copper in a similar way Fe(CNS)<sub>2</sub>+2Cu=2CuS + Fe(CN)<sub>2</sub> Conroy (Conroy, Hurter, and Brock, Eng Pat 3867, of 1896, J T Conroy, J. Soc Chem Ind 1898, 17, 98) then found that a mixture of a soluble sulphocyanide and a ferric or ferrous salt would do as well as pure ferrous sulphocyanide, and also that the sulphur is more rapidly removed by finely divided iron than it is by copper Using very finely divided iron, the reaction takes place quantitatively in 5 to 6 hours at a temperature of 135°-140° and a pressure of 50 to 60 lbs per square inch, in accordance with the empirical equation  $Fe(CNS)_2 + 2Fe = Fe(CN)_2 + 2FeS$  (Sodium or potassium sulphocyanide is acted on by iron under similar conditions very slowly and incompletely) The black mud produced is either boiled with caustic alkali to decompose the ferrous ferrocyanide, or the ferrous sulphide is first dissolved in hydrochloric acid. latter course is to be preferred, owing to the fact that ferrous sulphide combines with the alkalis, each 3 or 4 equivalents of it rendering 1 equivalent of either caustic soda or potash insoluble, a difficulty with which the manufacturers of ferrocyanide from nitrogenous organic matter had long been familiar

#### PROPERTIES OF THE FERROCYANIDES.

The principal characteristics of those salts which possess some technical interest are described here; further information is to be

found in Cyanogen Compounds, by H E Williams, 1915, or in the text-books of inorganic chemistry.

Ammonium ferrocyanide is readily prepared in aqueous solution from the free acid and ammonia, or from barium ferrocyanide and ammonium sulphate. The solid salt

$$(NH_4)_4$$
Fe $(CN)_6$ ,  $3H_2O$ 

is obtained by evaporating the solution in vacuó, or by precipitation with alcohol. The salt crystallises with difficulty in very pale yellow plates, which are isomorphous with potassium ferrocyanide (Bunsen, Pogg. Ann. 1835, 36, 404). The aqueous solution decomposes when boiled, yielding ammonium cyanide and an insoluble residue, which possibly consists of more or less oxidised ammonium ferrous ferrocyanide.

 $2(NH_4)_4Fe(CN)_6-(NH_4)_2Fe[Fe(CN)_6]+6NH_4(NH_4)_2Fe[Fe(CN)_6]$ 

From solutions containing much ammonium chloride a double salt of the formula

crystallises very readily in well-formed orange-yellow rhombohedra, which were at one time mistaken for the ammonium salt (Bunsen, lc)

Barium ferrocyanide Ba, Fe(CN), 60H<sub>2</sub>O is precipitated when strong solutions of barium chloride and sodium ferrocyanide are mixed, it separates in the form of a light yellow, crystalline powder. Its solubility (grains of anhydrous salt per 100 grains water) is 0.34 at 15.5°, and 1.01 at 100° (Williams). The water of crystallisation is not quite completely driven off even at 160° (Drechsel, J. pr. Chem. 1880, [2]. 21, 77). The dried salt, when heated in vacuo, or in an atmosphere of introgen, carbon, iron, and barium cyanide and cyanamide being formed. When oxygen is completely excluded the product contains about 2 molecules of cyanamide to 1 of cyanide (Ewan and Napier, J. Soc. Chem. Ind. 1913, 32, 467).

## Barium potassium ferrocyanide

 $BaK_2[Fe(CN)_6],3H_2O$ 

is precipitated as a pale yellow, crystalline powder when solutions of barrum chloride and potassium ferroeyanide are mixed in equivalent quantities. Its solubility in water (grams anhydrous salt per 100 grams water) is 2.1 at  $15.5^{\circ}$ , 6.08 at  $100^{\circ}$ .

Calcium ferrocyanide (Ca<sub>2</sub>Fe(CN)<sub>6</sub>,12H<sub>2</sub>O<sub>15</sub> extremely soluble in water, highly concentrated solutions are very viscous and crystallise with difficulty, yielding small yellow trichine crystalls of sp gr 1 68, which effloresce when exposed to the air and lose 11½ molecules of water at 40° H<sub>2</sub> G Colman (Journ of Gas Lighting, 1910, 583) finds that the whole of the water is removed by heating the salt in a current of air. The dry salt begins to decompose (in absence of oxygen) at 500° yielding calcium cyanamide thus

Ca<sub>2</sub>Fe(CN)<sub>6</sub>=2CaCN<sub>2</sub>+Fe+N<sub>2</sub>+4C (unpublished experiments made by the writer).

Calcium ammonium ferrocyanide

Ca(NH<sub>4</sub>)<sub>2</sub>Fe(CN)<sub>6</sub>

is a white, minutely crystalline powder, which becomes coloured on exposure to light; it is anhydrous; 100 grams of water, at 15°-17°, dissolve 0.258 gram of the salt (J. Campbell Brown, Chem. Soc. Trans. 1907, 91, 1826)

Calcium potassium ferrocyanide  $K_2\text{CaFe}(\text{CN})_6$  is a white powder consisting of small quadratic prisms which becomes slightly cream-coloured on exposure to light. It contains no water of crystallisation and, at  $15^\circ-17^\circ$ , 100 grams of water dissolve 0.41 gram of it (J. Campbell Brown, *ibid* 1907, 91, 1826).

According to Williams (lc), the solubities

According to Williams (lc), the solubilities of these salts are rather larger, viz 0 35 at 15°, and 0 39 at 100° for the ammonium salt, and 0 72 at 15°, and 0 80 at 100°, for the potassium salt

Cupric ferrocyanide Cu<sub>2</sub>Fe(CN)<sub>6</sub> 11H<sub>2</sub>O is a red-brown, amorphous substance, obtained by adding a solution of calcium ferrocyanide or hydroferrocyanic acid to excess of cupric chloride solution In general, when solutions of a cupric salt and of a soluble ferrocyanide are mixed, an insoluble substance is produced, in which only a part of the four valencies of the ferrocyanogen ion is saturated by copper, the remainder being occupied by the metal of the ferrocyanide employed A very large number of these substances have been described Colloidal membranes formed of one or other of these copper ferrocyanides have been much used in the study of osmosis. The precipitate formed from potassium ferrocyanide and a cupne salt  $K_4$ Cu<sub>4</sub>[Fe(CN)<sub>6</sub>]<sub>3</sub> was used as a pigment under the name Hatchett's brown. The black, silver deposit in an ordinary photographic 'bromide' print may be converted into a red one, consisting of a copper ferrocyanide, by immersing the print in a solution containing 6 6 grams of copper sulphate crystals, 88 grams of potassium citiate, and 5 8 grams of potassium ferricyanide per litre (Ferguson, Photographic Journ 1900, 133)

Cuprous ferrocyanide Cu<sub>4</sub>Fe(CN)<sub>6</sub> xH<sub>2</sub>O is a white precipitate, which differs from most of the ferrocyanides in evolving the whole of its cyanogen as hydrocyanic acid when boiled with dilute sulphune acid. J. Dewrance and H. E. Williams (Eng. Pat. 28074, 1908) utilise this property in order to obtain complete decomposition of alkali ferrocyanides by distilling them with acid in presence of a small amount of cuprous chloride.

Hydroferrocyanic acid H4Fe(('N)6 is prepared by adding concentrated hydrochloric acid to a saturated solution of potassium ferrocyanide (both should be free from dissolved oxygen) and then enough ether to saturate the solution acid crystallises out in colourless scales which are washed with water saturated with other W Williamson, Annalen, 1846, 57, 225; A Joannis, Ann Chim Phys. 1882, [5] 26, 484), and dried over sulphuric acid. It may be puilfied by dissolving in alcohol and precipitating with ether, it obstinately retains ether, which is removed by heating at 100° in a current of dry hydrogen (Browning, Chem Soc Tians 1900, 1234) A solution of the acid, saturated at 14°, contains 150 grams per little The solution is readily oxidised by atmospheric oxygen, especially at higher temperatures, forming a blue substance which is usually assumed to be Prussian blue. This reaction is utilised in calico printing for the production of blue patterns

Ferrous ammonium ferrocyanide Fe(NH<sub>4</sub>)<sub>2</sub>[Fe(CN)<sub>6</sub>]

is formed when a solution containing ammonium ferrocyanide and excess of an ammonium salt is exactly precipitated by a ferrous salt, oxygen being excluded It is also formed when ammonium ferrocyanide is distilled with an acid in absence of air.

Ferrous cyanide  $Fe(CN)_2$  is formed when a solution of hydroferrocyanic acid is boiled, H<sub>4</sub>Fe(CN)<sub>6</sub>=4HCN+Fe(CN)<sub>2</sub>, or when the dry acid is heated at 300° in an atmosphere of hydrogen (Addie and Browning, Chem Soc Trans 1900, 158; Browning, ibid 1900, 1234)

Ferrous potassium ferrocyanide (Everitt's salt) K<sub>2</sub>Fe[Fe(CN)<sub>6</sub>] is formed in a similar way if potassium salts are present, as, for example, in the preparation of hydrocyanic acid by distilling a mixture of potassium ferrocyamide and dilute sulphuric acid Everitt's salt is a paleyellow crystalline powder which is not identical with the white precipitate obtained by mixing solutions of potassium ferrocyanide and ferrous chloride, although this precipitate has the same composition.

The iron in these ferious salts is displaced by treatment with caustic alkalis, but not by alkalı carbonates.

Ferric ferrocyanides. The blue compounds of iron and eyanogen are all amorphous substances, which yield colloidal solutions in pure water. It is therefore extremely difficult to prepare them in a pure condition Owing to the well-known tendency of colloids to adsorb other substances from solutions, the precipitates require very prolonged washing to free them from adhering salts, and during this process their composition may, and very often does, change, owing to atmospheric oxidation or to hydrolysis On the other hand, the method of determining the composition of a precipitate by mixing known quantities of the reagents and estimating the excess left in the supernatant liquid, is open, in this case, to the objection that the precipitate is not a pure substance because it has adsorbed some of the salts in the solution in contact with This method has been used by E Muller the This method has been used by E. Muller and by P. Woringer (E. Muller and T. Stanisch, J. pr. Chem. 1909, [n.] 79, 81; \*ibid. 1909, [n.] 80, 153, E. Muller, G. Wegelin, F. R. Treadwell, and O. Diefenthaler, \*ibid. 1911, [n.] 84, 353; E. Muller, \*ibid. 1914, [n.] 90, 119, P. Woringer, \*ibid. 1914, [n.] 87, 51.) K. A. Hofmann and his contractor (K. A. Hofmann, O. Hoppe, and F. collaborators (K A Hofmann, O Heine, and F Hochtlen, Annalen, 1904, 337, 1, K A Hofmann and F. Resensheck, *ibid* 1905, 340, 267, 1905, 342, 364; Hofmann, Arnoldi, and Hiendlmaier, ibid 1907, 352, 54), have investigated more especially the composition and properties of the substances which remain after exhaustive purification

The freshly prepared blue precipitates contain indefinite quantities of water, but the whole of it cannot be removed by drying in iacu6 over phosphorus pentoxide until a constant weight is attained (which usually requires from 2 to 5 This residual water must be regarded as constitutional, because when the substances are heated to temperatures a little above 100°, they lose hydrocyanic acid, but not water The formulæ of the dry substanecs given in what follows refer to the very thoroughly washed substances after complete drying over PaOs in the way mentioned above

When cold neutral solutions of one molecule of a ferric salt and one molecule of potassium ferrocyanide are mixed together, a precipitate of ordinary soluble Prussian blue is formed. The dry substance has the composition

 $2[KFe^{III}(Fe^{II}Cy_6)]+3\frac{1}{2}H_2O$ 

The same compound is obtained by mixing equal molecular quantities of a ferrous salt and potassium ferricyande. This was first proved by Skraup (Annalen, 186, 371, 1877).

When a cold solution of potassium ferrocyanide is added gradually to a cold neutral (or ½ p c. hydrochloric acid) solution containing an excess of a ferric or of a ferrous salt, or when soluble Prussian blue is digested with a solution of ferric chloride, precipitates are formed which, after the prolonged exposure to air involved in washing them completely, are identical in com-position and properties. The dry substance is position and properties normal ferric ferrocyanide

 $\text{Fe}_{4}^{\text{III}}(\text{Fe}^{\text{II}}\text{Cy}_{6})_{3} + 10\text{H}_{2}\text{O}$ 

usually referred to as Prussian blue. 'Turnbull's blue, obtained by precipitating a solution of potassium ferricyanide with an excess of a ferrous salt, is, after exhaustive washing with exposure to air, identical with ordinary Prussian blue

The composition of the unwashed precipitates initially formed depends, as Muller has shown, on the relative quantities of reagents employed. Setting out from potassium ferrocyanide and ferric chloride, Prussian blue is precipitated when the ratio mols K4Fe(Cv6)/mols. Fe(13 is less than 0.75, for values between 0.75 and 0.92 the precipitate contains increasing quantities of KFeIII[FeIICy,], and when the ratio exceeds 0.92 the substance

K. FeII FeII Cv.

is also present Setting out from potassium ferrieyanide and ferrous chloride a precipitate of the composition KFeIIFe<sub>5</sub>III[FeIICy<sub>6</sub>]<sub>5</sub> is obtained when the ratio, mols K<sub>3</sub>FeCy<sub>6</sub>/mols FeCl, is less than 0.714 Between 0.714 and 0 75 a substance KFe<sup>II</sup>Fe<sub>3</sub><sup>III</sup>[Fc<sup>II</sup>Cy<sub>6</sub>]<sub>3</sub> is formed, which Muller regards as ordinary Turnbull's blue; from 0.75 to 0.9 increasing quantities of KFeIII[FeIICy,] are present, and finally, when the ratio exceeds 0.9 ordinary Prussian blue appears According to Muller, Turnbull's blue and Prussian blue are interconvertible

 $\begin{array}{c} \operatorname{Fe_4^{III}}[\operatorname{Fe^ICy_6}]_3 + \operatorname{K'} + \operatorname{Fe''} \\ & \stackrel{\textstyle \textstyle \succeq}{\rightleftharpoons} \operatorname{KFe^IFe_3^{III}}[\operatorname{Fe^ICy_6}]_3 + \operatorname{Fe'''}, \end{array}$ 

although a very large excess of ferrous salt is required to convert an appreciable amount of Prussian blue.

It is obvious that the reactions involved in the formation of the blues are not simple double decompositions, mutual oxidations and reductions take place, and adsorption phenomena are superadded, so that, in general, the precipitates first produced are mixtures either of definite insoluble compounds, as Muller supposes, or if insoluble substances with soluble constituents adsorbed from the solutions.

The precipitates obtained from hot acid solutions differ more or less from those formed at ordinary temperatures in their properties, and appear even when washed to be mixtures of different substances.

Both the soluble and insoluble Prussian blue, above described, are ferric ferrocyanides, and not ferrous ferricyanides. This is proved by the fact that they can be made by the action of hydrogen peroxide on a solution containing a ferric salt and potassium ferricyanide in suitable quantities. As Schonbein (J pr. Chem 1866, 98, 65) showed, hydrogen peroxide reduces ferricyanide to ferrocyanide, but has no action on a ferric salt, so that ferrous ferricyanide could not be produced by this reaction

Blue compounds, which are not identical with those described above, are also obtained by the oxidation of ferrous ferrocyanides

(a) By adding a cold neutral solution of 1 molecule of a ferrous salt to a solution of 1 molecule of potassium ferrocyanide, and then oxidising the white precipitate by means of atmospheric oxygen, which is greedily absorbed (or better by hydrogen peroxide), a blue compound of the composition KFe<sup>III</sup>(Fe<sup>II</sup>('y<sub>6</sub>)+H<sub>2</sub>O is obtained. This differs from ordinary soluble blue in being insoluble in oxalic acid solution. It is referred to in the table of properties of the blues as K. A. Hofmann's blue.

(b) By proceeding exactly as in (a), except that the solutions contain 1 p c of sulphuric acid, a white precipitate is obtained, Fe<sup>II</sup>K<sub>2</sub>(Fe<sup>II</sup>Cy<sub>6</sub>), which is rather less readily oxidised to a blue of the same composition as before:

FeIIK(FeIICy,)+H,O

but possessing rather different properties. Like ordinary soluble Prussian blue, it is soluble in water and oxalic acid, but differs from it in being much more stable to ammonia solution. This compound is referred to as 'stable soluble blue'

(c) When a solution of potassium feirocyanide, containing about 10 pc of sulphuric acid, is heated in an atmosphere devoid of oxygen, one half of the hydrocyanic acid is evolved and a pale-yellow crystalline powder of Eventt's salt K<sub>2</sub>Fe<sup>II</sup>(Fe<sup>II</sup>Cy<sub>6</sub>) remains behind. This salt is very much less readily oxidised than the precipitate obtained by mixing solutions of potassium ferrocyanide and a ferrous salt, it is not completely oxidised by several weeks' exposure to the air. It is readily oxidised by warm dilute intric acid (A W Williamson, Ann Chem Pharm. 1846, 57, 225), but a homogeneous product is prepared most certainly by suspending it in 1 pc sulphuric acid and adding about 10 times the theoretical quantity of hydrogen peroxide. The substance thus obtained is Williamson's blue (sometimes called violet), and it also has the composition

KFeIII(Fe Cy6)+H2O

Its properties are, however, totally different from those of the isomenic substances already described

PROPERTIES OF THE BLUE IRON-CYANOGEN COMPOUNDS.

Reagent	Ordinary soluble Prussian bluc (Turnbull'ssoluble blue)	Insoluble Prussian blue	K A Hof- mann's blue	Stable soluble blue	Williamson's blue or violet
Pure water .	Soluble, reddish-blue solution	Insoluble	Soluble, greenish-blue solution	Soluble	Insoluble, turbid green- ish-blue sus-
10 pc oxalic	Soluble, blue solution	Soluble, blue solution	Insoluble	Soluble	pension Insoluble
4 p c ammonia yields red- brown feiric hydroxide		In 10 seconds	In a tew se- conds	In 6 to 7 mm- utes	After some hours
Ferric chloride	Precipitates insoluble Prussian blue.	No change	(۶)	Gradually co- agulated in unchanged condition	No change
1 to 1½ p c. H <sub>2</sub> SO <sub>4</sub>	Slowly precipitated, moist precipitate soluble in water	No action	Quickly pre- cipitated, insoluble in water	Precipitated, insoluble in water	No action
Appearance of dry solids		Deep-blue pur- ple lustre			Purple-red

For comparison, the distinguishing properties of the different blues are tabulated above. The isomerism of these substances may be due to differences in the linking of the atoms—of the kind familiar in organic chemistry—or it may be due to differences in the size of the molecules, the data available do not appear to be sufficient to decide these questions with any degree of certainty.

In accordance with the commonly accepted view, the blues have been formulated as ferro-

cyanides P Woringer, in 1914 (let), however, advanced certain arguments in favour of the ferricyanide structure. He confirmed an observation, made by A W Williamson in 1846, that when ferric chloride acts on an excess of ferrocyanide the solution contains ferricyanide; he also states that Prussian blue, when decomposed by ammonium carbonate, yields 90 p.c. of ferricyanide. As E. Muller pointed out in 1909, it is easy to show, from the well-known oxidation potentials of solutions of ferrous

and ferric ions and of ferro- and ferri-cyanogen ions that, if these four ions could coexist in solution the ratio of their concentrations [Fe] [FeCy<sub>6</sub>"'] [Fe][Fe "Cy<sub>6</sub>""] would be about 10<sup>6</sup>, that is, ferrocyanide would be exidised

practically completely by ferric salts

Williamson, however, showed that the reverse reaction also takes place, ferricyanide being partly reduced by ferrous salts. It appears thus that ferri- and ferro-evanogen ions can coexist in approximately equal concentration, so that, in equilibrium, the concentration of ferric ion must be very small compared with that of ferrous ion These concentrations are determined by the solubilities of the precipitates formed, about which very little appears to be known; the mode of formation of a blue affords, therefore, no certain clue to its structure The same is true of its decomposition by strong alkalis, since alkaline ferricyanide solution oxidises ferrous hydroxide very readily mann's methods of preparation by means of hydrogen peroxide appear at present to be the best evidence available in favour of the ferrocvanide structure

It is of interest that precipitation from hot acid solutions always tends to produce substances (or mixtures) of a more stable type, approaching Williamson's blue more or less closely in properties, and also that all the blue iron cyanogen compounds contain both ferrous and feiric iion The compounds which contain only ferrous or only ferric iron are colourless or (compared with the blues) feebly coloured Solid pieces of ordinary Prussian blue assume a coppery lustre when rubbed on a smooth, hard surface The blue solution in oxalic acid was formerly used as an ink Prussian blue is reduced by hydrosulphurous acid, H2S2O4, or its sodium salt to white ferrous ferrocyanide (M. Rohn, Zeit Anorg Chem 1906, 49, 443) turns a suspension of Prussian blue in water green; the solution contains ferric chloride . the blue colour is restored by washing with water Concentrated sulphuric acid unites with Prussian blue to a white paste from which the original blue is obtained by diluting with water It is slowly decomposed by concentrated aqueous hydrochloric acid, but dissolves to a colourless solution in alcoholic hydrochloric acid, the solubility increases with the molecular weight of the alcohol used Water precipitates unchanged Prussian blue from these solutions By dissolving impure Prussian blue in a solution of hydrochloric acid in propyl alcohol, filtering, and reprecipitating with water, it may be separated from all commonly occurring impurities, such as barium sulphate, kaolin, alumina, and starch (C. Coffignier, Bull. Soc chim. 1904, 391) Potassium, sodium, and calcium hydroxides decompose Prussian blue into ferric hydroxide and a ferrocyanide, alkalı carbonates act more slowly dyed with Prussian blue gradually lose their colour in sunlight, but regain it in the dark When heated to about 170° in the air, it burns like tinder leaving a brown ash.

Colloidal solutions of Prussian blue (0.2 to 2 p.c) may be prepared by dialysis of the oxalic acid solution. The colloidal blue is negatively charged and is coagulated by cathions; hydrogen is especially active, otherwise the usual

increase of activity with valency obtains (W. Pappadà, Gazz. chim. ital. 1911, 41, ii. 454). K Å. Hofmann (Annalen, 1905, 340, 267) also observed that his soluble blue carried a negative charge, and did not depress the freezing-point

Commercial blues are usually mixtures of the compounds already described. J. G Gentele (Lehrbuch der Farbenfabrikation, 2nd ed. 1880) says that the best blues are obtained by precipitating a feirous salt with potassium ferrocyanide, and then oxidising the piecipitate Paris blue (the finest quality of commercial Prussian blue) is made as follows. 50 kilos of potassium ferrocyanide are dissolved in 250 kilos of boiling water, 42 5 to 45 kilos of green vitriol are dissolved in about the same quantity of water, preferably in the presence of scrap iron to prevent formation of ferric salt. The clear solutions are then run simultaneously into 250 kilos of water The almost white precipitate is allowed to settle and then drained on a cloth filter. The oldest method of 'blueing' the paste was to heat it to boiling in a basin, transfer it to a wooden tub, and add 25 5 kilos. of mtric acid (sp gr 1 23) and 18 kilos. of sulphuric acid (1 84 sp gr). Sometimes, red fumes were evolved only after some hours, but in presence of excess of green vitriol, they were always evolved at once and this gave the best colour After standing for 24 hours, the mixture was suspended in a large quantity of cold water and allowed to settle. The washing by decantation was continued until the blue was free from sulphuric acid. It was then collected on linen filters, pressed into thin cakes, and cut up into cubes which were dried in the an and finally at 30°-40°. The yield was 39 to 39 5 kilos of finished blue.

Another method was to treat the heated paste of ferrous ferrocyanides with ferric chloride, instead of with intire acid, until the solution contained ferric in addition to ferrous salts, but according to Gentele, the cheapest and best method is to acidify the paste slightly with hydrochloric acid and pass chlorine gas into it until the solution shows a distinct 'ferric' reaction with ferrocyanide.

From these descriptions it is clear that commercial Paris blue must contain, in addition to ordinary insoluble Prussian blue, one or more of the potassium ferric ferrocyanides presence of potassium appears to be of importance, a less satisfactory colour being produced when sodium ferrocyanide is used in its pre-paration H E Williams says that potas-sium may be replaced by ammonium with satisfactory results A Eibner and L Gerstacker (J Soc. Chem Ind 1912, 31, 1041) state that the blues with a high potassium content possess the technically desirable greenish shade, those poor in potassium being dull with a violet tinge. They have tound 12 p c K and 5 5 p c. H<sub>2</sub>O in 'steel-blue' and 'bronze blue,' 9 p c K and 105 pc H2O in 'Paris blue,' the lowest value found in a commercial blue being 15 p c K and 24 pc H<sub>2</sub>O. In the commercial pigments, Paris blue is diluted with starch, heavy spar, gypsum, zinc white, or burned and finely ground kaolin. The white admixture, after having been ground very finely by itself, is added to the Paris blue paste, and along with it passed through a colour mill. Very low qualities of

blue are faced by rotating the dry cubes in a cask with the fine dust of pure Paris blue.

Potassium ferrocyanide (yellow prussiate of potash, Blutlaugensalz, Ger) crystallises from aqueous solutions in large, well-formed, orangeyellow tetragonal pyramids of the composition  $K_4$ Fe(CN)<sub>6</sub>3 $H_2$ O; sp gr 186. The crystals do not change when exposed to the air at ordinary They begin to lose water at 60° temperatures and the finely powdered salt may be completely dehydrated at 100°, forming a white powder which begins to decompose, in absence of oxygen, slightly below a red heat to a mixture of potassium cyanide, carbon and iron, nitrogen escaping. The powder takes fire at a considerably lower temperature when heated in contact with air It is insoluble in alcohol The soluble ferrocyanides are completely decomposed by boiling with mercuric oxide

$$K_4$$
Fe(CN)<sub>6</sub>+3HgO+2H<sub>2</sub>O  
=3Hg(CN)<sub>2</sub>+FeO+4KOH

They are also decomposed by aurous evanide. hydroxide, and even by finely-divided gold in the presence of oxygen

$$\begin{array}{l} 3\mathrm{Au} + \mathrm{K_4Fe(CN)_6} + 2\mathrm{H_2O} + \mathrm{O_2} \\ &\quad = 3\mathrm{KAu(CN)_2} + \mathrm{Fe(OH)_3} + \mathrm{KOH} \end{array}$$

(E. Beutel, Zeitsch anorg Chem 1912, 78, 141) According to Etard (Ann Chim Phys 1894, [7] 2, 546), the solubility, expressed in grains of anhydrous salt dissolved by 100 grams of water, is 14 5 at 0°, 24 5 at 20°, 36 at 40°, 49 5 at 60°, 64 at 75°, 70 at 80°, 72 at 89°, 74 at 98°, and 88 at 157°. The salt is much less soluble in solutions of potassium salts The specific gravity of the aqueous solutions is given below (Schiff, Annalen, 1860, 113, 199) -

G1s  $K_4$ Fe(CN)<sub>6</sub>,3H<sub>2</sub>O  $\rbrace$  1 per 100 grs. of solution  $\rbrace$  1 Sp gr at 15° . . . 1 005 5 15 . 1 0058 1 0295 1 0605 1 0932 1 1275

Sodium ferrocyanide crystallises in large lemon - yellow monoclinic prisms or rhombohedra of the composition  $Na_4Fe((!N)_0, 10H_2O)$ (Conroy, J Soc Chem Ind 1898, 17, 103), which are permanent when exposed to the air at the ordinary temperature, but are completely dehydrated at a little above 100° The behaviour of the dry salt, when heated, resembles that of the potassium compound solubility in water (Conroy, J Soc Chem Ind 1898, 17, 103). grams anhydrous salt dissolved by 100 grams of water, is at 20°, 17 9; at 30°, 23 5, at 40°, 29, at 50°, 35 5, at 60°, 42 5, at 70°, 51 5; at 80°, 59 2; at 90°, 61, and at 100°, 63

Zinc ferrocyanide Zn<sub>2</sub>Fe(CN)<sub>6</sub>,6H<sub>2</sub>O is a white amorphous substance obtained by the action of calcium or hydrogen ferrocyanide on an excess of zinc chloride in aqueous solution With other soluble ferrocyanides the precipitate contains more or less of the metal of the ferrocyanide used, a fact referred to in more detail in the analytical section of this article. It is insoluble in dilute acids, completely soluble in caustic alkalis to a mixture of ferrocyanide and zincate, and only partially decomposed by alkali carbonates.

Potassium ferricyanide, red prussiate of potash, K<sub>3</sub>Fe(CN<sub>6</sub>), was discovered by Leopold Gmelin in 1822. He prepared it by passing chlorine into a solution of potassium ferro-VOL. II.-T.

eyanide until it no longer gave a blue colour with a ferric salt:

449

$$2K_{4}Fe(CN)_{6}+Cl_{2}=2K_{3}Fe(CN)_{6}+2KCl$$

The salt is always manufactured by this reaction; the ferrocyanide is sometimes used in the form of a dry powder, in which case a mixture containing potassium chloride is obtained Most commonly, a cold 10 pc. solution of potassium ferrocyanide is treated with chlorine until conversion is just complete. A little Prussian green is always formed by the action of chlorine on the ferricyanide, and as this passes through any filter it spoils the appearance of the product. The difficulty may be avoided by evaporating the solution to crystallisation and then adding a little caustic potash which decomposes the green into ferric hydroxide, which is easily removed by filtration, and potassium ferrocyanide, which is apt to crystallise out along with the ferricyanide The yield of ferricyanide, in this process, is 85 to 90 pc of the theoretical quantity

Λ W Williamson (Annalen, 1846, 57, 225) found that potassium ferric ferrocyanide (Wilhamson's blue), when boiled with potassium ferrocyanide, gives potassium ferricyanide and Everitt's salt

$$\begin{split} K_4 \mathrm{Fe}^{\mathrm{II}}(\mathrm{CN})_6 + K \mathrm{Fe}^{\mathrm{III}}[\mathrm{Fe}^{\mathrm{II}}(\mathrm{CN})_6] \\ = & K_3 \mathrm{Fe}^{\mathrm{III}}(\mathrm{CN})_6 + K_2 \mathrm{Fe}^{\mathrm{II}}[\mathrm{Fe}^{\mathrm{II}}(\mathrm{CN})_6] \end{split}$$

Since Everitt's salt is readily oxidised to Williamson's blue by warming with dilute nitric acid, the process might (as Dittmar suggested) be utilised to complete the conversion of ferrointo ferricyanide, chlorine being used up to the point at which Prussian green begins to be

Many other oxidising agents have been suggested in place of chlorine, such as bromine, lead peroxide, calcium plumbate, potassium permanganate and persulphate, and electrolytic oxidation (see Bertelsmann, Technologie der Cyanverbindungen)

Potassium ferricyanide crystallises from water in magnificent, deep - red, monoclinic prisms which may be very large. It is anhydrous. sp gr 1845. The aqueous solution has an intense yellow colour. The solubility (grams) of anhydrous salt dissolved by 100 grams of water) is 33 at 44°, 36 at 10°, 40 8 at 155°, 58 8 at 37 8°, 77 5 at 100°, and 82 6 at 104 4° (Wallace, Dingl. poly J 1856, 142, 52) Specific gravity of the solutions at 15° (Schiff, Annalen, 1860, 113, 199)

The aqueous solution is decomposed by light, potassium ferrocyanide being formed. caustic alkaline solution is a strong oxidising agent which decolourises indigo instantaneously The alkaline solution is also reduced by hydrogen

peroxide, oxygen being evolved

Ferric ferricyanide. The dark-brown solution, formed when solutions of a ferricyanide and of a ferric salt are mixed, contains ferric ferricyanide in the form of a highly disperse Ultramicroscopic examination show that the particles gradually increase in size, a green substance ultimately separating out

(R. Haller, Kolloid Zeit. 1917, 20, 76). The formation of a green substance by the action of chlorine on potassium ferricyanide was first noticed by Pelouze (Ann Chim Phys. 1838, [2] 69, 40). Green ferric ferricyanides are also obtained by boiling Prussian blue with strong nitric acid or by precipitating potassium ferricyanide containing a little ferrocyanide with ferric chloride (H E Williams)

# Potassium carbonylferrocyanide $K_3[Fe(CN)_5CO]H_2O$

is formed by the action of carbon monoxide on an aqueous solution of potassium ferrocyanide at 135°

 $K_4$ Fe(CN)<sub>6</sub>+CO+2H<sub>2</sub>O = $K_3$ Fe(CN)<sub>5</sub>CO+NH<sub>3</sub>+KHCO<sub>2</sub>

(Muller, Ann. Chim Phys 1889, [6] 17, 94) Carbonylferrocyanides are found in the cyanide mud and spent oxide of the gas works, in which from 2 to 5 p c of the cyanogen (exceptionally as much as 20 pc) is present in this form (H G. Colman, Analyst, 1908, 33, 261) When the caustic soda extract of these materials, mixed with 4 to 5 times its volume of methylated spirit, is allowed to stand for some hours, the sodium ferrocyanide crystallises out completely, leaving the carbonyl ferrocyanide in solution It is isolated, in the form of the amorphous, deep-violet ferrie salt, by adding ferric chloride to the acidified solution to the acidified solution. The deliquescent calcium salt  $\text{Ca}_3[\text{Fe}(\text{CN})_5(\text{O})]_2,8\text{H}_2\text{O}$  is obtained by boiling the ferric salt with milk of lime The carbonyl ferrocyanides resemble the corresponding ferrocyanides, but are paler in colour and more soluble in water The copper salt, a green precipitate soluble in excess of ammonia, is characteristic

Sodium nitroprusside  $Na_2[Fe(CN)_5NO]$   $2H_2O$  was first prepared by Lyon Playfair (Phil Mag 1850, [3] 36, 197) by the action of dilute intraction on a solution of a ferrocyanide. The primary products of the change, hydroferricyanic acid and nitric oxide, react thus  $H_3Fe(CN)_6+NO=H_2Fe(CN)_5NO+HCN$ . The salt is also formed by adding a concentrated solution of ferrous sulphate to a solution containing equal parts of sodium nitrite and potassium cyanide. It crystallises in dark-red prisms, soluble in  $2\frac{1}{2}$  parts of water at  $16^\circ$ . The solution is not precipitated by a ferric salt, and gives a deep violet colouration with alkali sulphides

#### Sodium aquopentacyanoferroate Na<sub>3</sub>[Fe(CN)<sub>5</sub>H<sub>2</sub>O],5H<sub>2</sub>O

is obtained by the action of reducing agents, such as phenylhydrazine or hydroxylamine, or of oxidising agents, such as potassium hypobromite and hydrogen peroxide on a solution of sodium nitroprusside at temperatures below 0° The action of hydrogen peroxide is analogous to its behaviour with a ferricyanide

 $2K_3$ Fe(CN)<sub>6</sub>+ $H_2O_2$ +2KOH

 $=2K_4Fe(CN)_6+O_2+2H_2O$ 

The salt crystallises in reddish-yellow needles, giving an intensely yellow solution, which does not yield a violet colouration with alkali sulphides.

## Sodium aquopentacyanoferriate

 $Na_2[Fe(CN)_5H_2O]$ 

is obtained by oxidising the foregoing compound with bromine. It is a deep violet-blue substance,

and Cambi has shown that the potassium perferricyanide obtained by Bong (Bull. Soc. chim 1875, [2] 24, 268) and Skraup (Annalen, 1877, 189, 368) by the action of potassium chlorate and an acid on a solution of potassium ferricyanide, is the potassium salt of the same acid. It is also obtained by the action of excess chlorine on potassium ferricyanide, a deep violet solution being formed, from which the salt is precipitated by alcohol.

A series of pentacyano iron compounds has also been prepared, which may be regarded as derived from the aquo-compounds just described by the replacement of the H<sub>2</sub>O group in the anion by one of the groups NO<sub>2</sub>, AsO<sub>2</sub>, SO<sub>3</sub>, NH<sub>3</sub>, or substituted ammonias.

Literature.—K A. Hofmann, Zeitsch anorg Chem 1896, 11, 31, and 278; *ibid.* 1896, 12, 146; Annalen, 1900, 312, 1; W Manchot, Ber 1912, 45, 2869; L Cambi, Gazz chim ital. 1911, 41, i. 157.

Hexamethyl carbylamine ferro saits By the action of methyl sulphate on dry potassium ferrocyanide or of methyl iodide on silver ferrocyanide, E G J Hartley has obtained a series of salts of the general formula [Fe(CNCH<sub>3</sub>)<sub>6</sub>]R<sub>2</sub>, R being a univalent acid radical, in which the iron cyanogen complex plays the part of cathion. When heated with concentrated sulphuric acid, or boiled with aqueous caustic soda, these salts are decomposed, the whole of the nitrogen appearing in the form of methylamine, or of methyl carbylamine, which proves that all the nitrogen atoms are linked to methyl

Tetramethyl ferrocyanide  $\left[ \operatorname{Fe}^{(\operatorname{CNCH}_3)_4}_{(\operatorname{CN})_2} \right]$  is formed when hexamethylcarbylamine ferrochloride [Fe(CNCH<sub>3</sub>)<sub>6</sub>]Cl<sub>2</sub> is heated at 140°-150° in a vacuum for 6-9 hours, methyl chloride being Two isomeric forms, with the same split off molecular weight, have been separated from The a-isomer is much the reaction product more readily soluble in water, alcohol, and chloroform than is the  $\beta$ -isomer, and from water it crystallises with two molecules of water of crystallisation, the β-isomer combining with six They also differ in their behaviour towards methyl iodide, with which the α-compound combines at  $100^{\circ}$ , yielding hexamethylcarbylamine ferroiodide, the  $\beta$ -compound remaining un-No method of converting the one changed form into the other has been discovered (E. G. J. Hartley, Chem Soc Trans 1910, 1066 and 1725. ibid 1911, 1549, ibid 1912, 705; ibid 1913,

Constitution of the Iron Cyanogen Compounds.—When dissolved in water these compounds dissociate electrolytically into a complex ion containing the iron, cyanogen, and other groups, the sign and magnitude of the electric charge of this ion depend on the nature of its constituent groups of atoms. The ions Fe(CN)<sub>6</sub>"', Fe(CN)<sub>6</sub>"'', Fe(CN)<sub>6</sub>"'', Fe(CN)<sub>6</sub>"'', Fe(CN)<sub>6</sub>"'', Fe(CN)<sub>6</sub>"'' are examples of this These ions possess considerable stability, but are probably dissociated to a very small extent For example, the ferrocyanogen ion is supposed to dissociate thus, Fe(CN)<sub>6</sub>""'\sqrt{Fe}"+6(CN)' (F. Haber, Zeit. Elektrochem. 1905, 11, 846; Foster, Chem. Soc. Trans. 1906, 912). In the dark the concentration of ferrous ion is too small to produce

a precipitate of FeS with an alkalı sulphide, but the complete decomposition of the ferrocyanogen complex by mercuric oxide shows that the minute concentration of (CN)' in a solution of a ferrocyanide is greater than that in a solution of mercuric cyanide. The dissociation of the ferrocyanogen ion is increased by light, especially the ultra-violet rays, so that in alkaline solutions, oxygen produces a precipitate of ferric hydroxide. The increased dissociation is, however, so small that no appreciable increase in the conductivity of the solution can be detected

Measurements of the osmotic pressure and electrical conductivity of solutions of the ferrocyanides of calcium and strontium have been made by the Earl of Berkeley, Hartley and Stephenson (Phil Trans. 1909, 209A, 319), the results of which are incompatible with the assumption of a molecule Ca. [Fe(CN)<sub>6</sub>] dissociating into three ions, it is suggested that the undissociated molecules may be double. The non-electrolyte, tetramethyl ferrocyanide, exists as single molecules in aqueous solution, and, from measurements of the freezing-point, Buchbock (Zeitsch. phys. Chem. 1897, 23, 157) arrived at the same result in the case of tetraethyl ferrocyanide.

Several cases of isomerism have been recorded Locke and Edwards (Amer Chem. J 1899, 21, 193 and 413) observed that the ordinary anhydrous, red potassium ferneyanide is converted into a greenish-yellow modification, crystallising with 1 molecule of water by treatment with acids This \$\beta\cdot \text{ferricyanide gives no}\$ precipitate with bismuth nitrate, the ordinary α-modification giving a straw-coloured precipitate. The \beta-ferricyanide is converted into the a-salt by treatment with alkalis Briggs (Chem. Soc Trans 1911, 1019) describes similar behaviour on the part of the ferrocyanides. Faintly acid solutions deposit more deeply coloured crystals of the B-isomer, which are readily converted into the a-form by treatment with potassium cyanide or alkalis. The lmenthyl ammonium salts of the two isomers have different specific rotatory powers in alcoholic solution, otherwise only minute differences in density and solubility were observed Bennett (Chem. Soc. Trans. 1917, 490) finds that the  $\alpha$ - and  $\beta$ -ferrocyamides are crystalographically identical, and explains the difference of colour by slight decomposition of the acid solutions


In addition to the isomeric tetramethyl ferrocyanides already mentioned, Hartley (Chem Soc Trans 1914, 521) has recorded the existence of two isomeric trimethyl cobalitivamides  $Co(CNCH_1)_3(CN)_3$  (attempts to prepare the ferricyanide derivative were unsuccessful), which differ considerably in solubility.

The synthesis of ferrocyanides from a feirous compound and of ferricyanides by boiling excess of ferric hydroxide with an alkali cyanide (Skraup, Annalen, 1877, 189, 376), show that these compounds and their direct derivatives contain divalent and trivalent iron respectively It is to be remembered that the iron cyanogen compounds here considered form one group of a great family of complex salts. Substances of the general formulæ

are known in which the metal is Co, Cr, Mn.

Ru, Rh, Ir, and Os, and in which the CN group may be replaced by SCN These are closely related in structure and chemical behaviour to large groups of metal ammonia and halogen complex salts Under the circumstances, Werner's formulation of these salts, although it lacks the definiteness of the structural formulæ of organic chemistry, appears to be the best available A Werner (Neuere Anschauungen auf den Gebiete der anorganischen Chemie, 3rd ed 1913) regards the complex salts as built up by the co-ordination of a maximum number of six groups of atoms round a central atom; carbon, nitrogen, and boron are exceptional, their maximum co-ordination number being four. The six groups are arranged symmetrically; at the angles of a regular octahedron, they are attached to the central atom, partly by the ordinary principal valencies, partly by secondary The nucleus so formed acts as a more or less stable whole, which may still be able to combine with other atoms at a greater distance from the centre or on an outer sphere. These more loosely held atoms dissociate electrolytically when the substance is dissolved; whether they are attached to the nucleus as a whole or to any particular atom in it is left undecided, although Werner inclines to the The number and nature of the former view ionogenic groups depends on the nature of all the atoms included in the nucleus For example, among the compounds described, we have

A further consequence of the theory is that complexes of the formulæ  $\left[ \begin{array}{c} M_{R'_2}^A \end{array} \right]$  and  $\left[ \begin{array}{c} M_{R'_3}^A \end{array} \right]$  should exist in the two stereoisomeric forms represented by '



Possibly Hartley's isomeric tetramethyl ferrocyamde and trimethylcobalticyamdes are examples of these forms of isomerism. Werner classifies the (somewhat uncertain) isomerism of the ferro- and ferricyamides as unexplained, but as Briggs has remarked (Chem. Soc. Trans. 1908, 1564), they would also come under the above heading if the potassium atoms are supposed to be attached to definite CN groups in the nucleus.

Manufacture of Potassium and Sodium Cyanides.—The methods of manufacture of these salts are most conveniently described together, since most of the processes used may equally well

be applied to making either salt. The most important technical processes have already been mentioned in the introduction to this article

Manufacture from Yellow Prussiate —Potassium cyanide was first prepared by Berzelius's method, which consists in heating dry potassium ferrocyanide to redness, when it decomposes thus:  $K_4 \text{Fe}(\text{CN})_6 = 4 \text{KCN} + \text{Fe} + 2 \text{C} + \text{N}_2$ . The loss of cyanogen, which occurs in this way, is partly avoided by fusing the ferrocyanide with potassium carbonate, as first described by F and E Rodgers (Phil Mag 1834, [3] 4, 91) Liebig showed that the product contains cyanate, formed thus:

 $K_4$ Fe(CN)<sub>6</sub>+ $K_3$ CO<sub>3</sub>=5KCN+KCNO+Fe+CO<sub>2</sub>

The fused salt is allowed to stand until the iron and carbon have settled out, and is then poured off into moulds. The comparatively small quantities of cyanide, which were used in electroplating and for other purposes up to about 1890, were made by this process. The product was sold in white, porcelain-like cakes, containing from 30 to 70 p.c. of KCN.

In 1876, E Erlenmeyer (Ber 1876, 9, 1840) pointed out that a very pure product, containing on the average 40 p c. of CN (equivalent to 100 p c KCN), was obtained by fusing dry prussiate with metallic sodium in the proportions required by the equation

$$K_4$$
Fe(CN)<sub>6</sub>  $+2$ Na=4KCN+2NaCN+Fe

This process was taken up first by the Deutsche Gold and Silber Scheide-Anstalt (F. Roessler, V. Inter Kongress für Angewandte Chemie, Berlin, 1903, 1, 638) in Frankfort-on-Main, in 1890, they obtained a practically quantitative yield by filtering off the spongy iron from the fused evanide and pressing the iron while red hot. The eyanide so obtained is east in cakes and forms a pure white crystalline mass. Since about 1905, sodium ferrocyanide has been used in this process in place of the potassium salt, and the product is almost pure sodium evanide. The excess of the gas works ferrocyanide over that required in the colour industries is converted into sodium evanide in this way.

It has been proposed to distil ferrocyanides with acids and to absorb the hydrocvanic acid in solutions of caustic alkalis. Grossmann (Eng. Pats 36 and 4513, of 1903, J Soc Chem. Ind 1903, 22, 1327) distils sodium ferrocyanide with 20 pc sulphuric acid in lead stills the Everitt's salt remaining is reconverted into sodium ferrocyanide by boiling it with caustic soda in a current of air The use of calcium ferrocyanide, which readily yields two-thirds of its hydrocyanic acid, was proposed by J. H. Paul (Eng. Pat. 72 of 1903). W. Feld (Eng. Pat. 24920 of 1901) converts insoluble ferrocyanides into mercuric cyanide, and then distils Dewrance and Williams this with an acid (Eng. Pat 28074 of 1908) distil the ferrocyanide, either soluble or insoluble, with 2-3 p.c. of its weight of cuprous chloride and a small excess of sulphuric acid, the whole of the hydrocyanic acid being evolved. None of these proposals is at present of technical importance.

The processes of Stepermann and Beilby were the first synthetic processes to produce cyanide in large quantities, they are most conveniently considered together because, although they differ

widely in the methods adopted in carrying them out, they are both based on the same chemical reaction which was also used by Scheele in the first synthesis of a cyanide ever made

In Siepermann's process (D. R. P 38012, March 10, 1886; Eng Pat 13697 of 1889; 9350 and 9351 of 1900) a dry mixture of about 2 parts of charcoal and 1 part of potassium carbonatethat is, a sufficient excess of charcoal to keep the mixture in the form of a dry unfused mass throughout the process—is heated in the upper part of a vertical iron tube to a dark-red heat and treated with dry ammonia gas, which is taken up, forming potassium cyanate. The product is then allowed to fall into the lower end of the tube, which is heated to bright red-At the higher temperature the cyanate is converted into cyanide, the carbon monoxide evolved escaping through the upper part of the tube along with the hydrogen formed from the ammonia, without taking any part in the reaction. The finished material is drawn off into air-tight receptacles where it is allowed to cool completely, after which it is systematically liviviated with water Potassium cyanide may be precipitated from the solution obtained (sp gr about 14) by adding excess of potassium carbonate, but according to the later patents it is preferable to evaporate it almost to dryness and extract the unchanged potassium carbonate with small quantities of The residue of potassium cyanide water at 66° and cyanato is then treated with water between -18° and +5°, which dissolves the cyanide and any residue of carbonate, leaving almost pure potassium cyanate The product of this process was first put on the market in 1892 (G. Beilby, potassium cyanate V Int Kongress angew Chem Berlin, 1903, 1, 630), but as the substance was in the form of a damp deliquescent mass, it could not be exported for use at the mines by itself but had to be fused with the product of the ferrocyanide process

In the Bellby process (Eng Pat 4820 of 1891), which began to produce in the same year, the quantity of charcoal used is very much smaller, very little more being employed than is necessary to convert the potassium carbonate into eyanide thus

$$K_2CO_3+4C+2NH_3=2KCN+3CO+3H_9$$

The charcoal is added gradually during the progress of the operation, so that the material is always in the form of a fused liquid through which the ammonia gas is forced under some This is continued until a sample pressure shows the conversion to be sufficiently complete: it is then only necessary to separate the small excess of charcoal by filtration in order to have a pure white saleable product Since the melting-point of pure potassium carbonate (890°) is inconveniently high, previously made cyanide was mixed with it in starting the process, or the more readily fusible mixture of sodium and potassium carbonates was used, so that the cyanide made by the process was very similar in composition to that produced by the Erlenmeyer process from potassium ferrocyanide. In 1899 the Beilby process contributed fully one-half of the total European production of cyanide

It is difficult to obtain a satisfactory yield on the ammonia in these processes, owing to the rapidity with which it decomposes into its

elements at temperatures above 500° and in contact with metals, especially iron (Ramsay and Young, Chem. Soc. Trans. 1884, 88; Beilby and Henderson, *ibid.* 1901, 1245). For this reason it is necessary to work at the lowest possible temperature. Siepermann, for example, obtained a yield of 86 p c on the ammonia when it was absorbed at a dark-red heat, but only 58 pc at a bright cherry-red heat Readman (J Soc. Chem Ind 8, 757, 1889) showed that almost theoretical efficiencies can be obtained at the low temperatures attainable in an ordinary organic combustion furnace, whilst ('onroy (ibid 15, 12, 1896), working at very high temperatures (1000°-1200°), could not get more than 30 to 33 pc of the ammonia in the form of cyanide. In practice the reaction velocity is cyanide also of great importance, in the Beilby process where the advantage of the large surface exposed by Siepermann's dry mass of alkalised charcoal is sacrificed, the reaction is too slow at tempera tures below 900°, the best results are obtained at  $980^{\circ}$ . In order to keep down the dissociation of the ammonia at this high temperature it is injected into the fused mixture through copperlined tubes of such diameter that it's time of contact with the hot metal is less than one-tenth of a second

The chemistry of the reaction has been investigated by J. Pfleger (F. Roessler, V. Int. Kongress angew. Chem. Berlin, 1903, 1, 638). Ammonia and potassium carbonate react, yielding potassium eyanate and hydroxide, a carbamate being most probably formed as an intermediate product.

$$\begin{array}{c} \text{CO(OK)}_2 + \text{NII}_3 = \text{CO} \\ \text{OK}^2 + \text{KOH} \\ \text{CO} \\ \text{OK}^{-1} = \text{KCNO} + \text{H}_2\text{O} \end{array}$$

When pure potassium cyanate is heated at 800°-900°, it decomposes, cyanide and carbonate being formed and introgen, carbon dioxide, and carbon monoxide evolved. The mechanism of this change is probably as follows. Drechsel (J. pr. Chem. 1877, [2] 16, 201) has observed that the cyanates of the alkaline earth metals decompose, when heated, into carbon dioxide and the corresponding cyanamide salts. Assuming that the alkali cyanates behave similarly, the first change will be

$$2KCNO = K_2CN_2 + CO_2$$

Potassium cyanamide, however, decomposes completely at lower temperatures than  $800^{\circ}$  into cyanide, potassium, and nitrogen .

$$K_2CN_2 = KCN + K + N$$

The carbonate and carbon monoxide observed are thus due to the reaction between the metallic potassium and carbon dioxide which are formed simultaneously. In presence of chatcoal in excess, the greater part of the cyanamide follows Drechsel's reaction (J pr Chem 1880,[2] 21, 77),  $\rm K_2CN_2 + C = 2KCN$ , instead of decomposing; so that the apparent induction of cyanate by carbon probably takes place in the two stages:

$$\begin{array}{c} 2KCNO\!=\!K_2CN_2\!+\!CO_2\\ \text{and } K_2CN_2\!+\!C\!=\!2KCN\\ CO_2\!+\!C\!=\!2CO \end{array}$$

The reactions occurring in the Beilby process are the same, except that, the whole process

being carried on at the higher temperature, the formation of cyanate and its conversion into cyanide take place simultaneously instead of being more or less sharply separated as they are in Siepermann's process

In the Siepermann process, it is easier to use the ammonia efficiently, owing to the lower temperature employed, but this is counterbalanced by the greater difficulty of working up the product into a marketable form.

The Castner Process—In 1894 H Y Castner (Eng Pat 12219 of 1894) patented a process in which sodium was fed in at the upper end of a column of red-hot charcoal through which a current of dry ammonia gas was passed from below, the reaction being expressed by the simple equation.

$$2Na + 2C + 2NH_{3} = 2NaCN + 3H_{2}$$

In a second patent (H Y Castner, Eng Pat 21732, 1894) it is shown that better results are obtained by first allowing the sodium and ammonia to react at a low temperature (300° to 400°), forming sodium amide NaNH<sub>2</sub>, and then running the amide on to an excess of red-hot charcoal with which it reacts thus:

Working in this way, it happened on one occasion that an excess of amide was accidentally run into the vessel containing the charcoal, and a product of unusual appearance was obtained, in which the writer found a large quantity of sodium cyanamide. The reaction, therefore, takes place in three stages, in which the introgenenters successively into more and more stable combinations. First sodium amide is produced, a compound which begins to decompose rapidly into its elements at 500° (Titherley, Chem. Soc. Trans. 1894, 504), this reacts either with carbon.

$$2NaNH2+C=Na2CN2+2H2$$

or, as Pfleger found, with sodium cyanide

forming sodium cyanamide, a compound which only decomposes slowly at 600°, and finally this takes up carbon, forming sodium cyanide which is quite stable at any temperature attained in the process

Like the Bellby process, this process yields a fused product containing very small quantities of solid impurities which are easily removed by filtration. After being cast in non moulds, the cyanide is obtained in the form of pure white cakes with crystalline fracture, containing 97½ to 98 p.c. of NaCN (equivalent to 129–130 p.c. of KCN)

The greater part of the world's cyanide requirements has been from its inception to the present date, 1920, supplied by this process in England. It was worked out, under Mr. Castner's direction, at the works of the Aluminium Co at Oldbury; the first manufacturing plant was erected, by the writer, at Frankfurton-Main, just before Mr. Castner's death in 1899, and the second at Glasgow in 1900. The experienced assistance of the staffs of the Frankfurt and Glasgow companies led to rapid improvement of working details A third installation started work near New York in 1901.

'Schlempe' Process -The juice of the sugar beet contains both inorganic and organic substances other than sugar, and these substances accumulate in the molasses In Germany, rather more than half of the molasses made are treated with strontia, which combines with the sugar to form an insoluble sucrate, the residue left after removing this, the concentrated essence of the impurities of the original juice, is known as 'schlempe' After concentration to sp.gr 14, it contains 75 pc of dry substances, 30 p c of which are inorganic (mainly potassium) salts and the remainder organic substances The 'schlempe' contains about 4 p c of nitrogen in the form of betaine

and decomposition products of the vegetable proteins. Of the molasses not treated by the strontia process, a large part is consumed as cattle food and a small part fermented to alcohol. The residue of the alcoholic fermentation is known in France as 'vinasses'

The first attempt to utilise the nitrogen of 'vinasses' was made by Vincent in 1877-1880 He distilled the 'vinasses' in horizontal iron retorts at a red heat, ammonia, a mixture of the three methylamines, monoethyl-, monobutyl, and monopropylamines, methyl alcohol, and combustible gases was evolved, and potassium carbonate mixed with some carbon remained behind (E Duvillier and A Buisme, Ann Chim Phys 1881, [5] 23, 289, H Ost, Zeitsch angew Chem 1906, 19, 609) In 1879 Ortheb and Muller devised a process for converting Vincent's methylamines into hydrocyanic acid, which was worked by the Société Anonyme de Croix (Eng. Pat. 3844, of 1879) The vapour of the methylamines was passed through a redhot retort, yielding ammonia, hydrocyanic acid, methane, and hydrogen The ammonia was absorbed by sulphuric acid, and the hydrocyanic acid by a suspension of ferious hydroxide in caustic potash. The manufacture of the methylamines was abandoned in 1881 for want of a market, and with it the production of cyanides.

The problem was again attacked by Bueb (Reichardt and Bueb, Eng Pat. 7171 of 1895, Bueb, Eng. Pat. 26259 of 1898) Instead of separating the products of distillation of the 'schlempe,' he passed them directly, after condensing some tar, into a series of firebrick channels which were heated to bright redness The ammonia and hydrocyanic acid were absorbed together in a solution of ferrous sulphate, and the product worked up in the same way as that made in the gas works The narrow flues were soon stopped up by deposits of carbon, so that they were replaced by chambers filled with fire-brick checker-work, which is heated up to 1000° by firing directly with producer gas, during this process, the deposit of carbon is burned Two such chambers are used alternately, the one being heated up whilst the gases from the retorts

chamber, the methylamines disappear and hydrocyanic acid is formed, the ammonia undergoing but little change. The ammonia is now removed from the gas by absorption in sulphunic acid, and the hydrocyanic acid absorbed in water in a suitable column. The solution of hydrocyanic acid is then distilled into a solution of caustic soda (A. Vasseur, Bull. Soc. Ind. Nord de la France, 1913, 41, 445) The solution of sodium cyanide is evaporated to crystallisation, the crystals dried, and briquetted by compressing them into blocks which are dried at 70° in vacuó (Eng. Pat. 18096, of 1907), and then contain sodium cyanide equivalent to 120 p.c. of KUN.

Of the nitrogen in the 'schlempe,' about onehalf is converted into nitrogen gas and so lost, a quarter is obtained as sodium cyanide and the remainder as ammonium sulphate

The decomposition of diand tri-methylamine, when passed through a heated porcelain tube filled with fragments of porcelain, fire-brick, or charcoal, has been studied by Voerskelius (Dissertation, Hannover, 1909). At temperatures lying between 800° and 1000°, as much as 98 p.c. of the trimethylamine decomposes into hydrocyanic acid and methane N(CH<sub>3</sub>)<sub>3</sub>=HCN+2CH<sub>4</sub>, the remaining 2 p.c. yielding ammonia N(CH<sub>3</sub>)<sub>3</sub>+3H<sub>2</sub>=NH<sub>3</sub>+3CH<sub>4</sub>. This result was obtained when the gaseous contents of the tube were changed once in 7½ seconds; with a slower current of gas and also with a contact material of large surface (fireday or charcoal) or with iron, a part of the hydrocyanic acid is decomposed into its elements. The rate of decomposition at 1000°, however, is small. At 720° the yield of hydrocyanic acid was only 78 p.c. and at 600° none was produced.

was only 78 p c, and at 600° none was produced. The formation of hydrocyanic acid from dimethylamine also begins between 600° and 700°, from 800° to 1100° the change takes place quantitatively in accordance with the equation NH(CH<sub>3</sub>)<sub>2</sub>=HCN+CH<sub>4</sub>+H<sub>2</sub>. If hydrogen is present, however, a second change occurs NH(CH<sub>3</sub>)<sub>2</sub>+2H<sub>2</sub>=NH<sub>3</sub>+2CH<sub>4</sub>. Using equal volumes of hydrogen and dimethylamine, this reaction is of no importance, but as the hydrogen present increases more and more of the dimethylamine is converted into ammonia until, with a mixture of 97 parts of hydrogen to 3 parts of dimethylamine, ammonia is the sole product

The behaviour of monomethylamine is probably analogous to that of dimethylamine A Wurtz (Ann Chim Phys. 1850, [3] 30, 443) found that, at a red heat, it decomposes in accordance with the equation

$$3CH_3.NH_2 = 2HCN + NH_3 + CH_4 + 3H_2$$

In practice, therefore, the losses of hydrocyanic acid and ammonia by decomposition will depend on the rate of passage of the gases and on the nature of the contact material—the deposit of carbon on the porous bricks diminishing the decomposition of hydrocyanic acid

this process, the deposit of carbon is burned. Two such chambers are used alternately, the one being heated up whilst the gases from the retorts are passing through the other. The gases has made practical application of these observations by passing the decomposing chambers contain 7 pc HCN; 7 pc, NH $_3$ ; 8 pc (mainly) CH $_4$ ; 12 pc H $_2$ ; 18 p.c. CO; 24 pc. CO $_2$ ; and 24 pc. N $_2$ . In passing through the decomposing

of the nitrogen is obtained as ammonia or hydrocyanic acid.

Manufacture of Sodium and Potassium Cyanides from Sulphocyanides.—The sulphocyanide is first conveited into ferrocyanide by one of the methods already described, and the ferrocyanide then worked up into cyanide; this is the method actually in use. Other more direct processes have been used to some extent

The earliest of these was D J. Playfair's process (Eng Pat 7764 of 1890) Playfair found that certain metals, lead and zinc especially, when fused with potassium sulphocyamide, abstract the sulphur, leaving potassium cyanide, which may be separated from the metallic sulphide by extraction with water. The fusion is carried out at 400°, and 70–80 p c of the sulphocyamide is converted into cyanide (D J. Playfair, J. Soc Chem Ind 1892, 11, 14, Conroy, *ibid.* 1896, 15, 8) The reaction is slower with lead, owing to the small surface offered by the fused metal, whilst with zinc the product is apt to be contaminated by the double cyanide of zinc and potassium. The process was worked about 1894 by the British Cyanides Co at Oldbury (Brussels Exhib 1910, Catalogue of British Chem. Section, p. 63).

of British Chem. Section, p. 63).

Raschen's process (Eng. Pats. 10476, 10956, and 21678 of 1895, 19767 of 1898, 12180 of 1900) is based on the oxidation of sulphocyanic acid to hydrocyanic and sulphuric acids which was first observed by E. A. Hadow (Annalen, 1858, 108, 380). Hadow found that permanganate, lead peroxide, manganese dioxide, and nitric acid bring about the same reaction fraction of the same reaction acid solutions only when an excess of the oxidising agent is present. If the oxidising agent is added to the sulphocyanide, solid yellow compounds are produced of variable composition, one of which, 'canaiin' (qv), has been used as a yellow dye

The Raschen process (J T Conroy, J. Soc Chem Ind 1899, 18, 432) was carried out by allowing a solution containing about 15 p e of sodium sulphocyanide to flow into dilute nitric acid kept at the boiling-point by means of live steam. The reaction:

#### NaCNS+2HNO<sub>3</sub>=HCN+NaHSO<sub>4</sub>+2NO

From 3 to 5 pc of the nitric acid was less completely reduced, N<sub>2</sub>O<sub>3</sub> being formed, so that the gases leaving the decomposer were first scrubbed with a little water The resulting solution of nitrie, nitrous, and hydrocyanic acids The gas, was returned to the decomposer which now contained about 1 volume of hydrocyanic acid to 2 volumes of nitric oxide, was passed through a solution of caustic soda (sp.gr. 137), which absorbed the hydrocyanic acid. It was found necessary to avoid saturating this solution completely with hydrocyanic acid because of the brown decomposition products which are formed from the free acid in presence of a cyanide The nitric oxide was then mixed with air and passed up-The nitric wards through towers packed with flints down which water trickled, in this way it was reconverted into nitric acid (sp gr. 1 12) which was returned to the decomposer. The solution of sodium cyanide was evaporated to dryness in vacuum pans without appreciable loss.

The reduction of sulphocyanides by hydrogen was studied by Playfaii (J. Soc Chem Ind. 1892, 11, 14). He found that hydrogen acts on potassium sulphocyanide at a dull-red heat, forming potassium cyanide and sulphide and sulphiretted hydrogen.

455

Conroy, Heslop, and Shores (J Soc Chem. Ind 1901, 20, 320) showed that the reaction takes place slowly at 500°, and fairly fast at 600°. Their results indicate that two reactions occur simultaneously:

 $KSCN+H_{2}=KCN+H_{2}S$ 

and

 $2KSCN + 2H_2 = K_2S + 2HCN + H_2S$ 

A little ammonia is also evolved, probably owing to the difficulty of drying the sulphocyanide completely. About 70 p c of the nitrogen is obtained in the form of cyanide and 20 p c as hydrocyanic acid. With barium and cuprous sulphocyanides, the reaction is similar to that represented by the second of the above equations, the principal introgenous product being hydrocyanic acid. With the cuprous salt, the reaction begins below  $300^\circ$ 

Rossiter, Crowther, and Albright (Eng Pats 4403 and 6226 of 1901) have patented a process based on the behaviour of the cuprous salt. It is mixed with finely divided copper and heated at temperatures rising from 150° to 650° in a current of hydrogen Hydrocyanic acid is evolved and cuprous sulphide remains behind

Synthesis of Cyanides from Uncombined Nitrogen. Up to the outbreak of the European War in 1914 the mtrogen of cyanides was almost exclusively derived either from organic matters of animal origin or from coal The persistent attempts to synthesise cyanogen compounds from atmospheric nitrogen, which date back to the first half of the nineteenth century, have been renewed during recent years, and important progress has been made. The deficiency of cyanide caused by war conditions has, to a small extent, been supplied by the product of one of the processes using atmospheric nitrogen, but it remains to be seen whether any one of them will be able to hold its own in competition with the older sources of supply under more normal conditions

The attacks on the problem of fixing free nitrogen in the form of cyanide have been made by several paths which sometimes intersect. For convenience of description, they may be arranged according to the raw materials used, the action has been studied of introgen on:—

- 1 a. Alkalı metal carbonates and carbon
  - b. Alkalı metals and carbon
- c. Barium carbonate and carbon
- 2 a. Carbides of metals.
- b. Acetylene
- 3. Metals forming nitrides, subsequently converted into cyanides.

#### 1 a. Action of Nitrogen on Alkali Metal Carbonates and Carbon.

Scheele may have been the first to fix atmosphere pitrogen in the form of a cyanide Writing in 1783, he says (Chemical Essays, 1901, reprint, 288). We know that charcoal powder, brought into fusion with fixed alkalis, yields likewise a kind of lixivium sanguinis, though weak.' The

source of the mtrogen is, however, doubtful, | since charcoal contains a little combined nitrogen. Desfosses (Ann Chim. Phys. 1828, 38, 158) mentions that notable quantities of cyanide are obtained by the action of nitrogen on a mixture of potash and carbon heated to redness, but the first thorough investigation of the reaction was made by Lewis Thompson, who was awarded the Gold Isis Medal of the Society of Arts, in 1838, for his improvements in the manufacture of Prussian blue. The chief improvement consisted in the substitution of atmospheric nitrogen for that contained in animal matters (Mechanics Magazine, 1839, 31, 92) (2 parts), coke (2 parts), and non tunings (1 part), were ground together to a coarse powder and heated in an open crucible, for half an hour, to a full red heat with occasional stirring, the end of the operation was indicated by the cessation of the jets of purple flame which arose from the surface of the mixture After cooling, the material was lixiviated, and Piussian blue precipitated in the usual way. The yield was better with larger quantities of material than it was with smaller batches, thus 6 oz of pearlash, containing 45 pc K<sub>2</sub>O, yielded 0.62 oz (295 grains) of Prussian blue, whilst 12 oz (1 lb) of the same material yielded 2.8 oz (1355 grains) of blue, 1 that is, 0 23 and 0 54 part of Prussian blue for 1 part of K2O used Assuming that the Prussian blue was dried at 30°-40', and contained 28 pc of water, the theoretical yield is 2 1 parts, so that Thompson cyanised from 11 to 26 p c of the potash used This calculation may underestimate the yield, because it is stated that the Prussian blue spoken of is the 'pure perferrocyanate of iron'. Coke was used (because wood charcoal burns away too quickly), and the 'carbonaceous matter may be worked over again many times, and is even improved by each operation.' The importance of the addition of iron is insisted on, when it is not employed a much higher temperature is required

The importance of Thompson's discovery of the catalytic action of iron was not fully appreciated for nearly three-quarters of a century, perhaps because the addition of iron for an entirely different purpose was a familiar feature of the current process of manufacture of ferrocyanide Contemporary criticism was concerned mainly with the source of the nitrogen, although Thompson's statement that carbonaceous matter improves on re-use would, if confirmed, have put the matter beyond doubt

G. Fownes, in a paper read to the British Association (The Chemist, 1841, 2, 261), insisted on the possible error due to the presence of nitrogen in the coke used, and found, like Scheele, that charcoal or coke yields 'abundance of cyanide' when heated with potash in a closed crucible at a moderate red heat, which is not the case when pure carbon is used When pure nitrogen was passed over a mixture of pure potassium carbonate and sugar charcoal heated, in a porcelain tube, he observed that carbon monoxide was evolved at first in much greater volume than that of the nitrogen passed in. When the evolution of carbon monoxide ceased he found in two experiments at a 'full red heat' that 111 and 121 pc. of the potash had been converted into cyanide. whereas at 'whiteness, much above the meltingpoint of copper,' the production of cyanide

appeared much greater

The repetition of Fownes' experiment by several chemists added little to our knowledge, and may be dismissed in a few words Erdmann and Marchand (J. pr Chem. 1842, 26, 407) observed only a doubtful trace of (yanide, Liebig and Wohler (Handworterbuch der Chemie, 1842, ii 384) could not confirm Fownes' result, and Delbiuck (J pr Chem 1847, 41, 161) also obtained a mere trace of cyamde. None of these authors gives details of then methods Bunsen and Playfan (But Ass. Rep 1845, 142), on the other hand, by heating a mixture of pure potassium carbonate and sugar charcoal in a gun bairel in a current of nitrogen at a temperature high enough to form potassium (about 1200°), obtained 6.982 grams of silver cyanide from an unmentioned quantity of potash It may be surmised that the different results noted were due to differences of temperature

In addition to these laboratory experiments, a most remarkable attempt was made to develop this process on a manufacturing scale Possoz and Boissière (Eng. Pat. 9985 of 1843, granted to their agent A. V. Newton) began their experiments at Grenelle, near Pans, in 1843, but transferred the plant in 1844 to the works of Messrs Bramwell & Co (one of the leading prussiate makers of that time) at Newcastleon-Tyne The experiments were carried on by Mr Bramwell and Mr F R Hughes until 1847, and a detailed account of them, furnished by Mi Hughes, was published by Giaham (Report of Junes, Exhib of 1851, 1, 95)

Wood charcoal was soaked in a solution of ordinary potash and dried, yielding a mixture containing about 20 pc K<sub>2</sub>O. This was, at first, heated in vertical fire-clay retorts of small diameter, which were fired externally. Owing to the low thermal conductivity of the retorts and of the charcoal, the heat penetrated the mass very slowly, and the output of a tube was  $_{
m small}$ It may be pointed out that this very serious difficulty is accentuated by the endo-thermic nature of the reaction, which was unknown at that time; it is not only necessary to heat the charge to the reaction temperature, but the large quantity of heat absorbed by the reaction

 $K_2CO_3+4C+N_2=2KCN+3CO-129$  Cals

must also be supplied

After a number of alterations, retorts were adopted, which were built of ordinary firebrick; they were 2 feet in internal diameter, and were heated for a length of 6 to 8 feet to a temperature sufficient to produce cyanide In an upper prolongation, the wet charcoal was dried and preheated, and in a lower, unheated extension, made of cast iron, the product was cooled down, and then discharged, by means of an arrangement which permitted a periodical withdrawal of a part of the charge, into a solution of a ferrous salt, into which the lower end of the cast-non tube dipped A ring of small slots was left in the brick retort at every third or fourth course of bricks, through which the hot

<sup>&</sup>lt;sup>1</sup> The bracketed figures are Thompson's; they are recalculated on the assumption that Troy weight was used.

mixture of nitrogen and carbon dioxide in the furnace was drawn into the retort by means of an aspirator attached to the upper end In this way the difficulty of heating the charge to its centre was overcome The highest temperature in the furnace was sufficient to soften a Stourbridge fireclay bink throughout its substance (1350°-1400°?) The hot gases were aspirated upwards, drying and preheating the fresh charge in the upper part of the retort As much as one-half of the alkalı in the cyanised charcoal produced was found to be in the form of cyanide, and 7 or 8 retorts of the dimensions given yielded 36-40 cwts of prussiate of potash per week, under favourable circumstances About onehalf of the charcoal used was consumed, and 3 parts of potash were used for 1 part of prussiate made About one-third of the potash was accounted for by the prussiate, one-third remained in the refuse charcoal, which could not be profitably lixiviated, and the remainder combined with the bricks of the retort, or was volatilised up the chimney. The financial failure of the process was due to the losses of potash and to the short life of the retorts

R Hoflmann (Report of Junes, Exhib of 1862, 59) also tried large-scale experiments at Occlenwald in the Black Forest, in 1858–1859, which may be referred to as a further illustration of the difficulties mentioned above. The alkalised coke was heated in wrought-iion tubes, not more than 1 foot in diameter, protected with fireday. The furnace was heated to a bright, white heat, which fused the common brick work behind the firebrick lining, but only minute quantities of cyanide were formed, the failure is ascribed to too low temperature, for it was found that the middle portion of the coke in a

tube was far from being white hot

In 1878, C A Fawsitt (Eng Pat 4544 of 1878) described a method of making ammonia by this process In his experiments, which were made at St. Rollox, Glasgow, he used sodium carbonate instead of potash (J Soc Chem Ind 1890, 9, 30) Using 3-inch or 4-inch wroughtnon tubes, he obtained as much as 50 p c of the theoretical yield of ammonia, but with 6-mch cast-iron tubes the yield was much smaller, because the east iron would not withstand the temperature needed. These experiments, therefore, indicate that a temperature exceeding 1100° (the approximate melting-point of cast iron) is required, although, in this case, the differences may have been due, in part, to the facts that a tube of larger diameter would give worse results even with the same external temperature, and that the nitrogen used with the 6-inch tubes contained from 3 to 5 p.c of

oxygen. The patents of Victor Alder of Vienna are of interest, because they contain a clear recognition of the catalytic rôle of iron, and of the necessity of obtaining intimate admixture of the reacting constituents, together with sufficient porosity to admit of free access of nitrogen Alder proposed to heat an intimate mixture of alkaline material and carbon in a closed vessel, in an atmosphere of nitrogen, and said. 'The addition of finely divided iron greatly promotes the conversion into cyanide, but is not absolutely essential to the success of the operation' (Eng. Pat. 1004 of 1880). In Eng. Pat. 5617 of

1881 he states that the reaction only succeeds in presence of carbonising gases, such as hydrocarbons or carbon monoxide, and that metals 'capable of fixing and transmitting the carbon, such as iron, chromium, manganese, nickel, or cobalt, greatly promote the conversion' The charge is prepared, for example, by mixing a solution of the alkali with charcoal and adding sawdust or other substance, which, when heated, will yield a porous mass, or a thin paste of the alkaline solution and finely ground charcoal containing some binding material is spread out on the surface of lumps of charcoal The 'car-bonising metal' is to be added as such, as oxide or in the form of a solution of a salt which, when heated in a current of steam, will decompose, leaving a deposit of the oxide, which, in the cyanising process, will be reduced to metal Alder appears, however, to have been under the impression that the success of a technical process depends on the employment of the cheapest raw materials, however unsuitable they may be, and a great part of the specifications consists of descriptions of the preparation of alkali-carbon mixtures from such substances as alkalı sulphates, limestone, and coal, as in the Le Blanc alkalı process, the failure of the process is not, therefore, surprising

The catalytic action of non was re-discovered by E Tauber in 1899 (Ber 1899 32, 3150, and more fully in Die Chem Ind 1903, 26, 26) Trials of different kinds of carbon and different alkalis showed that potash or soda and wood charcoal are the most active materials intimate mixture of alkali and charcoal was made by soaking sawdust in the alkaline solution and carbonising the mixture, non (reduced by hydrogen) in fine powder was added to this, 3 parts of iron to 1 part of soda giving the best The mixtures were heated, in iron boats enclosed in glass tubes, for 11 hours at temperatures between 'dark and moderate' red heats, a current of purified nitrogen being passed. In some cases more than 25 p.c. of the soda used was eyamsed, but usually a conversion of 10 p.c. was not exceeded. Magnesium, tungsten, chromium, and nickel were found to possess some catalytic activity, though much less than iron

Between 1843 and 1912 some 30 patents were granted in Great Britain for modifications of the process described by Possoz and Boissière, in which the blast furnace type of apparatus recurs frequently, no details of the results obtained, or of the causes of failure, appear to have been published Since 1912 the numerous patents of J E Bucher (especially Eng Pats 27713 of 1912, 23292 and 13332 of 1914; 2286 of 1915), which are being developed by the Nitrogen Products Co of Providence, Rhode Island, at their works at Saltville and Niagara Falls, have directed renewed attention to the catalytic action of iron Prof Bucher's results (J Ind Eng Chem 1917, 9, 233) surpass any which have previously appeared in print, this appears to be due partly to the fact that he has clearly realised that the necessary conditions, already known separately, must be fulfilled simultaneously, and partly to the ingenuity and care devoted to the detailed adaptation of means to this end

Carbon and iron oxide are separately ground

in a ball mill to pass 100-mesh sieve, and then re-ground together, after which dry sodium carbonate is added. The open-grained structure which is necessary to allow nitrogen to pass through the material is obtained by briquetting This is done by kneading the powder, at about 100°, with a little more water than is required to form the monohydrate Na<sub>2</sub>CO<sub>3</sub>,H<sub>2</sub>O, and squirting the paste into small rods about I inch long and 1 mch diameter These are dried at once, before cooling, because the expansion due to the formation of the decahydrate at temperatures below 35° causes them to fall to powder When heated above the melting-point of sodium carbonate (852°) these little briquettes become plastic to some extent, but retain their shape sufficiently well to permit free circulation of gases through a column several feet thick Mixtures of coke, non, and sodium carbonate in proportions from 2 2 1 to 2 2 2 give satisfactory results, the smaller proportion of sodium carbonate giving the fastest reaction owing to the greater accessibility of the active carbon-iron surfaces to nitrogen. The reaction is said to begin about 700°, and the eutectic temperature of iron-carbon (about 1150°) should not be exceeded because fusion of the particles of iron into larger globules diminishes their active The speed of the reaction, under surface suitable conditions, is considerable, for example, the 2 2 1 mixture heated, in 13 minutes, from 710 to 920 in a current of nitrogen, yielded a product containing 152 p.c. NaCN, 92 p.c. of the sodium being in the form of cyanide Sodium cyanide volatilises somewhat freely at 900°-950°, so that in a stationary column of material in a vertical tube, the upper layers are much richer than the lower ones, and a considerable loss of alkalı is observed, it is anticipated that this will not occur in a continuous process, since the incoming fresh charge will condense the evanide vapour.

In order to supply the necessary quantity of heat  $(Na_2CO_3+4C+N_2=2NaCN+3CO-138.5)$ Cals ) at a sufficient rate, Bucher proposes, for example, to use the hot charge itself as an electrical resistance (ef Readman's apparatus, below), or to use electrically heated molten iron saturated with carbon and to pass a mixture of nitrogen and sodium vapour through it

Producer gas may be used at high temperatures, but the cyanide disappears if the product is allowed to cool in an atmosphere containing carbon monoxide This may be due to the formation of carbon dioxide, as Buchei suggests, by the reaction 2CO=CO2+C, which is accelerated by iron, it is also possible that the reaction  $Na_2CO_3+4C+N_2=2NaCN+3CO$  is reversible, as has been proved in the analogous case of barium carbonate

Since this was written the reversibility of the reaction has been proved by J. B Ferguson and P. D. V. Manning (J. Ind Eng Chem 1919, 11, 946). Mixtures of sodium cyanide or of sodium carbonate (3 parts), charcoal (1 part), and iron (1 part) were heated in presence of definite mixtures of carbon monoxide and nitrogen. The composition of the product was not affected by the nature of the initial materials, but varied with the temperature and the com-position of the gas With 5 pc CO, for ex-ample, 75 p.c of the alkali was cyanised at

 $946^{\circ}$  and 80 pc. at  $1000^{\circ}$ ; with 48 p.c. CO, 27.5 pc and 59 pc were cyanised at the same temperatures Owing to volatilisation of cyanide it is probable that true equilibrium was not realised, the error will be greatest for products

rich in cyanide
C O Brown (J. Ind. Eng Chem 1919, 11, 1010) has described the plant which was erected by the Government of the United States in 1917-18 to produce 10 tons of sodium cyanide daily by the Bucher process Briquettes, made in the way described above, were heated at 1000° in vertical, gas-fired, steel, or mehrome tubes in a current of very pure nitrogen. The product, containing about 20 p.c. NaCN, was extracted with liquid, anhydrous ammonia; by this means the formation of ferrocyanide is avoided, and any unchanged alkali is left undissolved in the residues which are reground with sufficient carbon and soda and reused Evaporation of the ammonia solutions leaves sodium cyanide, containing 92 p.c. NaCN, as a white powder. The loss of ammonia was about 2 lbs per 100 lbs cyanide made. Each steel tube produced about 7½ lbs of sodium cyanido hourly, and had a life of 11-14 days. The work was abandoned after the armistice in 1918

It may be mentioned here that the Cassel Cyanide Co, of Glasgow, has used iron as a catalyst in the manufacture of cyanides from atmospheric nitrogen since the year 1909, when the writer first became fully aware of its great activity under suitable circumstances. The work in which sodium carbonate was employed as raw material was carried out entirely by Mr J. H Young Setting out from finely divided, precipitated ferric oxide and chaicoal (the purest and most active form of carbon available), the mixtures were made by grinding these substances separately and then together, finally adding the dry sodium carbonate The necessary porosity of the and re-grinding material to be treated with nitrogen was secured by adding a quantity of the very bulky charcoal obtained by carbonising shavings, the addition of 15 parts of this material to a ground mixture of 20 parts of sodium carbonate, 8 parts ferric oxide, and 15 parts charcoal, gave good results, 60-80 pc of the alkali being cyanised at 900°. The residual charcoal-iron mixture resulting from the lixiviation of the cyanised mass gave even better results than a fresh mixture. A larger scale, continuous plant was at work in 1913, in which the mixtures were heated in vertical steel tubes, of 6 inches bore and about 10 feet long, of which a length of about 5 feet was heated to the working temperature. The lower part of the tube served for cooling the product, which was withdrawn by means of a houzontal spiral conveyor, fresh material being fed in at the top of the tube Pure nitrogen was employed, and very careful measurements of the temperature showed that under satisfactory working conditions the temperature in the axis of the tube was 900°, near the inside wall of the tube it was 950°, whilst a thermocouple in the furnace in contact with the outer wall of the tube indicated 1010°. These measurements indicate quite clearly the necessity of using tubes of small diameter when external heating is relied on Temperatures higher than those mentioned were found to be undesirable

owing to their serious effect on the life of the steel tubes, and also on account of the danger of fusing part of the iron catalyst, and so diminishing its active surface. Each tube of the dimensions mentioned produced from 40 to 50 lbs of sodium cyanide in 24 hours, the product containing from 21 to 26 p.c. NaCN.

The results described, together with the older experience of the reaction between alkalis and nitrogenous organic matter, make it easy to understand the occurrence of cyanides in kelp, barilla, Le Blanc alkalı, and the residues from the preparation of potassium (which were known to Lewis Thompson in 1839), and especially in the blast furnace. Clark (Phil Mag 1837, 10, 729) described the occurrence of cyamde at the Clyde Iron Works, in a furnace using raw coal and hot blast, in quantities such that wheelbarrows full could be obtained Zincken and Bromeis (J pr Chem. 1842, 25, 246) found a quantity of potassium cyanide in the charcoal remaining in the hearth of a furnace in the Harz which had been closed down furnace had been worked with charcoal and a hot blast Redtenbacher (Annalen, 1843, 47, 150) described the formation of potassium cyanide in quantity (it was used for electroplating) in a charcoal non furnace at Mariazell in Steiermark, which was also worked with blast heated as high as 350° Bunsen and Playfair (But Ass Rep. 1845, 142) observed the production of more than 2 cwts per day of potassium cyanide in the coal-fired furnace at Alfreton, which worked with blast at 330°, they also showed that the potassium was derived mainly from the non ore used, the furnace charge containing sufficient potassium to produce 377 lbs of KCN per day The cyanide was found in the hottest part of the furnace, 2 feet 9 inches above the tuyeres, but some 7 feet higher there was very little Sir Lowthian Bell (J. Soc Chem Ind 1890, 9, 710) gives the following figures for the distribution of cyanogen in a more modern furnace at the Clarence Works, which was worked with coke and Cleveland iron ore and blast at about 700° .-Height escape

8 ft. 24 ft 52½ ft. 60 ft. pipe above hearth. Grams CN 49 06 15 76 7 67

5.94 4.73

per cb m gas It is generally assumed, probably correctly, that the blast furnace cyanide is produced from free nitrogen, but it is interesting to note that Sir L Bell's furnace consumed 50 tons of coke

per day, which probably contained at least I ton of combined nitrogen.

The mechanism of the reaction which occurs between carbon, alkalı carbonates, and mtrogen, and the rôle of iron, are still quite obscure. old view, that carbon and nitrogen ('nascent' or otherwise) unite to form cyanogen which is then fixed by alkalis, is extremely improbable in view of the nature of the equilibrium

$$2C+N_2 \gtrsim C_2N_2$$

(v. Cyanogen), and of the fact that no one has succeeded in isolating a trace of cyanogen gas made by the direct union of the elements. The possibility that a carbide may be an intermediate product is discussed later. It is, at least, an ascertained fact that some metallic acetylides do combine with nitrogen, although the behaviour of

the alkalı metal derivatives has hardly been investigated. The intermediate formation of free alkali metal has also been asserted to be necessary, but as the alkalı metals (except lithium) do not combine directly with nitrogen, whereas they do combine with carbon, this seems to be identical with the assumption of the intermediacy of a carbide. No valid experimental evidence appears to have been advanced in support of the existence of any intermediate compound

1b. Action of Nitrogen on Alkali Metals and Carbon.

This reaction appears to have been observed for the first time by R F Marchand (J pr Chem 1850, 49, 351) A statement that cast iron and steel contain from 05 to 12 pc of nitrogen induced him to test some 30 samples of cast iron by Lassaigne's method, that is, by heating the sample to redness, in a test tube, with potassium, extracting the product with water, and applying the Prussian blue test to the solution A copious precipitate was obtained in every case He found, however, that, when the residue of non was dried and again heated with potassium, there was a further production of Prussian blue, and that this continued so long that, with 8 grams of powdered iron, it was impossible to exhaust the reaction. This led Marchand to suspect that the nitrogen was derived from the air, and he proved that this was the case by carrying out the reaction in an atmosphere of hydrogen or of carbon dioxide, when no more than a doubtful trace of blue was obtained, and, further, by heating the mixture of cast iron and potassium (after drying it in a current of hydrogen) in an atmosphere of nitrogen in a closed tube, when he found that the nitrogen was completely absorbed He also showed that steel behaves like cast iron, but soft non Marchand did not obtain the reaction with a mixture of carbon and iron obtained by heating ferric succinate or benzoate, and therefore concluded that success depends on the carbon being in chemical combination with the iron.

The success of Lassaigne's test, when applied to organic substances containing sulphur, had been questioned by O Jacobsen, who recom-mended the addition of reduced non in such Making the test in this way, I. Remsen (Amer. Chem J. 1881, 3, 134) obtained Prussian blue from substances, such as pure cane sugar and sodium potassium tartrate, which contain no nitrogen After proving that the reduced iron employed was not the source of the nitrogen, Remsen heated a mixture of the tartrate, pure reduced iron and sodium, first in hydrogen to dry it, and then in nitrogen, obtaining a very large precipitate of Prussian blue. He also made the new observation that the iron powder loses its activity when it is superficially oxidised by exposure to air, which may perhaps explain Marchand's failure to obtain the reaction with his mixtures of iron and carbon

The reaction was re-discovered by E. Tauber (Ber. 1899, 32, 3150), whose experience coincided almost exactly with that of Remsen.

Y. Castner, in 1894, patented a process in which sodium was allowed to flow downwards over a column of red-hot charcoal in

an non vossel, a current of nitrogen passing upwards through it (Eng. Pat 12218 of 1894). It is doubtful whether this process received more than a cursory trial—It was taken up again by the Roessler and Hasslacher company about 1910—They find that a mixture of sodium vapour and nitrogen reacts with charcoal in the absence of non at 650°, and that the speed of reaction is sufficient for manufacturing purposes at temperatures above about 800°. (H. Philipp, Chem and Met Eng 1920, 22, 313, U.S. Pats 1249821, 1917, 1235887, 1917).

In 1912, J. E. Bucher applied for a patent

for the manufacture of evanide by the direct union of its elements in presence of iron (Eng. Pat. 11797 of 1913, U.S. Pats 1082845 of 1913, and 1116559 of 1914). A temperature of from 500° to 800° is said to be suitable, and iron and carbon are to be intimately mixed Like Marchand, Bucher also observed the removal of carbon from cast iron or steel by the action of sodium and nitrogen on them He also follows Marchand in adopting the view that iron acts as a solvent of carbon, the dissolved carbon being supposed to be capable of entering into combination with sodium and nitrogen When sodium carbonate is used, it is assumed that a little metallic sodium is formed by the action of carbon, even at the low temperature (700°) at which the reaction proceeds, the removal of this, by conversion into cyanide, distuibs the equilibrium and permits the decomposition of further quantities of sodium carbonate hypothesis appears to be open to the criticism that it makes the extremely rapid rate of formation of cyanide depend on the rate of solution of solid carbon in solid iron, it also fails to make clear why the usually meet mtrogen becomes so active

The Cassel Cyanide Co's experiments with this process were begun, as already stated, in 1909, in 1911 engineering difficulties had been so far surmounted that a unit plant producing about 50 lbs of cyanide hourly was at work. From the experience thus gained a larger industrial unit was designed and erected, which was producing cyanide for some time before war was declared in 1914.

# 1c. Action of Nitrogen on Barium Carbonate and Carbon.

In 1860 Margueritte and De Sourdeval (Eng Pat 1171, of 1860, Compt 1end 1860, 50, 1100) discovered that a mixture of barium oxide and carbon absorbs nitrogen readily forming barium cyanide. Ludwig Mond (Eng Pat 433 of 1882, J. Soc. Chem Ind 1889, 8, 505) experimented with this process in 1882. He found that a temperature of 1200° was required for the formation of cyanide, and that the reaction goes best about 1400°. The barrum compounds fuse readily at these temperatures and attack the fireclay retorts. To avoid this he briquetted them with an excess of carbon and pitch apparatus used was similar to that of Possoz and Boissière, and 40 p.c. of the bailum used was readily cyanised. The object of Mond's experiments was the production of ammonia, and he abandoned this process in favour of the more promising method of acting on coal with air and steam in large excess (see Ammonia).

As Mond pointed out, the reaction is endo-

$$BaCO_3+C=BaO+2CO-101\ 2\ Cals.$$
  
 $BaO+3C+N_2=Ba(CN)_2+CO-43\ 1\ Cals.$ 

The introduction of internal electrical heating by Readman (Eng Pat 6621 of 1894) therefore marks a very considerable step in advance His process was worked by the Scottish Cyanides Co at Leven in Fifeshire, between the years 1899 and 1907. Considerable quantities of sodium cyanide of excellent quality were produced, but the financial results were unfavourable. The following account of the process in its final form is believed to be substantially accurate.

Barium carbonate, produced in a later stage of the process, was intimately mixed with ground coal to a thick paste, which was dried and coked in ordinary gas retorts The granular mixture of barium oxide and coke so produced was then fed into an electric furnace, consisting of a vertical shaft of circular cross-section, lined with magnesia bricks A ring of carbon blocks in the wall of the shaft near its lower end formed one electrode, a carbon cylinder suspended in the mouth of the shaft formed the other The alternating current passed between these electrodes through the granular mass of coke and barium oxide, heating it to a high tempera-A current of producer gas with about 70 p c of mtrogen, passed upwards through the mass, which was constantly fed downwards into a cooling chamber by means of a plate and rotating arm which closed the lower end of the shaft, fresh material being fed in to the annular space found the upper electrode. The cooled product was lixivated with water, and the resulting solution, containing ballum cyanide, cyanamide, and hydroxide, treated with sodium bicarbonate After filtering off the barium carbonate, the solution of sodium cyanide was concentrated in racub and then passed over cooling coils depositing crystals of the hydrate NaCN,2H,O These were separated from the mother liquor in a centrifugal machine, and then dehydrated by heat and the powder briquetted The mother liquor, from which no further crystals could be obtained, was distilled with sulphuric acid to recover the residue of hydrocyanic acid

The chemistry of this process has been studied in some detail at temperatures up to 1150°, but little is known of it at higher temperatures. Undecomposed barium carbonate, as Finkelstein (Ber. 1906, 39, 1585) showed, does not melt at 1350°, but when heated alone or with carbon, it is partially converted into barium oxide, which yields readily fusible mixtures with the carbonate. Le Chatcher, for example, found that partly decomposed barium carbonate melts at 795°. The fusibility of the mixtures is increased by the formation of barium cyanide (which melts about 600°), so that the material under treatment in this process is a liquid phase of variable composition.

Kuhling and Berkhold (Ber 1908, 41, 28) found that when a mixture of barium carbonate and carbon is heated in a current of nitrogen under constant conditions, a little more than 40 pc. of the barium combines with nitrogen at 1150°, and that at lower temperatures the quantity combined diminishes, reaching zero at about 925°.

Ewan and Napier (J. Soc Chem. Ind. 1913, 32, 467) also found that the absorption of nitrogen begins at 925°. They showed, however, that it depends on the two reversible reactions  $BaCO_3+C \rightleftharpoons BaO+2CO$  and

$$\rm BaO + 3C + N_2 ~ \rightleftarrows ~ Ba(CN)_2 + CO$$

If carbon monoxide is allowed to accumulate, therefore, the change is brought to a standstill, if it is swept out with large quantities of pure nitrogen a considerable portion of the barium may be cyanised even at comparatively low temperatures Using 29 molecules of nitrogen to each molecule of BaO, for example, over 40 p c of the barium combined with nitrogen in 2 hours at 960°.

The pressure of carbon monoxide in equilibrium with pure barium carbonate and carbon is relatively large, it falls, however, to very low values for the dilute solutions of the carbonate in barium oxide or cyanide. For this reason the greater part of the barium carbonate decomposes before any cyanide is formed, but the decomposition is never complete.

When a mixture of BaCO3+BaO+C or of Ba(CN)<sub>2</sub> +BaCN<sub>2</sub>+C is heated in a mixture of carbon monoxide and nitrogen of definite composition until no further change occurs, the percentage of the barium in the product, which is combined with nitrogen (as cyanide or cyanamide), is independent of the composition of the initial mixture, and depends only on the temperature and on the composition of the gas used The maximum quantities of barium which can combine with nitrogen under different conditions, observed in this way, are given in the following table .-

c of CO	P c of the barum combined with n			
the gas	1000°	1100°	1150°	
0	50	50	50	
10	28	41	44	
20	5	32	38	
30		22	32	
40		13	26	
50		4	15	
70			9	

The curves drawn through the experimental points appear to be straight lines converging towards 50 pc. conversion for pure nitrogen, it is probable, however, that this is not their true form at very small concentrations of carbon monoxide, because a product in which 571 pc. of the barium had combined with nitrogen was obtained by continuing the heating for 82 hours at 1100° with nitrogen containing only 0.1 p c of GO For practical purposes, the rapid increase in the quantity of carbon monoxide required to reverse the reaction at higher temperatures is important

In addition to the two equilibria just mentioned, the reaction Ba(CN)<sub>2</sub>=BaCN<sub>2</sub>+C occurs Pure barium cyanide decomposes in this way even at 500°, the reaction is a slow one, which is accelerated by iron. The results appear to show that at temperatures between 600° and 1000°, and in absence of barium oxide, the decomposition ceases when about two-thirds of the

nitrogen is in the cyanamide form, in presence of barium oxide the change does not go so far. The relative quantities of cyanide and cyanamide produced by the action of nitrogen on mixtures of barium oxide and carbon are very erratic, anything from 0 to 70 p c. of the nitrogen fixed may be in the cyanamide form, the average of 36 experiments was 15 p.c.

The equilibrium pressure of carbon monoxide in the reaction  $BaO+3C \gtrsim BaC_2+CO$  was found to be 1 mm of mercury at  $1200^{\circ}$  Cyanide formation therefore proceeds freely in presence of quantities of carbon monoxide, which are quite incompatible with the existence of barium The suggestion that barium carbide is an intermediary in this process is, therefore, incorrect.

The effect of pressure was studied by Hempel (Ber. 1890, 23, 3388). The mixture of banum oxide and carbon was heated for 15 minutes by the same quantity of electrical energy in a closed bomb; the quantity of hydrocyanic acid obtained increased 16 times when the nitrogen pressure was increased from 1 to 60 atmospheres The improvement may have been due to greater reaction velocity of the more concentrated nitrogen or to a larger value of the ratio of nitrogen to carbon monoxide in the bomb Stahler (Ber 1916, 49, 2292), by a somewhat similar method, finds that with 60 atmospheres N pressure a product containing 74 p c of barium cyanide and no appreciable quantity of cyanamide is obtained.

The addition of barrum chloride or of potassium carbonate to the mixture of barium oxide and carbon neither reduces the minimum temperature nor increases the speed of the reaction

The effect of the addition of calcium and barium fluorides was studied by N (Zeitsch angew Chem. 1910, 3, 2405) at temperatures about 1100° and 1200° Since the Since the quantity of carbon monoxide in the nitrogen was not observed it is impossible to draw definite conclusions from the results, it is, however, of interest that in the 15 experiments recorded the quantity of the barium which combined with nitrogen never exceeded one-half of that

Very little information exists about the course of the reaction at higher temperatures From Eng Pats 2133 of 1907, and 22039 of 1906 (Badische Co), it may be gathered that 80 p c of the barium can be combined with nitrogen at 1500°, and, according to Kaiser (Fr Pats. 454237-8 of 1913), 90-95 pc of the theoretical quantity of nitrogen is absorbed, when barrum carbonate and carbon are heated '900° to 1400° ' in nitrogen and cooled in it under pressure It appears, therefore, that the inactivity of one-half of the barium, which has been shown to exist at temperatures below 1200°, may disappear at considerably higher temperatures; possibly because a different mechanism comes into play Ewan and Napier (lc), by applying Nernst's thermodynamic theorem to their measurement of the equilibrium pressure ( $p_{CO}$  in mm. of mercury) of the reaction BaO+3C  $\rightleftharpoons$  BaC<sub>2</sub>+CO, showed that, at absolute temperature T  $\log_{10} p_{CO} = -\frac{16250}{T} + 1.75 \log_{10} T + 5.48$ 

$$\log_{10} p_{00} = -\frac{16250}{T} + 1.75 \log_{10} T + 5.48$$

from which the value of  $p_{co}=100$  mm, at 1500° and 1026 mm at 1700°. M Kahn (Compt rend. 1907, 144, 913) found that barium carbide is formed from barium oxide and carbon heated in a well-closed crucible at the melting point of platinum (1760°), the equilibrium pressure at 1760° is, therefore, greater than I atmosphere, in approximate agreement with the calculated value It is therefore possible that barium carbide is an intermediate product at high temperatures.

# 2a Action of Nitrogen on the Carbides of Mctals.

F. R. Hughes (Report of Juries, Exhib of 1851, 1, 95) suggested that the first stage of the reaction between potassium carbonate and carbon might lead to the formation of potassium carbide, and that this might combine directly with nitrogen Potassium carbide is actually formed in this way, E Davy (Brit Ass. Report, 1836, n. 62), having discovered acctylene by treating with water, the black product of an unsuccessful attempt to prepare potassium Berthelot (Compt rend 1868, 67, 1141) was led to suggest the same mechanism by the analogy with his synthesis of hydrocyanic acid from acetylene and nitrogen. He also showed that the products of the action of potassium on potassium carbonate or on carbon monoxide contain potassium acetylide. The action of nitrogen on the alkali metal carbides has not been investigated in detail Frank and Caro (Eng Pat 15066 of 1895) mention that they combine at a 'dark red heat' forming cyanides

The absorption of nitrogen by the carbides of calcium, strontium, and barium has been much more completely studied. Moissan (Compt. rend 1894, 118, 503, and 684) found that pure, dry mitrogen did not react with the pure carbides even at 1200°, although the product obtained from barium carbide yielded a little ammonia when boiled with water In 1895 Frank and Caro (Eng. Pat 15066 of 1895), and T L Willson (Eng. Pat 21997 of 1895), discovered independently that the impure carbides absorb nitrogen According to Frank and Caro, moist nitrogen is necessary when the pure carbides are used, but not with impure substances, they also found that barium carbide yields the corresponding cyanide, but obtained an 'unsatisfactory result' with calcium carbide unless it was mixed with barium or sodium carbide Willson proposed to treat the carbides with nitrogen either during their formation in the electric furnace, or subsequently in a separate operation He also appears to have found difficulty with calcium carbide, and says that a cyanide or some more complicated nitrogen compound is formed. The fact that a large part of the combined nitrogen is in the form of a cyanamide salt was discovered by F Rothe, working under Frank's direction, in 1898 (Zeitsch angew Chem 1903, 658), and was published in Frank and Caro's Eng. Pat. 25476 of 1898. Frank and Caro at first attempted to apply

their discovery to the manufacture of cyanides.

Finely ground barium carbide, made in the electric furnace, was heated at about 700° in presence of pure mtrogen. The product contained about 11 p.c of nitrogen, 30 p.c of which was in the form of barium cyanide, the remainder

being barium cyanamide. In order to convert the cyanamide into cyanide, the material was heated with dry sodium carbonate and carbon. The sodium carbonate may convert the barium cyanamide into the sodium salt, which then takes up carbon in the way discovered by Dreschel,

$$\begin{array}{c} \operatorname{BaCN}_2 + \operatorname{Na}_2\operatorname{CO}_3 = \operatorname{Na}_2\operatorname{CN}_2 + \operatorname{BaCO}_3 \\ \text{and } \operatorname{Na}_2\operatorname{CN}_2 + \operatorname{C} = 2\operatorname{NaCN} \end{array}$$

By extracting the product with water, converting the sodium cyanide into ferrocyanide, and then fusing it with sodium, about 86 p.c. of the introgen fixed was obtained in the form of marketable sodium cyanide (G. Erlwein, V Int Kongress angew. Chem. Berlin, 1903, 1, 646) The process, however, even in this form, did not pay, and the cheaper calcium carbide was therefore tried. Commercial calcium carbide absorbs nitrogen at about 1100°, yielding a product containing from 20 to 23 5 pc. of nitrogen, which is wholly in the form of calcium cyanamide  $CaC_2 + N_2 = CaCN_2 + C$  Details of the manufacture of calcium cyanamide are given in the article on NITROGEN, ATMOSPHERIC, UTILISATION OF By heating this product with sodium chloride, from 90 to 95 p c of the cyanamide is said to be converted into sodium cyanido (G Erlwein, loc. ctt., G. Erlwein and Frank, U.S. Pat. 708333, 1902)

It is not easy to obtain such a high yield in the laboratory; Kıtawakı and Kusano, for example (J Chem Ind Tokyo, 1917, 20, 234), state that when 10 parts of calcium cyanamide (186 pc N) are heated with 16 to 24 parts of salt, and 2 parts of charcoal for 10 minutes at 900°-950°, and the fused product cooled in absence of oxygen, it contains more than 70 p.c. of the theoretical quantity of cyanide

Large-scale experiments were carried on in Germany with but little success; a description of these, up to 1914, has been published by W. S Landis (Chem. Met Eng. 1920, 22, 265) In the latest modification, from 2 to 3 parts of salt were fused separately, and 1 part of calcium cyanamide added gradually in order to control frothing. The highest temperature used was 960°, and from 40 to 75 p c of the nitrogen was

obtained in the form of cyanide In 1916 the process was installed at Niagara. It was soon found that the troublesome evolution of gas was almost suppressed by the addition of a little calcium carbide, and that much better results were obtained by raising the temperature to 1200°-1400° (H. P. Eastman, U.S. Pat. 1282395, 1918), and also that the cyanide disappears when the fused product is allowed to cool slowly, the reversion apparently occurring between 400° and 700° (H. Freeman, U.S. Pat. 1282405, 1918) In 1919 the process was, therefore, carried out by feeding a mixture of calcium cyanamide (1 part), salt (about 0.8 part), and calcium carbide (0.05 to 0.1 part) into a single phase electric furnace of the arc-resistance type with conducting hearth and a single suspended electrode. The furnace treats about 30 tons daily. The mixture fuses, and is run off almost continuously into a pan in which a water-cooled drum rotates, taking up a thin layer of the fused material which solidifies and is scraped off in thin flakes which contain cyanide (possibly the calcium salt) equivalent to 36 p.c. NaCN. The

efficiency on the nitrogen is about 90 p c. The equivalent of 975 tons of sodium cyanide was made in 1917, 1050 tons in 1918, and 1250 tons in 1919. The chemistry of this process is still incompletely known, Landis states that cyanide is formed even when little or no salt is used, which suggests that the principal change is represented by the reversible reaction

$$CaCN_2+C \ge Ca(CN)_2$$

The initial frothing may be due to the decomposition of sodium eyanamide

$$2NaCN_2 = 2NaCN + N_2 + 2Na$$

the graphitic carbon present not being sufficiently active at lower temperatures to convert it quickly into cyanide; the addition of calcium carbide is supposed to supply the necessary active earbon.

# 2b Action of nitrogen on acetylene

In 1863 Berthelot had shown (Ann Chim Phys 1863, [3] 67, 52) that acetylene is formed in the electric are between carbon electrodes in an atmosphere of hydrogen, and also that hydrocarbons in general yield acetylene when induction sparks pass through their vapours. The careful measurements of Bone and Jerdan (Chem Soc. Trans. 1901, 1042) indicate that, when equilibrium is established in the gas surrounding a carbon are burning in hydrogen or hydrocarbon vapour, it contains 90–91 p.c.  $H_2$ , 7.8 p.c.  $C_2H_4$ , 1.25 p.c.  $C_4H_4$  and 0.75 p.c.  $C_2H_6$ , no other

hydrocarbons being present In 1868 Berthelot (Compt 1end 1868, 67, 1141) discovered that acetylene and nitrogen combine, under the influence of strong induction sparks, to form hydrocyanic acid, and that the reaction  $C_2H_2+N_2 \gtrsim 2HCN$  is reversible. Taken together with his earlier observations on the formation of acetylene, this led him to the generalisation that hydrocyanic acid must be formed in the carbon are burning in mixtures of hydrogen and nitrogen or when sparks are passed through mixtures of nitrogen and hydrocarbons The cyanogen compound observed when the arc burns between carbon electrodes in an atmosphere of nitrogen is hydrocyanic acid, the hydrogen of which is derived from the electrodes or from water vapour A mixture of 10 vols C<sub>2</sub>H<sub>2</sub>, 14.5 vols N<sub>2</sub>, and 75.5 vols H<sub>2</sub> is suitable for the synthesis of hydrocyanic acid, since mixtures richer in acetylene deposit carbon. Dewar (Proc Roy Soc 1879, 29, 188, and 1880, 30, 85) confirmed Berthelot's observations on the formation of hydrocyanic acid in the are, and also showed that it is formed when a mixture of hydrogen and nitrogen is passed through a carbon tube heated externally by an electric The minute traces of hydrocyanic acid found in the slightly luminous flame of a Bunsen burner (Lüdeking, Annalen, 1888, 247, 122) are

Subsequent investigations add little to the information given by Berthelot, Hoyermann (Chem. Zett. 1902, 26, 70) passed a mixture of nitrogen (2 vols.) and acetylene (1 vol.) into the arc through tubular carbon electrodes and found that 60 to 70 pcoof the acetylene was converted into hydrocyanic acid. Gruszkiewicz (Zeitsch. Elektrochem. 1903, 9, 83) found that

doubtless due to this reaction

a mixture of carbon monoxide (2 vols ) hydrogen (1 vol ) and nitrogen (1 vol ) yielded small quantities of hydrocyanic acid when exposed to induction sparks between platinum points Lipinsky (Zeitsch Elektrochem 1911, 17, 761) passed mixtures of methane, introgen and hydrogen through a 2200 volt alternating current are between platinum electrodes. No carbon was deposited from a mixture containing  $\mathrm{CH_4}$  (20 pc),  $\mathrm{H_2}$  (10 pc), and  $\mathrm{N_2}$  (70 pc). Under the best conditions about 1.75 gram of hydrocyanic acid was produced per kilowatt hour. It is instructive to compare this with the 50 grams of nitric acid per kilowatt hour which Raleigh obtained, in apparatus of comparable dimensions, from mixtures of oxygen and nitrogen

E Biner and A Bacifuss (Helv Chim Acta 1919, 2, 663) have obtained as much as 7 39 grams HCN per kilowatt hour by circulating 8-10 litres per hour of a mixture of 1 part methane and 5 parts introgen through an arc at 505 volts between platinum electrodes 12 mm. apart.

According to H Philipp (Chem. Met. Eng. 1920, 22, 213), Lipinski's process was tried on a commercial scale in 1914 at Neuhausen, and a yield of 30 grams of HCN per kilowatt hour

attained.

Hutton and Smith (Eng Pat 23835 of 1906, and Trans Amer Electrochem. Soc 13, 359, 1908) showed that the temperature of the arc is not necessary, hydrocyanic acid being formed when a carbon rod is heated in a mixture of hydrogen and nitrogen to 1700°, and H v Wartenberg (Zeitsch anorg Chem 52, 299, 1907) carried out quantitative measurements with the following results (using a mixture of equal volumes of hydrogen and nitrogen)—

Temperature	P c HCN in the gas
1635°	1.95
1752°	3 10
1875°	4 70

Wallis (Annalen, 1906, 345, 353), by passing the same mixture through the electric arc at varying rates obtained a maximum of 33 5 p e of HCN in the gas. The estimated temperature of the arc is about 3250°

Cyanides from nitrides. At high temperatures, introgen combines directly with several elements to form intrides, and these, when fused with sodium carbonate and carbon, produce sodium cyanide. This has been shown to be the case with the nitrides of boron, calcium, magnesium, titanium, silicon, and aluminium, and with phospham PNH, Since none of these processes has emerged as yet from the experimental stage, it is sufficient to mention the possibility of producing cyanides in this way.

Hydrocyanic acid. Prussic acid (Ger Blausaure) HCN

This substance was discovered by Scheele in 1782 (Chemical Essays, London, 1786; reprinted 1901) He observed that Lixivium sanquims loses its power of forming Prussian blue when it is exposed to the air, and traced this effect to the atmospheric carbon dioxide. Other acids also produced it, and the blue-forming principle was found in the vapour given off. By distilling

the haverum with sulphuric acid, he obtained an aqueous solution of hydrocyanic acid. more convenient method of preparation was to heat potassium ferrocyanide with dilute sulphuric acid In doing this, he obtained a blue residue (oxidised Everitt's salt or Williamson's blue) Finally, he prepared mercuric cyanide by boiling Piussian blue with meicuric oxide; by treating this with iron and sulphuric acid, he decomposed it into mercury and hydrocyanic acid, which he distilled off In 1783, Scheele found that hydrocyanic acid is inflammable, its combustion yielding carbon dioxide proved that it contains nitrogen by synthesising it from ammonia Equal parts of charcoal and potassium carbonate were heated to redness, but not sufficiently to melt the potash, in a crucible and then some ammonium chloride stirred in (for 2 minutes only), the mass was then thrown into water and a solution obtained which gave 'a great quantity of Prussian blue,' whereas a similar experiment without the sal-ammoniac yielded a very small quantity.

exact quantitative composition of prussic acid was ascertained by Gay-Lussac (Annales de Chimie, 1815, 95, 156) He showed that it is similar in constitution to hydro-chloric and hydrodic acids, the hydrogen being united with a compound radical, CN, which he called 'cyanogène,' which plays the part of an element in so far as it passes through many transformations unchanged

Compounds of hydrocyanic acid are widely distributed in nature in plants The acid does not exist as such in the living plants, but is liberated after their death by the hydrolytic action of an enzyme on a glucoside, both of which are contained in the living cells, the glucoside may also be split up by boiling with dilute acids—Rarely a glucoside is found without an accompanying enzyme, as in the leaves of the common elder (Sambucus niger) chemistry of the cyanogenetic glucosides is described under GLUCOSIDES. Very different quantities of hydrocyanic acid are obtained from different parts of the same plant, the quantity also depends on the age of the plant (it is usually greatest in the young plants) and on the conditions of growth such as climate and rainfall, the use of nitrogenous fertilisers has been found to increase the quantity of hydrocyanic acid present. Plants raiely yield as much as one or two tenths of 1 pc of their weight of hydrocyanic acid, for example great millet (Sorghum vulgare) 0 013 to 0 044 pc in young plants, an exceptional North American sample gave 0 114 pc; wild cherry (Prunus padus) 0 03 pc, cherry laurel (Prunus laurocerasus) up to 0 2 pc m young leaves, cassava, sweet, 0 016 pc in the leaves, 0 043 pc, in the peel of the stem, 0 005 pc in the edible part of the root, the corresponding parts of the bitter variety contained 0.041 pc, 0.113 pc, and 0053 pc respectively, Rangoon beans (Phaseolus lunatus), 0 025 p c; linseed, 0 02 to 0 038 p c; Nandina domestica, 0 07 to 0 147 p c, bitter almonds 0 12 to 0 18 p c Cases of cattle poisoning are not infrequent in hot countries where sorghum is used as fodder, and linseed meal and white clover are also stance there is a certain temperature at which sometimes poisonous. The enzymes are detection the yield is a maximum. The following table

stroyed and the cyanogenetic glucosides rendered harmless by thorough boiling with water.

Minute traces of hydrocyanic acid have been observed in the products of incomplete combustion, for example, in the luminous flame of a Bunsen burner, in boiler flue gases, and in tobacco smoke

Preparation of Hydrocyanic Acid -Hydrocyanic acid was formerly prepared by distilling potassium ferrocyanide with dilute sulphuric A more convenient method, however, is to run a solution of sodium cyanide (97 to 98 pc NaCN) in from 2-21 times its weight of water into hot sulphuric acid of 75 p c. strength, about 11 mols of sulphuric acid to 2 mols sodium evanide is sufficient and the eyanide solution is preferably allowed to enter the acid a little below its surface, so as to avoid contact of the alkaline cyanide solution with hydrocyanic acid vapour.

The vapour may be dried by calcium chloride, kept at a temperature of about 30° and then condensed in a vessel cooled by ice, although the greater part of it is quite readily liquefied in an ordinary Liebig's condenser cooled by cold water. It may be purified from traces of water and ammonia by distillation over phosphorus pentoxide (Nef, Annalen, 1895, 287, 265)

Hydrocyanic acid is formed in many different ways, the synthesis from free nitrogen and acetylene has already been The oxidation of organic compounds by nitric acid often yields hydrocyanic acid in greater or smaller amount, and it has been proposed to manufacture hydrocyanic acid in this way The interesting reaction described by Huntingdon (Eng Pat 14855, 1895) may be mentioned A mixture of acetylene and nitric oxide is readily exploded by a spark, yielding very considerable quantities of hydrocyanic acid in accordance with the equation

$$2NO+2C,H_2=2HCN+2CO+H_2$$

When ammonium formate is distilled, water is first given off together with a little ammonia and formic acid, at 180° formamide begins to distil over, and, as the temperature gradually rises to 220°, this is accompanied by increasing quantities of carbon monoxide, but only traces of hydrocyanic acid are produced. In order to obtain a good yield of hydrocyanic acid, the vapour must be heated to much higher temperatures, and the result is greatly influenced by the nature of the substance with which it is in contact According to unpublished experiments by the writer, the principal changes which occur

$$\begin{array}{l} \text{H COONH}_4 = \text{HCO·NH}_2 + \text{H}_2\text{O (1)} \\ \text{H COONH}_4 = \text{NH}_3 + \text{HCOOH} \\ = \text{NH}_3 + \text{H}_2\text{O} + \text{CO (2)} \\ \text{HCO NH}_2 = \text{HCN} + \text{H}_2\text{O} \\ \text{H}_5\text{O} + \text{CO} = \text{CO}_5 + \text{H}_3 \end{array}$$

The ammonia and hydrocyanic acid in the product usually accounted for more than 95 p.c. of the nitrogen in the ammonium formate or formanide used. Under similar conditions, the yield of hydrocyanic acid obtained from formamide is much greater than that obtained from ammonium formate. For each contact sub-

contains the best temperatures and corresponding yields for several contact substances:—

	Amn	nonium formate	Formamide	
Contact, substance	Yield of HCN p c. of theoretical	Temperature	Yield of HCN p c of theoretical	Temperature
Porcelain chips . Copper gauze Iron wire . Charcoal .	41.5 43.0 39.0 21.0	700° just below 500° just below 500° 420°	75 78 — 55	800° 600° - 400°

With finely divided mickel, no hydrocyanic acid was obtained in any circumstances; at 273°, 92½ p.c. of the mitrogen was recovered as ammonia, but much less at higher temperatures

Formaldoxime CH<sub>2</sub> N OH, an isomeride of formamide, obtained by the interaction of formaldehyde and hydroxylamine, decomposes very readily when heated alone or with phosphorus pentoxide into water and hydrocyanic acid (Dunstan and Bossi, Chem. Soc. Trans 1898, 353).

Ammonia and carbon monoxide were said to yield hydrocyanic when passed over heated spongy platinum (Kuhlmann, Annalen, 1841, 38, 62), CO+NH<sub>3</sub>=HCN+H<sub>2</sub>O, and this apparently simple reaction has attracted several inventors. A careful investigation by Jackson and Laurie (Chem. Soc. Trans. 1905, 87, 433) has shown that a dry mixture of the two gases, when sparked, treated by the silent discharge or heated in presence of platinum (an electrically heated spiral of platinum wire gives the best result) yields ammonium cyanate thus:

and Kuhlmann's reaction does not take place. The reverse change,  $H(N+H_2O=CO+NH_3)$ , however, does occur very readily, even at  $200^\circ$ , when a mixture of steam and hydrocyanic acid is passed over broken brick as contact material, and at  $600^\circ$  it is practically quantitative. With iron oxide as contact substance, the conversion into ammonia is complete at  $285^\circ$ , and with Weldon mud at  $200^\circ$  (Carpenter and Linder, J. Soc. Chem. Ind. 1905, 63).

Cyanogen and hydrogen combine, forming hydrocyanic acid when a mixture of them is heated at 500°-550° (Beithelot, Ann. Chim.

Phys. 1879, [5] 18, 380).

Properties of Hydrocyanic Acid.—A colourless liquid possessing a characteristic smell (or rather taste), which produces a choking sensation in the larynx. It is one of the most poisonous substances known, the fatal dose for a full-grown man being about 0.06 gram of the acid or 0.15 gram of potassium cyanide. It may be absorbed through all parts of the body, even through the skin, and its action is very rapid, large doses causing death almost instantaneously. Air in which the smell of hydrocyanic acid is distinctly perceptible may be breathed constantly without ill effects. Experiments with cats showed that 3 to 4 volumes of the acid per 100,000 volumes of air could be borne, but 5 volumes

produced symptoms of poisoning In industrial cases of hydrocyanic acid poisoning the persons affected were usually working in continued spaces, for example, cleaning or repairing apparatus which had contained cyanides, removing damaged eargo from the holds of ships, or working in insufficiently ventilated buildings where cyanide solutions were used; in all these cases hydrocyanic acid is generated from alkali cyanide solutions partly by hydrolysis and partly by the action of carbon dioxide.

The first symptoms of poisoning are giddiness and darkness of the visual field (these symptoms disappear after a few minutes exposure to fresh air), they are followed by un-consciousness accompanied by muscular cramps and more or less complete cessation of respiration; the heart still acting, however, artificial respiration with oxygen may be successful at this stage. When a cyanide has been swallowed, hydrocyanic acid is liberated by the acids of the stomach. The simplest and most effective antidote is prepared by mixing solutions of ferrous sulphate and sodium carbonate (suitable quantities are 50 c c each of solutions containing 150 grams of FeSO<sub>4</sub>,7H<sub>2</sub>O and 57 grams of Na<sub>2</sub>CO
<sub>3</sub> per litie) Stimulants, such as ammonia, are also helpful The writer has seen a dog, which had swallowed some potassium cyanide, and was already partially paralysed in its hind quarters, recover in a few seconds when dosed with the above mixture which had been made up hastily from ferrous ammonium sulphate. Subcutaneous injection of hydrogen peroxide, which converts hydrocyanic acid into oxamide, has also been recommended

The action of hydrocyanic acid appears to consist partly of a direct effect on the nerve centres and partly of an inhibitory action on the enzyme of the red blood corpuscles, which brings about the transference of oxygen from oxyhaemoglobin to the oxidisable substances oxynacmognomic of the Galactic name of the body, the venous blood therefore becomes arterial, and death is due partly to a kind of internal suffocation Schonbein (J pr. 1868, 105, 202) showed that hydrocyanic acid inhibits the catalytic decomposition of hydrogen peroxide by plant ferments, and also by the red blood corpuscles, and suggested that its poisonous properties might be due to the inhibition of the oxidising power of the blood. Bredig and von Berneck (Zeitsch phys. Chem 1899, 31, 329) showed that hydrocyanic acid has the same effect on the inorganic ferment, colloidal platinum The ferments recover their activity when the hydrocyanic acid is removed. This property has been applied in the pieservation of fruit

The sp gr of the liquid acid is 0 7058 at 7° and 0 6969 at 18° (Gay Lussac) Freezing-point, -10° to -12° (Nef). The freezing-point is raised by pressure; it is -2 88° under 500 kilos per sq. cm. pressure (Tamman) The boiling-point at atmospheric pressure is 25° (Nef) The heat of evaporation is exceptionally large, 210 7 Cals. per gram. Heat of formation of the liquid acid from the elements (amorphous carbon) -218 Cals. per moi. (Thomsen), diamond -24 8 Cals. (Berthelot). Heat of solution, liquid acid in water, 0.4 Cal. per moi. The dielectric constant of hydrocyanic acid, 95, is the largest known,

with the exception of that of hydrogen peroxide (Schlundt). The acid is miscible in all proportions with water, alcohol, ether and many other substances; it is almost insoluble in light petroleum. It is an excellent solvent for many solid substances, among the inorganic salts the iodides are usually readily soluble. The solutions are good conductors of electricity, the conductivity of solutions of potassium iodide, for example, is about 31 times greater than that of the corresponding aqueous solutions

Hydrocyanic acid is one of the weakest acids; its dissociation constant in aqueous solution at 18°, 13×10<sup>-9</sup>, is 44 times smaller than that of sulphuretted hydrogen, and 230 times smaller than that of carbonic acid (Walker and Cormack, Chem Soc Trans 77, 16, 1900) In a 01N-solution, only 0011 pc. of the acid is dissociated into its ions, consequently it does not behave as an acid to indicators, and its salts are more or less completely decomposed by water; thus for example,

# KCN+H<sub>2</sub>O ≥ KOH+HCN

In 0 l N-solutions of the sodium or potassium salts, 1 1 p c is dissociated in this way, in N-solutions, only 0 3 p c at 24 2° (Shields, Zeitsch physikal. Chem. 12, 167, 1893). In solutions of pure sodium eyamide, containing no caustic soda, the hydrolysis is greater than that found by Shields. It may be calculated from the equilibrium constant K == [HaOH] HCN | NaCN |. the bracketed quantities representing concentrations of the substances in gram mols per htre The values of  $K \times 10^4$  are. at 0°, 0 135, at 5°, 0 15, at 10°, 0 16; at 15°, 0 20, at 20°, 0 27, at 25°, 0 484; and at 30°, 0 72 (F P. Worley and V R. Browne, Chem Soc Trans 1917, 1057) This is of importance it is required to concentrate an aqueous solution of a cyanide, the free hydrocyanic acid escapes until sufficient excess of the base has accumulated to reduce the hydrolysis to negligible proportions In the case of weaker bases, the result is a considerable loss of hydrocyanic acid; with calcium cyanide, for example, the quantity of lime formed is greater than can be retained in solution, consequently equilibrium is never attained, and when such a solution is evaporated, the whole of the hydrocyanic acid escapes, leaving a residue of pure lime

A large part (sometimes one-half) of the cyanide used in gold extracting is lost by the evaporation from the solutions of hydrocyanic acid set free by hydrolysis (H A White, J Chem Met and Min Soc. S. Africa, 1915, 16, 24)

Pure hydrocyanic acid, distilled over phos phorus pentoxide, can be preserved without change for considerable periods at the ordinary temperature; when heated to 100° or at the ordinary temperature in presence of alkaline impurities, such as ammonium or potassium cyanide, it changes into a solid, black mass from which a ter-molecular polymer, probably the nitrile of amino malonic acid, NH<sub>2</sub> CH(CN)<sub>2</sub>, can be separated Under imperfectly known conditions, the anhydrous acid cometimes explodes. G Salomone (Gazz. chim. ital Salomone (Gazz. chim. ital explodes. 1912, 42, 1 617) isolated, from the products of an explosion, colourless crystals of a substance having the composition of cyanic acid and a cyanic acid also resembles the nitriles rather

molecular weight, in benzene solution, corre-

sponding with (HUNO)<sub>5</sub>.

The aqueous acid also becomes brown rapidly in presence of alkali cyanides and deposits a dark brown amorphous substance, azulmın or azulmıc acıd, which contains carbon, oxygen, nitrogen, and hydrogen in variable proportions. The discolouration of solid alkalı

cyanide, when exposed to the air, is due to the hydrocyanic acid liberated at its surface under-

going this change

The only useful application of hydrocyanic acid, as an insecticide, was developed at the Agricultural Experiment Station of the University of California, largely by the labours of D W Coquillett and H D Bishop (California Station Bull. 122) They found that foliage is much less liable to injury at night than in sunlight, the hydrocyanic acid vapour is applied by electing a tent over the trees and generating the acid inside of it The safe concentration of hydrocyanic acid depends on the kind of tree and on the time of year In greenhouses less than 2 grams of hydrocyanic acid per 1000 cb. ft. of air space is said to kill green aphis, but as much as 30 grams may be needed for mealy bug or red spider Most plants are unaffected even by the larger quantity Since about 1918 98 p.c hydrocyanic acid has been placed on the market for the fumigation of orange trees; formerly the acid was always generated on the spot, from sodium cyanide and sulphune acid

The constitution of hydrocyanic acid has been the subject of much discussion arising largely from the observation that the reaction of potassium cyanide with an ester yields a nitrile, R-C: N, as principal product, whereas silver evanude yields a carbylamine, R-N C. On the hypothesis that the salts are formed by the substitution of an atom of metal for an atom of hydrogen without change of position, and that the alkyl derivatives are formed similarly, it is of course necessary to suppose that the hydrogen in hydrocyanic acid may be united either with carbon or with nitrogen. The cyanides are, however, in all probability more complex substances; silver cyanide, from its mode of formation, is probably silver dicyano argentaate, Ag[Ag(CN)2], and the reaction with methyl iodide is similar to that of silver ferrocyanide (described above) leading to the well-crystallised

substance  $\left[\mathrm{Ag}^{\mathrm{CN}}_{\mathrm{CNCH}_{3}}\right]$ , which, like the corre-

sponding ferrocyanide, splits off methyl carbylamine when heated or when acted on by a solution of potassium cyanide (see, for example, H. Guillemaid, Ann. Chim Phys 1908, [8] 14, 311)

The properties of hydrocyanic acid itself, however, leave little doubt that it is formonitrile; the isomeric substance H.N.C, has not been isolated. Like the nitriles, hydrocyanic acid is hardly affected by dilute mineral acids (it is frequently separated quantitatively by distilling acid solutions), the carbylamines are violently decomposed Anhydrous hydrocyanic acid and the nitriles do not dissolve silver cyanide, whilst the carbylamines dissolve it very readily its high dielectric constant, solvent and dissociating powers and physiological action hydro-

than the carbylamines (Wade, Chem. Soc.) Trans. 1902, 1613).

# THE PRINCIPAL CYANIDES.

Potassium cyanide KCN crystallises in anhydrous, colourless octahedra from aqueous solutions, or in cubes when the fused salt solidi fies; sp gr. 152 It melts at a dull red heat, and volatilises without decomposition at high temperatures It is readily soluble in water, the saturated solution contains 122 2 grams of KCN to 100 grams of water at its boiling-point (103 3°). At 19.5°, 100 grams of absolute alcohol dissolve 0 88 gram, 100 gr methylalcohol dissolve 4.91 grams (Landolt, Bornstein and Meyerhoffer, Tabellen, Berlin, 1905). It is moderately soluble in anhydrous ammonia at -38° (Fianklin and Kraus, Amer. Chem. J 1898, 20, 820) It deliquesces in moist air

Sodium cyanide NaCN crystallises from the fused liquid in apparently cubic crystals, from aqueous solutions below 35° large colourless plates of the hydrate NaCN,2H<sub>2</sub>O are deposited Joannis (Ann Chim Phys. 1882, [5] 26, 484) obtained a hydrate NaCN, H2O from cold 85 p.c. alcohol F Roessler (Eng. Pat. 279, 1899) showed that the hydrate NaCN,2H<sub>2</sub>O decomposes at 33° (34.7° according to the writer's measurements) into the anhydrous salt and a saturated solution, about 40 pc of the sodium cyanide separating in the anhydrous The solubility of the hydrate (in grams of NaCN to 100 grams of water) is, 35 4 at  $-20^{\circ}$ , 43 4 at  $0^{\circ}$ ; 58 3 at 20° and 82 at 34 7°. The solubility of the anhydrous salt is almost independent of the temperature (82 grams per 100 grams of water) Sodium cyanide appears to be more soluble in ethyl and methyl alcohol and anhydrous ammonia than the potassium salt The sp gr of aqueous solutions of sodium cyanide at 25° is given by the expression 1+0.00534p, where p is the percentage of NaCN Sodium cyanide (96 p c) melts at 540° The pure salt, when heated in a glass tube so as to be out of contact with metals, in a current of dry air free from carbon dioxide, remains entirely unchanged at temperatures up to 370°; if, however, it is in contact with iron or nickel, or if traces of the oxides of these metals are added to it, oxidation proceeds rapidly, cyanate and carbonate being formed and carbon dioxide evolved The carbonate is formed by the oxidation of the cyanate, which is the primary product Sodium cyanate behaves towards oxygen in exactly the same way as the cyanide, it is quite unaffected by dig air (free from CO2) even at 400°, but when mixed with 1 p c. of nickel oxide it oxidises rapidly, thus.

$$2NaCNO+3O=Na_2CO_2+CO_2+N_2$$

(writer's unpublished experiments)

Sodium cyanide absorbs moisture with great avidity from ordinary moist air The writer has found that it takes up water from purified air saturated with aqueous vapour at 20° at all temperatures below 44°, but not at higher temperatures. This is of importance in sampling and analysing the salt

Dilute solutions of potassium or sodium cyanide decompose slowly in confact with the air, the loss being due to evaporation of the hydrocyanic acid formed by hydrolysis, which

permits the decomposition to proceed further; absorption of carbon dioxide acts in the same direction by neutralising the caustic alkali

Liebig's statement that metallic non is dissolved by hot solutions of potassium or sodium cyanide with evolution of hydrogen is true of the finely divided metal, but iron vessels may be used for years for boiling concentrated solutions of cyanides without requiring repair, and iron tanks are regularly used in the treatment of gold ores with cyanide solutions

With the exception of lead and platinum, all ordinary metals are dissolved by cyanide solu-

tions in presence of oxygen

Zinc is slowly dissolved without oxygen, but if a piece of zinc is half immersed in a cyanide solution, it is eaten through at the surface of the liquid before any appreciable action has taken place on the submerged portion

Alkalı cyanıdes yıeld ammonia and a formate when boiled with water or treated with strong sulphuric acid, in the latter case the formic acid produced is at once decomposed

into carbon monoxide and water

Mercuric cyanide Hg(('N)2 was first prepared by Scheele, in 1782, by boiling Prussian blue with mercuire oxide, and, in 1783, by dissolving mercuric oxide in hydrocyanic and and evapo-rating to crystallisation. He also found that mercurous salts, when treated with soluble cyanides, yield mercuric cyanide and metallic mercury. The decomposition of the ferromercury The decomposition of the ferro-cyanides, as already mentioned, depends on the extremely small electrolytic dissociation of mercuric cyanide in aqueous solution For the same reason, it is not decomposed by caustic alkalis, and hydrocyanic acid can only be expelled from it by boiling with a strong acid The concentration of the mercury ions is, however, sufficient to permit of the precipitation of the extremely insoluble mercuric sulphide by sulphuretted hydrogen

Mercuric cyanide crystallises in anhydrous, colourless crystals of sp gr. 40 The saturated solution contains 93 grams per litre at 20°, 111 grams per litre at 25°, 100 grams of absolute alcohol dissolve 10 l grams of it at 195°, and 100 grams of absolute methyl alcohol take up

44 2 grams at 19 5°

Small quantities of mercuric eyanide are sometimes used in addition to alkali cyanide in the treatment of ores containing silver sulphide (Clennel, Cyanide Handbook, London, 1910, 297), the dissolution of which is accelerated, probably owing to the formation of mercuric sulphide:

 $Hg(CN)_2 + Ag_2S + 2KCN = HgS + 2KAg(CN)_2$ 

Calcium cyanide has never been prepared in the pure state An aqueous solution of lime in hydrocyanic acid decomposes completely into its constituents when boiled (Scheele) By evaporating such a solution in vacuô, Joannis obtained a crystalline compound, 3CaO Ca(CN)2,15H2O, which decomposed completely in a vacuum, leaving nothing but lime. Dry reactions, at high temperatures, yield calcium cyanamide, but no cyanide •

Barium cyanide Ba(CN)<sub>2</sub> When a concentrated solution, made by adding hydrocyanic acid to baryta in theoretical quantities, is evaporated in vacuô, crystals of the hydrate

Ba(CN)<sub>2</sub>,2H<sub>3</sub>O are deposited (A Joannis, Ann Chim. Phys. [5] 26, 484, 1882). This is a very deliquescent substance which, when dried *in* vacuó over sulphuric acid, yields an amorphous monohydrate which can be completely dehydrated by heating it very gradually to 100° in a good vacuum. The writer has failed to obtain a pure product in this way Some 7 or 8 p c of the hydrocyanic acid escapes during evaporation, and the product contains a corresponding quan-By drying pure tity of barium hydroxide Ba(OH)<sub>2</sub>,8H<sub>2</sub>O in a good vacuum at 100°, a very finely divided powder of Ba(OH)<sub>2</sub> is obtained which is suspended in dry light petroleum and treated with a small excess of dry hydrocyanic acid also mixed with light petroleum. In this way, the solid hydrated barium cyanide is obtained without evaporation:

$$Ba(OH)_2 + 2HCN = Ba(CN)_2 2H_2O$$

When dehydrated in vacuô, it still contains a little hydroxide, but by shaking it with a little hydrocyanic acid and light petroleum and again drying, practically pure anhydrous barium cyanide is fairly easily obtained

By saturating a solution of baryta in methyl alcohol with hydrocyanic acid, a compound Ba(CN)(OCH<sub>3</sub>) + CH<sub>3</sub>OH is obtained, which, when heated, first loses methyl alcohol and then methyl ether, leaving BaO + Ba(CN), (Diechsel, J. pr. Chem [2] 21, 77, 1880).

Dry methods of preparation yield mixtures of cyanide and cyanamide When barrum terrocyanide is heated it begins to decompose about 500°, yielding a mixture of barium evanide and evanamide (with carbon and iron) in which nearly 60 p c. of the nitrogen is in the form of cyanamide, practically the same product as that obtained by the action of nitrogen on barium carbide Even the action of cyanogen on barium amalgam yields a mixture (Drechsel), the reason being that barium eyanide decomposes into barium cyanamide and carbon, thus . Ba(CN)<sub>2</sub>=BaCN<sub>2</sub>+C. The change is very slow at 500° to 600°, but it is accelerated by iron (Ewan and Napier, J Soc Chem Ind 1913, 32, 467)

Barium eyanide is colourless, 100 grams of cold water dissolve about 80 grams of it (Joannis) It melts about 600° It is distinctly volatile, even at its melting-point, but when mixed with

barium oxide it is much less so.

The precipitate ob-Silver cyanide AgCN tained by adding excess of silver nitrate to a solution of a soluble cyanide is probably the silver salt of the complex acid  $H[Ag(CN)_2]$ The solubility of silver cyanide in pure water lies between those of the chloride and sulphocyanide, as the following table shows

> Solubility at 25° in gram-mols per litre AgI  $0.01 \times 10^{-6}$  $0.81 \times 10^{-6}$ AgBr AgSCN  $1.08 \times 10^{-6}$ 4 40  $\times$  10-6 AgCN AğCl 14 10×10-6

It is hardly attacked by cold dilute natric acid, but is decomposed quantitatively on boiling (Plimmer, Chem Soc Proc. 1903, 285) When

cyanogen C<sub>2</sub>N<sub>2</sub>. Further heating in presence of oxygen yields a residue of metallic silver, which usually contains a little carbon. It is slightly soluble in a solution of hydrocyanic acid, forming the unstable acid HAg(CN)2. It dissolves very readily in alkali cyanides to salts such as KAg(CN), in which the silver is contained in the complex amon Ag(CN)'<sub>2</sub>. Owing to the excessively small dissociation of this anion, all silver salts except the sulphide are dissolved by cyanide solutions, and in presence of oxygen, metallic silver and the sulphide are also dissolved.

Gold cyanides. Aurous cyanide and the soluble aurocyanides are of technical importance. The former, AuCN, a yellow crystalline powder, insoluble in water and acids, is formed by the action of hydrocyanic acid on auric hydroxide. It is readily soluble in solutions of alkali or alkaline-earth cyanides, yielding aurocyanides such as KAu(CN)2. solution of the potassium salt, which is used for electroplating with gold, is readily obtained by precipitating a solution of gold chloride with ammonia and dissolving the fulminating gold in a solution of potassium cyanide, or by precipitating gold chloride with magnesia, dissolving out the excess of magnesia in the precipitate with nitric acid, and then dissolving the auric hydroxide in potassium cyanide as before

Potassium aurocyanide is also formed when gold is dissolved in a solution of potassium cyanide in presence of oxygen or other oxidising

agent:

 $4KCN+2Au+O+H_2O=2KAu(CN)_2+2KOH$ 

(Clennel, Cyanide Handbook, 1910, 10-32 gives a full account of the discovery of this reaction.)

The dissolution of metallic gold is primarily due to the great stability of the aurocyanogen ion Au(CN),'. Metals, like all other substances, have a tendency to dissolve in water. In contact with its aqueous solution, a salt will dissolve if the solution is unsaturated, or crystallise out if it is supersaturated Metals behave sumilarly, with the important difference that they dissolve in the form of positively charged ions, so that when a metal is placed in contact with a solution containing fewer of its ions than corre spond with equilibrium, it gives out positively charged atoms, becoming itself negatively charged, until the electrical attraction stops further change, and vice versa. In this way, a definite potential difference always arises between a metal and a solution of it, the sign and magnitude of which depend on the nature of the metal and on the concentration of the positively charged ions of the metal in the solution

In order for a metal to dissolve at all, therefore. it is necessary that the solution shall contain less than the saturation quantity of positive ions; and for the dissolution to continue, the positive charges of the ions and the negative charge of the metal must be got rid of in some way.

These conditions are fulfilled in the case of the dissolution of gold in a cyanide solution containing oxygen, because the dissociation of the complex ion,  $Au(CN)'_2 \gtrsim Au' + 2CN'$  is extremely small, and solutions of it therefore dry silver cyanide is heated about one-half contain infinitesimal quantities of gold ion, and of the CN is given off in the form of further the electrical charges produced by the

dissolution of the gold are removed by the reactions:

and Au'+2(CN)'=Au(CN)' $O+H_2O+2(-)=2OH'$ 

the result of which, written in the ionic form, is  $4K'+4(CN)'+2Au'+O+H_2O+2(-)$  =  $4K'+2Au(CN)_2'+2OH'$ 

The potential difference between a metal like zinc (which has a greater tendency to dissolve than gold) and a cyanide solution is much greater than the natural potential difference between gold and the solution; hence gold ions are drawn out of the solutions to the zinc, where their positive charges are neutralised by the negative charge of the zinc. This disturbs the equilibrium between zinc and the solution and also between aurocyanogen ions and gold ions, causing fresh quantities of zinc to dissolve and of aurocyanogen ions to dissociate, processes which continue until the whole of the gold is precipitated.

As W. Skey (Trans. and Proc. New Zealand Inst. 8, 334, 1876) first showed, gold in cyanide solutions is electro-positive to silver and silver to mercury, so that gold will precipitate silver and mercury from their solutions in cyanides. This has been confirmed by Bodlander (Ber 1903, 36, 3933). This reversal of the usual order is, as will be understood from the above explanation, due to the increasing stability of the mercury, silver, and gold complex cyanogen

ions.

It is of interest to note that most of the minerals which accompany gold in its ores are electro-negative to gold in cyanide solutions, and therefore accelerate its dissolution in the way which is familiar in the solution of zinc in sulphuric acid in presence of copper or platinum

Barium platinocyanide BaPt(CN)<sub>4</sub> 4H<sub>2</sub>O is

Barium platinocyanide BaPt(CN)<sub>4</sub> 4H<sub>2</sub>O is most readily made by electrolysing a solution of barium cyanide (270 grams per litre) with platinum electrodes and an alternating current (0.4 ampere per sq. cm.), about 0.6 gram platinum is dissolved per ampere hour. After removing the excess of barium by CO<sub>2</sub> the solution is concentrated to crystallisation (A Brochet and J Petit, Bull Soc chim. 1904, [3] 31, 630) The salt crystallises in monoclinic prisms, which are yellow by transmitted, violetblue by reflected, light—It dissolves in 33 parts of water at 16°. By recrystallisation from a solution containing barium cyanide it is obtained in small crystals which are highly fluorescent when exposed to X-rays, and are suitable for making fluorescent screens for X-ray work

Copper cyanides. When potassium cyanide is added to the solution of a cupric salt, a brownish precipitate of cupric cyanide Cu(CN), is produced, which gradually evolves cyanide. When heated, this loses a further quantity of cyanogen and yields white cuprous cyanide CuCN, the constitution and reactions of which appear to resemble those of silver cyanide.

Cuprous cyanide is soluble in solutions of potassium cyanide, about 2½ molecules of KCN being required to dissolve 1 molecule of CuCN; the solutions contain the complex anions Cu(CN)," and Cu(CN)," (Kunschert, Zeitsch. anorg Chem. 41, 359, 1904) These anions are so little dissociated into copper ions that sul-

phuretted hydrogen will not produce a precipitate in the solutions, and therefore copper itself (in presence of oxygen) and all its compounds dissolve readily in cyanide solutions. Solutions of the sodium or potassium cupro-cyanides are used in the electro-deposition of copper on iron.

469

Zinc cyanide Zn(CN)<sub>2</sub> is a white substance, insoluble in water, but soluble in acids and in cyanide solutions. It is not easily prepared in a perfectly pure condition, because water decomposes it partially. When dry it is very stable, decomposing only at a bright-red heat.

The complex salt potassium zinc cyanide  $K_2Zn(CN)_4$  crystallises in large well-formed octahedra from aqueous solutions; 100 grams of water at 20° dissolve 11 grams of it. It is anhydrous, and may be fused at a red heat without decomposition. The sodium salt  $NaZn(CN)_3, 2\frac{1}{2}H_2O$  is very readily soluble in water. The complex zinc cyanogen ions are much less stable than those containing copper, gold, silver, and mercury; a small excess of an alkali sulphide precipitates the zinc completely.

Cyanogen (Ger Cyan) C<sub>2</sub>N<sub>2</sub> was discovered by Gay-Lussac It is prepared by heating perfectly dry mercuric cyanide at a dull red heat, Hg(CN)<sub>2</sub>=Hg+C<sub>2</sub>N<sub>2</sub> A part of the cyanogen is always polymerised to dark-brown, non-volatile paracyanogen. Paracyanogen appears to be the stable form at ordinary temperatures, at high temperatures it is converted into cyanogen. The velocity of the change is very small at temperatures below 300°, so that cyanogen gas can exist in the cold without change.

Cyanogen itself is a gas which burns with a peach-blossom-coloured flame Liquid cyanogen boils at  $-20.7^{\circ}$  under atmospheric pressure and freezes at -34 4° Its critical temperature and pressure are 128 3° and 59 6 atmos respectively (Cardoso and Baume, Compt rend 1910, 151, 141). Its heat of formation from graphite and nitrogen is 70,000 cals. per molecule Water dissolves 4 95 times its volume of the gas at 0°. At 18° the dissolved cyanogen is hydrolysed to hydrocyanic and cyanic acids, the latter yielding ammonia and carbon doxide so that no definite solubility is observed (Naumann, Zeitsch Elektrochem. 16, 772, 1910) After some time the solution deposits dark brown azulmic acid and contains also oxalic acid and urea dissolves in caustic alkalis, yielding cyanide and eyanate:  $2KOH + C_2N_2 = KCNO + KCN + H_2O$ . Silver nitrate solution has no action on it. By means of these two reactions, it may be separated from hydrocyanic acid and estimated quantitatively in mixtures of the two gases (Wallis, Anna-

len, 345, 353, 1906) Cyanogen is very poisonous.

A knowledge of the exact conditions under which cyanogen can be formed from its elements is of interest in the theory of several technical processes for the manufacture of cyanides

Wallis loc cit has shown that the electric arc burning between electrodes of very pure carbon in an atmosphere of nitrogen produces no cyanogen at all H. v. Wartenberg (Zeitsch anorg. Chem 52, 299, 1907) pointed out, however, that a very intense cyanogen spectrum is seen, and that carbon is deposited at some distance from the arc. Nernst's theorem indicates that very considerable

quantities of cyanogen should exist in equilibrium with carbon and nitrogen at the temperature of the arc:

Temperature	Per cent C <sub>2</sub> N <sub>2</sub> in the gas calculated from Nernst's theorem
2000°	0.01
2500°	0 9
3000°	17 0
39500	45.0

The probable explanation is that evanogen is essentially unstable at low temperatures, that it can exist at all is due to an excessively slow rate of decomposition. Berthelot (Force des Matières Explosives, Paris, 1883, 113) has Matières Explosives, Paris, 1883, shown that it can be detonated by mercury fulminate, and, when heated, its rate of de-1200°. composition becomes measurable at v Wartenberg supposes, therefore, that cyanogen is really formed at the high temperature, but that it decomposes again completely before it can cool to temperatures at which it can continue to exist. It is obvious from the figures given above that the direct union of carbon and nitiogen to evanogen can play no part in the formation of cyanides in processes which are carried out at ordinary furnace temperatures

Cyanogen chloride ('NCl is formed by the action of chlorine on mercuric cyanide (Serullas, Ann Chim. Phys 1827, 35, 291), zinc cyanide (A Held, Bull Soc chim 1897, [3] 17, 287), or aqueous hydrocyanic acid (Berthollet, 1787; A Wurtz, Annalen, 1847, 64, 367) Alkali cyanides are unsuitable, because, except in cold, very dilute solutions, they react violently with cyanogen chloride, yielding dark-brown, amorphous products. The preparation of cyanoamorphous products gen chloride in considerable quantities is described by T S Price and S J Green (J Soc Chem Ind 1920, 39) A 12 to 15 pc solution of pure hydrocyanic acid is treated with chlorine at about 40°, the slow absorption of chlorine is hastened by injecting it into the solution of a finely divided condition through porous carthenware. The cyanogen chloride distils off as it is formed, together with hydrocyanic acid, the product containing 10-20 p c. of the latter It is purified by retreatment with chlorine, or by shaking it with zinc oxide, which removes hydrocyanic acid almost completely Hydrochloric acid is removed by distillation over sodium bicarbonate

Cyanogen chloride is a colourless liquid, which boils at 12 66°, 760 mm, and solidifies at Its vapour pressure (mm) is  $-30^{\circ}$ ;  $148.2^{\circ}$ at  $-20^{\circ}$ ;  $270.5^{\circ}$  at  $-10^{\circ}$ ;  $441.1^{\circ}$ at 0°, 681 9 at 10°; 1001 9 at 20°; 1427 4 at 30°, 1988 0 at 40°, 2719 3 at 50°; 3664 2 at 60; 4873 2 at 70° (Regnault). The vapour attacks the eyes violently The cold, saturated aqueous solution contains about 6 pc CNCl. The heat of formation of liquid CNCl from its elements is -26.9 cals, per molecule. Dry cyanogen chloride may be preserved indefinitely when pure; in presence of hydrochloric acid it polymerises to cyanuric chloride C3N3Cl3. Aqueous hydrochloric acid (stronger than about 2N) hydrolyses it thus.

 $CNCl + 2H_2O = NH_4Cl + CO_2$ 

Its behaviour towards alkalıs is the same as that of cyanogen bromide described below. Hydriodic acid hardly attacks it in the cold, and it is without action on metals, such as iron and copper, at ordinary temperatures.

Cyanogen bromide CNBr. A solution containing cyanogen bromide in addition to a cyanide dissolves gold more rapidly than one containing cyanide alone (Sulman and Teed, Eng Pat 18592, of 1894, and J Soc Chem. Ind 1897, 961) Cyanogen bromide is therefore used in the treatment of a few refractory ores, the West Australian tellurides especially The final result of the reaction is expressed by the equation:  $2Au + 3KCN + BrCN = 2KAu(\tilde{C}N)_2 + \hat{K}Br$ 

Cyanogen bromide is formed by the action of bromine on a cyanide or on hydrocyanic acid. Owing to the difficulty of transporting either bromine or cyanogen bromide, it is always prepared at the place where it is to be used by liberating bromine from a mixture of a bromide and a suitable oxidising agent by the addition of sulphuric acid, and then adding a concentrated solution of an alkali cyanide. By the use of a suitable quantity of the oxidising agent, the whole of the bromine is utilised, with sodium bromate, for example, the reaction is (C Gopner, Zeitsch angew Chem 1901, 14, 355):

 $2NaBr + NaBrO_3 + 3NaCN + 3H_2SO_3$ 

 $=3Na_2SO_4+3BrCN+3H_2O$ 

Cyanogen bromide is a colourless, crystalline substance, which melts at 52° and boils at 61°; 100 grams of water dissolve from 5 to 6 grams of it at the ordinary temperature. It has an intensely irritating effect on the mucous membranes and eyes, and is said to be very poisonous The perfectly pure substance can be kept without change, but traces of bromine (or hydrobromic acid) cause it to polymerise rapidly to cyanuric bromide, partial decomposition always occurring It is very slowly decomposed by water: CNBr+H<sub>2</sub>O=HCNO+HBr, the cyanic acid decomposing further to ammonia and carbon dioxide Caustic alkalis bring about the same change instantaneously, alkali carbonates much more slowly, and bicarbonates have no action at all Cyanogen bromide and potassium cyanide react in aqueous solution, yielding cyanogen; thus:

CNBr+KCN=KBr+C<sub>2</sub>N<sub>2</sub> (private communication from Mr C. J Ellis). In solutions containing 0.2 to 08 pc of potassium cyanide and 01 to 02 pc of cyanogen bromide, the latter is completely decomposed in from 3 to 6 hours at the ordinary temperature

In aqueous solution cyanogen bromide reacts with hydriodic acid and with sulphuretted hydrogen according to the equations

$$\begin{array}{c} \text{CNBr+2HI=HCN+HBr+I}_2\\ \text{nd} \quad \text{CNBr+H}_2\text{S=HCN+HBr+S} \end{array}$$

The first reaction provides a convenient method for the quantitative estimation of cyanogen bromide. Chattaway and Wadmore (Chem. Soc Trans 1902, 191) have shown that these reactions are characteristic of compounds in which the halogen atom is directly linked with nitrogen, and that cyanogen bromide therefore has the constitution C. N Br.

Cyanamide NH<sub>2</sub> CN was first prepared by the action of ammonia on an ethereal solution of cyanogen chloride (Cloez and Cannizzaro, Compt rend 1851, 32, 62); ammonium chloride crystallises out and cyanamide remains when the ether is evaporated. The same reaction

occurs in aqueous solution. The formation of the disodium salt,  $Na_2CN_2$ , by the action of sodamide on carbon or on sodium cyanide has already been described The same substance is also produced when sodamide acts on sodium cyanate, sodium carbonate or carbon dioxide at a low red heat; it is also formed when either sodium cyanate or sodium cyanide is fused with sodium hydroxide.

The calcium and barium salts, CaCN, and BaCN<sub>2</sub>, may be prepared from nitrogen and the carbides, by heating the corresponding ferrocyanide or by heating the carbamates or cyanates to redness (Drechsel, J pr. Chem. 1877, [2] 16, 188, 1880, [2] 21, 77) thus:

and 
$$\begin{array}{c} \operatorname{Ca(OCONH_2)_2} = \operatorname{CaCN_2} + 2\operatorname{H_2O} + \operatorname{CO_2} \\ \operatorname{Ca(OCN)_2} = \operatorname{CaCN_2} + \operatorname{CO_2} \end{array}$$

Cyanamide is also obtained by the action of mercuric oxide on a solution of thiourea

$$CS(NH_2)_2 + HgO = NH_2CN + HgS + H_2O$$

Several guanidine derivatives yield evanamide when boiled with water or alkalis, for example, nitroso guanidine decomposes thus:

(Thiele, Annalen, 1893, 273, 136)

Cyanamide is easily prepared from the disodium salt by neutralising its cold, aqueous solution with nitric acid, and evaporating to dryness on the water bath in a vacuum, the dry residue is extracted with ether Werner (Chem Soc Trans 1916, 1323) describes a convenient method of preparing it from the readily accessible calcium salt; 100 grams are mixed with 125 gr of acetic acid and 120 gr of water, the mixture well kneaded and left for 24 hours The friable mass is then powdered and extracted with ether in a Soxhlet apparatus. Yield 94-96 p c of theory

Cyanamide forms colourless, deliquescent crystals which melt at 43°-44° (E. Colson, Chem. Soc. Trans. 1917, 555), it is readily soluble in water, alcohol and ether, very sparingly soluble in carbon disulphide, benzene, and chloroform. When heated it polymerises very readily to dicyanodiamide,  $\mathrm{NH}_2$  (C NH) NH CN, it boils at 140° at 19 mm, much melamine left behind; the neutral aqueous being solution possesses considerable stability, but polymerisation occurs in presence of either acids or alkalis. The process has been studied carefully by Morrell and Burgen (Chem. Soc Trans 1914, 576) and by Grube and Kruger (Zeitsch phys. Chem 1914, 86, 65), who find that in alkaline solutions the reaction takes place between undissociated cyanamide molecules and cyanamide ions,  $H_2CN_2+HCN_3'=H_3C_3N_4'$ . The rate of polymerisation is therefore a maximum when one-half of the cyanamide is in the ionised form, a condition which is practically attained when a quantity of a strong base (soda or lime) equivalent to one half of the cyanamide is added, the salts of a weak base such as ammonia being largely hydrolysed, the necessary concentration of cyanamide ion is not readily attained, and weak bases, therefore, have a much smaller polymerising action than strong ones.

The action of acids is more complicated,

polymerisation and hydrolysis to urea occurring

simultaneously. According to Carlson (Zeitsch. angew. Chem 1914, 27, 724) cyanamide, treated with 10 to 20 p.c sulphuric acid, is converted almost wholly into urea at the ordinary temperature, whereas near the boiling-point polymerisation predominates In acid solutions the reaction is further complicated by the hydrolysis of the dicyanodiamide to dicyanodiamidine


471

 $NH_{2}(C \cdot NH)NHCN + H_{2}O$ = NH<sub>2</sub>(C; NH) NH CO NH<sub>2</sub>

a strong base which neutralises the acid

Cyanamide is an extremely weak acid, its dissociation constant, for H2CN2=H+HCN2 is of the order 10-11 (Grube and Kruger) sodium salt NaHCN2 crystallises from a solution of cyanamide in absolute alcohol when sodium ethylate is added to it, but the salt Na<sub>2</sub>CN<sub>2</sub> can only be prepared by one of the dry fusion methods mentioned above. The calcium salt CaCN, also decomposes when treated with water yielding a solution of  $\text{Ca}(\text{HCN}_2)_2$ The silver salt Ag<sub>2</sub>CN<sub>2</sub> is a canary-yellow precipitate obtained by adding a solution of silver nitrate to a dilute ammoniacal solution of cyanamide, it is readily soluble in cold dilute nitric acid. The lead salt PbCN<sub>2</sub> is an orange-yellow precipitate Cyanamide, also combines with acids, a white crystalline substance H2CN2 2HCl is formed when dry hydrochloric acid is passed into a solution of cyanamide in dry ether (Mulder and Smit, Ber 1874, 7, 1634).

Cyanic acid HCNO is prepared by distilling cyanuic acid (Wohler) or by warming urea with phosphorus pentoxide (Weltzien, Annalen, 1858, 107, 219) and condensing the vapour at -18° It is a volatile liquid, the smell of which It is a volatile liquid, the smell of which resembles that of strong acetic acid It is extremely unstable, changing rapidly even at 0° into the white amorphous polymeric substance cyamelide The solution in dry ether is comparatively stable. The interesting restance cyamelide lationship of cyanic acid HCNO, cyanuric acid



 $(HCNO)_3$ , and cyamchide  $(HCNO)_x$ , has been studied by Troost and Hautefeuille (Compt rend. 1868, 67, 1340), van't Hoff and van Deventer (Studies in Chem Dynamics, 1896). The dia (Studies in Chem Dynamics, 1896) gram represents the approximate relative positions of the vapour pressure curves of cyanic acid (HCNO), cyanuric acid (HCNO)<sub>3</sub>, and cyamelide  $(HCNO)_x$ , all of which yield the same vapour.

(The few points determined experimentally are indicated by small circles.) At the triple

point A (150° and 50 mm.) the two solid phases cyamelide and cyanuric acid coexist together with the vapour phase. Above 150° cyanuric acid is the stable phase (vapour pressure curve AC), at lower temperatures cyamelide (curve FAB) The position of the vapour pressure ıs stable curve of cyanic acid itself (very roughly indicated by DE) shows that eyanic acid is quite unstable relatively to cyamelide. If the curve DE cuts the curve AB and AC the points of intersection would represent the melting-points of cyamelide and cyanuric acid respectively. They obviously he at temperatures at which these substances decompose.

In aqueous solution eyanic acid is hydrolysed, slowly in cold dilute solutions, rapidly when warmed, to ammonia and carbon dioxide,

HCNO+H<sub>2</sub>O=NH<sub>3</sub>+CO<sub>2</sub>.

The formation of cyanates by the action of cyanogen on alkalis, by oxidation of cyanides by the action of alkalis on cyanogen bromide, and from carbon monoxide and ammonia, has already been described

Ammonium cyanate NH4 CNO is prepared in the form of a white snow, by mixing the vapours of cyanic acid and ammonia well diluted with an indifferent gas. When warmed it is converted into urea NH<sub>4</sub>CNO=CO(NH<sub>2</sub>)<sub>2</sub> The same reaction occurs in aqueous solutions, in which, however, it is reversible, 3.2 p.c of the cyanate remaining unchanged at 32° and 65 p c. at 100' (Walker and Hambly, Chem Soc. Trans 1895, 746)

Potassium and Sodium cyanates KCNO and NaCNO are easily prepared by grinding together the dry cyanides and the theoretically necessary quantity of lead oxide, and heating the mixture gently until the lead is melted. The salt is then recrystallised from water or dilute alcohol According to Paterno and Pannain (Gazz. chim ital 1904, 34, ii 152) when a solution of 4 to 6 mols KCN and 1 mol. KOH per litre is electrolysed with 4-6 volts and 1-4 amps per sq dcm of anode surface and the solution at the anode agitated, the evanide is oxidised almost quantitatively to cyanate. Milbauer (Zeitsch anoig Chem 1904, 42, 433) observed that potassium sulphocyanide, when heated at 300° with lead oxide, gave cyanate thus: KSCN+PbO=KCNO+PbS

The decomposition of the alkali cyanates at high temperatures and their reduction by carbon are described above.

Potassium cyanate crystallises in thin transparent plates of sp gr 2 05, readily soluble in water, and anhydrous ammonia, insoluble in absolute alcohol, and fusible without decomposition below a red heat

Sodium cyanate. 100 grams of water at 16° dissolve 10.68 grams; it fuses without decomposition about 500°. Lead cyanate is a crystalline precipitate, which is formed when solutions of a soluble cyanate and of a lead salt are mixed The dry salt is stable, but it decomposes when boiled with water, yielding urea and lead carbonate.

Silver cyanate is very slightly soluble in cold water, 100 c.c. at 12° dissolve 0 006 gram, but it is much more soluble in hot water. It is readily soluble in aqueous ammonia, and sparingly soluble in cold dilute nitric acid

Sulphocyanides or Thiocyanates. (Ger. Rhodanide or Schwefelcyanide.)

At the present time sulphocyanides are obtained exclusively from the hydrocyanic acid of coal gas, but a modification of Gélis' synthetic process was working in 1907.

A. W Hofmann (Reports by the Juries, Inter Exhib of 1862, 59-63), said that at that time ammonium sulphocyanide obtained by washing the spent oxide of gasworks was only treated to recover its ammonia, the sulphocyanide being thrown away. In 1863 P. Spence (Eng. Pat 2364 of 1863 (provisional only)) proposed to separate the sulphocyanides in gas liquor (after distilling off the ammonia) in the form of cuprous sulphocyanide by adding copper and ferrous sulphates, and the idea has been revived in various forms since that time, but it does not appear to have become of any Gas liquor is really a very dilute importance solution of sulphocyanide, as the following figures, calculated from Linder's analyses of various English gas liquors (Chief Inspector of Alkalı Works, 42nd Rep 1905, 35), show :-

Grams per litre

HCN as ferrocyanide . 1 0 008 0 0 0 576 0 460 0 HCN as sulphocyanide . 1 088 4 457 1 238 1 029 1 042 HCN as hydrocyanic acid 0 680 0 0 0 000 0 050 0 355

These liquors had all been stored for some perfectly fresh gas liquor contains ammonium sulphide and cyanide By the action of the air, the former is oxidised to ammonium polysulphides, which convert the cyanide into sulphocyanide; thus:

$$(NH_4)_2S_2 + NH_4CN = NH_4SCN + (NH_4)_2S$$

The recovery of the whole of the hydrocyanic acid of coal gas in the form of a concentrated solution of ammonium sulphocyanide is possible by applying this reaction in a suitable way

A very large part of the gas liquor produced is distilled and the gases evolved, consisting of ammonia, carbon dioxide, sulphuretted hydrogen, and a little hydrocyanic acid, passed into sulphuric acid for the production of ammonium The unabsorbed gases, mixed with sulphate steam, are first cooled, which yields an aqueous solution of hydrocyanic acid and sulphuretted hydrogen, known as 'Devil liquor,' which is run to waste.

The cooled gases are passed usually into a heap of ferric hydroxide, in order to absorb the sulphuretted hydrogen, or occasionally they are burned in a Claus kiln, in which, as Carpenter has shown, the hydrocyanic acid is converted by steam into ammonia

The waste liquors from the ammonia stills contain relatively considerable quantities of calcium sulphocyanide (about 0 15 p.c SCN) and phenols, and are usually discharged into Frankland and Silvester (J Soc Chem. Ind 1907, 231), and Fowler, Ardern and Lockett (*ibid* '1911, 174), have published detailed accounts of the bacterial oxidation of these impurities in the septic tank system of sewage treatment. It appears that sulphocyanides are practically completely oxidised in the contact beds, if sufficient time is allowed Fowler and Holton (J. Soc Chem. Ind. 1911, 180) have shown that 92-98 pc of the impurities in the ammonia still effluent can be oxidised by allowing it, after dilution to one-tenth of its original strength, o trickle through a column of clinker inoculated with bacterial material from a contact bed of a sewage work; one cubic yard of filter is required

or 10 galls. per day of effluent.

Synthesis of sulphocyanides from ammonia ind carbon disulphide—In Gélis' original process A. W Hofmann, Reports by the Juries, Inter. Exhib of 1862, 59-63; A Gélis, Eng. Pat 1816 of 1860), a concentrated solution of ammonia and ammonium sulphide was agitated with arbon disulphide, yielding a solution of amnonium sulphocarbonate

$$(NH_4)_2S + CS_2 = (NH_4)_2CS_3$$

which was then heated to 90°-100° with potasium sulphide

 $2(NII_4)_2CS_3+K_2S=2KCNS+_22NH_4IIS+3H_2S$ I very large amount of sulphuretted hydrogen is produced in this way, the disposal of which would ccasion much trouble and expense Gunsburg nd Tchernkiae (Eng. Pat. 1148 of 1878, 1359 f 1879; 1261 of 1881) heated a 20 p c solution f ammonia with carbon disulphide in autoclaves provided with stirrers at 100° until a pressure f 15 atmos. was reached, which indicated the nd of the reaction As Conroy (J Soc Chem nd. 1896, 15, 8) has shown, ammonium sulphoyanide is formed quantitatively under these onditions, the large pressure developed being ue to the formation of the volatile ammonium ulphide  $4NH_3+CS_2=NH_4SCN+(NH_4)_S$ 

Subsequent improvements were all in the irection of diminishing the quantity of amionia required and reducing the pressure in the utoclaves The simplest and most effective ray of doing this is to add lime to the charge Prowther and Rossiter, Eng Pat 17846 of 893, Brock, Hetherington, Hurter, and taschen, Eng. Pat 21451 of 1893), the reaction hen taking place in accordance with the equation.

 $\begin{array}{l} \mathrm{NH_3} + 2\mathrm{CS_2} + 2\mathrm{Ca(OH)_2} \\ = \mathrm{Ca(SCN)_2} + \mathrm{Ca(SH)_2} + 4\mathrm{H_2O} \end{array}$ 

A certain excess of ammonia over the uantity required by the equation must be used prevent the formation of calcium sulpho-The excess of ammonia is distilled ff, the residual mixture treated with carbon ioxide to remove sulphuretted hydrogen (in the ay used in the treatment of Leblane alkali aste), and the calcium carbonate filtered off. he clear solution of calcium sulphocyanide is sily converted into other salts

#### THE PRINCIPAL SULPHOCYANIDES.

Potassium sulphocyanide KSCN crystallises anhydrous, striated prisms or needles of sp gr 9, which melt at 172 3°. It resembles saltetre in appearance and taste It is delinescent and very soluble in water, 100 grams of hich dissolve 177 2 grams at 0°, 217 grams at )°, and 239 grams at 25°. It is also soluble in thydrous ammonia and in many organic liquids he solubilities (in grams per 100 grams of olvent) are:

Acetone, 20.8 at 22°; 20 4 at 58°. Ethyl acetate, 0 44 at 0°; 0 4 at 14°; 0 2 at

Amyl alcohol, 0 18 at 13°; 2.14 at 100°.

Pyridine, 6.75 at 0°; 4.97 at 58°, and 3.21 at 115°.

The fused salt becomes deep blue at 430°; the colour is said to be due to the liberation of sulphur. The sulphocyanides are plant poisons. Frankland and Silvester found that gold-fish were not affected in 24 hours by quantities of potassium sulphocyanide up to 260 parts SCN per 100,000 of water, but even as little as 10 parts of SCN per 100,000 is sufficient to stop the multiplication of, and eventually to kill, Bacıllus coli communis.

Ammonium sulphocyanide NH<sub>4</sub>SCN crystallises in anhydrous, colourless leaflets of sp gr. 1 3, which melt at 159°, 100 grams of water dissolve 122 grams at 0° and 162 grams at 20°, it is readily soluble in alcohol When heated a little above its melting-point, it is partly converted into the isomeric thiourea. At 220° decomposition takes place, ammonia, sulphuretted hydrogen, and carbon disulphide being evolved, and guanidine sulphocyanide remaining

Thiocyanogen is obtained by the action of a solution of a halogen in an anhydrous solvent on a suitable thiocyanate, a dry ethereal solu-tion of bromine cooled to 0° and lead thiocyanate give good results By cooling an approximately N/2-solution in carbon disulphide to  $-70^{\circ}$ thiocyanogen is obtained in almost colourless crystals, melting at  $-3^{\circ}$  to  $-2^{\circ}$ , and decomposing at the ordinary temperature chemical behaviour thiocyanogen resembles the halogens, it is at least as strongly electronegative as iodine. The molecular weight has not yet been determined, but the constitution of thiocyanogen is probably CN-S-S-CN (E Soderback, Annalen, 1919, 419, 217)

# ANALYSIS OF CYANIDES.

The methods of analysis described here are those commonly employed in technical labora-

Detection of hydrocyanic acid -The material is distilled with an acid and the distillate ex-The Prussian blue test may be applied amined by evaporating the solution (after the addition of a little sodium hydroxide) in vacuô at temperature below 70°, to 1 cc, 02 cc. of 3 pc FeSO<sub>4</sub> and 0 05 gram KF are added, and, after 10 minutes, a small excess of 30 p c. HNO<sub>3</sub>. The blue colour is compared with that produced by a known quantity of KCN under similar conditions (A Viehoever and C O Johns, J Amer Chem Soc 1915, 37, 601). The sulphocyanide test is also characteristic; distillate containing 0 1-8 milligrams HCN is made alkaline with KOH, treated with 1 cc. ammonium polysulphide solution and evaporated to dryness. The residue is extracted with acetone, which dissolves KSCN only. After removing the acetone by evaporation the aqueous solution of the residue is treated with 2 c c. of 0 5 p.c FeCl<sub>3</sub>, and the colour matched by means of a standard sulphocyanide solution (M O Johnson, J Amer Chem Soc. 1916, 38, 1230). Ferric sulphocyanide may be extracted from the aqueous solution by ether; meconic acid gives a similar colouration in ether, which, however, is not destroyed by 10 pc. AuCla (P. Lavialle and L. Varenne, J. Pharm. Chim. 1915, 12, 74)

Other sensitive colour tests have the disadvantage that they may be produced by

substances other than HCN. An alkaline solution of pieric acid yields a reddish-violet colouration with NaCN, which is due to sodium iso-purpurate C<sub>6</sub>H<sub>2</sub>(NO<sub>2</sub>)<sub>2</sub> N(CN)<sub>4</sub> ONa, when excess of cyanide is used, but to amino-nitro-phenols when pieric acid is in excess, as is usually the case (A. C. Chapman, Analyst, 1910, 35, 469) The liquid to be tested is mixed with an equal volume of a solution containing 0 05 pc pierre acid and 0.5 p.c. Na<sub>2</sub>CO<sub>3</sub>, and digested at 40° for 1 hour. The colour is then compared with that produced by a standard HCN solution, I part of HCN in 500,000 parts water can be detected (A. D Waller, Proc Roy Soc. 1910, B 82, 574)

The cyanogen liberated by the action of HCN or a cupric salt oxidises many substances to coloured products, the tests are highly sensitive, but uncertain owing to the difficulty of excluding other oxidising agents For details, see Thiery (phenolphthalin), J Pharm Chim 1907, 25 Chem. Zeit 1913, 37, 609, G W Anderson (gualacum), J Soc Chem Ind 1917, 36, 195, J Moir and J Gray, J Chem Met and Min Soc S Africa, 1910, 11, 152, and J Moir, Chem Soc Proc 1910, 26, 115 (hydrocœrulignone, tolidine, dianisidine)

Owing to the hygroscopic Alkalı cyanıdıs nature of sodium and potassium cyanides when cold, samples, taken from different parts of a cake, should be transferred at once and without touching them with the hands to a hot iron mortar, broken up coarsely, and transferred to a hot perfectly dry bottle If the sample has to be kept for any length of time before it is analysed, it is preferable to use glass-stoppered bottles and to grease the stopper with a little Errors due to absorption of moisture in sampling are more common and more serious than those due to maccuracies in testing.

Estimation of cyanogen -5 grams of the sample are weighed out and made up to 500 c c; 25 c c of the solution are diluted to 200-300 c c and titrated with a solution of pure silver nitrate until the faintest visible turbidity results. The silver nitrate must be neutral and free from lead and zine especially venient strength of solution is obtained by dissolving 13 l grams of silver nitiate in 1 litie of distilled water 1 e c of this solution is equivalent to 0.01 gram of KCN The solution is standardised by means of a solution of pure sodium chloride (dried by heating to redness before weighing) containing 4 487 grams per litre, 1 c c of the silver solution should exactly precipitate 1 c.c. of the sodium chloride solution using pure potassium chromate as indicator impure solutions of cyanides, the end point is often more easily observed if a little caustic soda (or a few drops of ammonia) and a crystal of potassium iodide are added before running in the silver nitrate solution Sulphides interfere with the test, but they are easily removed by shaking the solution with a little lead carbonate and filtering it.

Estimation of sulphide -10 grams of the finely powdered sample are dissolved in 15 c.c. ot ordinary water and a solution of lead nitrate (2 4 grams per litre) run in gradually. The mixture is stirred until the colloidal lead sulphide at first formed has coagulated into brown flocks;

a drop of the solution is then placed on blotting paper beside a drop of the lead nitrate solution, and the two allowed to run together. When no difference can be detected between the colour of the paper where the drops have mixed and where they have not, the titration is considered to be at an end; 1 cc of the lead solution=0 001 gram of K<sub>2</sub>S (T Ewan, J. Soc Chem Ind. 28, 10, 1909, Rossiter, ibid 30, 583, 1911)

Water is estimated by weighing out about 1 gram of the coarsely powdered sample into a small distilling flask (10-20 cc) which is then evacuated by a good water pump or preferably by a mercury pump The sample is then warmed, gently at first, and finally to 200°-300°. The whole of the moisture is expelled in a few minutes without appreciable decomposition of

the cyanide

Carbonate is conveniently estimated by precipitating a suitable portion of the solution with barium chloride in the cold (to avoid decomposition of cyanate). The precipitate is washed and titrated, together with the paper, with normal hydrochloric acid and methyl

orange

The solution is neutralised with Chloride dilute nitric acid, and methyl orange and boiled for about 1 hour (in a draught cupboard, of course) to drive off the hydrocyanic acid and decompose evanate The chloride remaining (Richards and Singer, Amer Chem. J. 27, 205, 1902) is then precipitated by a known excess of standard silver nitrate solution, the silver chloride filtered off and the excess of silver titrated with ammonium sulphocyanide and iron alum in the usual way.

If ferrocyanide is present, it may be precipitated with excess of silver nitrate in ammoniacal solution; by acidifying the filtrate with nitric acid, silver chloride is precipitated and may be

weighed

Cyanamide in presence of alkali cyanides is easily estimated by exactly neutralising the solution of 1 gram of the substance in 100 to 150 cc of water with nitric acid, and then boiling for 20 minutes to expel hydrocyanic acid. After cooling, 2 or 3 drops of ammonia are added, and then excess of silver nitrate canary-yellow precipitate of silver cyanamide  $\mathrm{Ag}_2\mathrm{CN}_2$  is filtered off and dissolved in cold dilute nitric acid (which usually leaves a small residue of undissolved silver cyanide). The silver in the solution is then titrated by means of ammonium sulphocyanide, using ferric alum as indicator The method is applicable to most substances containing cyanamide, the boiling with acid being, of course, omitted when little or no cyanide is present.

When cyanamide is the principal constituent this method is insufficiently accurate, because the composition of the precipitate is not exactly that required by the formula Ag<sub>2</sub>CN<sub>2</sub>, the silver content depending to a small extent on the excess of silver nitrate used in precipitation. Caro (Zeitsch. angew Chem 1910, 23, 2405) has shown, however, that it contains the whole of the cyanamide nitrogen, which is very readily

estimated by Kjeldahl's method

Cyanate The solution is acidified with nitric acid and boiled for 20 minutes at least to expel hydrocyanic acid, and convert the cyanate into ammonium nitrate and carbon dioxide

The ammonia is then distilled off, after adding | treated with 4 to 5 times its volume of methyexcess of caustic soda, and estimated by titration with N-hydrochloric acid as usual. In presence of cyanamide, the method is not available, because some ammonia is formed from the cyanamide when it is boiled with caustic soda In this (rather uncommon) case, the cyanate must be estimated by collecting the carbon dioxide evolved when the sample is boiled with acid (Ewan, J. Soc Chem Ind 1904, 23, 244)

Ferrocyanide is most conveniently estimated, either alone or in presence of cyanides, by titration with a standard solution of zinc sulphate until a drop of the solution no longer gives a blue colour when brought in contact with a drop of ferric solution on paper A zinc alkalı ferrocyanide of somewhat variable composition is precipitated, and H G ('olman (Analyst, 33, 261, 1908; and J Soc Chem Ind 27, 806, 1908) has shown that the quantity of zinc solution required to precipitate a given quantity of ferrocyanide depends, to some extent, on the nature of the metal in the ferrocyanide used effect can be eliminated almost entirely by adding an excess of potassium sulphate. The adding an excess of potassium sulphate method is carried out as follows a solution containing 10 2 grams of ZnSO<sub>4</sub>,7H<sub>2</sub>O and 10 c c of concentrated sulphuric acid per litre is standardised against pure potassium ferroevanide, 25 c c of a solution containing 10 grams of K4Fe(CN)6 3H2O per litre, mixed with 25 c c of a saturated solution of potassium sulphate, 25 c c of water, and 10 c c of 1 10 sulphuric acid, are convenient quantities Equal volumes of the standard zinc and ferrocyanide solutions are required, the precipitate formed having the composition Zn<sub>3</sub>K<sub>2</sub>(FeCy<sub>6</sub>)<sub>2</sub> (R Meurice, Ann Chim anal 1913, 18, 342) Subsequent titrations should be made as nearly as possible under the same conditions

A more accurate method is that of W Feld (Journ fur Gasbeleuchtung, 47, 565, 1903). Ferrocyanide, equivalent to 03 to 05 gram of yellow prussiate is boiled for 5 minutes with 10 c.c. of N-NaOH and 15 c.c. of 3 (molecular)  $\widetilde{N}$ -HgCl<sub>2</sub> If eyanide is present, the boiling is continued for 10 minutes and the hydrocyanic acid distilled into caustic soda and titrated with silver intrate with potassium iodide as indicator 100 c.e of boiling 0.1N-Hg(I<sub>2</sub> are now run in, and the boiling continued for 10 minutes to convert ferrocyanide into mercuiic This is then distilled with 30 cc of cvanide 3N-sulphuric acid for 20-30 minutes, and the hydrocyanic acid collected in caustic soda and The conversion of the ferrotitrated as before cyanide into mercuric cyanide is avoided by H. E Williams' procedure (J Soc Chem Ind 1912, 31, 468). After distilling off hydrocyanic acid in the way just described (lead nitrate may be used instead of magnesium chloride), 0 05-0.1 gram of pure cuprous chloride dissolved in HCl is added, and the hydrocyanic acid distilled off by I hour gentle boiling Even insoluble ferrocyanides are completely decomposed

The estimation of the valuable constituents of gasworks products (cyanide and ferrocyanide) is performed by triturating 30 to 40 grams of the material in a mortar with excess of a cold solution of caustic soda for several hours. A little ferrous sulphate may be added to convert cyanide into ferrocyanide. The filtrate is

lated spirit. Carbonyl ferrocyanide (from 2 to 5 p c. of the cyanogen is usually present in this form, but exceptionally as much as 20 pc may be present), sulphocyanide, sulphide, and excess of caustic soda remain in solution, whilst the whole of the sodium ferrocyanide is precipi-tated. The precipitate is then treated by one of the methods described above (H G. Colman, Analyst, 33, 261, 1908, and J Soc Chem. Ind. 27, 806, 1908, F. W. Skirrow, *ibid.* 29, 319, 1910)

Sulphocyanide. Ferrocyanide, if present, is first removed from the solution by adding iron alum after acidifying slightly. The filtrate is treated with acid sodium sulphite in excess and then with copper sulphate and boiled for 1 or 2 minutes. The precipitate of cuprous sulphocyanide, which may be mixed with copper sulphide, is filtered off and washed with boiling water It is then digested with excess of caustic soda which gives cuprous hydroxide and a solution of sodium sulphocyanide. This is separated by filtration, acidified with nitric acid, and titrated with silver nitiate, using a ferric salt as indicator (Linder, 43rd Alkali Report,

The small quantities of cyanogen compounds in concentrated ammonia liquor may be estimated by converting them into ferric sulphocyanide and measuring the depth of colour by means of the tintometer (P. E. Spielmann and H. Wood, J. Soc. Chem. Ind. 1919, 38, 43T, and 369T) Sulphocyanide is first estimated by acidifying the diluted sample, in the cold, with sulphuric acid, and removing ferrocyanide by means of a small excess of non alum; a large excess of non alum is added to the filtrate and the depth of colour measured Cyanide is converted into sulphocyanide by digestion at 30°-35° with ammonium polysulphide, and a second estimation made as above described. Thiocarbonate is converted into sulphocyanide by digestion at 70°-75° for 45 minutes, cyanide and ferrocyanide remaining unaffected

Alkalı metals ın commercial cyanides 1 gram of the substance is evaporated to dryness with excess of hydrochloric acid. Traces of iron and calcium are removed by re-dissolving, adding a drop of ammonium carbonate, and filtering The solution is again evaporated to dryness, and the residue is gently fused and weighed The mixture of alkaline chlorides is dissolved in water and an aliquot portion titrated with silver nitrate, using chromate as indicator If n gram of cyanide has yielded a gram of the pure mixed chlorides containing b gram of chlorine, the percentage of potassium in the cyanide is given by the formula

$$\left(\frac{2\ 431a-4\ 012b}{n}\right)$$
100

This method is sufficiently accurate for most purposes, but the gravimetric estimation of the potassium as potassium platinichloride must be resorted to if a greater degree of accuracy is T. E. required.

CYANIBIN, CYANIN v Anthocyanins CYANINE and CYANINE DYESTUFFS v. QUINOLINE COLOURING MATTERS

CYANITE v KYANITE. CYANOMACLURIN. C15H12O6, a colourless compound existing in Jakwood is closely allied to the catechins, and may possibly belong to the same group. In its general reactions, indeed, it is very similar to these compounds, but differs from them in that by alkali fusion it yields  $\beta$ -resorcylic acid and not protocatechuic acid.

A full description of the properties of this compound is given in the article on JAKWOOD,

 $\hat{\mathbf{C}}$ YANOSINE v Triphenylmethane colouring matters.

Iso-CYANURIC ACID v. FULMINURIC ACID.
CYCLOFORM. The trade name of isobutyl ester of p-amino benzoic acid

# NH2.C6H4.COOC4H9

used as a local an esthetic. A white crystalline powder, mp. 65°. Sparingly soluble in water (0 022 pc), readily soluble in alcohol and other.

CYLINDRITE. (Kylindrit, Ger) A complex sulphide ore of tin, Pb<sub>3</sub>FeSn<sub>4</sub>Sb<sub>2</sub>S<sub>14</sub>, containing, according to this formula, Sn 25 85 p.c It forms very peculiar cylindrical rolls of thin folia, the concentric shells parallel to the axis of the cylinder readily separating under pressure The smooth, bright surfaces of separation no doubt represent the cleavages of cylindrically curved crystals The colour is blackish leadgrey, with a brilliant metallic lustre. mineral marks paper, and is difficult to pulverise, in these respects resembling graphite. The ore consists of numbers of these rolls, which measure up to 3 cm in length and 1 cm in diameter, confusedly aggregated together, or sometimes with a sub-parallel grouping, the interspaces being filled with massive material It has been found in some quantity in the Santa Cruz mine, at Poopó, Bolivia. The above formula, written in the form

# 3PbSnS<sub>2</sub>+SnFeSb<sub>2</sub>S<sub>8</sub>

brings out a relation between cylindrite and the allied Bohivian minerals, teallite (PbSnS<sub>2</sub>) and franckeite (3PbSnS<sub>2</sub>+Pb<sub>2</sub>FeSb<sub>2</sub>S<sub>3</sub>) Analyses I and II by G T Prior (1904), III. by A Frenzel (1893)

	I	II	111
Pb	. 35 24	34.58	35.41
Fe	. 281	2 77	3 00
$\mathbf{A}\mathbf{g}$	. 0 50	0.28	0.62
Sn	$25\ 65$	25 10	26.37
$\mathbf{S}\mathbf{b}$	12 31	12 98	8 73
$\mathbf{s}$	. 23 83	$23 \ 88$	24 50
	100 34	$99\ 59$	$98 \ 63$
Sp gr.	. 546	5 49	542
. 0			LJS

CYMARIN. The physiologically active constituent of the root of Apocynum cannabinum and A and rosoemi folium, is a glucoside,  $C_{30}H_{46}O_{10}$ , colourless glistening prisms, mp  $135^{\circ}-140^{\circ}$ , which, when boiled with alcohol containing hydrochloric acid, is resolved into cymarose  $C_7H_{14}O_4$ , colourless prisms, mp  $88^{\circ}$ , a substance resembling digitoxose, of which it is probably a methyl ether, and cymarigenin  $C_{23}H_{10}O_5$ , a lactone which appears to be identical with apocynamarin (Moore, Chem. Soc Trans. 1909, 95, 734); it crystallises from aqueous methyl alcohol in rhombic tablets, mp  $171^{\circ}$  (decomp). With a solution of a diazonium salt, cymarigenin forms dyes; it reduces ammoniacal silver nitrate, forms a

benzoyl derivative  $C_{30}H_{36}O_{7,\frac{1}{2}}H_{2}O$ , pearly leaflets, m.p. 230°, and when treated with hydrogen chloride in chloroform solution, is converted into anhydro cymaragenin  $C_{23}H_{28}O_{4}$ , rosettes of crystals, m.p. 246° (decomp) (Windaus and Hermanns, Ber. 1915, 48, 979).

According to Impens (Pfluger's Archiv. 1913, 153, 239), cymarin is not a glucoside. It corresponds in activity with digitalis, being slightly more potent as a diuretic and slightly less so as a cardiac stimulant. The 'cynotoxin' of Finnemore and 'apocynamarin' are probably decomposition products of cymarine. V. Digitalis and Strophanthus.

CYMENES. Cymols. The name cymene was first applied to the hydrocarbon isolated by Gerhardt and Cahours (Ann Chim. Phys [3] 1, 102 and 372) from Roman oil of cumin, which is the ethercal oil of Cuminum cyminum. Until 1891 this hydrocarbon was wrongly called p-methylpropylbenzene, but in that year Widman (Ber 1891, 439, 970, 1632) proved it to be the isopropyl-compound (see also Filet, J. pr Chem [2] 44, 150) Frequently, however, the various benzene hydrocarbons of the formula C<sub>10</sub>H<sub>14</sub> are grouped together under the collective name of cymenes No fewer than 22 of these are theoretically possible, of which 18 have been prepared; but only the most important, namely, cymene proper (p-methylisopropylbenzene) and the tetramethylbenzenes are described here

p-Methylsopropylbenzene (Cymene)

$$^{\prime}\mathrm{CH_{3}})_{2}\mathrm{CH}\ \mathrm{C_{6}H_{4}}\ \mathrm{CH_{3}}\ (1,\ 4)$$

occurs in Roman oil of cumin (Gerhardt and Cahours, lc.; Bruhl, Ber 1892, 149; Wolpian, Chem Soc Abstr 1897, 1 357), in the volatile oil from the seeds of the water-hemlock (Cicuta virosa) (Trapp, Annalen, 108, 386); in the oil from Phychotis arowan (H. Muller, Ber 2, 130); in oil of thyme (Lallemand, Annalen, 102, 119; Labbé, Bull Soc. chim. 1898, [111.] 1009) along with thymol C<sub>10</sub>H<sub>14</sub>O and thymene C<sub>10</sub>H<sub>16</sub>, in eucalyptus oil (from Eucalyptus Globulus) (Faust and Homeyer, Ber. 7, 1429); in oil of caraway (Wolpian, Chem Soc Abstr. 1897, i 17), in bitter fennel oil (Tardy, J. Pharm. Chim. 1897, 98), in oil of lemon grass (Barbier and Bouveault, Compt. rend, 121, 1159; Stiehl, J pr Chem. 1898, 51); in oil of origanum (Gildermeister, Arch Pharm 1895, 174); in geranicl (Semmler, Ber 1891, 205); and in American petroleum (Mabery, Amer Chem. J 19, 419). Considerable quantities of crude cymene, known as sulphite turpentine, collect on the surface of the liquid in the separator used in recovering sulphur dioxide in the manufacture of wood May be obtained from camphor by pulp abstracting from it the elements of water

$$C_{10}H_{16}O = C_{10}H_{14} + H_2O$$

by heating it either with phosphorus pentoxide (Dumas, Ann. Chim Phys 50, 226; Delalande, *ibid.* [3] 1, 368) or with phosphorus pentasulphide (Pott, Ber 2, 121), or with phosphorus pentachloride (Lippmann and Lougunine, Bull Soc. chim. [2] 7, 374; *see also* Bruhl, *l.c.*; Tiemann, Ber. 1895, 1089). Various isomerides of camphor yield cymone by the same treatment. The terpones C<sub>10</sub>H<sub>16</sub> may be converted into cymone by withdrawing two atoms of hydrogen; this

CYSTINE. 477

may be effected in the case of oil of turpentine, for example, by distilling it with bromine (Oppenheim, Ber. 5, 94 and 628), or better, with iodine (Kekulé, ibid 6, 437). Old turpentine generally contains cymene (Wallach, Chem. Zentr. 1898, 1 572; Kondakow and Schindelmeier, Chem. Zeit 30, 722). It has been obtained synthetically from p-bromisopropylbenzene by means of methyl lodide and sodium (Widman, l.c.). To prepare cymene, camphor is warmed with an equal weight of phosphorus pentoxide, the cymene is poured off, treated again twice with a little phosphorus pentoxide, and finally rectified from sodium. The yield varies from 50 to 80 pc (Fittica, Annalen, 172, 307) It can also be prepared from isopropyliodide, and p-bromotoluene (Reuter, Chem Zentr 1892, 1 625, Bruhl,  $l\,c$ ) or from Liquid boiling at 177°-1775° (corr); sp gr. 0 8670 at 7.9°/4° (Perkin, Chem Soc Trans 1896, 1125; 1900, 267) With chromic acid it yields paratoluic acid (CH<sub>3</sub>  $C_6H_4$  CO<sub>2</sub>H) and terephthalic acid  $C_6H_4$ (CO<sub>2</sub>H)<sub>2</sub> (1, 4); with potassium permanganate, hydroxyssopiopylbenzoic acid (CH<sub>3</sub>)<sub>2</sub>C(OH)C<sub>6</sub>H<sub>4</sub> CO<sub>2</sub>H (1, 4) Taken internally, it appears in the urine as cumic (p-isopropylbenzoic) acid

$$(CH_3)_2CH \cdot C_6H_4 CO_2H (1, 4)$$

The preparation of cumene (q v) from propylbromide, benzene, and aluminium chloride, and that of cymene from p-bromotoluene, normal propylbromide, and sodium (Fittig, Schafer, and Konig, Annalen, 149, 331) was for a long time the cause of great confusion in ascertaining the constitution of members of the cumene and cymene groups, as it was not realised that in each case the propyl- group changed into the isopropyl- group owing to the temperature of the reaction not being regulated

Tetramethylbenzenes

Consecutive tetramethylbenzene (Prehmene)  $C_6H_2(CH_3)_4$  (1, 2, 3, 4) When sodium durene-sulphonate

 $C_6H(CH_3)_4SO_3Na(CH_3$ -groups 1, 2, 4, 5, v infia) is dissolved in concentrated sulphuric acid and allowed to stand for 12 hours, prehnitenesulphonic acid is formed. The mixture is diluted with water, saturated with barium carbonate, the barrum salts converted into sodium salts, and these, by treatment with phosphorus pentachloride and ammonia, transformed first into sulphonic chlorides and then into sulphon-Prehnitenesulphonamide is separated amides by means of its sparing solubility in alcohol, and is then hydrolysed by heating with concentrated hydrochloric acid to 170°, when it yields prehiitene (Jacobsen, Ber 19, 1213; see also V. Meyer and Molz, Ber. 1897, 30, 1278). Also obtained by heating bromocumene

 $C_6H_2Br(CH_3)_3(CH_3,Br,CH_3,CH_3:1,2,3,4)$ with methyl iodide, benzene, and sodium, in a sealed tube at  $150^{\circ}$  for 12 hours —Liquid boiling at 204° (corr); solidifies in a freezing mixture, and melts at -4° Dilute nitric acid oxidises it to prehnitylic acid C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>CO<sub>2</sub>H.

2. Unsymmetrical testamethylbenzene (Isodurene) C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>4</sub> (1, 2, 3, 5). From bromomesitylene (v Cumenes), methyl 10dide and

sodium in presence of benzene (Jannasch, Ber. 8, 356). Best by passing methyl chloride into a mixture of mesitylene and aluminium chloride.

$$\substack{ \text{C}_{\pmb{6}}\text{H}_{\pmb{3}}(\text{CH}_{\pmb{3}})_{\pmb{3}} + \text{CH}_{\pmb{3}}\text{Cl} = \text{C}_{\pmb{6}}\text{H}_{\pmb{2}}(\text{CH}_{\pmb{3}})_{\pmb{4}} + \text{HCl} \\ (1, \, 2, \, 3, \, 5) }$$

(Jacobsen, Ber 14, 2629) -Liquid boiling at  $195^{\circ}$ – $197^{\circ}$ , solidifies in a freezing mixture. With dilute nitric acid it yields  $\alpha$ -isodurylic acid,  $\beta$ -isodurylic acid, and mesitylenecarboxylic acid, isomeric acids of the formula

and dimethylisophthalic and dimethylterephthalic acids, isomeric acids of the formula C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>(CO<sub>2</sub>H)<sub>2</sub> (Jannasch and Weiler, Ber. 1894, 3441, 1895, 531), potassium permanganate ultimately converts it into mellophanic acid

C<sub>6</sub>H<sub>2</sub>(CO<sub>2</sub>H)<sub>4</sub> (1, 2, 3, 5) Symmetrical tetramethylbenzene (Durene) C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>4</sub> (1, 2, 4, 5) (Schulze, Ber 18, 3032) Occurs in coal tar By the action of sodium on a mixture of bromopseudocumene  $(^1_6H_2(\mathrm{CH}_3)_3\mathrm{Bi}$  (1, 2, 4, 5), methyl iodide and sodium (Jannasch and Fittig, Zeitsch Chem 1870, 161) From toluene and methyl chloride in presence of aluminium chloride (Friedel and Crafts, Ann Chim Phys [6] 1, 461) o- or p-xylene, methyl chloride and aluminum chloride (Jacobsen, Ber 14, 2629) From pseudocumene, methyl iodide, and aluminium chloride (Rugheimer and Hankel, ibid 1896, 2173) - Crystalline mass, smelling like camphor, melte at 79°-80°, and boils at 189°-191° (Jacob-Yields by oxidation with potassium permanganate, finally pyromellitic acid

 $C_8H_9(CO_3H)_4$  (1, 2, 4, 5)

CYMOLS v CYMENES.

CYMOPHENOL v CAMPHORS CYPRIAN VITRIOL. Copper sulphate (v.

COPPER'

CYPRIDOL. Trade name for a suspension of mercuric iodide in sterilised oil.

CYRTOLITE v ZIRCON

CYSTAMINE, CYSTOGEN. Syns for hexamethylenetetramine

CYSTAZOL. An additive compound of hexamethylenetetramine and sodium benzoate. Used as a urmary antiseptic

CYSTINE, B-disulphido-a-aminopropionic acid HO<sub>2</sub>C CH(NH<sub>2</sub>)·CH<sub>2</sub> S S·CH<sub>2</sub> CH(NH<sub>2</sub>) CO<sub>2</sub>H first discovered by Wollaston in a urmary calculus (Phil. Trans 1810, 223), is a product of proteid metabolism under certain pathological conditions (Blum, Beitr chem Physiol Path. 1903, 5, 1, Wolf and Shaffer, Amer J Physiol. 19, Loebisch, Annalen, 182, 231, Niemann, *ibid* 187, 101; Thiele, Trans Path Soc 1907, 788, 255), which, according to Abderhalden (Zeitsch. physiol. Chem. 1903, 38, 557), are hereditary. Dreschel found cystine as a normal product of metabolism in the liver of horse (Arch. Anat Physiol 1891, 243) and porpoise (Zeit Biol 1896, 33, 85) It was found by Kulz (*ibid.* 1890, 27, 415) among the soluble products of an artificial pancreatic digestion of fibrin; and is a constant product of hydrolysis of keratın (Mörner, Zeitsch physiol. Chem. 1899, 28, 595): Buchtala (*ibid*. 1907, 52, 474) obtained the following percentages of cystine from the materials mentioned: human hair, 13-14.5; human nails, 5.15; horsehair, 7.98;

478 CYSTINE.

horse's hoofs, 3 20, ox hair, 7 27, hoofs of oxen, 5 37; pigs' bristles, 7 22; pigs' hoofs, 2 17
According to Folin (J. Biol. Chem. 1910, 8,

According to Folm (J Biol Chem 1910, 8, 9), cystine is most readily prepared from wool by boiling it in a reflux apparatus for 5-6 hours with twice its weight of stiong hydrochloric acid, or until it ceases to give the biuret reaction, on removing the free hydrochloric acid by the addition of solid sodium acctate, a dark heavy precipitate of cystine is obtained, this is purified by decolourising a solution in 3-5 pc hydrochloric acid with bone-black, and precipitating the cystine in the hot filtrate by hot concentrated sodium acctate solution (see also Denis, J Biol Chem 1911, 9, 305)

The cystine of urinary calculus is identical with that obtained from protein hydrolysis (Rothera, J. Physiol. 1905, 32, 175., Alsberg and Folin, Amer. J. Physiol. 1905, 14, 54; Fischer and Suzuki, Zeitsch. physiol. Chem. 1905, 45, 405), and was proved to be \$\beta\delta\subsymbol{u}\text{bulletholo-a-amino-propionic acid.}\$ by Friedmann (Bett. Chem. Popular Cystine from ethyl monobenzovlserine, obtained by reducing ethyl monobenzovlserine, obtained by reducing ethyl formylhippurate, the product of the condensation of ethyl formate and ethyl hippurate. On heating othyl monobenzoylserine with phosphorus pentasulphide, and hydrolysing the product, cystein is obtained and this is readily converted into cystine by oxidation with air in an ammoniacal solution:

H CO₂Et+NHBz CH₂ CO₂Et →CHO CH(NHBz) CO,Et

→OH CH, CH(NHBz)CO,Et →SH CH, CH(NHBz) CO,Et

 $\rightarrow$ SH CH<sub>2</sub> CH(NH<sub>2</sub>) CO<sub>2</sub>H  $\rightarrow$ CO<sub>2</sub>H CH(NH<sub>2</sub>) CH<sub>2</sub>S S CH<sub>2</sub> CH(NH<sub>2</sub>) CO<sub>2</sub>H

(Erlenmeyer, Ber 1903, 36, 2720) Fischer and Raske (ibid 1908, 41, 893) also obtained incystine from dl-α-amino-β-chloropropionic acid hydrochloride by heating it with barium hydrosulphide at 100° in sealed tubes, and oxidising the product by atmospheric oxygen in the presence of ammonia From the natural leserine the authors obtained, by a similar process, an optically active cystine which is identical with the l-cystine obtained from natural sources. Removal of sulphur by ammonium copper solution gives dl-α-aminopropionic acid, and removal of sulphur by zine dl-alanine (Mauthner, Zeitsch physiol Chem. 1912, 78, 30–32)

For the importance of cystine in nitrogen metabolism, see Abderhalden (Zeitsch physiol Chem 96, 1-147, Ber 1916, 49, 2449-2473), and in humin formation, Roxas (J. Biol. Chem.

1916, 27, 71-93)

Cystine crystallises in thin colourless hexagonal plates; it has no melting point, but decomposes slowly at  $258^{\circ}-261^{\circ}$  (Nouberg and Mayer, Zeitsch physiol Chem 1905, 44, 472) It is sparingly soluble in water, insoluble in alcohol or acetic acid, soluble in acids or alkalis. The heat of combustion at constant pressure is 4 137 calories (Emery and Benedict, Amer J Physiol 28, 301–307). The cystine molecule contains two asymmetric carbon atoms, and lavo-cystine, the naturally occurring variety, has  $[a]_{D}-97$  5° in ammonia,  $-223^{\circ}$  in hydrochloric acid solution (Gaskell, J. Physiol. 1907, 36, 142), or  $-205\cdot28^{\circ}$ 

in a hydrochloric acid solution containing 112 grams hydrochloric acid in 100 cm (Mauthner, ibid 29) l-Cystine is converted to the extent of 50 p c into r-cystine by heating with excess of hydrochloric acid of sp.gr. 1124, at 165° for 12–15 hours, this decomposes at the same temperature as the active isomeride is three times as soluble in water and can be resolved into the active components by the action of Aspergillus niger; the l-compound being consumed

Cystine is readily reduced to cystein (a aminoβ-thiolpropionic acid) HS CII, CH(NH.) CO<sub>2</sub>H (Baumann, Zeitsch physiol. Chem. 1882, 8, 300, Mauthner, Zeit Biol 1901, 42, 176), which can be converted by the action of bromine water into cysteic acid (a-amino-B-sulphopro-pionic acid) HO<sub>3</sub>S CH<sub>2</sub> CH(NH<sub>2</sub>) CO<sub>2</sub>H, and from this, by loss of carbon dioxide, taurine NH2 CH2 CH2 SO3 II is obtained (Friedmann, Chem Physiol Path 1902, 2, 433). According to Von Bergmann (ibid 1903, 4, 92), the normal taurine in the bile originates from the cystin groups contained in the ingested protein Cystem yields cystine on oxidation the change occurring spontaneously in alkaline solution, although the rate is increased by the addition of a mixture of ferric chloride and potassium cyanide (Mathews and Walker, J. Biol Chem 1909, 6, 289) For the behaviour of cystine on oxidation, see Dems (J. Biol Chem. 1911, 9, 373), and for its breakdown under the action of certain bacteria, see Burger (Arch Hyg 1914, 82, 201-211, from the abstr. Amer. Chem Soc 1916, 10, 2911)

Slyke's reaction (Ber 1910, 43 3170) for  $\alpha$ -amino acids does not give absolutely quanti-

tative results in the case of cystine

Cystine and tyrosine can be quantitatively separated by means of absolute alcohol saturated with hydrogen chloride (Plimmer, Bio-Chem J. 1913, 7, 311-317).

When cystine is given with food, it is completely oxidised to sulphate and excreted as such (Rothera, J. Physiol 1905, 32, 175). Baumann and Preusse (Zeitsch physiol Chem. 1881, 5, 309, Ber 18, 258) found that when bromobenzene was administered to animals it was excreted as bromophenyleystein.

a compound crystallising in glittering needles or plates , m p  $180^{\circ}-182^{\circ}$ , and yielding a blue crystalline copper salt (C<sub>9</sub>H<sub>9</sub>BiSNO<sub>2</sub>)<sub>2</sub>Cu

Derivatives.—Cystine forms sparingly soluble salts with copper,  $C_6H_{10}O_4N_2S_2Cu$ , or mercury

$$\begin{array}{c} {\rm C_6H_8O_4N_2S_2Hg_2,HgCl_2}\,;\\ {\rm 2C_6H_{10}O_4N_2S_2Hg,HgCl_2,7H_2O}\,;\\ {\rm 2C_6H_8O_4N_2S_2Hg_2,Hg(NO_3)_2} \end{array}$$

With 30 p c nitric or hydrochloric acid (at room temperature) it yields compounds

$$\rm C_6H_{12}O_4N_2S_2$$
 2HNO3 and  $\rm C_6H_{12}O_4N_2S_2{\cdot}2HCl$ 

(Morner, Zeitsch physiol Chem. 1914, 93, 203-208) Benzoylcystine  $C_6H_{10}O_4N_2S_2Bz_2$ , m.p.  $156^\circ-158^\circ$ , crystallises in needles and forms a sparingly soluble sodium salt (Goldmann and Baumann, Zeitsch. physiol. Chem. 1888, 12, 254). Cystinephenylhydantoin is a well-defined

crystalline compound, and is used for identifying cystine. Cystine ethyl ester hydrochloride

S<sub>2</sub>[CH<sub>2</sub>·CH(NH<sub>2</sub>) CO<sub>2</sub>Et]<sub>2</sub>,3HCl

forms colourless needles, mp 185°, with decomposition (Friedmann, Beitr. Chem Physiol Path. 1902, 3, 1) Numerous acyl-derivatives of cystine have been described by Fischer and Suzuki (Zeitsch physiol Chem 1905, 45, 405; Fischer and Gerngross, Ber 1909, 42, 1485; Abdeihalden and Wybert, Ber 1910, 2449-2473) Of these dr-l-leucyl-l-cystine S<sub>2</sub>[CH<sub>2</sub> CH(CO<sub>2</sub>H)NH CO (CH(NH<sub>2</sub>)(<sup>2</sup>4H<sub>0</sub>] is a granular powder, turns yellow at 200°, and decomposes at a higher temperature, has  $\left[\alpha\right]_{D}^{20}$  -136 6°, or, when obtained in a crystalline

form, has  $[a]_D^{20}$ -111 4°, it gives a reddish-violet colour with copper sulphate, is precipitated by ammonium sulphate, and may be termed M A W

CYSTOGEN. Syn for hexamethylenetetramine

CYSTOPURIN. A compound of sodium acetate and hexamethylenetetramine

**CYTISINE**  $C_{11}H_{11}ON_2$ , prisms, m.p. 153°, b.p.  $218^{\circ}/2$  mm,  $[\alpha]_D^{17^2} - 120^{\circ}$ , readily soluble in water, alcohol, or chloroform, insoluble in ether and petroleum, is a diacid base forming well-crystalline salts It is found in the seeds of Cytisus laburnum, Ulese curopaus, Baptisia spp, Sophora spp, Genisla spp, and Anagyris fætida in amounts varying from 1 to 2 pc

Cytisine is a secondary-tertiary base Electrolytic reductions gives a base, tetrahydro deoxycytisine  $({}^{\circ}_{11}H_{18}N_2)$ . Hydrodic acid and phosphorus at 230° yields cytisoline  $({}^{\circ}_{11}H_{11}ON, \beta$ -cytisolidine  $({}^{\circ}_{11}H_{11}N)$  (this on reduction giving a cytisolidine  $({}^{\circ}_{11}H_{11}N)$ ), hydrocarbons, and ammonia The constitutions of  $\alpha$ - and  $\beta$ cytisolidines have been established by synthesis (Ewins, Chem Soc Trans 1913, 103, 97) B-Cytisolidine is 6 · 8-dimethyl quinoline, αcytisolidine the corresponding tetrahydro quino-Cytisoline, which on distillation with zinc-dust gives \(\beta\)-cytisolidine, is 2-hydiexy-6 8-dimethyl quinoline and has I cen synthesised 244, 20)

by Spath. On these grounds it has been suggested that cytisine is constituted by the fusion of three rings.

The following constitution is considered the most probable of the fourteen possibilities -

Cytisine is a violent poison. Cases of infantile poisoning from eating laburnum seeds are by no means infrequent Dale and Laidlaw (J Pharm Expt Therap 1912, 3, 502) have shown that the physiological action of this alkaloid is qualitatively and almost quantitatively indistinguishable from that of nicotine See Spath, Monatsh 1919, 40, 15, 93, Partheil, Ber 1890, 23, 3201, 1891, 24, 634; Freund and Friedmann, A 1901, 1 288, Freund, A 1904, 1. 263, Freund and Horkheimer, A 1906, 1 302, Freund and Gauff, Arch Pharm 1918, 256, 33

Methylcytisine  $C_{12}H_{16}ON_2$ , prismatic needles, m p 137°,  $\lfloor \alpha \rfloor_D - 221$ 6°, readily soluble in water and organic solvents, occurs in Caulophyllum thalictioides (Linn) (Power and Salway, Chem. Soc Trans 1913, 103, 191), it may also be obtained by methylating cytisine. The salts are crystalline

The physiological action of the base is similar to that of cytisine (q v), but the alkaloid is less potent

Caulophylline (Lloyd, Proc Amer Pharm.

Assoc 1893, 41, 115) is methyl cytisine

Anagyrine C15H22ON2 (1), amorphous, occurring along with cytisine in the seeds of Anagyris fortida, is possibly butyl cytisine (Klestermann, Diss Marburg, 1898, Chem Zentr 1899, 1 1130; Partheil and Spasski, Apoth Zeit 1805, 10, 903, Goessmann, Arch Pharm 1906, AJE

D.

DAGUERREOTYPE v PHOTOGRAPHY.

DAHL'S ACIDS. 1-Naphthylamine-4-7- and 4 6 -disulphonic acids v. NAPHTHALENE.

DAHLIA. Syn Hofmann's la (v. Triphenylmeihane Hofmann's violet, Pri-COLOURING mula (v. MATTERS).

DAHMENITE v Explosives

occurs in the seeds of Nigella damascena (Linn.) (Schneider, Pharm. Centrb 1890, 31, 173) and N. arıstata (Keller, Arch Pharm. 1908, 246, 1). Its constitution was established by synthesis (Ewins, Chem. Soc. Trans. 1912, 101, 544), and shown to be methyl-2-methylamino 3-methoxy benzoate ·

Keller's 'methyl damascenine' is identical with damascenine and 'damascenine acid' C<sub>9</sub>H<sub>11</sub>O<sub>3</sub>N, said by Pommerchne (Arch. Pharm. 1900, 238, 531) to be isomeric with damascenine, is obtained on hydrolysis of the alkaloid. Damascenine is mildly narcotic. A. J E.

DAMMAR RESIN, DAMMARYL, DAM-MARYLIC ACID v. RESINS.

DANAITE v. MISPICKEL.

**DANALITE.** A rare sulpho-silicate mineral, with the formula  $3R_2\mathrm{SiO_4}$  RS, where R represents Fe, Zn, Be, Mn, containing BeO about 14 pc. The flesh-red or grey cubic crystals are octahedral or tetrahedral in habit; sp gr 3.35-3.43; H.= $5\frac{1}{2}-6$  It occurs in granite at Rockport and Gloucester, in Massachusetts, and large crystals have been found in Connwall Closely allied is the tetrahedral-cubic mineral helvite, which contains about the same amount of beryllium, but with a preponderance of manganese and no zinc L J. S.

DANDELION ROOT. Taraxacı radıx, B.P; Taraxacum, USP. (Pissenlit, Fr.; Lowenzuhnwurzel, Ger) The root of the common dandelion, Taraxacum officinale (Weber) (Bentla Trim 159), is used in medicine for its mild laxative and tonic properties. It is administered either in the form of expressed juice or weak spirituous extract. The roots employed should

be collected in the autumn

The latex contains two crystalline bitter principles, laraxaxin and laraxacerin (Polex, Arch Phaim. 19, 50, Kiomayer, ibid. 105, 6). The root further contains inulin, 24 pc in October, according to Dragendorff (Sachsse's Farbstoffe, 125), but only 1.7 p.c in Maich, when, however, 17 pc of uncrystallisable sugar and 19 pc of levulin is present. The latter carbohydrate is soluble in cold water to a sweet solution, without any action on polarised light.

The leaves and stalks, but not the root of the dandelion, contain inosite (Marmé, Annalen, 129, 222)

G B

**DAPHNETIN.** Daphnetin is present in the form of its glucoside daphnin in the bark of the Daphne alpina (Linn) and D mezereum (Linn)

The concentrated alcoholic extract is digested with boiling water, filtered, and treated with lead acetate solution. The precipitate is removed, basic lead acetate added to the filtrate, the lead compound of the glucoside decomposed with sulphuretted hydrogen, and the solution evaporated (Zwenger, Annalen, 115, 8).

Daphnin  $C_{15}H_{16}O_{9},2H_{2}O$  crystallises in prisms, melting at 200°, and dissolves in alkaline solutions with a yellow colour. It is hydrolysed by boiling dilute acids with formation of daph-

netin and glucose.

C<sub>16</sub>H<sub>16</sub>O<sub>9</sub>+H<sub>2</sub>O=C<sub>0</sub>H<sub>6</sub>O<sub>4</sub>+C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>
Daphnetin C<sub>9</sub>H<sub>6</sub>O<sub>4</sub> forms pale yellow needles, is soluble in alkalis with a yellow colour, and gives a yellow precipitate with lead acetate solution. On acetylation, a diacetyl-derivative C<sub>9</sub>H<sub>4</sub>O<sub>4</sub>(C<sub>2</sub>H<sub>3</sub>O)<sub>2</sub> is obtained, and by means of ethyl iodide a diethyl ether C<sub>9</sub>H<sub>4</sub>O<sub>2</sub>(OEt)<sub>2</sub> is produced. When daphnetin diethyl ether is boiled with caustic soda solution, the diethyl other of daphnetinic acid

$${\rm (OEt)_2C_6H_2} \begin{array}{c} OH \\ CH = CH COOH \end{array}$$

is produced in the form of its sodium salt, which when ethylated gives the *triethyl* derivative:

By oxidising daphnetic acid triethyl ether triethoxybenzaldehyde is formed, which is readily converted into triethoxybenzoic acid. The

silver salt of this acid, when destructively distilled, gives pyrogallol triethyl other. Daphnetin is accordingly a dihydroxycoumarin, and has the following constitution:—

(Will and Yung, Ber 1884, 17, 1081).

It has been synthetically prepared by heating pyrogallol with malic acid in the presence of sulphuric acid (Pechmann, Ber 17, 933):

 $C_4H_6O_5+C_6H_3(OH)_3=C_9H_6O_4+CO_2+2H_2O+H_2$  and by the condensation of pyrogalic aldehyde with sodium acetate (Gattermann and Koebner, Ber 1899, 32, 287):

C<sub>6</sub>H<sub>2</sub>(OH)<sub>3</sub> CHO+CH<sub>3</sub> COOH=C<sub>9</sub>H<sub>6</sub>O<sub>4</sub>+2H<sub>2</sub>O
Daphnetin is a yellow colouring matter, and
gives with mordanted woollen cloth the following
shades .—

Chromium Olive-yellow Pale olive-yellow
Trn Iron

Very pale yellow Olive-black (Perkin and Wilson, Chem Soc Trans 1903, 83,

DAPICHO or ZASPIS. The South American name for the caoutchouc which exudes from the

roots of Hevea guianensis (Aubl) **DARWINITE** v DOMEYKITE

DATE. The fruit of the date palm, Phænix dactylifera A palm of the tropics of Africa and the East Indies It furnishes food to a large population in Egypt and Arabia

The fruit grows in bunches of from 15 to 20 lbs weight, and a tree usually produces about 20 bunches annually. The date palm also yields a sweet sap, which flows, when incisions are made in the trunk, almost continuously from November to March, the sap is clarified and boiled down, when a brown sugar, known as jaggery, is obtained (Bied Zenti 1884, 284).

Analyses of the flesh of dates give .

Dried dates, according to Konig, have the mean composition.

Water Protein Fat Sugar acid hydrates Fibre Ash 185 19 06 672 13 43 38 18

Among the 'soluble carbohydrates' is included about 3 p c of pentosans, while the date stone contains about 2.7 p c. of the same substances (Wittmann, Zeitsch landw Versuchs. Oesterreich 1901, 4, 131)

König gives, as the composition of the date stone:

Water Protein Fat free extract Fibre Ash 7.7 5 2 9 0 53 0 24 0 1.1

7.7 52 90 530 240 1.1

Dates are sometimes used as coffee substitutes, the whole fruit being occasionally employed, but more usually the stones only.

König gives as the composition of 'date-coffee':

Water Protein Fat Sugar hydrates Fibre Ash 6 6 5 5 7.9 2 2, 48.7 278 1.3 yielding about 14 p.c. of its weight to water H I.

#### DATHOLITE v DATOLITE.

DATISCA CANNABINA (Linn). This is a tall, erect herb, resembling hemp, belonging to the Datisceæ, and met with in the temperate and sub-tropical Western Himalayas from Kashmir to Nepal The root, known as Akalbir, has been extensively used in Kashmir and throughout the Himalayas as a yellow dyestuff chiefly on alum-mordanted silk; but the colouring matter appears to be present in the whole plant, for the leaves and twigs can also be employed for the same purpose

The leaves of the Datisca cannabina were first examined by Braconnot (Ann. Chim Phys [2] iii 277), who isolated from them a crystalline substance which he named datiscin. Somewhat later it was shown by Stenhouse (Annalen, 1856, 98, 167) that this compound is a glucoside, and when hydrolysed is converted into a sugar, together with a yellow colouring matter, datis-

cetin C15H10O6

Schunck and Marchlewski (Annalen, 278, 261), however, considered that the formula C<sub>15</sub>H<sub>12</sub>O<sub>6</sub> was preferable, and showed that, on fusion with alkali, datiscetin gives salicylic acid By the action of dilute nitric acid, a nitrosalicylic acid was produced, whereas with strong nitric acid, pieric acid, as already indicated by Stenhouse, was formed Datiscetin appeared to contain methoxyl groups, and was at that time considered by the authors to be most probably a xanthone derivative

Korczynski and Marchlewski (Ann. d. Akad Krakau, 1906, 95) have shown that pure datiscetin  $C_{15}H_{10}O_6$ , however, does not contain methoxy-groups It consists of yellow needles, methoxy-groups It consists of yellow needles, mp 268°-269°, soluble in alkaline solutions, with a yellow colour, and gives, with sulphuric acid, a yellow liquid possessing a green fluorescence When boiled with strong alkaline solutions, phenol and salicylic acid are produced

Tetraacetyldatiscetin C15H6O6(C2H3O)4 forms colourless needles, and melts at 138°, and

tetrahenzon/klatiscetin C<sub>15</sub>H<sub>6</sub>O<sub>6</sub>(C<sub>7</sub>H<sub>5</sub>O)<sub>4</sub> gives colourless needles, mp 190°–191°.

Leskiewicz and Marchlewski (Ber 1914, 47, 1599) by the action of bromine on datiscetin in boiling acetic acid solution obtained tribromphloroglucinol, and as a result suggest that this colouring matter is the 1 3 1'-trihydroxyflavonol -

A substance indeed of this constitution in the form of its trimethyl ether has been synthesised by Bargellini and Peratoner (Gazetta, 1919, 49, 64) by the usual methods of flavonol synthesis Thus 2 hydroxy 4:6-dimethoxy phenyl methyl ketone (phloracetophenone dimethyl ether) condensed with salicylic aldehyde methyl ether yields 2 hydroxy 4:6:2' trimethoxy phenyl styryl ketone I :-

I. MeO 
$$\stackrel{\mathrm{OH}}{\longrightarrow}$$
 CO—CH : CH— $\stackrel{\mathrm{OMe}}{\longrightarrow}$  Vol. 11.— $T$ .

which by means of dilute alcoholic hydrochloric acid gives 5 7:2' trimethoxy flavanone II :-

The iso-nitroso derivative of this latter when heated with hydrochloric acid in acetic acid solution gives the trimethoxy flavanone

crystallising in pale yellow needles, mp 158°-160°, and giving with sulphuric acid an intensely green fluorescent solution. This is probably a trimethyl ether of datiscetin

Datiscin  $C_{21}H_{24}O_{11}$  (S and M), the glucoside crystallises in silky needles, which melt at about 190°, and dissolve in alkaline solutions with a yellow colour Hydrolysis with dilute sulphuric acid yields datiscetin and rhamnose, not datiscetin and glucose, as stated by Stenhouse (lc)

The dyeing properties of datisca root on wool are very similar to those of old fustic. It is, however, slightly inferior in dyeing power, On cotton it dyes more like quercitron bark, in so far that the olives obtained on iron mordant are bluer in shade, as though some tannin matter It appears to have about the were present same colouring power as quercition bark, but gives a brighter yellow with aluminium mordant Applied to wool, it differs from querestron bark by giving with chromium mordant an oliveyellow which is not of such a reddish tinge, and with tin mordant a bright yellow instead of an orange It is an excellent dyestuff, and worthy to rank with the natural yellow mordant colouring matters at present in use (Hummel and Perkin, J Soc Chem Ind 1895, 14, 458) A G. P.

DATISCIN v Datisca Cannabina. DATISCETIN v DATISCA CANNABINA

**DATOLITE.** A native boro-silicate of caleium HCaBSiO<sub>5</sub>, containing B<sub>2</sub>O<sub>3</sub> 218 p.c. Monochnic crystals, with a profusion of small brilliant faces, many with five-sided outlines, are of common occurrence; these are usually transparent and colourless, or with a faint greenish tinge, sp gr 30. The mineral most greenish tinge, spgr 30 often occurs associated with zeolites in the amygdaloidal cavities of basaltic rocks, e g in the neighbourhood of Edinburgh and Glasgow, and at Bergen Hill and West Paterson in New It, however, presents several other Jersey modes of occurrence, viz in granite at Baveno in Piedmont, in diorite at Rosskopf in Baden; in gabbro with copper ores at Monte Catini in Tuscany; in 'limurite' (an axinite-pyroxenite) at Dundas in Tasmania; in serpentine in the Lizard district in Cornwall, in calcite veins in sandstone at Sonthofen in Bavaria; in veins of silver ore at Andreasberg in the Harz Mountains; in the Lake Superior copper mines (here in a compact, opaque, white form resembling unglazed porcelain); in beds of magnetite in gness at Arendal in Norway L. J. S.

21

# DATURA STRAMONIUM v TROPEINES.

DAUCINE C11H18N2, colourless, oily base, b.p.  $240^{\circ}-250^{\circ}$ , occurring in carrot leaves (Pictet and Court, Ber 1907, 40, 3771), together with pyrrolidine

**DEAD DIPPING.** The process of producing a pale-yellow dead surface on ornamental brass-The brasswork, after the final stamping. with its adhering black scale from the annealing oven, is placed in dilute nitric acid until the scale is detached, when it is removed and washed with water. It is next plunged into stronger nitiic acid, and when its surface is covered with minute gas bubbles, it is washed in a solution of argol and diled in hot saw-

**DEAD OIL.** The higher boiling fractions of shale oil, from which the greater portion of the solid paraffin has been crystallised out

**DECACYCLENE**  $C_{36}H_{18}$   $\Lambda$  hydrocarbon obtained together with fluorocyclene  $C_{48}H_{28}$ and chlorenc C<sub>48</sub>H<sub>26</sub> by heating accnaphthene with lead oxide at 370°-380° (Dziewonski and Suknarowski, Ber 1918, 51, 457)

n-DECAHEXANEDICARBOXYLIC ACID CO<sub>2</sub>H[CII<sub>2</sub>]<sub>16</sub>CO<sub>2</sub>H Prepared by the electrolysis of an aqueous solution of the potassium salt of the monocthyl ester of suberic acid, and hydrolysing the diethyl ester thus formed (Brown and Walker, Annalen, 261, 125), m p 118°.

DECAMETHYLENEDICARBOXYLIC ACID CO2H [CH2]10 CO2H Prepared by heating the methyl ester of bromundecylic acid with alcohol and potassium cyanide, and hydrolysing the product with alcoholic potassium hydroxide (Nordlinger, Bei 1890, 2357), mp 124 5°-125 5°

DECATOIC ACIDS C<sub>10</sub>H<sub>20</sub>O<sub>2</sub>.

1 Capru acid CH<sub>3</sub>(CH<sub>2</sub>)<sub>8</sub>COOH is found as a glyceride in the butter of the cow and goat, and in cocoa-nut oil, in association with caproic and caprylic acids Many fusel oils also contain it (Fischer, Annalen, 118, 307, Gimm, *ibid* 157, 264, Rowney, *ibid* 79, 236), and it is found as isoamyl capitate in certain Hungarian wines, and is formed by the oxidation or distillation of oleic acid (Redtenbacher, ibid 59, 45), and by heating octylacetoacetic ester with alcoholic potash (Guthzeit, ibid 264, 5). A white crystal-line substance of m p 31 3°; b p 268 4° (corr), sp gr 0 8858 at 40°/4°, possessing a goat-like smell; insoluble in cold water, but slightly soluble in boiling water

# 2 Dusobutylacetic acid

# [CH(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>]<sub>2</sub>CH COOH

Obtained by heating dissobutylmalonic acid (Bentley and Perkin, Chem Soc Trans 1898, 62), bp 225°-230° (730 mm)

DECHENITE v VANADIUM

DECOIC ACID, CAPRIC ACID v. DECATOIC

DECOLOURISING CARBONS. This term has been applied to nearly pure forms of carbon, possessing remarkable decolourising powers, and manufactured only within the last ten In this article it will be convenient to groups these special carbons separately from those which have been employed as decolorants for over a century

# (a) Vegetable Charcoals

The decoloursing property of wood charcoal was first observed by Lowitz in 1791, and was used for treating beet-syrup An improved charcoal, produced by washing and grinding, was patented by Louis Constant in 1812, and employed in the refining of sugar. Wood charcoal is manufactured by burning wood in special ovens, or in heaps, air being excluded Oven charcoal has the composition: carbon, 81 p c, hydrogen, 4 p c; oxygen and mineral matters, 15 p c Charcoal burnt in heaps has the composition carbon, 90 pc; hydrogen, 2:5 pc, oxygen and mineral matters, 75 p.c. Klasen and Bergh (Chem. Zeit 1910) state that the oven-charcoal absorbs 15 pc. by weight of oxygen at 100° C. The decolourising power of most vegetable charcoals is very slight, but superior qualities are obtained from cherry stones, cocoanut, and certain grasses wood be treated with certain chemicals before being carbonised, the resulting charcoal is a very efficient decolorant (see group (c) below)

# (b) Animal Charcoals.

In 1811, Figurer proved that bone charcoal was a far more effective decolorant than wood In 1815, Messrs P and J Martineau charcoal patented the use of bone charcoal for refining and clarifying sugar solutions In 1817, De Cavaillon patented the calcining of bone charcoal after it has been used as a decolorant, rendering it 'superior to new charcoal, and the operation may be repeated any number of times with equal effect' Up to the year 1828, bone charcoal was used in the form of fine powder, mixed with the liquid to be decolourised, and, finally, separated therefrom by filtration Payen, in 1828, introduced an important improvement, the 'bone-char' being produced in a granular form, permitting the syrup (or other liquid) to be filtered through the bone-char in 'char cisterns' or 'char filters,' first designed by Dumont. The granular charcoal acted as efficiently as the powder, and avoided the tioublesome filtration to separate the powder from the liquid. The granular form was also more suitable for burning in order to restore its activity after it becomes charged with organic impurities absorbed from the liquid under treatment Dumont's 'char filter' was of the open type, operating by gravity. The modern vessels are closed, and operate under pressure, and the same liquid may pass through two or even three filters, if necessary

Formerly, bone-char was employed for decolourising both beet and cane juices, in order to produce white sugar direct from the plant. Less costly methods of purifying these juices have led to the abandonment of bone-char in the raw sugar factory, and its application in the refinery

Manufacture of Animal Charcoal The bones employed should be carefully selected, hard, and free from extraneous matter Whale and fish bones are not of a suitable character, as they yield a soft char Bones which have been exposed to atmospheric action for a long time, or which have been buried in the ground, cannot be successfully employed, as, owing to the alteration in their composition, they produce a char deficient in carbon

Before proceeding to carbonise the bones, the fat is removed by boiling, or by means of a suitable solvent such as benzene By the first-named treatment from 4 to 5 p c of fat is extracted, and by the second from 8 to 12 p c

The crushed bones are carbonised in apparatus similar to that used in a gasworks, in vertical or horizontal iron retorts of a round, oval, or D section, the latter by preference The length of the horizontal retort is usually from 8 to 10 feet, the long diameter being 18 inches and the short diameter of the oval retort 12 inches There are usually five or seven retorts in each The retorts are connected in the same way as gas retorts, with a hydraulic main, and this again with condensers or scrubbers filled with coke, in which the bone oil separates, and from which the gases are exhausted The gases are then forced through a series of washers containing water to remove the ammonia, the residual gases, which are employed for heating and lighting purposes, finally passing into a gasholder Charges of about 2½ cwt of bones are carbonised in vertical, and 5 cwt in horizontal retorts, the operation taking about 6 to 8 hours with the former, and 8 to 10 hours with the latter The bone oil, collected in the condenser, amounts to from 3 to 5 p c on the bones carbonised, and the ammonia in the ammoniacal liquor is equal to about 8 pc of ammonium sulphate, into which it is converted in the usual way the bones are completely carbonised, the charcoal is removed from the retorts and cooled in strong sheet-iron canisters, which are at once covered with closely fitting lids and luted round the edges, either with charcoal paste, or a water The char, when quite cold, is crushed in a suitable mill (any iron it may contain being removed by passing over the poles of an electromagnet), and then sifted into various sizes to suit the requirements of the sugar-refiner is usually moistened before guinding, to keep down the dust Good bones yield about 65 p c of char, but from 20 to 30 pc of this is dust, which fetches a lower price than the larger grist

The following is the approximate composition of a good sample of new bone charcoal

Carbon		10 51
Calcium and magnesium pho	s-	
phates, calcium fluoride, &c		81 21
Calcium carbonate		7 30
Calcium sulphate		0 17
Ferrie oxide		0.12
Silica .		0.34
Alkaline salts		0.35
		100 00
Moisture originally present		8.00

Space occupied by one ton, 48 cubic feet Weight of char left on sieves of various degrees of fineness -

Above 10	holes	to linear	ınch		0
10 to 20	٠,	,,	,,		28
20 to 30	,,	,,	,		32
30 to 40	,,	••	,,		27
40 to 50	,,	• ,,	,,		11
Through 50	,,	,,	or	dust	2

New charcoal of good quality should, in the dry state, contain not less than 9 and not more than 11.5~p~c of carbon The silica should not exceed 0.5~p~c, the oxide of iron 0.15~p.c, the calcium sulphate 0.2 p.c., and the moisture Its weight should not exceed 52 lbs per cubic foot It should be of a dull black colour, and, when incinerated, leave an ash of a uniform white or cream colour, the presence of grey or reddish particles indicates that the sample has been mixed with old charcoal When brought in contact with the tongue, it should adhere to it somewhat strongly The size of grain depends upon the use to which it is to be applied grain charcoal is preferable for strong liquors, such as that used for washing loaf sugar Refiners, as a rule, prefer small grain, if, however, it be very small, it impedes the passage of liquor and also gives considerable trouble in washing (har may be used over and over again, and sometimes lasts several years, being revivified from time to time by reburning in special kilns (see Sugar) When, by continued use, it has lost its power of removing colour, it is known as spent char, and is then used as a manure, generally after treatment with sulphuric acid, so as to form superphosphate of lime

The carbon always contains a certain proportion of nitrogen, amounting to about onetenth of its weight, also a minute proportion The nitrogen present decreases of hydrogen whilst the bone charcoal is being used for sugarrefining

When char is repeatedly reburned, it becomes less porous and shrinks in volume, so that a ton, which measures 48 to 54 cubic feet when new, may be reduced to 28 cubic feet after being reburned many times, in other words, its apparent density may be nearly doubled

Wallace has, however, shown that the real sp gr varies but little, thus, a new char, occupying 50 6 cubic feet per ton, or having an apparent sp gr of 071, has a real sp gr of 2 822, whilst a moderately old sample, occupying 35 cubic feet per ton, or having an apparent sp gr of 1 03, has a real sp gr of 2 857, or only a trifle over that of the new.

Another proof that char loses its porosity by long-continued use and reburning, is afforded by the fact, pointed out by Wallace, that dry new char will absorb from 80 to 100 pc of its weight of water, whereas old char will only retain

from 30 to 45 p c

Properties —Although the decolourising property is the most important, bone-char absorbs other organic and mineral substances from their Tilhot was the first to prove that the absorption of organic matter increases with the temperature, and Kohlrausch first demonstrated that the absorbed colouing matters remain chemically unchanged, and can be extracted from the charcoal by washing it with ammonia, in which the colouring matters dis-It was proved by Graham that various mineral substances are removed from solution by animal charcoal, thus the lime is taken up from lime water, and metallic salts are absorbed from their solutions in water. According to Chevallier, lead nitrate and acetate are completely removed by animal charcoal. Weppen has shown that this action extends to a great variety of metallic salts, including cupric, zinc, chromic and ferrous sulphates, nickel, cobalt, silver, mercurous and mercuric nitrates, tartar ometic, stannous chloride, and ferric acctate

As the result of a number of experiments on the absorption of different salts by animal charcoal, Bodenbender arrived at the following conclusions

1 The power of absorbing salts is a physical

- 2 A given weight absorbs a larger proportion of salts from a concentrated than from a diluted solution, on the other hand, the proportion absorbed from a constant quantity of salts is more considerable when this quantity is in a dilute, than when it is in a concentrated, solution.
  - 3 The presence of sugar has only a slight

influence on the absorption of salts

4 The salts of potassium are absorbed in smaller proportion than those of sodium

5 Among the salts experimented with, the amount taken up is in the following order, beginning with that least absorbed · potassium chloride, sodium chloride, potassium nitrate, sodium nitrate, potassium acetate, sodium acetate, potassium sulphate, sodium sulphate, magnesium sulphate, potassium carbonate, sodium carbonate, sodium phosphate

6 A chemical action of charcoal has been observed with respect to some carbonates, oxalates, and other salts, being brought about by the presence of calcium sulphate and phosphate

in the charcoal

7 Charcoal, saturated with one salt, is capable, within certain limits, of withdrawing another salt from solution

8 There is less absorption of a salt when the contact is brief than when the contact is prolonged. The difference due to time of contact ceases when the contact has lasted some hours

Investigations by W. B. Clark (J. Soc. Chem. Ind 1912, 262) indicate that the decolourising power of bone-char depends very largely on the quantity of mineral matter and reducing sugar in the liquor treated in the sugar refinery first portion of syrup leaving the char filter contains only about 60 pc of the reducing sugar, and 30 p c. of the mineral matters originally in After these have been absorbed by solution the char, they are subsequently set free, so that the final portions of syrup contain even more reducing sugar and mineral matters than were originally in solution At this stage there is a gradual decrease in the decolourising power of After the syrup inlet has been closed, the char hot water is passed through the char to displace In this wash-water, the reducing the syrup sugars and mineral matters are present in con-This washing treatment siderable quantities restores the normal activity of the char, although the washed char still contains much organic matter, which can be extracted by means of caustic soda solution. Clark remarks that the behaviour of char closely resembles diffusion, the char itself acting as a dialyser, permitting the crystallisable sugar to diffuse through it more rapidly than the other substances present. Bonechar attains its maximum efficiency after being used, washed, and burnt several times. porosity is thereby reduced, as also is the percentage of organic matter other than carbon, and the percentage of calcium carbonate is reduced from about 9 to 3 p.c. The residual calcium

exists as a basic phosphate, which is partially decomposed by heating the char with dilute acids, causing an increase in decolourising power

New char exerts a slight oxidising action on colouring matters which are not absorbed by Degener and Lack state that freshly ignited bone-char, moistened with water, and exposed to light and air, produces a perceptible quantity of hydrogen peroxide, and is thus rendered more active than the untreated char If milk of lime be sprinkled over the char, calcium peroxide is produced

The high decolourising power of bone-char relative to wood charcoal was formerly supposed to be due to the presence of 5-7 p c. of nitrogen in the former, and only traces in the latter; but this view has now been abandoned in favour of a physical, or purely mechanical, action

Charcoal prepared from flesh has about the same decolourising efficiency as that made from Blood charcoal is less efficient charcoal is the subject of a patent by Adler, 1913. The finely minced fish is boiled with water, carbonised, and the resulting charcoal washed with hydrochloric acid.

# (c) 'Decolourising Carbons'

As stated under (a), very active carbons can be obtained from various woods when these are chemically treated before being carbonised The earliest patent appears to be that of R von Ostrejko, in August, 1900, for treating wood powder with calcium chloride or magnesium chloride before carbonising, and subsequently treating the carbon with acid to extract the lime In a later patent, October, 1900, he employs calcium acetate or chloride, and starch mucilage Subsequent patentees employ sulphuric acid at 200° C (Halse, 1902); earthy lignites treated with kaolin (Goldstein, carbonaceous matter of animal or vegetable origin is ignited in the presence of chlorine gas (Molenda and Wunsch, 1912); peat treated with hydroxides or carbonates of potassium, sodium, calcium, or barium (Kruszewski, 1912); wood treated with zinc chloride (Wunsch, 1913); wood treated with calcium oxide or carbonate (Bonnard, 1914)

Two very successful carbons are the Dutch

'Norit' and the German 'Eponit.'

'Norit' is a fine black powder, containing 94-96 p c carbon, and 4-6 p c mineral matters. It possesses about 30 times the decolourising power of bone-char, and about 100 times that of wood charcoal The activity increases in the presence of acids, but diminishes in the presence of alkalıs.

Eponit' is also a fine powder, containing carbon, 86 5 p c.; mineral matters, 4·1 p c; and moisture, 9.4 pc Experiments by Strohmer indicate that it possesses 10 times the decolourising power of bone-char, and is not influenced by the acidity or alkalinity of the liquid treated. From his chemical and microscopical examina-tion he concludes that 'Eponit' is a carbonised vegetable substance, probably wood.

More recently, a Committee of the Royal Society of London has investigated this carbon, and attempted to reproduce it by carbonising various kinds of wood. The conclusion arrived at was that 'Eponit' is made from a mixture of coniferous wood, such as pine or cedar, and angiospermous wood, such as poplar or willow.

Following Ostrejko's patent (see above), the Committee carried out the following experiments:—

(a) A mixture of pine and willow woods was treated with calcium acetate and staich, carbonised at a white heat, and the resulting carbon boiled with hydrochloric acid, washed with water, and heated to a red heat in a retort.

(b) Cedar wood was treated with calcium acetate alone, before carbonising, and the same

subsequent treatment

Both (a) and (b) proved superior to 'Eponit' Further experiments with cedar and willow were conducted as follows: (a) untreated, (b) treated with milk of lime; and (c) with calcium acctate before carbonising at a white heat, subsequent treatment as before. The following results were obtained (a) both charcoals were useless for decolourising purposes, (b) and (c) both woods gave very active charcoals, equal to 'Eponit,' (c) being slightly superior to (b). The Committee concluded that the type of wood has little or no influence on the result, and that probably any porous wood, after similar chemical treatment, would yield an active charcoal. Further, that the treatment with milk of lime is less costly than that with calcium acctate, and the resulting carbon is practically equal in decolourising power.

Pellet (Bull Assoc Chim Sucr 1916, 33, 220 227) has studied the relative decolourising powers of numerous carbons and charcoals relative to 'Epomit' taken as standard. He employed a dilute solution of molasses, previously clarified with basic lead acctate and the clear solution acidified with (a) hydrochloric, and (b) sulphurous acids. As some of the carbons gave slightly different values in the presence of these different acids, the two following series were obtained, in which 'Epomit' is the most

efficient —

(a) With hydrochloric acid Eponit, 1. Littoral, 2, Flandrac, 3, Flaming, 4; special carbon for wines, 5, bone-char, 6, special carbon, 7, bone-char washed with acid, 8, cherry stone carbon, 9.

(b) With sulphurous acid Eponit, 1, Flandrac, 1, Littoral, 2, special carbon, 3, special carbon for wines, 5, bone-char washed with acid, 6, Flaming, 7; bone-char, 8, cherry-stone carbon, 9

'Noist' and 'Eponit' resemble bone-char in absorbing organic and mineral matters from the solutions treated by them, and thus gradually losing their activity. This can be restored by two methods. (a) carbonising, so as to destroy the organic matter, and (b) by boiling with dilute caustic soda, which dissolves the organic matters, after which the carbon is separated in a filter-press, and washed with hot water until free from alkali

In the former method, a special type of kiln must be employed (Wijnberg's Patent, 1916). The second method was patented by Wijnberg and Sauer in 1911 and 1912, and requires no special plant, but appears to have been applied to bone-char many years previously. In another patent by Wijnberg, in 1915 Norit is employed to facilitate filtration of sugar juices through cloth-filters, owing to its power of absorbing gums and pectins.

# (d) Mineral Carbons.

Coal, or other bituminous substance, is impregnated with an alkali, or alkali carbonate, and some time later with an acid, or first with acid, and later with alkali, before ignition in retorts (Lotz, D. R. P. 1911). 'Culm,' a bituminous mineral present in the alum slate of Sweden, is dry distilled, the residue leached with acid, and washed with water. It can be regenerated by washing and re-heating (Hellsing, Eng. Pat. 1913). Coal is heated with excess of anhydrous zine chloride in an open vessel until the chloride begins to volatilise. The mass is cooled, crushed, washed in a filter press with dilute hydrochloric acid, and finally with water (Wunsch, Eng. Pat. 1914). The residue from shale-oil distillation is carbonised (Catlin, U.S. Pat. 1917).

#### (e) Carbons produced from Industrial By Product

Clacher claims that an active carbon can be prepared from 'bagasse, 'or crushed sugar cane, after the extraction of the juice (Int Sugar Journ 1914, 64 66) The Pan Chemical Co, USA, manufacture an active carbon from the by-product obtained in the manufacture of salts of tartaric acid from argol This residue contains 50 pc moisture, and 50 pc crude fibre with lime and sand It is dried and heated in retorts (U.S. Pat. 1912) Mathéus (Papierfabrikant, 1911, 9, 1435) evaporates sulphite waste lyes with milk of lime to a thick syrup, and finally in flat vessels, until it solidifies, forming a brittle mass This is powdered and heated in a closed vessel

'Black ash residue,' obtained by evaporating the alkali separated from wood pulp, consists mainly of carbon with 1–5 p.c. of ash. The percentage of ash can be reduced to about 0.8 p.c. by washing the residue with hydrochloricacid. The washed carbon has ten times the decolourising power of bone-chai (McKee, U.S. Pat. 1914).

DÉGRAS, or SOD OIL, is the waste fat obtained in the chamoising process, and is used for currying purposes, i e dressing bark-tanned or chrome-tanned leather The skins which are to be converted into chamois leather, are first 'limed,' the hair is then removed and the unhaired hides are 'drenched,' i e placed in a sour bath,' in which the free lime is neutralised through the development of an acid fermenta tion, caused by Bacillus furfuris The skins are then stretched and well rubbed with whale oil, or cod (liver) oil, in the United States with menhaden oil, and worked in 'stocks,' so as to become thoroughly saturated with the oil The skins are then exposed to the air, and the process of rubbing with oil and stamping in the stocks is repeated until enough oil has been absorbed By exposure to the atmosphere, a portion of the oil becomes changed, and is most likely converted into the glycerides of 'oxidised' acids In order to promote the oxidation of the oil, the skins are heaped together in a warm room, and covered carefully with canvas, so as to keep the generated heat in the heap whereby a large proportion of the oil is converted into oxidised 'oil The bulk of the oil is expressed from the skins by pressure, the remainder of the

oil retained by the skins being removed by one of the two following methods, known respectively as the English and German method, or the French method In the English and German method, the skins are washed with alkaline lyes; the emulsion thus obtained is acidified with sulphuric acid, when the fatty matter separates and is skimmed off This fatty substance forms the 'sod oil' of commerce. In the French method, the skins are 'stocked,' aired, and 'fermented' for a shorter period than in the English or German method, so that a large proportion of the oil can be obtained from the skins by steeping them in warm water, and wringing or pressing in hydraulic presses. The oil thus obtained is termed 'première torse moellon' The oil still retained by the skins is recovered by washing with alkaline lyes, as is done in the English and German method, and the fatty matter so recovered is usually added to the moellon An addition of other oils is also made to moellon, and for this reason the commercial product contains less fibre and mineral matter than sod oil

The most characteristic constituents of dégras

are the oxidised fatty acids

According to Jean, the characteristic oxidation product is a resin-like body melting at This substance is insoluble in petroleum spirit, but dissolves in alkaline solutions, from which it is again precipitated on the addition of acid. It has been termed dégras former by Simand, and has been shown by Fahrion to be composed of a mixture of hydroxy acids and anhydrides

Dégras is largely admixed with fish oils, blubber oils, tallow of low quality, &c ing to trade custom, such products are considered as a genuine product. On the other hand, the numerous substitutes of dégras, or artificial degras, which occur in commerce, consist of highly adulterated dégras, the adulterants being mineral oils, wool fat, or blown blubber oil, and 10sin Analyses of dégras and sod oils of different kinds are given m J Soc Chem Ind 1891, 10, 557, 1892, 11, 639, and J Amer Chem Soc 1899, 21, 291

DEHYDROTHIO-p-TOLUIDINE, DEHYDRO-THIO-p-TOLUIDINE SULPHONIC ACID, vPRIMULINE AND ITS DERIVATIVES

DEHYDROTHIO-m-XYLIDINE

$$\begin{array}{c|c} CH_3 & \\ \hline -N & \\ -S & C \\ \hline -NH_2 \\ \hline \end{array}$$

is prepared by heating m-xylidine with sulphur until hydrogen sulphide ceases to be evolved (cf. Dehydrothio-p-toluidine, Ait Primuline) The excess of m-xylidine is removed by distillation and the residue treated with hydrochloric (m p. 121°), formed simultaneously, is insoluble, m.p. 107°, b p  $283^{\circ}/14$  m.m. water, readily soluble in hot alcohol (Anschutz and Schultz, Ber. 1889, 22, 582; Paul, Zeitsch angew Chem. 1896, 9, 679).

DELPHINE BLUE v OXAZINE COLOURING

DELPHINIDIN. DELPHININ, v. ANTHO-

**DELPHININE** C<sub>31</sub>H<sub>49</sub>O<sub>7</sub>N is one of a series of ill-defined poisonous alkaloids, including delphisine, delphinoidine, staphisagroine, and staphisagroidine, found in stavesacre seeds (Delphinium Staphisagna (Linn). Official in the B. P and U.S. P. (Kora-Stojanoff, Chem. Zentr 1890, [n ] 628, and Ahrens, Ber 1899, 32, 1581, 1669) Delphinium consolida and D. ajacis, according to Keller (Aich Phaim 1910, 248, 468; 1913, 251, 207), also contain alkaloids. From various Delphinium spp Heyl has obtained 'delphocurarine,' an amorphous mixture of bases, including a crystalline substance, mp 184°-185° (Chem Soc Abstr 1903, i. 650) 'Delphocuratine' resembles curarine in

physiological action (Lohmann, Pflug. Arch

1902, 92, 398)

DELPHINIUM CONSOLIDA. Delphinium consolida is a common European plant belonging to the Larkspur family; its name refers to its powers, real or imaginary, of healing or consolidating wounds. The blue flowers were examined by Perkin and Wilkinson (Chem Soc Trans 1902, 81, 585) to determine if these yield the same colouring matters as those previously isolated from the flowers of the *D* zahl (bhd 1898, 73, 267) The presence of kaempferol only could, however, be detected For its isolation an aqueous extract of the flowers was digested at the boiling-point with addition of sulphuric acid, and the brown resinous product which separated on keeping, extracted with alcohol and the extract evaporated to a small bulk Addition of ether to this solution caused the precipitation of resinous impurity, and on evaporating the ethereal liquid a semicrystalline residue of the clude colouring matter was obtained The product was crystallised from dilute alcohol, converted into acetyl derivative, and this after purification retransformed into colouring matter in the usual manner The yield was approximately 1 p c

Kacmpferol  $C_{15}H_{10}O_6$  consists of yellow needles, mp  $276^{\circ}-277^{\circ}$ , soluble in alkaline solutions with a yellow colour. These liquids on exposure to air are slowly oxidised with

development of a brown tint

Tetra-acetylkaempferol  $C_{15}H_6O_6(C_2H_3O)_4$ , when crystallised from methyl alcohol forms colourless needles, and when heated commences to melt at 116° and becomes completely fluid at 120° On further heating, however, gradual solidification ensues and the product subsequently melts at 181°-182° This peculiarity of acetylkaempferol, which is not apparent when the substance is crystallised from ordinary alcohol, affords a convenient method for its detection

Tribromkaempferol $\rm C_{15}H_7Br_3O_6$  forms yellow needles, m p  $275^{\circ}\text{-}277^{\circ}$ ; haempferol sulphate needles, mp 275-271; https://doi.org/10.1016/html.006/hg804, orange-red needles; kaempferol hydrodude  $C_{15}H_{10}O_{8}HI$ , needles, and monopotussium kaempferol  $C_{15}H_{9}O_{8}K$ , orangeyellow pusmatic needles.

Kaempferide, a monomethyl ether of kaempferol, exists in Galanga 100t, and a monomethyl ether, apparently distinct from this latter, has been isolated from the berries of the Rhamnus catharticus The glucosides at present known of kaempferol are robinin, which occurs

in the flowers of the Robinia pseudacacia, kacmp-feritrin, found in the leaves of the Indigofera arrecta; and kaempferin, which has been isolated from senna

Fused with alkali, kaempferol gives phloro-

glucinol and p-hydroxybenzoic acid

To kaempferol v Kostanecki assigned the constitution of a trihydroxyflavonol (Ber 1901, 34, 3723) —

and its synthesis was subsequently effected by v. Kostanecki and Tamboi (Ber 1904, 37, 792)

2-Hydroxy-4·6 4'-trimethoxychalkone (1) when digested with boiling dilute alcoholic sulphuric acid gives 1 3 4'-trimethoxy flavanone (2) —

and the latter by means of amyl nitrite and hydrochloric acid is converted into iso-nitroso-1 3 4'-trimethoxyflavanone (3).

When a solution of this substance in acetic acid is boiled with 10 pc sulphuric acid 1-3-47-trimethoxyflaronol (4) is produced, from which by the action of hydriodic acid kacipperol (5) is readily prepared—

(3) 
$$CH_3O$$
  $CH_3O$   $CH_3O$ 

Kaempferol possesses well-defined dyeing properties, and gives with mordanted woollen cloth the following shades which closely resemble those given by morin:—

Chromium
Brownish-yellow
Tin
Lemon-yellow
Deep olive-brown.

It is also present in the Impatiens balsamina (Chantili Pass), the Eightrina stricta (vernacular name 'Kon kathet') (Perkin and Shulman, Chem Soc Proc 1914, 30, 177), the berries of the Rhamus catharticus and together with quercetin, both apparently as glucosides, in the flowers of the Prunus spinosa (Perkin and Phipps, Chem Soc. Trans 1904, 85, 56) For the separation of the two colouring matters a fractional crystallisation from acetic acid was

employed, kaempferol in these circumstances being the more sparingly soluble A G P

**DELPHLEGMATOR.** An apparatus used in fractional distillation. The term is also comployed in distillation for an apparatus for freeing alcohol from water (phlegma)

DELTA ACID. Disulpho acid S 1-Naph-

thylamine-4 · 8-disulphonic acid

DELTAPURPURÎN v. Azo- colouring matters.

DENATURANTS. This term is applied to substances which are added to articles usually subject to revenue imposts, and primarily intended for human consumption (as spirits, tobacco, &c) in order to render such articles unfit for human consumption, but without impairing their efficacy as raw material for use in various arts and manufactures. Articles thus denatured 'may safely (from a revenue point of view) be released from revenue control without payment of duty

The principles underlying the choice of suitable denaturants are mainly as follows—

(a) They should be inexpensive, so as not to

add materially to the cost of the denatured article,
(b) Then use should cause the minimum
amount of inconvenience to the manufacturer,
both in regard to his processes and the necessary
revenue supervision involved,

(c) The denaturant should not be incompatible with the process of manufacture or the subsequent use of the denatured article,

(d) It should be of such a nauseous and repugnant character as to remove any tempta tion to would-be consumers of the denatured article, and

(e) The denaturant should have such a relation to the original article both in quantity and character as to preclude its removal from the mixture by any process which could be profitably carried out on a commercial scale, and it should be capable of being readily detected even if present in very small quantity. In the United Kingdom the dutable articles

In the United Kingdom the dutiable articles allowed to be denatured are tea, tobacco, wine, and spirit, the last being by far the most important, both by reason of the large amount of revenue derived from it, and the great variety and extent of its applications in the arts and manufactures

Tea, before being admitted for consumption in the United Kingdom, is examined by the Customs under sect 30 of the Sale of Food and Drugs Act, 1875, and, if found satisfactory, is passed on payment of duty. but if otherwise, it is refused admission as 'tea' Since, however, it still possesses a certain commercial value as a raw product for the extraction of theme or caffeine, the Commissioners of Customs and Excise usually permit it to be admitted for this purpose duty free, after it has been effectively denatured in such a way as to make it unsuitable for use as a bevorage, but without interfering with its efficacy as a source of caffeine The same applies to those teas which have been accidentally damaged and are voluntarily withdrawn by the importers from sale as 'tea

In the case of tea, the denaturing process consists in grinding and mixing the tea with lime and asafectida in the proportions of 1 part of asafectida and 100 parts of lime to 1000 parts of tea

Tobacco is dutied on the whole leaf, of which certain portions, as the midribs or so-called 'stalks,' are by manufacturers treated as 'stalks,' are by manufacturers treated as waste or 'offal,' and the same applies to the unused portions of manufactured tobacco, known as 'horts' or 'smalls' On these the duty which has been paid is returned as 'drawback' to the manufacturer on condition, either that the tobacco offal is surrendered to the Government for destruction, or (if intended for such purposes as the manufacture of nicotine, insecticides, sheep dips, &c ), that it is denatured according to certain prescribed formulæ, and under the supervision of revenue officers. Denaturing is equally necessary in the case of offal from tobacco manufactured in bond (although no duty has been paid nor drawback given), in order to prevent its fraudulent use as a substitute for duty-paid tobacco

The following formulæ have been officially approved in the United Kingdom for denaturing certain preparations having tobacco as a basis

(a) Sheep wash To every 100 lbs. of tobacco are added 10 lbs of blue vitriol, 15 lbs. of common salt, 2 lbs of oil of turpentine.

(b) Tobacco extract manufactured in bond may be denatured by adding 30 parts of soft

soap to 70 parts of extract
(c) Hop powder, To every 75 lbs of snuff or (c) Hop powder. To every 75 lbs of snuff or tobacco are added 28 lbs of sulphur, 5 lbs of

asafætida, and 3 lbs of sago flour.

(d) Funiquet for horticultural purposes every 100 lbs of tobacco or snuff are added 10 lbs of ground hellebore, 18 lbs of saltpetre, 6 lbs of asafætida, 4 lbs of cayenne, 2 lbs of lampblack, 10 lbs of sago flour

Wine which has become unsound and sour (owing to the formation of acetic acid) so as to be no longer suitable for use as a beverage, is released from revenue custody after being denatured with a view to its conversion into wine vinegar The denaturant used in this case 18 20 p c of commercial vinegar, or an equivalent quantity of acetic acid.

(For methods of denaturing alcohol, v Methylated spirits and Industrial alcohol, art ALCOHOL.)

**DENITRIFICATION** v FERMENTATION AND

DENSIMETER v Specific gravity.

**DENTALONE.** A solution of chloretone in essential oils

DERMATOL. Bismuth subgallate BISMUTH, ORGANIC COMPOUNDS OF, also SYN-THETIC DRUGS)

DERMOGEN. A trade name for zinc peroxide.

DERMOL. Bismuth chrysophanate. yellowish - brown powder, said to have the composition Bi(C<sub>15</sub>H<sub>9</sub>O<sub>4</sub>)<sub>3</sub>Bi<sub>2</sub>O<sub>3</sub>. According to Merck, it is a mixture of impure chrysarobin and bismuth hydroxide (v Bismuth ORGANIC COMPOUNDS OF, CHRYSAROBIN and SYNTHETIC DRUGS

**DERRIS ULIGINOSA** (Benth.) The stem of this species of derris, used in the Far East as a fish poison, contains a poisonous resin (Proc Amer Pharm Assoc 1902, 50, 296).

DESAEGIN ('Colloidal chloroform'). An

albuminous substance containing chloroform.

DESCLOIZITE. A basic lead and zinc vanadate, (Pb,Zn)2(OH)VO4, crystallising in the

orthorhombic system, and isomorphous with olivenite (basic copper arsenate). A small amount of copper is sometimes present. The small crystals are deep reddish-brown in colour, and the streak is a paler yellowish-brown. It occurs in veins of lead ore in association with vanadinite, &c, and has been mined to a limited extent in New Mexico and Arizona, as a source of vanadium. It also occurs in considerable quantities in the Sierra de Cordoba in Argentina, and at Broken Hill in North-Western Rhodesia L. J. S

DESICCATION AND DRYING. The removal of mechanically admixed water from substances. The removal of chemically combined water is usually termed dehydration, but it is not always possible to draw a sharp distinction between

the two processes.

A substance may be termed 'dry,' even though it still contains mechanically admixed water, provided that this water is not apparent to the touch, the term 'desiccation' refers. strictly speaking, to the complete removal of admixed water.

The importance of drying and desiccating will be appreciated by a consideration of the numerous reasons for which these processes have to be employed, among which the following may be mentioned

1 To preserve material from undergoing physical and chemical alterations, eg leather,

tobacco, fruit

2 To remove water needed in preceding operations,  $e\,g$  drying coal after washing, woollen and cotton goods after dyeing, &c.

3 To facilitate accurate sampling, since substances are much more readily and uniformly mixed in the dry state than in the moist condition

4 To diminish the weight of an unnecessary ingredient and thereby diminish cost of packing, carriage, and freight; e.g. timber, peat, &c.

5 To retain desirable moisture in a substance by drying its surface, thereby rendering the outer layer impervious to moisture, eg. the drying of soaps and chocolates

6 To prevent injury to health or goods, as in the case of the moisture-laden air of textile

factories, evaporating rooms, &c

7 To prevent waste of heat in vapourising water when combustibles are burnt

A slight consideration of the various purposes served by drying processes is sufficient to bring out the wide range of degrees of dryness, and the various conditions to which the term 'dry' is applied This vagueness arises partly from the difficulty of stating quantitatively the amount of water present in a given instance, and partly owing to the varying capacity that substances possess for absorbing moisture. Thus it is that in ordinary conversation substances are variously described as 'soaked,' 'wet,' 'moist,' 'damp,' 'dry,' 'dried,' 'desiccated' or 'dehydrated,' in order to indicate qualitatively their degrees of dryness.

The drying processes in use may be classified under the following headings, corresponding with the degrees of drying to which they lead

A. Gravitational drying, by draining, filter-

ing, absorbing, and condensing.

B Mechanical drying, by pressure and centrifugal force.

C. Drying by evaporation.

A. Gravitational drying. The processes included under this heading are usually only preliminary to the employment of method C above.

Drainage is a natural process occasionally used for the purpose of removing suiface water, e.q. from washed coal The procedure requires

no special description.

Decantation is a process by which solid substances mixed with a large amount of water are separated from the greater portion of that liquid by allowing them to settle, and running off the liquid from above the sediment practice it is often the custom to arrange settling tanks' in series, stepwise, and gradually feed the mixture of liquid plus suspended solid into the upper vessel by means of a pipe The overflow runs into leading to the bottom the second vessel, and so on throughout the series. Arrangements are provided for emptying each tank as required China clay, chalk, and barytes are separated in this manner

Filtration is a process of accelerated drainage Thus wet pigments prepared by precipitation are drained on calico sheets stretched loosely over a skeleton framework Filtration is greatly accelerated by partially exhausting the air from the chamber into which the liquid drains vacuum filter has been designed to work con-tinuously, and consists of a hollow drum, mounted with its axis horizontal, and partly submerged in a tank containing the suspension to be filtered. The periphery is covered with filter cloth, and the water is drawn by suction, applied to the interior of the drum, through this cloth, leaving the solid matter as an outside By slowly revolving the dium, a continuous coating of solid is thus formed, and, after it reaches a certain thickness, an adjustable skimmer removes the surplus arrangement can be adapted to many drying operations, and consumes very little power

Absorption is a drying process in which the material is dried by causing it to yield its moisture to other substances which are capable of annexing it. This process is not often applied on a large scale, although it is commonly employed in the laboratory Instances are furnished, however, in the preliminary drying of large clay ware in dried plaster of Paris moulds, and in the drying of the highly water-laden atmospheres of evaporating rooms, dye houses, &c, by pumping warm, fresh air into them Warm air having a much higher capacity for holding water vapour than cold an, the result is naturally to effect a drying of the atmosphere, besides warming up the room and fittings

In the laboratory, substances are frequently dried by placing them in a desiccator This is simply a closed glass vessel containing in a suitable receptacle some chemical compound, such as anhydrous calcium chloride, concentrated sulphuric acid, or phosphoric anhydride, which has a remarkable affinity for water, so that it will, by absorption, remove water vapour from the interior even when the tension of the aqueous vapour therein is exceedingly small The substance to be dried being placed in the desiccator, in the interior of which the tension of aqueous vapour is almost nil, it gives off its moisture, and this is continuously absorbed by the desiccating agent employed. The latter must expose a large surface for rapid absorption | nected with a pump. Resting on the funnel is


The velocity of drying in a desiccator to occur may be greatly increased by exhausting the air from the interior (vacuum desiccator), since the rate at which moisture can diffuse from the material to the absorbing agent is thereby largely augmented The degree of drying attained cannot, however, be increased, since that is limited merely by the pressure of the aqueous vapour when it has reached equilibrium with the absorbing agent. The rapidity of drying is also increased by placing the material to be dried at the bottom and suitably airanging the desiccating material above it This arises from the fact that moist air has a smaller density than dry air at the same temperature and pressure; consequently, the natural process inside the desiccator is for the moist air to rise

The error in design exhibited in most laboratory desiccators. in which the above arrangement is reversed, was pointed out by Hempel (Ber 1890, 23, 3566), to whom desiccator shown in Fig 1 is due In this desiccator, the drying agent is placed in the trough of the lid

It is usual to evacuate laboratory desiccators by means of the water-pump, but it may be easily accomplished by simple chemical means (J Amer Chem Soc 1906, 28, 834; Amer Chem J 1902, 27, 340)



A desiccator for drying by means of a current of air has been described by Pratt (J Amer. Chem. Soc 1917, 39, 271) It consists of a Buchner funnel connected below, through



cock s', to a safety flask, which in turn is con-

a heavy brass disc, B, on the lower surface of which is cemented a ring or sheet of rubber, R, to make a tight joint. Pressure is brought to bear on the disc by means of three screws acting between the disc and a metal collar, c. Heavy rubber tubing, R', prevents the collar from exerting unequal pressure on the funnel. In the centre of the disc is set a brass tube, to which is attached, by means of heavy rubber tubing, R'', the cock s'. This cock is further attached to a system of drying-bottles or towers. The material to be desiccated is either filtered into or placed in the funnel (Analyst, 1917, 156).

Condensation -This is employed as a means of drying materials which must be handled at low temperatures, and is a method adopted in cold storages. The principle of the process is, briefly, that the moist air of the rooms is drawn out and passed into chambers maintained at a very low temperature, where it is made to deposit the greater part of its moisture as a liquid The dry air is then returned to the drying rooms, either cold, when it waims itself at the expense of the materials drying, or after being previously heated This circulation of air goes on continuously The method adopted in order to maintain a low temperature in the cooling rooms is to allow liquid ammonia, carbon dioxide, or ether to evaporate at the requisite rate through long coils of metal pipes over the outer surfaces of which the moist air is drawn The latent heat of the ammonia (or other) vapour is obtained at the expense of the heat of the moist air, which is thereby cooled, the vapour condenses on the pipes and the water trickles down into troughs and flows away. The ammonia (or other) gas is repeatedly used in a cycle of operations, being successively liquefied and evaporated This process is highly efficient, but requires careful handling for economical working

B Mechanical drying. These methods are efficient when water forms a large proportion of the total weight, and when the solid material to be separated is porous in texture. They leave the solid damp or moist

Drying by pressure—There are very tew cases in which direct mechanical pressure is used for removing water from substances, the most important of which is the direct squeezing of peat in specially designed machines, in order

partly to remove the surplus water.

Filter presses are used for partly drying semi-solids, and leave them still containing about 20 pc of moisture They are efficient in removing the large bulk of the water from granular substances in a state of thick mud Essentially a filter press consists of a number of flat sections or chambers formed of iron plates covered with filtering cloths, and supported in a strong horizontal frame, in which they can be squeezed up tightly together The chambers are made either round or square, and usually each has its own outlet cock, while the washing arrangement can be made reversible The sludge to be filtered is fed in under moderate pressure, and all the outlet cocks opened After water has ceased to flow out, the solid in the press is washed, if desirable, and the press then taken to pieces to remove the moist solid cakes from its interior. This process is used for filtering, among other things, sewage sludge, gold slimes, and precipitated pigments.

Drying by centrifugal force.—This method is largely employed for drying clothes in laundries, when the centrifugal machines are termed hydro extractors. It is also used for drying wool, cotton yarns, piece goods, &c., after they have left the dye houses, and for separating and drying sugar, gun-cotton, and staich.

The process consists in placing the material to be dired in a basket containing numerous holes, and rotating it in a houzontal plane at a considerable speed. The water is thrown off from the material very effectually, much more effectually than when simple filtration or filtration under moderate pressure is adopted.

A centufugal machine consists essentially of a circular basket, rotated on a vertical central shaft and enclosed in an outer stationary casing. The circular basket is drum shaped, having an open top to introduce the material and sometimes a valve at the bottom to form an outlet The periphery is perforated for dried crystals with holes of a suitable size. These baskets are made of steel, bionze, copper, aluminium, silver, or china, as may be suitable, and are fined with rubber, cnamel, or lead when dealing with acid liquids. The outer stationary casing serves as a guard in case of accident and as a The successful working collector for the water of a centrifugal machine depends upon the skill with which the vertical shaft is mounted and driven, since the apparatus must have sufficient freedom when running for the axis to take up the position it requires, otherwise vibrations will lapidly wreck the machine

C Drying by evaporation. This is by far the most efficient method of drying, indeed, several of the preceding processes are only employed as preliminary steps to the application

of this more powerful method

Drying by evaporation is a process in which the water is removed by first converting it into vapour and then conveying the vapour away. Water always has a tendency to pass into the gaseous state, and is constantly giving off vapour from its fice surface. If this process occurs in a closed space, some of the vapour finds its way back into the liquid, and a state of equilibrium is set up when the rate at which vapour is leaving the suiface has become equal to the rate at which it is passing back into the To the state of equilibrium at any particular temperature there corresponds a definite vapour pressure, which is a measure of the tendency of water at that temperature to pass into the gaseous state, and this vapour pressure rapidly increases with rise of temperature. When evaporation takes place in free space, no condition of equilibrium is reached, since, owing to diffusion of the vapour, the rate at which it passes back into the liquid can never equal that at which vapour leaves it. Evaporation therefore goes on until no more liquid remains From these considerations, it follows that the two main practical principles in drying by evaporation are (1) to warm the material and thereby increase the tendency of its contained water to vapourise, and (ii) to remove the vapour as it is produced from the neighbourhood of the substance, in order that the evaporation shall proceed continuously.

In applying these principles it is necessary to bear in mind that the rate of drying permissible in any given instance is restricted by (a) the physical and chemical nature of the material, which put a limit to the rate at which heat may be safely applied; and (b) by the cost of drying, since the plant mechanically most sound may not be most economical in practice. It must always be remembered that the nature of the material governs the method of drying to be employed, and must be considered before the question of mechanical efficiency.

The various methods for applying heat and for removing vapour are indicated below, and in order to explain more definitely how these methods are employed in practice, bird descriptions follow of various types of drying installations in actual use, and of the purposes

for which they are used

Application of heat. In the majority of cases hot air is employed for conveying heat to the material, and its use is attended with many advantages. Air, although an extremely poor conductor, is an excellent absorber of heat, the increase in volume resulting from this process naturally tends to disperse this heat and to warm surrounding materials by convection. The mobility of the air enables it to come into intimate contact with the surfaces of materials, thereby supplying the heat necessary to vapourise the moisture. Most important of all, the heated air can be thus supplied in large or small amounts, at any desired rate and at uniform pressure, while any of these factors may be rapidly and easily altered at will

Of the various sources of heat, the cheapest Owing to its unis, of course, solar heat reliability so far as quantity and continuity are concerned, its direct use is attended by considerable risks Indirectly, however, it is of great value, in being the primary cause of natural air currents. Drying directly by furnace heat obtained by fuel combustion is economical, but the method is restricted in use to drying those substances which are not sensibly affected by high temperatures For the commonest method of drying, by means of a warm blast of an, heat is usually supplied by passing the air over the surfaces of radiators. These are simply iron vessels, built to expose as much surface as possible, and through the insides of which steam, hot water, or hot gases are passed is conducted through the non-to-the outer surface, and waims the surrounding air mainly by radiation

The dangers encountered in the application of heat result from the nature of the material, and the factors involved, which are briefly discussed below, naturally affect various

materials to very different degrees

(a) Rate of application of heat Although the amount of heat required to vapourise a definite quantity of water is fixed, yet if this heat be applied at a rate faster than that to which the material can accommodate itself, injury will result, e.g. wet bricks will crack unless the greater part of their water is first driven off at a slow and uniform rate.

(b) Change in power of absorbing heat with loss of water As a material dries, its heat

capacity usually diminishes, and hence the risk of overheating increases as drying progresses

(c) Uncertainty as to degree of dryness,

which is often unavoidable

(d) Local application of heat. This is the usual cause of distortion and cracking, since most materials shrink as they dry, and local shrinking sets up strain. This makes it advisable to keep wet material from actual contact with the source of heat, and to agitate the material to continually expose fresh surfaces, since evaporation is a purely surface phenomenon.

Methods for removal of vapour. The natural method by which vapour is removed is by means of an currents, and, excepting in the case of the vacuum process, this is the method upon which all artificial processes for vapour removal

are based

The ordinary atmospheric process for drying utilises the natural movements of the air, eg by utilising the action of the wind on floors, pans, gantries, sheds, &c It requires no motive power, but, being dependent upon varying atmospheric conditions, it is not reliable

In the exhaust process the moisture-laden and is drawn away from the wet material quicker than gravity would perform the removal. For this purpose a propeller of fan is usually employed. Since the operation is carried out under slightly reduced pressure, vapourisation proceeds quicker than it otherwise would, on the other hand, the exhaust carries away the heat-bearing medium from the material to be heated, and unless proper control is exercised, this defect may more than counterbalance the increased rate of removal of vapour.

In the racuum process the material is enclosed in an air-tight receptacle, from which the air may be exhausted to any desired extent Heat is then applied, and the vapour pumped off as fast as it is produced ('onsiderable experience is necessary in the application of this process, which, properly applied, is highly

efficient and rapid

In the plenum process the vapour is removed by supplying sufficient warm air to create an augmentation in an pressure. A fan is usually employed for this purpose, and drying must, of course, take place in an enclosed room with regulated egress for the air. This process is usually the best to employ, since it can be maintained strictly under control.

Drying installations. Drying floors. Natural floors are not often used, since serious difficulties arise as to heating, handling the material, and removing the vapour Haymaking affords a

good instance of its employment

Artificial floors warmed by the products of combustion from a furnace, by hot gases, or by steam passing in flues or pipes beneath, are a considerable improvement on natural floors. Advantage is taken in constructing them of any convenient sandy or porous ground floor; sand is put down if the natural ground is clayey

These floors are used for drying china clay, calcium phosphate, barytes, chalk, cement, and calcined quartz, materials which are not readily injured by excessive or irregular heating.

Drying gantries or racks. These are employed in order that as much surface as possible may be exposed to air currents. Their construction assumes various forms. According to

circumstances, they may be either skeleton shelves, or racks, or props, &c Occasionally hot water or steam pipes are used as supporting racks, but their use is to be avoided as uneconomical, and tending to produce uneven drying and distortion

Such substances as damp corn, peat, raw rubber, chalk, bricks, and clayware are frequently dried on open racks, while pigments in colour works are often dried in open gantries

of skeleton shelves

Drying sheds. These are merely drying floors or gantries provided with roofs to keep off rain and snow These sheds are best constructed of light open framing in order to facilitate the removal of vapour, and the more open it is on all sides the better, provided that the substance is not such that direct sunlight must be excluded from it, eg rubber Such sheds are highly efficient in producing 'air-dried' Bricks are often dried in sheds erected on drying floors, while Chile saltpetre is dried in sheds without the application of artificial heat, as also are china clay and othre in the quarry state

Drying pans and troughs. Drying pans, correctly designed and handled, are very useful appliances The oldest commercial use of such pans is in the evaporation of salt brine, which, as usually carried out, is not at all an eco-The pans are usually rectnomical process angular, and each stands in its own shed They are directly heated, three or four fires being beneath each pan

Shallow earthenware pans are of great use in the colour trade, the pigments being readily dried if the pans are carefully arranged on gantries in well-heated and ventilated chambers

Semi-cylindrical iron pans heated from below by gas or steam give good results in drying soft materials, especially powders They are most efficient when provided with revolving grids for mechanically stirring the wet material and with valve outlets at the bottom for automatically discharging their dried contents

Granulated sugar is dried in a type of troughed pan, through which passes a hollow conveyer' carrying hollow paddles. Steam is blown through the conveyer and paddles. The trough is jacketed on one side, and through this jacket air is blown into the bottom of the trough. The sugar is fed into the hot end, and automatically carried by the heated paddles to

the outlet at the cool end of the trough

Kilns. These are buildings containing a furnace, the direct heat of which is used for drying materials by allowing the products of combustion to pass through them For substances which may be strongly heated without damage, kiln-drying is an economical method of procedure, since the walls of the kiln are very effective in saving heat that would otherwise be lost; but care must be taken to provide proper exits for the vapour

A primitive type of kiln is one in which calcium phosphate is dried. It consists of a plain rectangular brick building with a wooden roof. The floor is piled with wood, and the phosphate placed upon it. Flues admit air for combustion and circulation Such a kıln wıll burn for from 2 to 5 days and dry from 1000 to 1200 tons of phosphate.

Kilns for drying clays, fuller s earth, barytes, and chalk are constructed upon the same principle, the fire being placed so as to prevent the ashes mixing with the material, the latter is sometimes supported within the kiln on iron racks.

A rotating bed dryer is in extensive use for drying materials which allow of direct firing The material is placed on a circular cast-iron bed, which slowly rotates on a vertical shaft The bed is covered by a firebrick arch, into which the exit flue leads Heating is done directly by fires, and stationary blades fixed in the arch stir the material as it is drying blades are fixed at such an angle that the material is slowly moved from the centre, above which the feed is situated, to the edge, where it is automatically discharged a common type of drying kiln for ores

Drying cylinders. This term includes those dryers in which the material to be dried is contained in a cylindrical tube, which is slowly rotated The essential feature of their construction is that the rotation of the cylinder affords the means of moving the material along. With suitable materials and the necessary power, cylinders will automatically deliver large quantities of uniformly dried product, with the minimum of labour and fuel consumption They are consequently extensively used, and many types of drying cylinders have been

devišed For materials that withstand high temperatures, the heating may be carried out by passing the hot gases from a furnace through the tube The latter is constructed of steel, and is mounted nearly horizontally (slope of 1 in 12), with the furnace at the lower end The feed is situated at the higher cold end, and the gases pass out through a flue at this end, connected with a fan or chimney As the material gravitates down the tube it is dired, and on reaching the lower end, it drops out, quite dry, into trucks A modification of this construction consists in enclosing the steel tube in a horizontal brickwork chamber and heating it externally by passing the furnace gases into this brickwork chamber, whence they pass through a series of external hooded openings into the steel cylinder itself. This arrangement allows of a more gradual and systematic heating, while dust and other objectionable parts of the products of combustion may be arrested prior to the gases entering the cylinders these two arrangements, many others have been

devised, each having its particular advantages Cylinder dryers are very suitable for heating by steam, thus meeting the requirements of such substances as sugar, salt, soda, cocoa, coffee, &c., provided they contain not more than 25-30 p.c. of moisture. The drying cylinder may be either steam-jacketed externally, or internally, or by both methods united exhaust is best taken by means of a fan at the feed end of the cylinder. In using such dryers for sugar, they are employed in conjunction with a conveyer trough dryer (v. supra), into which the sugar falls after passing through the steamheated cylinder.

Drying rollers. These are heated rotating cylinders, with the outside periphery of which the material to be dried is brought into contact. They are made of steel, cast iron, copper, or tinned iron, and are usually heated by passing steam or hot air through their Another method of heating consists in coating the cylinder externally with felt and rotating it in a polished iron bed, against the under surface of which steam is blown. The material to be dried is automatically passed between the roller and its bed, and the rollers are usually arranged in series of two to eight

The latter method is largely employed in laundry work, and the use of drying rollers is extensive in the paper and textile trades

Drying tunnels. These consist essentially of a stationary passage through which the material passes either continuously or intermittently as the drying proceeds Drying in a tunnel is progressive from one end to the other Heat may be applied either by direct firing or by heated air, which passes into the tunnel from beneath one end and travels along to the other. The warm air current usually travels in the opposite direction to that in which the material moves, but not always, as, for instance, in the drying of wool, where the pressure of the warm blast is utilised in blowing the material forward The vapour-laden air may be removed either by chimney, exhaust, or by pressure fan, while the material may be handled in a very large number of ways, according to circumstances This method of drying is economical, but the initial cost of plant is often considerable

In drying timber, the material is carried on special trucks running on a tram track air is heated by means of a heater coil fed with steam and situated under the track at the The air ascends, dries the discharging end timber, thereby becoming cooler and denser, and then, pressed forward by the incoming air, it gravitates into a large space beneath the feeding end of the tunnel and passes up the Wool is dried in a tubular tunnel, being blown along by the warm air. Soap is dried by passing it on trolleys through narrow tunnels, usually arranged in fours, through which warm air is driven by means of a propeller. In particular, bricks and clayware are dried by passing them through drying tunnels, for which purpose a very large number of types of plant have been devised

Drying rooms. These are enclosed spaces, which are warmed by means of hot air, and into which the loader or discharger has to enter When not constructed of brick or stone, the material of the walls should be chosen for its insulating qualities

The expense attached to drying by this method is due largely to the manual labour required in handling the material, which arises, in most cases, from the fact that the peculiar properties of the material render it impossible to utilise mechanical methods. Since manual labour in a heavily moisture-laden atmosphere is unhealthy, desiccation in drying rooms is restricted in practice to those materials for which the process is unavoidable. Such products as hops, confectionery, leather, paper, textile fabrics, pottery, and rubber are dealt with by this method.

The materials are supported in the most convenient manner within the chambers, and warm air circulated either through or over them. The warming of the air may be accom-

plished either within or without the compartment, by means of radiators, and, if desired, the same air can be used continuously in a circuit (compare the condensation process)

Drying ovens. These are entirely enclosed drying chambers, the necessary heat for the purpose is not generated within them, neither is it necessary to enter them in order to charge or remove the material to be dried Ovens can be adapted to suit a wide range of conditions, and some of the most scientific and economical dryers are of this type

A great variety of methods are available for the purpose of heating, since the heat is generated outside the oven Products of combustion are readily excluded from the material no necessity for the plant to cool down before the dried product is removed, and hence heat losses are greatly minimised. Heat is applied to the material either by means of hot gases, usually hot air, or by direct conduction through metallic plates, as, for instance, in steam ovens. Vapour is usually removed mechanically, either by pressure or exhaust, but it may be pumped off, as, for instance, in all vacuum machines. Ovens are more efficient than drying rooms, since the latter are frequently used for other purposes than drying, whereas the former are designed exclusively for drying. Only a few types of drying ovens can be here described

Laboratory ovens are rectangular boxes, constructed of copper, iron, or aluminium, which may be either steam- or air-jacketed, or in simple ovens not jacketed at all The substances to be died are supported on shelves inside. With air ovens heat is applied directly at the bottom of the oven or of the outer jacket. A ventilator situated at the bottom of the oven door allows air to enter, and one or more chimneys in the lid allow the warm vapour-laden air to escape. A convenient method of heating, which can be readily controlled, consists in heating electrically, by passing a current through a number of resistance coils embedded in the jacket surrounding the oven (See also Analysis, Vol I) Removable tray dryers are largely used for drying tea, coffee, cocoa, cinchona bark, &c The trays are arranged in the oven in a series of lows. Each tray has a separate slide, and is independent of the others, and each column of trays has its own regulating damper for the hot air method of working is progressive, the wet material being first placed in the top tray of the column and passed downwards stepwise to the The heat required is supplied from a multitubular furnace, which also produces the air current which aerates the material The hot air may be allowed to rise through the material, or be drawn down by two centrifugal fans on either side of the centre through the substance. The latter arrangement is more readily controlled, gives a larger output, and effects a considerable saving in height A modification giving an even larger output consists in causing each horizontal line of trays, commencing with the lowest, to automatically tilt and discharge its contents into the base of the dryer. For drying other substances than those mentioned above, many other types of tray ovens have been devised.

Sliding-panel dryers are used in laundries, and can also be adapted for drying yarns, piece goods, &c. They consist of a row of vertical sliding panels, built up with hanging rods to form rigid horses for the goods. They are readily movable on runner wheels, and each panel may be withdrawn independently of the others, with the minimum loss of heat, as the back of the panel closes the outlet and thus reshuts the oven A hot-air blast is supplied from above, and after passing through the oven it discharges into the open.

The Lennox plate-oven dryer is a type which has many advantages, is compact, and occupies a small floor space The outer vertical cylindrical casing is built up in sections to allow for extension and renewal A concentric cylindrical At each external drum slowly rotates inside section there is a horizontal metal plate within the apparatus, and each section plate contains a hole, these holes are not vertically above one The material to be dried is fed in at the top, falls on the first section plate, and is slowly stirred round by horizontal arms on the rotating centre drum till it reaches the hole in the plate It then falls through on to the next plate, is slowly stirred round till it falls on to the third plate, and so on, till it is discharged at the base Hot air is blown in at the base, and passes upwards into the central drum, from which it issues, through numerous holes in its sides, passes through the material, and thence upwards and outwards

A horizontal steam-jacketed cylindrical oven, containing a very powerful and thorough agitating gear inside, is used for drying wet pasty materials like meat, fish, &c. The material is fed in at the top, and discharged as a dry, mealy powder through a manhole in the bottom The vapour is removed from the interior through a large outlet duct operated either by a fan or pump according to circumstances In vacuum ovens the removal of vapour by means of air or other gases in motion is discarded in favour of pumping away the vapour from the warmed-up material, operating in a closed space and diminishing the pressure within it to any desired The oven must be strongly built, and all joints must fit tightly to maintain the diminished pressure. The material may be supported inside in any suitable manner, for instance, on steam-heated grids The necessary accessories are a vacuum pump, a condenser for the vapour and a receiver for the condensed The latter is provided with a sight glass in order to tell when no more water is given off by the material in the oven effective working of such a plant depends largely upon the observance of various mechanical and engineering points, the chief feature of the vacuum process is the saving of time that it offects. The oven may be designed to suit either solids or liquids, and provision may also be made for effectively stirring the material during the drying process

Desiccators. (Laboratory desiccators have

already been described)

A large-scale desiceator is any type of drying machine, which, working by evaporation, carries the degree of drying to the utmost possible extent. In some of the types previously described, it is difficult to prevent thorough desiccation, and, in fact, to avoid dehydration and calcination, but a number of machines have been devised for the effective desiccation of liquids and thin

pastes, which work upon a common principle that may be briefly explained. It consists in spreading the liquid in a thin film over a heated metallic surface kept in motion, and, when the water has been removed, skimming off the solid residue with a knife. By suitable arrangements. this process can be made continuous, and it is employed on a large scale for desiccating milk to a dry powder A modified method of removing the desiccated product is to cause the liquid to spread out on the upper part of an endless horizontal metallic belt, working over two rollers contained in a heated chamber under reduced The heated liquid, being fed on to the belt at one roller, is desiccated by the time it arrives at the other, and, on the belt passing over that roller, the dry powder falls into the delivery hopper beneath.

Literature - Marlow's Drying Machinery and Practice, which contains a bibliography of the subject See also a paper by F. A. Alliott, J. Soc Chem Ind 1919, 38, 173 T.

DESICHTOL. Trade name for ichthyol ammonia deodorised by hydrogen peroxide DESOXYCHOLEIC ACID v Cholic ACID

DESPYRIN. Salicyl tartaric ester DETONAL. Diethyl acetyl urethane

**DETONATORS** v Explosives

DEVARDA'S ALLOY. An alloy of alumi-

nium 45, copper 50, and zinc 5 parts

DEVELOPED COLOURS. Colours formed on the fibre by coupling an azo dye after fixing with a second component. V. PRIMULINE.

DEXTRAN v. Gums

**DEXTRINS**  $n(C_6H_{10}O_5)$  When diastase acts on starch paste (gelatinised starch), maltose is the final product, but, intermediate between the starch and maltose, various dextrins are formed as the reaction proceeds in successive stages These dextrins are colloidal amorphous compounds, and do not form characteristic derivatives They are soluble in water, and precipitated by alcohol Only maltodextrin, which is possibly the simplest of them, has been prepared in anything like a state of purity, and most of the other dextrins described must be regarded as mixtures

As diastatic hydrolysis proceeds, the optical rotatory power diminishes, and the cupricreducing power increases The change is rapid to a certain point, but then becomes arrested; this behaviour is interpreted as proof of an essential difference between the higher and lower dextrins The constants of this stage of hydrolysis are [a]<sub>J</sub>=1625°, K (cupric-reducing power) = 48 3, and correspond to the conversion of four-fifths of the starch into maltose remaining dextrin is hydrolysed with greater difficulty According to H. Brown, it cannot have a simpler formula than  $40(C_6H_{10}O_5)H_2O$ .

The dextrin isolated from a starch conversion stopped at a stage which gives a deep brownishred colouration with iodine is termed erythrodextrinAt a later stage indine ceases to give a colouration, and the dextrin present is termed achrodertrin. These dextrins have little or no cupric-reducing action, this point being still undecided.

Dextrins are also formed by the action of hot dilute mineral acids, or of heat alone on starch The final product in this case is dextrose instead

These methods are used for the of maltose commercial preparation of dextrin

Malfitano and Moschkoff (Compt. rend. 1912, 154, 443-446) have found that starch may be converted into dextrin merely by desiccation When kept in vacuo at 25° over phospholic oxide 28 l p c of the starch was found to have become soluble in the course of 20 days authors conclude that the starch molecule is composed of dextrin complexes linked together by water molecules

Oscar von Friedrichs (Arkiv. Kem Min Geol 1913, 5, No 3, 1-14) has studied the action of various yeasts upon the dextrins Generally speaking, dextrins with higher molecular weights are not so readily attacked as those with lower.

Yeasts which split up α-methyl glucoside are capable of fermenting dextrins to a very limited extent, while Sacchsia Suaveolens, which is known to ferment  $\beta$ -methyl glucoside, is found to ferment achroodextrins energetically results support the view that  $\beta$ -glucoside linkings are present in the starch and dextrin molecules.

Crystalline dextrins may be prepared by the action of Bacillus macerans upon starches (Schardinger, Centr. Bakt Par 1911, ii 29, 188-197) By this method two dextrins designated as  $\alpha$  and  $\beta$  have been isolated, both in crystalline form.

According to Langhans (Ber. 1912, 45, 2533-2546), both forms are acctylated by acetic anhydride in presence of zinc chloride, but seission of the molecules occurs simultaneously; under this treatment the  $\alpha$ - compound gives a hexa-acetate of a diamylose, the B- a nona-acetate of a triamylose, hydrolysis of the acetates yields a diamylose and triamylose respectively, both crystalline compounds

The properties of these crystalline dextrins have been further investigated by Pringsheim and Eissler (Ber 1913, 46, 2959-2974, see D R P 279256, 1913), in which the dextiins are produced by the action of Bacillus maccians on starch and isolated by precipitation with light petroleum

In presence of a little platinum black as catalyst, l-glucosan, heated to 180°, undergoes polymerisation, and is transformed into a white amorphous substance, which exhibits the characteristic properties of a dextrin and yields dextrose when warmed with dilute sulphuric acid (Pietet, Helv. Chim. Acta, 1918, 1, 226)

Maltodextrin  $6(C_6H_{10}O_5)H_2O$  has  $[\alpha]_J+193$  6°, [a]<sub>D</sub>+181°; cupric-reducing power, 42 These figures agree with a mixture of 1 part maltose to 2 parts dextrin; but the facts (1) that it cannot be separated into maltose and dextrin by dialysis, (2) that it is not fermentable by yeasts, (3) that it is completely converted into maltose by diastase, prove maltodextrin to be a definite chemical compound To prepare it, a starch transformation is carried out at 60°-65° until  $[\alpha]_J = +198^\circ$ The solution is boiled, concentrated to a spgr of 106, and the maltose removed by fermentation. The solution is filtered, evaporated to a syrup, and digested with alcohol first of 90 p c, and then of 85 p.c This last liquid is decarted hot, and the alcohol distilled off, when maltodextrin remains; it is purified by further treatment with alcohol.

On oxidation of maltodextrin with mercuric oxide, until the cupric - reducing power has disappeared, maltodextrinic acid is obtained This behaviour proves maltodextrin to contain an aldehydo group, but the change goes further than the oxidation of this group to carboxyl (CO,H), since it splits up the terminal C12 group into a C<sub>5</sub> acid residue, which remains attached to the rest of the molecule, and a C<sub>6</sub> residue which is eliminated

495

(For a full account of the decomposition

products of starch, v STARCH )

Manufacture - The commercial products known as solid dextiin, British gum, starch gum, &c, are mixtures containing various dextrins, together with carbohydrates (copper oxide reducing substances). They consist mainly of erythrodextrin, and accordingly give a brown iodine colouration

They are prepared (1) by the action of heat on dried starch; (2) by the action of heat on acidified dried starch

The starch is first freed as much as possible from combined water by drying in suitable ovens, and then submitted to a temperature of 212°-275° in rotating sheet-iron drums heat is supplied by hot oil, rape oil being fairly When the oilgenerally used, or by direct fire bath is used, the control over the temperature is greatest, and the colour of the product can be varied at will from white to dark brown the drums are heated by direct fire, they are made to rotate on slightly inclined axes, the dried starch being fed in at the higher end of the revolving cylinder and the product discharged at the other Sheet-non ovens are also used they are constructed singly or in sets heat is supplied by a furnace, the heated an from which is drawn over the top of the ovens The material is kept continually stirred by flat iron oars. Boxes and trays are also used as converters, and, indeed, the maker has only to consider the variety and quality of product required, and to remember that the direr the starch and the lower the temperature of conversion, the whiter will be the product, and, further, that different starches yield different products, ie behave differently when submitted to the action of heat, to enable him to construct an apparatus that will suit his require-It must also be borne in mind that the conversion takes place more slowly at a low temperature

The alternative method, in which less heat is required to yield a satisfactory product, is to spray or moisten the starch either with a mixture of dilute nitile and hydrochloric acids, or with each acid singly, or with dilute oxalic acid The paste is dired and heated at 100°-120° or to 150°, till the transformation is complete, as determined by the iodine test, it must then be arrested promptly.

Payen recommended 1000 lbs starch to be moistened with 30 gallons water containing 2 lbs nitric acid (sp gr 138), the mixture to be well kneaded, made up into balls, and dried in a hot-air chamber; then powdered, spread on brass trays to the depth of 1½-2 inches, and submitted to a temperature of from 110°-120° in a hot-air oven.

The heat method is that which was most commonly used on the Continent, whereas the

496 DEXTRINS.

use of weak nitric acid found most favour in this country.

The products thus obtained appear in commerce in the form of powders varying in colour from almost white to dark brown, through every shade of light yellow and brown. They are reduced to this state after roasting, by crushing in pans or passing through rollers, and then sifting through a rotating silk screen such as is used in flour mills.

Dextrins prepared from powdered starch are used by textile manufacturers and calico printers as vehicles for colours. They are also used for all kinds of gum, paste, sizings, and in the manufacture of paper boxes, wall paper, and for a variety of other purposes.

For a method of examination of commercial dextim and related starch products, v Babington, Tingle, and Watson, J Soc Chem Ind

1918, 37, 257 T.

On heating dry starch with formic or acetic acids, the starch nucleus is broken down and esters of dextrins are formed, which are soluble in cold water The proportion of acid fixed depends on the time of treatment After 15 hours' heating with glacial acetic acid at 90°, the product just ceases to give a blue iodine colouration, and the products of the further heating are soluble in cold water When care is When care is taken to exclude moisture, and the heating is continued for 2 or 3 hours only, using equal weights of starch and glacial acetic acid, an acceylated starch, known commercially as Feculose, is obtained. This behaves like starch towards boiling water, but the jelly does not revert on standing It forms very clear flexible films, and is said to give better results than dextrin as a textile finish (see Traquair, J Soc Chem Ind 1909, 28, 288)

Destrin symps find employment in a variety of industries. They are prepared from staich by heating it with hydrochloric or sulphuric acids in precisely the manner already described for dextrose (v. Carbonydbrates), with the exception that conversion is stopped at a much earlier stage. Thus in the modern American process of manufacture of corn syrup from maize starch, less than 10 minutes' heating is required. The product evaporated to 42°-45° B forms a colourless viscous syrup, baving the composition water, 19 pc; dextrose, 38 5 pc; dextrin, 42 0 pc., ash, 0 5 pc

In America corn syrup is used extensively in making confectionery, preserves, and table syrups, being preferable to sugar or cane syrup,

as the goods are not likely to grain

According to Yano (J Chem Ind Tokyo, 1918, 21, 865), dextrin prepared by adding 10 parts dilute nitric acid (5 parts of acid of sp.gr. 1-4 and 95 parts water) to 100 parts of sweet potato starch, drying the resulting paste at about 50°, and heating it for an hour at about 150°, is a suitable substitute for gum arabic as an adhesive

Dextrin may also be prepared by the action of malt extract on gelatinised starch. E F A.

**DEXTROFORM.** A condensation product of dextrin and formaldehyde.

DEXTROSE v. CARBOHYDRATES.

DHAK GUM v Kino.

DHAURA. A gum obtained from Anogeissus latifolia (Wall), It is extensively used in India in

calico-printing; it occurs in clean straw-coloured elongated masses, sometimes honey-coloured or even brown from impurities. As an adhesive gum it is inferior in strength to gum arabic. In India, the reputation of this gum stands high with the calico printers, especially of Lucknow, and it is probable it possesses some specific peculiarity justifying the preference, since it is used with certain dyestuffs, such as with haldi (Curcuma longa [Linn]), whilst gum arabic or 'babul' is used with madder (Rubia cordifolia [Linn]) (Watts' Economic Products of India).

DHURRIN v GLUCOSIDES.

DIABASE. This name was introduced by the French mineralogist A Brongniart, in 1807, for those varieties of basalt in which the light and dark minerals, respectively felspar and augite (or as he erroneously supposed, felspar and hornblende), are distinguishable to the unaided eye As actually defined by him, the term is therefore synonymous with the diorite of Hauy (1822), whilst with the correction of augite for hornblende it becomes a synonym of the dolerite of Hauy (1822) In the latter sense it is used by some authors at the present time (v. Dolerite) Unfortunately, the term was re-defined by German petrographers, and it came to be applied to the older, or pre-Tertiary, dolerites, which differ from those of more recent date only in showing correspondingly more alteration, with the development of secondary minerals, such as chlorite In consequence of this, the name 'diabase' is now applied by most English authors to a partially altered or weathered dolerite In this sense it has an economic bearing, since rocks of this class, being of wide distribution, are much quarried for road mend-For this purpose the fresher dolerites are somewhat brittle, whilst diabases are tougher and more durable, providing the decomposi-tion has not proceeded too far There is also a There is also a tendency to use the name 'diabase' in a loose sense, like the popular terms 'greenstone,' whinstone,' 'toadstone,' and 'trap-rock'

L J S.

**DIABETIN.** A trade name for a preparation of lævulose

DIACETYL v KETONES

DIACETYLDIOXIME, DIMETHYLGLYOX-IME v Ketones

DIACETYL MORPHINE v OPIUM.

**DIAL-CIBA.** A trade name for diallyl-barbituric acid.

DIALKYL BARBITURIC ACIDS v. Pyrimi-

DIALLAGE. An important rock-forming mineral belonging to the monoclinic series of the pyrovene group. It has the same composition as diopside and hedenbergite,

# Ca(Mg, Fe)(S1O<sub>3</sub>)<sub>2</sub>

but with the addition of alumina passes into augite. It differs from the other members of the pyroxene group in possessing a fine lamellar structure, due to secondary twinning and partial alteration of the material. On the surfaces of foliation the mineral displays a shining metallic sheen, and on this account it is sometimes used as an ornamental stone. The colour is grey, brown, or green. Diallage is of common occurrence as a constituent of gabbro.

L. J. S.

**DIALOGITE.** Manganese carbonate (v. Manganese and Rhodochrosite).

DIALURIC ACID, Tartronylurea

prepared by reducing alloxan with zinc and hydrochloric acid, or with sulphuretted hydrogen at the temperature of boiling water (Liebig and Wohler, Annalen, 26, 276), or with ammonium sulphide, when ammonium dialurate is obtained and is readily converted into the acid by the action of hot dilute hydrochloric acid (Gregory, Chem. Soc. Mem. 1845, n. 11). A mixture of potassium dialurate and oxalurate is obtained by adding potassium carbonate to a solution of alloxan containing potassium cyanide (Strecker, Annalen, 113, 53), it is prepared by reducing alloxantin with sodium amalgam, or dibromobarbituric acid with sulphuretted hydrogen (Baeyer, Annalen, 127, 12; 130, 133) Magnier de la Source showed that dialuric acid is obtained by prolonged boiling of uric acid with water (Bull Soc. chim 1875, 23, 529), or it can be obtained from uric acid on gentle heating with hydrochloric acid and potassium chlorate (Biltz and Damm, Ber. 1913, 46, 3662–3665) Grimaux effected its synthesis by treating tartronic acid and urea with phosphorus oxychloride (Compt rend 1879, 88, 85)

Dialuric acid forms colourless short foursided prisms, or narrow thin leaflets with rounded ends, sparingly soluble in cold, more readily so in hot water, turns red at 180°, m p 214°-215° (corr) (Biltz and Damm, lc 3664). The moist crystals or the aqueous solution rapidly absorbs oxygen from the air to form alloxan, which combines with the dialuric acid, forming alloxantin The rate at which the change takes place is illustrated by the results obtained by Hartley (Chem Soc Trans. 1905, 87, 1804), in an unsuccessful attempt to obtain the absorption spectra of dialuric acid; 0 144 gram of pure dialuric acid was dissolved by agitation and warming in 100 cc of water, and immediately afterwards the solution yielded the same series of absorption spectra as a solution of 0 322 gram of alloxantin, dissolved in 250 cc of water Alloxantin is also formed by mixing solutions of equivalent quantities of dialuric acid and of alloxan (cf Richter, Ber 1911, 44, 2155-2158). Dialuric acid is monobasic, and, contrary to the statements of Menschutkin (Ber. 1875, 8, 760; Annalen, 1876, 182, 70), forms only one series of salts of the type  $C_4H_3O_4N_2M'$  The sodium, polassium, ammonium, and barium salts are crystalline (Koech, Annalen, 1901, 315; Behrend and Friederich, Annalen, 1906, 344, 1)

crystallises in white prisms or leaflets; m.p.  $210^{\circ}-212^{\circ}$ ; gives a violet precipitate with barium hydroxide; forms the potassium salt  $C_6H_5O_5N_2K,H_2O$ , and combines with alloxan to form acetylalloxantin  $C_{10}H_8O_9N_4,H_2O$ , m.p.  $263^{\circ}-265^{\circ}$ ; benzoyldialuric acid  $C_{11}H_8O_5N_2$ , crystallises in prisms, m.p.  $209^{\circ}-210^{\circ}$ , yields with barium hydroxide a white precipitate, turning violet, and combines with alloxan to form benzoylalloxantin  $C_{16}H_{10}O_9N_4,H_2O$ , or  $1\frac{1}{2}H_2O$ , m.p.  $253^{\circ}-255^{\circ}$  (Behrend and Friederich, Annalen, 1906, 344, 1-18)

Substituted dialuric acids of the type  $CO \stackrel{\text{NH CO}}{\sim} C(OH)R$  are obtained by the condensation of alloxan with (1) aromatic ketones, eg phenacyldialuric acid, where  $R = C_6H_5$  CO  $CH_2 -$ ; (2) aromatic amines, eg. p-aminophenyltartronylurea, where

$$R = NH_2 C_6 H_4 - ;$$

(3) pyrazolone bases, eg phenylmethylpyrazolonetartronylurea, where R = | C - |

(4) phenols, e.q p-hydroxyphenyltartronylurea, where R=OH C<sub>6</sub>H<sub>4</sub>-. For the preparation and technical application of these compounds, see art ALLOXAN.

Methyl dialune and C<sub>5</sub>H<sub>6</sub>O<sub>4</sub>N, H<sub>2</sub>O may be prepared by reducing dimethyl alloxantin with sodium amalgam, crystallises in round clusters of silky needles or narrow lancet-shaped leaflets, in p. 167°-168° (corr.), or directly from theobromine on careful treatment with finely divided potassium chlorate in presence of hydrochloric and crystallises in bundles of colourless glistening filaments, many of which widen to slender narrow leaflets. The molecule of water is driven off at 110°-120° From alcohol the and crystallises in anhydrous, stout, lustrous rhombohedra, in p. 184°-185° (Biltz and Damm, Bei, 1913, 46, 3665-3667)

Dimethyldialuric acid
CO NMe CO CH OH

obtained by reducing amalic acid (tetramethylallovantin) with sulphuretted hydrogen (Maly and Andreasch, Monatsh 1882, 3, 92), or with sodium amalgam (Techow, Ber 1894, 27, 3082), forms colourless compact crystals which become red at 100° and decompose at 170° (cf Biltz and Damm, thid 3667–3668), the acid crystallises with one molecule of water, which is partly lost in vacuum and completely at 110°. The hydrated acid has mp just under 100°, and decomposes about 210°. The anhydrous acid begins to discolour at 190°, and has mp 218° (corr) with decomposition. It forms crystalline alkali salts, and reduces cold solutions of silver and copper salts.

180-Dialuric acid

the action of bromme water on 150-barbitume acid CO NH CH CO C(OH) or aminouracil or hy-

droxyxanthine (Behrend and Roosen, Ber 1888, 21, 999), crystallises in trimetric prisms sometimes a centimetre long (Weinschenk, Annalen, 1889, 251, 243), is readily soluble in water, less so in alcohol, it loses one molecule of water of crystallisation at 100°. The other molecule, which is regarded as water of constitution, is lost at 140°-150°, at which temperature the compound decomposes Isodialuric acid is converted into dialuric and by the action of bases (Koech, Annalen, 1901, 315, 246). It yields unclaim of silver salts, and combines with hydroxylamine to form two isomeric oximino- compounds. The

a- oxime forms shining plates or prisms, becomes red and decomposes at 100°; the \$\beta\$- oxime crystallises in needles which do not become red at 110° (Behrend and Roosen, Annalen, 251, 244).

Methylisodialuric acid C5H8O5N2, prepared by oxidising methylisobarbituric acid (Lehmann, Annalen, 1889, 253, 80) with bromine water, crystallises in rosette-like aggregates from water, and in contradistinction to isodialune acid, contains only I molecule of water, which 13 regarded as water of constitution M A. W. DIAMIDOGEN BLACK, -BLUES v Azo-

COLOURING MATTERS

DIAMINE BETA BLACK, -BLACK, -BLUE, -BRONZE, -BROWNS, -CUTCH, -FAST RED, -GOLD, -GREEN, -PINK, -ROSE, -SCARLET, -VIOLÉT,-YELLÓW v Azo-colouring matters, Primuline

DIAMINES v AMINES.

DIAMINOACRIDYLBENZOIC ACID ACRIDINE DYESTUFFS

DIAMINOBENZENES, DIAMINOTOLUENES v. AMINES

4.4'-DIAMINOBENZOPHENONE Ke-TONES

4 4'-DIAMINODIPHENYL - 3 . 3' - DISUL-PHONIC ACID. Benzidine disulphonic acid v. DIPHENYL

# 2.6-DIAMINOPHENOL-4-SULPHONIC ACID



is obtained by reducing a solution of sodium 2 6-dinitrophenol-4-sulphonate by zine dust and hydrochlone acid Used in the manufacture of azo dyes (Farb vorm Meister, Lucius, and Biuning, 1) R P 148212)

#### DIAMINOSTILBENEDISULPHONIC ACID

$$NH_2$$
  $CH \cdot CH$   $NH$   $SO_3H$ 

may be prepared by adding zine dust to sodium p-nitrotoluene sulphonate in a hot solution of caustic soda, until the liquid is decolourised, filtering and acidifying with hydrochloric acid The precipitated diaminostilbenedisulphonic acid may be purified by re-solution in sodium carbonate, and reprecipitation with acid (Leonhardt and Co D R P 38735, cf Bender and Schultz, Ber 1886, 19, 3234) May also be obtained by reducing dinitrostilbenedisulphonic acid with ferrous hydroxide and ammonia (Green and Wahl, Eng Pat 5351, 1897, Ber 1897, 30, 3100)

**DIAMOND.** Carbon crystallised in the cubic The contrast presented by the physical system characters of the two forms of crystallised carbon—diamond and graphite—is very striking Diamond is the hardest of minerals, and, indeed, of all known substances, but at the same time it is very brittle, on the other hand, graphite is one of the softest of minerals, and it is flexible and inclastic The one is colourless and transparent and a bad conductor of electricity, whilst the other is black and opaque and a good conductor. There is also a considerable difference in density, diamond, 3 52; graphite, 2 25 "I'he name diamond, in French and German Diamant, and similar in most other European

languages, is a corruption of the Latin Adamas, which was used for this mineral by Manilius (A.D. 16) and Plmy (A D 100) This came from 'Aδάμας, meaning invincible, a word applied by the Greeks to hard metals and stones similar name adamantine-spar was used by the chemist Joseph Black for coundum. Although diamond and colundum are numbered respectively 10 and 9 in the mineralogist's scale of hardness, and with no other mineral falling between the two, yet, as shown by abrasive tests, diamond is 140 times harder than corundum, and more than a thousand times harder than quartz (No 7 on the scale) It is on this high degree of hardness that all the technical applications of diamond depend, and a faceted gem never shows any signs of wear. Lapidanes assert that stones from Borneo and New South Wales are harder than those from other localities.

Crystalline Form -Crystals of diamond are almost always found singly and with faces developed on all sides, thus indicating that they grew freely from individual centies in some surrounding medium. By far the commonest form is the regular octahedron, the faces of which are often delicately marked with minute triangular pits or 'etched figures,' whilst the edges are nearly always rounded and often marked with a pronounced furrow Other forms are a hexakis-octahedron (six-faced octahedron), and less often the rhombic-dodecahedron and the cube, while crystals of distinctly tetrahedral habit are rare Nevertheless, it is probable that crystals of diamond are really hemihedral (tetrahedral-cubic), the apparent octahedron consisting of two tetrahedra interpenetrating in twinned position Distinctly formed twinned crystals are not uncommon. These are contact-twins with a face of the octahedron as twin-plane, and the two individuals flattened parallel to this plane, giving a triangular plate with re-entrant angles at the three corners These are known to the Dutch cutters as 'naadsteenen' (suture-stones). In addition to the rounded edges, noted above, the faces themselves are often curved, sometimes to such an extent that the crystals are almost spherical in form. This rounding, as well as the etched figures, indicate that the crystals have been corroded subsequent to their growth An important crystallographic character of diamond is the existence of perfect cleavages in four directions parallel to the faces of the (On the crystallography of diaoctahedion Fersmann and V Goldschmidt, mond, v ADer Diamant, Heidelberg, 1911)

Physical Characters - In its optical characters, diamond is remarkable for its very high refractive and dispersive powers, the indices being for red light (B) 2 40735, yellow (D) 2 41734, violet (H) 2 46476, and the dispersive value (H-B) 0 05741 The critical angles of total reflection corresponding with these indices are 24° 33′, 24° 26′, and 23° 56′, respectively. The angles between the facets of the brilliant form of cutting adopted for the diamond are so arranged that a ray of light entering the gem by the front facet will meet the back facets at a greater angle than the critical angle, and will be internally totally reflected, passing out again from the front of the stone, and with a chance that rays of different colours will be separately

DIAMOND 499

ected A cut stone, therefore, shows much illiancy or 'fire,' together with flashes of ismatic colours Corresponding with the high ractive index, diamond displays a high degree lustre, which is of the quality known as damantine 'lustre Uncut stones, especially ien not quite clear, and with rough rounded rfaces, present a peculiar and very characteramond is without colour and perfectly transrent The best stones are described as of the ust water' or 'blue-whites.' More often ere is a tinge of colour ('off-coloured' stones), ually pale yellow, brownish, grey, or greenish. ones of a good canary-yellow, coffee-brown, black colour are not uncommon, but those of h shades of green, blue, or red are very rare, no nature of the colouring matter is uncertain, it it is probably inorganic (eg iron and anium in the brown stones) (hanges in lour may be effected by the action of heat or / exposure to radium emanations ('rystals diamond being cubic, they should be optically otropie, but when examined under the polarisg microscope they are frequently seen to possess iomalous double refraction, especially around iclosures in the stone This is due to a state strain, which in some stones is so great that ey burst Phosphorescence is readily produced diamond by the action of sunlight, ultraolet, Rontgen, cathode, and radium rays, or mply by rubbing (triboluminescence) Diaand becomes positively electrified by friction is very transparent to the Rontgen rays; hilst glass imitations are opaque – Bem∉ a ood conductor of heat, it can be distinguished om a glass mutation by touching with the tip the tongue, diamond feeling much colder an glass. The specific heat is less than that f graphite, and it increases rapidly with the emperature

Chemical Characters — The combustibility of iamond was first proved experimentally by the lorentine Academicians in 1694 Lavoisier, in 772, found that the presence of an was necesirv, and that 'fixed air' iesulted, but it was mithson Tennant, in 1797, who first proved hat equal weights of diamond and carbon ielded equal weights of carbon dioxide onvincing experiments in this direction have een made by Davy, Dumas and Stas, A Crause (1890), and H. Moissan (1893). Powdered hamond burns readily in air at a red heat 101ssan determined the temperature of ignition n oxygen to be 690°-790°, according to Joly, t is 850° in air It burns with a small paledue flame, and in oxygen is able to support ts own combustion It is infusible even in the lectric arc; and it can be heated at high emperatures (1500°) without alteration in various gases (hydrogen, nitrogen, sulphur lioxide and trioxide, and nitrous oxide) earbon dioxide, however, at 1200° the crystals ire corroded, with formation of carbon monoxide t is unattacked by acids, caustic alkalis, iodic inhydride, hydrogen potassium sulphate, or a nixture of sedium chlorate and nitric acid, but s oxidised when heated to 180°-230° in a mixture of potassium bichromate and sulphuric acid In fused potassium nitrate or sodium carbonate at about 900° small etched figures are slowly produced; at 1200° the action is more rapid

carbon dioxide being liberated. It is acted upon by sulphur vapour at 900°, and it combines with iron, yielding steel ('iystals are etched by molten olivine (magnesium silicate) The fact that diamond gives with oxidising agents only carbon dioxide, whilst graphite yields graphitic oxide has led to the suggestion that these two forms of carbon are chemical isomerides Observations at high temperatures are somewhat contradictory; a mere blackening of the surface has often been observed, but not always confirmed C Doelter (1911), with crystals packed in charcoal or thorium oxide and heated to 2500°, obtained only a superficial blackening without any alteration in the physical characters of the material Moissan (1893), however, records the conversion into graphite at the temperature of the electric arc (about 3600°); and Parsons and Swinton (1907) found that a diamond placed in the focus of cathode rays un vacuó swelled up into a coke-like mass when the temperature reached 1890° In this connection the occurrence of cubes of graphite (cliftonite) in meteorie irons is significant, these have been supposed to be paramorphs after diamond

The ash remaining when diamond is buint amounts, for colourless crystals, to 0.02–0.05 p.c., but in the less pure carbonado it may reach 4.8 p.c.. It consists mainly of non-oxide and silica, with some lime, magnesia, and titanium. The non-and titanium may represent the colouring matter present in the stone, but the frequent presence of minute enclosures in diamond is not to be overlooked. In addition to liquid enclosures (carbon dioxide and water), the following materials have been recorded, black carbonaceous matter, ilmenite or hematite, quartz, juitle, non-pyrites, gold, chlorite (1), apophyllite, &c

Varieties—Differing in certain points from the general characters enumerated above, there are two varieties of diamond—bort and carbonado—which, for technical purposes, are more important than the puricity stals used for gens. These are crystalline aggregates, as distinct from single well-developed crystals.

Bort (boart or boort) consists of a number of individual crystals with a radial grouping around a common centre, giving rise to rounded extenor forms These are sometimes quite spherical ('shot-bort'). They are usually rough on the exterior, it being only here that the fibres, or individual crystals, have been free to develop crystal faces—in the interior these have interfered with each other's growth. Although each individual possesses the cleavage characteristic of diamond, yet for the whole mass there is no continuous cleavage. For this reason, bort is tougher than the single crystals, which readily split along the cleavage directions, and is thus better adapted for boiling, turning, and cutting tools This, no doubt, is the explanation of the common statement that bort is harder than diamond; further, since in ciystals the degree of hardness varies with the direction, a chance section of bort will present both maximum and minimum values. The rounded pieces of bort are translucent to cloudy and opaque. They are grey or black with a gleasy to metallic lustre, and, usually, a peculiar leaden appearance; sp.gr. 3.50. In the trade

DIAMOND. 500

the term bort also includes fragments (such as those obtained by bruting), and all crystals and stones that are useless for gem purposes. Such material is of value as an abrasive, but not for tools.

Carbonado, or 'black diamond' (known in the trade as 'carbon' or 'carbonate'), is a finely granular crystalline material, dull, black, and opaque, and often more or less porous, with somewhat the appearance of coke. It forms irregularly-shaped masses, and consists of a confused aggregate of minute crystal grains of diamond. It therefore presents no cleavage as a whole, and is consequently tougher and less frangible than crystals It is less pure than ordinary diamond, yielding more ash (up to 48 pc) when burnt. An analysis of the ash gave Fe<sub>2</sub>O<sub>3</sub>, 53 3, SiO<sub>2</sub> 33 1, CaO 13 2 pc, MgO trace The sp gr is less, 3 15-3 34 (the value for crystals being between the limits 3 516-3 525) The largest known piece of carbonado was found in Bahia, Brazil, in 1895 It weighed 631 9 grams, that is, slightly more than the famous 'Cullinan' diamond (which weighed 6212 grams) Both carbonado and bort show considerable variations in texture, and with increasing coarseness of grain they may pass insensibly into ordinary diamond

Occurrence -India, from very remote times until the middle of the eighteenth century, was the only source of diamond (with the unimportant exception of Borneo) Since the discovery of the Brazilian and South African deposits the output has gradually fallen off, amounting in 1916 to only 20 carats. The amounting in 1916 to only 20 carats secondary deposits have, however, only been worked by native methods, and the original deposits have not been discovered. Indian diamonds, being of the best gem-quality, there is thus a possibility of further developments. They occur associated with pebbles of jasper and vein-quartz in thin beds of conglomerate in the ancient sedimentary rocks (sandstones and shales) of the Vindhyan system (probably pre-Cambrian), which rest directly on the crystalline rocks With the weathering of these strata, the diamonds are left on the surface or are accumulated in the alluvial deposits of the The districts where mining present rivers has been done fall into three main groups: a southern group in the basins of the Pennar, Kistna, and Godavaii rivers, an eastern group in the Mahandi valley, and a northern group near Panna in Bundelkhand A doubtful occurrence of diamond in pegmatite has been described from Wajra Karur near Bellary, and, unfortunately, nothing is known as to whence the diamonds found in the ancient sedimentary rocks were derived (On Indian diamonds, v Manual of the Geol of India, vol 3, Economic Geology, 1881, by V. Ball, vol. 4, Mineralogy, 1887, by F. R. Mallet)
In Brazil, diamonds were discovered about

1725 in the alluvial gold-washings at Tejuco (now Diamantina) in Minas Geraaes, and, although they have been since found in several other districts, notably in Bahia, this has remained the principal diamond-mining centre. Here, as in India, the diamond-bearing alluvial deposits have been derived from conglomerates and sandstones (including the flexible sandstone known as itacolumite) of considerable geological Finally, the stones are cleaned by boiling in a

age, and the original rock from which these were derived is not known. The Brazilian deposits are, however, remarkable for the variety of minerals found as well-worn pebbles ('favas') in association with the diamond; and the nature of these minerals, or indicators, has led to the suggestion that they, together with the diamonds, originally came from quartz-veins. (On Brazilian diamonds, see Mineralogical Papers by O. A Derby and by E Hussak, also E Hussak, Os Satellites do Diamante, Rio de Janeiro, 1917, H. Preston, J. Soc. Arts, 1909, lvm 101.)

In South Africa, diamonds were first discovered in 1867 near Hopetown on the Orange River. In 1868 the important workings ('river diggings') on the Vaal River were commenced; and in 1870-1871 the diamonds were traced to deposits of an unique type, in the neighbourhood of which the town of Kimberley in Griqualand West, Cape Province, very quickly sprung up. Here are situated the famous mines De Beers, Kimberley, Dutoitspan, Bultfontein, and Wes-Other mines of the same type are the Jagersfontein and Koffyfontein mines in Orange Free State, and the great Premier mine (discovered in 1902) near Pretoria in the Transvaal. The diamond-bearing rock of these mines fills volcanic pipes, 200-300 yards (in the case of the Premier mine half a mile) across, penetrating vertically the surrounding beds of basalt, shale, diabase, and quartzite, and extending to un-known depths The rock, known as kimberlite, or locally as 'blue ground,' consists mainly of a hydrated magnesium silicate with much the same composition as serpentine It contains boulders and broken fragments of various rocks (basalt, shale, sandstone, quartzite, granite, mica-schist, eclogite, &c ) and minerals (ilmenite, pyrope, enstatite, chrome-diposide, iron-pyrites, &c), and is of the nature of a volcanic agglomerate. Diamond is present in this rock on an average of one part in forty millions, and in the richest part of the Kimberley mine one part in two millions Numerous pipes of a similar character, but containing few or no diamonds, are scattered over a wide area in South Africa The diamond was brought from below up with the igneous material, which must have been derived from a basic magma rich in olivine. Although the origin of diamond has been the subject of much discussion, there seems no reason to doubt but that it crystallised from this basic magma The rare presence of diamonds embedded in boulders of eclogite (a garnetpyroxene rock) found in the blue ground has led to the suggestion that this is the motherrock; but it is quite conceivable that the mineral crystallised from more than one kind of basic magma

The early open workings ('dry diggings') at the Kimberley mines were soon replaced by a regular system of underground mining. The blue ground, when brought to the surface, is either directly crushed or is spread on floors and exposed to the action of the weather for about a year, when it is crumbly enough to enable the heavy minerals to be separated by washing. From this concentrate the diamonds are separated by means of a mechanical sorting table coated with grease, to which the diamonds adhere, whilst the other minerals pass over. DIAMOND. 501

solution of caustic soda and in aqua regia or hydrofluoric acid Diamonds also occur in gravels in the Somabula Forest in Southern Rhodesia, and kimberlite pipes have been described as present in this region. In 1908 large numbers of small diamonds were found in the sand-dunes on Luderitz Bay in South-West Africa; in this case the source of the diamonds have not yet been traced. The output of South African diamonds amounted in 1913 to over five million carats (approximately one ton), valued at 11,389,807l (On South African diamonds, v. G. F. Williams, The Diamond Mines of South Africa, 2nd edit. New York, 1907, P. A. Wagner, The Diamond Fields of Southern Africa, Johannesburg, 1914.)

Other diamond-producing countries of less importance are Borneo, New South Wales, and British Guiana. Finds have been reported from many of the states of North America, in British Columbia, Mexico, Dutch Guiana, Russian Lapland, the Ural Mountains, Siberia, China, Western Australia, South Australia, Queensland, Tasmania, Belgian Congo, Gold Coast, and Liberia. In most of these cases the mineral has been found as a few scattered stones in alluvial gold washings. But in Arkansas and British Columbia it has been found in an altered (scrpentinised) peridotite of much the same character as the South African' blue ground'


The presence of diamond in certain meteorites is of special interest. It was first observed in the stone which fell on September 4, 1886, near the village of Novo-Urei on the Alatyr river in Govt Pensa, Russia This stone is composed of olivine (675 pc) and augite (238 pc), together with nickel-iron, troilite, chromite, black carbonaceous matter, and about 1 pc of diamond, the last as minute greyish grains As dull black grains it has also been found in the meteoric stone of Carcote, Chili In several, though not in all, of the masses of meteoric iron from Cañon Diablo in Arizona, it has been found as colourless crystals, and as black and colourless grains, the largest 1 mm across It is also recorded from the meteoric iron of The cubic form of graphite Magura, Hungary (cliftonite), probably a pseudomorph after diamond, has been found in the meteoric irons of Youndegin (Western Australia), Smithville (Tennessee), Crosby's Creek (Tennessee), and Magura (Hungary)

Origin and Artificial Production —The occurrence of diamond in acid igneous rocks and in quartz-veins is very doubtful. On the other hand, its presence in basic igneous rocks is well established; eg in kimberlite and eclogite in South Africa, in serpentinised peridotite in Arkansas and British Columbia, and in diabase in New South Wales; whilst in meteoric stones the matrix is also of the same general character In these cases the diamond is very probably present as a primary mineral, and one of the first constituents to crystallise out from the In addition, we have basic silicate magma the occurrence of diamond in the nickel-iron of meteorites This may be regarded as a magma of a still more basic type, and it is thus probable that diamond may crystallise from basic magmas of widely varying composition

Many attempts have naturally been made to constrain the abundant element carbon to

crystallise as the rare and valuable diamond. but, as yet, with no great measure of success. The results have in many cases been doubtful, and have often not been confirmed. The products are always microscopic and their identification a matter of difficulty In some cases the very hard carbides of silicon, &c , have probably been mistaken for diamond The experiments of I Friedlander (1898), and of R. von Hasslinger (1902-1903), in which carbon was dissolved in molten magnesium silicate (olivine or kimberlite), bear some relation to the natural occurrences, but doubts have been expressed as to whether the microscopic crystals so obtained were really diamond E de Boismenn (Fabrication synthétique du diamant, Paris, 1913) states that he obtained crystals up to 2 mm across by the electrolysis of fused calcium carbide The only method, however, that has been amply confirmed is that first used by R S Marsden (Proc R. Soc. Edinburgh, 1881, x1 20) and elaborated by Moissan (1893-1896, Le Four électrique, Paris, 1897, and The Electric Furnace, London, 1904), in which carbon is dissolved in silver or iron at a high temperature and the mass suddenly cooled by immersion in water or lead A contracting crust is so formed and the molten interior subjected to great pressure Sir C A Parsons (Bakerian Lecture, Phil Trans 1919, A 220, 67, and J Inst Metals, 1918, xx) from a long series of experiments, however, concludes that pressure is not a necessary condition, and that the diamond has been formed from gases, probably carbon monoxide, occluded in the iron. O. Ruff (Zeitsch. anorg Chem 1917, 99, 73), in a systematic repetition of the various methods, also obtained negative results except by Moissan's method The presence of microscopic diamonds in certain kinds of hard steel should finally be noted.

Applications—The value of diamond as a gem depends on its high degree of hardness, enabling it to resist wear, and its high refractive and dispersive power. These optical characters are, however, only brought into full play when the stone is faceted in a particular manner;



and for this reason, old stones cut by Indian lapidaries have often been re-out. The form most suited for this purpose is the brilliant-cut, and stones so cut are known as brilliants. The rose-cut, or rosette, is only used for quite small stones usually mounted in number as a surround. The accompanying figures show a brilliant

502 DIAMOND.

viewed on the large front facet, or table, and a side-view, and the rose-cut, viewed from above and the side. Being the hardest known material, diamond can only be worked very slowly and tediously with its own powder. The groundform of the brilliant is approximately an octahedron, and this can be obtained, if necessary, by cleavage, or more safely by the modern method of sawing with a thin metal disc chaiged with diamond dust. The brilliant is then roughly shaped out by jubbing two diamonds to other—a process known as bruting. The final grinding and polishing of the facets is performed on a rapidly revolving cast-iron disc fed with diamond dust and olive-oil.

The large historical diamonds with their associations of mystery and romance are mostly of Indian origin, but these are now eclipsed by the numerous, much larger stones found in The largest Indian stone still South Africa preserved is the 'Koh-i-noor,' which was re-cut in England in 1852 as a brilliant of 106 to carats 1 (=21 786 grams). The largest Brazilian stone, the 'Star of the South,' found in 1853, weighed in the rough 254½ 'carats,' and when cut 125 'carats' The largest known crystal of diamond is the 'Cullinan,' which was found in 1905 in the Premier mine, Transvaal This showed a large cleavage surface, and was only a portion, probably about half, of the complete crystal The portion as found weighed 6212 grams (3106 metric carats, or nearly 1 lb 6 ozs avoirdupois) Being too large to cut as a single stone, it was cleaved and cut as mine larger gems (weighing from 106 04 to 0 88 grams) and ninety-six small brilliants, the yield of cut material amounting to 34½ p.c. The 'Cullman,' although the largest crystal, is not the largest piece of diamond that has hitherto been dis-A mass of carbonado weighing 631 9 grams was found in 1895 in Bahia, Biazil

As an abrasive, diamond powder is the most effective agent available It is the only material with which diamond itself can be worked, and it is much used by lapidaries for cutting other hard stones A thin disc of soft iron charged on the edge with diamond powder is used for sawing stones, both for ornamental purposes and in the preparation of thin microscopic sections of rocks. The small steel tools used by engravers of cameos and intaglios, and also the steel points used for boring precious stones, &c, are charged with diamond dust. The material used for abrasive purposes, though included in the trade under the term bort, consists largely of the waste material from shaping the better stones as brilliants, and also the less clear and spotted crystals not suitable for cutting as gems It is crushed in a diamond-mortar, splinters suitable for other purposes being picked out under a magnifying-glass, and the remainder reduced to a fine powder

The glazier's diamond is a crystal or fragment bounded by two rounded crystal-faces meeting in a curved, or cusped, cutting edge For writing, drawing, and engraving (e.g. fine scales and rulings) on glass, metal, stones, ivory, &c) fine splinters are used, or the diamond may be cut to a rounded point or chisel-edge. Diamond points are also used for drilling hard stones, glass, porcelain, teeth &c.

drilling hard stones, glass, porcelain, teeth, &c Carbonado and bort, by reason of their absence of cleavage, are much more suitable than ordinary diamond for mounting as small fragments (2-3 carats in weight) as cutting tools which may be subjected to shocks These are mounted in the crowns of rock-drills, used in tunnelling, mining, and sinking bore-holes for artesian wells, &c , and on the edges of blade, band, circular, or wire saws for sawing large blocks of marble, granite, &c Fragments, either in a rough form, or fashioned to a suitable shape, are mounted as lathe tools for working stone, metal, &c For example, for turning electric-light carbons, the edges of watch-glasses, finishing accurate turning work of the hard steel axles of instruments and machines of precision, boring cannon, forming the sharp-edged furrows on the gunding surfaces of millstones, for dressing emery wheels, &c

Small cleavage plates or flat crystals of diamond drilled from each side with fine conical holes are used for drawing fine wire, down to 0.01 mm in diameter. For the tantalum, tungsten, and osmium filaments of electric lamps the metal is forced by pressure through the hole. Bored diamonds are also used for the pivot-supports of delicate instruments, such as chronometers, electric meters, &c.

The application of diamond for optical purposes is limited by the high cost and difficulty of working Diamond lenses would need to have a much slighter curvature than those of glass, and they would be immune from scratching Microscope objectives with such lenses were constructed by Pritchard under the direction of C. R. Goring, in 1824–1826. A hemisphere of diamond would also increase considerably the range and durability of refractometers.

References — In addition to the references quoted in the several sections above, the following are given as being of a more general character M Bauer, Edelsteinkunde, Leipzig, 1896, 2nd edit 1909, Eng. transl (Precious Stones), by L J Spencer, London, 1904; C. Hintze, Handbuch der Mineralogie, Leipzig, 1898, vol. 1; C Doelter, Handbuch der Mineralchemie, Dresden and Leipzig, 1912, vol. 1; Sii W. Grookes, Diamonds, London, 1909, Sir H A Miers, Diamond, article in Encyol Brit 11th edit 1910, E Boutan, Le Diamant, Paris, 1886, E W Streeter, The Great Diamonds of the World, London, 1882, W R Cattelle, The Diamond, London, 1911; Annual Reports on Precious Stones, by G. F Kunz, The Mineral Industry (New York), and by D B. Sterrett, in Mineral Resources, United States Geol Survey

DIAMOND BLACKS, -FLAVINE, -GREEN, -YELLOWS v. Azo- colouring matters.

DIANA. The alchemistic name for silver DIANIL BLACK, -BLUE, -PONCEAU, -YELLOWS v. Azo- COLOURING MATTERS;

o-DIANISIDINE v. DIFHENYL.
DIANISIDINE BLUE v. Azo- colouring
MATTERS.

¹ This was the old English carat of 205 409 mg A later value of the English carat, as defined by the Board of Trade in 1888 and 1889, was 205 304 mg The carativeight has now been standardised in all countries as the metric carat of 200 mg (5 carats = 1 gram) This is the only legal value of the carat in the British Isles since April 1, 1914

DIASTASE. 503

DIANOL BRILLIANT RED, -RED v Azo-COLOURING MATTERS.

DIANTHINE v AZO- COLOURING MATTERS DIAPHTHERIN, DIAPHTHOL, DIASPIRIN, SYNTHETIC DRUGS

DIARSENAL. Canadian name for salvarsan DIASPIRIN. Trade name for succinyl salicylic acid.

DIASPORE. A native hydrated alumina, Al<sub>2</sub>O<sub>3</sub> H<sub>2</sub>O or AlO(OH), crystallising in the orthorhombic system and isomorphous with goethite and manganite. It is usually found as scales or foliated masses with a perfect cleavage in one direction on which the lustre is pearly but it is readily distinguished from other tohated minerals (mica, &c) by its much greater hardness  $(H = 6\frac{1}{2} - 7)$ . Sp gr 34 When heated before the blowpipe it decrepitates violently, breaking up into pearly white scales (hence the name diaspore, from  $\delta i\alpha\sigma\pi\epsilon i\rho\epsilon\nu$ , to scatter) In colour, it is usually grey or yellowish-brown, but pale-violet crystals are also found usually found as an alteration product of corundum and emery, but occurs also as a mineral of contact-metamorphic origin in crystalline limestones It is of importance as a constituent of bauxite, and in a particular type of bauxite, called diasporite, occurring in crystalline limestone in the Bihar Mountains, Hungary, it constitutes the bulk of the material

DIASTASE or Amylase is the name given to the enzyme or mixture of enzymes which converts starch and glycogen into maltose. In France, the word 'diastase' is used as a general term for all enzymes, a custom which causes considerable confusion, and which it is to be hoped will lapse.

Diastase is very widely distributed in plants and in animals—it is found in blood serum, in muscle, liver and panereas, and particularly in the saliva, this latter enzyme being long known under the name ptyulin

The power of genminated bailey to hydrolyse starch was discovered by Kirchoff in 1814, and the enzyme preparation was first made by Payen and Persoz in 1833. Diastase is found very generally in the leaves and twigs, in the germinating seeds, in the bark, in the pollen grains, and in the latex of plants. It occurs in yeast and in most of the mould fung, the enzyme of Aspergillus orizæ, the so-called Taka duastase being well known on account of its industrial use in the manufacture of the Japanese beverage, koji.

Diastase, like all other enzymes, has not been prepared in anything approaching a pure state. Considerable uncertainty exists also as to whether it is a single enzyme or a mixture of enzymes acting in turn to convert starch into dextrin and dextrin into maltose. Modern evidence favours the view that it consists of a starch-liquefying enzyme or amilopectase, and a saccharifying enzyme able to hydrolyse soluble starch, but without much action on raw starch until this has been rendered soluble by the amylopectase. The two enzymes are found together, but in unequal proportions. Wheaten flour, for example, frequently has a deficiency of liquefying enzyme

There are well-marked differences between the enzymes present in resting barley and in malt (cf. Lintner and Eckhardt, Zeitsch. Ges. Brauw. 1899, 389; J. L. Baker, Chem. Soc. Trans. 1902, 81, 1177), one of the chief of these being that the disatase of raw grain possesses little if any starch-liquefying power. Baker has shown that the products of hydrolysis of starch by alcohol-precipitated enzymes from the two sources, are not the same. Brown and Morris have termed the enzyme of resting barley 'translocation disatase,' that of malt 'secretion disatase' (Chem Soc Trans 1890, 58, 458. 1893, 63, 604) They regarded these as not only different in their actions, but as formed from different materials. This latter assumption has been disputed (cf. Ling, J. Inst. Brewing, 1908, 14, 494, Ford and Guthrie, Ic., 1908, 14, 61)

Some authors consider the saccharifying enzyme to be a mixture, attributing a different enzyme to each stage in the simplification of the starch molecule. It is impossible at present to make any definite statement on this point. In support of it, it appears that exposure to a temperature of 68° acts more injuriously on the conversion of dextrin to maltose than on that of starch into dextrin. Heating diastase does not affect the liquiditying power, though the saccharifying action largely disappears

The quantity of diastase contained in the bailey grain increases gradually in the course of germination—it is said to reach a maximum in about 8 or 9 days, when the plumules are twice as long as the grains.

Diastase as it exists in cereals is partly present as an insoluble zymogen in combination with protein substances which are capable of attack by pepsin; in solution the enzyme is free and unaltered by pepsin. The dissolved enzyme is more active as the amount of nitrogen in solution increases

Preparation —A great variety of methods have been proposed for the preparation of diastase, the majority of these starting from malt According to O'Sullivan (Chem Soc Trans 1884, 45, 2), finely ground pale barley malt is saturated with water, adding enough to slightly cover it and after 3-4 hours as much of the solution as possible is expressed by means of a filter press. After filtering, alcohol (sp gr. 0 83) is added, as long as a flocculent precipitate forms, the addition being discontinued as soon as the liquid becomes opalescent or milky. The precipitate is collected, washed with alcohol of increasing strength until dehydrated, pressed between cloth and dried in vació over sulphuric acid.

Lintner (J pr Chem 34, 378) treats 1 part of green barley malt or flour of air-dried malt with 2 to 4 parts of 20 pc alcohol for 24 hours, and precipitates the extract with 2 the volumes of absolute alcohol. The precipitate is washed with alcohol and ether to dehydrate it, and dried in ració

Loew extracts germinated barley with 4 p c. alcohol, and precipitates the extract with basic lead acctate. The precipitate is suspended in water, lead removed with hydrogen sulphide, and the enzyme precipitated with a mixture of alcohol and ether.

Wroblewsky extracts finely powdered malt, first with 70 pc and then twice with 45 pc alcohol Alcohol is added to the two last extracts until 70 pc is present, the precipitate formed is collected, washed with alcohol and ether, and dried.

For further purification, both Wroblewsky

504 DIASTASE.

and Osborne and Campbell dissolve the enzyme in water, and precipitate it with ammonium sulphate, which is added drop by drop, until the liquid contains 50 p.c. On standing, a very active yellowish flaky precipitate is obtained

Green barley malt may also be extracted with glycerol, and the diastase precipitated by alcohol Effront extracts with water and induces alcoholic fermentation in the extract so as to remove carbohydrate and protein matter. It is then filtered and the diastase precipitated by alcohol

Animal diastase is best obtained from saliva Cohnheim uses freshly prepared calcium phosphate as precipitant, extracts this precipitate with water, and adds alcohol to obtain the enzyme. Alcohol is also used for direct precipitation, likewise ammonium sulphate. Von Wittich extracts pancreas diastase with glycerol.

For the preparation of an active diastatic extract for laboratory use, ground malt should be extracted with twenty times its weight of distilled water at 21° for 3 hours and filtered bright.

The diastase of Aspergillus origae is sold in commerce as Taka-diastase. A pure form can be prepared from the commercial article by extracting with water, precipitating with ammonium sulphate, dissolving the precipitate in water, dialysing and finally precipitating fractionally with alcohol

Preparations of diastase obtained as described above are colourless, amorphous, and friable, readily dissolving in water, but if allowed to dry without being dehydrated, they are horny and not easily penetrated by water. They invariably contain ash, which consists of potassium, calcium, and magnesium phosphates, with a trace of sulphate. They give a deep blue colour with a 1 pc alcoholic solution of gum guaiacum and hydrogen peroxide.

Linther's diastase gave the reactions of the proteins, and was found to contain C, 46 66 pc; H, 7 35 pc; N, 10 42 pc; S, 1 12 pc, calculated on the ash-free product. Osborne (Ber. 1898, 31, 254) found for his most active preparation, which had a diastatic power of 600 and contained 0 66 pc of ash, C=52 5 pc, H=672 pc, N=161 pc, S=19 pc; and Wroblewsky gives N=165 pc for his purest diastase. Both authors regarded diastase as a protein combined, according to Osborne, with a proteose

The most recent work of Frankel and Hamburg (Betr Chem Physiol n. Path, 1906, 8, 389) leads to the conclusion that diastase contains neither protein groups nor reducing carbohydrates in its molecule. Impurities were removed from the enzyme solution with lead acetate, the solution sterilised by filtration, and further purified by fermentation with a yeast which had been rendered nitrogen-hungry. This preparation could be divided by dialysis into two distinct enzymes, one of which—the saccharifying—passed through the paper, while the other—the liquefying—was left behind. It showed absence of typical protein reactions, but contained a pentose in small amount.

The purified diastase preparations obtained by Pribram (Biochem Zeitsch 1912, 44, 293) from malt extract contained 77 p c of nitrogen and 15 p c of ash; one-sixth of the nitrogenous matter separated as a coagulum on heating. The filtrate on hydrolysis with sulphuric acid

yielded a reducing substance which was considered to be a polycarbohydrate acid which existed in the enzyme in combination with a simple polypeptide. The purified diastase was inactive, but is activated by the addition of traces of lactic acid.

Particularly active preparations have been obtained by Sherman and Schlesinger (J. Amer. Chem Soc 1913, 35, 1617) by extracting ground malt with two and a half times its weight of water or very dilute acid phosphate solutions at as low a temperature as possible, with subsequent dialysis, followed by fractional precipitation with alcohol or acetone. The products obtained had a diastatic power of 1800–2200 on the Lintner scale

The most active preparations contained most nitrogen—about 14 pc in maximum. When tested by the Van Slyke method they contained combined nitrogen in all the usual eight forms, the proportions being within the range of variation shown by typical protein substances. The optimum hydrogen ion concentration expressed by Sorensen's exponent was between  $p_{\rm H}^+$  4 2 and 4·6 (J Amer. Chem. Soc. 1915, 37, 623) The purified amylases obtained from the pancreas and from malt are similar in physical properties, show the same qualitative reactions

the pancreas and from malt are similar in physical properties, show the same qualitative reactions and have the same percentage of nitrogen distributed in the same way among the different types of amino acid radicals (J. Amer. Chem. Soc. 1915, 37, 1305)

The activity of diastase is very greatly influenced by the nature of the substrate on which it acts, more particularly as to whether this is alkaline or acid Certain substances. such as the amphoteric amino- acids and asparagine, also ammonium and calcium phosphate, increase the activity, asparagine being especially effective in some instances. Vegetable diastase exhibits the maximum activity in presence of traces of weak organic acids. The very smallest quantities of alkali stop action, though this begins again as soon as the alkali is neutralised. In reality, the above accelerating substances are all of acid nature, and the apparently favourable effect of acid is due not to the acid per se, but to its neutralising alkaline impurities present. Maximum activity is attained in neutral (in the sense that water is neutral) solution. Minute traces of metallic impurities have an adverse influence. The temperature of maximum activity of diastase is about 55°; heated above this, it begins to show decomposition.

Estimation of diastatic power - Since it has not been obtained in a pure state, the absolute estimation of diastase is an impossibility; moreover, the very great influence of external conditions on diastatic activity, to which allusion has been already made, makes it difficult to do more than compare the relative activities of a number of materials under precisely similar As this is an important operation conditions in brewery control, the methods of analysis have been accurately laid down. The diastatic power is often termed the Lintner value, and expressed in degrees Lintner. The standard is 100° when 0 1 c.c. of a 5 p.c. solution of malt, acting on a 2 p.c solution of soluble starch for 1 hour at 70°F, produces reducing sugar equivalent to 5 c.c. of Fehling's (Barreswil's) solution. To obtain accurate results, great stress must be laid on the purity of the distilled water and soluble starch used: commercial starches are frequently alkaline. All work must be carried out in hard-glass apparatus (cf. Ford, J. Soc.

Chem Ind 1904, 23, 8)

The method adopted by the Malt Analysis Committee of the Institute of Brewing (J Inst Brewing, 1906) for malt, is as follows: 25 grams ground malt are extracted with 500 cc, of distilled water for 30 hours at 70°F, and filtered bright 3 cc. is allowed to act on 100 cc of 2 pc. soluble starch at 70°F for 1 hour Diastatic action is then stopped by the addition of 10 cc N/10-sodium hydroxide, and the liquid made up to 200 cc. Sugar is then determined either volumetrically or gravimetrically. The results are calculated by the formula

### Diastatic power=1000/xy

where x=the number of c.c of malt extract contained in 100 c c. of the fully diluted starch-conversion liquid, and y=the number of cc. of the same liquid required to reduce 5 c c. of Fehling's solution. When the diastatic capacity exceeds 50°, 2 c c or less of malt extract are taken, particularly for the malts over 80° used by distillers and vinegar makers. The diastatic power of English brewing malt ranges between 35° and 40°, that of green malt from 110°-125°, showing that a large proportion of the enzyme is destroyed during the kilning

Lintner tests solid preparations of the purified enzyme by dissolving 0.2 to 0.5 gram in 50 c c water and adding to a series of test-tubes, each containing 10 c c of 2 p c soluble starch solution, 0.1, 0.2, &c, up to 1 c c of the enzyme solution. After an hour, each tube is boiled in the water-bath with 5 c c of Fehling's solution, and that determined in which the copper solution is just completely reduced. A second series enables this point to be ascertained with

greater accuracy.

Similar methods have been based on the colourations with iodine, shown by starch and dextrin solutions. Wohlgemuth sets increasing quantities of enzyme in a series of tubes with 5 c c of a 1 p c starch solution under precisely similar conditions. After 30 to 60 minutes at 40°, the tubes are cooled in ice and a drop of N/10 iodine solution added. Colours varying from dark-blue, blue-violet, reddish-yellow, and yellow are obtained. Wohlgemuth selects the tube in which the violet shows as indicating complete conversion to dextrin.

Diastase, like other enzymes which act on carbohydrates, is much affected by the products of its activity with which it forms adsorption compounds—hence the peculiarities exhibited in a starch conversion. Provided that the temperature and other conditions selected are such that the enzyme is not destroyed, the diastase recovered at the end of a reaction is just as active towards a second quantity of starch as an equal portion of fresh enzyme. Diastase is thus a true catalyst (cf. Van Laer, Bull Soc. chim. Belg. 1912, 26, 223).

Uses —Diastase plays a most important part in the brewing, distilling, and baking industries Malt extracts, concentrated in vacus at a low

temperature to viscid syrups, are largely used by bread bakers and in some food preparations. Considerable quantities of such active preparations, made from malts of high diastatic power, also find employment in laundries for the purpose of removing starch from stiff collars, shirts, &c. (See also articles STARCH. FERMENTATION; HYDROLYSIS)

E. F. A.

DIATOMITE (Diatomaceous Earth, Infusorial Earth, Tripolite or Tripoli; Ger Kieselguhr, Bergmehl) A form of opaline silica consisting of the siliceous remains of diatoms (Diatomaceæ) These are unicellular, free-swimming plants belonging to the large group of Algæ. The cell-wall, or frustule, of each minute individual consists of two shields or valves of siliceous material with elaborate and beautiful ornamentation More than 10,000 species of diatoms have been described, they vary considerably in form—fusiform elliptical, discoidal, globular, and accoular Material composed of individuals of the acicular form, the length of which varies from 0 093 to 0 156 mm and the thickness 0 00156-0 00468 mm, is best suited for technical purposes Diatoms multiply extremely rapidly and inhabit both fresh and salt water, being more abundant in cold latitudes On death, their remains accumulate, under certain conditions, at the bottom of swamps, lakes, or the open ocean Besides being formed at the present day, such deposits are also met with in strata of Pleistocene and Tertiary age

Diatomite is a loose, pulverulent material with a mealy or earthy texture, resembling soft chalk or dried clay in appearance. It is friable between the fingers with a harsh feel, and grates between the teeth In colour it is white, grey, yellowish, brownish, or greenish. It is very light, floating on water, sp gr of air-dried material 0.24 and of calcined 0.34. Its high degree of porosity depends on the fact that the shells of the diatoms are hollow material absorbs four or five times its weight of water Further, it is a bad conductor of heat, electricity, and sound, and is resistant to heat and most chemicals. Of acids, only hydrofluoric attacks it, but it is slowly dissolved by a hot solution of caustic alkali Chemically it consists of hydrated silica, mixed with various impurities, such as sand, hydroxides of iron, clayey, calcareous, and organic matter p c. of SiO2 may be present

Analysis I, is of white material (dried at 105°) from Auxillac, Cantal, Auvergne; II, of green material from Auxillac; III, from Loch Cuthir (or Quire), Skye, Scotland; IV, from Richmond, Virginia, and V, of Tripolite from Bilin, Bohemia (including also SO<sub>3</sub> 0 12, P<sub>2</sub>O<sub>5</sub>

0.24, NH<sub>4</sub> 0 03).

	I.	II.	III.	IV.	v
$S_{1}O_{2}$	91.61	88.56	88.73	75.86	74.20
TiO2	0.10	0 18			-
Al <sub>2</sub> O <sub>3</sub>	1.53	2.16		9 88)	6 81
Fe,O3	$2\ 22$	0 73	0.67	2 92∫	
CaO	0.18	0.26	0 04	0 29	041
MgO	0.29	0.02	)		
K <sub>2</sub> O	trace	0 47	}	1.63	0.02
Na,O	trace	0 44	}		0.30
H,Ŏ	2 33	3 01	6 40	8 37	13 30
Organ mat	1.74	4.17	4.15		4 20

506 DIATOMITE.

Tripolite (Ger. Tripel, Polierschiefer=polishing shale) differs from diatomite in being more compact with a laminated structure and consequently less light (sp gr 186) and porous merely represents the same material of a greater

geological (Tertiary) age

The most extensively worked deposits of typical diatomite (Kieselguhr) are those on the Luneburger-Heide (heath) in Hanover beds with a thickness of 10-15 metres occur in The uppermost basins in Pleistocene sands layers are white, below they are grey, and at the bottom of the deposit green. The green material contains from 15 to 30 p.c. of organic matter, sufficient for calcination without A bed of 5-7 metres covered addition of fuel by Tertiary clay and basalt is worked at Vogels-berg near Altenschluf in Hesse Similar deposits of recent to Miocene age are also widely distributed in the volcanic region of central France (Departments of Cantal and Puy-de-Dôme) the material is known as randannite and ceyssatite, from Randanne and Coyssat respectively, and it is worked commercially under the name tellurine (On the French and German deposits, see C Schmidt, Ann des Mines, Paris, 1910, xvii. 370) Deposits are met with in peat-bogs at several places in Scotland, and beds up to 40 feet in thickness of good quality material are now worked at Loch Curthin (or Quire) in the parish of Kilmuir in Skye (W. I. Macadam, Min Mag 1884, vi 87, 1886, vii 30, 35, 1889, viii 135 Special Reports on the Mineral Resources of Great Britain, Mem Geol Survey, 1916, vol v) Similar deposits also occur in Co. Antrim (J H Pollok, Sei Proc R Dublin Sec 1899, ix 33) Extensive deposits, many of them of Tertiary age, are of wide distribution in the United States, especially in California, (H. Ries, Economic Geology, New York, 1916). Other localities are Monte Amiata in Tuscany, Victoria, 1 Norway, Denmark, Hungary, Algeria Chili, &c The typical tripolite forms yellowishwhite bands 0.5-1.5 m thick in beds of clay, gypsum, and 'Saugschiefer' (absorbentshale) in a Tertiary basin near Bilin in Bohemia

In the preparation of the material for the market, that of the purer white quality, after being dug out from open pits, is simply air dried in the open or in sheds Organic matter is removed by calcination. Less pure material containing sand, &c, is treated by washing and sedimentation Iron is removed by the action of acids

The varied uses of diatomite depend on its peculiar properties of porosity, lightness, nonconductivity, resistance to heat and acids It was used by the Greeks and Romans for making light fire-proof tiles, and in the middle ages was mixed with meal for bread, under the impression that it had a medicinal value an absorbent it has been much used with nitroglycerine for the manufacture of dynamite. The so-called dry sulphuric acid is prepared by saturating calcined diatomite with three or four times its weight of acid Such material can be transported with little risk Diatomite packed around carboys of acid will soak up leakage in case the vessels become cracked It has also been used as an absorbent of bromine, dis-

On Australian deposits, see Bull Imp Inst., London, 1916, xiv 40.

infectants, and liquid manures. On account of its non-conductive properties it is used, sometimes with asbestos, as a light packing material for boilers, steam-pipes, cookingstoves, safes, refrigerators and refrigerating apparatus, fire-proof and sound-proof floors and partitions of buildings, bulkheads of ships. Mixed with clay, lime, sawdust, &c, it can be used in the manufacture of fire-proof bricks and tiles, and having a low coefficient of expansion such material is specially adapted for making tubes, models, &c. It is used in the manufacture of various cements and artificial stones. Mixed with magnesia cement it gives a fire-proof cement for stoves, with shellac, scaling-wax and gramophone records are made It is also used as a filtering material for liquids and gases, and various liquids (wine, beer, acids, oil, syrups) can be clarified by stirring with 1 p c of diatomite and allowing to settle. As a filler it is employed in the manufacture of rubber, cotton fabrics, paints, safety matches, paper, papier-mâché, soap, scouring-powders, tooth-powders, &c Fused with calcium and magnesium borates it yields a non-cracking glass for glazes and enamels, which can also be used for painting on glass. As a source of silica, it is used in the preparation of sodium silicate, ultramarine, smalt, &c Finally, as an abrasive (qv) it, particularly the tripolite variety, is used for polishing metals and stoves (Ref. B. Dammer and O. Tietze, Die Nutzbaren Mineralien, Stuttgart, 1913, vol 1, Diatomaceous Earths (Kieselguhr) and their Utilisation, Bull. Imp. Inst London, 1905, m 88-103 L J S

DIAZINE BLACK, -BLUE, -GREEN v Azo-COLOURING MATTERS

DIAZO- COMPOUNDS. These organic substances, like the azo- compounds, contain the characteristic group  $-N_2$ -, but whereas, in the latter series, this bivalent radical is invariably attached to two carbon atoms, CN, C, in the former it is generally, although not invariably combined with only one carbon atom, the other valency being satisfied by (1) an electro-negative 100  $\overline{A}c$  in the diazonium salls  $CN_2\overline{A}c$ , (2) an oxygen atom in the diazohydroxides ('diazohydrates')  $CN_2$  OH and diazo-oxides ('diazotates')  $CN_2$  OR, (3) a nitrogen atom in diazoamines (diazoamino- compounds)

CN, NHR

C·N.·NH·C;

and diazoimines (azimino-compounds)

(4) a sulphur atom in diazosulphonates

CN2 SO3K

and diazosulphides C  $N_2$  S  $N_2$  C The diazocyanides C  $N_2$  CN contain the azogroup attached to two carbon atoms, one of which belongs to a cyanogen group, and these compounds which, in general, are distinctly coloured, form an intermediate group between the foregoing diazo- derivatives and the azo- compounds

Certain aliphatic diazo- compounds (v infra) contain the group -N2- attached by two active valencies to the same carbon atom.

# I. DIAZONIUM SALTS.

These diazo- derivatives are generally, but not exclusively, prepared from aromatic amines, and the process, which is termed diazotisation, is carried out with the undissociated salt of the amine, and not with the free base. In a hydrolysing solvent like water, it is therefore customary to diazotise aniline and its derivatives with sodium nitrite (1 molecule) and excess (2½ to 3 equivalents) of mineral acid.

#### AROMATIC DIAZONIUM SALTS.

The diazotisation of the benzenoid and naphthalenoid amines is frequently carried out on a manufacturing scale owing to the employment of aromatic diazonium salts in the production of azo-colouring matters. More than 1000 tons of para-intrandine are diazotised annually in the production of 'para-intrandine red'.

The aromatic base is dissolved in dilute hydrochloric or sulphune acid (3 equivalents in 10-20 p.c. aqueous solution) and the solution treated with an equivalent quantity (1 molecule) of sodium nitrite in 10-20 p.c. solution. The diazotisation is completed when the liquid contains a slight excess of nitrous acid as indicated by starch and rodide paper.

With anilme and its homologues, the temperature of the diazosolution is generally maintained at 0° 5° to prevent, so far as possible, the decomposition of the unstable diazonium salt, but in other cases, eg the intrainlines, naphthylamines, benzidine, and other diphenyl bases, the diazotisation is carried out most readily at 10°. The amino anthraquinones are diazotised at 30°-40° (Schaarschmidt, Bei 1916, 49, 2678)

In the manufacture of soluble azo- dyes, the diazonium safts are generally employed in aqueous solution, but solid diazo- compounds, in dry or paste form, have been introduced to enable dyers to prepare 'ingrain' azo- colours which are developed on the textile fibres (v. Eng. Pat. 18429, 21227, 1894, 8989, 11757, 1895, 1645, 1896, 15353, 1897, D. R. PP. 85387, 88949, 94280, 1894, 89437, 1896)

In alcoholic or acetic acid solution, diazotisation can be effected on the salt of the amine without using an excess of acid, and ethyl or amyl nitrite may, with advantage, be substituted for sodium intrite. Gaseous methyl nitrite, which has been employed in certain diazotisation (Chem. Soc. Trans. 1913, 103, 1398), is very readily prepared and is a cleaner reagent than amyl nitrite.

Benzenediazonium chloride  $C_gH_g$  N<sub>2</sub>Cl, the simplest example of an aromatic diazonium salt, is prepared by adding the calculated quantity of amyl nitrite to aniline hydrochloride suspended in glacial acetic acid at  $10^\circ$ , it is precipitated from the clear solution thus obtained by the addition of ether

Benzenediazonium sulphate C<sub>6</sub>H<sub>5</sub> N<sub>2</sub> HSO<sub>4</sub> is prepared in a similar manner from aniline sulphate suspended in glacial acetic acid, the precipitation being effected with alcohol and other

Benzenediazonium nitrate C<sub>6</sub>H<sub>5</sub> N<sub>2</sub>·NO<sub>3</sub>, which was first precipitated from aqueous solution by ether and alcohol (Griess, Annalen, 1866, 137, 39), can also be produced in alcoholic or glacial acetic acid solution (Knoevenagel, Ber 1890, 23, 2995; Bamberger, *ibid* 1896, 29, 446, Hirsch, *ibid*, 1897, 30, 92, Hantzsch and Jochem, *ibid* 1901, 34, 3337).

Diazonium salts have been obtained with a large variety of amons—platinichlorides, aurichlorides, stannichlorides, ferrit yanides, nitropriestates (Ber 1879, 12, 2119, 1885, 18, 965), chromates (Jahresberichte, 1867, 915), Chem Soc Trans 1905-87, 1), carbonates and nitrites (ibid, 1906, 87, 921), fluorides (Ber. 1903, 36, 2059), prichlorides (ibid 1906, 39, 2713, 3146), picrates (Chem Soc Trans 1907, 91, 1316), azides (ibid 1910, 97, 1697), Ber 1903, 36, 2056), thioxidiphates, tungstates (J. Soc. Dyers, 1901, 17, 279), thioacetates (Monatsh. 1907, 28,247), benofluorides (D. R. P. 281055), phospho molyblates and tungstates (Chem Soc. Trans. 1909, 95, 1319), arylsulphinates (Ber. 1911, 44, 1415), zinichlorides, Rev. Gén. Mat. Col. 1914, 18, 4, \$\beta\$ naphthol-1-sulphonates, Eng. Pat. 8989, 11757, of. 1895, D. R. P. 93305

All diazonium salts are hable to explode in the dry state, for they are endothermic compounds (Berthelot and Vieille, Compt. rend. 1881, 92, 1076. Sventoslavsky, Ber. 1910, 43, 1479) and in some cases the explosion point has been determined (Ber. 1899, 32, 1691., 1900, 33, 527). The diazonium chromates and perchlorates decompose with considerable violence, so that the former have been suggested as explosives (Bull. Soc. chim. 1867, [2], 7, 270., Fr. Pat. 73286).

Bisdiazonium salts (Tetrazo- salts) These compounds are produced from aromatic diamines which do not contain their aminogroups in ortho- or peri- positions (v. Cyclic diazo- compounds)

When the two amino- groups are in the same aromatic nucleus, complete diazotisation is effected only with some difficulty Metaphenylenediamine hydrochloride and sodium nitute furnish the azo- colouring matters included under the term 'Bismarck brown' (Bei 1897, 30, 2111, 2899, 1900, 33, 2116), but when a solution of the base in excess of hydrochloric acid is added quickly to a large excess of well-cooled aqueous nitious acid, complete diazotisation takes place and benzene-m bisdiazonium chlorede Calla N, Cl. 18 produced When the nitrite is added suddenly to the diamine hydrochloride, nitroso m phenylenediamine results Para-phenylenediamine hydrochloride, when diazotised in the usual way, gives a mixture of diazonium and bisdiazonium chlorides, but when the hydrochloride is added to excess of nitrous acid, benzene-p-bisdiazonium chloride is obtained

In the naphthalene series, 14-naphthylenediamine is diazotised with difficulty, owing to its oxidisability, but in producing its azo- derivatives, this difficulty is surmounted by working with its monoacyl-derivatives and diazotising in two stages with an intervening hydrolysis The heteronucleal (Eng. Pat. 18783, 1891) naphthylenediamines, excepting the 18-compound, can be completely diazotised, yielding naphthalenchisdiazonium salts (Ewer and Pick, D R PP 45549, 45788, Badische Anilin und Soda Fabrik D R P 130475, Morgan and Micklethwait, Chem Soc Trans 1910, 97, 2558) Benzidine, tolidine, dianisidine, and other similar heleronuclear diamines of the diphenyl and ditolyl series are readily diazotised, furnishing bisdiazonium salts (e.g. diphenylbisdiazonium chlorude Cl N2 C4H4 C4H4 N2 Cl), which are employed in the manufacture of substantive azo- dyes (v. Congo red and Benzopurpurin; art. Azo-colouring matters) There are certain substituted para-diamines in the benzene diphenyl and naphthalene series which have hitherto resisted all attempts at complete diazotisation (Ber 1884, 17, 603; 1896, 29, 2285, 1905, 38, 3769; Eng Pat 2946 of 1896)

### VELOCITY OF DIAZOTISATION

Under the usual experimental conditions, diazotisation proceeds with great rapidity, but in N/1000 solutions, the velocity of this reaction has been determined by a colorimetric method for estimating the free nitrous acid The reaction is of the second order, and the values of the constant  $C = \frac{x}{t(a-x)}$  are 0.036 for aniline, 0 038 for p-toluidine, and 0 041 for m-xylidine, with one molecular proportion of free acid at 0°. At higher temperatures, the rate is increased (Hantzsch and Schumann, Ber 1899, 32, 1691; 1900, 33, 527; Tassilly, Compt rend 1913, 157, 1148; 1914, 158, 335, 489) Both colorimetric and electrolytic determinations of the velocity of diazotisation show that, in the absence of disturbing influences, all the aromatic amines are diazotised at the same rate.

# ALTERNATIVE METHODS OF PRODUCING AROMATIC DIAZONIUM SALTS

Nitrosyl chloride or bromide has been suggested as a diazotising agent (Tilden and Millar, Chem Soc Proc 1894, 59; Kastle and Keiser, Amer Chem J 1895, 17, 91) Nitrosyl sulphate has been used with advantage (Pabst and Girard, D R P 6034; Ber 1879, 12, 365, Morgan and Micklethwait, Chem Soc Trans 1910, 97, 2558) This reagent or sodium nitrite dissolved in concentrated sulphuric acid is employed in diazotising slightly basic amines (Morgan and Evens, Chem. Soc. Trans. 1919, 116, 1130, cf. J. pr. Chem. 1906, [ii.] 74, 275) Barium nitrite, when employed in conjunction with sulphuric acid, furnishes diazo- solutions free from inorganic salts (Witt, Ber 1903, 36, The last-named investigator has brought forward an interesting process which consists in dissolving the base in cold concentrated nitric acid and adding the calculated amount of potassium metabisulphite required to produce sufficient nitrous acid for the diazotisation (Ber 1909, 42, 2953)

Diazonium bromides are formed by the inter action of hydrazines and diazonium perbromides

$$2R N_2Br_3+RN_2H_3=3R N_v Br+3HBr$$

(Chattaway, Chem Soc Trans 1908, 93, 958) An electrolytic process for the production of diazonium salts has been patented (Boehringer & Sons, D. R. P. 152926, and Eng. Pat. 2608 of 1904)

Diazonium salts can also be prepared from compounds not containing amino- groups; mercury diphenyl and nitrous fumes give benzenediazonium nitrate (Ber 1897, 30, 509), and the same salt is produced from nitrosobenzene and nitric oxide in chloroform solution (*ibid* 1898, 31, 1528) Fuming ritric acid reverses the coupling process and regenerates the diazonium salt from its azo- derivatives (Meldola, Chem. Soc Trans 1889, 55, 608; 1894, 65, 841; and O. Schmidt, Ber. 1905, 38, 3201).

Diazonium chlorides are regenerated from azo-dyes by decomposing the latter with chlorine or hypochlorous acid (J. pr. Chem. 1912, [n.] 85, 235.

Certain complex or highly substituted amines resist diazotisation unless treated under pressure (D R P. 143450) or with excess of nitrous and mineral acids (Annalen, 1873, 165, 187; 1891, 260, 224; Ber 1900, 33, 520)

(For indirect methods of forming diazonium salts, v Annalen, 1879, 199, 320; 1888, 243, 282; 1892, 270, 116; Ber. 1893, 26, 2190, 1899, 32, 1809; 1900, 33, 1718; J pr. Chem. 1896, [Ii.] 54, 496; Gazz chim. ital. 1900, 30, 1526; Bull. Soc. ehim. 1891, [Iii.] 6, 94)

#### REACTIONS OF AROMATIC DIAZONIUM SALTS.

In addition to their power of coupling with phenols, aromatic amines, and their sulphonic acids to form the technically important azocolouring matters, the diazonium salts are synthetical agents of the greatest value, and the following is a synopsis of the principal reactions in which they take part:—


l Decomposition of Diazonium Salts by Water.

(Cf Rer 1874, 7, 1061; 1894, 17, 2704; 1899, 32, 1773; 1890, 23, 3635; 1902, 35, 89, J pr Chem 1881, [n | 24, 476, Chem Soc Trans, 1902, 81, 77; 1903, 83, 688; 1905, 87, 5; 1906, 89, 19.)

The decomposition of soluble diazonium salts by water is a unimolecular reaction, and is represented by the well-known equation.

$$C = \frac{1}{t} \log_{A-x}^{A}$$

The relative stability of the diazonium salts is ascertained by comparing the values of the constant C for a given temperature. The most successful method of making this determination consists in heating solutions of the diazo-compounds at a fixed temperature, and measuring the rate at which nitrogen is evolved. A convenient apparatus, which can be used in this determination and also for studying the decomposition of diazo-compounds in general, is indicated in the following figure.



These quantitative decompositions show that the diazonium salts from aniline and orthoand meta-toluidine are among the least stable of the series, whereas considerably greater stability is exhibited by the diazonium sulphonates derived from the benzenoid and naphthalenoid aminesulphonic acids.

The rate of decomposition of diazo- comunds increases with the temperature and in cordance with Arrhenius's formula for the nperature coefficient

$$\mathbf{C}t_1\!=\!\mathbf{C}t_0\epsilon^{\mathbf{A}(\mathbf{T}_1-\mathbf{T}_0)/\mathbf{T}_1\mathbf{T}_0}$$

The rate of decomposition is independent of e quantity of mineral acid present, and equivait solutions of benzenediazonium chloride, omide, sulphate, nitrate, and oxalate decomse with the same velocity; this change is, wever, accelerated by colloidal platinum or ver, owing to catalytic action (Hantzsch, Ber. 00, 33, 2517; Hausser and Muller, Bull Soc. m. 1892, [m] 7, 721; 1893, 9, 353; Compt. d. 1892, 114, 549, 669, 760, 1438; Schwalbe, itsch. Farb. Ind 1905, 4, 433; Ber 1905, 38, 96, 3071; 1909, 42, 196; Cam, Chem. Soc. ans 1902, 81, 1412; 1903, 83, 206, 470, 1905, 2511; Euler, Annalen, 1902, 325, 292; Voley d Cam, Trans. Faraday Soc. 1909, 5, 1)

In general, the decomposition of a diazonium It by water leads to the formation of a phenol

$$R N_2 \cdot HSO_4 + H_2O = R OH + H_2SO_4 + N_2$$

The sulphate is the best salt to employ in is reaction, because the chloride and nitrate e apt to lead respectively to alternative or condary changes. Benzenediazonium sulate and its homologues furnish phenol and e cresols and xylenols; diphenyl-4 4'-bisazonium sulphate gives rise to 4 4'-dihydroxyphenyl Aurin is similarly obtained from azotised rosaniline (Annalen, 1878, 194, 306)

In certain cases, more especially in the phthalene series, the product is contaminated azo- compounds, resulting from the coupling the hydroxy- derivative with a portion of the ill undecomposed diazonium salt (Chem Soc ans 1903, 83, 221) To obviate this difficulty, far as possible, excess of acid is used in the

composition The manufacture of the aphtholsulphonic acids (e.g. OH SO<sub>3</sub>H=1 4 id 1.8) from the corresponding a-naphthylninesulphonic acids furnishes typical examples the technical application of the process

Secondary reactions occur even in the simest cases (Ber 1890, 23, 3705), and in many stances, where the boiling is protracted owing the great stability of the diazo- compound, t a trace of the hydroxy- derivative can be olated (Amer Chem J. 1889, 11, 319, Cam, iem Soc Trans 1903, 83, 688) When the ienolic product is volatile in steam, the normal ange is facilitated by operating with solutions uling above 100° A good yield of guaracol is stained by adding the diazo- salt of orthousidine to a mixture of sodium sulphate and lute sulphuric acid boiling at 135°-145°, the

olatile product being removed at the moment formation (Kalle & Co, Eng. Pat 7233 1897; cf. Annalen, 1889, 253, 281; Ber 1874, 1008; D. R. P 167211, Chem Soc Trans. 06, 87, 19; cf. Willstatter and Schudel, Ber. 118, 51, 782.

(For other exceptional decompositions of azo- compounds, v. Ber. 1874, 7, 1061; 1884, , 2704, 1899, 32, 1773; 1902, 35, 89; Chem. oc Trans 1896, 69, 1327; 1902, 81, 77, 1905; , 5; 1906, 89, 19).
The diazo- compounds are very sensitive to

tht, which induces the phenolic decomposition

even in certain cases where the action of hot water or acids is very slight (Chem Soc Trans 1907, 91, 35; cf. Compt. rend. 1881, 92, 1074; Ber. 1890, 23, 3131; 1901, 34, 1668; D. R. PP 53455, 56606; J. Soc. Chem Ind. 1890, 9, 1001) 2. Decomposition of Diazonium Salts by Alcohols.

The reaction between diazonium salts and alcohols may proceed in two different ways. 1 R N<sub>2</sub> X+C<sub>2</sub>H<sub>5</sub> OH=R H+C<sub>2</sub>H<sub>4</sub>O+N<sub>2</sub>+HX 2 R·N<sub>2</sub>·X+C<sub>2</sub>H<sub>5</sub>OH=R O C<sub>2</sub>H<sub>5</sub>+N<sub>2</sub>+HX

The former of these changes was originally regarded as the general reaction, but more recent researches have shown that the second is the The factors determining the normal change course of the reaction are, however, numerous and complicated, involving (1) the nature of the alcohol; (2) the influence of substituents in the diazonium salt, (3) the pressure and temperature at which the decomposition is affected.

The tendency to replace the diazo- group by hydrogen increases with the molecular weight of the alcohol employed. Benzenediazonium chloride or sulphate and methyl alcohol give anisole exclusively With ethyl alcohol, these salts yield chiefly phenetole, but benzene is also pro The diazonium sulphates from p-chloroduced and p-bromo-aniline furnish only p-chloro- and p-bromo benzene with ethyl alcohol, but with methyl alcohol they give use mainly to the methyl ethers of p-chloro- and p-bromo-phenol (Amer Chem J 1898, 20, 229, cf Annalen, (Amer Chem J 1898, 20, 229, cf Annalen, 1871, 159, 191, Ber 1901, 34, 3337, 1903, 36, 2061, Amer Chem J 1893, 15, 518; 1897, 19, 531, 547, 561)

The diazo- salts of the simpler amines (aniline and its homologues) react chiefly in the direction indicated by the second of the foregoing equations, and yield mixed ethers, but the presence of acid substituents

# (Cl, Br, NO<sub>2</sub>, CO<sub>2</sub>H, NH SO<sub>2</sub>R)

favours the predominance of the first change. The intramlines (ortho-, meta-, and para-) give respectively 87, 51, and 40 pc of introbenzene the 1 2-, 2.1-, and 1 4-intronaphthylamines furnish the corresponding nitronaphthalenes, whereas  $\alpha$ - and  $\beta$ - naphthylamines yield  $\alpha$ - and β-ethoxynaphthalenes (Remsen and others, Ber 1885, 18, 65, Amer Chem J 1886, 8, 243, 1887, 9, 587, 1891, 13, 153, 1892, 14, 15)

The yield of alkyl ether increases with rise of pressure, but the addition of zinc-dust or an alkalı (sodium hydroxide or ethoxide, potassium carbonate) inhibits almost entirely the formation of mixed ether and leads to the substitution of the diazo- group by hydrogen (Amer. Chem J 1893, 15, 105, 301, 320, 379; 1894, 16, 235, 1895, 17, 454; 1897, 19, 163, 531; 1898, 20, 298; 1904, 31, 119)

3. Replacement of the Diazo- Group by

Hydrogen.

In addition to the action of alcohol, the diazo- group can be eliminated by the use of a variety of reducing agents

(1.) The diazonium salt is reduced to the corresponding hydrazine and the hydrazino- group NH NH, removed by boiling with copper sulphate (Baeyer and Pfitzinger, Bei 1885, 18, 90, 786)

(ii) Reduction of the diazo- compound with alkuline sodium stannite (Friedlander, Ber. 1889, 22, 587).

(iii) By various other reducing agents, stannous formate (Pci 1885, 18, 357, J pi Chem 1889, [ii ] 40, 97), hypophosphorous acid (Ber 1902, 35, 162), sodium hyposulphite (ibid 1907, 40, 858), formic and and copper powder (ibid 1890, 23, 1632), aluminium powder in

alcohol (Morgan and Evens, lc).

4 Replacement of the Diazo- Group by Halogen, Cyanide, Cyanate, and Thiocyanate Radicals

Fluorine may be introduced into the atomatic nucleus in place of the diazo- group by treating the diazonium salt with hydro-fluoric acid Diazoamino- compounds (v. infra) are also employed in bringing about this replacement, and when treated with concentrated hydrofluoric acid they yield the corresponding fluoro- derivative, fluorobenzene is thus produced from diazoaminobenzene or diazobenzenepiperidide (J. pr. Chem. 1870, [n. ] 1, 395., Ber. 1875, 8, 893, 1879, 12, 581, 1889, 22, 1846)

The first really successful process for replacing the diazo- group by chlorine (or bromine) was discovered by Sandmeyer, it consists in adding the diazotised solution to cuprous chloride (or bromide) dissolved in warm hydrochloric (or hydrobromic) acid, when nitrogen is evolved and the corresponding chloro- or bromoderivative is produced (Bct 1884, 17, 1633, 2650, 1885, 18, 1492, 1496, 1886, 19, 810, 1890, 23, 1628, 1880, Annalen, 1893, 273,

The first stage in the Sandmeyer reaction is considered to be the formation of an additive compound of diazonium salt and cuprous chloride, the decomposition of this intermediate product proceeds readily only above a certain temperature, which is different for each diazonium salt (Eidmann, Annalen, 1893, 272, 141, of Votoček, Chem Zeit Rep. 1896, 20, 70, Hantzsch and Blagden, Ber. 1900, 33, 2544) The substitution of biomine for diazonitrogen follows a similar course (Phil Trans 1864, 154, 673, Annalen, 1866, 137, 49, Amer Chem J 1891, 13, 486, Gazz chim ital 1890, 20, 631)

The foregoing substitutions are also effected by the addition to solutions of diazonium chlorides or bromides of finely divided copper or the copper bronze of commerce (Gattermann, Bei 1890, 23, 1218, 1892, 25, 1071, Ullmann, ibid 1896, 29, 1878, if Angeli, ibid 1891, 24,

Iodine is readily substituted for the diazo complex by treating the acid diazo- solutions with potassium iodide even in the absence of

copper or cuprous salts

The Sandmeyer reaction includes the important synthetical operation of replacing the diazo- group by a cyande radical, the diazo-nium salt being added to a warm solution of potassium cuprocyanide The Gattermann process is also applicable to this synthesis, the addition of copper powder to a mixture of diazonium salt and potassium cyanide inducing a similar replacement (Ber 1890, 23, 1218) The cyanate radical is introduced in the place of the diazo- group by adding successively to a solution of diazonium salt, potassium, cyanate, and copper powder (Ber 1890, 23, 1220, 1892, 25, 1086). The thiocyanate group is introduced by adding potassium thiocyanate and cuprous throcyanate to a solution of diazonium sulphate (1814 1890, 23, 738, 770) , 5 Replacement of the Diazo- Group by Sul-

phur and Sulphur-containing Radicals.

These reactions are of importance as furmshing methods for the synthetical production of sulphonic acids

Throphenols or disulphides are obtained by treating diazonium salts with potassium xanthate and then hydrolysing the resulting aromatic vanthate (Leuckart, J. pr Chem. 1890, [n | 41, 179], cf. Ber 1887, 20, 349, Chem Soc Proc 1895, 11, 141). Phenyl sulphide is produced by treating benzenediazonium salts with hydrogen sulphide or ammonium sulphide (Ber. 1882, 15, Benzenoid diazonium salts yield sul-1683)phides on treatment with sodium cupious thiosulphate, but diazotised α-naphthylamine only gives a-azonaphthalene (Bei 1901, 34, 3968)

The replacement of a diazonium by a sulphinic group is effected by saturating with sulphur dioxide an acid solution of diazonium sulphate and adding copper powder, introgen is evolved and the resulting sulphinic acid extracted with other (Gattermann, Ber 1899, 32, 1136, Eng Pat 26139 of 1896, D R P 95830)

The foregoing thiophenols, disulphides and sulphinic acids yield the corresponding sulphonic acids on oxidation (D. R. P. 70286). Eng. Pat. 11865 of 1892)

6 Replacement of the Diazo- Group by Nitro-, Nitroso-, and Amino- Radicals.

The introduction of a nitro- group is brought about either by heating the diazonium salt with nitious acid and cuprous oxide or by mixing the diazonium sulphate with cupro-cupric sulphite and adding excess of alkali nitrite latter process, 2.4.6-tribromobenzenediazonium sulphate gives 65 p c of 1-nitro 2.4 6-tribromobenzene and diazotised β-naphthylamine gives 25 p.c. of \(\beta\)-nitronaphthalene, a substance prepared with considerable difficulty by other processes (Ber 1887, 20, 1497, 1900, 33, 2544). The introduction of a nitroso- group is effected by treating a diazonium chloride with alkaline potassium ferrocyanide (Ber. 1893, 26, 471) The addition of hydroxylamine to a solution of diazonium salt results in the regeneration of the original amine (Ber 1892, 25, 372; 1902, 35, 2593, 3920)

#### 7 Formation of Azoimides.

The introduction of the azimino- group  $N_3$ into the aromatic nucleus is effected by the following reactions: (1) action of hydroxylamino on diazonium sulphates (Bei 1892, 25, 372, 1893, 26, 1271, Forster and Fierz, Chem Soc Trans 1907, 91, 855, 1350); (2) interaction of ammonia and diazonium perbromides; (3) direct action of sodium azide on diazonium salts in excess of sulphuric acid (Nolting and Michael, Ber 1893, 26, 86, Forster and Fierz, Chem. Soc Trans 1907, 91, 1942). The last of these processes is carried out so readily that the production of an azimino- compound may be used to estimate the number of diazo- groups in a diazonium salt (Meldola and Hawkins, Chem Soc Proc 1892, 8, 133, Morgan and Micklethwait, Chem Soc. Trans. 1910, 97, 2560). Aromatic azoimides are ilso produced by the interaction of chloroamine, NH<sub>2</sub>Cl, and diazonium chlorides (Forster, Chem. Soc. Trans. 1915, 107, 260), and by removing water from the nitrosoarylhydroxylamines

These aryl azomides can be employed synthetically as a means of transforming an amine into the corresponding aminophenol when warmed with moderately concentrated sulphuric acid, two-thirds of the introduced into the nucleus (Ber. 1894, 27, 192; Morgan and Porter, Chem Soc Trans 1915, 107, 651).

Miscellancous substitutions, v Chem Soc Trans 1888, 53, 460; Amer. Chem J 1888, 10, 368, Bei 1902, 35, 1964.

For the replacement of the diazo- group by assinic and stibinic complexes consult Organic Arsenicals and Organic Antimonials.

#### CONDENSATIONS EFFECTED BY MEANS OF AROMATIC DIAZONIUM SALTS

Azo- compounds are formed, to a slight extent, during the Sandmeyer reaction, and this condensation takes place to a greater extent when the usual order of mixing is reversed and the cuprous chloride added to the cold solution of diazonium salt Aniline, o-chloraniline, and o- and p-toluidine give considerable quantities of azo- derivatives, but the nitianilines and their homologues furnish diphenyl compounds (Niementowski, Ber 1901, 34, 3325, Ullmann and Forgan, *ibid* 3802, D R P 126961) Cuprous oxide dissolved in ammonia or hydroxylamine also promotes the diphenyl condensation (Annalen, 1902, 320, 122), diphenyl itself being readily prepared by adding copper, zinc, or non powder to benzenediazonium sulphate in dilute alcohol (Ber 1890, 23, 1226), ββ-dinaphthyl is produced from B-diazonaphthalene similarly sulphate (Chem Soc. Trans 1895, 67, 653) 2 2'-Dimethyl-1 1'-dianthraquinonyl results from the condensation of 2 mols of 2-methylanthraquinone-1-diazonium sulphate when stirred with acetic anhydride and copper bronze powder (Scholl and Kunz, Ber 1907, 40, 1697, Eng. Pat 14578 of 1905)

# INTERCHANGE OF GROUPS IN DIAZONIUM SALTS.

The diazonium salts of the more highly substituted amines frequently undergo remarkable transformations, some of which have been suggested for use in the technical production of azo- compounds.

The dinitroanisidines, containing nitrogroups either in ortho- or para- positions with regard to the amino- group, exchange these radicals for hydroxyl during the process of diazotisation. In certain instances, the methoxyl group present in these compounds undergoes demethylation, &c, with the production of internal diazo- oxides (quinonediazides) The nitro- group is set free in the form of nitious acid, which continues the diazotisation process on further quantities of the base (Meldola and others, Chem Soc Trans 1900, 77, 1172, 1901, 79, 1076; 1902, 81, 988, 1905, 87, 1205) Methoxyl or chlorine radicals in para-positions with respect to the diazo-complex are rendered labile by contiguous nitro- groups so that paradiazo-oxides are produced (Klemenc, Ber. 1914, 47, 1407; Morgan and Porter, Chem Soc. Trans 1915, 107, 653), but an accumulation of acidic (nitro) groups favours the removal of adjacent nitro- groups and the formation of oitho-diazo oxides (Meldola and Reverdin, Chem Soc Trans, 1910, 97, 1204). A sulphonic group is replaced by hydroxyl in the diazotisation of m-phenylenediammedisulphonic acid (Eng Pat 18283 of 1903), and certain of the dimitronaphthylamines, when diazotised, yield internal diazo-oxides, owing to the initial replacement of a intro-group by hydroxyl (Bei 1894, 27, 2211, 1895, 28, 1951, of Chem Soc Trans 1895, 67, 908)

The replacement of an acidic (electronegative) substituent by hydroxyl frequently occurs on treating the diazonium salt with alkalis

$$\begin{array}{c} \cdot \\ \cdot \\ \text{CIN}_2 \\ \\ \cdot \\ \text{SO}_3 \\ \text{H} \end{array} \rightarrow \begin{array}{c} \text{OH} \\ \\ \cdot \\ \text{OH}_2 \\ \\ \cdot \\ \text{SO}_3 \\ \text{H} \end{array} \rightarrow \begin{array}{c} \text{O} \\ \\ \cdot \\ \cdot \\ \text{N}_2 \\ \\ \cdot \\ \text{SO}_3 \\ \text{H} \end{array}$$

A sulphonic acid group may be similarly eliminated as in  $\beta$ -naphthylamine-1 5 7-trisulphonic acid which, when diazotised and treated with weak alkali, yields naphthylene 2-diazo-1-oxide-5 7-disulphonic acid (D. R. P. 141750), Eng. Pats. 16811, 20551, of 1901, 1561, 6615, 23993, of 1902, 16995, 27372, of 1903, 4997, of 1904).

These replacements sometimes take place even when the diazo-solutions are diluted with water, or spontaneously when the diazonium base is present as the salt of a weak and (acctate, oxalate, carbonate, &c) (v. Orton, Proc. Roy. Soc. 1902, 71, 153., Chem. Soc. Trans. 1903, 83, 796., 1907, 91, 1554., Nolting and Battegay. Ber. 1906, 39, 79)

 $p\hbox{-}{\rm Chlorobenzene diazonium}\quad {\rm thioeyanato}$ 

undergoes a remarkable molecular transformation when dissolved in alcohol containing a trace of hydrochloric acid. The thiocyanate and chlorine radicals change places, and on adding other to the solution, p-thiocyanobenic ne diazonium chloride is precipitated (Hantzsch, Ber 1896, 29, 947) Similarly, 2 4 6-tribromobenzenediazonium sulphate in presence of excess of potassium thiocyanate is converted into 2.4.6-trithiocyanobenzenediazonium thiocyanate and many brominated diazonium chlorides pass into chlorimated diazonium bromides (Bei 1897, 30, 2334, 1898, 31, 1253, 1900, 33, 505, cf Chem Soc Trans 1902, 81, 1376) The last of these transformations has been studied quantitatively, and the following generalisations have been established -

- (1) Bromine atoms are replaced by chlorine only when present in the ortho- or para- position with respect to the diazonium group, those in ortho- positions being most readily removed. A bromine atom in the meta- position is not affected
- $(\pi)$  The ease of transformation increases with the number of bromine atoms present
- (iii) The transformation constant, calculated from the equation for a unimolecular reaction  $K=1/t(\log A/A-x)$ , increases with the temperature and is also influenced by the solvent, having its minimum value in water, and becoming greater as the series of alcohols is ascended.

(iv) Diazonium salts containing two bromine atoms are stable when dry, but are rapidly transformed in ethyl alcohol; 2 4.6-tribromobenzenediazonium chloride becomes transformed in the dry state

These changes do not occur either with iodinated diazonium chlorides or with brominated diazonium fluorides (Hantzsch, Ber. 1903, 36,

zuoe).

CONSTITUTION OF AROMATIC DIAZONIUM SALTS.

Aromatic diazonium salts are generally soluble in water to a neutral solution, thus showing, by an absence of hydrolytic dissociation, that the diazonium base is a stronger base than the aromatic amine from which it was derived. The strength of the diazonium base has been confirmed by physico-chemical measurements.

Determinations of the electrical conductivity of solutions of benzenediazonium chloride and mirate show that the benzenediazonium radical is strictly comparable with other quaternary ammonium ions. The rate of migration of the benzenediazonium ion at 25° is 457, the corresponding constants for the methylpyridinium and tetramethylammonium ions being 443 and 436 respectively. The molecular electrical conductivity of solutions of diazonium salts increases with dilution, just as in the case of the corresponding potassium and ammonium compounds.

The affinity constant of benzenediazonium hydroxide at 0° is 70 times greater than that of ammonium hydroxides, and somewhat exceeds that of piperidine. The affinity constants of p-methoxybenzenediazonium and \(\psi\)-cumidine-diazonium hydroxides are even greater, and approximate closely to those of the alkali hydroxides A comparison of the electrical conductivity experiments, with the results obtained in the hydrolysis of ethyl acetate by benzenediazonium hydroxide, shows that in N/128 solutions at 0°, approximately 33 p.c. of the base exists in the ionised condition (Ber. 1890, 23, 3220; 1895, 28, 1737, 1898, 31, 1612)

Benzenediazonium (Diazobenzene) behaves like an alkali or a complex ammonium cation

Benzenediazonium hydroxide (Diazobenzene hydrate or hydroxide) is only known in solution, and is produced by treating aqueous benzene-diazonium chloride with the calculated amount of silver oxide, or by adding barium hydroxide to a solution of benzenediazonium sulphate (Hantzsch, Ber 1898, 31, 340) The solution is strongly alkaline, and gives at once an azocompound with \$\beta\$-naphthol, the dissolved base decomposes, however, fairly rapidly even at 0°

The striking similarity in the physical and chemical properties of the diazonium and quaternary ammonium salts suggests the existence of quinquevalent introgen in the former compounds, and on these grounds the following formula, formerly proposed by Blomstrand (1869), Strecker (1871), and Erlenmeyer (1874), has now been generally adopted for benzenediazonium chloride and its homologues:

$$C_6H_5$$
— $N^V$ — $Cl$ 

although, for many years, the alternative configuration,  $C_6H_6$ : N=N CI, suggested by Kekulé, was usually employed.

The close analogy existing between the diazonium and alkali salts is shown further by the facility with which the former yield double salts with cobaltic nitrite and with the chlorides of platinum, gold, and mercury. Diazonium perhalides are also known, corresponding with the perhalides of potassium, rubidium, cæsium, and quaternary ammonium bases (Ber. 1895, 28, 1736, 2754).

The benzenoid diazonium salts with colourless anions (nitrates, sulphates, chlorides) are generally colourless, but the corresponding bromides, iodides, and thiocyanates are coloured, and the deeper the colour the more explosive the salt. When cooled in liquid air, these coloured salts become less coloured; moreover,

their aqueous solutions are colourless

The development of colour and increase in explosibility are considered by Hantzsch to indicate a change in constitution, the coloured substance being regarded as an equilibrium mixture of diazonium salt and syn-diazocompound (v infra). This equilibrium is disturbed in the sense of forming more colourless diazonium salt by lowering the temperature or by dissolving the substance in water (Hantzsch, Ber 1897, 33, 2179, 1900, 33, 2179, Euler, Ber. 1895, 31, 4168) But although in the benzeno series the diazonium chlorides, nitrates, and sulphates are colourless, yet in the diphenyl and naphthalene series, many diazonium salts with these amons are coloured and comparatively stable. Moreover, this colour (yellow or orange) is retained either in aqueous solution or at the temperature of liquid air. Apart from their colour, these salts have all the properties of the simpler diazonium salts (neutral solutions, &c). These facts coupling with phenols, indicate that increase in the complexity of the organic portion of the diazonium cation leads to the development of colour in normal diazonium

(Coloured diazonium salts, v. Hantzsch, Ber 1900, 33, 3183, 1901, 34, 4168, Annalen, 1900, 312, 126, Cain, Chem. Soc. Trans. 1905, 87, 5; Morgan and others, ibid. 1907, 91, 1311, 1505; 1908, 93, 614, 1909, 95, 1319)

The idea embodied in Blomstrand's diazonium formula (v. supra) fails to express one important point, namely, the dependence of the stability of the diazo-complex on the presence of an unsaturated group. Yet this is a matter of fundamental importance, for hitherto no amine in which the basic nitiogen is attached to a fully saturated ring or complex has yielded a diazonium salt.

Cain has suggested an alternative formula, (2), giving expression to this circumstance (Chem Soc Trans 1907, 91, 1051), in which the diazonium salt is represented as having a parahemiquinonoid constitution, and this conception has been extended by Morgan (*ibid.* 1908, 93, 617; 1910, 97, 2563) so as to include the equally admissible ortho-hemiquinonoid configurations (1) and (3):

$$\begin{array}{c|c} X & & H \\ H & & H \\ H & & H \\ \end{array} \xrightarrow[N]{H} H \end{array} \rightleftharpoons \begin{array}{c} X & & H \\ H & & H \\ H & & H \\ \end{array} \xrightarrow[N]{H} H$$

Euler has also advocated a para-hemiquinonoid structure for diazonium salts (Ber 1908, 41, 3979).

#### Non-aromatic Diazonium Salts.

The foregoing hypotheses of the constitution of aromatic diazonium salts, which bases their existence on the unsaturated nature of the aromatic complex, accord with the fact that certain non-aromatic primary amines containing unsaturated organic groups manifest, in some degree, the property of diazotisability. Among those bases are aminotriazole (1) and its derivatives, and aminoantipyrine (II):

(Thiele and Manchot, Annalen, 1898, 303, 33: Knorr and Stolz, *ibid*. 1896, 293, 67, *cf* Ber 1900, 33, 1158, 1906, 39, 2925, Annalen, 1900, 312, 133). In addition to amino derivatives of the pyrazoline group (Michaelis, Annalen, 1906, 350, 288), the 4- and 5-amino derivatives of the pyrazole group give rise to remarkably stable diazonium salts (Morgan and Reilly, Chem Soc. Trans 1914, 105, 436; Michaelis and others, Annalen, 1911, 385, 1; 1915, 407, 229, 274; Mohr, J. pr Chem. 1914, [11] 90, 509) 2 Aminothiazole, a heterocyclic base containing sulphur and nitrogen in the ring, furnishes diazonium salts of the oxyacids (Morgan and Morrow, Chem Soc Trans 1915, 107 1291). Distinct evidence of diazotisability has been adduced in the indole and pyrrole series (Angeli and d'Angelo, Atti. R. Accad Lincei, 1904, [v.] 13, 1. 258; Angelico, Ic 1905, 14, 11 167) It has also been shown (K. A. Hofmann, Hock and Roth, Ber 1910, 43, 682, 1087) that aminoguanidine gives rise to diazonium salts, derived, however, not from the salts of aminoguanidine itself, but from a more complex molecule containing two guanidine residues, and having a greater degree of unsaturation. The diazonium greater degree of unsaturation nitrate, for example, is represented by the formula ·

$$NH \underbrace{\begin{array}{c} NO_2 \cdot N_2 \ NH \ C(\ NH) \\ C(NH_2) \ NH - NH \ N \end{array}}_{} N$$

containing a diazo- complex associated with an open chain having three unsaturated linkings

#### II. CYCLIC DIAZO- COMPOUNDS.

In certain instances, the action of nitrous acid on substituted aromatic amines leads to the formation of compounds having their diazonitrogen in a cyclic complex, this result being due to the subsequent interaction of a substituent group.

#### 1. INTERNAL DIAZONIUM SALTS.

Sulphonated ammes of the benzene and naphthalene series furnish diazo- derivatives which are generally misnamed diazo-sulphonic acids. These compounds, however, do not contain ionic hydrogen, and are really internal diazonium salts; No 1, the so-called 'diazobenzenesulphonic acid,' being benzenediazonium-p-sulphonate, whilst No. 2 is naphthalene-1-diazonium-5-sulphonate

These internal diazonium sulphonates are of great technical importance in the manufacture of azo-dyes (r. Azo-colouring matters). They are prepared either by adding sodium nitrite to an acid suspension of the amino sulphonic acid or by acidifying a solution containing nitrite and the alkali salt of the aminosulphonic acid (Rev. prod. chim. 1917, 20, 21).

Of a similar nature are the cyclic diazo-

Of a similar nature are the cyclic diazoderivatives obtained by diazotising aromatic aminocarboxylic acids (cf. Bulow and Haas, Ber, 1911, 44, 601), e.g. 'diazo-anthranilic acid' or benzeneduzonium-o-carboxylate

#### 2. Cyclic Dia7o- Compounds due to Internal Condensation

Aromatic ortho- and pen-diamines do not yield diazonium salts, masmuch as their interaction with introus acid leads at once to the formation of cyclic diazonimes (Arimino-compounds)

Phenylene-o-diazoimine (1) (Aziminobenzene) and naphthylene-1 8-diazoimine (2) (1 8-Azimino-naphthalene) are typical members of this series of diazo-compounds

The triazole rings thus set up are remarkably stable, and in the case of the ortho-derivatives (No 1) are not opened by mineral acids without complete rupture of the molecule. The peridiazonimes (No 2) occupy a position intermediate between the ortho-diazonimes (No. 1) and the para-diazonimes (v infra)

The 12- and 23-naphthylene-diamines furnish respectively 12- and 23-naphthylene-diazomines (12- and 23-aziminonaphthalenes). (Constitution of ortho-diazomines and their acyl- derivatives, Griess, Ber. 1882, 15, 2195; Kekulé, Lehrbuch, 11739, Morgan and Godden, Chem. Soc Trans 1910, 97, 2557; cf Ber 1874, 7, 316; 1876, 9, 221, 1887, 20, 2999, 1894, 27, 874; Annalen, 1889, 255, 344. Chem. Soc Trans 1906, 89, 4) Although Griess's formulation for the azimino-compounds

$$C_6H_4$$
 $N$ 
 $NH$  still persists in chemical litera-

ture, it has been disproved completely, and that of Kekulé substantiated by the preparation of three pairs of isomerides having respectively the following formulæ, where R=acetyl, benzovl, or benzenesulphonyl

This form of isomerism could not exist with Guess's symmetrical configuration (Morgan, Micklethwait and Schaiff, Chem Soc Trans 1913, 103, 1394, 1914, 105, 117). Although these diazoimines do not couple with phenols and aromatic bases to form azo-dyes, yet a synthetic employment has been found for them in the production of carbazole derivatives which are obtained on heating these cyclic dia/o-compounds (Ullmann, Ber 1898, 31, 1697, 1914, 47, 380).

Similar stable cyclic diazo- compounds are obtained by diazotising the ortho-aminothiophonols, o-phenylenediazosulphide being a colourless crystalline substance with a sweetish odour and feebly basic properties

$$C_0H_4{<}_N^S{>}N$$

 $^{\text{Cl}_{0}}\text{H}_{4}{<}_{N}^{\text{S}}{>}N$  (Jacobsen, Annalen, 1893, 277, 209, 218, 232, 237)

3 CYCLIC DIAZO-COMPOUNDS DUE TO COMBINED INTERNAL CONDENSATION AND SALT FORMA-

In this group are found certain diazoderivatives of amines containing slightly acidic substituents in ortho- or para-, but not metapositions with respect to the aminic nitrogen

(1) Internal diazo- oxides are produced from ortho- and para- aminophenois and their derivatives În some cases, an intermediate diazonium salt is formed which condenses to the diazo-oxide on treatment with silver oxide or an alkalı (Schmitt, Ber 1868, 1, 67, Hantzsch and Davidson, *ibid* 1896, 29, 1522, Cameron, Amer Chem J 1898, 20, 229) The first diazocompound prepared by Griess (Annalen, 1858, 106, 123), namely, dinitrobenzenediazo-oxide

$$(NO_2)_2C_6H_2$$
  $\bigcup_{N_2}^{O}$  belongs to this class; it is

produced by diazotising pieramic acid, and has been utilised in the production of azo-colours (D R. P 151332). The 12-, 2.1-, 23-, and 1.8 aminonaphthols and their sulphonic acids similarly give rise to internal diazo-oxides which are so stable that they can be nitrated. These internal diazo-oxides are readily transformed into diazonuim salts by mineral acids, and are therefore available in the production of azo-dyes (Eng Pat 28107, of 1897; 10235, 15025, of 1904; Fr Pat 353786 of 1905; D R. PP 155083, 164665, 171024, 172446, 175593, 176618, 176619, 176620).

Internal diazo-oxides can also be produced by leaving the diazonium salts of highly substituted amines (eq 246-trichloroaniline) in aqueous solutions containing no free mineral acid (Orton, Proc Roy Soc 1903, 71, 153; Chem. Soc Trans. 1903, 83, 796)

Thes, diazo- derivatives have been formulated either as cyclic diazo-oxides (1) or as quinonediazides (II.):

$$\begin{array}{ccc}
O \cdot C_6 H_4 \cdot N_2 & O \cdot C_6 H_4 < N \\
I & II.
\end{array}$$

(cf. Wolff, Annalen, 1900, 312, 126, Morgan and Micklethwait, Chem. Soc. Trans 1908, 93, 607)

Repeated attempts to prepare internal diazooxides from m-aminophenol and its nitroderivatives, which have been quite unsuccessful, strengthen the view that these diazo-derivatives are to be regarded as quinonediazooxides (II), with the reservation that similar compounds are producible from 18-aminonaphthol, 'H acid,' and other peri-derivatives (Morgan and Porter, lc). In the case of aminophenolsulphonic acids the NH: OH=1.2 compounds give these coloured o-quinonediazides, the NH<sub>2</sub>OH=14 compounds yield diazonium sulphonates changing to p quinonediazides on partial neutralisation, the NH<sub>3</sub>:OH=13 compounds give rise only to diazonium sulphonates which lose diazo-nitrogen and furnish complex azo derivatives on treatment with alkali (Morgan and Tomlins, Chem. Soc Trans. 1917, 111, 497).

(11) Para diazoimines and diazoimides.

The simplest example of the para-diazomines is obtained by the action of alkalis on the diazonium salts of p-aminodiphenylamine (Annalen, 1888, 243, 282; Ber 1902, 35, 895) The following alternative formulæ (III and IV) have been given to this product -

corresponding respectively with the foregoing formula (1 and II) for the internal diazo-oxides. Successive nitration of the phenyl group gradually increases the stability of these pdiazoimines (Chem Soc Trans 1908, 93, 604).

The para-diazomides are produced from the acyl- and arylsulphonyl- derivatives of the para-diamines, and similar products are obtained the arylsulphonyl-I · 8-naphthylenediamines. The production of these diazoimides takes place in two stages, the diazonium salt first produced being decomposed by sodium acetate or mild alkali, yielding the internal condensation product

These para-diazoimides (VI.) are yellow sparingly soluble substances readily reconverted by strong acids into the corresponding diazonium They combine additively with phenols, naphthols, naphthylamines, and their sulphonic naphonous, naphonylamines, and their sulphonic acids, yielding azo-colouring matters (Morgan and Micklethwait, J. Soc. Dyers and Colourists, 1909, 25, 107, cf. Chem. Soc. Trans. 1905, 87, 74, 921, 1302, 1906, 89, 1162; 1907, 91, 1509, 1908, 93, 615. Badische Anilin- und Soda-Fabrik D. R. P. 205037)

The foregoing general reaction has been extended further by the device of diazotising with liquid nitrogen trioxide in an anhydrous solvent such as acetone. In this way the simplest members of the series Ac N.C. H.: N2 have been obtained containing formyl, acetyl, and benzoyl groups (Morgan, Upton and Cleaze, Chem. Soc. Trans. 1917, 111, 187; 1918,

#### III. DIAZOAMINES.

The diazoamines or diazoamino- compounds have the general formula XN<sub>2</sub> NHY, and may be either aromatic, mixed aromatic-aliphatic, or aliphatic in character, according as to whether X and Y are aromatic or aliphatic hydrocarbon radicals.

#### 1 AROMATIC DIAZOAMINES.

The aromatic diazonmines are produced (1) by the action of nitrous acid (1 molecule) on two molecular proportions of a primary benzenoid amine, (2) by coupling a diazonium salt with a primary or secondary benzenoid amine.

Diazoaminobenzene C<sub>6</sub>H<sub>5</sub>·N<sub>2</sub> NH C<sub>6</sub>H<sub>5</sub>, yellow crystals, mp 98°, is produced in quantitative yield by dissolving 10 parts of aniline in two parts of water containing 12 parts of hydrogen chloride, diazotising with 8 parts of sodium nitrite, and adding the diazo-solution to a solution of 10 parts of aniline in the calculated amount of hydrochloric acid. The diazoamine is precipitated on the addition of sodium acetate. (Absorption spectrum of diazoaminobenzene, Purvis, Chem. Soc. Trans. 1914, 105, 590.)

When warmed with aniline containing aniline hydrochloride, diazoaminobenzene is transformed into p-aminoazobenzene (v Azocolouring matters), the velocity of transformation obeying the law of unimolecular reactions. When the transformation occurs below 40° a small proportion (about 4 pc) of o-aminoazobenzene is produced (F H Witt, Ber 1912, 45, 2380, 1913, 46, 2557)

Diazoamino p-toluene C<sub>7</sub>H<sub>7</sub> N<sub>2</sub> NH C<sub>7</sub>H<sub>7</sub> is prepared in a similar manner, and undergoes transformation into an ortho-azo-compound, but the velocity of this change is only about one-ninth of that of the preceding transformation to para-azo-derivative

The coupling of diazonium salts with metaphenylenediamine, diphenylamine, the naphthylamines and their alkyl derivatives, leads directly to the production of azo-derivatives, in these cases, intermediate diazoamines have not been isolated, except when the reactive ortho- or para-positions are already substituted (Morgan, Chem Soc Trans, 1902, 81, 91; 1907, 91, 370)

Chem Soc Trans. 1902, 81, 91; 1907, 91, 370)

When an aromatic base, XNH<sub>2</sub>, is coupled with a diazonium salt YN<sub>2</sub>Cl, the resulting diazoamine is identical with that produced from XN<sub>2</sub>Cl and YNH<sub>2</sub>, and is generally regarded as being an equilibrium mixture of the two isomerides XNH·N<sub>2</sub>Y and XN<sub>2</sub>·NHY This explanation assumes the migration of the diazogroup which has been observed to occur in several instances (Ber 1882, 15, 2190; 1896, 29, 287; 1897, 30, 1412).

(For the constitution of the unsymmetrical aromatic diazoamines, XNH·N<sub>2</sub>Y, v McIdola and Streatfelld, Chem Soc Trans 1886, 49, 624; 1887, 57, 102, 434; 1886, 53, 664; 1889, 55, 412; 1890, 57, 785; cf. Forster and Garland will 1909, 95. 2051)

## 2. ALIPHATIC-AROMATIC DIAZOAMINES.

The diazoamines containing one aliphatic and one aromatic group can be produced by (1) coupling a diazonium salt with an aliphatic amine (Ber. 1888, 21, 1016, 1112; 1889, 22, 938,

1302); (2) by treating an azide with Grignard reagent:

R N
$$\stackrel{N}{\downarrow}$$
 +R' MgBr=R N(MgBr) N NR'  
H<sub>2</sub>O  $\Rightarrow$  R NH N NR'+MgBr OH

(Dimroth, Ber 1903, 36, 909; 1905, 38, 670, 2328; 1907, 40, 2390). Diazoamines present the possibility of stereoisomerism, but their relative stability and the fact that they couple only slowly with  $\beta$ -naphthol, confirm the view that they are generally anti-diazo-compounds.

that they are generally anti-diazo- compounds. (For stereoisomeric diazoamines, v J Russ Phys. Chem Soc 1906, 38, 578; Zeitsch angew Chem 1900, 13, 762; 1902, 15, 1209)

Aromatic bisdiazoamino- compounds, e.g. (C<sub>6</sub>H<sub>5</sub> N.), N. C<sub>6</sub>H<sub>5</sub>, have been obtained by the action of 2 molecular proportions of a diazonium salt on one of an aromatic amine (Ber. 1894, 27, 1863, 2597, 1895, 28, 170)

Mixed aliphatic-aromatic bisdiazoamino- derivatives (v. Ber. 1888, 21, 1016, 1889, 22, 933; 1907, 40, 2390)

Diazo- $\psi$ -semicarbazines (v. Forster, Chem. Soc. Trans. 1906, 89, 223)

#### 3. ALIPHATIC DIAZOAMINES.

Dimroth's general method may be applied to the production of purely aliphatic diazo-amino-compounds. In this way, the simplest member of the series, diazoaminomethane (dimethyltriazene) CH<sub>3</sub>N·N NH CH<sub>3</sub>, has been prepared from methyl azide and magnesium methyl iodide. This substance, which is isolated by distilling its copper derivative with diazoaminobenzene under reduced pressure, is a colourless liquid, solidifying at --12° and boiling at 92° (Dimroth, Ber. 1905, 38, 1575, 1906, 39, 3905)

#### IV DIAZO- OXIDES

When a diazonium salt couples with a phenol the product is generally an ortho or a parahydroxyazo- compound (the C-azo- derivative), but in all probability, the O-azo- derivative is first produced, for in a few instances, when the velocity of transformation has been lessened by substitution, the intermediate O-azo- compound or diazo-oxide has been isolated p-Bromobenzenediazonium chloride and p-introphenol give p-bromobenzenediazo-4 oxynitobenzene

#### CaHaBr N N O CaHa NO,

which, at 80°, becomes transformed into its isomeride, p-bromobenzene-2-azo-4-nitrophenol

C<sub>6</sub>H<sub>4</sub>BrN N C<sub>6</sub>H<sub>4</sub>(NO<sub>2</sub>)OH

(Ber 1908, 41, 4027, 4304)

Occasionally traces of these intermediate diazonmines and diazo-oxides can be detected in the commercial azo-colouring matters (Vaubel, Zeitsch Farben Textil Ind 1902, 1, 3).

# V. METALLIC DERIVATIVES OF DIAZO-COMPOUNDS.

In 1894 Schraube and Schmidt (Ber 27, 514) found that on adding a 10 p.c. solution of p-nitrober zenediazonium chloride to 18 pc aqueous sodium hydroxide at 50°-60°, a sodium diazo-derivative separated in yellow bronzy leafiets, to which they gave the formula

NO2 C6H4 NNa NO, H2O

regarding the compound as the sodium derivative of p-nitrophenylnitrosamine. This view of the constitution of the product was confirmed by the action on the compound of methyl iodide when p-nitrophenylmethylnitrosamine.

was obtained

This sodium salt, which was regarded by Hantzsch as having the constitution of an antidiazo-compound, sodium p-nitrobenzenediazo-nxide (v infra), has been manufactured for use in the production of 'introsamine red' (Badische Anilin- und Soda-Fabrik D. R. PP. 78874, 80263, 81134, 81202). The addition of excess of mineral acid to this metallic diazo-derivative, determines the regeneration of the original diazonium salt, but treatment with dilute acetic acid leads to the formation of a yellow product, p-nitrophenylnitrosamine. NO<sub>2</sub> C<sub>6</sub>H<sub>4</sub> NH NO, a substance showing little tendency to couple with β-naphthol

The silver derivative obtained from the foregoing sodium salt yields on methylation an oxygen ether NO<sub>2</sub> C<sub>6</sub>H<sub>4</sub> N N O CH<sub>3</sub>, a reaction suggesting the possibility of tautomeric change in many instances, two isomeric modifications of these metallic diazo-deit-

vatives have been isolated

Benzenediazonium chloride and cold concentrated potassium hydroxide yield the normal labile form of potassium benzenediazo-oxide (I), which readily couples with β-naphthol. The stable modification of potassium benzenediazo-oxide (II) is produced by heating the strongly alkaline solution of the normal or labile salt at 130°–140°; this isomeride couples only very slowly with β-naphthol

According to Hantzsch, these isomerides have respectively the following formula —

$$\begin{array}{cccc} C_8H_8 & N & C_8H_8 & N \\ & \parallel & & \parallel \\ & KO & N & N & OK \\ I & \textit{Syn-} & (labile form) & II & \textit{Anti-} & (stable form) \end{array}$$

Dobbie and Tinkler (Chem. Soc. Trans. 1905, 87, 273) have found, however, that these isomeric potassium salts have totally different ultra-violet absorption spectra But as stereo-isomerides, like the benzaldoximes, have identical absorption spectra, it seems probable that the above isomeric salts are structurally dissimilar. Confirmation of the stereochemical theory would be obtained by isolating the antidiazohydroxide from the stable (anti-) potassium salt, but on treating this with an acid, the nitro-Moreover, the product supsamine separates posed by Hantzsch and Pohl to be an antidiazohydroxide (Ber. 1902, 35, 2964) was shown by Orton to be a mixture of quinonediazide and a hydroxyazo- compound (Proc Roy Soc 1903, 71, 153; Chem Soc Trans 1903, 83, 796)

The existence of the syn-diazohydroxides is doubtful, and diazonium hydroxides (v. supra) are only known in solution.

Diazo-anhydrides RNNON·NR or R·N·ON.NR are very unstable explosive com-N

pounds produced by the action of acetic acid on the metallic sym-diazo-oxides. The anti-diazooxides, when thus treated, give nitro samines.

Aromatic diazoic acids Ar N<sub>2</sub>O<sub>2</sub>H. Benzene-

diazoic acid, produced by oxidising potassium benzenesyndiazo-oxide or -isodiazo-oxide with potassium ferricyanide in alkaline solution, is obtained in sparingly soluble white leaflets (mp 46°) dissolving readily in organic solvents or alkalis. With mineral acids it is transformed into o- and p-introamline, and with hypochlorites it yields 4-chloro-2-introamline lits sodium salt gives the  $\alpha$ - or N-ester C<sub>6</sub>H<sub>5</sub>, N(CH<sub>2</sub>) NO<sub>2</sub>, the silver salt furnishes the  $\beta$ - or O-ester, C<sub>6</sub>H<sub>5</sub> N NOO CH<sub>3</sub>. Since the acid is also obtained by the action of introgen pentoxide on aniline it is probably a tautomeric substance C<sub>6</sub>H<sub>5</sub> N N NO<sub>2</sub>=C<sub>6</sub>H<sub>5</sub> N NO<sub>2</sub>H (Bamberger, Ber 1893, 26, 471; 1894, 27, 359, 914, 584, 1273, Hantzsch, Ber. 1902, 35, 258)

#### VI DIAZO- CYANIDES.

The addition of the calculated amount of potassium cyanide to a cold acid solution of a diazonium salt determines the formation of a sparingly soluble diazo-cyanide. In many instances, two modifications of the diazocyanides can be distinguished, and these varieties are regarded by Hantzsch as being stereoisomeric forms p-Chlorobenzenediazonium chloride, from p-chloraniline, yields first the yellow labile syn-p-chlorobenzenediazoeyanide (1.), which readily evolves nitrogen, yields

$$\begin{array}{cccc} C_6H_4CIN & C_6H_4CIN \\ \parallel & & \parallel \\ CN\cdot N & & N\cdot CN \end{array}$$

I. Syn-diazocyanide II. Anti-diazocyanide p-chlorobenzontrile on treatment with copper powder, and passes into the stable anti-p-chlorobenzenediazocyanide (II) The latter substance, which is not affected by copper powder, may be distilled in steam without decomposition.

Confirmation of this stereochemical theory of the constitution of the diazo-cyanides has been obtained by a study of the cyanides derived from p-methoxybenzenediazonium chloride. This salt with potassium cyanide in alcoholic solution yields the syn-diazocyanide (III), an orange-red insoluble substance (mp. 51°), which couples with \(\beta\)-naphthol and slowly changes into the non-coupling anti-diazocyanide (IV), a brownish - red compound, melting at 121°.

The existence of a third isomeric cyanide is indicated by evaporating at the ordinary temperature in the presence of excess of hydrocyanic acid, an aqueous solution of p-methoxy-benzenediazonium hydroxide. The colourless crystalline product has the composition:

CH<sub>3</sub>O C<sub>6</sub>H<sub>4</sub>·N<sub>2</sub>·CN,HCN,2H<sub>2</sub>O

and possesses all the properties of a true metallic salt; it is very soluble, and its solution is an electrolyte. Moreover, this double salt couples with  $\beta$ -naphthol, and is converted by alkalis into the yellow syn-diazo cyanide. These properties correspond with those of the normal diazonium salts, and the foregoing soluble cyanide is regarded as having the following constitution:

N (Hantzsch, Ber. 1900, 33, 2161; 1904, 34, 4166).

The two pairs of diazo- cyanides from pchlorandine and p-anisidine were examined spectroscopically by Dobbie and Tinkler, who found that each pair gave almost identical ultra-violet absorption spectra, whereas the above soluble diazonium cyanide gave an entirely different spectrum These results are in accordance with Hantzsch's view of the configuration of the diazo- and diazonium cyanides (Chem Soc. Trans. 1905, 87, 273)

It has, however, been suggested by several workers in this field that the isomerism of the diazo- cyanides is structural, the syn- compounds being iso-cyanides R N N NC and their antiisomerides, cyanides R N N CN (Ber 1895, 28

861; Chem Soc. Trans 1903, 83, 805)

#### VII DIAZO- SULPHONATES.

Although the stereochemical theory of the constitution of diazo- cyanides is supported by physical as well as chemical evidence, yet it is significant that the only other series of salts in which this isomerism has been detected is one derived from sulphurous acid, a substance resembling hydrocyanic acid in giving rise to organic isomeric derivatives which are structurally dissimilar

The diazo-sulphonates, prepared by adding potassium sulphite to aqueous solutions of benzenoid diazonium chlorides, frequently exist in two differently coloured modifications, but in most cases the syn- isomeride is too unstable to be isolated in a pure state

Potassium syn-2 4-diiodobenzenediazosulphonate (I.) is an orange substance, whilst the antisalt (II) is yellow:

The diazotised naphthylamines behave exceptionally, yielding only syn-diazosulphonates, which, on warming pass, not into their antiisomerides, but into the corresponding azonaphthalenes (Ber 1897, 30, 71).

The syn- and anti- modifications of potassium benzenediazosulphonate  $C_0H_5$  N N  $SO_3K$ , were found by Dobbie and Tinkler (lc) to have identical ultra-violet absorption spectia fact supports the stereochemical theory of their structures (cf. Ber 1894, 27, 1726, 2099, 2586, 2930, 3527; 1895, 28, 242, 447, 834, 861, J pt. Chem 1894, [n] 50, 239; Meyer and Jacobsen, Lehrbuch der Org Chem 11 303)

# VIII. ALIPHATIC DIAZO- COMPOUNDS.

The amines of the aliphatic series, when treated with nitrous acid, generally lose their basic nitrogen, the amino- group becoming replaced by hydroxyl In some cases, however, the nitrogen is retained and an aliphatic diazocompound is produced, having the general

formula RH C  $\stackrel{\cdot}{\underset{N}{\bigcup}}$  with the azo- group attached

entirely to the same carbon atom

An alternative formulation has been suggested for the aliphatic diazo- compounds in the general formula becomes >('N' N (Thicle, Ber. 1911, 44, 2522; Forster and Cardwell, Chem Soc. Trans. 1913, 103, 861; cf Ber 1912, 45, 1654; 1916, 49, 1884

Although glycine is converted by nitious acid or alkyl nitrites into glycollic acid, yet its ethyl ester gives 1180 to ethyl diazoacetate  $N_2$  CH CO<sub>2</sub> C<sub>2</sub>H<sub>5</sub>, a yellow liquid freezing at 24 and boiling at  $143^{\circ}-144^{\circ}/72$  mm (Curtius,

Ber 1883, 16, 2230)

One kilogram of ethyl aminoacetate hydrochloride and 750 grams of sodium nitrite are added successively to 2 lities of water containing 5 grams of sodium acetate. The mixture is shaken until the temperature falls to 0°, 5 cc. of 10 p c sulphuric acid and 500 c c of ether are added and the liquids thoroughly agitated The ethereal layer is separated and the treatment with dilute acid and ether repeated until red fumes are evolved The ethereal extracts are washed with dilute aqueous sodium carbonate till alkaline, dried with calcium chloride, and the solvent removed on the water-bath. The yield of ethyl diazoacetate is 94 7 p c of the calculated quantity (Silberrad, Chem Soc Trans 1902, 81, 600)

With iodine, ethyl diazoacetate yields duodoacetic acid, and with aldehydes it condenses to form ketonic esters

$$\begin{array}{c} {\rm R\cdot CHO + N_2\ CH\ CO_2\ C_2H_5} \\ \longrightarrow \\ {\rm R\ CO\ CH_2\ CO_2\ C_2H_5} \end{array}$$

One of the most remarkable reactions of this diazo- ester is its condensation with benzene and its homologues, giving rise to a series of esters containing seven-membered hydrocarbon rings (Bucherer, Ber 1896-1903, 29, 106, 30, 632, 1949, 31, 399, 402, 2004, 2241, 2247, 32, 705, 33, 184, 36, 3509; Annalen, 1908, 358, 1)

Ethyl diazoacetate, when heated with alkalis undergoes a series of complex polymetic changes (Curtius, Ber. 1885, 18, 1283, 1906, 39, 1383, 3398, 4140, 1907, 40, 84, 815, 1176, 1194, 1470, 1908, 41, 3116, 3140, 3161, J pr Chem. 1888, [2] 38, 408, Hantzsch, Ber 1900, 33, 58, cf Chem Soc Trans 1902, 81, 598)

Diazo- derivatives are not obtained from free a-amino-carboxylic acids, and only from those amino-esters of aliphatic acids containing the amino- group in the a-position with respect to alkylated carboxyl group. The esters of those polypeptides which contain the aminogroup in the α-position to a carboxyl- group can also be transformed into diazo- derivatives.

$$\begin{array}{c} \mathrm{NH_2\,CH_2\,CO\,NH\,CH_2\,CO_2Et} \\ \quad \boldsymbol{\to} \quad \mathrm{N_2\,CH\,CO\,NH\,CH_2\,CO_2Et} \end{array}$$

The  $\alpha$ -aminoketones (e.g.  $\alpha$ -aminoacetophenone  $C_6H_5$  CO  $CH_2$   $NH_2$ ) give diazo- compounds (Ber 1904, 37, 2080), and so also do certain urie acid compounds, eg ammomethyluracil.

Diazomethane 
$$CH_2 \stackrel{N}{\swarrow}_N$$
, or  $CH_2 \cdot N : N$ , the

simplest aliphatic diazo- compound, is prepared by adding methyl-alcoholic potassium hydroxide to an ethereal solution of nitrosomethylurethane

and distilling the mixture on the water-bath, when the distillate consists of an ethereal soluwhich one nitrogen is quinquevalent, so that tion of the diazo- compound, the yield being 50 p.c. of the calculated quantity. This substance is also produced by the action of hydroxylamine on dichloromethylamine

 $CH_3 NCl_2 + H_2NOH = CH_2 N_2 + H_2O + 2HCl$ 

Diazomethane is a poisonous yellow gas condensing in snow and calcium chloride to a yellow liquid boiling below 0° It is a powerful methylating agent, converting hydroxyl groups into methoxyl, and alkylating primary and secondary bases. With iodine, diazomethane gives nitiogen and methylene iodide, and aldehydes are converted into methyl ketones (cf Monatsh 1905, 26, 1295, 1311, Ber 1907, 40, 479, 1908, 41, 3199, 1909, 42, 2559; Chem. Soc. Trans. 1908, 93, 242).

Diazomethane combines additively with unsaturated compounds; with acetylene it gives pyrazole (Ber 1898, 31, 2910), and with ethyl fumaiate it furnishes ethyl pyrazolinecar-

boxylate

Diazoethane CH3 CH . N2 and phenyldiazomethane C6H8 CH N2 have both been prepared (Ber. 1902, 35, 897) The former closely resembles diazomethane, the latter is a dark-red

Substitution increases considerably colour of these diazo- derivatives dimethyldiazomethane  $C(CH_3)_2$   $N_2$  is red and diphenyl-diazomethane is obtained in bluish red needles melting at 20' (Staudinger, Ber 1916, 49, 1884)

Diazomethane should not be confused with azoniethane CH<sub>3</sub> N H CH<sub>3</sub>, the simplest azocompound which is produced by oxidising symmetrical dimethyllightazine (hydrazomethane) with chromic acid (Thiele, Bei 1909, 42, 2575) Azomethane is a colourless gas condensing to a pale yellow liquid (b p 15°).

Diazomethanedisulphonic acid, noteworthy example of an aliphatic diazo-compound, results from the interaction of potassium cyanide and potassium bisulphite in presence of caustic potash. The solution, acidified and treated with nitrous acid, yields successively aminomethanedisulphonic acid and the diazo- compound Sulphurous acid and diazomethanedisulphonic acid yield an additive compound which, on boiling, furnishes hydrazine (V. Pechmann, Ber · 1895, 28, 2374, 1896, 29, 2161)

Diazoacetone CH<sub>3</sub> COCH N<sub>2</sub>, a pale yellow liquid, is obtained from aminoacetylacetone by converting this base into diazoacetylacetone anhydride and treating this product with aqueous caustic alkalı (Wolff, Annalen, 1912,

394, 23)

Metallic diazo- derivatives of the aliphatic series are produced by treating nitrosoalkyl urethanes with concentrated caustic potash solution or ethereal potassium ethoxide Potassium methyldiazo-oxide CH3 N B OK, H2O separates in white crystals, when nitrosomethylurethane is added to concentrated aqueous caustic potash at 0°. Potassium benzyldiazo-oxide

C<sub>6</sub>H<sub>5</sub> CH<sub>2</sub> N N OK,H<sub>2</sub>O

is similarly prepared from nitrosobenzyluiethane These metallic derivatives are very unstable, they are decomposed by water with explosive violence, yielding respectively diazomethane and phenyldiazomethane (Hantzsch and Lehmann, Ber 1902, 35, 897)

Bibliography - Hantzsch, Die Diazoverlindungen, Ahren's Sammlung, 1902, Morgan,

Our Present Knowledge of Aromatic Diazo-Compounds, Brit Assoc. Report, 1902; Eibner, Zur Geschichte der Aromatischen Diazoverbindungen, 1903, Cain, The Chemistry and Technology of the Diazo- Compounds, 1920

GT. M DIAZOACETATE ETHYL vDIAZO- COM-POUNDS

DIAZOAMINOMETHANE vD1A70- COM-POUNDS

DIAZOANTHRANILIC ACID Diazo-COMPOUNDS

DIAZOBENZENE, Renzenedrazonium DIAZO- COMPOUNDS

DIAZOBENZENE SULPHONIC ACID

DIAZO- COMPOUNDS DIAZOIMIDES, DIAZOIMINES v DIAZO-COMPOUNDS

DIAZOMETHANE v Diazo- compounds

**DIAZURINE** r Azo- colouring matters. DIBASIC ACIDS. Oxalic acid series numerous and important group of acids, having the general formula  $(nH_{2n-2}O_4)$  The lowest member of the series is oxalic acid C2H2O4, and the highest member at present known, dicetyladipu acid C38H74O4 As the radical CnH2n is capable of existing in a variety of ways, it follows that these acids may exist in many isomeric forms. Unlike the fatty acids, the lowest members of the oxalic acid series are solids

The naturally occurring acids exist partly free and partly as salts or esters. Thus oxalic acid occurs chiefly as the potassium salt in different varieties of Oxalis and Rumex, malonic acid, as the calcium salt, is found in the sugar beet, succinic acid in amber, certain varieties of lignite, resins, turpentine oils, and animal fluids, rocellie acid in Rocella tinctoria (D. C.) Other members of this series are obtained by the oxidation of naturally occurring substances suberic acid is obtained by boiling cork with mitie acid, azelaic acid by the oxidation of castor oil, adipic acid by the oxidation of suct or. tallow, whilst homologues of succinic, glutaric and adipic acids are frequently obtained in the oxidation of derivatives of the terpene compounds. The higher members of the series are prepared by the oxidation of members of the fatty acid series

General methods of preparation

(1) By the oxidation of the diprimary glycols, primary hydroxyaldchydes, dialdehydes, primary hydroxy- acids, or aldehydeeg glycol, glycollic acid, glyoxal, or glyoxylic acid -> oxalic acid

(2) By the oxidation of the fatty acids or of acids of the oleic acid series, e.g. erucic acid

→ brassylic acid

(3) By the reduction of unsaturated diearboxylic acids, eg fumanic acid  $\rightarrow$  succinic

(4) By the action of silver (powder) on monoiodo- (or bromo)- fatty acids;  $e.g. \beta$ -iodopropionic acid -> adipic acid

Exception —The reaction is abnormal when a-bromisobutyric acid is used, di- and tri- alkyl-

glutaric acids being produced.

(5) By the conversion of monohalogen. substituted fatty acids or of the halogen addition products of the alkylenes  $C_nH_n$  into cyanderivatives and saponification of the latter, e.g. cyanacetic acid -> malonic acid; ethylene dicyanide -> adipic acid.

(6) By the replacement of the hydrogen atoms of the CH<sub>2</sub> group in the malonic esters by various alkyl-groups; e g sodium malonic ester and methyl iodide  $\rightarrow$  180-succinic ester.

(7) By the blectrolysis of concentrated solutions of potassium alkyl salts of the dicarboxylic acids, eg potassium ethyl malonate  $\rightarrow$ diethyl succinate

(8) By the introduction of acid residues into the acctoacetic esters, and decomposition of the product with alkali,  $\epsilon g$ , acctosuccime ester  $\Rightarrow$  succime acid

(9) By the decomposition of tricarboxylic acids containing two carboxyl- groups attached to the same carbon atom,  $e\,g$ , ethanetri-

earboxylic acid  $\Longrightarrow$  succinic acid

The acids of this series behave differently on the application of heat, depending upon the position of the carboxylic groups Oxalic acid, the first member of the series, breaks down on heating mostly into carbon dioxide, carbon monoxide, and water, and partly into carbon dioxide and formic acid. This latter decomposition is typical of those homologues of oxalic acid in which the two carboxyl- groups are attached to the same carbon atom, they are the \$-dicarboxylic acids, and of these malonic acid is the type On application of heat, malonic acid splits into acetic acid and carbon The next class, the γ-dicarboxylic acids, in which the two carboxyl- groups are attached to adjacent carbon atoms, when heated, lose a molecule of water and pass into anhydrides Succinic acid is the type of these acids. Acids, such as glutaric acid, in which the earboxyl- groups are attached to two carbon atoms separated by a third, behave similarly to the  $\gamma$ -dicarboxylic acids, but form anhydrides less readily When the carbon atoms to which the carboxyl- groups are attached are separated by two carbon atoms, eq adipic acid, they do not influence one another on the application of heat For these reasons, the paraffin dicar-boxylic acids are arranged in different groups, the general properties of which will be discussed under Malonic Acid, Succinic Acid, and Glu-TARIC ACID, which are types of the several groups mentioned above

(For melting-points of the dibasic acids, v Massol, Bull Soc chim [3] 21, 578, melting-points of their esters, v. Schneider, Zeitsch physikal Chem. 22, 228, electrical conductivity of the acids, v Walden, ibid 8, 448, volatility in steam, v. Auwers, Annalen, 292, 159)

DIBENZIL v. KETONES.

DI-150-BUTYLACETIC ACID v DECATOIC ACIDS.

#### 2 6 DI 150-BUTYLPIMELIC ACID

 $CH_2[CH_2 CH(C_4H_9) CO_2H]_2$ 

Prepared by heating 1.5 di-180-butylpentanetetracarboxylic acid at 200°-220° (Perkin and Prentice, Chem. Soc Trans 1891, 842); mp 82°-84°.

**DICENTRINE**  $C_{20}H_{21}O_4N$  occurs with protopine in *Dicentra pusilla* (Sieb and Zucc) and possibly in *D. formosa* (Walp.). Prismatic crystals, m r  $168^{\circ}-169^{\circ}$ ,  $\lceil \alpha \rceil_D+62^{\circ}$  in chloroform Salts crystalline (Heyl. Arch. Pharm. 1903, 241

313; Asahina, tbid. 1909, 247, 201). Produces slight narcosis (lwakawa, Arch expt. Path Pharm. 1911, 64, 369) Gadamer has stated (Arch. Pharm. 1911, 249, 680) that dicentine resembles glaucine in its properties, and suggests that it is glaucine in which the two—OMe groups in positions 5 and 6 are replaced by a dioxymethylene group

by a dioxymethylene group

DICETYLADIPIC ACID C<sub>38</sub>H<sub>74</sub>O<sub>4</sub>. Propared by heating dicetylbutanetetracarboxyle acid to 205° (Lean, Chem Soc. Trans. 1894, 1016) Two isomerides are thus produced, which may be separated by crystallisation from alcohol, α acid, m p 42°-43°, β-acid, m.p 32°-34°

**DICETYLIC ACID** ('H(('1<sub>8</sub>H<sub>32</sub>)<sub>2</sub>COOH is obtained by heating dicetylmalome acid; m.p. 69°-70° (Guthzeit, Annalen, 206, 365)

#### DICETYLMALONIC ACID

The ethyl ester is formed by treating diethyl-malome ester with sodium methoxide and cetyl rodide, the ester is decomposed with a concentrated solution of caustic potash, mp 86°-87° (Guthzeit, Annalen, 206, 362)

**DICHLORAMINET.** Toluene-p-sulphone di-

chloramme.

# 2 5-DICHLOROANILINE



Prepared by reducing 2–5-dichloronitiobenzene with non-and hydrochloric acid; mp 50°, b p 246°/744 mm. Used in making the chloramine dyes

# 2.5-DICHLOROANILINE-4-SULPHONIC

2 5-dichloroaniline Soluble in hot water. Solum salt with 5 molecules H<sub>2</sub>O readily soluble (Noclting and Kopp, Ber 1905, 38, 3513)

# 1.5-DICHLOROANTHRAQUINONE



is obtained by boiling an aqueous solution of alkali anthraquinone-1 5-disulphonate with hydrochloric acid and sodium chlorate (Ullmann and Knecht, Ber 1911, 44, 3125, Farbenfab. vorm F Bayer & Co, D R P 205195) Also by suspending 1 5-diaminoanthraquinone in hydrochloric acid, diazotising at 5°-10°, and treating the solution with cuprous chloride dissolved in hydrochloric acid (idem D R. P. 131538) Citron-yellow needles, readily soluble in hot toluene, sparingly soluble in alcohol, mp 251° Used in making indanthrene violet (q v art Indanthrene).

#### 3 3'-DICHLOROBENZIDINE

$$NH_2$$
 $Cl$ 
 $NH_2$ 
 $NH_2$ 

Prepared by chlorinating a sulphuric acid solution of diacetyl benzidine (made by boiling benzidine with glacial acetic acid) with bleaching powder or sodium hypochlorite solution, filtering, adding hydrochloric acid, and hydrolysing the product (Levinstein, Eng. Pat. 25725, 1896) May also be obtained by reducing o-chloronitrobenzene in an alkaline solution by means of zinc dust (cf. Cohn, Ber. 1900, 33, 3551).

DICHLOROCARBAMIDE v HYDRAZINES. DICHLOROPHTHALIC ACIDS v. PHTHALIC

2' . 5'-DICHLORO-4'-SULPHO-1-PHENYL-3-METHYL-5-PYRAZOLONE. Prepared by condensing ethyl acetoacetate with 2 5-dichlorophenylhydrazine 4-sulphonic acid (D. R. P. 222405). Used in making Xylene yellow (see PYRAZOLONE COLOURING MATTERS)

DICHROITE v IOLITE.

DICYANDIAMIDE, Cyanoquanidine

# NH2(NH )C NH C+N

Obtained by the polymerisation of cyanamide in aqueous solution on long standing or on warming, especially after addition of a drop of ammonia (Belstein and Geuther, Annalen, 108, 99, 123, 241, Haag, ibid 122, 22), by treating thiourea with mercuire and ine (Montecchi, Gazz chim ital 28, ii 434) Prepared by heating ammonium sulphate and sodium cyanamide on the water-bath, cyanamide separates as a vellow oil, which, after some hours, with frequent shaking, changes into dicyandiamide (Hermann, Monatsh 1905, 1025) Soll and Stutzer (Ber 1908, 4534) make crude calcium nitride into a paste with water and heat to boiling The hot liquid is filtered by suction, and dicyandiamide separates out from the filtrate Crystallises in needles; mp 205° (Pohl, J pr Chem 77, [2] 533) By warming with dilute acids, it is transformed into dicyandiamidine (quanyl urca) NH<sub>2</sub>(NH )C NH CO NH<sub>2</sub> It is used in quantitative analysis to separate nickel from cobalt The solution containing mickel and cobalt is made alkaline, and to it is added a small quantity To this is added dicyandiof sugar solution amidine sulphate (prepared by dissolving di-cyandiamide in dilute sulphure acid), whereby the nickel salt of dicyandiamidine is precipitated, the cobalt remaining in solution. The precipitate is filtered through a Gooch crucible, dued at  $115^{\circ}$ – $160^{\circ}$ , and weighed as anhydrous mekel dicyandiamidine Ni(C<sub>2</sub>H<sub>5</sub>N<sub>4</sub>O)<sub>2</sub> (Grossmann and Schuck, Analyst, 1910, 247, 1909, 455 . 1907, 273, 394) Dievandiamide is reduced by zine and dilute hydrochloric acid at the ordinary temperatures to methylamine and guanidine, the methylamine being formed by the reduction of the prussic acid first produced (Bamberger and Seeberger, Ber 1893, 1583), a mixture of nitric and sulphuric acids converts it into nitrodicyanodiamidine (Thiele and Uhlfelder, Annalen, 303, 107); it condenses with dialkyleyanacetic or malonic esters yielding pyrimidine derivatives, thus with diethylevanacetic ester a condensation product is obtained which yields diethylbarbituric acid on treatment with sulphuric acid (Bayer & Co, D R P 165223; Chem Zentr 1906, 1 514) All manures containing calcium cyanamide contain also dicyandiamide, formed by the spontaneous polymerisation of the calcium cyanamide; it is not poisonous to plants if not used in too large quantities (Perotti, Atti R. Accad. Lincei, 15, 48; Chem Zentr. 1909, 1 1497; Inouye, J. Soc.

Chem Ind. 1909, 1054; Aso, ibid.). The Cyanide-Gesellschaft (Chem. Zentr. 1904, u. 1079; D.R P. 154505) treat calcium carbide or a carbideforming mixture with nitrogen, lixiviate the mass with water, and use the product, which contains calcium cyanamide, dicyandiamide, &c, as a manure (v. also Frahk, Zeitsch angew Chem 19, 835) The constitution of dicyandi-Chem 19, 835) amide is regarded as NH<sub>2</sub>(NH.)C·NH·C·N or the tautomeride (NH<sub>2</sub>)<sub>2</sub>C N C·N (Bamberger, Ber 1883, 1459, 1891, 899; Biltz, J pr. Chem 77, In 1533, Prianichnikov, J. Soc Chem. Ind 1909, 724)

DIDYMIN v SYNTHETIC DRUGS

DIDYMIUM v Praseodymium and Neo-DYMIUM

DIETHYLACETIC ACID. Pseudo-ca proic acid (v CAPROIC ACID)

DIETHYLALLOXAN v ALLOXAN.

**DIETHYLAMINE** v. under Ethylamines, art ETHYL

p-DIETHYLAMINOBENZOIC ACID

$$CO_2H$$
  $N(C_2H_5)_2$ 

Prepared by the action of carbonyl chloride on diethylaniline or by ethylating p-aminobenzoic acid, mp 188°.

DIETHYL-m-AMINOPHENOL

$$\underbrace{OH}_{N(C_2H_{5})_2}$$

Formed by heating m-aminophenol hydrochloride with ethyl alcohol under pressure (Bad Aml und Soda Fabrik, D. R. P. 44002), by heating resorcinol with diethylamine (Leonhardt & Co, D R P 49060), by alkalı fusion of diethylaniline-m-sulphonic acid (Soc Chem Ind Basle, D R P. 44792) For details of the last-named process as used on the large scale, see Cain, Intermediate Products for Dyes, mp 78° bp 201°/25 mm Used in the manufacture of rhodamine B

DIETHYLANILINE  $\langle N(C_2H_5)_2 \rangle$ Ob-

tamed by heating aniline hydrochloride with ethyl alcohol under pressure, or, better, by heating aniline hydrobromide with (thyl alcohol at 145°-150° (Stadel, D. R. P. 21241), or by ticating aniline with ethyl alcohol in presence of iodine at 235° (Knoll & Co, D. R. P 250236), m p -381°, b p 2165°, spgr 0930 at 18°. DIETHYLANILINE-m-SULPHONIC ACID

$$\underbrace{\begin{array}{c} \mathrm{SO_3H} \\ \mathrm{N}(\mathrm{C_2H_5})_2 \end{array}}$$

Is prepared by acting on diethylaniline with fuming sulphuric acid

DIETHYLDIACETOSUCCINATE v KETONES. DIETHYL OXALACETATE v KETONES DIETHYLSAFRANIN v. AZINES

**DIETZEITE.** A double rodate and chromate of calcium crystallising in the monoclinic system, discovered in 1891 in the sodium nitrate deposits of Atacama, Chile Analyses lead to the formula  $7\text{Ca}(\text{IO}_3)_2$ ·8CaCrO<sub>4</sub>, but the simple double salt formula  $\text{Ca}(\text{IO}_3)_2$ ·CaCrO<sub>4</sub> appears more probable Distinct crystals are rare, the mineral usually forming crystalline fibrous aggregates of a dark gold-yellow colour. Sp gr. DIFFUSION. 521

3.698 It is soluble in hot water; on cooling crystals of hydrated calcium iodate  $(Ca(IO_3)_2 + 6H_2O)$  separate, leaving the calcium chromate in solution. The simple calcium iodate  $(Ca(IO_3)_2)$  called lautarite occurs in the same deposits.

L. J S DIFFUSION. In any gaseous mixture of liquid solution, even when protected from mechanical disturbances and kept at a uniform temperature throughout, the composition ultimately becomes the same at all points, whatever the original distribution of the gaseous or dissolved substances may have been The process by which this condition is reached is termed 'diffusion,' and it depends, not on the circulation of the unequally distributed substances as a whole, but on molecular movement According to the kinetic theory, a mass of gas which, as a whole, is at rest, is yet the scene of great activity on the part of the constituent molecules are endowed with a high velocity, and alter the direction of their motion only when they collide with one another or with the walls of the containing vessel If this point of view is adopted, then the occurrence of diffusion in a mixture of irregularly distributed gases becomes intelligible There are, moreover, many experimental grounds for extending the kinetic theory to the liquid state, and for the view that the molecules of a liquid also are in a state of constant motion With this conception the phenomenon of diffusion of a dissolved substance is obviously in harmony

#### Diffusion of Gases.

When two vessels containing different gases at the same temperature and pressure are put into free communication, each gas penetrates the other until its partial pressure is uniform in both vessels. The experimental investigation of this phenomenon on quantitative lines is not easy, but it appears to be well established that the diffusion coefficient of each gas varies from point to point of the diffusion system (see Loschmidt, Sitzungsbei. Akad Wiss Wien, 1870, 61, 367, 1870, 62, 468, von Obermayer, 500 dt 1880, 81, 1102, 1882, 85, 147, 748, 1883, 87, 188; 1887, 96, 546, Waitz, Ann Physik, 1882, 17, 201, 351, Lomus, 1001, 1909, 29, 664)

When the concentration of a gas is maintained constant at one point of a gas system and zero at another point, a static condition of affairs is established analogous to what is observed when one end of a metal bar is kept at a high temperature and the other end at a low temperature. If, for instance, a tall cylinder, communicating freely with the atmosphere, contains at the bottom a layer of caustic alkali, there is a regular concentration gradient from the top downwards, the carbon dioxide being completely absorbed at the surface of the alkali Provided the concentration difference between the top and bottom of the cylinder is maintained constant, a steady condition of diffusion is attained, and the rate of flow of the carbon dioxide, as deduced from the amount absorbed by the alkalı, is inversely proportional to the length of the diffusion column (see Brown and Escombe, Phil. Trans B, 1900, 193, 223, also Winkelmann, Ann Phyrik, 1884, 22, 1, 152)

Much attention has been devoted to the phenomenon of diffusion across porous dia-

phragms, through capillary tubes, and from Graham showed (see Cheminarrow apertures cal and Physical Researches, p 44) that the exchange of two gases across a perous diaphragm, the pressure on the two sides remaining the same, takes place in such a way that the volumes of the gases diffused in a given time are inversely proportional to the square roots of their densities. This observation is of very great interest in connection with the kinetic theory, from the assumptions of which it follows that the mean velocity of the molecules of a gas is inversely proportional to the square root of its Giaham's law is not accurate except when the diaphragm is extremely thin, but it is nevertheless the basis of a method for separating Thus, for two gases of different densities instance, when a mixture of hydrogen and oxygen in the proportion of 2 1 by volume is passed at a suitable rate through a train of claypipe stems, the gas collected at the further end will ignite a glowing splint By experiments of this kind it was shown that atmospheric nitrogen is a mixture of at least two gases of different density (see Rayleigh and Ramsay, Phil Trans 1895, 186, 206). The operation of separating gases by this method is termed 'atmolysis'

In the passage of a gas through a capillary tube—' transpiration,' as it is sometimes called—friction generally plays a considerable part, but it has been found that if the diameter of the tube is small in comparison with its length and with the mean free path of the gas molecules, then the quantity of gas passing through a given tube in unit time is inversely proportional to the square root of the density (Knudsen, Ann. Physik, 1999, 28, 75)

The time required for the escape or 'effusion' of a given volume of gas through a fine aperture in a thin plate into a vacuum is approximately proportional to the square root of the density of the gas. The thinner the plate and the smaller the aperture, the more nearly is this relationship fulfilled (see Graham, Chemical and Physical Researches, p. 88., Bunsen, Gasometry, p. 121, Timofécif, Zeitsch physikal Chem. 1890, 6, 586, Donnan, Phil. Mag. 1900, 49, 423, Emich, Monatsh. 1903, 24, 747, Knudsen, Ann. Physik, 1909, 28, 999). An apparatus for determining the relative density of two gases, based on this principle, has been described by Bunsen (lee) (v. Specific gravity)

The passage of a gas across a diaphiagm takes place, not only when the diaphragm is porous in the ordinary sense, but also when it has the power of dissolving or absorbing the gas Thus, at high temperatures, platinum and palladium absorb considerable quantities of hydrogen, and this gas readily diffuses through the walls of a vessel made of other metal Winkelmann (Ann Physik, 1901, 6, 104, 1902, Nicol & Parnell, Phil Mag, 1901, 6, 104, 1902, Nicol & Parnell, Phil Mag, 1901, 8, 1, Schmidt, Ann Physik, 1904, 13, 747) has shown that the rate of diffusion of hydrogen through glowing platinum or palladium is not, as might be expected, proportional to the pressure of the hydrogen, at lower pressures diffuses relatively more rapidly. the gas The experimental data seem to indicate that it is the atoms, not the molecules, of hydrogen which can pass through the metal

522 DIFFUSION.

On the ability of hydrogen and the inability of other gases to diffuse through heated palladium, there is based the osmotic cell described by

Ramsay (Phil Mag 1894, 38, 206)

The rate of diffusion of gases across a layer of water or aqueous solution depends mainly on the solubility and the density of the gas. According to Exner, who made experiments on the diffusion of gases across a film of soap solution, the velocity is, cateris paribus, directly proportional to the absorption coefficient of the gas, and inversely proportional to the square root of its density (Exner, Sitzungsber. Akad. Wiss Wien, 1874, 70, ii. 465, cf. Muller, Ann Physik, 1891, 43, 554). Other workers, who have studied the diffusion of gases through a jelly, through a layer of water supported by a plate of hydrophane, or through a rubber membrane, find that Exner's law is only an approximate statement of the facts (see Wichelbewsky, Pogg. Ann. 1876, 158, 545, Ann Physik, 1879, 8, 29, Hufner, Zeitsch physikal Chem. 1898, 27, 227, Hagenbach, Ann. Physik, 1898, 65, 673). Except for oxygen, the diffusion soefficients of gases are smaller in gelatine solution than in pure water.

#### Diffusion of Dissolved Substances.

The foundation of our knowledge of this subject was laid by the classical experiments of Thomas Graham (see Chemical and Physical Researches, pp. 444-600). In his experiments, 700 cub cm of water were placed in a cylindrical jar, and then 100 cub cm of the solution to be diffused were carefully conveyed to the bottom of the jar by a pipette. After a suitable time, portions of 50 cub cm were successively drawn off, these portions were evaporated, and the amount of substance that had diffused into each layer was thus ascertained Graham's comparative experiments on the diffusive power of different substances in aqueous solution led him to draw a distinction, which has become of the highest importance, between the behaviour of easily crystallisable substances, and that of substances which are marked by the absence of the power to crystallise So far as diffusibility is concerned, the distinction in question is based on the following figures, representing approximate times of equal diffusion hydrochloric acid, 1, sodium chloride, 23, sucrose, 7, magnesium sulphate, 7, albumin, 49, caramel, 98. The difference in diffusive power of crystal loids and colloids is very evident from these figures

The significance of the quantitative results obtained by Graham was emphasised by Stefan (Sitzungsber Akad Wiss Wien, 1878, 78, ii 957; 1879, 79, ii. 161), who showed that they were in harmony with Fick's diffusion law, that is, the diffusion of substances in aqueous solution is comparable with the distribution of heat in conductors. Fick's law is expressed algebrai-

cally by the formula  $dS = -kq \frac{dc}{dx} dt$ , where dS is

the amount of substance crossing a section of the diffusion column in time dt, the sectional area of this column being q sq. cm, and dc/dx being the concentration gradient, k is known as the 'diffusion coefficient'

The diffusion of dissolved substances, notably electrolytes, has been the subject of repeated

nvestigation, but the method employed has not differed materially from that of Graham, described above (see Scheffer, Zeitsch. physikal. Chem. 1888, 2, 390; Arthenius, sbid. 1892, 10, 51; Abegg, sbid. 1893, 11, 248; Euler, Ann. Physik, 1897, 63, 273; Abegg and Bose, Zeitsch. physikal Chem. 1899, 1, 17; Thovert, Compt. rend. 1902, 134, 826; Graham, Proc. Roy. Soc. 1903, 72, 212; Oholm, Zeitsch. physikal. Chem. 1904, 50, 309) The general result of these investigations has been to substantiate the validity of Fick's law, although the opposite opinion also has been expressed (see Wiedeburg, Zeitsch. physikal. Chem. 1892, 10, 509)

Special interest has attached to the diffusion of electrolytes since the enunciation of Arthenius's dissociation theory, according to which the ions of an electrolyte in aqueous solution are to be regarded in many respects as separate units The researches of Kohlrausch have proved that the contribution which the one ion of an electrolyte makes to the equivalent conductivity is independent of the other ion associated with it, and Nernst has shown that the mobility which is thus characteristic of each ion must be a determining factor in the diffusive power of any electrolyte of which the ion forms a part (Zertsch physikal Chem 1888, 2, 613). He has further deduced a theoretical relationship between the diffusion coefficient of a binary electrolyte and the conductivities of the two ions, and the values of the diffusion coefficient calculated on this basis are in good agreement

with the results of experimental work. By allowing substances to diffuse into a jelly the effects of convection currents and mechanical disturbances are eliminated and the difficulties attending the experimental study of liquid diffusion are diminished. The rate of diffusion is, at most, very slightly lower in a jelly than in water at the same temperature. Experiments, in which the ends of cylinders of agar-agar jelly were immersed in aqueous solutions of various substances, have shown that the total amount of substance diffusing into the jelly is proportional to the square root of the time during which diffusion has taken place. This is a result in harmony with Fick's law (see Voigtlander, Zeitsch physikal Chem 1889, 3, 316; cf. Chabry, Journ de Physique, 1888, 7, 115, Morse and Pierce, Zeitsch physikal Chem 1903, 45, 589, Hausmann, Zeitsch anorg Chem. 1904, 40, 110).

Experiments have been made also on the diffusion of various electrolytes and non-electrolytes in alcohol (Kawalki, Ann Physik, 1894, 52, 166, 300, 1896, 59, 637), of various metals in mercury, fused lead, &c (Meyer, ibid 1897, 61, 225; 1898, 64, 752, Humphreys, Chem Soc Trans 1896, 69, 243, 1079, Roberts-Austen, Phil Trans 1896, 187, 383, von Wogau, Ann. Physik, 1907, 23, 345), and of chlorine, bromine, and iodine in some organic solvents (Euler, Ann. Physik, 1897, 63, 273, Walden, Zeitsch. Elektrochem 1906, 12, 77). In the last-mentioned connection, Walden has shown that, when a given halogen is considered, the product diffusion coefficient × viscosity of solvent' is a constant, independent of the nature of the solvent. The value of the product for the different halogens is inversely proportional to the square root of the molecular.

DIGITALIS. 523

weight (cf Pissarjewsky and Karp, Zeitsch physikal. Chem. 1908, 63, 257).

The different rates of diffusion found by Giaham for different substances suggested to him that the process of diffusion might be useful in separating the constituents of an aqueous solution In this connection, he found that the difference in diffusive power between crystalloids and colloids, referred to above, was greatly accentuated by the interposition of a colloid membrane between solution and solvent, and the method of separation by diffusion through a septum of gelatinous matter he termed dialysis' A sheet of parchment paper, for instance, is tied tightly over the end of a cylindrical or bell-shaped glass vessel, provided with a flange, and into the receptacle so formed, a 'dialyser,' as it is called, the mixed solution of crystalloids and colloids is poured The outside of the parchment membrane is bathed with water, which is frequently renewed, and the result is that the contents of the dialyser, after a few days, are found to be practically free from crystalloids Instead of the apparatus just described, a sausage-skin dialysei may be employed the parchment tube is charged with the solution of crystalloid and colloid, and is hung up by its ends in a vessel of water

Collodion may be used as a membrane instead of parchiment, and it is obtainable in any desired form by the evaporation of the ordinary collodion solution (see Lillie, Amer J Physiol 1907, 20, 127, Schryver, Proc Physiol

Soc 1908, 23)

Dialysis plays a considerable part in the hineal production of beet sugar. In the technical production of beet sugar diffusion process for the extraction of the sugar, beet cuttings are treated with hot water treatment kills the protoplasm, and the cell membrane, which in the hving state is impermeable to sugar, now allows this substance to pass out into the water. The high tempera-The high temperature also favours rapid diffusion. The albuminous substances in the beet juice however, cannot readily penetrate even the dead protoplasm, which acts as a differential membrane, and thus leads to a dialytic separation of the sugar from some of the non-sugar constituents of the cell sap In the treatment also of beet sugar molasses dialysis with parchment membranes is employed. The result of this procedure is to reduce the salt content of the molasses, and so make possible the crystallisation of a further portion of the sugar (v SUGAR)

#### Diffusion of Solids.

The efficiency of certain cementation processes indicates that one solid can slowly penetrate another The occurrence of diffusion in solids has been proved and investigated by Roberts-Austen (Phil Trans 1896, 187, 383), who finds that at temperatures between 100° and 200° there is an appreciable diffusion of gold into solid lead in 30 or 40 days, although the penetration is much slower than into the fluid metal Experiments lasting over four years have shown that gold diffuses into solid lead, even at the ordinary temperature (Roberts-Austen, Proc. Roy Soc. 1900, 67, 101) J. C P.

DIFORMIN v FORMINS. DIGALENV. DIGITALIS and SYNTHETIC DRUGS

DIGALLIC ACID (Gallotannic acid, tannin) v. TANNIC ACIDS.

DIGESTER v AUTOCLAVE.

DIGIFOLIN, DIGIPAN, DIGITALEIN, DI-TALIN, DIGITALOSE, DIGITALYSATUM, GITALIN. DIGITALOSE. DIGITOFLAVONE, DIGITONIN, DIGITOPHYL-LIN, DIGITOXOSE v DIGITALIS

DIGIPOTENE. A preparation of digitalis

DIGIPURATUM : DIGITALIS DIGITALINUM VERUM v DIGITALIS

Digitalis. Foxylore Digitalis folia, B.P. Digitalis, U.S.P. (Digitale, Fr., Fingerbut, Get ) The foxglove, Digitalis purpurea (Linn ), had been for some centuries a popular remedy for dropsy, when it was introduced into scientific It is medicine by Withering, about 1775 principally used for its action on the heart, it slows the beats, and makes them more powerful and regular, so that the output is increased Only the leaves are official, the seeds have a similar action, due, however, to different brinciples

Foxglove is a biennial plant, and the Brussels conference agreed that only the second year's leaf should be used, but, except for the greater difficulty of identification, there is no objection to using the first year's rosettes, which are as good as the flowering plant (Straub, Arch Pharm 1917, 255, 198) Cultivated plants up to the beginning of the flowering period, are fairly constant in activity, variations appear when the seeds upon—It would probably be the more economical to cultivate only first-year plants Mature wild plants differ greatly in various localitics (Straub). The time for gathering these is July and August, the British Pharmacopa ia prescribes leaves from plants beginning to flower. For further detail, see Nativelle (J. Pharm. Chim. [iv | 20, 81), and for culture, harvesting, and conservation, Newcomb (Amer J. Pharm 1912, 84, 201). Thapsil, which has a similar action to D purpurea, but is three times as strong, see Farwell and Hamilton, Amer J Pharm 1917, 89, 147

The active principles of digitalis (as of other plants with a like action, such as strophanthus and squill) are complex glucosides, often amorphous, and this explains the great difficulties and numerous contradictions with which the chemistry of digitalis is beset fusion is increased by the use of the same name (digitaline) for different substances, and by the application of different names to the same substance According to our present knowledge, cold water extracts from the leaves a mixture of glucosides, but hardly dissolves digitoxin, which may be subsequently extracted by alcohol From the aqueous extract chloroform removes the 'gitalm' fraction, and leaves behind one or more plucosides, designated as the 'digitalein' fraction, which is less active than either gitalin or digitoxin, and has not been obtained even approximately pure torically, the first moderately pure, but amor phous, a tive preparation was 'digitaline' of Homolic (J. Phaim Chim 1845, [iii.] 7, 57), undoubtedly a mixture Nativelle (J Pharm. Chim [v] 1869, 9, 225, 1872, 16, 430 1874, 20, 81) later prepared a 'digitaline cristallisée,' a commercial specimen of which Schmiedeberg 524 DIGITALIS.

(Arch. exp. Path. Pharm. 1874, 3, 16) declared to consist very largely of a substance to which he gave the name digitoxin, and as this substance was afterwards principally investigated in Germany, under this name, Nativelle's designation has been displaced, in spite of Fiench protests (e.g. Arnaud, Compt. rend. 1889, 109, 679) Kilmin, who has published something like forty papers on the chemistry of digitalis (Bei 1890, 23, 1555 to 1918, 51, 1613; and Arch Pharm 1892, 230, 250 to 1916, 254, 255), at one time did not agree with the identification of Nativelle's digitaline with Schmiedeberg's digitoxin, and considered the former substance identical with a crystalline leaf glucoside described by himself as digitophyllin (Arch Pharm 1897, 235, 425) Later, Kihani (Ber. 1898, 31, 2454) considered digitaline cristallisée and digitoxin very closely similar, but doubtfully identical. An attempt to restore the historically correct name for this glucoside would lead to still greater confusion, especially as German writers mean by digitalin a totally different glucoside from the seeds

The 'gitalm' fraction is, on account of its solubility in water and its physiological action, probably of most importance therapeutically, it was little studied until Kraft (Schweiz Wochenschr Chem Pharm 1911, 49, 161, 173, 236. Arch Pharm 1912, 250, 118) described the isolation of a crystalline glucoside gitalm Kiliam (Arch Pharm 1914, 252, 13) has declared Kriaft's preparation to be a mixture, but in any case it contains one or more crystalline substances of great activity, having the advantage over digitoxim of solubility in water, at least in the crude state. Indeed, the gitalm fraction seems to have been utilised some years before Kraft by Cloetta (Muench mediz Wochenschi 1904, 51, 1466), whose 'digitoximum solubile' is not digitoxim in a water soluble form (Kiliam, Ber 1907, 40, 2996), but consists largely of gitalm (Gottlieb, c.g. Arch. exp. Path. Pharm 1918, 82, 117).

1918, 83, 117)

For the preparation of their substances, both Homolle and Nativelle extracted the powdered leaves with 50 pc alcohol, and purified the extract with lead acetate. After removal of the excess of lead, Homolle evaporated to a syrup and precipitated in the cold with tannic acid The tannic acid precipitate was washed with tepid water, mixed with an equal weight of lead oxide, dried, and extracted with 90 p c alcohol After treatment with charcoal and evaporation, the residue was extracted with ether to remove impurities Tannin, apparently introduced by Homolle, was later applied to the separation of k-strophanthin (q v), and seems to be still in use for such commercial preparations as digipuratum (see below), but it is, of course, only a general colloidal precipitant, and cannot be expected to act specifically Nativelle moistened the powdered leaves with a solution of neutral lead acetate, then extracted with 50 pc alcohol, and purified the extract repeatedly with charcoal and lead acetate finally extracted with chloroform, and thus separated off the 'digitalein' fractica, which enabled him to obtain a crystalline substance Schmiedeberg (lc) introduced the preliminary extraction of the leaves with cold water, thus removing the gitalin and digitalein fractions,

and then, after pressing the leaves, he extracted with 50 pc. alcohol. After precipitation with basic lead acetate, and removal of the excess with ammonia, the alcohol was evaporated, while the solution was kept neutral. A partly crystalline substance separated, which was dried on the water-bath and extracted with chloroform. The residue, after evaporation of the chloroform, was washed with ether (which removes fat, luteolin, &c.), or with boiling petroleum ether, and then crystallised the digitoxin from 80 pc alcohol with charcoal. Kiliani's method of preparing digitoxin (Arch. Pharm 1895, 233, 311) is similar

Digitaline cristallisée, Digitoxin  $C_{34}H_{54}O_{11}$  (Kiliam) forms hydrated needles and leaflets from aqueous alcohol, which become anhydrous and sinter at  $145^{\circ}$ - $150^{\circ}$ , and anhydrous prisms from methyl alcohol and chloroform, melting at  $245^{\circ}$  ( $243^{\circ}$ - $245^{\circ}$  Arnaud) Kraft (Arch. Pharm 1912, 250, 118) could not obtain the hydrated form, and concludes Kiliam's supposed

hydrate was contaminated with gitalin

The yield of digitoxin is said to be 0.1 pc of the leaves, but Straub (lc) has concluded from physiological evidence that the amount present is very variable, and may considerably exceed 0.1 p.c. Merck's commercial preparation is stated by Kiliani to be about 90 pc pure 0 65 gram digitaline (digitoxin) dissolves at 14° in 100 cc absolute alcohol (Arnaud) Digitoxin dissolves readily in chloroform, only slightly in ether, not in water. It gives a characteristic colour reaction, glacial acetic acid and concentrated sulphuric acid are employed, 100 cc of each being previously diluted with 1 cc of 5 p.e. aqueous ferrie sulphate solution trace of digitoxin is dissolved in 1 cc of the acetic acid, and 2 cc of the sulphunc acid are allowed to flow underneath, in a narrow test tube. The zone of contact becomes duty brown, and after a few minutes the acetic acid This, Keller's begins to become indigo blue reaction, is still positive with 0 l mg in 1 c c of acetic acid Concentrated hydrochloric acid dissolves digitoxin in the cold to a colourless solution, becoming bluish-green on warming Alcoholic sodium hydroxide hydrolyses to digitozime acid  $\rm C_{34}H_{56}O_{12}$ , mineral acids hydrolyse to two molecules of digitoxose

#### CH, CHOH CHOH CHOH CH, CH O

m p  $101^{\circ}$ , and one of digitoxigenin  $C_{22}H_{32}O_4$ , m p  $230^{\circ}$  (v. e.g. Kihani, Arch Pharm. 1913, 251, 562) Digitoxose is a peculiar sugar, and gives Keller's reaction, digitoxigenin does not

Gitalin C<sub>18</sub>H<sub>48</sub>O<sub>10</sub> (?), mp 150°-155°, is amorphous, and was obtained by Kiaft by a method analogous to that of Homolle (v. supra), and extracting the glucoside from aqueous solution by shaking with chlorofoim, after drying, the chlorofoim solution is precipitated by light petroleum. Gitalin is soluble in 600 parts of water, and very readily in chloroform. By dissolving in 15 parts of alcohol and adding 0.75 part of water, a crystalline gitalin hydrate C<sub>28</sub>H<sub>48</sub>O<sub>10</sub>,4H<sub>2</sub>O, mp. 75°, is separated. Solutions of gitalin deposit on keeping mixtures of gitalin and anhydrogitalin C<sub>28</sub>H<sub>46</sub>O<sub>9</sub>, mp. 255°, crystallising from dilute alcohol and almost insoluble in chloroform. On hydrolysis by acids, gitalin, its hydrate, and its anhydride all

DIGITALIS.

yield digitoxose (see above), and hence they all stance, such as digitoxin (Rapp, Ic.), or ggive Keller's reaction; the other fission product is anhydrogualigenin C<sub>2.</sub>H<sub>34</sub>O<sub>5</sub>, mp 216°-219°, crystallising from alcohol As already stated, Kiliani considers gitalin to be a mixture, of anhydrogitalin with other substances, but this does not dispose of Kraft's contribution to the subject

Digitaleïn (Schmiedeberg) differs from gitalin and digitoxin in not producing the typical

systolic arrest of the heart

Commercial Preparations of Digitalis.—A considerable number of these have been introduced of late years, mostly in Germany Digalen (Hoffmann-Laroche & Co), introduced in 1904, has been one of the most successful Cloetta's 'digitoxinum solubile'=impure gitalin' Digifolin and digipan are similar (see above) impure active principles Digipuratum (D. R. P. 227552, 245193, 246571, of Knoll & Co ) is a purified leaf extract freed from digitonin, and apparently consists of the tannin compounds of the gitalin fraction From adigan the digitonin has been removed by cholesterol (see below) Digitalysatum, from fresh piess juice, and other preparations, are purified by dialysis Digita-Ione (Parke, Davis & Co ) is a sterile extract of For a physiological examination of these preparations, see Rapp, Pharm Zentral-H 1914, 55, 961, 978

Methods of Assay.—There is a great variation in the content of active leaf glucosides Straub (lc) found, in percentages of the air-dry leaf

Gitalin + digitalein, cultivated, 0 4-0 64, wild, 0 22-0 89, digitoxin, cultivated, 0 053-0 11, wild, 0 01 0 15

This variability has led to the introduction of standardised leaves ('folia digitalis titiata') and to methods of assay of pharmaceutical preparations (official in the USP for digitalis, strophanthus, and squill) A chemical assay (of digitoxin?) has been described by Keller (Ber Deut pharm Ges 1897, 7, 125), but Ziegenbein (Arch Pharm 1902, 240, 454) has shown that there is no relation between physiological activity and the 'digitoxin' found by Keller's method Barger and Shaw (Pharm. J 1904, [iv] 19,7249) have confirmed this, and shown that of added crystalline digitoxin less than half is recoverable by Keller's method The chloroform extract, weighed in this method, consists, according to Kraft (lc), largely of The assay must therefore be physiological, and is generally based on Houghton's method (J Amer med Assoc 1898), by which the minimum amount is determined necessary to arrest the frog's heart in systole, in, say, I hour (for length of time, cf Gottlieb, below) The isolated frog's heart may also be employed, with graphic registration of its action (Schmiedeberg, Arch. exp Path. Pharm. 1910, 62, 307) For earlier investigations, see Edmunds and Hale (Bull No. 48, Hygienic Lab Washington, 1909) and Focke (Arch. Pharm. 1910, 248, 365); for the later ones, Holste (Zeit. exp. Path. Ther. 1917, 19, 153), Focke (ibid. 1914, 16, 443), Stretch (Arch. 2012, 2012). Straub (Arch. exp. Path. Pharm. 1916, 80, 52), and Gottheb (ibid 1918, 83, 117) The susceptibility of frogs is greater in winter than in summer, and it has been suggested that they should themselves be standardised with a pure sub-

strophanthin (ouabain) (qv), suggested by Heinz, and adopted by the USP. This picsupposes a constant ratio of sensitiveness to dissimilar preparations, which has not been Indeed. Gottlieb (lc.) states that the sensitiveness of frogs to q-stropanthin is fairly constant throughout the year

525

Digitalis seeds contain entirely different glucosides, which have been examined chiefly From an alcoholic extract, purified with lead acctate, the mixed glucosides are precipitated with tannic acid (cf Homolle's method) This gives the commercial so-called digitalinum germanicum, which may be separated, according to Kiliani (Ber 1918, 51, 1613), into 'digitalinum verum,' or digitalin, digitanin,

gitonin, and a fourth glucoside

Digitalin C<sub>35</sub>H<sub>56</sub>O<sub>14</sub>, mp 210°-217°, is less active than digitoxin, and forms amorphous granules, soluble in 1000 parts of water It dissolves in sulphuric acid with a yellow colour, which, on standing, or in the presence of ferric salts or other oxidising agents, becomes cherry-red and purple It is hydrolysed by acids to equimolecular proportions of glucose, digitalose  $C_7H_{14}O_5$ , and digitaligenin  $C_{22}H_{30}O_3$ , needles, m p 210° 212° Digitalose is a syrup, and is oxidised by nitric acid to methoxydihydroxyglutaric acid (Kiliani, Ber 1905, 38, 3622. 1916, 49, 709)

Digitonin ( ${}^{1}_{54}H_{9}$ ,  ${}^{1}_{0.28}$  or ( ${}^{1}_{55}H_{91}O_{8}$ , mp 225°-235°, [ $\alpha$ ]<sub>0</sub>= -49 25°, is obtained in the crude state in a yield of about 45 pc of the 'German digitalin' employed. It is important as an example of a crystalline saponin, and probably also occurs in the leaves ilt has no stimulant action on the heart, and is thera peutically undesirable Digitonin dissolves to a clear solution in 50 parts of cold 50 pc alcohol, is more soluble on heating, and crystal-lises on cooling in fine needles. It is hardly soluble in chloroform, and forms an opalescent solution in water, which froths Keller's Acids hydrolyse to two reaction is negative molecules of glucose, two of galactose, and one of digitogenin  $C_{30}H_{48}O_6$  or  $C_{31}H_{50}O_6$ , needles, mp 240° Like other saponins, digitonin combines with cholesterol On mixing a (preferably hot) solution of 1 gram digitonin in 100 c c of 90 p c alcohol with 0 4 gram cholestorol in 60 cc of 95 pc alcohol, the addition compound C<sub>55</sub>H<sub>94</sub>O<sub>28</sub>C<sub>27</sub>H<sub>16</sub>O crystallises (Windaus, Ber 1909, 42, 238) Similar compounds with lower alcohols and phenol had already been observed by Houdas (Compt. 1end. 113, 648) This reaction has met with wide application On the one hand, digitonin can be removed from digitalis preparations (adigan, above), on the other hand, sterols can be estimated in fats, for as little as 0.1 mg of cholesterol in 1 c c of 90 p c alcohol may be detected (Windaus) Panzer, Zeitsch physiol Chem 1912, 78, 414, Marcusson and Schilling, Chem Zeit 1913, 37, 1001; Berg and Angershausen, ibid 1914, 38, 978; Prescher, Zeitsch Nahr Genussm 1914, 33, 7)

Gitonin.—Commercial digitonin (Merck) contain 5-14 pc. of a closely similar glucoside git nin C<sub>49</sub>H<sub>80</sub>O<sub>23</sub>,7H<sub>2</sub>O, mp 225° (Kraft, lc., Windaus and Schneckenburger, Ber 1913, 46, 2628), which may be crystallised (Kiliani, ib.d. 1916, 49, 701), and also forms cholesterol addition compounds.

Straub (Biochem Zeitsch 1917, 82, 48) finds that on germination the first pair of foliage leaves already contains the typical leaf glucosides, and in 4 months the maximum of 1 pc. of total glucoside is already nearly attained. This is made up of gitalin fraction, ½ pc, digitalein fraction, ½ pc, and digitoxin fraction, ½ pc. For variations, see above.

Digitalis leaves contain luteolin (= digito-flavone') C<sub>15</sub>H<sub>10</sub>O<sub>6</sub>,H<sub>2</sub>O, which imparts a yellow colour to ethereal extracts (Kiliani and Mayer,

Ber 1901, 34, 3577)

Other Drugs with a Digitalis-like Action — Schmiedeberg (Arch exp Path Pharm 1883, 16, 149, especially 162) has enumerated a considerable number of these. Strophanthus and squill (q v) are theiapeutically the most important; see also enonymus, hellebore, and the alkaloid crythrophleme.

Cymurin, a glucoside from the root of Apocynum cannabinum has been obtained crystalline (D. R. P. 255537, of Bayer & Co.) 1000 parts of the root are extracted with boiling carbon totrachloride until no longer bitter After evaporation, the residue is dissolved in 1000 parts of alcohol, and water at 50° is added, until no more resin is precipitated After filtration, basic lead acetate is added, and the excess of lead is removed, after a second filtration, by means of hydrogen sulphide evaporation to 100 parts, the solution is extracted with chloroform, and the chloroform dried with sodium sulphate. Impurities are then precipitated with other, and next the cymain with light petroleum - It is crystallised from methyl alcohol, and then has the composition

 $[a]_D^{20} = +23.5^{\circ}$  in chloroform. It gives Liebermann's cholesterol reaction (v. Strophanthidin), and Koller's digitoxin reaction (above), and is hydrolysed by acids

$$C_{30}H_{44}O_9 + H_2O = C_{23}H_{30}O_5, H_2O + C_7H_{14}O_4$$

into cymarigenin (=strophanthidin, qv) and cymarose, which is probably the methyl ether of digitoxose (Windaus and Hermanns Ber 1915, 48, 979), where the earlier investigations on Apocumum are referred to

Bulagin C<sub>18</sub>H<sub>14</sub>O<sub>4</sub>, from the parotid gland of the Central-American toad Bulo agua, has also been obtained crystalline and pure (Abel and Macht, J. Pharm exp. Therap 1912, 3, 319). It is not a glucoside—For bulotalin from European toads, see Wieland and Weil (Ber 1913, 46, 3315)

Convallamarin C<sub>12</sub>H<sub>44</sub>O<sub>12</sub> and convallarin C<sub>34</sub>H<sub>62</sub>O<sub>11</sub> are crystalline glucosides from Convallaria majalis (Linn) (Lily of the Valley), which plant was deleted from the U.S.P. in 1916 (Walz, Jahrb f Chem 1858, 518)

\*\* a-Anharm  $C_{27}H_{42}O_{10}$ ,  $4H_{2}O$  and  $\beta$ -anharm are crystalline glucosides from the latex of Anharm toxicaim, the Upas tree of Java (Kiliani, Ber. 1910, 43, 3574) Other crystalline glucosides with digitals action are thevetin  $C_{54}H_{84}O_{24}$ , from the seeds of Thevetia nevisfolia (Juss), crystallising readily (de Vrij, Pharm J. 1881, 457; Warden, ibid 1881, 417), and echujin

(C<sub>8</sub>H<sub>8</sub>O<sub>2</sub>)n, from the latex of Adenium Bæhmianum (Schinz), a South-West African arrow poison (Boehm, Arch. exp Path Pharm 1899, 26, 165) The latter substance has an activity intermediate between that of digitoxin and strophanthin

Tanghinin C27H46O8, mp. 182°, from the fruits of Tanghinia venenifera (Madagascar), is crystalline, hardly soluble in water, readily in alcohol and in other, and is not a glucoside (Arnaud Compt iend 1889, 108, 1255) Adonin C24H40O9, from Adonis amurensis (Reg. et Radd ) (Tahara, Ber 1891, 24, 2579), adonadm from A vernalis (Lum) (Cervello, Arch exp Path Pharm 1882, 15, 235), and oleandrin from Nerum oleander (Linn) (Schmiedeberg, lc) are amorphous Many of the above substances are used as arrow poisons The NO Apocynaceae seems to be especially rich in substances with a digitalis-like action (Apocynum, Strophanthus, Thevetra, Adenium, Nerium oleander). Windaus and Hermanns (Ber 1915, 48, 991) have suggested that there is a more or less close chemical relationship between many of the above sub-The non-sugar portion of the glucosides stances genin') seems to be derived from a cholesterollike complex by various degrees of oxidation, the sugars are peculiar and sometimes identical or closely related, and often contain less oxygen than carbohydrates The genins are less active than the glucosides (v k-strophanthmand Straub, Brochem Zeitsch 1916, 75, 132)

DIGITALON v DIGITALIS
DIGITOXIN v DIGITALIS

DIHEPTYLACETIC ACID (C<sub>7</sub>H<sub>15</sub>)<sub>2</sub>CH ('OOH is obtained by treating diheptylacetoacetic ester with potassium hydroxide (Jourdan, Annalen, 200, 116), mp 26°-27°

DIHYDRAZINES v HYDRAZINES
DIHYDROCAMPHOKETONE v KETONES
DIHYDROXY- ACIDS v HYDROXY- ACIDS.
DIHYDROXYANTHRAQUINONES. See ALI-

ZARIN AND ALLIED COLOURING MATTERS.
DIHYDROXYBENZOIC ACIDS v Proto-

CATECHUIC ACID
DIHYDROXYCARBOXYNAPHTHALENE
SULPHONIC ACIDS. See Naphthalene

5 5'-DIHYDROXY-7 7'-DISULPHO-2 2'-DINAPHTHLY CARBAMIDE

Prepared by the action of carbonyl chloride upon an aqueous solution of 2-amino-5-naphthol-7-sulphonic acid containing sodium carbonate, acidifying with hydrochloric acid, and adding common salt (Farbenf. vorm F Bayer & Co., D R. P 116200)

1 5-DIHYDROXYNAPHTHALENE. See

NAPHTHALENE
DIHYDROXYNAPHTHALENE SULPHONIC
ACIDS. See Naphthalene

DIHYDROXYTARTARIC ACID

CO<sub>2</sub>H C(OH)<sub>2</sub> C(OH)<sub>2</sub> CO<sub>2</sub>H

is made by the action of a mixture of fuming sulphuric acid and nitric acid upon dry tartaric at 20°, cooling with ice and neutralising with sodium carbonate when the sparingly soluble sodium salt separates out. Used in making

tartrazine · v Pyrazolone colouring matters and Ketones

DI-HODOFORM v SYNTHETIC DRUGS

DIKA-BREAD. An edible cake made by the natives of Sierra Leone and the Gaboon from the fruit of the mango tree, indigenous to most tropical climates. The fruit iesembles chocolate in appearance, and contains a large quantity of fatty matter. It has occasionally ripened under hot-house cultivation in this country.

DIKA FAT GROUP. Under this name may be comprised several fats which are characterused by high saponification values, like the fats of the Cocoa Nut Oil group, but which differ from them by the absence of volatile fatty acids high saponification value indicates a preponderant proportion of lauric acid. These fats are further characterised by low iodine values These fats are The most prominent of this group are . (1) Dika fat, obtained from the seed-kernels of various kinds of Irringia: I gabonensis (Baill), I Barteri (Hook, fil ), indigenous to West Africa The seeds yield about 65 pc of a yellowish-white fat, which the natives separate by expressing the kernels after boiling with water The fat melts at about 30°, and has an iodine value of about 31 (2) Tangkallak fat, obtained from the fruits of Lepidadenia Wighthana (Nees; BI), Cyhcodaphne sebrfera (BI), Tetranthera calophylla (Miquel), Tetranthera laurifolia (Jacq ), Litsea Schifera (Pers.), Schifera glutinosa (Loui.) (3) Irvingia butter. Cochin China wax, or Cáycáy fat, the fat obtained from Irvingia Oliveri (Pierre), Irvingia malayana (Oliv), Irvingia Harmandiana (Pierre), Buchanania fastigiata (H Baillon) The Cay-cay tree-' candle tree '-is a tropical tree indigenous to Cochin China and Large quantities of the fat are used ves as candles The freshly prepared Cambodia by the natives as candles fat is employed for edible purposes (4) Kusu oil, contained in the fruit of the camphor-yielding tree, Cunamomum camphora (Nees), Lauracea. From its chemical characteristics, the conclusion may be drawn that it consists of pure laurin with a few per cent of olem

DIKETONES v KETONES

DIKETOSUCCINIC ACID v KETONES

DILATOMETER. A term originally used to denote an apparatus for measuring the thermal expansion of liquids Subsequently given by Silbermann to an instrument for determining the strength of aqueous alcohol

DÍLITURIC ÁCID, núrobarbituric acid, nitro-malonylurea, CO NH CO CH NO<sub>2</sub>,3H<sub>2</sub>O, obtained by treating alloxantin first with hydrochloric acid and subsequently with nitric acid (Schlieper, Annalen, 56, 24); by warming hydurilic acid with nitric acid (Baeyer, *ibid* 127, 211), by nitrating barbituic acid (Baeyer, *ibid* 127, 211), by nitrating barbituic acid (Baeyer, *ibid* 130, 140), by oxidising violuric acid (nitrosobarbituric acid) (Ceresole, Ber. 16, 1134); or by oxidising with nitric acid the compound C<sub>16</sub>H<sub>26</sub>O<sub>11</sub>N<sub>12</sub>S<sub>2</sub> obtained by condensing *iso*dialuric acid with thiocarbamide (v. Vogel, Annalen, 315, 259; Bartling, *ibid*. 339, 27).

Dilituric acid crystallises in colourless quadratic prisms or places, is sparingly soluble in cold, more readily so in hot water, is sparingly soluble in alcohol and insoluble in ether. It

forms chloropierin when warmed with bleaching powder, is reduced by hydriodic acid to uramil (aminobarbitunic acid), and suffers partial reduction when heated with glycerol, yielding violuria acid. Dilitunic acid is tribasic, but most of the salts are acidic, and contain only one equivalent of the metal, they are sparingly soluble in water, and are not generally decomposed by mineral acids. The ammonium salt

the sodium salt  $C_4H_2N_3O_5N_3, 4H_2O$ , potassium salt  $C_4H_2KO_5N_3$ , calcium salt

$$Ca(C_4H_2O_5N_3)_2,4H_2O$$

the barium compound  $BaCl(C_4H_3O_5N_3),H_2O_5$ ferrous salt  $\operatorname{Fe}(\mathrm{C_4H_2O_5N_3})_2,8\mathrm{H_2O}$ , and the silver salt  $C_4H_2AgO_5N_3, 2H_2O$ , are colourless, the ferric salt  $Fe(C_4H_2O_5N_3)_3, 0H_2O$  is bright yellow, the copper salt  $Cu(C_4H_2O_5N_3)_3, H_2O$  is greenish, the dipolassium salt  $C_4H_5O_5N_3$  is bright-yellow (Schlieper, lc), and the triargentic salt  $C_4Ag_3O_5N_3$  is lemon-yellow (Baeyer, Annalen, 130, 140) According to Holleman (Rec trav chim 1897, 16, 162), dilituric acid and its homologues are aliphatic nitro- compounds in which the iso- form (cf. Hantzsch and Schultze, Ber. 1896, 29, 699, 2251, Hantzsch and Veit, ibid 1899, 32, 607) is the stable modification. The aqueous solution of dilituric acid is faintly yellow and decomposes carbonates, the molecular conductivity for I gram-molecule dissolved in v litres of water is 323 4 (v=32), 341 4 (v=512), ( $\mu_{\infty}$  should be 356), the acid is therefore almost completely dissociated in aqueous solution; the addition of one equivalent of a base does not appreciably change the colour of the solution, but on adding a second equivalent, the solution assumes the deep-yellow colour characteristic of the aqueous solutions of the salts of this acid Assuming that the formula CO < N = C(OH) > C N OH re-

presents the constitution of the acid, Holleman states that the anion

is colourless and slowly hydrolyses into the yellow anion CO NH CO ... C N O-.

Methyldilituric acid, 1 · methyl · 5 · nutrobarbituric acid CO NH CO CH NO<sub>2</sub>,1 5H<sub>2</sub>O,

prepared by oxidising methylvioluric acid with concentrated nitric acid (Andreasch, Monatsh 1900, 21, 281), crystallises from aqueous solution, and the anhydrous acid is colourless; it melts and decomposes at 143°, is readily soluble in hot water, less so in cold water, alcohol, or acctone, and sparingly soluble in ether. The aqueous solution of the acid is intensely yellow, and decomposes carbonates, the aqueous solutions of the salts are also yellow, but the solid sales are colourless or pale-yellow and sparingly soluble, the potassium salt  $C_5H_4\mathrm{KO}_8\mathrm{N}_3$ , the sodium salt  $C_5H_4\mathrm{NO}_6\mathrm{N}_3$ . He odium salt  $C_5H_4\mathrm{NO}_6\mathrm{N}_3$ , are colourless, the barium salt  $(C_5H_4\mathrm{O}_8\mathrm{N}_3)_2\mathrm{Ba}_3\mathrm{H}_4\mathrm{O}$ , the

cadmium salt (C5H4N3O5)2Cd,2H2O, are paleyellow, the strontium salt  $(C_5H_4O_5N_3)_2Sr,3H_2O$ is colourless, but becomes yellow on losing its water of crystallisation Methyldilituric acid is stable with bases, it reacts readily with chlorine to form methylchlordilituric acid

$$CO \stackrel{NMo}{<} \stackrel{CO}{<} CCI(NO_2)$$

m p 121°; the corresponding bromo-compound melts and decomposes at 138°-139°

Dimethyldilituric acid, 13 - dimethyl - 5 nitro barbituric acid CO NMe CO CH NO. obtained by oxidising dimethylvioluric acid with concentrated nitric acid (Techow, Ber 27, 3085, Andreasch, Monatsh 16, 26), forms colourless microscopic crystals from acctone; dissolves in 5.98 parts of water at 14°, is sparingly soluble in alcohol, melts at 152° and changes into dimethylallovan (Holleman, Rec trav chim 16, 162), and is reduced by hydriodic acid to dimethyluramil

Dimethyldilituric acid is intensely yellow in aqueous solution, and this is due to the presence of the coloured anion of the isonitro- form CO NMe CO C NO OH, for on adding con centrated nitric acid, the colour almost entirely disappears, further, the molecular conductivity of the acid for I gram-molecule dissolved in v litres of water is 3133 (v=32), and 3376(r=1024),  $(\mu_{\infty}$  should be 352), so that in each solution the acid is almost completely ionised (Holleman, Rec trav chim 16, 162) salts of dimethyldilituric acid are coloured, the salts of dimenylamouric acid are coloured, one sodium salt  $({}^c_6H_6N\alpha O_5N_3, H_2O)$  is yellow, the potassium salt  $({}^c_8H_6N\alpha O_5N_3)$  is greenish-yellow, the magnesium salt  $(({}^c_8H_6O_5N_3)_2Mg, 4H_2O)$  is yellowish-green, the calcium salt  $(({}^c_8H_6O_5N_3)_2Ca)$  is bluish green, the strontium salt  $(({}^c_8H_6O_5N_3)_2Ca)$ is pale-green, the barrum salt

(C6H6O5N3)2Ba,2H2O

is rose-red, and the silver salt C<sub>8</sub>H<sub>8</sub>AgO<sub>8</sub>N<sub>3</sub> is greyish-violet (Andreash, lc, and Monatsh 21, 295)

Dimethylchlordilituric acid

prepared by passing chlorine into an aqueous solution of dimethyldilitume acid, melts and decomposes at 150°, the corresponding bromoderivative melts partially at 152° (Andreasch, lc) Dimethyldilituric acid is readily decomposed by the action of bases forming water, carbon dioxide, and dimethylnitromalonamide

M A W BP The DILL FRUIT. Anethr fructus, BP dried ripe fruit of Peucedanum graveolens (Benth and Hook)

DILL OIL v OILS, ESSENTIAL

DIMETHYLALLOXAN v ALLOXAN

p-DIMETHYLAMINOBENZOPHENONE KETONES

**DIMETHYLAMINONAPHTHAPHENAZINE** v Azines

#### >N(CH₃)₃. DIMETHYLANILINE

Made by heating a mixture of aniline and methyl alcohol containing sulphuric acid in an autoclave. For details of mode of manufacture,

see Walter, Chem Zeit 1910, 34, 641, et seg; and Cain's Intermediate Products for Dyes Instead of sulphuric acid iodine may be employed (Knoll & Co, D R P 250236) Or a mixture of aniline and milk of lime and methyl chloride may be heated under pressure (Grandmougin, Rev prod (him 1917, 20, 68)

Dimethylaniline bods at 192 6°/738 mm; mp 2 07°. spgr 0 96 at 15° (see Amines)

DIMETHYLDIAMINODI - o - TOLYLME-THANE

Obtained by treating methyl-o-toluidine, or a mixture of mono- and dimethyl-o-toluidine with formaldehyde (40 pc solution) and hydrogen chloride, diluting with water, adding sodium and steam distilling to expel the excess of the methylo-toluidine On cooling the base crystallises, and may be purified by crystallisation from alcohol (Bad Aml and Soda Fabrik, D R P 67478), mp 87°, bp 255°/9 mm

DIMETHYLDIAMINOTOLUPHENAZINE v

Azines

2 : 2'-**DIMETHYL**-1 1'-DIANTHRAQUIN-ONYL

Obtained by diazotising 1-amino-2-methylanthraquinone dissolved in sulphuric acid, pouring the product on ice, separating the diazonium sulphate, and treating it with acetic anhydride and copper powder. The precipitate thus formed is mixed with water to remove acetic anhydride, and with dilute nitiic acid to free it from copper Can also be prepared by treating 1-chloromethylanthraquinone with copper powder (Bad. Anılın and Soda Fabrik Used in D R P 180157), mp 366°-367° making Indanthrene Golden yellow v. Pyranthrene, art Indanthrene

DIMETHYLETHYL ACETIC ACID v CA-PROIC ACID

**DIMETHYLGLYOXIME** ( $\beta \gamma$ -droximinobutane) CH<sub>3</sub> C . N OH was originally obtained by CH3 C NOH

Schramm, in 1883, by acting upon methyloximino-ethyl ketone with aqueous hydroxylamine hydrochloride (Ber 16, 180) It is readily prepared in quantity by the following method (Gandurin, J pr Chem 1908, 77, [u] 414) --

850 c c of commercial methyl ethyl ketone are mixed with 30 cc of hydrochloric acid of sp gr 1 19, and 1 litre of amyl nitrite is allowed to run in slowly with vigorous stirring This takes 11 hours, and the temperature should be maintained between 40°-50° When the reaction is over, 800 grams of ic are added, followed by 378 grams of sodium hydroxide in 800 cc. of water, and the mixture is shaken for half an hour. After standing, the layer of amyl alcohol adding the oxime and acetate (Brunck, Zeitschis removed, and the aqueous liquid twice extracted with ether. 100 c.c of the alkaline liquid are treated with the calculated amount of advisable to work with solutions containing not hydroxylamine hydrochloride, and the dimethylglyoxime produced is collected and weighed. If the original volume of the alkaline liquid is  $v \in c$ , a grams the weight of dimethylglyoxime obtained from the 100 cc of liquid, then the amount of hydroxylamine hydrochloride in grams required for the remaining liquid is equal to 650(v-100)a/109500 This weight of hydroxylamine hydrochloride is dis-, solved in a little water, added to the alkaline liquid, and the whole warmed on the water-bath for an hour The liquid is acidified with acetic acid, and the dimethylglyoxime, which separates, is filtered, washed, and dried This product is sufficiently pure, but it may, if desired, be iecrystallised from alcohol Yield, 650-700 grams

(For details, see Adams and Kamm, J. Amer

Chem. Soc. 1918, 40, 1281)

(For other methods of preparation, v Biltz, Zeitsch anal Chem 1909, 48, 164, Tschugaeff, Ber 1905, 38, 2520)

Properties —Dimethylglyoxime forms colourless, glistening needles, and is insoluble in water, but soluble in alcohol It melts at 245°-246° (Meyer-Jacobsen, I (ii ) 826, 1913) Its dibenzovi derivative forms rhombic plates, melting at 223° (Diels and Stein, Ber 1907, 40, 1629).

On oxidation with potassium ferrieyanide or nitrogen peroxide it yields furovane (Scholl, Ber 1890, 23, 3498), this is reduced to the dioxime again when treated with zinc and acetic acid (Rimini, Gazz chim ital 1895, 25, When heated with aqueous amfn | 266) monia at 170°-180° it yields the anhydride dimethylfurazan (Wolff, Ber 1895, 28, 69)

Application in Analysis -- Dimethylglyoxime is now an important analytical reagent, owing to its property of forming a bright scarlet insoluble nickel derivative It thus becomes possible to estimate nickel simply and accurately, in the presence of iron, aluminium, chromium, cobalt, zinc, or manganese, all of which form soluble derivatives In addition, an excellent qualitative test for nickel in the presence of cobalt is supplied. For the latter purpose, a pinch of the oxime is added to the solution containing excess of ammonia or sodium acetate, and the liquid boiled. A scarlet precipitate is obtained even when there is present only one part of nickel in two million parts of solution (Tschugaeff, Ber 1905, 38, 2520; Compt rend 1907, 145, 679, Klaut, Zeitsch angew Chem 1906, 19, 1793) 01 milligram of nickel can be detected in the presence of 5000 times as much cobalt

For the quantitative estimation of nickel, the acid solution is heated nearly to boiling, a slight excess of 1 p c alcoholic dimethylglyoxime is added, and then a small excess of either am-The precipitate is monia or sodium acetate filtered while hot on to a Gooch crucible, washed with hot water, and dried at 110°-120° Its formula is (CH<sub>3</sub> C NO)<sub>2</sub>N<sub>1</sub>+ (CH<sub>3</sub> C NOH)<sub>2</sub>, and it contains 20 31 pc of nickel, it has no definite melting-point, but sublimes at 250° Ammonium salts do not interfere; if the acetate method is employed, the original solution should be nearly neutralised with ammonia before

Vol. II.-T.

angew. Chem. 1907, 20, 1844). As the precipitate of nickel glyoxime is very voluminous, it is more than about 0 l gram nickel, owing to the difficulty of filtration, and again in order to avoid waste of material, the cobalt, if present, should not exceed 0.1 gram, otherwise a large excess of ammonium hydroxide and dimethylglyoxime is necessary in order to prevent its precipitation

By either of the above methods, cobalt and zine may be separated from nickel, but about twice the theoretical amount of reagent is Manganese, of course, can only be necessary separated by the acetate method. Iron, aluminium, and chromium may be separated by the ammonia method, if tartaire acid be previously added (and, in addition, ammonium chloride in the case of chromium) Iron may also be separated m acctic acid solution, provided it is in the ferrous state (Brunck, Zeitsch angew Chem 1907, 20, 1845, Prettner, Chem Zeit 1909, 33, 396, 411; Ivanicki, Chem Zeit Rep 1908, 32, 644, Rhead, Analyst, 1910, 35, 97, v also Analysis)

The oxime is sufficiently expensive to make its recovery desirable This may be effected by dissolving the nickel derivative in aqueous potassium cyanide and acidifying with acetic acid The precipitated dimethylglyoxime is purified by recrystallisation from hot alcohol, after decoloursation with charcoal (v Ketones).

DIMETHYLMALONIC ACID v GLUTARIC ACID

DIMETHYLOXALIC ACID v Hydroxy-BUTYRIC ACIDS

aa DIMETHYLPROPIONIC ACID v VALERIC ACID

DIMETHYLPYRROLE v Bone on. DIMETHYLSAFRANINES & AZINES

DIMETHYLTRIAZINE, Diazoaminomethane

v Diazo- compounds

DINAPHTHAZINES v AZINES DINITROANTHRAQUINONES.

1.5-Dinitroanthraquinone (in p. 330')

is made, together with the 1-8 isomeride, by adding sodium initiate to anthraquinone dissolved in sulphune acid. By extraction with alcohol or acctone the 1 8 dinitroanthraquine (m p 312°) NO. CO NO.

is removed

Used in the manufacture of anthracene See Alizarin and Allied colouring blues MATTERS

DINITROBENZENEDIAZO-OXIDE 1 DIAZO-COMPOUNDS

DINITROBENZENES v. BENZENE AND ITS HOMOLOGUES

4-DINITRO-4'-HYDROXYDIPHENYL-AMINE

Made by boiling together in water containing chalk a mixture of 4-chloro-1:3 dinitrobenzene and p-aminophenol in molecular proportions, mp. 190° Used in the manufacture of sulphide dyes (q v)

DINITRONAPHTHALENES v NAPHTHALENE.

DINITRONAPHTHALENES v NAPHTHALENE.
DINITROPHENOLS v NITROPHENOLS AND
THEIR HOMOLOGUES

# DINITROSTILBENEDISULPONHIC ACID

$$NO_2$$
 CH  $CH$   $O_2$   $O_3H$   $O_4$ 

Obtained by waiming a solution of sodium p-nitrotoluene sulphonate containing sodium hydroxide with sodium hypochlorite when sodium dinitrostilbenedisulphonate separates on cooling. The free acid is obtained by treating the sodium salt with hydrochloric acid. Pale yellow needles

**DINITROTOLUENES** v BENZENE AND ITS HOMOLOGUES, NITRATION OF.

# DIOCTYLMALONIC ACID

 $(C_8H_{17})_2C(CO_2H)_2$ 

Prepared from diethylmalonic ester, sodium methoxide and octyl iodide, and subsequent decomposition of the ester thus formed with caustic potash (Conrad and Bischoff, Annalon, 204, 163), m p 75°.

**DIOFORM.** Trade name for symm 1-2-di-CHCl

chlorethylene (acetylene dichloride) || CHCl

(q.v) Suggested as an anæsthetic

**DIOGEN.** Trade name for sodium aminonaphthol-disulphonate.

DIOGENAL. Trade name for dibromo-propyl diethylbarbituric acid

DIONINE. Ethyl morphine hydrochloride,

v OPIUM and SYNTHETIC DRUGS

DIOPSIDE. A double salt meta-silicate of calcium and magnesium in which a part of the magnesium is usually replaced isomorphously by an equivalent amount of ferrous oxide, the formula being Ca(Mg,Fe)(SiO<sub>3</sub>)<sub>2</sub>. It is a member of the monoclinic series of the pyroxene group, and is of importance as a rock-forming minicial, while occasionally, when transparent and of a rich bottle-green colour, it is employed as a gem-stone. The depth of colour, sp gr (3 2-3 3), and other physical characters vary with the percentage of 4ron present, which in material of gem quality, such as the crystals from Ala in Piedmont, and the Zilleithal in the Tyrol, amounts to about 3-4 pc FeO. The hardness is 6.

DIOPTASE. Hydrous silicate of copper, H<sub>2</sub>CuSiO<sub>4</sub>, crystallising in the rhombohedral system with parallel-faced hemihedrism. The water is expelled only at a red heat, and the formula has consequently been variously interpreted as an acid ortho-silicate, or as a basic meta-silicate. F. Zambonin (1908) concludes from dehydration experiments that the formula is CuSiO<sub>3</sub> H<sub>2</sub>O, with the water held in solid solution in the meta-silicate. The mineral is of a characteristic emerald-green colour (hence the name emerald-copper; Ger, Kupfer-Schmaragd), and transparent crystals have occasionally been cut as gem-stones. For this purpose, however, the hardness (H = 5) is rathalous (spiralling) and the limestone hill of Altyn-Tübe in

the Kirghiz Steppes, Siberia, and in the copper mines near Mindouli in the French Congo.

An allied mineral, planchette (of A Lacroix, 1908,=shattuckite of W. T. Schaller, 1915), occurs as blue, fibrous concretions associated with dioptase in the French Congo. This has the composition 2CuSiO<sub>3</sub>·H<sub>2</sub>O, representing a partly dehydrated dioptase. L. J. S.

DIORITE. An igneous rock composed essentially of plagioclase felspar (oligoclase to labrado-11te) and hornblende, and with a coarse- to medium-grained crystalline-granular texture. It belongs, with granite, to the series of plutonic 10cks, and is of intermediate composition (SiO, 52-60 pc), falling between syenite and gabbio Its volcanic equivalent is andesite Other minerals may also be present as essential constituents of the lock, and accordingly the following varieties are distinguished. blende-diorite (diorite proper), augite-diorite, enstatite-diorite, mica-diorite (the mica here being biotite), quartz-mica-diorite, quartz-diorite (or tonalite) The colour is usually dark grey or greenish-black, and for this reason the rock is included under the general term 'greenstone' The sp gr ranges from 2 8 to 3 1; the porosity is low, 0.25 p.c., and the crushing strength 1640 to 1830 tons per square foot Under 1 is given Biogger's average of sixteen analyses of typical diorites, analysis II is of dionite from Strathspey, Scotland; III. of quartz-diorite from Spanish Peak, Plumas Co, California (including BaO) 0.04, CO $_2$  0.20,  $P_2O_5$  0.25,  $SO_3$  0.03), IV homblende-augite-diorite from Inchnadampf, Sutherlandshire (including CO<sub>2</sub> 0 54).

	1.	11	111	1V
$SiO_2$	. 56 52	53 22	59.68	$52\ 47$
T10,	. 0 25		0.65	
$Al_2O_3$	16 31	16 84	17.09	12 15
$Fe_{3}O_{3}$	4 28		2.85	347
FeO	. 5 92	9.27	2.75	523
('a()	. 694	8 53	6.62	9.71
MgO	$4\ 32$	6.48	3.54	9.94
K,0	1 44	1.39	1 31	2.26
Na <sub>2</sub> O	3 43	3.54	3 87	2 81
Η <sub>Σ</sub> Ο	. 103	0.95	1 15	1 62
	100 34	$\bar{100} \ \bar{22}$	100 03	100 20

Diorites are of wide distribution and are very common in association with granitic rocks in the southern uplands and highlands of Scotland. They are extensively quarried for road-metal and paving-setts in the Channel Islands, Guernsey, Jersey, and Alderney, and at Brazil Wood near Mount Sorrel in Leiestershire. Owing to the difficulty of working the rock and its dull colour, it is not much used as a building stone, except locally. Certain coarse-grained varieties are, however, polished for ornamental purposes. For example, a variety from Corsica, known as napoleonite, coisite, or ball diorite, shows a spheroidal structure with alternating concentric layers of light felspar and dark hornblende.

L. J. S.

DIOSCORINE ('13H19O2N, greensh-yellow plates, mp 435°, occurs in Diogorea hirsuta (Blume) tubers of Japan Ritter and poisonous (Gorter, Rec trav. chim. 1911, 30, 161) Has an action like picrotoxin

DIOSPHENOL, Buchu camphor v. Buchu DIOXINE. Gambine H. 1-nitroso-2: 7dihydroxynaphthalene, v. QUINONEOXIME DYES. DIPHENYL.

DIOXOREA. A genus of plants of the Dioxoreaceous order, the tubers of which are known as yams, and are largely eaten in the West Indies and other tropical countries D. batatas (Deene) is the plant producing tubers known as 'sweet-potatoes,' which are cooked and eaten as a substitute for potatoes

DIOXYGEN. Trade name for a 3 pc

solution of hydrogen peroxide

**DIPHENAL**, Duminohydroxydiphenol, used as a photographic developer

DIPHENAZINE v Azīnes.

o-DIPHENETIDINE v DIPHENYL.

**DIPHENYL.** This hydrocarbon, which has the molecular formula  $C_{12}H_{10}$ , is structurally composed of two benzene nuclei joined together by a bond linking two carbon atoms, one in each ring. The carbon atoms are numbered for the purposes of nomenclature in the following way.—

Although of minor importance, diphenyl is nevertheless the parent substance of a large number of important bases which have found application in the preparation of the so-called substantive cotton dyes, that is to say, of colouring matters which will affix themselves to the vegetable fibres without the aid of a mordant. These bases are, however, not derived directly from the hydrocarbon, but are prepared by an inducet method which will be described later.

Diphenyl was originally obtained by Fittig (Annalen, 121, 363) from an ethereal solution of bromobenzene by the action of metallic sodium  $C_6H_5Br+BrC_6H_5+2Na \rightarrow C_6H_5-C_6H_5+2NaBr$  but it is more conveniently prepared by the method discovered by Berthelot (Ann Chim Phys [4] 9, 453, [4] 12, 185), who obtained it by passing the vapour of benzene through a redhot tube

$$('_{6}H_{5}H + H('_{6}H_{5} \rightarrow ('_{6}H_{5} - ('_{6}H_{5} + H_{2}))))$$

The following procedure is recommended (Schultz, Ber 9, 547, Hubner, Annalen, 209, 339; Smith and Lewcock, Chem Soc Trans 1912, 101, 1453; [apparatus] La Coste and Sorger, (Annalen 230, 5) The vapour from a quantity of benzene (free from thiophen), contained in a large flask heated on the waterbath, or produced by dropping the liquid into a copper flask heated to 150°, is allowed to pass through an iron tube, filled with pumice, which is heated to about 720° in a combustion furnace, the lower end of the iron tube is attached to a condenser and receiver, in which the products are collected. When all the benzene has distilled over, the distillate is again placed in the flask, and the vapour passed anew through the red-hot tube, or the process may be made continuous. The ultimate product is then fractionally distilled, and the portion boiling between 250° and 260°, which solidifies on cooling, is purified by recrystallisation from alcohol The yield is about 50-60 p.c. The pyrogenic preparation can also be effected by means of an incandescent filament of carbon, platnum, or nickel (Löb, Zeitsch. Elektrochem. 1902, 8, 777) Diphenyl separates from alcohol as large, lustrous lamine which melt at 70 5°. It boils at 254° (corr.), and is insoluble in water. The

hydrocarbon, when oxidised by chronic acid passes into benzoic acid.

531

Bennett and Turner (Chem. Soc Trans. 1914, 105, 1057) found that chromic chloride reacted quantitatively with magnesium phenyl bromide to form diphenyl Krizewsky and Turner (Chem Soc Trans 1919, 115, 559) have observed that anhydrous cupric chloride behaves similarly, yields of about 85 p c of the theoretical amount of diphenyl being thus obtained

Derivatives of Diphenyl and their Orientation -The more important derivatives of diphenyl are not prepared directly from the hydrocarbon, but from certain nitio- delivatives of benzene in a manner to be described later The hydrocarbon yields, however, halogen, nitro- and sulphonic acid derivatives analogous to those prepared from benzene, substitution taking place in either one or both rings according to the conditions of the experiment The method of orientation adopted for these compounds may be briefly described as follows (Schultz, Annalen, 174, 201, Schmidt and Schultz, ibid 207, 320). When diphenyl is treated with bromine, it yields a monobromodiphenyl, a substance which, on oxidation, is converted into p-bromobenzoic acid, thus  $\cdot$ 

$$BI \leftarrow \rightarrow Br \leftarrow CO_2H$$

It follows, therefore, that in bromodiphenyl the bromine atom must have entered in the paraposition with respect to the linking carbon atom. When bromodiphenyl is treated with intrie acid, it is converted into a monobromomonomitrodiphenyl, a substance which, on oxidation, yields a mixture of p-bromobenzous acid and p-introbenzous acid. It is therefore apparent that the formula of this bromomitroderivative must be.

The same compound is also obtained from dinitrodiphenyl by first reducing this to intro-aminodiphenyl (mp 198°) and then replacing the amino-group by bromine by means of the diazo-reaction. Hence dinitrodiphenyl and its reduction product are di-para-derivatives. Since, however, introaminodiphenyl, on reduction, furnishes the diamino-derivative, benzidine, the latter is also a di-para-compound, of 4'-diaminodiphenyl, the preparation of which is described latter.

The transformation of the simplest atomatic hydrocarbon benzene into diphenyl by the action of heat has already been mentioned. This transformation can, however, be effected much more readily by the oxidation of certain derivatives of benzene, and especially those which, like the phenols, have a reactive hydrogen in the para-position with respect to the substituting group. Thus phenol itself and also resorcin readily pass into derivatives of diphenyl when fused with potash (Barth, Annalen, 156, 95). With hydroquinone, the transformation is still more readily effected and the action of the mildest oxidising agents is sufficient for the purpose (Nietzki and Bernard, Ber. 31, 1334).

The primary bases of the benzene series, such as aniline, do not lend themselves to the production of derivatives of diphenyl, but the territary bases readily do so; thus dimethylaniline yields tetramethylbenzidine when oxidised (Michler and Pattinson, Ber. 14, 2162 and 2164)

DIPHENYL. 532

It has also been found that certain azoelerivatives of benzene can be transformed by oxidation into tetrazo- derivatives of diphenyl, and in this way Congo red can be formed by benzeneazonaphthionic acid with manganese dioxide and sulphuric acid.

$$\begin{array}{c} C_{6}H_{5}N\!=\!N\!-\!C_{10}H_{5}(NH_{2})SO_{3}H \\ +O \\ C_{6}H_{5}N\!=\!N\!-\!C_{10}H_{5}(NH_{2})SO_{3}H \\ C_{6}H_{4}\!-\!N\!=\!N\!-\!C_{10}H_{5}(NH_{2})SO_{3}H \\ \to \downarrow \\ C_{2}H_{4}\!-\!N\!=\!N\!-\!C_{10}H_{5}(NH_{2})SO_{4}H \end{array}$$

Derivatives of diphenyl can also be prepared from certain diazonium salts of the benzene Thus Kuhling has shown (Ber 28, 41, 29, 165) that when an isodiazo- salt is treated with an aromatic hydrocarbon in the presence of an acid chloride, the following reaction ensues :-

$$\begin{array}{c} \mathrm{NO_2} \xrightarrow{\mathrm{C_6H_4N_2OK+C_6H_6+CH_3COCl}} \\ \xrightarrow{\hspace{0.5cm} \rightarrow} \mathrm{NO_2C_6H_4-C_6H_5+N_2+KCl+CH_3CO_2H} \end{array}$$

This reaction has been simplified by Bamberger (Ber 28, 403, 29, 272 and 465), who uses free isodiazobenzene hydroxide, and shows that it passes, in the presence of benzene, without any condensing agent, into a derivative of diphenyl Many derivatives of diphenyl can also be formed by the method of Ullmann, in which halogen derivatives of benzene are treated with finely divided

copper (Ber 43, 2082, Annalen, 1904, 332, 38)

Benzidine and the 'Benzidine Rearrangement'—The benzidine rearrangement, which has been widely applied to the preparation of a large number of important bases used in the production of the substantive cotton dyes, is the name given to the change which an aromatic hydrazo- compound undergoes when treated with acids or ceitain acid reagents. The change is effected by the wandering of the hydrogen atom which is in the para-position in each ring on to the nitrogen atom attached to that ring, at the same time, the link between the nitrogen atoms is broken, and the two aromatic nuclei are joined together It is apparent, therefore, that only those hydrazo- denvatives of benzene which have their para- position free will undergo this transformation

$$\begin{array}{c} \stackrel{\longleftarrow}{\underset{NH-C_6H_4H}{\longleftarrow}} \\ \downarrow \\ NH-C_6H_4H \end{array} \rightarrow \begin{array}{c} NH_2-C_6H_4 \\ NH_2-C_6H_4 \end{array}$$

The history of the discovery of this rearrangement is briefly as follows. Zinin (J. pr. Chem 36, 93), in the first instance, showed that azobenzene  $C_6H_5-N=N-C_6H_5$ , when treated with alcoholic ammonium sulphide, passed into a crystalline compound which, when dissolved in alcohol and boiled with sulphuric acid, gave the sparingly soluble sulphate of a base. This base sparingly soluble sulphate of a base he called benzidine

It soon became apparent that the substance responsible for the formation of benzidine must be an intermediate product formed by the action of the reducing agent on azobenzene This view was confirmed experimentally by A. W Hofmann (Jahresbericht, 1863, 424), who prepared the intermediate product, hydrazobenzche, and for I that it passed with remarkable case muo benzidine when treated with mineral acids.

Finally, Fittig (Annalen, 124, 280) showed that benzidine was identical with the base formed by the reduction of 4-4'-dinitrodiphenyl, and hence established the constitution of the base as 4-4'-diaminodiphenyl.

Manufacture of Benzidine - A jacketed iron pan fitted with an agitator, reflux condenser, &c., is charged with 100 kilos of nitrobenzene and 50 kilos of alcohol This mixture is heated to boiling, and 150-160 kilos of zinc dust are added. The stirring must be very efficient during the next operation, which requires 5-6 hours, of adding gradually a mixture of 100 kilos of alcohol and 40 kilos of addium hydroxide solution (36°-40°B). The reaction is vigorous and is regulated by the rate of addition of the alkalı and, if necessary, by cooling. When all the alkalı has been added the mixture is stirred until it is of a uniformly grey colour, and to obtain this more zinc may have to be introduced. The alcohol is then distilled off by means of steam, and the residue is thrown on a fine sieve which retains the hydrazobenzene and allows the zine dust, zine oxide, &c, to pass through Another method of working up consists in diluting the reduction mixture with water, carefully neutralising with hydrochloric acid, and filtering off the hydrazobenzene

The latter is transformed into benzidine by mixing it with ice and acidifying with concentrated hydrochloric acid. After some time water is added and steam introduced so as to dissolve the benzidine hydrochloride, the solution is filtered, and the benzidine sulphate precipitated by adding sodium sulphate solution This is filtered off, washed, and the base obtained by boiling with sodium hydroxide solution, cooling, collecting, and distilling in a vacuum The yield is about 80 p c. of the theoretical.

The above process can also be carried out without the use of alcohol, or solvent naphtha

can be employed instead.

The electrolytic reduction of nitrobenzene to hydrazobenzene is the subject of a number of patents, but it would not appear that this method is yet applied on the large scale (cf Lob, Ber 1900, 33, 2329)

The production of benzidine from azobenzene is easily effected by an electrolytic method with titanium salts or vanadium trioxide used as catalysts The azobenzene is suspended in 30 p c sulphuric acid containing 10 p c titanous sulphate; a lead cathode is used with a current of 3-4 amp. per square dem at 3-4 volts (D R PP 168273 and 172654)

The presence of small quantities of hydriodic acid facilitates the reduction of azobenzene to benzidine by means of sulphurous acid (D. R. P. 172569)

Benzidine, when pure, forms colourless lustrous laminæ, which melt at 127 5°-128°; it boils at 400°-401° (740 mm) The base is characterised as a para-derivative by the formation of quinone when it is oxidised by manganese dioxide and sulphuric acid It can be identified by a number of colour rechlorine, bromine, potassium permanactions ganate, potassium ferricyanide, and chromic acid, producing characteristic shades. most characteristic salts of benzidine are the sulphate  $C_{12}H_{12}N_2, H_2SO_4$ , and the chromate  $C_{12}H_{12}N_2, H_1CrO_4$  Both salts are sparingly soluble in alcohol and in water.

When treated with nitrous acid, benzidine hydrochloride is transformed into a tetrazonium salt of diphenyl

 $C_6H_4N_2Cl$   $C_6H_4N_2Cl$ 

which reacts readily with the usual second components of the azo-colouring matters, forming azo-compounds, the solutions of which possess the property of dyeing the vegetable fibres without the aid of a mordant. The number of colouring matters of this kind prepared from benzidine is largely increased owing to the insulable character of the diazonium salt.

$$C_6H_4N_2X$$
 $C_6H_4N_2CI$ 

in which one molecule of a second component, XH, has combined with the tetrazonium salt derived from benzidine. These diazonium salts, when combined with some other second component, YH, yield mixed azo-colouring matters of the formula.

$$\begin{array}{c} C_6H_4N_2X \\ \dagger \\ C_6H_4N_2Y \end{array}$$

Benzidine in the presence of hydrogen peroxide is used as a test for blood, with which it gives a blue colouration. The test solution is made by taking 10 drops of a concentrated solution of benzidine in glacial acetic acid, and adding 2 c c of a 3 p c solution of H<sub>2</sub>O<sub>2</sub> Sensitiveness—I part in 200,000 (Utz, Chem Zeit 1907, 31, 737, Schumm, Pharm Zeit 1907, 52, 604, McWeeney, Sci. Proc. Roy Dubl. Soc. 1909, 12, 216, Bordas, Compt. rend 1910, 150, 562, Gigli, Boll chim, farm. 1910, 49, 955, cf. O. and R. Adler, Zeitsch. physiol Chem. 1904, 41, 59.)

Homologues and Derivatives of Benzidine used in the Preparation of the Sushtantive Cotton Dijes—Although a great many of the derivatives of benzidine yield azo- colouring matters which are substantive to the vegetable fibres, yet this property is largely dependent on the position of the groups substituting the hydrogen atoms of the diphenyl molecule—Thus it has been found that when the substituting groups are in the meta-position to the coupling bond, as in

$$\begin{array}{c|c} \operatorname{CH_3} & \operatorname{CH_3} \\ \operatorname{NH_2} & & \operatorname{NH_2} \\ \\ \operatorname{OCH_3} & \operatorname{OCH_3} \\ \end{array}$$
 and 
$$\operatorname{NH_2} & & \operatorname{NH_2} \\ \\ \bullet \cdot \operatorname{Dianisidine} \\ \\ \bullet \cdot \operatorname{Dianisidine} \\ \end{array}$$

the azo- colours formed are substantive to cotton. If, however, groups are in the orthoposition, as in the contract of the c

the property of the azo-colours of dyeing unmordanted vegetable fibre is lost. The orthooderivatives nevertheless yield substantive cotton dyes if ring formation takes place between the substituting groups in those positions, thus

Benzidine sulphone

yield azo- dyes substantive to cotton.

Benzidine-o-disulphonic acid is prepared by heating benzidine sulphate with 2 parts of sulphuric acid for 36-48 hours at 210° (Ber 1889, 22, 2464, Eng Pat 1074, 1884, D R P. 27954)

o-Dichlorobenzidine is prepared from ochloronitrobenzene, and is also produced by chlorinating diacetylbenzidine (Eng. Pat. 25725, 1896; Fr. Pat. 265155, D. R. P. 94410, U.S. Pat. 625174). It melts at 133°

m-Nitrobenzidine, obtained by treating a solution of benzidine in sulphuric acid with one equivalent of intile acid or potassium nitrate (Tauber, Ber. 1890, 23, 796, Eng. Pat. 13475, 1892; Fr. Pat. 223176, D.R. P. 72867, U.S. Pat. 493583), melts at. 143°; when two equivalents of the intrating agent are used, m-dinitrobenzidine, in p. 214°, is produced. Monointration of diacetylbenzidine or phthalylbenzidine, and subsequent hydrolysis of the product, furnishes o-nitrobenzidine, in p. 190° (Koller, Ber. 1904, 37, 2883, Fr. Pat. 203468), whilst two different o-dinitrobenzidines, in p. 275° and 233°, are obtained by the dinitration of diacetylbenzidine and diphthalylbenzidine respectively (Trans. Chem. Soc. 1912, 101, 2298)

The following compounds, prepared from the corresponding nitro-derivatives of benzene, are of importance —

o-Toldine, 4 - 4' - diamino - 3 - 3' - dimethyl diphenyl  $\mathrm{NH}_2$  ( ${}^{}_{6}\mathrm{H}_3(\mathrm{CH}_3)$ —( ${}^{}_{6}\mathrm{H}_3(\mathrm{CH}_3)\mathrm{MH}_2$  from o-nitrotoluene, forms glatening plates, melting at 129° It is spaningly soluble in water, but readily soluble in alcohol and in ether. The sulphate is spaningly soluble in water

o-Diamsidine, 4-4'-diamino-3-3'-dimethoxy-diphenyl NH<sub>2</sub> ('aH<sub>3</sub>(O('H<sub>3</sub>)—('aH<sub>4</sub>(O('H<sub>3</sub>)NH<sub>3</sub>, from o-nitroanisole, forms colourless leaflets, melting at 135° (For its manufacture, see Jansen, Zeitsch Farb Ind 1913, 12, 247)

o-Diphenetidine, 4-4-diamino-3-3-diethoxy-diphenyl NH $_2$  C $_6$ H $_3$ (OC $_2$ H $_5$ )—C $_6$ H $_3$ (OC $_2$ H $_5$ )NH $_2$ , from o-nitrophenetole, forms needles or leaflets melting at 117° J F T

DIPHENYLAMINE was first obtained by Hofmann, in 1864, by distilling amline blue (Annalen, 132, 163), and in small quantity by distilling rosamline and leucaniline. De Laire, Girard, and Chapoteaut discovered an easier method of prepaining it by heating amiline with the hydrochloride at 210°-240° (Zeitsch Chem. 1866, 438, Compt. rend. 63, 91)

 $C_6H_5NH_2+C_6H_5NH_2HCl=NH(C_6H_5)_2+NH_4Cl$ 

This reaction is greatly facilitated by the use of certain catalysts. It may also be obtained by heating phenol with the double compound of

and line and zine chloride at 250°-260° (Merz and Weith, Ber. 13, 1298, 1880)  $^{\circ}$ 

$$C_6H_5NH_2+C_6H_5OH=(C_6H_5)_2NH+H_2O$$

and by heating aniline and diazoaminobenzene to 150° till the evolution of nitrogen ceases. In this last reaction, ortho- and para- aminodiphenyl are formed at the same time (Hirsch, D. R. P. 62309). Diphenylamine has also been prepared by the action of aniline on bromobenzene in presence of cuprous iodide (Goldberg, D. R. P. 187879, 1906). Aniline, when heated with ferric chloride, finely divided copper, and iodine, is converted into diphenylamine. Other catalysts can also be used (Rogers and de Nemours, U.S. Pat. 1314538).

Manufacture —Aniline (1 mol ) is heated with aniline hydrochloride (1 mol ) for 30-35 hours in an autoclave at 220°-230°. The product is extracted with hot dilute hydrochloric acid which dissolves the aniline hydrochloride and converts the diphenylamine into the free base, forming an oil which solidifies on cooling. The cake of diphenylamine is washed and then distilled alone or in a current of steam. Yield, 60-70 p.c. of the aniline employed. It is used in the manufacture of diphenylamine blue, diphenylamine orange, and aurantia (v. infra).

Properties — Monoclinic laminæ, mp 54°, b p 310°, almost insoluble in water, soluble in alcohol, benzene, and ether—It is feebly basic, and the salts are hydrolysed by an excess of

water

Reactions—The hydrogen of the imino-group is replaceable by alkyl or acyl radicals (Boseken, Roc trav chim 1912, 31, 350) With methyl rodide it gives a methyl derivative, with benzyl chloride, a benzyl derivative With formic or oxalic acid it yields formyldiphenylamine, mp 74°; with acetic acid or the acid chloride, an acetyl derivative, mp 103°, and with benzoyl chloride, a benzoyl derivative, mp 180° When heated with zinc chloride, the acyl derivatives lose water and form acridine derivatives. Formyldiphenylamine gives acilding

$$C_{6}H_{5} \stackrel{N}{\underset{CHO}{\mid}} C_{6}H_{5} = C_{6}H_{4} \stackrel{N}{\underset{CH}{\mid}} C_{6}H_{4} + H_{4}O$$

The acetyl compound yields methylacridine (Bernthsen and Bender, Ber 1883, 16, 767) Diphenylamine also gives acridine by heating with chloroform and zine chloride or aluminium chloride

$$(\mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{5}})_{\mathbf{2}}\mathrm{NH} + \mathrm{CHCl}_{\mathbf{3}} {=} \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{4}} {<} \bigvee_{\mathrm{CH}}^{\mathrm{N}} {<} \mathrm{C}_{\mathbf{6}}\mathrm{H}_{\mathbf{4}} {+} 3\mathrm{HCl}$$

(O Fischer and Korner, Ber 1886, 17, 101) With excess of oxalic acid, diphenylamine is converted into diphenylamine blue

$$3(C_{6}H_{5})_{2}NH+CO_{2}=C \underbrace{C_{6}H_{4} \ NH \ C_{6}H_{5}}_{C_{6}H_{4}=NC_{6}H_{5}}+2H_{2}O$$

Carbon tetrachloride reacts like oxalic acid, and gives the same blue colour With sulphur, thiodiphenylamine, the parent substance of methylene blue, is produced

$$(C_{6}H_{5})_{2}NH + 2S = C_{6}H_{4} \underbrace{\begin{array}{c} NH \\ S \\ \end{array}} C_{6}H_{4} + H_{2}S$$

(Berntheon, Ber 1883, 16, 2897) When the varour of diphenylamine is passed through a red-hot tube, carbazole is formed.

$$C(_{\mathbf{0}}\mathbf{H}_{\mathbf{5}})_{2}\mathbf{N}\mathbf{H} = \begin{bmatrix} \mathbf{C}_{\mathbf{6}}\mathbf{H}_{\mathbf{4}} \\ \mathbf{C}_{\mathbf{6}}\mathbf{H}_{\mathbf{4}} \end{bmatrix} \mathbf{N}\mathbf{H} + \mathbf{H}_{2}$$

(Graebe, Annalen, 167, 127); and the latter is also obtained from thiodiphenylamine and copper powder (Goske, Ber 1887, 20, 233).

Tests —A solution of diphenylamine in cone sulphuric acid gives a deep indigo-blue colour with nitric, introus or chloric acid (Ber 1912, 45, 2641, 1913, 46, 2286, Chem Zeit 1912, 26, 1239, Bull Soc Chim 1914, (14), 15, 186. The commercial product is a pale-yellow solid, melting not much below 54°, it should contain no oil, possess no unpleasant odour, nor turn blown on exposure to air

Derivatives -By passing chlorine into a solution of diphenylamine in acetic acid, a tetrachlorodiphenylamine is formed 4- chloro-2.6 dinitrodiphenylamine has been prepared (Ullmann and Shinang, Ber 1911, 44, 3730 Bromine in alcoholic solution produces a tetrabromodiphenylamine, in acetic acid, a hexabromo-derivative, and at a high temperature in presence of rodine, an octo- and deca-bromocompound. Diododiphenylamine is formed by the action of iodine in alcoholic solution in presence of mercuric oxide Bromine derivatives of nitrodiphenylamine are also known (H and P Ryan, Proc Roy Irish Acad 1919, By nitration in acetic acid solution, a tetranitro-derivative is formed, whilst direct nitration yields a hexanitro- compound, crystallising in yellow prisms, and melting at 238° The latter has strongly acid characters, and gives an ammonium salt which is used as a yellow dye known as aurantia (q v) Nitrosodiphenylamine is obtained by acting upon the base with amyl

Alkyl Derivatives—Methyldiphenylamine is obtained by heating together diphenylamine (100 parts) cone hydrochloric acid (68 parts), and methyl-alcohol (24 parts) for 10 hours to 250° in an autoclave (Girard, Bull Soc. chim [11]23, 2) It is a liquid, b p 282° The ethyl derivative is obtained in a similar way, b p 293° The amyl derivative boils at 330°–340° Benzyl chloride produces a benzyl derivative which is used in the preparation of viridine and alkali green

Amino- Derivatives —2-Chloro-5-ntrobenzenesulphonic acid combines with p-phenylenediamine to form 4-nitro-4-aminodiphenylamine sulphonic acid SO<sub>2</sub>H

which, on reduction, gives p-diaminodiphenylamine sulphonic acid (Baeyer & Co., D. R. P. 86250, 1894, Farb vorm Meister, Lucius and Bruning, D. R. P. 265197, 1914). Similar derivatives are obtained by the action of p-aminophenol, animophenol sulphonic acid, and p-aminosalicylic acid. The sulphonic acid group may be removed by heating with dilute acids under pressure (Meister, Lucius, and Bruning, D. R. P. 112180, 1899). Paraphenylenediamine and phenol in presence of copper salts and a solution of hypochlorite are oxidised to p-amino-phydroxydiphenylamine (Aktien Gesell. D. R. P. 204596, 1907). Aminohydroxydiphenylamine is also obtained by the electrolytic reduction of m-introdiphenylamine (Piccard and Lassen, J. Amer. Chem. Soc. 1918, 40, 1079). Diphenylamine derivatives prepared by the action

primary aromatic bases such as p-phenyleneamine, p-aminophenol, and aminosalicylic acid (1.34)-dintrochlorobenzene, (1.346)-ditrodichlorobenzene, and (1.43)-nitrochloronzene sulphonic acid, are used in the preparaon of sulphide colours, of which the most imrtant is dinitroxydiphenylamine

 $C_6H_3(NO_2)_2NH C_6H_4(OH)$ 

tained from p-aminophenol and dinitrochloronzene. Phenoxydinitrodiphenylamine is obnied from 1 3 dichloro 4 6 dinitrobenzene, tassium phenate and amine (Borsche, Ber 17, 50, 1339)

Diphenylaminedicarboxylic acid is obtained the action of anthramilic acid on o-chloronzoic acid in presence of finely divided copperher amino-acids behave similarly (Meister, icius, and Bruning, D. R. P. 148179, 1902, irgotti, Gazetta, 1914, 44, 1, 386)

Diphenylamine combines with formaldehyde, d the product forms an additive compound the sodium bisulphite

 ${}^{1}_{6}{}^{1}_{5}{}^{1}_{2}N]_{2} + {}^{1}_{1}SO_{3}Na$ 

=(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>N CH<sub>2</sub> SO<sub>3</sub>Na + (C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>NH ad. And Sod Fab D R P 158718, 1904) r nitro- derivatives, see Borsche and Rantelf, Annalen, 1911, 379, 152, B Linke, prakt Chem 1915, n 91, 202, Eckert and einer, Monatsh 1914, 35, 1153, Meldola, ster and Brightman, Trans 1917, 111, 546, yan and Glover, Proc Roy Irish Acad 1918, [B] 97, 194, J Soc Chem Ind 1918, 762, A, n Duin and van Lennep, Rec trav chim 19, 38, 358, Hofman and Dame, J Amer em Soc 1919, 41, 1413 J B C DIPHENYLAMINE BLUE v TRIPHENYL-


**DIPHENYLAMINE BLUE** v TRIPHENYL-THANE COLOURING MATTERS

DIPHENYLAMINE ORANGE v Azo- COLOUR-

**DIPHENYLBISDIAZONIUM CHLORIDE** *v* AZO- COMPOUNDS.

DIPHENYL BROWN, -CATECHINE, -CHRY-IENES, -CITRONINE, -FAST BLACK, -FAST LLOW, -GREEN, -ORANGE, -RED v Azo-LOURING MATTERS

### DIPHENYL-DIMETHYLOLID



e dilactone of dihydroxy-diphenyl dicarboxylie d, is the mother substance of a small series colouring matters, of which but one natural resentative, ellagic acid, is as yet known to st. The methylolid grouping —CO—O—sent in the compounds has a feebler chromoric character than the carbonyl itself, and this account the dyeing properties of the mbers of this class are inferior as regardsingth to those of the benzophenone, flavone, I allied dyestuffs. That the diphenyl linkage exerts an influence is certain, for gallonin containing as it does digalloyl nuclei:

)H)  ${}_{1}C_{6}H_{2}$ —CO—O—C( ${}_{6}H_{2}(OH)_{2}$ —CO—O—

JH)  $(C_6H_2 - CO - C_6H_2(OH)_2 - CO - C_6H_2$ 

for whereas the former is colourless, the latter possesses a strong yellow tint.

Though diphenyl-dimethylolid is at present unknown, diphenyl-methylolid:

needles, melting at 92.5°, has been prepared by various methods, and of these that of Richter (J pr Chem [2] 28, 294), who obtained it by the action of phosphorus oxychloride on ethyl salicylate

is the simplest

Colouring matters derived from diphenylmethylolid itself do not appear to exist in nature, though certain of them can be readily prepared by the action of alkali on diphenyl dimethylolid compounds, in which case hydrolysis followed by the elimination of a carboxyl group occurs:

By a more diastic action of the alkali, hydroxydiphenyl derivatives are produced

Interesting is the fact pointed out by Graebe (Ber 1903, 36, 214) that by distillation with zine-dust diphenylmethylolid yields fluorene.

in addition to diphenyl and methyldiphenyl Formerly the production of the first-named hydrocarbon from ellagic acid proved a stumbling-block in attempts to obtain a clear conception of its constitution.

A. G. P.

DIPHENYLENEMETHANE v FLUORENE.
DIPHENYLHYDRAZINE v HYDRAZINES.
DIPHENYLHYDRAZONES v HYDRAZONES
DIPHENYLKETONE v BENZOPHENONE and
KETONES

**DIPHENYLMETHANE** was first obtained by Jena, in 1870 (Annalen, 155, 86), by the dry distillation of barium diphenylacetate

According to Zincke (Annalen, 159, 374), it is produced by the action of finely divided copper, non, or zinc or their chlorides, on a mixture of benzyl chloride and benzene or by heating benzoylbenzoic acid with soda-lime Friedel and Crafts (Bull Soc chim [n] 41, 324) and Schwartz (Ber. 1881, 14, 1526) obtained it by the action of benzene on methylene chloride in presence of aluminium chloride; E & O. Fischer (Annalen, 194, 253), as by-product in the formation of triphenylmethane from chloriform and benzene in presence of aluminium chloride, the Friedel-Crafts reaction applied to arominate

alcohols in presence of benzene (Huston and Friedemann, J. Amer Chem Soc. 1916, 38, 25): and Hirst and Cohen (Chem Soc. Trans 1895, 67, 826), by the action of the aluminium-mercury couple on a mixture of benzyl chloride and benzene It is also formed by the reduction of benzophenone (Zincke, Thorner, Ber. 1877, 10, 1473, Staedel, Annalen, 194, 307; Graebe, Ber. 1874, 7, 1624), and by the action of phosphorus pentoxide on benzyl ethyl ether Meyer, J prakt chem 1910, n. 82, 538) But But the methods which have found a technical application in the production of diphenylmethane derivatives, as well as of the parent hydrocarbon, are those of (1) Meyer and Wurster, which consists in heating a mixture of benzyl alcohol, benzene and conc sulphuric acid (Ber 1873, 6, 693), and of (2) Baeyer, in which benzene, formaldehyde, or methylal are shaken together in presence of cone sulphurie acid (Ber 1873 6, 221)

$$CH_2O + 2C_6H_6 = CH_2(C_6H_5)_2 + H_2O$$

Properties -- Diphenylmethane crystallises in long, colourless, prismatic needles, possesses a fragrant smell, melts at 26°-27°, and distils without decomposition at 261°-262° solves easily in alcohol, ether, and chloroform

Reactions -On oxidation with chromic acid, it yields benzophenone On passing the vapour through a red-hot tube, it is converted into o-diphenylenemethane. On heating it with sulphur to 250°, it gives tetraphenylethylene, and on prolonged chlorination with iodine chloride at 350° it breaks up into hexachlorobenzene and carbon tetrachloride (Ruoff, Ber 1876, 9, 1485)

Nitro Derivatives — On nitration, diphenylmethane yields two dinitro- derivatives, one of which melts at 183°, and the second, which is the chief product, at 118° Other dinitro- derivatives are obtained from nitrobenzyl chloride, benzene and aluminium chloride, nitrobenzyl alcohol, benzene and sulphuric acid or nitrobenzene, formaldehyde and sulphuric acid, in a similar manner to the parent hydrocarbon Also nitrohydroxy- derivatives can be prepared by condensing nitrophenols and their ethers with formaldehyde and cone sulphure acid (D. R. P. 72490) The prolonged action of mitric and sulphuric acids gives rise to a tetranitro-derivative

Amino- Derivatives — The only substances possessing at the present time an industrial value are the amino- derivatives of diphenylmethane. They may be obtained by the reduction of the nitro- derivatives by tin and hydrochloric acid, but the more common method is to effect condensation between the aromatic base and formaldehyde, a general process for which a number of patents have been granted in recent years. Thus, by heating amline, aniline hydrochloride and formaldehyde, 40 pc, to 100°, 44'-diaminodiphenylmethane is produced (D. R. PP 53937 and 55565). The reaction appears to occur in two stages, for on heating aniline, formaldehyde, and a strong alcoholic solution of potassium hydroxide, the following change takes place :-

 $\begin{array}{l} \mathrm{CH_2O} + 2\mathrm{C_6H_6NH_2} \\ = \mathrm{C_6H_6NH\ CH_2\cdot NH\ C_6H_6 + H_2 O} \end{array}$ 

and the latter when heated with aniline and its h, in mioride passes into the diphenylmethane

derivative. A tetramethyldiamino- compound is obtained in a similar manner from dimethylaniline and formaldehyde in acid solution Similarly, a diphenyldiamino- derivative

C6H5NH C6H4 CH2 C6H4NH C6H5 has been prepared from diphenylamine and formaldehyde (D R P. 58072), a tetramethyldiaminodihydroxy- derivative

 $(CH_3)_2N(OH)C_6H_3CH_2CH_3(OH)N(CH_3)_2$ from dimethyl-m-amino phenol (D R PP 63081 and 67001), and a diethoxydiaminodiphenylmethane NH<sub>2</sub>(C<sub>2</sub>H<sub>5</sub>O)C<sub>6</sub>H<sub>3</sub> CH<sub>2</sub> C<sub>6</sub>H<sub>3</sub>(OU<sub>2</sub>H<sub>5</sub>)NH<sub>2</sub> from o-phenetidine and formaldehyde in presence of hydrochlone acid (D. R. PP. 70402, 72490, 73951, 73946)

Other methods for obtaining 44'-diaminodiphenylmethane are (1) the action of p-aminobenzyl alcohol on aniline hydrochloride in aqueous solution (D R P 96762), and (2) heating p-aminobenzylaniline with hydrochloric acid (D. R. P. 107718)

 $\begin{array}{c}
\text{acta} \text{ Cir} \\
\text{NH}_2\text{C}_6\text{H}_4 \text{ CH}_2 \cdot \text{NHC}_6\text{H}_5 \\
= \text{NH}_2\text{C}_6\text{H}_4 \text{ CH}_2 \cdot \text{C}_6\text{H}_4\text{NH}_2 \\
\text{has been}
\end{array}$ The 4.4 diaminodiphenylmethane has been prepared from p-aminobenzylaniline by heating with aniline and similar aromatic bases

 $\begin{array}{l} \mathbf{NH_2C_6H_4\ CH_2NH\ C_6H_5+C_6H_5NH_2} \\ = \mathbf{NH_2\ C_6H_4\ CH_2\ C_6H_4\ NH_2+C_6H_5NH_2} \end{array}$ 

(v Braun and Kruber, Ber 1913, 46, 3952) For diaminoditolyimethane see Farb vorm. Meister, Lucius and Bruning, D R. P 252916,

A 1 1' dichloro 2 2' diamino diphenylmethane is described by Duval, Compt rend 1912, 154, 780, and by Meyer, Ber 1914, 47, 1161, who prepared it from formaldehyde and o-chloraniline

A tetrahydroxy derivative was obtained by Ghosh and Watson (Chem. Soc Trans. 1917, 111, 824) from quinol and formaldehyde

Aryl derivatives of diphenylmethane, prepared by the Grignard reaction, are described by Boudroux, Compt rend 1915, 161, 131

Sulphonic acids of diphenylmethane have been obtained by Wedekind and Schenk, Ber 1911, 44, 198, and Staudinger and Pfenninger, Ber 1916, 49, 1941.

Diphenylmethane mono- and dicarboxylic acids have been prepared by Mitter, Ber 1912, 45, 1207, by the action of oxalyl chloride in presence of aluminium chloride on diphenylmethane J. B C

DIPHENYLMETHYLAMINE. Methyldiphenylamine v Diphenylamine.

DIPHENYLTOLUQUINOXALINE v Azines.

DIPICOLINIC ACID v BONE OIL.

DIPLOSAL. The salicylic ester of salicylic acid OII C<sub>6</sub>H<sub>4</sub> COO C<sub>8</sub>H<sub>4</sub> COOH, a white crystalline powder, mp 147°; almost insoluble in water and dilute acids. Used in the treatment of rheumatic diseases (v. SYNTHETIC DRUGS).

DIPPEL'S OIL V BONE OIL

DIPROPÆSIN. A derivative of urea containing the residues of 2 mols of proposin.

DIPROPYLACETIC ACID v OCTOIC ACIDS.

**DIPYRIDINE** v Bone oil.

DIPYRIDYL v BONE OIL

DIRECT BLUE, -BROWN, -GREY, -INDONE BLUE, -ORANGE, -ROSF, -YELLOW v. Azo-COLOURING MATTERS,

DISALOL. Trade name for phenyl salicylsalicylate

DISAZO- and TETRAZO- COLOURING MATTERS. The terms 'disazo' and 'tetrazo' are both used to indicate the presence of two azo- groups (N.N) in the molecule of a colouring matter, but at the same time they indicate the mode of preparation of such dyostuffs rather than, strictly speaking, their constitution. Tetrazo- colouring matters may be regarded as a special division of the disazo-colouring matters, as will be seen from the following classification.

Primary disazo-colouring matters are obtained by the action of two molecules of a diazo-compound (derived either from two molecules of an amine or from two different amines) on one molecule of an amine (e.g. m-phenylenediamine, m-tolylenediamine), a phenol (e.g. phenol, resorcinol, a dihydroxynaphthalene) or an amino-phenol (e.g. aminonaphtholsulphonic acids). The position taken up by the diazo-groups may be illustrated by the commonest cases (see also Azo-colouring matters) m-Phenylenediamine. The azo-groups occupy the two positions parawith respect to the two amino-groups. When one of these para-positions is occupied, as in m-tolylenediamine, the second azo-group enters the ortho-position with respect to the second amino-group; thus

$$CH_3 \underbrace{\hspace{1cm} N_2 - N_1}_{NH_2 - N_2} NH_2$$

Phenol, a-naphthol The para- and orthopositions with respect to the hydroxy- group are occupied

Resorcinol Two isomeric disazo- dyestuffs may be formed. When the combination is effected in alkali hydroxide solution, the symmetrical compound (I) is obtained, but when

the reaction takes place in the presence of sodium acctate or sodium carbonate, the unsymmetrical

compound (II) is produced

Dihydroxynaphthalene enter the usual position with respect to the hydroxy- groups, as would be expected if these groups represented two independent naphthols (of Azo- Colouring Matters)

Aminonaphtholsulphonic acids. In this case also, the compound may be regarded both as an independent naphthylamine and as a naphthol, one molecule of the diazo-compound is combined in acid solution, when the position taken up by the azo-group is the same as if the substance were a naphthylamine, and the second molecule is combined in alkaline solution as if the substance were a naphthol

Secondary disazo- colouring matters are obtained by the action of a diazoazo- compound on an amine or phenol; their general formula is  $X:N_2:Y:N_2:Z$ , the essential difference from the primary class being that Y does not contain auxochrome groups  $(NH_2 \text{ or } OH)$ .

Tetrazo- colouring reatters are produced from diamines, both amino- groups of which are diazotised (i.e. the diamine is tetrazotised) and combined with two molecules of the same or

different components (amines or phenols) These are mostly derived from benzidine, tolidine, dianisidine, &c; some, however, are prepared from p-phenylenediamine, and in this case direct tetrazotisation cannot conveniently be effected, but the monoacetyl compound is first diazotised and combined, the acetyl-group being then eliminated and the free amino- group is diazotised and combined with the second molecule of a component 1 When the first combination has been effected with an amine (in practice nearly always a naphthalene derivative), and in consequence a diamino- compound is produced on saponification, treatment with one molecule of nitrous acid leads to the diazotisation of the benzenoid amino- group, that attached to the naphthalene ring not being attacked

For the sake of completeness, it is necessary here to describe those dyestuffs which contain

three or more azo- groups

Trisazo- colouring matters are obtained in the following ways (1) Combination of three molecules of a diazo- compound with a phenol method is of only theoretical importance, and has hitherto been applied only to the cases of phenol (Wallach and Fischer, Ber 1882, 15, 2814, Grandmougin and Freimann, ibid 1907, 2814, Grandmougin and Freimann, volu. 1907, 40, 2662, J pr Chem 1908, [n ] 78, 384; Heller and Notzel, volu. 1907, [n ] 76, 58; Heller, volu. 77, 189), phloroglucinol (A G Perkin, Chem Soc Trans 1897, 71, 1154), and resorcinol (Orndorff and Ray, Ber 1907, 40, 3212; Amer Chem J 1910, 44, 1) (2) Combination of one molecule of a diazozo- and a diazo- compound with a pharmal course and a diazo- compound with a pharmal course and a diazo- compound. with a phenol, amine, or aminophenol diazoazo- compound may be produced, for example, by diazotising an aminoazo- compound such as p-sulphobenzeneazo-a-naphthylamine, or the intermediate product, obtained by combining a tetrazotised diamine (benzidine, &c) with one molecule only of a component (such as salicylic acid, &c ), may be used, or the formation of the diazoazo-compound may be left to the last stage of the operation, for example, a tetrazotised diamine (benzidine) may be combined with one molecule of an aminonaphthol (H acid) in acid solution, a diazo-compound (diazobenzene) now added in alkaline solution and the diazodisazo-compound coupled with a component (m-phenylenediamine or phenol) (Ene direct black GX and Ene direct green DT). (3) Combination of a diazotised tetrazo-colouring matter (which must, of course, contain a diazotisable amino- group) with a phenol or amine, either in substance of on the fibre bination of a triamine (eg magenta), having each amino- group diazotised, with three molecules of a component

Tetrakisazo- colouring matters, containing four azo- groups, are obtained (1) by the combination of two molecules of a diazoazo- compound with a phenol or amine (e.g. Anthracene acid brown B, v. Azo- colouring matters). (2) by combining two molecules of a diazo- compound with one of a tetrazo- colouring matter (e.g. Benzo brown G); (3) by combining a tetrazo- compound with two molecules of a monoazo-colouring matter (e.g. Hessian brown BBN);

<sup>1</sup> An alternative method is first to combine diazotised p-nitroniline with the component, then to reduce the nitro- group with sodium sulphide, and diazo<sub>v.</sub> <sup>1</sup> amino- group thus formed.

(4) by tetrazotising a tetrazo- colouring matter (containing two diazotisable amino- groups) and combining with two molecules of a component (e.g. Dianil black PR), (5) by diazotising a trisazo-colouring matter (containing one diazotisable amino- group) and combining with a phenol or amine either in substance or on the fibre

Colouring matters, containing more than four azo-groups, can be prepared, but, generally speaking, their functorial power is little or no greater than those described above. J. C. C.

DISINFECTANTS. A disinfectant is agent that will kill lower organisms which act injuriously on the higher forms of life though most disinfectants are also antiseptics and deodorants, a clear difference between the three classes lies primarily in their uses and Disinfectants are germicides, anti septics may only retard or inhibit the growths It is not essential for a disinfectant to effect absolute sterility in practical disinfection, there is nearly always a residue of highly resistant and non-pathogenic organisms destroy these involves great additional time and cost, and may even be disadvantageous when they act as natural scavengers in removing objectionable organic matter and antagonising dangerous forms The choice of an agent will obviously depend on considerations of safety, convenience, and economy in use

#### INORGANIC SUBSTANCES,

Ordinary oxygen acts slowly by the aid of Its activity is slightly inaerobie organisms creased by compression; it becomes exceedingly energetic when converted into Ozone. Since ozone is decomposed by heat and absorbed by most metals and by otherwise mert organic matter, the chief conditions for success and economy in working are (1) avoidance of sparking; (2) keeping the gas and the apparatus cool; (3) straining the air, and also partially drying when it is desired to prevent the formation of oxides of nitrogen, (4) parts in contact with ozone must be made of, or lined with, unoxidisable materials. (5) the grosser organic impurities, as well as sulphuretted hydrogen and ferrous non, should be removed before disinfection. The use of ozone on the large scale for punifying water is now well known, and the principal types of apparatus have been described by Don (Inst Mechan Figureers, Jan and Feb. 1909, see also S Rideal, J. R. San Inst. 30, 1, 1909, Rideal found at St Maur Waterworks, Paris, that with the sand-filtered (Marne) river water, containing a fluctuating number of bacteria, 28-320 per c.c, with a mean frequency of coli organisms of 1 per 100 cc (in more than onethird of the tests it was 1 in 40 cc), ozone in the proportion of about 0.6 gram per cubic metre of water (6 lbs per million gallons) destroyed all bacteria except an average of I per c c. of innocuous spore-bearing organisms of the subtilis type. The ozonised air was applied in a De Frise tower at the rate of about 40 p.c of the volume of water treated, and contained 1.4-18 grams of ozone per cubic metre (determined at frequent intervals by neutral potassium iodide) An advantage of ozone is that in disinfecting it adds no mineral proved that where water is not too contaminated

with organic matter, sulphuretted hydrogen, or ferrous iron, ozone will destroy all pathogenic organisms at a reasonable cost. In the dry state, like chlorine, it has very little action on micro organisms, and therefore does not act powerfully on bacilli in air. S grams per cubic metre of air in a room ordinarily closed accomplish disinfection in 3 hours. Its use as an adjunct to ventilation has often been proposed, but its irritant action on the lungs prohibits its use in these quantitites. It is most active when moist. Ozonised air, under pressure, has been used successfully for destroying moulds and bacteria in brewers' foul casks, and has been introduced into wort in the latter part of the fermentation 'to destroy noxious germs' (Fng. Pat. 22355, 1898) Will and Beyersdorfer (Zeit ges Brauw 1912, xxxv 73) showed that air containing 0 6-0 7 gram of ozone per cubic metre destroyed organisms mimicable to the On account of its rapid consumption by oxidisable substances, it can only be effective in disinfecting the surfaces of organic solids the Pittsburg Homoropathic Hospital,  $\mathbf{At}$ ozone, besides purifying the water supply, is used for sterilising instruments, bandages, &c and for dressing wounds and ulcers storage it would probably be useful for destroying injurious organisms and odours, thereby possibly lessening the amount of refrigeration required Many attempts have been made to utilise it for disinfecting, combined with bleaching, textiles, feathers, and other surface-contaminated articles, and have been only limited by questions of expense

Patent 4153 of 1902, for 'fixing ozone in liquids,' states that pure acetaldehyde dissolves 2823 times its volume of ozone, that 10 pc solutions of acetaldehyde in water dissolve 536, in alcohol 748, in ether 859 volumes of ozone; that these solutions are very stable, and have

been successfully used in surgery.

Hydrogen peroxide. The first experiments (Ber 15, 1585) showed that this was a bactericide which, in the dilute state, appeared mert to enzymes such as diastase,1 ptyalin, pepsin, and pancreatin; did not injure foods; and is also innocuous and non-corrosive These properties suggested its use for killing pathogenic organisms, and those that cause change in articles of diet, especially milk (Renard), and also as an internal disinfectant. Altchoefer (Centr Bakt 1890, viii 129) and most later observers give 1 per 100 of H<sub>2</sub>O<sub>2</sub> as required to destroy pathogenic organisms in 24 hours. In 1903 S. Rideal found that in milk raised to 50°, then treated with 0.6 gram of H<sub>2</sub>O, per litre, and the heat maintained at 52°-55° for 8 hours, pathogenic organisms and even the spores of *B anthracis* and subtilis were killed, the milk was not physically altered, and kept for over a month unchanged in the closed receptacles (Budde's patent 10903, 1903) Similar results have been obtained by Lewin and others, by Hewlett (Lancet, Jan 27, 1906), and Reichel (Zeit f Hyg 1908, lxi 49) So that hydrogen peroxide aids pasteurisation, and enables it to be efficient The solution is at a lower temperature.

The enzyme catalase (present in malt, yeast, milk, and many organic liquids) and hydrogen peroxide mutually decompose one another (Van Laer, J Soc. Com Ind 1906, 489, 1909, 950), but the above statement remains practically true. (See also that 1909, 1267.)

frequently used as a spray, gargle, or dressing, and has had a wide application for war service in dealing with gas gangrene infected wounds (see reports by Capt M Page, R.A M S., Lancet, Sept 1914, Col Makins, ibid. August, 1914)
Peroxide of hydrogen is present in 'Sanitas' (qv) 'Perhydrol' is a 30 pc or 100 volume solution of  $H_2O$ . The use of peroxide of hydrogen in medicine depends partly on its bactericidal action Registered names of solu-tions are 'Dioxogen' (3 pc), 'Hydrozone,' 'Glycozone' and 'Pyrozone' (dental practice), and 'Peroxols' are mixtures of  $H_2O_2$  with various disinfectants Peroxides of sodium, magnesium ('Hopogan'), zinc ('Ektogan' or 'Dermogen') and mercury, generate oxygen or H<sub>2</sub>O<sub>2</sub>, and have analogous use Hydrogen peroxide is more stable in ether than in water, 'ozonic ether' is an ethereal solution containing about 12 pc of H2O2, which has been used in whooping-cough and scarlet fever 'Perhydrit' or 'Hyperol' dressings, containing H<sub>2</sub>O<sub>2</sub>, urea, and a trace of citric acid, have recently been introduced

Foregger and Philipp (J Soc Chem Ind. 1906, 298) treat at length on the applications of peroxides, and suggest calcium peroxide ('Bicalzit') instead of H<sub>2</sub>O<sub>2</sub> for sterilising milk, as more lasting in effect. For water sterilising, tablets of magnesium perovide and citiic acid are made; Park found that 01 gram of this peroxide killed upwards of 2 million typhoid organisms in 150 cc of water in 30 minutes, and 0.2 gram gave the same result in 1 minute Although Croner (Zeit f Hyg 1908, lvin 487) obtained sterility in water with the use of 07-10 gram MgO, per litre, more recent investigations by Young and Sherwood (J. Ind. Eng Chem 1911, 3, 495) have indicated that over five times this quantity must be used to ensure sterility Patents for similar uses of  $H_2O_2$  and other peroxides are very numerous.

Nitric acid and oxides of nitrogen. The corrosive action of nitric acid limits its value as an active disinfectant. J. C. Smith used it with success in 1780 for fumigation in a violent outbreak of typhus fever in the British fleet (Vallin, Désinfectants, 1882, 265). Disinfection of rooms by nitrogen peroxide was carried out during the siege of Paris. 48 hours were occupied, and the cost was very high. Severe bronchitis and several deaths from poisoning have resulted from breathing this gas, and intre disinfection has been replaced by safer methods, except in the local destruction of bacteria as in laboratories

Sulphur. Dusting with sulphur is used for killing insects and fungi on plants, as sulphurous acid is gradually formed. The use of sulphur internally is due to its forming sulphuretted hydrogen in the system, and thereby destroying or enfeebling micro-organisms and higher parasites. A solution containing calcium polysulphides, made by boiling milk of lime with sulphur, has been extensively used as an insecticide (see Agric. Bull. 319 and 320, New York, 1909)

Sulphur dioxide disinfection is still a subject of controversy, but certain of the unfavourable results adduced have been due to faulty application. Spores are not generally killed by it, and the dry gas has very little effect on microorganisms, but it destroys all vermin, and therefore is of special value in combating insect-borne

As to its effect on the lungs, about diseases 5 pc in the air has produced fatal results, but pc. can be tolerated for some time, and a larger amount with the protection of a wet 1 lb. of towel alkalised with washing soda sulphur burnt in a room of 1000 cubic feet produces theoretically an air-content of 115 pc SO,, and requires 0 56 lb. of water to turn it into sulphurous acid Such an amount of moistening is, however, not necessary, on account of the natural moisture of the air. Prescriptions as to the proper quantity of SO, have varied greatly, partly because in some cases it has been considered sufficient to destroy only a special organism Klein, Houston, and Gordon (Report, Med Off LCC 1902) found that with 31 lbs. S per 1000 cubic feet in 24 hours, B typhosus, diphtheria, pyocyaneus, and Staph p aureus were killed, but not anthrax spores nor B tuberculosis in dried sputum In America, 5 lbs per 1000 cub feet have been found sufficient to kill all spore-bearing organisms after 16 hours' exposure (Rosenau) S Rideal's results with the liquefied gas were as follows Silk threads and paper slips infected with B coli and Staph p aureus were sterilised after 24 hours' exposure in a sealed room of 1500 cub feet, into which 20 ozs of  $SO_2$  were passed Subtiles spores were not killed With 10 ozs, the threads were again sterilised, but the paper shps were not shallow pans of water were exposed in the room to supply moisture, the SO<sub>2</sub> content of the air was actually lowered after the 24 hours (through absorption by the water), from 0.5 p.c. present in the previous experiment to 02 pc, and the organisms were still living Therefore moisture should only be supplied by washing or lightly spraying the surfaces In chemical disinfection, it is generally recommended that walls should be previously stripped and fabrics spread out on lines, but spreading the infection must be carefully avoided The room must be sealed by obvious means, and, observing this condition, heating is advantageous Liquid SO, in canisters is more convenient than burning sulphur, and avoids risk of fire one pound of SO<sub>2</sub> has the liquid = 5} cub feet of the gas the disadvantage that it attacks metals and organic substances from the production of sulphuric acid; a good deal is also absorbed by plaster It has little penetrating power, and therefore bedding and thick articles must be removed and sterilised by heat The 'Clayton method 'of disinfection is much used for the holds of ships, and has been favourably reported on by Wade and Eyre for the Local Government Board (Rep 232, 1906, and Med Off Rep 1903-4, 330). Sulphur is buint in a special apparatus, and the gases, consisting of air, SO,, and a visible cloud of sulphuric acid, are forced in till the SO, reaches about 10 pc of the air in the infected space (For experience of its use, see Rev d'Hyg Oct 1902, NSW. Agric Gaz 1900 and 1901)

Sulphurous acid is used beneficially as a germicidal spray or wash in various parasition affections (eq. diphtheria), also internally in cholera and gastric fermentation, for rectal injections, and other allied objects 'Thiocamf' is a liquid obtained by the action of the gas on camphor It can be preserved without pressure in well-closed vessels, and on exposure to '1. a 6-oz bottle should evolve about 20 litres of SO<sub>2</sub>.

Sulphurous acid and bisulphites are widely used for checking fermentations and preserving food. Lehmann states that an amount equivalent to 0 004 p.c of SO<sub>2</sub> may be allowed without injury to health The Report of the British Departmental Committee on Preservatives, 1901, 26, leaves the matter open. Some commercial preparations are 'Meat preserve crystals,' 'Freeze-em,' and 'Hawke's anticrystals,' 'Freeze-em,' and 'Hawke's anti-ferment' For objections to the use of sulphites in food, see Thresh and Porter, Preservatives, 1906, 70-77, also an article by Durham in the J of Hygiene, April, 1909 The latter recommends the fixing of a maximum legal limit, not exceeding 0.01 p.c. of 'total sulphur dioxide, with a declaration of the amount, but less than 0 001 or 0 002 pc might be simply declared as

Sulphuric acid, like acids generally, is antiseptic, and in some cases disinfectant (which is a palatable strength) is fatal to Sp. choleræ in 15 minutes Stutzer found that this strength disinfected iron pipes, cleaning out sust and sediment without sensibly attacking the metal With 0.04 p.c. Ivanhoff destroyed cholera organisms in Berlin sewage, and with 0 08 pc in that from Potsdam In the laboratory it was found that 0 035 p c kills B typhosus in 30 minutes, and 0 07 pc in 15 minutes, in impure water infected ordinarily with typhoid, but with very heavy infections the latter strength requires 45 minutes He recommended 0 09 p c. as having advantages over heat sterilisation for dealing with the drainage from hospitals and other infected areas. the acidity would speedily be neutralised on mixing with the bulk of the sewage (But Assoc 1901)

Sodium bisulphate has been introduced by one of the writers and Dr. Parkes, as a means of sterilising drinking water for armies in the field and travellers 15 grains of NaHSO4 to a pint of water in 15 minutes destroys B typhosus and enteritedrs, Spirillum cholera, and internal parasitic worms. In effervescent tabloids, yielding a slightly acid solution, it was used with success in the South African, Russo-Japanese, and the European wars

Persulphates give off oxygen when moist, and are useful as hand disinfectants (Lancet, 1905,

n 1106)

Carbon disulphide, if it were not for its other properties, would be a valuable disinfectant and preservative. As well as its alkaline compounds, the xanthates, it has been found useful against fungi, and insect parasites such as the vine phylloxera. Ckeandi Bey's lamp burns it safely

for generating SO<sub>2</sub>.

Boric acid is not a disinfectant, but has a power of restraining bacteria which act injuriously in foods Rideal and Foulerton found that I in 2000 of a boric mixture, containing 3 parts of H<sub>3</sub>BO<sub>3</sub> and 1 part of crystallised borax, keeps milk sweet for 24 hours without appreciable effect on digestion The British Departmental Committee already referred to, recommend that the only preservatives allowed to be used in cream, butter, and margarine be boric acid, or mixtures of boric acid and borax, not exceeding 0 25 p c of H<sub>3</sub>BO<sub>3</sub> for cream, and 0.5 p c for butter and margarine Later, as this quantity appeared insufficient in the cream tande an official report by Hamill advised that 0 4 p.c H<sub>3</sub>BO<sub>3</sub> should be allowed in cream from

May to October inclusive (Food Report 10, LG.B 1909) Soluble borates are used as a larvicide in manure heaps. It has no action on the crops as they are rendered insoluble (Cook and Wilson, J Agric. Research, 1917, 10, 591).

The Halogens, chlorine, bromine, and iodine, act as disinfectants in several ways: (1) they can combine with H2 and liberate O, in water; (2) they can combine directly with organic matters, or replace the hydrogen in them, precipitating albuminous substances and rendering them imputrescible, killing organisms by combining with and coagulating protoplasm, removing their food or rendering it unsuitable, acting toward them as irritants or direct poisons, also producing substances which have that effect

Chlorine and the hypochlorites, like other oxidisers, are consumed by otherwise mert organic and inorganic matters present, but in ordinary disinfection leave behind them as substitution compounds which maintain germicidal power—the chloramines, hydrazine, and the chloroproteids (Rideal, J. R. San Inst. 31, 2, This paper shows that in dilute solution of hypochlorites the Rideal-Walker coefficient of 2 18 for 1 pc of available Cl is increased to 6.36 by the addition of an equivalent of ammonia, remains near this level for 24 hours, and even after 72 hours has an enhanced value Therefore, since the coefficient of chlorine itself averages 220 units, that of ammonia less than 0.7, and ammonium chloride nil, that of the chief product, chloramine NH Cl, must be over 600 units, hence this substance, but for its instability, would probably be the most valuable of disinfectants. It has a pungent odour, hitherto often mistaken for chlorine or hypochlorous acid, and gives the blue reaction with KI and starch With excess of ammonia, as in sewages, it gradually disappears, forming a salt of hydrazine, which still has a germicidal value (R W. coefficient of the base at least 24). The action of chlorine on nitrogenous compounds gives compound chloramines, many of which are insoluble, they have a tendency to become fixed on cellulose, and in this way attack the envelopes of organisms.1

Various substituted chloramines have been introduced by H D Dakin for use as sterilisers and disinfectants Amongst the more important may be mentioned chloramine-T, p-toluene sodium sulphochloramide, p-sulphodichloraminobenzoic acid, and dichloramine T (BMS Aug. 1917) The former, under the title Halozene, has been used for water sterilisation on a small scale.

The addition of ammonia solutions to bleaching powder solutions preparatory to the sterilisation of water has effected a considerable economy in the quantity of materials used in the public water supply at Toronto, Canada.

The chloramine is about five times as strong as the corresponding molecular concentrations

of sodium hypochlorite

The bromamines of the benzene series, the naphthaline chloramines and the protein chloramines, are all strong germicides, but suffer from lack of stability in solution.

Chloride of lime, or 'calx chlorinata' (about 33 p c. available chlorine), has some advantages

<sup>1</sup> Also see Cross Bevan and Bacon, Chem. Soc. Trans 1910, 244.

from being a solid, but is quickly deliquesced; and decomposed by air, takes some time to dissolve, and gives a quantity of insoluble waste Klein found that 1 p c of bleaching powder only killed less resistant organisms in 24 hours There is, in many respects, a preference for the sodium solutions. chlorinated soda or 'Eau de Labarraque' (often called 'Eau de Javel,' which was originally the potash salt), contains about 25 pc av Cl, 'Chloros,' 10 pc av. Cl, 'Antiformin' is a solution of caustic soda and hypochlorite used in breweries. Electrolytic preparations are 'Oxychloride,' up to 115 pc av 'Cl, and 'Hermite fluid,' used successfully at Poplar In the latter solution magnesia is employed as an adjunct in the electrolysis contains free hypochlorous acid av Čl 42 to The effective strength for local dis-48 pc infection is 0.35 p.c av Cl, which kills non-aatring organisms in 5 to 10 minutes, and anthrax and enteritidis spores in  $1\frac{1}{2}$  hours Chloride of lime has the objection that it leaves the surface for a long time damp from the presence of calcium chloride The same salt imparts its bitter taste, and increases the hardness, when chloride of lime is applied to the purification of drinking water, while the soda preparations tend to soften the water and are not sensible in the flavour. In trials at Guildford in 1904 (J. R. San. Inst. xxvi. 891) with electrolytic hypochlorite in the proportion of 0.75 to 1.0 part per nullion of av (1, one of the writers found that water containing an average of 90 organisms per c c was practically sterilised, while after standing 11 hours the odour and taste were normal, and there was no need for further treatment Candy's patent, 4212, 1909, removes excess of av Cl by filtration through coke, Thresh's, 3023, 1909, by treatment with metallic iron Other methods of removal are suggested. (1) by the addition of ferrous salts, subsequently removing the ferric salt formed by a 'Permutit' filter, (2) by the addition of gaseous sulphur dioxide, (3) by the addition of sodium bisulphite After the Maidstone typhoid epidemic, 1897, the infected reservoir, mains, and pipes, were sterilised with 1 pc solution of chloride of lime during 48 hours · no corrosion of the non was observed During a similar outbreak at Lincoln, in 1905, the water supply was sternised by sodium hypochlorite (av. Cl, 1 per million) with most satisfactory results In 1910 Toronto (reated its water supply with 0 14 part per million of av Cl as chloride of lime Since June, 1909, the writers have been applying 'Chloros' in the proportion of 1 to 2 parts av Cl per million to 100,000 gallons daily of the drinking water of a town in the north of England: the liquid is made practically sterile, coli is absent from 75 to 85 c c, and there is no effect on the taste At Cambridge, in 1910, a portion of the supply (lower chalk) was treated with one in 4 to 8 millions of av Cl as chloride of lime, and Houston found that this minute quantity destroyed 66 to 98 p c of the bacteria, and coli, which was always present before, was afterwards absent in 500 cc. This water, however, contained hardly any organic matter. America, treatment of the entire water supply of Jersey City (about 40 million gallons per la ger works, considerably reduce the cost, and day, from the Rockaway river) with chloride of lis cheaper than sand filtration. Itellerman, lime, has been adopted since the beginning of Pratt, and Kimberley (U.S. Dept. of Agric.

1909: the proportion used is now 0.2 part per million of av Cl, and the daily examination shows that coli is absent from 10 cc of the treated water In England, the 'De-Chlor system was applied at Reading to the prefiltered river water in 1910. Since the war chlorine treatment has been adopted for the London supply (11th Annual Report, M W B.)

Recently an impulse has been given to the sterilisation of water by means of gaseous chlorine injected by means of suitable dosing apparatus, owing to the convenience of handling and transporting chlorine in the liquid state contained in steel cylinders of 100 lbs capacity. Injection into the water may be direct through diffuser, as in the Wallace and Tieman apparatus, or indirect through the addition of a minor volume of more concentrated chlorine water to the bulk volume, as in the Electrobleach and Patterson plants NH<sub>3</sub> does not seem to enhance the germicidal value in the absence of alkaline bases

With sewage, of course, a larger dose is required, and the first experiments at Nice, Brest, and Worthing, in 1894, with the Hermite fluid, showed that, as with ozone and other disinfeetants, the agent should be applied, not to a raw liquid (except in special cases), but as a finisher' to one that was partially purified 1898, using an electrolytic brine (Woolf's 'Electrozone') for purifying an effluent at Maidenhead, one of the writers found that with three parts per million of av Cl, and a contact of 5 minutes, the bacteria were reduced to 10 per 50 cc, and that 17 parts per million killed all the organisms in 15 minutes. His long series of trials at Guildford commenced in 1904, using 'Oxychloride,' and proved that with 2.5 parts per million of av CI, tertiary effluents could be practically sterilised, but that the raw sewage and septic effluents required 30 to 70 parts, the contact period was  $4\frac{1}{2}$  hours (J. R. San Inst. xxvi 7, 1905) The oxygen consumed from permanganate in 5 minutes in the cold, as measuring the H2S and other easily oxidisable substances, was a guide to the amount of the agent used, and in the earlier experiments it was found that when the av (I added was 1.7 times this figure, the availability of the chlorine was sufficiently prolonged to ensure the desired sterilising effect, and the subsequent experiments frequently yielded excellent results with an amount equal to this oxygen figure, that is, one to one (Sec further, Rideal, Trans Farad. Soc iv 3, 1909) An important point shown was that the necessary factor of safety could be ascertained by simply testing with KI and starch, and noticing the persistence after 4½ hours of the blue colour, due in this case, as has been mentioned, not to chlorine, but to chlor-Schumacher (1905), at Hamburg, found that with raw sewage sufficient sterilisation is attained if after 4 or 6 hours the test shows a reaction equal to 19 and 17 parts per million of av Cl respectively Investigations on sewage filter effluents with chloride of lime were continued in America by Phelps and Carpenter (U.S. Technological Quarterly, Dec. 1906), who remark that the use of electrolytic chlorine manufactured on the spot would, in the

Bull 115, Oct 1907) conclude that with a hypochlorite, disinfection costs less than with copper sulphate. Later American experiments at several centres have proved that 2 to 5 parts per million of av Cl destroy 95 to 98 pc of the bacteria in 2 hours, and that 3 parts are sufficient for the trickling filter effluents tried (Mass Inst Technol 5, 1909; 6, 1910).

The Royal Commission on Sewage obtained results at Guildford which were in agreement with those of the writer (Appendix iv 1910, 87, also *ibid.* 84-171, and Rep. 1908, 195-201)

Sod. hypochlorite solutions are the best means of sterilising swimming baths, they remove from the surfaces growths of alga and fungi which harbour dangerous organisms. (Also see Rideal and Orchard, J. San. Inst. 556, 1906.)

'Ferrochlore' (Duyck) is the name of a treatment with 8 parts per million of ferric chloride and 0.5 part of chloride of lime, which at the same time clarifies water and liberates

hypochlorous acid 1

Hypochlorous acid has been employed as an antiseptic for wounds by Dakin, Daufresne, and Lorraine Smith, and is also known as "Eupid" and 'Eusol'. In these solutions hypochlorous acid is obtained by double decomposition of bleaching powder with sodium carbonate, with the addition of bone acid or sodium bicarbonate. The presence of sodium borate or sodium carbonate in the resulting solutions when freed from the precipitated calcium carbonate ensures the stability of the solution owing to the 'buffer' action of the easily hydrolysable salt.

Chlorine peroxide, obtained from potassium chlorate and sulphutic acid, has been successfully employed in the Howatson-Bergé process at Brussels, Ostend, and Lectoure (South France), for sterilising (previously clarified) polluted waters. The writer found that 20 parts of CIO, per million did not kill B typhosius in a vigorous pure culture in 2 hours, but that 50 parts killed it in half an hour. (See also Reychler, Bull Soc chim [3] xxv 13, Chem

News, 1901, 316)

On a large scale, chlorine and ozone are both in successful use at present for sterilising water. The latter seems preferable for drinking supplies, the former is suitable for effluents. From questions of efficiency as well as cost, both require that the liquid should be previously clarified, and as free from organic matter as

possible, that is, they are finishers

Time as in all disinfections, is an important factor, in an effort to get the work done quickly, it is sometimes forgotten that a longer time is advantageous, and economises the reagent. For rapid sterilisation of a small quantity, as in the field, a larger proportion of reagent is required, and from questions of portability, bromine and iodine have been used instead of chlorine, but are attended by residual products having a taste and physiological effect. These, in the quantity required for dealing with the moderate pollution ordinarily met with, have been proved to be negligible, although, with heavier contamination, the amount of the halogen consumed by the organic matter before a sterilising excess is present, may become in-

jurious in its products. Cash showed that the quantities of chlorine, bromine, and iodine required for disinfection of anthrax and tubercle were nearly in the ratio of their atomic weights

Bromine is a powerful disinfectant, but its is obviously limited by its danger. In 1897 use is obviously limited by its danger Altmann patented a solution of Br in KBr for sterilising water, using 40 parts per million of Br with 5 minutes' contact, and removing excess by ammonia The solution is, however, bulky, and the use of a potassium salt has disadvantages. Braithwaite introduced 'Biomidine,' a dry mixture of sodium or potassium biomido and bromate with sodium bisulphate, which with moisture liberates bromine Schumberg recommended free bromine in thin sealed tubes for the same purpose (Zeitsch f Hyg 1900, 53) in the proportion of 60 parts per million with 5 minutes' contact, removing excess by thio-sulphate (Braithwaite advised 57 parts) Schoder (ibid xxvii 306) and others have stated that this treatment is insufficient to sterilise, and the presence of residual bromides is against the process (see also Parkes and Rideal, Trans. Epidem Soc. xx 1901, Fraser, Publ Health, Sept 1902) It was tried in the Soudan campaign of 1898 20 parts per million of Br sterilises typhosus (in water) in less than half an houi It is more rapidly absorbed by organic matter than iodine

India owes many of its uses to its germicidal action, as in parasitic diseases. Grossich, in 1908 (Centi f Chiruig No 4), adopted the tincture as a cutaneous disinfectant before operations, the surface being previously cleansed by a 1 p c solution of iodine in benzene (Bogden, abid Jan 15, 1910) Instead of the tineture, Chassevant (Lancet, March 26, 1910) suggests the solution in chloroform, as it does not, like the former, cause desquamation, its use is specially advised prior to hypodermic injections (see also But Med J Feb 6, 1909, and Aug 14, 1910, Lancet, April 16, 1910) Solutions of iodine and iodine trichloride, both as field diessings, and enclosed in friable glass compounds as emergency first-aid sterilisers, have been extensively employed during the war (see B. M. S. Jan 2, 1915, Lancet, Dec. 12, 1914) During the Sikkim and Thibet expedition, rodine (in some cases as tincture) followed by sodium sulphite, was successfully employed for sterilising drinking water, and Nesheld recommends triple tablets of (A) include and indate, (B) citrie or tartain acid, followed after 2 minutes' contact by (C) sodium sulphite, and states that used in this way 3 8 to 5 parts per million of free iodine sterilises the organisms of typhoid, cholera, and dysentery. One of the writers, in 1905, confirmed this result as regards distilled water infected with 20,000 per e.c. of B typhosus, the A and B tablets used gave 4.75 parts of 10dine per

Iodine trichloride was introduced by Von Langenbach for sterilising the hands and instruments and for other surgical uses 50 parts per million are required to kill typhoid (in water) in 30 minutes fodine cyanide is said, by Kobert, to be universally destructive to lower forms of life, and is suggested for preserving biological specimens.

Organic substitution compounds of chlorine, bromine, and iodine, as a class, inhibit organisms,

<sup>&</sup>lt;sup>1</sup> For an account of trials at Marseilles of ozone, chlorine, and ultra-violet rays, see Engineering, Jan. 27th and reb. 3rd, 1911.

and may sometimes kill them. The germicidal power of a large number of these has been determined by Bechhold (Zeitsch Hyg. Infect 1909, lxiv 113), he finds that a Br atom gives a greater power than a sulphonic group \*Chloroform\* is often used to preserve perishable specimens and infusions Fairel and Howles (J Soc Dyers, 1908, 24, 109) state that CCl<sub>4</sub> kills typhoid, and that pure CHCl<sub>3</sub> does not kill it, whist 0 62 p c aqueous CHCl<sub>3</sub> kills it in less than 30 minutes

Iodoform was formerly much employed in surgery, and a large number of other rodine compounds have been introduced into medical and surgical practice. Many of them are non-irritant, but their bactericidal power is much inferior to that of dilute solutions of the element Funigation of sick rooms with rodine, sometimes in conjunction with other disinfectants, by burning prepared candles or lamps, has been tried, but apart from the fact that the method is unsatisfactory, no chamber can be disin-

fected while occupied.

The organic iodo compounds owe most, if not all, their germicidal power to the gradual liberation of iodine by the alkaline wound secretions. Iodoform itself gives rise to symptoms of poisoning owing to the too rapid evolution of iodine. Amongst the more important organic iodine derivatives may be mentioned hexamethylenetetramine compounds, chrysoform, iodoformin, phenol, and homologous compounds, aristol, iodothymoloform, curophen, traumatol, isophon, iodoamsol, iodocol, iodised albumen, tannin, gelatine, casein, gluten peptone lecithin, and various iodised oils have been protected under fancy names.

Fluorides, like bone acid, check bacteria, but do not kill them As food preservatives, they are only moderately efficient, and are not innocent physiologically, as they hinder pancreatic digestion Sodium silicofluoride ('salufer ') and borofluoride ('pyricit') have been tried Effront introduced the use of hydrofluorie acid and its acid salts for cheeking injurious fermentation, and J. Brand (Zeitsch. f. Brauw 1904, xxvn 115) states that 0.5 to 1 pc solution of acid ammonium fluoride is largely used for disinfecting rubber hose in breweries (see also Hehner, Analyst, 1902, 173, Richmond, Dep Comm on Preservatives, appendix xxxII, Thresh and Porter, Preservatives, 1906, 87) Hydrofluosilicic acid is the active constituent of 'Montanin' (obtained from by-products in the pottery industries), and 'Keramyl' (about 25 p c H SiF<sub>6</sub>) (J Soc Chem Ind 1904, 34, 1909, 904) Frank (Wasser u. Alwasser) recommended the use of aluminium silicofluoride for water sterilisation. They are stated to have no injurious action on health (Public Health, Oct 1908)

Carbonic acid seems to have a certain amount of disinfectant action, since 'aerated' beverages have been shown to be in many cases stoilised, and CO<sub>2</sub> under pressure preserves food (c<sub>I</sub> J Soc Chem Ind 1907, 1290)

Cyanogen and hydrocyanic acid are more destructive to insects than to lower organisms Funnigation of fruit trees with HCN is extensively practised in America, and is considered better than straying. For greenhouses, zinc capsules containing sodium cyanide are made; they are placed in dilute sulphuric acid,

and while the zinc dissolves the operator has time to retire.

Acids. An acid medium is unfavourable to the growth of most bacteria, and acids generally possess considerable disinfectant power Kitasato found that the growth of B typhosus in nutrient broth or gelatine was entirely prevented by the following percentages of different acids. 0 08 H<sub>2</sub>SO<sub>4</sub>; 0 2 HCl or HNO<sub>3</sub>; 0 28 SO<sub>2</sub>, 03 to 04 phosphone, acetic, carbolic, formie, oxalie, or lactic, 0 476 tartarie, citrie, and malie, 166 tannie, and 27 bone One of the writers has shown that in infected waters. 0 035 p c of H<sub>2</sub>SO<sub>4</sub> kills typhosus in 30 minutes, that in terms of acidity nitro-hydrochloric was slightly more active, and that with tartaric and citric at least three times the amount of acidity was required (Epidemiological Society, Jan 1901) But like other disinfectants, the activity of dilute acid solutions is considerably modified by the presence of added matters, for instance, in 1 pc peptone broth, Sp. cholera is killed by 0.01 pe hydrochloric acid in half an hour, but in 2 pc peptone broth, 004 pc HCl is required

Paul, Kronig, and Sawey (Phaim J 1900) conclude that the germicidal activity of acids generally, depends upon the degree of ionisation in solution, exceptions being HF, HNO<sub>1</sub> and trichloracetic, which excises specific toxic influences. In normal solutions, the three latter killed anthrax spores exposed on gainets in 24 hours, while H<sub>2</sub>SO<sub>4</sub>, HCl, HBi and HClO<sub>3</sub> were less active. Working with more dilute acid solutions, Winslow and Lochridge (Mass Inst. Tech. 1906) have compared the effect on B coli and tuphosus in tap water and in a 1-pc peptone water. With upwards of 10,000 organisms per ccc, sterilisation was obtained by the following percentages of each acid after

In agreement with the results of other workers, larger amounts of each acid are required to disinfect in the presence of peptone, with the organic acids, benzoic and acetic, this diminished activity is not so pronounced. Winslow and Lochridge find that the disinfectant power of the acids is not proportional to their strength in terms of normality, but, in the case of the mineral acids, hydrochloric and sulphuric, to the degree of ionisation of the solution, 99 to 100 pe reduction of B coli or typhosus being effected with either acid at the same concentration of dissociated hydrogen The evidence tends to show that the alteration in ionisation in a dilute acid solution brought about by the presence of otherwise mert substances in solution, as, for example, neutral salts of the acid, accounts for the variation in germicidal activity In practical disinfection, the subject is often rendered still more complex by a further interference -the absorptive effect on solids in Therefore the acidity, as found by titration, does not give the germicidal value of an acid solution unless these disturbing factors are considered.

Metallic derivatives. Warm alkaline leys are the most ancient of disinfectants: potassium and sodium hydroxides are sterilisers in the proportion of not less than 2 to 5 pc, their carbonates in 5 to 10 pc. At 60° a contact of 5 minutes ensures the death of most organisms (Zeitsch. f. Hyg. 1903, 349), hence ordinary soaps have a slight disinfectant value. Strong bline is preservative, but not disinfectant, and the practice, formerly frequent, of washing out milk cans and other utensils with it, is not a safe one. Lode shows (Chem. Zentr. 1902, i. 1122) that 50 pc. salt solution does not kill the spores of common moulds.

Lime, as quicklime, is fatal to bacteria when it comes in contact with them in an almost dry condition, and has been much used where there is a considerable space and bulk of material to be dealt with. It is not safe, however, to trust entirely to the old method of burying infected bodies in lime, as, on disinterring, the live spores may again be diffused, unless a long period has clapsed The disinfecting power of milk of lime and whitewash has been much overrated, as to milk of lime, Liborius stated (Zeitsch f Hyg n 25) that in the proportion of 74 parts per million of CaO, it destroyed typhoid bacilli, and that 246 parts per million were required for cholera organisms, while 2 p c of dry lime was needed for cholera discharges, with in each case 'a few hours' 'contact When lime is added to sewage, a great number of bacteria are carried down, but are not killed, and the supernatant liquid is not sterrlised. Dr Houston (7th and 8th Reports of the Metropolitan Water Board, 1911 and 1912) found that 0.02 pc of quicklime effectively destroyed B coli in raw Thames water in from 5 to 24 hours Zinc chloride, 'Burnett's fluid, formerly in high repute as a disinfectant, is not reliable m its effect, and ranks below copper sulphate and mercune chloride The sulphate has still less energy (R W coeff 0 09)

Aluminium chloride, 'chloralum,' was classed by Miquel between zinc chloride and copper sulphate. Iron salts, especially ferrous sulphate, were at one time in vogue for sanitation, but are ineffective

Permanganates and manganates were first introduced as Condy's red and green fluids oxidisers generally, they are largely consumed by otherwise mert matters before attacking organisms, and as complete disinfection cannot be ensured unless an excess of reagent remains, evidenced by a pink colour, which is not always easily visible owing to the brown oxide of manganese produced, their use may be very costly and its execution frequently imperfect writer carried out experiments in street-watering of two similar areas of ordinary soiled asphalt roadway (a) with water alone, (b) with permanganate solution 1 in 5000 (six times the amount then customary), and examined the liquid from the surface. The permanganate was almost immediately decolourised (b) was nearly inodorous, on keeping became much less foul than (a), and the reduction in the number of organisms was 96 pc, but the survivors included a large proportion of the dangerous forms (Rideal's Sewage, 1906, 117) For polluted water Rosenau recommends addir; permanganete drop by drop till the pink colour

persists for 24 hours. 'Pinking' of wells was many years ago adopted by Hankin in India as a precaution against cholera Permanganates, used in various ways, are still being patented, chiefly for water purification.

Chromic acid, although a powerful oxidiser and instantly coagulating albumen, is excluded by its cost and poisonous and corrosive nature from the list of useful disinfectants or of pre-

servatives.

Arsenic, like cyanogen, has more effect on higher than on lower forms of life, and is the basis of valuable insecticides, such as the sheep-dip solutions of arsenious oxide or sulphide in soda, frequently associated with tar products and mootine. See Chemist and Druggist, June 11, 1904, Rep. Dept Comm B of Agric, leading to the Sheep-Dip Act of 1903, and Quibell, J Soc Chem Ind 1907, 1266 The copper compounds, Paris green' (acctorisemite) and others, are used for spraying or dusting trees, combining the actions of arsenic and copper Several organic arsenic compounds have been introduced as medicinal parasiticides (see Lancet, Brit Med J, &c, 1910 and onwards)

Arsenicals and similar organic metallic compounds are frequently selectively germicidal Salvarsan, p-p-dihydrosia, m-m-diamino arsenobenzene has a strong antiseptic action on anthrax, eryspiclas, and glanders, as well as syphilis. The selective action is more pronounced in scrum (Zeitsch f Hyg u Infeht Krankh 1914, 77)

Among the more important arsenicals may be mentioned .—

Salvarsan

Cacodylic acid and its salts
Atoxyl sodium aminophenylarsenate
Asyphil mercury ammophenylarsenate
Arrhenal methyl disodium aisenate
Aisacetin p-acetyl aminophenyl sodium
aisenate
Sommi sodium aisenate

Soamin sodium arsanilate Neosalvarsan sodium 3 3-diamino-4 4'-dihydioxy arsenobenzene methanal sulphoxylate

Some metals in the free state are antagonistic to germs. Dievert states (Compt rend 136, 707) that agitation with granulated zinc kills B typhosus and coli in a few hours, and Baeyer's method of treating polluted waters with zinc-dust, charcoal, and lime causes, according to Margosches (Leip: Monats Textilind 1901, vi), practical sterilisation Rankin (Proc Roy Soc 1910, B, 82, 78) shows that clean strips of Al, Zn, or Cu, in air-free water, do not reduce the number of col, whilst when air is bubbled through there is a great reduction after 1 hour with Cu, peroxide of hydrogen was produced, but the amounts of this and of the metal dissolved were too small to account for the germicidal effect, which is ascribed to the metal and oxygen together Metallic iron has long been used for purifying water, as in Anderson's process (J. Soc Arts Feb 14, 1896), in this case, galvanic action and the oxides formed come largely into play, but Frankland proved that the metal itself was, to a certain extent, bactericidal Old Hindoo writings direct water to be kept in vessels of copper; among modern observers,

Nageli, Galeotti, and Israel and Klingman hold that there is formed a colloidal solution which renders the water toxic to many algae and bacteria. In 1904 the U.S. Dept. of Agriculture reported that I sq. cm. of bright copper per 100 c.c. of water was sufficient to exterminate Uroglena and some forms of Spirogyra, and Kraemer showed that the same treatment destroyed colon and typhoid bacilli in 4 hours, whilst colloidal copper was quickly fatal to these organisms For household purification, he commended the use of strips of copper, about 3½ sq inch to each quart (2 sq cm to 100 c.c.), immersed in the water for 6 to 8 hours ln a similar way, Rideal and Baines (J. San. Inst. 1904, 594) obtained sterilisation of typhosus, coli, and S py aureus in less than 24 hours Bassett Smith (J Prev Med July, 1904) found that in a bright copper vessel, B typhosus was still living at 12 hours, but was dead at 24, and that of ordinary water organisms, 1020 per cc at first, only 8 per ce (none of them liquefying) were left after 24 hours He concluded that clean iron or zinc was nearly equal to copper in first effect, but soon lost the power by oxidation

Copper salts have a distinctly poisonous action on lower organisms, and are now used in many places to retard algae development early in their seasonal growth in water reservoirs Kroncko considered cuprous chloride to be the most active of the salts he treated Elbe water containing 40 to 50 thousand organisms per c c. with 50 parts per million of Cu Cl<sub>2</sub> and 20 parts of ferrous sulphate, allowed 6 hours' contact, then agitated with 10 parts of lime After settling and filtration through sand, the water was sterilised, clear, and colourless, and free from iron and copper Schumberg (Chem Zentr 1900, n 203) corroborated the sterility at the end of 6 hours The efficiency of soluble cupite salts is generally dependent on their percentage of copper (Green, Zeitsch f. Hyg 1893, 495, Rideal and Baines, lc), but the sulphate is commonly used as most convenient For disinfecting sowage effluents (and inferentially for ieservoirs), American investigations conclude that copper sulphate 'is not so efficient as chlorine compounds, is more seriously affected by carbonates, and is much more expensive' (U.S. Geol Survey, Water Supply Paper 229, 1909, 32; also see Bull U.S. Dept. of Agric No 100, 1906)

Disinfection of excreta and morbid products, even with considerable quantities of copper salts (e g 5 to 10 pc of the sulphate), is not For killing agricultural parasites, their use is well known, including that of the sulphate for soaking soeds and against potato blight, also verdigris (acetate), 'Paris green' (arsenite), copper-lime-sugar and copper-soap washes for trees, and Bordeaux mixture' (copper sulphate and lime) The last named was introduced for vines, but has been since extended to other crops. In reference to the general practice of employing the metallic compound in the precipitated or solid form, instead of in solution, the main points stated are (1) that, in the former way, it is not liable to injure the plant or contaminate its tissues; (2) that the copper should remain as an external coating, because the spores germinate on the surface: (3) that uncertainty of composition

and inequality of application are overcome by certain piecautions and definite prescriptions. See British Board of Agriculture leaflets 23 and 225; the latter (May, 1909) gives a method of preparing and using Bordeaux mixture: 'for a 3 pc solution, 3 lb copper sulphate and 2 lb. freshly burnt quicklime to 10 gallons of water. See also J Soc Chem Ind 1896, 332; 1907, Chuard Compt 1end 1911, 152, 532 (ibid 1910, 150, 839), estimating that 12,000 tons of copper are thus consumed annually in France, expresses anxiety as to the result of its accumulation in the surface soil, and with a view to reducing the quantity states that a half pe aqueous mixture of the oxychloride is as adhesive and effective as the usual 2 pc paste of the Cupile sulphate impregnation for preserving or 'kyanising' timber has been superseded by creosote oils, since the former washed out of the wood. 'Microsol,' sold as a soluble disinfectant for stables and drains, contained 75 p.c of copper sulphate with sulphocarbolate and some free sulphur dioxide.

Mercury -Mercuric chloride, 'corrosive sublimate,' has long been recognised as one of the most powerful of disinfectants, and a 1 per 1000 solution was the earliest standard for companisons Apart from cost, points limiting its utility to special purposes are: (1) the poisonous effect on higher animals and plants; (2) its precipitation by so large a number of substances, such as hard water, alkalis, and numerous salts, metals, sulphides, and many organic bodies, causing its action to be liable to great variations and inconveniences often form a pellicle over organisms without killing them, and it is inapplicable to the disinfection of sputum, excreta, and the like, from its producing a coagulum which prevents further Sodium or ammonium chloride is penetration sometimes added to increase the stability of the aqueous solution, but considerably diminishes the germicidal power The Local Government Board recommended for disinfecting purposes  $HgCl_2$   $\frac{1}{2}$  oz , HCl I fluid oz , aniline blue 5 grs (a colouring for safety), water 3 gallons (I in 962)

Kionig and Paul in 1897 showed that mercury salts followed a general rule that the germicidal power depends on the metallic ion, and is in proportion to the degree of ionisation, so that a 1 in 500 solution of Hg(1, is much less than twice as active as 1 in 1000 moreover, the chloride is more active (in equivalent solutions) than the bromide, and 4 times more active than the cyanide, which is almost non-ionised tions of the orycyanide are used in surgery, are distinctly alkaline, and only slightly precipitate albumen, a strength of 1 in 1500 is antiseptic. does not really attack instruments, and is not Mercuric iodide, dissolved in potassium ırrıtant iodide, is a powerful disinfectant, and is made up as 'iodic hydrarg' It is less affected by albuminoids and less irritant than the chloride, and is incorporated in a disinfectant soap. Mercury-zinc cyanide, 'Lister's antiseptic,' has the disadvantage of being of variable composition. A great number of organic mercurial compounds, introduced into medicine and surgery, trace their main effect to an antiseptic or garmicidal action 'Sublamin,' mercuricethylenediamine sulphate (43 p.c. Hg), is easily

soluble in water, is much less irritant than mercuric chloride, does not coagulate albuminous solutions nor precipitate soap, and is a valuable hand-disinfectant 'Protectyl' contains about 96 pc water, 1 pc salicylic acid, 0.2 pc.

mercury, and about 3 p c gelatin

Sodium hydriodic mercurobenzoate was carefully investigated as a disinfectant by Schreller and Schranth (Zeitsch f. Hyg u Infect Kranh 1911) They found that the germicidal powers could be enhanced by the introduction of halogens, methyl, methoid, or a second hydroidic mercury group into the nuclous, whilst a corresponding depression was caused by sulphonation or the introduction of an amino grouping, sodium oxymercury orthonitrophenolate 'Mercurophen' is said to have a carbolic acid coefficient by the R W method of over 10,000, to be less toxic than mercuric chloride, and to exert less than one-quarter of the precipitating power which this salt has on blood scrum (S Schamberg, J Amer Med Assoc 1917)

Silver nitrate stands next to mercuric chloride in germicidal power, and has similar limitations in its use, with the additional one of being precipitated by chlorides. The writer found that 1 in 1000 killed B coli and S py aureus in 24 hours, but a small quantity of chloride or organic matter prevented the effect Among other silver salts are the citrate, 'Itrol,' recommended as a non-irritant antiseptic dusting for wounds (Pharm Zentr 38, 460); 'Actol,' the wounds (Pharm Zentr 38, 460); 'Actol,' the lactate, for antiseptic injections 'Tachiol silver fluoride, easily soluble, and a powerful non-toxic bactericide specially useful for the urmary tract (1 in 1000 to 1 in 5000 is suggested by Paterno and Cigolani for sterilising water in a concentration of 1 in 400,000. Chiefly with the object of avoiding irritant action, incompatibility with chlorides, and coagulation of albuminoids, a large number of organic preparations of silver have been introduced, such as 'argentamine' (ethylenediamine-silver-phosphate), and the proteid compounds 'argonine' (casein), 'argyrol' (gluten), 'largin' (albumen), and 'protargol' (protein) A report to the British Med. Assoc in 1906 gives the time in minutes required to kill S. py aureus by the preparations used as · silver nitrate (1 to 2 p c), 2 to 5; protargol (2 to 4 p.c), 3 to 5; largin (10 pc), 2 to 5; argonine (5 pc), 3 to 6; whilst argyrol and 'collargol' (colloidal silver, J. Soc. Chem. Ind 1903, 315) had very little bactericidal power (Lancet, 1907, 1. 675). Many of these are destructive to gonococci and to ophthalmic disease organisms.

Osmic acid was found by Koch and Klein to be powerfully bactericidal; it is very poisonous and a strong oxidiser. The former stated that an extremely dilute solution of potassium auricyanide, 1 part  $Au(CN)_3$ , in 2 millions, checked the growth of B tuberculosis

Chick and Martin (J of Hyg 1908, 634) observe that for spores, metallic salts rank highest as germicides; with mercuric chloride, the reduction of efficiency caused by blood srum is much greater than with phenol.

## ORGANIC COMPOUNDS.

Compounds related to the alcohols. It has long been known that 'wood spirit,' crude

methyl alcohol, in a dilution as low as about 5 p.c., can kill insects and most micro-organisms, and preserve perishable organic materials owing to the tar products that it naturally contains. The alcohols, by themselves, are only efficient in a very much higher strength by coagulating albumen and (when absolute) by withdrawing water Wirgin (Zeitsch Hyg 1904, 46, 149) has tested methyl, ethyl, butyl, and amyl alcohols with anthrax spores, and S py aureus, and concludes that the disinfecting power is low, that it increases with the molecular weight, and that there is scarcely any action on dry spores It has been suggested to add alcohol vapour in steam disinfection, but Seige (Chem Zentr. 1902, 1, 130) finds no advantage Hand disinfection with alcohol is not reliable (Goenner; Harrington, Boston M J. May 21, 1903), although Hansen found that epidermal bacteria, e.g eczema, ere, when moist, destroyed in one minute by absolute, and in most cases by 50-60 p c alcohol (Centr Bakt 1907, 466) This statement is supported by A Zabludowski (Deut. Med. Wochenschrift, 1911, 405), who advised the addition of 5 pc of tannic acid. Sterility of the hands could be obtained by 2 minutes' immersion Glycerol in strong solutions (25 pc and upwards), is antiseptic, but in weaker ones it rather promotes the growth of bacteria Allyl isothiocyanate, mustard oil, has figured in a number of patents, and is capable of killing many organisms, but not economically.

Formaldehyde is found in commerce as the 40 pe solution 'formalin' or 'formol'; it is antiseptic and germicidal, and is official in the German, Austrian, Belgian, United States, and Formalin, diluted Japanese pharmacopæias 10 times, is used for embalming and for preserving bodies for dissection and museum specimens, but for antisepsis generally, a much weaker solution is efficient. The Board of Agriculture (Dec. 1906) found that fruit of all kinds immersed for 10 minutes in a solution of 3 pints formalin to 10 gallons water (= $1\frac{1}{2}$  p c CH<sub>2</sub>O), then allowed to drain and dry, keeps 10 to 21 days longer than when untreated, and that one lot of liquid can be used for a number of batches S Rideal and Dr Foulerton have . proved that 1 in 50,000 of formaldehyde (=1 in 20,000 of formalin) suffices to keep milk sweet for 24 hours, even in warm weather, without injury to health (Pub Health, May, 1899; Lancet, 1899, 1427, 1571). Apparently because experiments with larger doses showed effects on nutrition, the British Dept. Committee on Preservatives, in 1901, recommended 'that the use of formaldehyde or formalin or preparations thereof in foods or drinks be absolutely prohibited' And yet its agency in the preservation of food is as ancient as that of salt. The disinfectant and preservative power of woodsmoke is in part due to formaldehyde, which is present on the surface of smoked provisions,1 and was found by Trillat in amounts varying from 10000 to 100000 of the weight of the substance consumed, in smoke from wood and cellulose In tobacco smoke, he found 0 05 to 0 12 pc., and in soot from house chimneys, 0.28 to 0 35 p.c.

<sup>1</sup> Perrier finds zobon to zoobono in bacon, ham, sausages, and herrings; remarks that a strict regulation against formaldehyde would forbid these foods; and contends that a limit should be fixed instead of prohibition.

as paraformaldehyde Typhoid and cholera organisms were killed by products from the combustion of 2 grams of sugar passed into a 12 litre vessel, and by burning 4 kilos. of sugar in a room of 100 cub metres 'even very resistant organisms were destroyed.' (Rouguier has made this the basis of Fr. Pat 360476, 1905) With straw, the best results were obtained by burning it in a heap of alternate dry and wet layers using 18 kilos, of straw in a room of 140 cub. metres (4944 cub feet), B. coli and diphtheriae were destroyed after contact with the smoke for 12 hours; the room temperature rose to 35°, and up to 2 grams of formaldehyde were generated per kilo of straw (Compt rend 138, 1613; 139, 742, 140, 797, 141, 215; 150, 339)

H Will (Zeitsch. Brauw. 1905, 28, 330, 347) gives formalin a very high place among brewery disinfectants. He notices, like other observers (Kingzett, Slater and Rideal, Lancet, April 21, 1894), that it is more powerful against bacteria than against yeasts and mould fungi. Limits recorded as preventing development are sareina and bacteria, 0 003 to 0 031 pc, yeasts, 0 007 to 0 062 pc; moulds, 0 031 to

0 125 pc.

Advantages in formaldehyde disinfection are. (1) that it can be easily applied in the state of gas or vapour for fumigation, in the liquid condition as spray, wash, dipping, or in higher dilutions as an antiseptic, or in the solid form as its polymers, compounds, or mixtures; (2) that unlike mercuric chloride and many other agents, it is not thrown out of action by albuminous matter or by most chemicals, (3) that it is effective in strengths which are not irritant or poisonous, and does not injure fabrics or metals At quarantine stations, large quantities of bulbs, roots, nuts, fruits, &c, coming from plague or cholera-infected regions are disinfected by immersion in a 5 p c solution of formalin, which does not injure the food value and much retaids decay. In the Linley process, meat is treated with formaldehyde vapour, 'I oz formalin per cub foot of space in the chamber,' the gas after a time removed by a fan, and the meat frozen for shipment Buchanan and Schryver (L G.B Food Report, No 9) state that the gas commonly penetrates for about 20 mm under a thin layer of connective tissue, that the amount absorbed often reaches 1 in 3500, that it is reduced or removed by boiling or roasting, but that grilling seems to make it penetrate further

In surgery, formaldehyde and its preparations have been of wide service. Amongst the more important may be mentioned the following formaldehyde and formaldehyde ammonia (hexa-

methylene tetramine) compounds:-

With formaldehyde
Inorganic compounds Glyceroborate
or boroform
Alumnum sileate or dreiaform

Aliphatic derivatives Iodoform or eteasodoform Acetamide or formicin With hexamethylene tetramine
Triborate or
borovertin
Dichromate or
chromoform
Acid phosphate
or hexanitrine
Dibromoduodo
chrysoform
Sodium acetate
cystopurin
Iodoform iodoformin

With forms	aldehyde	With hexamethy- lene tetramine
Aromatic deri-	Phenol	Resorcinol
vatives	Resorcinol	Guaiacol
	Guaracol	Salicylates
	Thymol	Salicyl-sulpho-
	Cresols and	nates
	potash or	
	tysoform	
	Tar and pitch	
Carbohydrates	Lactose	Mılk sugar
January and January	Dextrin	sugar
	Milk sugars	
Miscellaneous	Starches	Silver albumin
	Casem	Camphorate
	Gelatin	
•	Dextrin Milk sugars Starches	Ü

Geronzi found that the addition of 5 pc of sodium carbonate to the 5 pc formalin which he successfully used for obstinate ear disease mereased the tolerance and enhanced the germicidal power (Arch. Ital. di Otologia, 1903).

Formaldehyde antiseptic powders for dressing are 'Amyloform' (starch), 'Proteol' (casein); 'Steriform' and 'Sterisol' (lactose), 'Glutol' (gelatin) These, especially the last, when moistened as in a wound, slowly regenerate formaldehyde. Formalised gelatine is now largely used with success instead of surgical collodions Formamint tablets, 'for infectious diseases and as a prophylactic,' are said to contain 0.01 gram of ('H<sub>2</sub>O in each, combined with milk sugar (Eng. Pat. 2672, 1906), Pharm J. 1907, in 838) M. Jones states that with B. diphtheriæ the R. W. coefficient is only 0.01 as against 0.30 for cyllin pastiles

Paraform, paraformaldehyde, trioxymethylene, or 'triformal,' is a solid polymer, only slightly soluble in cold water, and dissociated into CH<sub>2</sub>O gas on heating. It was introduced as an internal antiseptic and for disinfecting instruments, and said to be as active as β-naphthol, but its chief use is for evolving formaldehyde either alone or in mixture, for which a large number of methods have been proposed, many of them patented. Mixtures with Na, Ba, or Sr peroxide, when brought into contact with water, yield formaldehyde and hydrogen peroxide. It is stated that 'Autan' contains I part of paraform to 2 of BaO<sub>2</sub> (Lancet, 1908, 1–139), and that 50 grams mixed with water kills all organisms in a cubic metre of air bubbled through it (Fr. Pat. 366605, 1906).

Room disinfection with formaldehyde cannot be economically effected by simply evaporating the solution, under ordinary circumstances, only a portion of the substance is volatilised, the remainder polymensing and remaining behind as a solid, while reliable disinfection requires a definite large volume of gas to be produced in a short time. For this object, very numerous methods and apparatus have been devised, which may be grouped as follows:—

been devised, which may be grouped as follows:—

A. Direct production of CH<sub>2</sub>O gas, by passing methyl alcohol vapour and air over hot surfaces of platinum (Hofmann) or partly oxidised copper (Loew). The various lamps on this principle have had the faults, among others, that the oxidation is incomplete and inconstant. Kenv ood found that 1½ litres of methyl alcohol were required to disinfect a room £ 2000 cub. feet.

B Heating the solution (formalin) under pressure was used by Trillat, calcium chloride being added to raise the boiling-point In his autoclave, 1 lb formalin was required for 1500 cub feet By this method, Kenwood obtained sterilisation of diphtheria swabs, and S. Rideal with various articles infected with B typhosus, diphtheria, coli, S py aureus, and anthrax spores, found that in 24 hours all were sterilised, whilst fabrics, furniture, metals, and leather were (See also J R San Inst 1903, 508) not injured

C Formalin, mixed with glycerol, is distilled at nearly the ordinary pressure, as in the Trenner-Lee, Lentz, and Lingner apparatus The latter, according to the Russian National Health Society's Journ 1900, was mainly instrumental in extinguishing the plague at Astrachan, in 1899 Klein (report of 1902), by 3 hours' exposure in a sealed room with the Linguer generator, sterrised anthrax spores and tuberculai sputum Houston and Newman also report favourably (Practitioner, Sept. 1902)

An important point to notice is that in method (B), the CH<sub>2</sub>O is chiefly liberated at the beginning of the operation, in (C) at the end, mixed in the latter case with steam and glycerol spray, which deposits as a film on the surfaces, and assists in the penetration and disinfectant

D Formaldehyde has also been used in conjunction with acetone (Eng. Pat. 10398, 1908), SO<sub>2</sub> (Fr Pat 379998, 1906), alcohol, and

some fatty acids

E Evolution by mixing with other chemicals Dehydrating agents disengage very little CH,O gas from formalin, since most of it polymerises With quicklime, S Rideal has only obtained 8 p c of the theoretical yield (cf Eng Pat 8259. 1899, Fr Pat 323041, 1902, and U.S. Pat 790468, 1905) Carteret observes (Compt. rend. 1908, 146, 819) that 1 part formaldehyde and 2 parts bleaching powder added to 3 parts water evolve 70 p c of the theoretical yield (D R P 217941 of 1908) Potassium primanganate behaves similarly, generating great heat, U.S. Pat 885223 relates to a mixture of this Major Munson (Trans Inter Congress Hygiene, Washington, 1913, 756) has obtained 86 p c of the formalin vapoused, and S Rideal (Blackpool Congress, 1914) used a paper bag for sterilising clothes, using 40 grams per-manganate to 100 c c formalin. The action of The action of peroxides has been already mentioned

F. The simplest process is heating paraform by a small flame (avoiding ignition), when it melts and is resolved into CH<sub>2</sub>O. The 'Alformant,' or Schering lamp, uses 1-gram tablets placed in a perforated cup, and it is directed to use 10 tablets per 1000 cub feet with 6 hours' exposure S Rideal found that this quantity killed B coli, typhosus, diphtheria, and S py aureus in 20-24 hours, and that double the quantity in the same time killed moist, but not always dry anthrax spores He recommended spraying with  $\frac{1}{2}$  pc solution of formalin before using the lamp, as the gas is evolved with insufficient moisture. The latter defect is remedied in the 'hydroformant' lamp, in which 12 ozs. of water in an annular vessel is simultaneously evaporated. Rosenau considers this method of generating CH<sub>2</sub>O useful fc the disinfection of closets and small enclosures of in this way.

less than 100 cub feet, with not less than 12 and preferably 24 hours' exposure, and not less than 2 ozs. of paraform per 100 cub feet (about 2 grams per cub metre). Klein, Houston, and Gordon (London County Council, 1902) record that with 23 grams of paraform per 1000 cub feet, B typhosus, diphtheriæ, pyocyaneus, and S py aureus were killed in 5 hours, but not anthrax spores nor (with certainty) tuberculosis in sputum. It will be noticed that the time is not even that given in the directions issued with the lamp, and the humidity and temperature of the room are not stated Allan, Newman, and Cribb had previously obtained generally satisfactory results with 10 grams paraform per 1000 cub feet, under moist conditions (Brit Med J Aug 13, 1898) Kenwood's experiments pointed to an average of 25 grams per 1000 cub feet being necessary Werner and Bonhoff (Berl Klin Woch 1904) find that tubercular sputum is certainly disinfected by 5 grams per cub metre (141 grams per 1000 cub. feet) acting for 7 hours, which is double the quantity Flugge first recommended In Breslau a large number of formaldehyde disinfections are carried out for scarlet fever and measles. and the disease has in no case recurred in the same room

Spraying with the solution should be the invariable practice except with the large CH,O generators, which are capable of filling the room also with steam and spray When spraying is used alone, there is a danger of not penetrating into crevices, and the solution must be of higher strength, Dr Mackenzie's extensive experience in the disinfection of houses and hospital wards, proved that 1 p c CH2O (4 ozs formalin per gallon) is attended with success, he adds 5 ozs glycerol per gallon of formalin to prevent too-rapid drying Thresh and Sowden in 3 to 4 hours sterilised diphtheria, typhosus, cholera, pyoq aweus, and produgosus with ½ p c CH,O, but pyocyaneus (on whitewash) only with 2 p c, they, therefore, recommend the latter strength

Where the space has to be occupied soon afterwards, any residual formaldehyde can be neutralised by volatilising ammonia The product, under the name of Urotropine, is a serviceable antiseptic in cystic affections, and is administered to typhoid 'carriers' to destroy the bacilli of the disease in the urine Urotiopine is the basis of a large number of antiseptic

preparations (see above)

Disinfection of excreta is attained by mixing with 3 to 5 pc of formalin and keeping in a closed vessel for at least an hour Rosenau states that faces are rendered sterile at the end of 10 minutes by an equal volume of a 4 pc solution of formalin Houston prescribes 1 to 2 pints (according to bulk) of 5 p c formalin, acting for at least an hour (Practitioner, Sept. 1902).

Formaldehyde, as an insecticide, is much inferior to SO2, although Brough (J. Mass. B. of H. March, 1898, 51) observed, in his disinfections with the former gas, that all the flies were invariably killed, and generally the bed-bugs. It

13 an excellent deodorant.

No acidity is produced by continuous spraying or distillation of formaldehyde, therefore it is not readily converted into formic acid

Formic acid.—We have discussed the germicidal action of acids generally in an earlier section of this article. Formic acid has this power to a considerable extent, and is a strong antiseptic Kitasato's results, as summarised by Horrocks, gave acetic acid a slightly higher activity in nutrient media, stating that with B typhosus the amounts pc in cases of (1) growth, (2) growth restrained, (3) no growth, were respectively formic acid, 0 22, 0 278, and 0 356, acetic acid, 0 2, 0 225, and 0 3, with Sp cholerw, formic, 0 11, 0 167, 0 22, acetic, 0 1, 0 153, 02. But very different results are obtained in water and most organic solutions. The writer has found that typhosus is killed by 0.5 p.c. of formic acid in less than 15 minutes and by 0.1 p.c. in about 30 minutes, and in 1907 he made the R W typhosus coefficient of formic acid many times higher than that of acetic also found the preservative action to be higher with vegetable substances, I pe of formic acid was equal in effect to 5 pc of acetic (time of trial, 2 months) Raw fish were mounted in 053 pe forme, in 5 pe acetic, and in 03 pc boric, acids, heated for 20 minutes to 80°, and the jar closed with cotton wool, in a week, the last was brownish, disintegrated, and uncatable, though not distinctly putrid, while the first two kept sweet and of natural characters for 3 months Experiments with bread-pulp infected with a mixed mould-growth proved that 01 pe of forme acid entirely inhibits the growth for several days, even under the most favourable conditions of culture But in the destruction of the spores, formic and is less effective than formaldehyde Lebbin (Chem Zert 1906, 30, 1009) effectively preserved different classes of foods by 0.15 p  $\epsilon$  of formic acid, and others have proved that in the quantities mentioned above it is physiologically harmless, the taste is more pleasant than acetic acid, and the odour is not noticeable B H Smith (J Amer Chem Soc 1907, 1236) preserved vegetables with formic, with salicylic and with benzoic acid, in unprotected jars Under conditions in which the untreated samples became sour and mouldy in 2 days, 01 pc formic acid preserved them for 7 days, 0 3 p c for 12 to 18 days, 05 pc for 25 days, 07 pc and 1 pc for 45 to over 85 days, and 2 pc for over 85 days He states that 0 1 p c of benzoic acid kept his sample for 85 days, whilst with the same quantity of salicylic acid it was sour in 9 days. In ordinary preserving practice, as we have seen, the articles would have remained good for the longer time with the smaller quantities of formic acid, which, even if it must be applied in a somewhat higher proportion, is from physiological and physical reasons preferable to salicylic or benzoic The chief formic acid preservatives used in commerce, mainly for fruit preparations, are 'Werderol' and 'Fructol' (10-14 p.c. formic acid, '1-1} p.c. of the liquid to be added'); and 'Alacet' (made synthetically from CO and NaOH at the Nitritfabrik, Kopenick, about 50 to 60 pe formic acid, 'use 03 p c. of the liquid') The result in the three cases would be about 0 14-0 18 p c of the acid In an investigation of these articles by Croner and Seligmann (Zeitsch. f. Hyg. 1907, 56, 387), they find that inhibition of moulds, yeasts, and acid-forming organisms

begins at 0 15 p c formic acid, that sterilisation is effected in 24 hours by 02 pc., and in 10 to 30 minutes by 1 p c

Only the free acid is active against organisms. although the salts do not favour the growths as do the alkaline acetates Sodium formate helps the solubility of many antisepties without lessening their power, therefore figures as an adjunct in a number of patents

Acetic acid is one of the most anciently used of preservatives, 'aromatic vinegar,' a more or less concentrated acid containing essential oils, had a partially-justified reputation against infection. The writers find that  $B \ coli$  is killed by 5 pc HA in 5 minutes, by 25 pc in 15 minutes, while with 05 and 01 pc it is alive after 40 minutes Pyroligneous acid, or crude wood vinegal, owes it's antiseptic power chiefly to the presence of creosote and formaldehyde

Acetyl peroxide and benzoyl-acetyl peroxide are stated by Freer and Novy (Amer Chem J 1902, 27, [3] 161) to be strongly germicidal, since solutions corresponding to 0.037-0.074 p.c. of the former, or 0.056-0.112 p.c. of the latter, =0 005-0 01 pc of active oxygen, destroyed within a minute all known disease-producing bacteria, and even spores of B mescutericus, which are not killed by 5 p c phenol, whilst  $H_2O_2$  solution containing 0.05 p c of active O was without action on many bacteria even in Benzoyl peroxide had no oxidising action, and was without effect on bacteria solution of acetyl peroxide was introduced under the name 'acetozone,' but the title is now applied to a more convenient and stable mixture of the benzoyl-acetyl compound with infusorial earth, said to be useful in typhoid, dysentery, and cholera (Lancet, 1904, n. 1160, Brit Med J 1907, 1 634)

Propronic acid has been tested comparatively by Duggan (Amer Chem. J 7, 62) with the highly resistant B subtiles He gives as the strengths necessary for killing tormic acid, 7 pe; acetic acid, 9 pe, propionic acid, 12 pe (numbers proportional to the molecular weights). According to A C Jordan (Practitioner, Sept 1902, 297), 01 pc of butynic acid in broth tubes infected with vigorous B coli or typhosus caused them to remain sterile, S pyog aureus required 0.2 p.c. Valeric acid is a teeble antiseptic, but has been introduced along with a little crossote, as an internal antiseptic called 'geosote

Oleic acid, the fats, 'Lanolin,' &c, have but slight antiseptic power in themselves, but when applied to the skin may prevent infection by the exclusion of germs Reichenbach (Zeitsch f Hyg u Infect 1908, 59, 296) showed that the alkali salts of the palmitates had the strongest bactericidal action of the soaps investigated. The potassium salts of the saturated fatty acids are comparatively strong bactericides, but those of the unsaturated acids, with the exception of elaidic, are not Vicario, however, observed that the fatty matters sometimes aheady contain germs, and Baldas (Giorn R S Ital d'Igiene, Feb. 1901) proved that B coli and typhosus, S pyog aureus and albus, retained their virulence in oils for 2 months; therefore ointment bases require to be s'errlised by heating to 100°, or preferably to 120°, as various bacilli survive ir oil far longer than in aqueous solution. In fatty media antiseptics and disinfectants, as a rule, are much less active than in water, hence the ineffectiveness of carbolised oil as shown by Koch and Breslauer Lanoline and cold cream contain water, so can enable agents in aqueous solution to come in contact with micro-organisms Some metallic cleates are parasiticide. cupric cleate in ringworm and favus (Lancet, 1907, 1510); mercuric cleate for pediculi, zinc cleate is said to cure chronic eczema.

Vegetable acids have the general power of inhibiting bacteria which prefer a neutial to an acid medium, but their effect is much less than that of mineral acids, and their solutions quickly become mouldy in warm weather. Citric acid was formerly recommended for sterilising water: Dr. Christmas in 1892 stated that cholera organisms were killed by 0.08 p.c. and typhoid by 0.10 p.c. Kitasato's figures for tartaric, citric, and malic acids added to neutral media containing typhosus are in p.c.'s growth, 0.338; growth restrained, 0.384, no growth, 0.476. Succime acid is slightly antiseptic. 'Alphogen' or 'Alphozone,' succinyl peroxide, is said to be a powerful germicide (Lancet, 1905, 1.367).

Aromatic acids.—The introduction of —COOH groups or aliphatic acid radicals into the benzene ring yields acids of relatively low antiseptic power, increasing with the molecular weight of the extense acid.

the entering acid Benzoic acid is strongly antiseptic in solution and vapour, and to a rather less degree in its soluble salts  $\,$  It dissolves in about 400 parts of cold water, and the saturated solution kills B typhosus in a few minutes  $\,$  When molecular proportions, i e sodium benzoate 0 026 p.c. potassium benzoate 0 029 p.e., benzoic acid 0 022 p.e., are separately mixed with milk, the writer observes that all retard the souring, the acid more than its salts, but the effect comes to an end sooner than with inorganic salts like the fluorides, since benzoic acid is itself decomposed by some organisms (Analyst, 1907, 32) Not being poisonous, it has been much used as a food preservative, particularly in countries where salicylic acid was forbidden. Following a report of the Referee Board of the US Board of Agric that doses of benzoates under 0.5 gram per day were not injurious to health, and that even 4 grams per day did not act as a poison, the US. Board, in Pamphlet 104, 1909, have announced that they will make no objection to the use of sodium benzoate in food, provided it is plainly labelled on each package, both as to presence and quantity. The taste and odour are difficult to mask. It has long been used as an antiseptic in surgery, e g. in 'Friar's Balsam' (tinct benzoini co) Benzoic aldehyde is official in the US Pharmacopæia, and in the form of commercial oil of bitter almonds owes its parasiticide power in ointments mainly to the hydrocyanic acid it contains, which, of course, necessitates care in Benzoyl-sulphonic-imide, 'Saccharin' 'Gluside,' has some preservative power, and has been given internally to stop decomposition of urine in cystitis, but its use for sweetening is less on account of this power than because it

is itself unfermentable.

Salicitic acid.—The saturated aqueous solu-

tion contains 1 in 500. tested with typho pyog., aureus, and anthrax, Westcott found t in 3 hours only the former was killed. A food preservative, it is about equal to benz their relative activities varying under differ circumstances Its physiological effect is n distinct than that of boric or formic acid formaldehyde in the small quantities requi but there is no clear evidence that it can injury to adults if the amount is limited British Depart Committee's suggested rest tion is that it 'be not used in a greater propor than 1 grain per pint in liquid food and 1 g per lb in solid food,' or 0 0114 and 0 014 respectively, which practically would amoun a prohibition, since moulds and ferments only reliably inhibited by 0 1 pc, the amo commonly employed, though often exceed Addition of the acid to foods is forbidder France, Austria, and some other count Traces are present in a large number of fr (Analyst, 1903, 149). Dr Linsa G. Andei used extensively a saline gelatined salicylic : paste containing 2 p c of the acid as a wo dressing with satisfactory results Compoun Compoun with an equal weight of boric acid it is used as a first dressing under the name Borsul 'Salol' is phenyl salicylate, an ternal antiseptic which is decomposed in duodenum into phenol and salicylic a According to Lowenthal, when mixed a pancreatic juice, it rapidly kills Sp cho (Compt. rend 107, 1169) 'Salophen,' aminophenyl acetyl salicylate, is said to hav 1ather stronger antiseptic power (Practitio March, 1907). A large number of derivat of salicylic acid have been introduced, and m of them patented

Cinnamic acid, at 2 p c, prevents the glo of bacteria, and at 4 p c. destroys them. I contained in Peru and storax balsams, where parasiticide in skin diseases. Fr 371091 preserves butter by enclosing it in wrapper impregnated with cinnamic acid, where the prevents butyric fermentation. Acetylomaric acid, "Tylmarin," is another internal a septic, said to have a R W coefficient of The use of halogen substituted hydrosic act especially 6-chlor-3-hydroxy-p-toluic acid, been suggested for disinfectant purposes Soc Chem Ind 1914, 33, 807)

Tannin is antiseptic to proteids through property of forming with them insoluble c pounds such as leather, but none of the tai extracts have even the power to preserve t own solutions.

Phenols and phenoloids. The introduc of aliphatic side chains into the nucleus of phraises the germicidal activity; the o- and positions for the entering groups have a gre influence than the m. Morgan and Co-found the following CA coefficients for t cresols, o=21, p=24, m=20. Whilst methol trimethyl methoxy phenol is said have a RW. coefficient of 40 when properulsified.

For general purposes of disinfection, 'car' fluids, having as their chief active stituents bodies allied to phenols, are rextensively employed. They are usually cayrupy liquids smelling of coal tar, and maroughly divided into two classes: (1) t

which on the addition of water form a more or | less clear dilution; and (2) preparations miscible with water but producing an emulsion. former usually consist of saponified tar-acids: the liquor cresoli saponatus, Ph G, consisting of equal parts of potash soap and crude cresol heated together, is of this nature. In the manufacture of these fluids, potash soap, glycerol, alcohol, and other agents are employed to aid the solubility in water. With the second group, the mostly insoluble, active ingredients are emulsified by the aid of soda, soaps, resin, neutral oils, or gelatin

Polyhydric phenols and naphthols. antiseptic powers of the di- and trihydroxy-

benzenes are less than those of phenol.

Resorcinol, m-dihydroxybenzene, is a powerful medicinal antiseptic, and a 1 pc solution kills most organisms 'Andeer's lotion' contains 40 grains in 1 oz 'water 'Anusol' is a bismuth iodo-resorcin sulphonate. Quinol or hydroquinone, p-dihydroxybenzene, is said to be a stronger antiseptic than the last, and has been suggested for typhoid in doses of 3 to 8 grains. The ortho-compound, catechol or pyrocatechin, is also antiseptic. Duggan finds (Amer. Chem J. 7, 62) that the amounts of the three isomerides required for preventing development of B subtilis in broth are ortho-20, meta-25, para- 30; phenol being 20 and pyrogallol 15

Morgan and Cooper (8th Int Cong App Chem. 1912, viii cl) give the following figures for the carbolic acid coefficients of the =0 48,

meta-=0 29, para-=1 1.

Pyrogallol, tri-hydroxybenzene, is antiseptic, and a 3 p c solution kills most bacteria, particularly aerobes, but it stains, is unstable, and very

poisonous

Guaracol (methyl-catechol) and creosol(methyl-guaiacol) are the active ingredients in wood creosotes, and have been widely useful as antiseptics and parasiticides, especially tuberculosis. Guaiacol has an antiseptic power (but not bactericidal—see Table) 21 times that of phenol, and a ½ to 1 pc solution destroys B. tuberculosis in 2 hours (Marfori, Ann di A one in 150 solution of (Morson's) Chim 12, 3) creosote kili pyog aureus and typhosus in 3 hours (Westcott) 'Little's Soluble Phenyle' contains wood creosote, and Corfield's experiments give it a high value for wounds and against Kenwood and Hewlett (J R San Inst 1906, 13) make its R W. coefficient 1.2

Beta-naphthol is official in the British (1898) and in a number of other Pharmacopæias, the dose in the former being 3 to 10 grains in a cachet for internal antisepsis, especially in typhoid and in cholera as a preventative. Its solubility in water is increased by boric acid Schneider observes that naphthols dissolved in alkaline carbonates have greater disinfectant power than alkaline naphtholates, the former solutions contain naphthol in the free state. He states that staphylococci and typhoid are killed in a short time by 05 to 10 pc of \$\beta\$-naphthol alkaline carbonate solution (Zeitsch. Hyg. 1906, 52, 534)

Ointments containing 10 to 15 pc are containing and psoriasis 'Betol,' efficient in scables and psoriasis β-naphthol salicylate, is less soluble than naphthol, and is now seldom used. 'Alphol' is its analogue from a naphthol, and is very similar.

A number of easily soluble compounds have been introduced, such as the \$\beta\$-naphthol sulphonates, 'Asaprol' or 'Abrastol' (calcium), and 'Alumnol' (aluminium). Helbing states that an aqueous solution of the latter, 1 in 250, prevents all growth of gonococci, pus cocci, and allied bacteria Bechold (Zeitsch angew. Chem. 1909, xxu. 2033) finds the odourless and nonpoisonous halogen-substituted naphthols more powerful than any disinfectant except HgCl2, and that tribromonaphthol kills staphylococci in dilutions of 1 in 250,000

Dihydroxynaphthalene The 23-substituted compound has a relatively high germicidal activity, with a ('A coefficient of 44 The 2.7- derivative has a coefficient of 2.8

Amidobenzene derivatives. The introduction of -NH<sub>2</sub> groups into the ring produces substitutes of low germicidal activity

Aniline is open to the same objection as the nitro- derivatives; it has a carbolic acid coefficient of 0 57, whilst o-toluidine, the next homologue, has a coefficient of 100, and m-toluidine one of 1 30

The dyestuffs. A number of amiline dyes are germicidal, owing to their power of penetiating and combining with living tissue. This action is very variable, but it is stated that I pe prevents development of organisms, whilst I in 500 to 4000 retards their growth

The investigation of the parasiticidal powers of the dyestuffs was conducted by Ehrlich and his pupils in their search for compounds containing strongly developed parasitotropic groupings, but only with feeble organotropic groupings It was found that tetrazo colours derived from naphthalene-3 6-disulphonic acid, basic dyes of the triphenylmethane series, and certain azocolours were equally effective Symmetrical colours were found to be less organotropic than unsymmetrical ones

'pyoktanın,' Methyl violet or violet, certain auramines, safranine, methylene and toluidine blue have been used in surgery and in malaria Malachite green, injected into the blood, kills trypanosoma in 48 hours (Birt Med J 1904, ii 1449, 1645, Deutsche Med. Woch Leipzig, 1906, 21, 463) Methylene blue, with atropine and benzoic acid, according to Lelean (Lancet, May 3, 1902), was the only drug he found of use against Bilharzia.

Derivatives of acridine, especially acriflavine-3.6-diamino-10-methyl acridinium sulphate and proflavine-3 6-diamino acridine sulphate, have been shown to be somewhat superior to malachite green when used for wound dressing Originally prepared for Ehrlich, their selective action on bacteria was first tested at the Middlesex Hospital, London. For wound dressing, the derivatives are made up as soaps, gelatine, or starch mucilages

Linked benzene derivatives. Both the chemical constitutions and the germicidal activities of the higher boiling tar acid fractions are as yet unknown. The direct linkage of two phenoloid groupings or linkage through =CH<sub>2</sub>, =CHOH, =CHOX where X is an aliphatic group, in some cases elevates the germicidal powers. Linkage through =CO or =SO<sub>2</sub> groupings is said to depress the coefficient.

Aromatic nitro- and halogen substituted compounds, such as nitro-benzene, have often been proposed as disinfectants, but are precluded by their odours, poisonous action, and sparing solubility. Trinitrophenol, 'pieric acid,' is rapidly fatal in saturated solution (about 1 p c) to bacteria and spores, but causes yellow stains, is irritant and very poisonous Cotton wool, soaked in the solution, is used very successfully in the aseptic treatment of burns, hence it is recommended by H M Inspector of Explosives to keep handy for this purpose The R W. coefficient is about 6 Potassium dinitro-orthocresol, 'Antinonnin,' 1 in 400, is destructive to all common injurious parasites, and does not injure plants.

Trichlorphenol has been used at Petrograd as a disinfectant It is said to be 25 times

stronger than carbolic acid

Hydrocarbons.—Petroleum, mainly consisting of hydrocarbons of the parafim series, has long been a well-known insecticide, and is mentioned by Marco Polo in the 13th century as used to kill acarı ın mange

For protecting burns and wounds from exposure, various substitutes for 'ambrine,' consisting essentially of paraffin wax or asphalt with a fatty oil have been suggested (BMJ 1917, 789)

A J Hull introduces into these burn dressings 'scarlet red,' 'bulliant green,' 'flavine,' and dichloramine-T on account of their antiseptic

Naphthalene is feebly antiseptic, but is not a disinfectant It is useful by its odour to drive

away domestic insects

It has been stated that articles are dismfected by dry-cleaning with petroleum-benzine, or by benzenc Farrel and Howles (J. Soc. Dyers, 1908, 24, 109, 166), however, find that acting on infected threads for 61 hours, neither benzene (C<sub>6</sub>H<sub>6</sub>) nor petroleum-benzine killed typhosus, whereas carbon tetrachloride killed it in 30 minutes, whilst 'benzine soap,' dissolved in benzine, has no germicidal power

Ether is used for sterilising catgut

Essential oils and camphors. Perfumes, chiefly composed of terpones and their oxidised products (eg the camphois, menthol, thymol, and eucalyptol), have a limited sanitary value All of them are very sparingly soluble in water, but impart to it a more or less antiseptic effect Early estimates of their power have not been concordant, chiefly owing to their having been tried variously in aqueous or alcoholic solution, or in suspension A saturated solution of thymol in water (0 3 pc) arrests fermentation and putrefaction Peppermint (containing menthol and menthane) has had a high reputation, and a number of recipes for 'plague water,' have it as a basis The use of hops in brewing partly rests on the power of the essential oil to check fermentation 'Terebene,' made by the action of H,SO<sub>4</sub> on terpentine, was once reputed as a disinfectant, but now hardly figures except as terebene soap Among other artificial products from turpentine are terpin hydrate (said by Colpi to arrest the growth of tubercle bacilli in 0 25 pc. strength), and terpineol (anthrax is stated to be killed by 1 p.c., and staphylococci by 10 p.c in 60 minutes. (H. Maix, 5. infra, also D R P 207576, 1906))

Camphor is much spoken of as a personal

the vapour to sterilise the air. It is an ingredient in many remedies for cholera and diarrhoea, and in several 'block disinfectants' such as 'Sanoscent,' 'Camphortar, &c Eucalyptus globulus and other species which have long been famous as antimalarial, are largely planted on the Continent in marshy districts. The essential oil is decidedly germicidal, having a carbolic acid coefficient of 12, and has been given with some success in typhoid, and also against tape- and threadworm.

Pinol is a liquid of similar character from Pinus pumilo H Marx has investigated the action of various perfumes on anthrax and S pyog aureus (Centr Bakt 1903, [1] 33, 74). Kobert (Chem Zentr 1907, 1 419) finds that the terpenes have, by themselves, in general only a weak antiseptic action The proportions needed for sterilising exceed the solubility in water, and can only be reached by a spirituous solution or

an emulsion

But it has been recognised, since Schonbein pointed out the fact about 1840, that essential oils have a disinfectant power conferred on them by their slow oxidation in the presence of an and moisture, whereby ozone is produced, that as long as air and oil are present, the ozone is continually regenerated and absorbed, forming oxidising and antiseptic products, which dissolve in the water, along with peroxide of hydrogen This was the principle of the 'Sanitas' patent 274, 1876

Pyridine, quinoline. and related bodies. Pyridine is volatile, mixes freely with water, and is strongly insecticide, but has a very unpleasant odour, which has militated against its use in horticulture. Tobacco smoke, indeed, owes its powers, not to nicotine, which is almost entirely decomposed by the heat, but to the products pyridine and its homologues, which are formed Tessman destroyed the bacilli of cholera and pneumonia by tobacco smoke passed for 10 to 30 minutes through a tube containing infected nutrient gelatin. Wynter Blyth killed the bacillus of nasal catarrh by 1 pc solution of pyridine and its homologues (from bone oil), or by tobacco smoke There is proof that smokers and workers in tobacco enjoy comparative immunity from epideinics, but pyridine inhalations (eg in asthma and whooping cough) require care, as it is a cardiac depressant The above experiments confirm the general view that disinfectant preparations containing the basic constituents of coal tar are to be preferred to those which only include the phenols

Tobacco juice, of course, contains nicotine, and in France is sold as an insecticide under an official guarantee as to its nicotine strength In England, a solution of the pure base in water is sometimes used as more definite Richard's cakes for fumigation are nicotine and camphor. the writer found a sample in 1909 to contain 1 17 p c of nicotine and to be completely volatile

at a gentle heat.

Chinosol is potassium oxyquinoline sulphonate  $C_9H_6NO$  OSO $_3K$ , a vellow powder easily soluble in water to an acid yellow solution, which does not stain, has only a slight aromatic odour, is permanent on keeping, is not poisonous, and does not coagulate albumen Cumphor is much spoken of as a personal has been patented as a disinfectant, and more prophylactic, but there cannot be sufficient of especially as an antiseptic. Klein compared its action on S pyog. aureus, B coli and anthracis with that of earbolic acid, and reports that a solution of the latter of 1 in 20 strength is required to kill aureus or coli in 5 minutes with certainty, whereas chinosol does this in a strength of 1 in 150 Anthrax spores are not appreciably affected in 48 hours by 5 to 6 p c carbolic, whereas chinosol of 1 p c acts germicidally on them in 5 minutes. In the writer's laboratory, B coli was not sensibly affected after 4 hours by 1 in 1000 chmosol, whilst 1 in 2000 of HgCl, produced sterility under the same conditions in 15 minutes, so that it is difficult to explain the much higher powers that some observers have found for chinosol Thresh and Sowden state that chinosol spray of not less than 1 pc, is a reliable disinfectant (J San Inst Jan 1903) Houston did not succeed in killing B coli in excreta by 1 in 75, and on linen by 1 in 62 (Practitioner, 1902, 331) On account of its acidity, it acts on non and steel Alkaline solutions interfere with its power the writer showed that a soap containing 5 p c of chinosol was inferior to a 0.5 mercuric iodide soap F G Muller reports encouraging results with chinosol in leprosy (Zeitsch Oest Apoth Vei xxxv 858) 'Cresochin' (Pharm Centialhalle, xxxvii 247) is said to be a neutral sulphonate of quinoline and tricresol, which is recommended for disinfecting instruments, as it does not darken the steel

Ichthyol --- Various derivatives from sulphurated tars and fossilised fish or petrolcums have been suggested for mild disinfection and the treatment of skin diseases Ichthyol ammonium ichthyol sulphurated has been compounded with various substances, such as albumen, formaldchyde, and metals, such as silver and mercury, under a variety of names

Quinine, which has a quinoline nucleus, probably owes its action against fevers as much to its antiseptic power as to its effect on the

nervous system

A Raal (Pharm Zeit 1900, xlv 569) asserts that strongly fluorescent bodies much increase the action of sunlight on organisms, whilst they are almost mert in the dark. A 1 in 20,000 solution of acridine killed infusoria exposed to sunshine in 6 minutes, in the dark they survived for 24 hours Similar results were noted with quinine sulphate and with cosin inference is that the effect is caused by the fluorescent rays Fluorescein has been patented as an antiseptic by Turpin, chiefly for toilet preparations (Fr Pat 346363, 1904) The biomofluoiesceins, particularly eosin, are found to have greater power. He claims a large number of other phthalein compounds as disinfectants

Certain faults of many antiseptics and disinfectants in the free state have led to an immense number of attempts to improve them by converting them into various derivatives Although the effort to produce new proprietary substances, has given rise to a multitude of often misleading titles, many of these preparations have been proved to have utility The modifications may be generally classified as follows

1. Metallic salts.—Compounds with the alkalis or earths are generally less active than the free substance Other metals have been introduced, with the object of adding their special properties, as Al for astringency, Zn, Cu, Ag, and Hg for developing germicidal effect; but it must be remembered that a combination of this kind, though it may have some lateral advantages, is commonly weaker than either agent singly The same remark applies to the two next

2 Substituted compounds —Those with iodine are exceedingly numerous, including a large number which have been improvements on iodoform when it was unduly valued. Where organic radicals (basic like methyl or ethyl, acid like acetyl or benzoyl) are introduced, the physiological properties are usually entirely altered

3 Esters, sulphonic, and other colligated compounds which, while less poisonous or irritant than the parent bodies, slowly yield the latter by gradual decomposition Several of

these have proved of great value

already indicated, the proprietary articles of this nature bearing fancy names are very numerous Since the Privy Council Orders of 1900 and 1902, the uncontrolled sale of liquids containing upwards of 3 p.e. carbolic acid or its homologues is prohibited, and the germicidal value of the more insoluble and higher boilingpoint tar acids has been recognised generality of commercial disinfectant fluids are now practically free from carbolic acid, and, in comparison with the latter, are non-poisonous These disinfectant preparations should not be employed to destroy resistant spores without prolonged exposure, according to the nature and amounts of active ingredients present, they are usually germicidal with non-sporulating organisms in dilutions far greater than is the case with absolute phenol. However, the strength of these proprietary articles varies considerably, and beyond the requirements of germicidal value specified in contracts with various authorities, there is at present, unfortunately, no legal control over the efficiency of disinfectants as sold to the public. There are instances of so-called disinfectants which enjoyed great popularity and even were advocated by the medical profession, yet possessed little or no germ-killing power under the conditions in which they would be employed

Since the adoption of the R W test, there has been a steady rise in the efficiency of com-

mercial disinfectants

The Mechanism of Disinfection -King and Paul, in 1897, first drew attention to the relationship between germicidal action and the ionic concentration of certain germicidal ions such as mercuiion, argention, hydrion, and formulated the rule that the germicidal activity of a solution was proportional to the ionic concentration of the germicidal ions recent work, such as that of Kitasato, Horrocks, and Winslow, on the common acids, and more especially the development of the organometallic group of disinfectants, indicated that this generalisation was by no means correct Ehrlich and his pupils attributed to disinfectants the possession of at least two specific properties, the presence of an organotropic and a parasitotropic grouping indicating the 'affinity' of the germicide to living tissue and the parasite or bacterium respectively

Neither of these theories will serve to

explain the phenomena observed in disinfecting | with ionised and unionised solutions, with emulsions and other colloids. Bacterial sus-censions must be regarded as suspensoids coated by, or containing, an emulsoid, and are o be classed as definite colloidal systems

The reactions between agglutin and the bacteria, similar to the reactions between toxins and antitoxins, resemble in many respects weak base, a view advanced and supported by Arrhenius ('Immunochemie'), or the usual absorption phenomena observed at colloidal surfaces. Although experimental evidence (notibly that of Marsden, Nyman, and Chick, sarried out at the Lister Institute) is not suffinently accurate to distinguish between the two, ret the other general characteristics of bacterial suspensions lead us to infer that absorption is the first stage of disinfection We can thus distinguish between the following types of action .

(1) Absorption of cations (ionic disinfectants) with simultaneous anion absorption, which may affect the activity

(2) Absorption of substances in solution, e g

phenol.

(3) Mutual action between bacteria and other colloids, such as emulsions (tar acids) or suspensoids (colloidal metals) The next stage, common to all, is penetiation and chemical attack by the absorbed disinfectant. It is evident that the germicidal activity of a substance after absorption is determined by its penetrative or intradiffusive powers as well as its chemical reactivity or 'parasitotropic' properties. The enhanced germicidal powers of the chloramine is doubtless due to their possession of a reactive -NH<sub>2</sub> group, but it is by no means clear that dyestuffs are germicidal on account of their possessing certain parasitogroupings or that their activity is not due to a medium diffusivity accompanied by gradual decomposition into germicidal constituents It appears probable that more light will be shed on this somewhat complex subject by a close study of the phenomena of absorption and the mechanism of intradiffusion of substances in colloidal material rather than by the development of the concept of ionic activity or postulation of certain parasitotropic groupings or valencies endowed with exceptional and unexpected properties

Testing of Disinfectants -Before 1881, when Koch introduced the use of pure cultures, tests depending upon the prevention of putrefactive changes in various organic infusions were mainly employed, and to a great extent the antiseptic or preservative value was noted rather than the actual germ-killing power (see first edition of this dictionary) Since then, an increasingly large number of investigators have worked upon practical methods of testing With the phenolic disinfectants, valuations were made upon the total amount of tar acids present, but the widely differing activity of these bodies, which is also modified according to their physical condition on dilution (in solution, tricresol has only one-third the germicidal value of that in emulsion), has led to the general adoption of bacterio-

logical tests

In the 'hread method (Koch), silk threads

are impregnated with the test organism, as exposed for a definite time to the action of the diluted agent, and then, after the removal adhering disinfectant, sub-cultures are inoc lated from the threads to determine the surviv or death of the organism (v. Lancet, July 3 1900; and J R. San Inst. 1907, xxviii). place of threads, small garnets can be employe (Kronig and Paul), stones of uniform size beil coated from cultures of the organism, and adde directly to the disinfectant, sterility beindetermined by inoculating sub-cultures intervals (Zeitsch angew Chem 1901, xiv. xv The chief points to be considered are (1) time (2) age of culture, (3) choice of mediur reaction of same, (4) temperature of incubation and (5) of disinfection, (6) variation in vit resistance of same species and (7) of differe species; (8) proportion of culture to disi In order to facilitate the compariso of disinfectants, Rideal and Walker (J. R. Sa Inst 1903, 424) suggested their carbolic ac coefficient method as the basis for a standar In this test, which has been extensive adopted in England and abroad, the efficience of the disinfectant is expressed in multiples absolute phenol performing the same wor It is briefly as follows: add 5 drops of a : hours' blood-heat culture of the organism both to 5 cc of a particular dilution of tl disinfectant in sterile distilled water, shall and take sub-cultures with a platinum loc every 21 minutes up to 15 minutes; Incuba sub-cultures at 37° for 48 hours Allowu 30 seconds for each operation, four differen dilutions of the disinfectant with one carbol control are tested against the same culture ar under strictly comparable conditions example.

Minutes 
$$2\frac{1}{2}$$
 5  $7\frac{1}{2}$  10 12;

Disinfectant 1.500 + - - - - -

,, 1 550 + + - - - -

,, 1 600 + + + - - -

Carbolic acid 1 110 + + - - -

Carbolic acid coefficient  $\frac{550}{110} = 5$ 

This test has for many years been official used by the War Office, Admiralty, India Office and other Government departments at hon and in the Colonies It is impossible in ar one test to include the greatly varying condition of practical disinfection, but numerous modific tions of the RW coefficient test have bee suggested, mainly with the object of testing the presence of considerable quantities organic matter. Sterilised sea-water, urin sewage, starch, and gelatine emulsions, nutriei broth and milk, have been incorporated with tl diluting water—also testing with organisms their natural environment, as with coli sewage, in place of pure cultures mentation test for emulsified fluids allows 1 p c solution to stand 24 hours before preparii final dilutions for testing (Lancet, Sept 19, 1908 The Lister test' (U.J. Martin) is made in the presence of 3 p.c. of fæces, previously sterilise dried, and powdered. Sims Woodhead at Ponder (J. Path. and Bact. 1909, 148) replace tl moculating loop by platinum spoons, used with a special sterilising apparatus, and extend the

time to 30 minutes. A mean coefficient is taken for two values obtained at the beginning and end of the experiment with B. coli and subculturing into McConkey's bile-salt fluid. The Lancet (Nov. 1909) has suggested, in conjunction with the above method, a chemical analysis for the valuation of emulsified fluids as follows 10 grams of disinfectant with 100 cc. of water and 15 grams baryta, are heated for 30 minutes at 100° under a reflux condenser, then cooled and decanted through asbestos, washing with warm baryta water; the tar acids are extracted with ether from an aliquot portion of the filtrate, after adding calcium chloride and acidifying with HCl, dried at 38°, weighed, and their bromine absorption determined hydrocarbon oils are extracted from the baryta residue with acetone, and the insoluble soaps and resins treated with HCl and shaken out with For estimation of water, 25 grams of the sample are acidified with 10 cc. of 10 pc H,SO4 and shaken with 25 c c of white petroleum spirit, taking the volume less 10 c c of the aqueous layer after standing. In presence of the higher phenoloids, the weight of tar acids obtained is too low. The coefficients obtained for B coli were found to be approximately onethird of the difference between the percentage of tar acids and the bromine value

In the United States, the National Hygienic Laboratory method of L Anderson and T B McClintic (1912) is now widely adopted The method, which follows closely the RW in technique, differs in the following points of detail. The standard temperature of medication of 20°C, instead of 18°C, is adopted 01 cc of broth culture is added to 5 c c of disinfectant solution instead of 5 drops. The coefficient is determined from the mean of the following two

Weakest strength of disinfectant which kills in 2½ minutes

Weakest strength of phenol that kills in the same time

and the similar ratio for death in 15 minutes From time to time various modifications

have been introduced in which solid media such as nutrient agar have been substituted for broth An emulsion of an agar slope culture of an organism is made with the various disinfectant dilutions and phenol controls, whilst the sub-culturing is either performed direct into agar slopes or spread out on agar plates, thus permitting counts to be made Nutrient agar made up with fresh meat is certainly less inhibiting than standard broth, whilst certain advantages attend the use of plates, giving approximate data as to the number of organisms surviving after specified periods of disinfection, as against the 'end point' method of the fluid media tests These advantages are nearly entirely negatived by the difficulty of obtaining a uniform agar streak growth emulsion for medication, and the extra labour involved in plating a number of tests

The germicidal power expressed in terms of carbolic acid with B typhosus is shown for a number of disinfectant substances in the following table; while most of the figures are Rideal-Walker carbolic acid coefficients, some of the results have been obtained by modifications of colloid mercury.

the original test as noted, and these are obviously not comparable with the others.

not comparable wi	ith the othe	ris.
Agent Ger	rmicidal valu	e Observer
Absolute alcohol l		F. 1905
Acetic acid .	06	R 1907
Acetyl-coumaric		
acid	4 5	
Andma	0 57	
. m . 1 1		
o-Toluidine .	1 00	
m-Toluidine .	1 3	
Benzoic acid . a	bout 50	R 1910
	ss than 0 1	W. 1904
Bromine water	64	R. 1910
Cadmium chloride		R. 1911
Cadmium sulphate	10	R 1911
Chinosol .	0 15-0 30	Various observers
Chlorine water	28	R. 1910
Componental batal	20	10. 1010
Copper sulphate)	0.0	D 1010
(calculated }	$2 \cdot 0$	R 1910
on CuSO <sub>4</sub> )		
Cresylic acid		•
commercial.	3 7	R. 1908
		16. 1500
o-Cresol	2 1	
p-Cresol .	24	M & C. 1912
m-Cresol .	20	
Eucalyptol .	1 2	L. 1904
Formaldchyde	0 55	R 1907
Ditto	0 75	W 1904
Formic acid	57	R 1907
Guaiacol	0.9	L 1904
	V 0	2 1001
Trimethylome-	4.0	***
thoxy-phenol	4 0	W
Quinol	11	М & С.
Catechol .	0.48	M & C
Hydrochlorie		
acid	1 58	fæccs and urine)
Hypochlorites		
(calculated	140 000	37 1
on 'available	146-220	Various observers
(hlorine')		
Ditto with 50		
pe urine .	8	K 1906
lodine water .	100	R 1910
Iodine trichlor-	200	20 1010
	0.4	D 1000
ıde .	94	R 1909
Lactic acid	18	R 1910
Mercuric chloride	400-3540	Various observers
2 3 Dihydroxy		
	4 4	
naphthalene	**	
2 7-Dihydroxy-		
naphthalene	28	
Potassium per-		
manganate.	42	do
	60	uo.
Pierie acid ;		M W D 1000
Pyrogallic acid	0 22	M. W. B 1906
Resorein .	0.30	do
Silver nitrate .	158	R 1910
	4.1	D 1000
phate .	41	R. 1908
Sulphate .	0.09	
17 11 . 1		
Zine entoride.	0 15	W 1906
Zinc chloride .	0 15	W 1906 , F M , Firth and Mac-

M W B, Meredith Wynter Blyth, F M, Firth and Macfadyen, F, Major Fowler, K, Klein; L, Lodwidge; M & C, Morgan and Cooper, R, Rideal, W., Walker.


The extremely high values once given for mercuric chloride were due to the powerful inhibitory action of the traces carried over with the sub-cultures in the inoculations.

S & E R. DISPARGEN. Trade name for a form of 556 DISPERMIN


DISPERMIN. Trade name for piperazine DISSOCIATION v. CHEMICAL AFFINITY DISTHENE v. KYANITE.

DISTILLATION. The object of distillation is the separation of a volatile liquid from a nonvolatile substance, such as pure water from seawater, or more commonly the separation of two or more liquids of different volatility

The apparatus required consists essentially of four parts a still or boiler, a (Fig 1), in which



the liquid is boiled, an upright tube or head, B a condensing tube, c, sloping downwards and cooled by suitable means—a Liebig's condenser, L, through which a current of water is passed, being generally employed in the laboratory,



lastly, a receiver, R. to collect the dis-tilled liquid The tilled hquid form of apparatus shown in Fig 2 is convenient for many purposes, the still, the head, and the condensing tube are scaled together in one piece

The stılÎ usually heated by the flame of a Bunsen burner, for very volatile and inflammable liquids a water-bath is generally used, but the rate of distillation can be better regulated by a very small flame in contact with the bottom of the still, the Liebig's condenser is unnecessary for liquids which boil above 160° or 170° if the condensing tube is fairly long. For many purposes, notably for distillation under reduced pressure, it is convenient to employ an electrically heated coil of platinum, German silver, or nichrome wire as the source of heat (Richards and Matthews, J. Amer Chem Soc 30, 1282, 31, 1200, Beckmann, Zeitsch physikal. Chem 64, 506, Bailey, J Amer Chem Soc 33, 447, Rosanoff and Easley, ibid 31, 964)

If the temperature at which the liquid boils is to be ascertained with accuracy—and this is generally the case in fractional distillation, especially in the final purification of a liquid—the following points should be attended to

1 The thermometer should be carefully calibrated, and it would be a great advantage if all thermometers were compared with an air thermometer, for different kinds of glass expand very differently when heated, and therefore two thermometers, constructed of different varieties of glass, will give different readings, especially at high temperatures

2 The zero point of the thermometer should be redetermined from time to time. It is a good plan to heat the thermometer for several hours to about 300° or 360° before calibrating it; the subsequent clanges of zero point, which other-

wise might be considerable at high temperatures, are thereby greatly reduced (Crafts, Comptrend. 95, 910, also several earlier papers) Thermometers treated in this manner may now be purchased

3 All thermometers, except those which do not register temperatures higher than 60° or 70°, should contain nitrogen to prevent volatilisation of mercury in the heated stem; if the thermometer is provided with a small bulb at the top, rise of pressure owing to compression of the gas

in the thermometer is avoided

4 The thermometer should, if possible, be so placed in the apparatus that not only the bulb but also the column of mercury in the stem is surrounded by the vapour of the boiling liquid, otherwise the following correction, which, at the best, is somewhat uncertain, must be applied -N = length of column of mercury in scale degree

not heated by vapour T = temperature of vapour (the approximate temperature directly read on the thermo-

meter is usually sufficiently accurate) mean temperature of column of mercury not heated by vapour This temperature may be approximately ascertained by attaching a small thermometer to the other by means of thin indiarubber bands or wire, with its bulb at about the middle of the unheated column

Correction add 0 000143(T t)N

 The coefficient 0 00016 is usually employed. but Holtzmann pointed out that this is too high, and adopted the number 0 000135 however (Chem. Soc. Trans. 37, 1 Thorpe, 159), has shown that the coefficient 0 000143 gives the best results 1. The table for the correction of thermometer readings is taken from Thorpe's paper, and shown in next column

For further information v. article on Ther-

MOMETERS

5 When the amount of liquid in the bulb is very small, the vapour is hable to be superheated by the flame, and unless the bulb of the thermometer is thoroughly moistened with condensed liquid, too high a temperature will be registered If a very little cotton wool or, for temperatures above 230°, a little fibrous asbestos be wrapped round the bulb of the thermometer, it remains thoroughly moist, and with a pure liquid the theirmometer registers a perfectly constant temperature until the last trace of liquid in the bulb has disappeared (Ramsay and Young, Chem Soc Trans 47, 42)

6 The barometer should always be read, and corrected to 0°, and in a long distillation or in unsettled weather it should be read frequently, for the boiling-point varies greatly with the pressure The observed boiling-point is corrected to that under normal pressure by means of the formula  $\theta = c(700 - p)(273 + t)$ , where  $\theta$  is the correction in centigrade degrees to be added to the observed boiling-point,  $t^{\circ}$ , c is a constant; and p is the barometric pressure. For the

<sup>1</sup> Rimbach (Bet 22, 3072) has constructed tables of ¹ Rimbach (Bet 22, 3072) has constructed tables of corrections by interpolation from experimental data up to 220° for two kinds of thermometer graduated in degrees, and up to 100° for thermometers graduated in tenths of a degree — All these thermometers were constructed of Jena glass, and from the results it would appear that the form of the their nometer, and doubtless also the nature of the glass, have a notable influence on the magnitude of the correction. majority of liquids the value of c may be taken. without serious error, to be 0 00012; but for water and the alcohols it is about 0.00010, and for a few liquefied gases it is considerably higher. A table of corrections was first given by Crafts (Ber 20, 709) The values of c for a large number of substances have been collected and

	200	0.29 0.557 0.657 1.11 1.12 1.72 1.72 1.72 1.72 1.72 1.7
-	180	00 00 00 00 00 00 00 00 00 00 00 00 00
	180	3 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
	170	0.454 0.433 0.433 1.466 1.466 2.92 2.92
	160	22.50 22.20 22.20 22.20 23.20 24.11 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20 25.20
	150	00000000000000000000000000000000000000
	140	00001111111111111111111111111111111111
1	130	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0
	130	0.17 0.34 0.51 0.69 0.69 1.03 1.20 1.37 1.34 1.34 1.34
	110	0.16 0.31 0.47 0.63 0.79 0.79 1.26 1.26 1.58 1.58 1.69
Z	100	0 14 0 0 14 0 0 0 2 1 0 0 0 1 1 0 0 0 1 1 0 0 1 1 0 0 1 1 0 1 1 1 1
	06	0 13 0 26 0 39 0 52 0 52 0 64 0 77 1 0 9 1 12 1 12 1 12 1 15 1 15 1 15
	80	0 11 0 22 0 34 0 47 0 60 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
	7.0	0 10 0 20 0 30 0 40 0 60 0 60 0 60 0 60 0 60 0 60 0 6
	9	0000 0000 0000 0000 0000 0000 0000 0000 0000
	20	0 0.7 0.14 0.28 0.28 0.43 0.50 0.50 0.71 0.79
	40	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
	30	0004 00113 0025 0025 00443 0647
	20	00000000000000000000000000000000000000
	93	0.00 0.05 0.05 0.05 0.05 0.05 0.11 0.11
	- <del></del>	100 100 100 100 110

tabulated by Young (Chem. Soc Trans. 81, 777; Fractional Distillation, 15)

The various parts of the simple apparatus (Figs 1 and 2) may require modification under certain circumstances.

The still -For larger quantities of liquid, a tin or copper vessel is preferable to glass.


The head -The modifications of this part

of the apparatus will be considered under Fractronal distillation.

The condenser .- For very volatile liquids, the condensing tube must be cooled by ice or a

freezing mixture (pounded ice and salt or ice and concen. trated hydrochloric acid, &c) In this case, a spiral or worm' tube should be used (Fig 3). Condensation of moisture in the receiver is prevented by the drying tube a V CONDENSERS.

The receiver -If a liquid boils at a very high temperature, or if it suffers decom-






Fig. 4

distillate is to be collected in separate portions. as in fractional distillation, the removal of the receiver would necessitate the admisapparatus and a sion of air into the fresh exhaustion after each change This inconvenience may be obviated by the employment of stopcocks arranged in such a manner that air may be admitted into the receiver and a fresh one put in its place while the distillation bulb remains exhausted (Thorner, Ber 9, 1868, F D Brown, Chem Soc. Trans 35, 554, L T Thorne, Ber 16, 1327 (Fig 5),



Fig 5

Lothar Moyer, ibid 20, 1834, Wade and Merriman, Chem. Soc. Trans. 99, 984; Rosanoff and Easley, lc; Rosanoff, Bacon and White, J. Amer Chem. Soc 36, 1806) Or again, a series of receivers on a revolving stand placed under an exhausted bell-jar may be empl-yed (Gorboff and Kessler, Ber. 18, 1363, Bruhl, *ibid.* 21, 3339 (Fig. 6)).

Many liquids absorb moisture from the air or are acted upon by oxygen, and in these cases special precautions must be taken, the moist air in the apparatus being expelled and replaced by dry air, carbon dioxide, nitrogen, or hydrogen;




Fig. 6.

the receivers must also be protected by drying tubes (For example, v Thorpe, Chem. Soc Trans 47, 120 , Young,  $\imath bid$  59, 923)

Theory of distillation.—Investigations of the vapour pressures of liquids have thrown much light on the subject of distillation If a pure liquid be confined in the space above the mercury in a barometer tube, it will evaporate until the pressure of its vapour-measured directly by the depression of the mercury column—has reached a definite amount, which, for a given substance, and, if liquid be still present, depends solely on the temperature From the readings of pressure at various temperatures we may construct a curve to illustrate the relations of temperature to pressure for a given substance, but this curve has a twofold meaning, for it represents not only the vapour pressures at definite temperatures, but also the boilingpoints of the liquid under definite pressures. Thus the vapour pressure of pure water at 100° is 760 mm of mercury at 0°; and under a  $100^{\circ}$  is 760 mm of mercury at  $0^{\circ}$ ; and under a pressure of 760 mm water boils at  $100^{\circ}$ . Again, the vapour pressure of water at 0° is 46 mm, and if the pressure in a distillation bulb containing water be reduced to 4.6 mm, the water will boil at 0°. The truth of this general statement was conclusively proved by Regnault (Mém de l'Academie, 21 and 26)

Regnault also investigated the behaviour of a few mixed and heterogeneous liquids, which he divided into three classes: (1) liquids which are quite or almost insoluble in each other; (2) liquids which are miscible only within certain limits; (3) liquids which mix in all proportions. Regnault employed both the statical and dynamical methods of investigation; that is to say, he determined, on the one hand, the vapou pressures of the liquids at vanous temperatures

in the barometer tube, and, on the other hand, the boiling-points of the liquids under various

The behaviour of two liquids, A and B, when brought into contact with each other, must be affected (a) by the attraction of the like molecules; those of A for each other, and those of B for each other, and (b) by the mutual attraction of the molecules of A and B If the attraction of the molecules of A and B tion of the unlike molecules is relatively very slight, it may be expected that the liquids will be non-miscible, or nearly so, as in the case of benzene and water If the attraction of the unlike molecules is greater, but still relatively small, miscibility within limits may be expected, as, for example, with aniline and water. In such cases there is frequently slight expansion and slight absorption of heat on admixture. Comparing together various pairs of liquids, as the mutual attraction of the unlike molecules increases relatively to that of the like molecules, the following changes may be expected; (a) increasing and finally infinite miscibility; (b) slight expansion, diminishing to zero, and followed by increasing contraction; (c) diminishing absorption of heat, changing to increasing These changes do not, in many heat evolution cases, run strictly pari passu, and among liquids which are miscible in all proportions, it is not unusual to find a small amount of contraction attended by a slight absorption of heat, as, for example, when a little water is added to normal propyl alcohol, but when certain compounds, such as chlorobenzene and bromobenzene, which are chemically closely related, are mixed together, there is no appreciable change of volume or temperature For such substances it is probable that the different molecular attractions, A for A, B for B, and A for B, are very nearly equal, and that the relation suggested by Galitzine (Wied. Ann 41, 770; and by D Berthelot, Compt rend. 126, 1703), namely, that  $a_{AB} = \sqrt{a_A a_B}$ , holds good, where  $a_{AB}$  represents the attraction of the unlike molecules, and  $a_A$  and  $a_B$  the respective attractions of the like molecules

It would appear, then, that there are two very simple cases I that in which the liquids are non-miscible; II that of two infinitely miscible and closely related liquids which show no heat or volume change on admixture

I. It was found by Regnault that when two non-miscible liquids are placed together over the mercury in a barometer tube, the observed vapour pressure is equal to the sum of the vapour pressures of the two liquids when heated separately to the same temperature. It is only necessary that both substances should be present in sufficient quantity, and that the tube should be shaken or sufficient time allowed for the evaporation of the heavier liquid. liquids evaporate independently of each other, and Dalton's law of partial pressures is therefore applicable to the distillation of non-miscible liquids. Each vapour behaves to the other as an indifferent gas, and the boiling-point of each liquid depends on the partial pressure of its own vapour. The temperature is necessarily the same for both liquids, and the total pressure, if the distillation is carried out in the ordinary manner, is equal to that of the atmosphere. The boiling-point is, therefore, that temperature it which the sum of the vapour pressures of the components is equal to the atmospheric pressure. For example, at 90·23° the vapour pressure of water is 530·1 mm; and that of chlorobenzene is 210.1 mm; total, 740·2 mm; and it was ound that when chlorobenzene and water were listilled together under a barometric pressure of 740·2 mm, the temperature varied only between 90·25° and 90·35°, until there was carcely any chlorobenzene left in the residue, when it rose rapidly to 100°. The boiling-point of a pair of non-miscible liquids is necessarily lower than that of the more volatile component, and may be far lower than that of the less volatile.

The process of distillation with steam is very frequently employed in the case of subtances of high boiling-point, which are insoluble in only slightly soluble in water, such as amline in introbenzene (For the application of steam listillation to the preparation of ethereal oils, von Rechenberg, Theorie der Gewinnung ind Trennung der atherischen Oele durch Destillation, Schimmel & Co. 1910)

The composition of the vapour—and thereore of the distillate—from two non-miscible iquids, like the vapour pressure and boiling-point, is independent of the relative amounts of the components, provided that both are present n sufficient quantity and that evaporation can ake place freely Calling the vapour densities  $D_A$  and  $D_B$ , and the vapour pressures at  $t^\circ$   $D_A$  and  $D_B$ , there will be, in a litre of the mixed vapour, 1 litre of A at  $t^\circ$  and  $p_A$  mm., and 1 litre of B at  $t^\circ$  and  $p_B$  mm. The weights of vapour vill therefore be:

$$-\frac{0.0899\times {\rm D_{A}}\times 273\times p_{\rm A}}{(273+t)\times 760} {\rm and} \, \frac{0.0899\times {\rm D_{B}}\times 273\times p_{\rm B}}{(273+t)\times 760}$$

espectively, and the relative weights will be  $O_A \times \underline{P_A}$  (Naumann, Bor. 10, 1421, 1819, 2014,  $O_B \times \overline{P_B}$  (Soumann, Bor. 10, 1421, 1819, 2014, 1999; Brown, Chem. Soc. Trans. 35, 547)

The vapour density of chlorobenzene is 562, and that of water is 9; and at 9023° the elative weights of vapour will be

$$\frac{m'_{A}}{m'_{B}} = \frac{562 \times 210 \cdot 1}{9 \times 5301} = 2475,$$

nd the percentage weight of chlorobenzene rill be 71.2. In the actual experiment to which eference has been made, the percentage of hlorobenzene in the distillate was found to e 71.4. Both boiling-point and vapour omposition agree well with the calculated alues

II The vapour pressure of a mixture of we infinitely miscible liquids which are chemially closely related to each other, or, probably, or which  $a_{\rm A}_{\rm B} = \sqrt{a_{\rm A}.a_{\rm B}}$ , is given by the formula  $00p = {\rm M}p_{\rm A} + (100 - {\rm M})p_{\rm B}$  where M is the moleular percentage of A, and  $p, p_{\rm A}$ , and  $p_{\rm B}$  are he vapour pressures of the mixture and of and B respectively, at the same temperaure,  $t^{\circ}$ . In other words, the relation between apour pressure and molecular composition is epresented by a straight line (Van der Waals,

Proc Roy Acad. Amsterdam, 3, 170; Young, Chem. Soc. Trans. 81, 768; Young and Fortey, 1b1d. 83, 45; Zawidski, Zeitsch physikal. Chem. 35, 129; Vanstone, Chem Soc Trans. 97, 429; Price, 1b1d 115, 1116) The relation has been found to hold accurately for chlorobenzene and bromobenzene, and with very slight error for other pairs of closely related liquids

In order to calculate the boiling-points of all mixtures of two such liquids under a given pressure, p, the vapour pressures of each must be known at all temperatures between their respective boiling-points under that pressure The percentage molecular composition of mixtures which would exert the vapour pressure p must then be calculated at a series of temperatures between these limits by means of the for-

mula  $M=100\frac{p_B-p}{p_B-p_A}$ . Lastly, the values of M must be plotted against the temperatures, when the curve drawn through the points will give the required relation between boiling-point and molecular composition. In the cases examined, the agreement between the observed and calculated results is quite satisfactory.

The relation between the composition of a liquid mixture and that of its vapour (distillate) has been the subject of many experimental investigations (Brown, Chem Soc Trans 35, 547; 39, 304, Lehfeldt, Phil. Mag. (v.) 46, 42, Zawidski, Zeitsch physikal Chem 35, 129, Carveth, J Phys Chem 3, 193; Winkelmann, Wied Ann 39, 1; Linebarger, J Amer Chem Soc 17, 615; Gahl, Zeitsch physikal Chem 33, 179; Rosanoff, Lamb, and Breithut, ibid 66, 349; Rosanoff and Easley, J Amer Chem. Soc. 31, 953, Rosanoff, Bacon and White, ibid 36, 1803; Rosanoff, Bacon and Schulze, *ibid* 36, 1993; Rosanoff and Bacon, *ibid* 37, 301 and 1072) Special attention may be directed to the last four of these papers. Unfortunately, however, it is only in a very few cases that the mixtures investigated consisted of closely related liquids (Young, Sci Proc Roy Dubl Soc 15, NS, 667) In the discussion of their results, Lehfeldt and Zawidski made use of formulæ which they derived from a general equation proposed independently by Duhem and by Margulès, and it is to be noted that both formulæ, in the simplest cases, can be reduced to one originally given by Brown,  $\frac{m'_{A}}{m'_{B}} = c \frac{m_{A}}{m_{B}}$ , where  $m'_{A}$  and  $m'_{B}$  are the relative weights of the two substances in the vapour,  $m_A$  and  $m_B$  the relative weights in the liquid, and c is a constant. From

Zawidski's formula it would follow that  $c = \frac{p_A}{p_B}$ . So far as experimental evidence is available, it would appear that Brown's formula is applicable to mixtures of liquids for which  $a_{AB} = \sqrt{a_A a_B}$ .

and that for closely related liquids  $c = \frac{p_A}{p_B}$ .

Another formula, which gives good results, is proposed by Rosanoff, Bacon and Schulze (loc. cit.)


When the components of a liquid mixture are not chemically closely related, the relation between vapour pressure and molecular composition is not, as a rule, represented by a straight

line, but by a curve, and, in all probability, the form of the curve depends on the relation between  $a_{\rm AB}$  and  $\sqrt{a_{\rm A}}\,a_{\rm B}$ , as shown in Fig. 7. That is to say, when

$$a_{AB} < \sqrt{a_A a_B}$$
,  $100p > Mp_A + (100 - M)p_B$ , and vice versa

The greater the difference between  $a_{A \ B}$  and  $\sqrt{a_A} \, a_B$  the greater will be the curvature, and it will be seen that, for any given values of  $p_A$  and  $p_B$ , if the deviation from straightness exceeds a certain limit, there must be a point of maximum or minimum pressure on the curve. It is obvious, also, that the smaller the difference between  $p_A$  and  $p_B$ , the smaller will be the deviation required to give a maximum or minimum pressure. If  $a_{A \ B}$  is so much smaller than  $\sqrt{a_A} \, a_B$  that the two liquids are only partially miscible, the maximum pressure will not be represented by a single point, but by a horizontal line forming part of the curve

The influence of chemical relationship is well seen by the behaviour of the monhydric aliphatic alcohols towards water. These alcohols may be regarded as derivatives of water, formed



by the replacement of a hydrogen atom by the group  $C_nH_{2n+1}$ , the smaller the alkyl group, the closer is the relationship of the alcohol to water At the ordinary temperature methyl alcohol is miscible with water in all proportions, as also

are ethyl and propyl alcohols, normal butyl and isobutyl alcohol are only partially miscible with water, and the solubility of the higher alcohols in water diminishes with rise of molecular weight, cetyl alcohol, for example, being practically insoluble. The volume and heat changes on admixture with water similarly indicate that the attraction of the unlike molecules diminishes as the molecular weight of the alcohol increases. Lastly, the deviation of the vapour pressure curves from straightness increases (Konowalow, Wied Ann. 14, 34), as may be seen from Fig. 8


(The temperatures are such that the vapour pressure of each pure alcohol is 400 mm)

The boiling-points of liquids which are not closely related cannot, as a rule, be calculated by means of the formula  $M=100 \frac{p_B-p}{m_B-p}$  If,

as is usually the case, the vapour pressures of the mixtures are higher than are given by the formula  $100p = Mp_A + (100 - M)p_B$ , the observed boiling-points will be lower than the calculated; if the vapour pressures are lower, the boiling-points will be higher. Moreover, when there is a point of maximum pressure on the curve which shows the relation between vapour pressure and molecular composition, there must be a point of minimum temperature

on the boiling-point—molecular compositic curve, the composition of the mixture whice exerts a maximum vapour pressure, p, at being the same as that of the mixture which has the minimum boiling-point,  $t^{\circ}$ , under the pressure p So, also, two substances capabof forming a mixture of minimum vapour pressure can also form a mixture of maximum boiling-point

Such mixtures of either minimum or maximum boiling-point, when distilled, boil at a constant temperature without change of composition like pure substances, and they have frequently been mistaken for chemical compounds. The composition, however, depends on the pressur which would not be the ease with a definition compound, and, moreover, mixtures of minimum boiling-point are formed owing to the relatively



small attraction of the unlike molecules for each other A very striking example of th influence of pressure on the composition of th mixture of constant boiling-point has bee observed by Wade (Chem Soc Trans 99, 997 in the case of ethyl alcohol and water normal pressure the mixture of minimum boiling point contains 4 43 pc of water At highe pressures the percentage of water is slightly higher, at lower pressures it is smaller, an it is remarkable that the diminution in th percentage of water becomes more and mor rapid as the pressure falls, so that below 80 mm no mixture of minimum boiling-point is forme at all V also Merriman, ibid. 103, 628 and 1790.

In the case of three liquids, if each of th three possible pairs is able to form a mixture of minimum boiling-point, it may happen that particular mixture of all three liquids will be constantly at a lower tem perature than will an of the pairs or single liquids. Thus benzene an water form such ternary mixtures of minimum boiling-point with ethyl, n-propyl, isopropyl, and tertiary butyl alcohols (Young, Chem. Soc Trans. 81, 707, Fortey and Young, ibid 81, 739)

A table of mixtures of constant boiling-point is given in Young's Fractional Distillation, Other mixtures have since been discovered by Wade (Chem Soc Trans 85, 938, 87, 1656, v also 95, 1842), and by other observers The subject has been exhaustively studied by Lecat, La Tension de Vapeur des Mélanges de Liquides, l'Azéotropisme (Brussels, Lecat has collected the data published by other chemists, and has himself investigated more than 2000 mixtures, and has discovered and tabulated avery great number of binary mixtures of minimum boiling-point, and a moderate number of binary mixtures of maximum boiling-point, and of ternary mixtures of minimum boilingpoint. He has also observed the dependence of the composition of the azeotropic mixtures on the pressure in a considerable number of cases.

The composition of the vapour from a

The composition of the vapour from a mixture of liquids which are not closely related, cannot, as a rule, be calculated from the vapour pressures of the components and the composition of the liquid, but for purposes of interpolation the formulæ of Lehfeldt, Zawidski or Rosanoff may be found useful (lc) Rosanoff, Bacon and Schulze (lc) have discovered a simple and very useful solution between the partial pressures (number of molecules) of the components in the vapour and the vapour pressures of the mixture and of the two pure substances. They give a formula which is applicable even when mixtures of maximum or minimum vapour pressure are formed

Fractional distillation. It has been stated that Brown's formula,  $\frac{m'_A}{m'_B} = c \frac{m_A}{m_B}$ , is applicable to mixtures of all closely related liquids so far investigated, and that, in all probability  $c = \frac{p_A}{p_B}$ . The value of c is necessarily such that the vapour is richer than the liquid in the more volatile component. The first portion of distillate obtained by condensation of the vapour will, therefore, be richer, and the

residual liquid will be slightly poorer in that component. Continuing the distillation, the vapour evolved from the residual liquid will again be richer than the liquid in contact with it in the more volatile component, though poorer than the first portion of vapour. If successive portions of vapour be condensed, the percentage of the more volatile component in the distillates or fractions will steadily diminish, until, if the constant of differs greatly from unity, the residue will consist of the less volatile liquid in a pure or very nearly pure state. As the distillation proceeds, the temperature will, therefore, rise until, if c has a high value, the boiling-point of the less volatile component is reached

If the fractions obtained in the first distillation were redistilled, each of them could be separated into smaller fractions, the first richer and the last poorer in the more volatile component, and by a process of systematic fractional distillation it should, theoretically, be possible to effect a separation of both components

A mathematical and experimental investigation (Barrell, Thomas, and Young, Phil Mag (v) 37, 8) of the behaviour on distillation of mixed liquids which follow Brown's law, showed that with two components it is the liquid of higher boiling-point which is the easier to separate; and that if there are three or more components boiling at fairly equal intervals of temperature, the substances of intermediate boiling-point are the most difficult to separate, and the least volatile substance is the easiest See also Loid Rayleigh, Phil Mag [5] 4, 527, and Rosanoff, Baker and Schulze, J. Amer. Chem. Soc. 36, 1993.

The method employed for the fractional distillation of a mixture of two highest can best be explained by taking a concrete example. A mixture of 100 grams of benzene (b.p. 80.2°) and 100 grams of toluene (b.p. 110.6°) was slowly distilled, and the distillate was collected in a convenient number of fractions, the receivers being changed when the boiling-point reached certain definite previously arranged temperatures, the actual readings being corrected for barometric pressure and for criois in the thermometer scale. The results of the first four fractionations, and also of the ninth, are given in Table I

TABLE 1.


Number of	Final	Weight of fraction = $\Delta W$			t	ΔW	
receiver	temperature (corrected) - t	I	11	III.	1V.	13	κ.
Pure benzene	•			_	_	80.2	10.20
1	812		l	12.9 <b>5</b>	31.55	₹0 3	45 00
2	83.2		3 80	24.80	23 90	80.6	17.55
3	86.2		33 85	22.75	16 20	81.4	7.20
4	89.2	9.75	22 30	13.50	9 55	83.5	4.50
5	92 2	<b>51</b> 80	19 6 <b>5</b>	11 80	8 00	88 0	<b>3·3</b> 0
6	95 4	28.85	13 60	9.15	5.80	95.4	2.75
7	986	21 20	12.95	7.30	5 35	102 8	<b>2 6</b> 0
8	101.6	$12 \ 80$	9.05	6 75	4.65	107 3	2.85
9	104.6	11.45	8.90	6 30	3 85	109.4	3 15
10	107 6	14.15	10 80	7 95	5.85	110.2	3.45
11	109 6	13 45	9 60	8 95	7 40	110 5	7.00
12	110.6	24.90	30.75	33 05	30 50	1106	5.75
Pure toluene	(110 6)	10 90	22 95	31 35	42.10	(110.6)	72.10
	'		l		ł	1	

In the first distillation the temperature rose almost at once to 86° (corr), and the first portion of distillate was therefore collected in receiver 4 On the other hand, the temperature reached 110 6° before the whole of the liquid had come over; the residue, therefore, consisted of pure toluene

The distillate in receiver 4 was now placed in a smaller flask and distilled, fractions being collected in receivers 2, 3, and 4 When the temperature reached 89 2°, the flame was removed, and the contents of receiver b were added to the residue in the still The distillation was recommenced, and fractions were collected in receivers 3, 4, and 5, the flame being removed when the temperature reached 92 2° The distillation was continued in a similar manner until, after addition of the contents of receiver 12, the temperature rose to 1106°, when the residue of pure toluene was added to that from the first fractionation. The third and fourth fractionations were carried out like the second In the subsequent fractionations the temperature ranges of the middle fractions were gradually increased and those of the lowest and highest fractions diminished. It was not until the minth fractionation that pure benzene began to be collected Eventually 81 4 grams of pure benzene and 88 8 grams of pure toluene were recovered.

Full details of the systematic fractional distillation of mixtures of two liquids and three liquids, and of more complex mixtures, are given


'Fractional Distillation,' 114-143
If the components of a mixture are not closely related, the composition of the vapour cannot, as a rule, be calculated by Brown's formula, but even in such cases the percentage of the more volatile component in the vapour will be greater than in the liquid, and the separation by fractional distillation is theoretically possible, unless the deviation of the vapour pressure-composition curve from straight-



ness is so great that a mixture of maximum or mınımum vapour pressure is formed.


For mixtures to which Brown's law is applicable, it may be stated quite generally that the greater the difference between the boiling-points of the compounds, the more readily can a separation be effected by fractional distillation, but for other mixtures the form of the curve representing the relation between boiling-point and molecular composition, if known, must be taken into account. If the curve for two components were of the form a

(Fig. 9), the substance of higher boiling-point could be separated very easily, but the more volatile component only with great difficulty, if at all On the other hand, if the curve were of the form b, nearly horizontal at the higher temperature, the more volatile component would be comparatively easy, and the less volatile difficult to separate. The first case is frequently met with, the second seldom. With curves of the form c and d (Fig. 10) it would not, under any conditions, be possible to separate



both components by distillation of any given mixture, but it would be possible to separate the mixture of minimum of maximum boilingpoint from that component which was in excess In a case represented by curve c, where the minimum temperature corresponds to the molecular composition q, the distillation might proceed in either of three ways (1) If the liquid mixture had the composition q, it would distil unchanged in composition at the constant minimum temperature, (2) if the molecular percentage of A were less than q, the liquid would tend to separate into two components, the more volatile of which would be the mixture of minimum boiling-point, and the less volatile the substance B, (3) if the molecular percentage of A were greater than q, the components separable by distillation would be, first, the mixture of minimum boiling-point, and, second, A well-known case of this kind the liquid A is that of normal propyl alcohol and water; the boiling-points under normal pressure are: water, 100°; n-propyl alcohol, 97.2°; mixture of minimum b p. 87.7°. This mixture contains 43.2 molecules p.c. of the alcohol; it was described for many years as a hydrate of propyl alcohol. If the curve had the form d, the mixture would distil like a pure liquid if it had the composition r, or it would tend to separate into two components, the first being either A or B, according as the molecular percentage of A in the original mixture was greater or less than r, and the second the mixture of maximum boiling-point. Formic acid and water behave in this manner, as shown by Roscoe (Chem. Soc. Trans. 13, 146; 15, 270). Boiling-points: water  $100^\circ$ ; formic acid,  $99.9^\circ$ ; mixture of maximum b.p.,  $107.1^\circ$ ; molecular percentage of acid in mixture,  $56.7^\circ$ .

If, as with ethyl alcohol and water (Fig. 11), the mixture of constant boiling-point distils at nearly the same temperature as one of the two pure liquids, it is impossible to separate that liquid in a pure state by distillation. In this



case, as the curve is exceedingly flat at the lower temperature (Noyes and Warfel, J Amer Chem Soc. 23, 463), it is practically impossible to separate the mixture of minimum boiling-point (containing 4 43 pc by weight of water) in a pure state even when water is in excess, although it is quite easy to separate pure water from dilute spirit.

The correct interpretation of the results of the fractional distillation of a complex mixture may be rendered very difficult by the following

causes. (1) the presence of two components boiling at nearly the same temperature; (2) the presence of one or more components in relatively very small quantity, (3) the formation of mixtures of constant boiling-point It is only by plotting the total weights of distillate against the temperatures, or by dividing the weight of each fraction,  $\Delta w$ , by its temperature range and tabulating the ratios  $\Delta w/\Delta t$ , that the results are likely to be understood, and in any case, the existence of a mixture of constant boiling-point may not be discovered, and it may be difficult to decide whether a distillate collected at a practically constant temperature consists of a single pure substance or of two liquids boiling at nearly the same temperature

All the above difficulties are met with in the fractional distillation of petroleum, which contains homologous and isomeric paraffins, polymethylenes and their alkyl derivatives (naphthenes) and aromatic hydrocarbons, Russian petroleum poor in naphthenes and aromatic hydrocarbons. When American petroleum was carefully distilled, it appeared at first as though the distillate coming over between the ordinary temperature and 40° contained only a single component, boiling at about 33°; but by systematic fractional distillation with a very

efficient still-head, it became evident (Fig. 12) that two components, isopentane, bp. 2795, and normal pentane, bp. 362°, were present (Young, Chem Soc Trans 71, 440) Again, in the early fractionations there appeared to be no indication of the presence of any component boiling between 40° and 60°, but by tabulating the values of  $\Delta w/\Delta t$ , it was seen that a very small quantity of a liquid boiling at about 50° was present, and a distillate coming over almost constantly at 495° was eventually obtained. A chemical and physical investigation of this distillate showed, however, that it was a mixture of pentamethylene and a hexane of low boilingpoint The presence of benzene in American petroleum is indicated by the formation of dinitrobenzene, when the distillate from, say, 60° to 80° is treated with nitric and sulphuric acids (Schorlemmer, Annalen, 127, 311) when the petroleum was carefully fractionated, it was found that the small fraction coming over at about 80° yielded no dinitrobenzene, the greater part being obtained from the fractions This anomaly is explained by the fact that benzene forms a mixture of constant boiling-point with n-hexane, coming over at almost the same temperature as the paraffin The amount of benzene being relatively small, the whole of it distils over at the low temperature In Russian petroleum the percentage of benzene is so much greater that some of it comes over at its true boiling-point, 80 2° (Young, Chem Soc Trans 73, 905)

(For a detailed discussion of the interpretation




Fig. 12.

of the results of the fractionation of complex mixtures, v Fractional Distillation, chap 19)

Improved still-heads. It would be quite impossible to effect really difficult separations by means of the ordinary apparatus employed for distillation, and various modifications of the still-head have been devised in order to increase its efficiency. Even when the separation is an easy one, much time and material may be saved by the employment of efficient still-heads. Great progress was made in the improvement of the still-head in commerce—notably for the better sep\_ration of alcohol from weak spirit—before

any advance was made in the laboratory. It will, however, be more convenient to consider first those forms of apparatus which are suitable

for laboratory purposes

In choosing or constructing a still-head, the first point to be considered is its efficiency in separating the components of a mixture. It frequently happens, however, that the quantity of liquid available is small, and it is therefore important that the quantity of condensed liquid in the still-head at any moment shall be as small as possible. If the residual liquid is to value, it is also of importance that the condensed liquid in the still-head shall all return to the still after the source of heat is removed.

In comparing the efficiency and usefulness of different still-heads, it is necessary always to distil a mixture of the same composition, and the rate of distillation must always be the same, because the efficiency of any still-head is greatly diminished by increasing the rate. In the comparisons mentioned in this article, equal weights of pure benzene and pure toluene were employed, and the distillate was collected at the rate of one drop per second. Unless otherwise stated, 50 grams of the mixture were distilled

Of all possible forms, the plain vertical stillhead is the least efficient During distillation, as the vapour rises up the cylindrical tube, the outer part of it comes in contact with the con densed liquid flowing down the sides of the Since the tube is constantly losing heat by radiation and by conduction to the surrounding air, this liquid is constantly cooled and fresh condensation constantly goes on, the outermost layer of vapour probably condensing almost completely with little change of composition The central portion of vapour uses rapidly up the tube, and can only reach the liquid by diffusion or by convection currents, and much of it may pass through the still-head without reaching the liquid at all, the condensed liquid, on the other hand, flows rapidly back to the still In order, however, that a satisfactory separation of the components may take place, as much of the vapour as possible should come in contact with the hot condensed liquid, so that equilibrium, as regards composition, may be brought about all through the tube

The efficiency of a plan vertical still-head is, roughly, proportional to its length. For tubes of equal length, the efficiency is lowest when the internal diameter is about 14 mm. On the other hand, the quantity of liquid and vapour in the still-head increases regularly with the width of the tube; it is therefore better to use very narrow rather than very wide tubes.

wide tubing


Fig 13

Better contact between liquid and vapour, and therefore greater efficiency, is ensured by sloping the middle part of the tube, the extremities being kept vertical, and the efficiency is further improved by bending the sloping portion so as to form a spiral In Young's 'rod and dise' still-head (Fig 13), the downflow of part of the condensed liquid is

greatly retarded by the discs on the central glass rod, and this liquid is protected from the cooling

action of the air, at the same time, eddies and cross-currents are produced in the vapour. The increase in efficiency caused by the introduction of the easily constructed rod with discs is very marked. Wurtz (Ann. Chim. Phys. iii. 42, 129) recommended a vertical tube with a series of bulbs blown on it, and Young has shown that the efficiency is somewhat increased and the amount of liquid in the still-head is greatly diminished by blowing pear-shaped instead of spherical bulbs on the tube (Fig. 14). The 'rod and disc' and 'pear' still-heads are specially suitable for liquids of high boiling-point.

Greater efficiency, combined with diminished condensation of liquid in the still-head, is attained in Young's 'evaporator' column, one section of which is shown in Fig. 15. The



vapour 118CS up the inner tube, passes down between the inner and middle tubes, and up again between the middle and outer tubes, and so reaches the section above. The liquid condensed in the upper part of the still-head falls in drops from the depression at the top of the section on to the flattened top of the inverted middle tube, and spreads itself over its surface, falling again in drops from the bottom of this tube. The temperature of the vapour passing through the inner tube of any section being higher than the boiling-point of the condensed liquid falling on to the inverted tube, re-evaporation takes place to some extent, and, if there is much difference between the boiling-points of the components of the mixture being distilled, buildition may actually occur. The quantity of condensed liquid in the still-head is thereby greatly reduced. The 'evaporator' still-head is specially suitable for accurate work where great efficiency is required (Chem. Soc. Trans. 75, 679. Wade and Meriuman, stud. 99, 984)

75, 679, Wade and Meruman, ibid 99, 984)

The 'Hempel' still-head (Zeitsch. anal. Chem. 20, 502) consists simply of a wide vertical tube, filled with glass beads of special construction, and constricted below to prevent the beads from falling out A short, narrower vertical tube with side delivery tube is fitted by means of a cork into the wider tube. The advantages of the Hempel apparatus are simplicity and efficiency, but the amount of liquid in the still-head is excessive. Raschig employs thin rings

glass beads, the depth of a ring being about equal to its diameter. Lessing uses split rings, the metal being also bent inwards across the These rings are easier to con- are given in Table II. ring, thus

of copper or some other metal in place of the | struct and give a larger surface; they give very good results Rings are much lighter than beads

The results obtained with various still-heads


TABLE II.

						11.							
Nature of still-head or dephlegmator	Plan vertical, 5 1 mm. diameter	Plam vertical 14 0 mm. diameter	Plam vertical 25 7 mm. diameter	'Rod and disc' 14 mm. diameter	'Pear'	'Evaporator'	Evaporator'	' Evaporator'	'Hempel' with 200 beads	Glinsky	Le Bel-Henninger	Young and Thomas	Young and Thomas
Number of discs, bulbs, or sections		_		20	13	3	5	8		3	3	3	18
Vertical height in em	62	62	62	62	62	46	62	78	58	30	43	51 ¹	130
Final temperature of fraction	-				Perc	centage	weight o	of distill	ate				
83 2 86 2 89 2 92 3 95 4 98 5 101 6 104 6 107 6 110 0 110 6 Pure toluene by difference	$ \begin{array}{c}  -05 \\  22 2 \\  165 \\  92 \\  72 \\  58 \\  55 \\  59 \\  71 \\  89 \\  11 2 \end{array} $	01 128 215 110 89 74 62 73 84 90 74	$ \begin{array}{r}     \hline       0 2 \\       20 0 \\       18 8 \\       9 2 \\       7 2 \\       4 7 \\       5 9 \\       6 \cdot 4 \\       10 0 \\       7 2 \\       10 4 \end{array} $	1 4 18 1 15 8 9 7 6 1 4 2 3 8 4 2 4 3 8 0 6 6	3 0 26 2 11 0 5 8 5 2 2 4 2 4 2 8 3 6 5 0 11 0 21 6	12 80 21 15 7 50 4 80 3 35 2 70 2 30 2 50 3 50 6 50 10 50 22 40	26 00 12 35 5 60 3 25 1 95 1 30 1 80 1 50 2 45 4 20 11 40 27 90	35 85 6 50 2 90 2 15 1 40 1 15 1 05 1 16 3 95 9 60 32 70	20 6 15 4 7 2 3 4 3 2 2 2 1 8 2 9 3 2 6 8 8 4 24 9	1 1 11 7 17 3 11 9 7 5 5 6 5 7 5 6 8 9 13 7 2 5	0 5 17 8 15 9 8 8 7 2 4 3 4 8 4 6 5 5 9 8 6 8	0 6 21 8 14 3 7 8 5 0 4 7 4 0 3 6 5 2 7 3 9 8 15 9	38 8 4 1 1 9 1 8 1 3 1 2 0 9 1 2 2 0 4 0 8 1 34 7
Weight of liquid and vapour in still-head.	1 05	1 55	3 15	2 35	26	2 70	4 55	6 25	7 85	28	5 85	28	12 1

In many of the still-heads employed on the large scale, for example in the Coffey still (Fig 28), the condensed liquid is caused, by means of suitable obstructions, to collect into shallow pools, and the ascending vapour has to force its way through these pools, very good contact is thus brought about between vapour and liquid at definite intervals. The excess of liquid is carried back from pool to pool, and finally to the still by suitable trapped reflux tubes. It is convenient to reserve the term 'dephlegmator for this class of still-head The Le Bel-Henninger' (Fig 16) and the Glinsky' (Fig 17) dephlegmators have been very frequently employed for laboratory work, but they do not give good results when small quantities of hquid are distilled; the dephlegmators of Brown (Chem Soc Trans. 37, 49) (Fig 18) and of Young and Thomas (ibid 75, 673) follow the principle of the Coffey still more closely, and are much better in this respect.

In a comparison of the Glinsky, Le Bel-Henninger, and Young and Thomas dephlegmators, each with three sections, it was found that when 400 grams of the mixture of benzene and toluene were distilled, there was not a great difference in efficiency, but the Glinsky

was the least satisfactory With 50 grams of the mixture the differences (Table II) were



quite noticeable, and with 25 grams they were greatly accentuated. With the Glinsky 1 Unnecessarily long

column the highest temperature reached was only 107.6°, and the residue, when distilled from a very small bulb, came over between 102.2° and 1104° No pure toluene was therefore obtained, and the residue left in the still was distinctly less pure than the last portion of distillate collected With the Le Bel-Henninger column the highest temperature was only  $107.25^{\circ}$ , and the residue began to boil at  $105.7^{\circ}$ with the Young and Thomas column the temperature just reached 110 6°, and the residue began to boil at 110 4°. The 'rod and disc,' the 'pear,' and the 'evaporator' still-heads gave better results when only 25 grams of the mixture were distilled, the residue in each case consisting of pure toluene. These results show consisting of pure toluene that for small quantities of liquid the Young and Thomas dephlegmator gives much better results than the Glinsky or Le Bel-Henninger, but that none of them is so satisfactory as the other still-heads.


In order that a dephlegmator may give the best results, the following requirements should be fulfilled · 1 It should be possible to greatly increase the number of sections without seriously increasing the fragility of the apparatus the amount of condensed liquid flowing back at any level is greatest at the bottom and least at the top, it follows that, in order to form a pool, more complete obstruction is required at the top of the column than at the bottom 3 Since the amount of condensed liquid in the still-head should be as small as possible consistently with efficiency, the reflux tubes should not be widerin that part which is filled with liquid during distillation-than is necessary to carry back the liquid freely, and the traps should be no deeper than is required to prevent the ascending vapour from forcing its way through them. On the other hand, the upper part of the reflux tubes should be wide, so as to facilitate the entrance 4 There should be a rapid flow of the liquid of condensed liquid through the reflux tubes, otherwise the change in composition of the liquid in these tubes will lag behind that of the distillate 5 There should be a reflux tube connecting each section with the one below it, so that the change in composition may be regular from top to bottom of the column In this respect the Glinsky dephlegmator is entirely wrong in principle 6 When the distillation is completed, the residual liquid should return as completely as possible from the still-head to the still In this respect the dephlegmators as a class are less satisfactory than the other stillheads.

By surrounding the still-head with water or any other liquid, the temperature of which is kept as little above the boiling-point of the more volatile component as will allow of vapour passing through, a considerable improvement in the separation is effected. The temperature of the bath, however, requires very careful regulation if the boiling-points of the components are near together, or if one component is present in large excess, for in either case a fall of a fraction of a degree would cause complete-condensation of the vapour, while a slight rise of temperature would prevent any condensation from taking place, and there would be no fractionation whatever. In constructing a regulated tempera ure still-head, it is advisable to bend the tube into

the form of a spiral, because its effective length may thus be greatly increased without unduly adding to the height of the bath, and also because the spiral form is more efficient than the vertical But Rosanoff and Bacon (J. Amer. Chem Soc 37, 301) find that an annular form of regulated temperature still-head is far more efficient and quite as simple as the spiral tube

form This class of still-head was first recommended by Warren (Annalen, Suppl 4, 51) A better apparatus was described by Brown (Chem. Soc Trans. 37, 49; 39, 517) (Fig 19) A combination of a regulated temperature still-head and dephlegmator was employed by Young and Thomas (ibid. 71, 440) for the separation of normal and iso-pentane from petroleum.

Experiments to test the relative efficiency of these various forms of apparatus have been made by Kreis (Annalen, 224, 259, Claudon, Bull Soc chim 42, 613; Brown, Chem Soc Trans 37, 49; Young, ibid. 75,



679, Fractional Distillation, chaps. 10, 11, and 14; Friedrichs, Zeitschr Angew Chem, 32, 341)

Continuous distillation of wort has long been carried out on the large scale by means of the Coffey still (Fig. 28) In 1902 Carveth (J Phys Chem 6, 253) and Lord Rayleigh (Phil Mag. vi 4, 521) independently suggested apparatus intended for use in the laboratory The liquid to be distilled (or its vapour) is introduced continuously into the middle part of a long spiral or other still-head, the lower part of which is prevented, by means of a liquid or vapour-bath, from cooling below the boiling-point of the less volatile of the two components, while the temperature of the upper part is prevented, by means of a liquid bath, from rising above the boiling-point of the more volatile component. Lord Rayleigh obtained good results in the distillation of weak spirit. It is probable that the method might be made applicable to the distillation of mixtures of three components (Fractional Distillation, 189) See also Rosanoff, U S Pat 1171464 Constant temperature zones are utilised to obtain products of the desired composition


As many of the stills used in the arts are described under the headings Alcohol, Benzene, Paraffin, &c, it will be unnecessary here to give a minute description of all of them, but an account of the principles underlying the different methods will not be out of place

The still is usually made of copper or some other metal, but for special purposes other materials must be employed. Thus for the distillation of sulphuric acid, vessels of glass or platinum are used; for nitric acid, cast iron is found to be the best material; stoneware or glass stills are employed for bromine and iodine; and so on

The still may be heated by the free flame


of a furnace or by steam. Fig. 20 shows an arrangement for heating a still with steam, which may be above the atmospheric pressure.

When only a rough separation of the constituents of a complex mixture is required, as in



the preliminary distillation of coal tar, dephlegmators or rectifiers are not employed, but the vapours pass directly from the still to the condenser


For the better separation of the components of a mixture, modified still-heads are employed A simple apparatus, devised by Mansfield for the distillation of tar oils, is shown in Fig 21, the vapours are partially condensed in the still-head a, which is cooled by water, and the condensed liquid flows back into the still B, the purified vapour passing on to the condenser c




A more perfect separation is effected by Joupier's apparatus (Fig 22) In addition to the column a, there is a series of bulbs, B, B, placed in containing brine, which may be heated by steam by means of the pipe c. The vapour—of benzene, for example—partially freed from ess volatile substances in the column, is further purified by partial condensation in the bulbs, the ondensed liquid flowing back by the tubes D, D nto the column, while the purified vapour passes in to the condenser by the pipe E.

A very similar arrangement is seen in the rench column apparatus (Fig 23) The liquid heated by a steam coil in A; the vapour is

partially purified in the rectifier B; it is further freed from the less volatile components in the




condensing apparatus c, and finally reaches the condenser p



In Savalle's apparatus also the vapours pass first through a dephlegmator or analysing

column, then through a  $\mathbf{or}$ multitubular condenser, which is provided with a water supply, regulated that its temperature is about that of the boiling-point of the liquid required, the liquid condensed in the regulatedtemperature still-head returns to the dephlegmator, and the purified vapour passes on to the cold condenser


In most of the still-heads employed on the large scale, the vapour first passes through a dephlegmator, the construction of which varies considerably in different cases Dubrunfaut's dephlegmator, employed in



Frg. 24.


the French column apparatus, is shown in Fig 24. Two others, devised by Savalle, are represented in Figs. 25 and 26.

In the Ergot dephlegmator an arrangement somewhat similar to the dome (Fig 24) is used, but a large number of these domes are placed in each tray, and the condensed liquid is made to follow a zigzag course from the circumference to the centre of each tray



The principle of the still-head in the Pistorius still is quite different. The vapour, entering a section (Fig 27) of the still-head, is deflected from centre to cucumference by the flat dome, it then passes back to the centre above the dome, and is partially condensed by the water in the shallow vessel above—It may be pointed Kubierschky (Ger Pat 269995, 1913)—has out that if, in the Pistorius still-head, the 'roof' devised an excellent column in which the liquid

of the section were depressed at the centre instead of the circumference, the condensed liquid would drop on to the flat dome below, and the arrangement would then be somewhat



similar to that of the 'evaporator' still-head (Fig. 15). In all probability the efficiency of the still-head would thereby be greatly increased.




Fig 28

falls in drops through perforated plates, whilst | between two concentric cylinders. The vapour the vapour passes freely upwards and downwards, travelling from centre to circumference and from circumference to centre alternately, coming into intimate contact with the falling drops.

A simple and efficient still-head has been described by Foucar (Eng. Pat. 1999) of 1908) The column consists essentially of a helical

from the still passes up the long spiral passage before reaching the condenser. The inner cylinder can, if desired, be used as a thermostat, the column then acting as a regulated-temperature still-head Dutton (J. Soc. Chem Ind 38, 45 T) has described a still-head, suitable for laboratory work, somewhat similar in principle to Foucer's A spiral of ware is wound round an septum which traverses the annular space inner closed tube which fits quite loosely in the DIURETIN. 569

outer tube of the still-head. But excessive condensation, with consequent priming, must be prevented by covering the tube with a thick layer of cotton wool The still-head is stated to

give very good results

A Hirsch (J. Ind. Eng Chem 1910, 2, 409) recommends the use of a differential condenser in the still-head, in the form of a spiral of metal tubing, through which a current of water of regulated initial and final temperature is caused to flow. The apparatus works easily and well so long as the temperature of the vapour leaving the still-head is considerably lower than that of the vapour entering it, but the difficulty of regulation increases greatly as this temperaturedifference diminishes

For work on the large scale Hirsch recommends the introduction of a differential condenser between the lower and upper parts of

a rectifying column

In England, the Coffey still is largely used, it differs considerably from those described, inasmuch as there is no large boiler, and the heat is supplied, not from an outer fire or steamjacket, but by the condensation of steam in the wort, moreover, the process is continuous essential parts of the apparatus are shown in Fig. 28(p 568) The wort is pumped from a reservoir A, up the pipe B, and passes down the rigzag pipe cc, where it is heated by the ascending vapours. then up the pipe D, from which it is discharged into the column E It is prevented from falling through the perforated plates of the column or analyser by the ascending vapours, but falls through the pipes F, F, from one plate to another until it reaches the bottom, when it is allowed to escape Meanwhile, steam is passed into the analyser E by the pipe H, and causes the wort to boil, so that by the time it has reached the bottom it is completely deprived of alcohol The ascending vapours force their way through the perforated plates of the analyser, a portion of the aqueous vapour being condensed by each washing, and the descending wort being thereby On reaching the top of the column, the concentrated alcoholic vapour passes down the pipe k into the rectifier, and there ascends through similar perforated plates; but in the rectifier it is not washed by wort, but by the weak alcohol formed by the previous condensa-This dilute alcohol, on reaching tion of vapour the bottom of the rectifier, flows off into a reservoir, from which it is pumped into the top of the analyser, where it mixes with the descending In the upper part of the rectifier there are simply shelves, which compel the vapour to take the same zigzag course as the pipe which conveys the wort downwards The purified and concentrated vapour then passes through the pipe L to the condenser In modern continuous stills of the Coffey type there are special arrangements for the separation of (a) volatile impurities such as aldehyde, (b) pure rectified spirit, (c) fusel oil. For details the articles on ALCOHOL, etc., should be consulted S. Y.

**DISTYRENE** C<sub>18</sub>H<sub>18</sub>. Produced by heating styrene (q.v.) with hydrochloric acid at 170° (Erlenmeyer, Annalen, 135, 122) Also, together with distyrenic acid and carbon dioxide, by prolonged boiling of cinnamic acid with diluted sulphuric acid (Fittig Ber 12, 1739-1744; Erdmann, Annalen, 216, 179-199).

A colourless optically inactive oil, with a blue fluorescence, which gradually disappears; sp gr. 1016 at 15°; b.p. 311° On prolonged boiling yields toluene, isopropylbenzene, and some styrene; whereas metastyrene, produced from styrene on keeping, or heating in sealed tubes at 200° (Kovalevsky, Annalen, 120, 66), or with aqueous sodium bisulphite at 110° (Miller, *ibid.* 189, 341), passes almost quantitatively into styrene at 320° (Stobbe and Posjnak, ibid 371, 259-286)

Distyrene yields benzoic acid on oxidation with chromic acid; and a crystalline dibromide C<sub>16</sub>H<sub>16</sub>Br<sub>2</sub> (m p. 102°) on brominating in carbon disulphide solution

According to Stobbe and Posinak (Annalen, 371, 287-302), distyrene is  $\alpha \gamma$ -diphenyl- $\Delta^{\alpha}$ -butene

$$C_6H_5-CH=CH-CH(CH_3)-C_6H_5$$

Distyrene disulphonic acid (dicinnamene disulphonic acid) C16H14(SO3H)2, described by Lange as obtained on heating  $\gamma$ -truxillic acid with sulphuric acid (Ber 27, 1410-1415), is therefore not a derivative of distyrene

Solid distyrene; mp 124°; obtained by distilling cinnamic acid or its calcium salt (Engler and Leist, Ber 6, 256; Miller, Annalen, 189, 340), and by passing C<sub>6</sub>H<sub>5</sub>C<sub>2</sub>H<sub>3</sub>Br, over red-hot lime (Radziszewski, Ber 6, 494), has been shown by Erlenmeyer (Annalen, 372, 249-251) to be stilbene.

DISÚLPHO ACID S. Delta acid, 1-Naphthylamine-4 8-disulphonic acid v NAPHTHALENE.

DITA BARK. Satween (Ph J'Trans [3] 7, 110) The bark of Alstonia scholars (R Br), an apocynaceous tree found in India, the Philippines, Eastern Australia, and tropical Africa (Ber 11, 2234)

According to Harnack (ibid Chem Soc Trans [2] 36, 332), it contains only one base, ditaine, a basic glucoside, having the composition C<sub>22</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub>; but Hesse (Annalen, 203, 144; Ber 13, 1750, Pharm J [3] 11, 251, 331) found in it three bodies ditamine,  $C_{16}H_{19}NO_2$ ; echitamine (cchitammonium hydroxide),  $C_{22}H_{28}N_2O_4$ ,  $H_2O$ , melting with decomposition at 106° (this base is identical with Harnack's ditaine), and echitenine C<sub>20</sub>H<sub>27</sub>NO<sub>4</sub>, a brown amorphous mass, melting above 120°, soluble in chloroform, alcohol, and ether dissolves in nitric acid with the production of a purple-red colour which quickly passes through green into yellow

The milky juice of this tree has the same properties as guttapercha; when inspissated it softens in boiling water, is soluble in turpentine and chloroform, and receives and retains impres-A decoction of the bark is used as an sions astringent tonic.

DITAINE v DITA BARK
DITHION. Trade name for sodium dithiosalicylate: used as an antirheumatic.

DITTANY. The root of Dictamnus albus (Linn) (ord Rutacea, a native of the south of Europe It contains a volatile oil, and also resin, bitter extractive matter, and gum, and was formerly employed in medicine, especially for the treatment of epilepsy (Aldis, Lond. Med. Gaz 19, 142).

'Cretan Ditany' is the common name for Origarum dictamnus (Linn ), a native of the Levant DIURETIN. A compound of sodium 570 DIURETIN.

theobromine and sodium salicylate (v. Synthetic DRUGS)

DIVICINE v. VICINE.

DIVI-DIVI. Divi-divi consists of the dried pods of the Cæsalpınıa corıarıæ (Willd), a tree 20-30 feet high, found in the West Indies and Central America The pods are about 3 inches long and 3 inch broad, are very thin, and frequently resemble in shape the letter S From 40 to 45 pc of tannin is present, which consists of ellagitannin and probably also gallotannin. Extracts of this material have a somewhat unfortunate tendency to ferment, with simultaneous development of a deep-red colouring matter, but this can be prevented to some extent by the use of antiseptics Divi-divi is largely imported for the preparation of leather, and is also employed for black dyeing, but its use is far more limited in this latter respect than myrabolans

DOCIMACY or DOCIMASY. (Docimasie, Fr.) Originally the act or art of examining by test (δοκιμασία, a test), applied in metallurgy to experiments made to determine the nature and purity of a metal; in pharmacy, to the testing of medicines and poisons, in medical jurisprudence, to the different proofs to which the lungs of a new-born child are subjected, for the purpose of ascertaining whether it was born alive or dead (Dunglison) Usually employed in chemistry as synonymous with assaying

## n-DODĚCANĚDICÁRBOXYLIC ACÍD

CO2H(CH2)12CO2H

Prepared by hydrolysing the diethyl ester obtained by the electrolysis of an aqueous solution of the monoethyl ester of suberic acid (Brown and Walker, Annalen, 261, 123), or by the oxidation of chaulmoogric acid, obtained from chaulmoogra seeds (Barrowcliffe and Power, Chem Soc Trans 1907, 568), mp 123°.

DODECATOIC ACIDS C<sub>12</sub>H<sub>24</sub>O<sub>2</sub>.

1. Lauric acid, found by Marsson in laurel fat, a glyceride obtained from Laurus nobilis (Linn) (Marsson, Annalen, 41, 330), also present in cocoa-nut oil (Gorgey, ibid 66, 295), in pichurin beans, in Dika bread, and in the fruit of Cylicodaphne sebifera (Oudemans, Zeitsch 1867, 256); mp 435, (mp 475°-48°, Levene and West); spgr 0883/20° and 0875/436°; bp 2255°(100 mm)

2 Horderc acid is found among the products of the distillation of dried barley with dilute sulphuric acid; forms crystalline plates; m p 60° (Beckmann, J. 1855, 513). DOG-TOOTH SPAR. The scalenohedral form

of calcspar (v Calcite or Calcspar) **DOLERITE.** An igneous rock composed essentially of plagioclase-felspar (labradorite to anorthite) and augite, and of the same general chemical and mineral composition as the coarsegrained plutonic gabbros and the fine-grained volcanic basalts, but intermediate in texture between these two extreme types. Other minerals may also be present, especially olivine, giving the varieties olivine-dolerite, enstatite-dolerite, hornblende-dolerite, mica-dolerite, and quartz-dolerite. Accessory minerals, usually present, are magnetite and ilmenite, and secondary minerals, produced by alteration, include quartz, calcite, epidote, chlorite, ind serpentine. Dolerites occur as intrusive veins,

dykes, and sills; and very often they pass imperceptibly into basalts. The texture may be coarse enough for the individual mineral to be distinguishable to the unaided eye, or it may be so fine that these can only be distinguished in thin sections under the microscope In the latter case the rock is compact and dark greenish-black or black in colour. Often, owing to the presence of secondary minerals, the colour is distinctly green, and the rock is then frequently included under the general term 'greenstone.' The specific gravity ranges from 2 8 to 3 0, and the crushing strength is about 1500 tons per square foot The following analyses are of dolerite from I, Rowley Regis, Staffordshire; II, Clee Hill, Shropshire; III, quartz-dolerite from the Whin Sill, Co Durham (including MnO 0·16, CO<sub>2</sub> 0·19, P<sub>2</sub>O<sub>5</sub> 0·25, FeS<sub>2</sub> 0·49), IV, olivine-dolerite from Skye (including MnO 0 17, BaO 0 06, P<sub>2</sub>O<sub>5</sub> 0 17)

	I.	II.	III.	IV.
SiO,	490	48 4	$51\ 22$	$50 \ 33$
$T_1O_2$	28	3 1	242	181
$Al_2\bar{O}_3$	$15\ 3$	13 4	14 06	19 97
$\text{Fe}_2\text{O}_3$	26	40	4 32	281
FeO	8 2	8 5	8 73	623
CaO	82	86	8 33	$8 \ 03$
MgO	49	6.5	4 42	324
K <sub>2</sub> O	26	2 1	1 25	1 19
Na <sub>2</sub> O	25	3 1	2.55	4 30
$H_2O$	3 3	2 2	1 28	1.86

Dolerites are of wide distribution and are much quarried for road-metal and paving-setts. Being difficult to work and of a dull colour, they are not much used as building stones (v DIABASE) L J. S

This term includes both dolo-DOLOMITE. mite-spar and dolomite-rock, both of which are often referred to simply as dolomite They bear the same relation to one another as calcite (calcspar) to limestone, and as quartz to quartzite; these being cases in which a single mineral species builds up a rock-mass. The name dolomite, after the French geologist, D. G. Dolomieu (1750-1801), was first applied to the rock, being later extended to the crystallised mineral The rock is now often referred to as dolomitic limestone or magnesium limestone.

The pure mineral (Dolomite-spar) is a double salt of calcium and magnesium carbonates, CaMg(CO<sub>3</sub>)<sub>2</sub>; containing CaCO<sub>3</sub> 54·28, MgCO<sub>3</sub> 45·72 p.e., or CaO 30·42, MgO 21·86 p.e. It is a member of the isomorphous series of rhombohedral carbonates (calcite group), and has an angle of 73° 45' between the surfaces of the perfect rhombohedral cleavages It, however, differs from the other members of the group in the degree of symmetry of its crystals, which are hemihedral with parallel faces. It further differs from calcite in the fact that it never shows lamellar twinning on the obtuse negative f The mostusual form of rhombohedron crystal is the primary rhombohedron, the faces of which are parallel to the cleavage directions (this being an unusual form in calcite, except in combination with other forms). The faces are frequently curved, sometimes to such an extent that the crystals are saddle-shaped. The most perfect crystais, such as those found 11 the crystalline dolomite rock of the Binnenthal DOLOMITE.

in Switzerland, are colourless and transparent with a vitreous lustre. More often the crystals are white or creamy-white and opaque with a pearly lustre; hence the popular name pearl-spar The mineral is harder (H. 3½-4) and denser (sp gr 2 85) than calcite 1t may be readily distinguished from the more commonly-occurring calcite by the fact that it is only slightly attacked by cold dilute acid; in hot acid it dissolves with a brisk effervescence Dolomite is much less soluble in water containing carbon dioxide (3 parts in 10,000 at 18°) than is calcite Various micro-chemical tests have been devised for the purpose of distinguishing dolomite and calcite, especially in thin sections of rocks Lemberg's solution contains aluminium chloride and a staining material (haematoxylin or an aniline dye); a stained film of aluminium hydroxide deposits on calcite, but not on dolomite composition of dolomite may be varied somewhat by isomorphous replacement Iron and manganese may replace magnesium, forming a passage through brown spar to ankerite (qv) Zinc or cobalt is also sometimes present in small amount. Any excess of calcium over the double salt ratio may perhaps to some extent be present in isomorphous mixture, but more often it is probably present as mechanically admixed calcite Simple ratios of CaCO<sub>3</sub> MgCO<sub>3</sub> other than the 1·1 of 'normal dolomite,' probably do not exist Several trivial names have been applied to varieties of dolomite showing pecuharities of structure or supposed differences in composition, the analyses have not, however, always been made on well-crystallised material

The best crystallised specimens of dolomite are met with in mineral veins and other metalliferous deposits. For example, with iron ores at Traversella in Piedmont, and Cleator Moor in Cumberland; with lead and zinc ores at Laxey in the Isle of Man, and Jophin in Missouri, with silver ores at Schemnitz in Hungary, and Guanaxuato in Mexico Crystalline masses occur in serpentine, tale-schist, and other magnesian silicate rocks. As a rule, only quite small crystals are found in dolomite rocks, but occasionally good crystals are found in cavities.

Dolomite rock, dolomitic or magnesian limestone, is, as a rule, more indefinite in its characters and composition, since here the dolomite mineral is mixed with other mineral substances, such as calcite, carbonates of iron and manganese, oxide and hydroxides of iron, argillaceous matter, quartz, etc. With increasing calcite the rock may pass into ordinary limestone, and some authors restrict the term 'magnesian limestone' to rocks intermediate between 'normal dolomite' and ordinary limestone. The presence of calcite in a magnesian limestone containing a molecular excess of calcium over magnesium may be demonstrated by etching a polished surface or a thin section with cold dilute acid, when the calcite will be readily dissolved out leaving pits; or by means of the staining test with Lomberg's solution mentioned above

The purest dolomite rocks are the snow-white crystalline dolomite marbles of the Swiss Alps, Norway, &c, which have sometimes the character of a statuary marble and can be used for ornamental purposes More often, however, the rock is yellowish, grey, creamy, or brown, and very often cavernous in texture. It is

usually minutely crystalline, consisting of an aggregate of tiny rhombs of dolomite-spar more or less firmly held together, but at times quite loose and sandy Dolomite rock is of wide distribution and is met with in sedimentary strata of all geological periods In England a long narrow outcrop extends along the eastern slopes of the Pennines from Nottinghamshire to the Durham coast. This has long been extensively quarried as a building stone, the principal quarries being near Anston in Yorkshire, Bolsover Moor in Derbyshire, and Mansfield Woodhouse in Nottinghamshire Here the rock is of Permian age In the Dolomite mountains of southern Tyrol the rock is of Triassic age, whilst that of Swabia belongs to the Jurassic formation

571

The origin of dolomite rock has been a subject of much speculation and experimental work of a chemical character. A review is given in Doelter's Handbuch der Mineralchemie, 1912, vol 1 (here also 1s given a large collection of analyses, methods of analyses, &c) That of Permian age, which is usually associated with beds of gypsum and rock-salt, is supposed to have been deposited as the double carbonate by chemical precipitation from the salt waters of mland seas In some other cases the rock has been formed by the action of magnesium-bearing solutions on limestone, either contemporaneously and progressively with the formation of the limestone (as proved by the dolomitisation of coral-limestone-the magnesium in this case being derived from sea-water), or long subsequently by the action of subterranean mineralwaters percolating along the bedding and joint planes (as in the Carboniferous Limestones of Derbyshire and South Wales) Or again, it may have been formed by weathering and the solution of the more soluble calcium carbonate from limestones containing a small proportion

The following are analyses of dolomite rock I, from Bolsover, Derbyshire, II, Mansfield, Nottinghamshire, III, Anston, Yorkshire; IV, Southern Tyrol—

i			I	II	III	IV
CaO			28.6	29.0	30.7	31 4
Mg()			19.2	20 4	20 6	192
Fe,O,	, Al,	$_{2}O_{3}$	18		0.73	12
SiO			3 6	3 7	0.56	10
co,		. 4	43 5	44 9	46 7	46 l
H,Ö			3 3	2 1	0.75	

The economic applications of dolomite are confined to the rock, since this is much more plentiful and of wider distribution than the crystallised mineral Its principal use is as a building stone For this purpose it is more suitable than ordinary limestone, being harder and more resistant to solvents, but it needs to be carefully selected It is also used for paving and road-metal The Houses of Parliament are built of stone from the three English localities noted above (Anal I-III) The stones from these quarries have a crushing strength of 484-883 tons per square foot, absorption of 4.6-7.5 pc, and weigh 134-152 lbs per cubic foot Certain varieties of dolomite rock (dolomite marble) find an application as an ornamental Dolomite rock is also extensively used 8.one for the basic linings of Bessemer converters and

572 DOLOMITE

open-hearth furnaces in the manufacture of For this purpose the rock is calcined until it commences to sinter, crushed, mixed with gas-tar, and moulded in the furnaces or The best result is obtained shaped into bricks when the rock contains 3-4 p c of Fe<sub>2</sub>O<sub>3</sub> and Al<sub>2</sub>O<sub>3</sub>, and about  $2\frac{1}{2}$  p c of SiO<sub>2</sub> (On British sources of dolomite rock for use as a refractory material, v Mineral Resources of Great Britain, Mem Geol Survey, 1918, vol v1) The rock is completely burnt or half burnt below a red heat (giving MgO-+CaCO<sub>3</sub>) for the preparation of cements and plasters, when it contains quite a small amount of argillaceous matter it gives a good hydraulic cement It is also burnt for agricultural purposes, and is said to be especially beneficial for leguminous crops It is a readily available source for the manufacture of magnesium compounds, eg Epsom salts, magnesium chloride, magnesia alba, but in this direction it was not able to compete with the large quantities of soluble magnesium salts obtained as by-products from the Prussian salt-deposits Various processes have been suggested for extracting the magnesia for use as a refractory material In one, the calcined dolomite is forming mixed with magnesium chloride, calcium chloride which is readily removed in In another, the calcined dolomite is treated with a solution of sugar, when the calcium passes into solution as saccharate, from which the sugar can be recovered by precipitating the calcium as carbonate with carbon dioxide When half burnt to  ${\rm MgO+CaCO_3}$ the magnesia can be extracted by carbonated water under pressure

The pure white dolomite marble is used for the manufacture of pure carbon dioxide for use in the manufacture of aerated waters LJS

**DOMEYKITE.** A copper arsende, Gu<sub>3</sub>As, corresponding to copper 71 7 and arsenic 28 3 p c It occurs in reniform, botryoidal, and compact masses, of white or grey colour, with metallic lustre, rapidly tarnishing H=3-3.5, sp.gi 7-7.5 Readily fusible before the blowpipe, with emission of arsenical vapours Soluble in nitrie acid, but not in hydrochlorie acid found at Coquimbo and Copiapo in Chile, at Cerro las Paracatas in Mexico, and at Zwickau in Saxony

Several other arsenides of copper, more or slike domeykite, are known The Algodonite less like domevkite, are known of F Field is Cu, As, and occurs at the Algodones mine, near Coquimbo. Whitneyite, from Houghton Co, Michigan, US, described by F A Genth, is Cu<sub>2</sub>As, and D. Forbes's Darwinte, named after Charles Darwin, is a similar mineral from near Copiapo Condurrite, from Condurrow mine, Cornwall, appears to be a mixture of arsemide of copper with various oxidised products derived from the alteration of tennanite or some kindred species The names Keweenawite, Mohawkite, and Ledouxite have been applied to various copper arsenides from Michigan, in which the copper is in part replaced by nickel and cobalt L J S.

**DONARITE.** An explosive consisting of a mixture of ammonium nitrate (80 pts), trinitrotoluene (12 pts), nitroglycerine (3 8 pts),

gun-cotton (0 2 pt, flour (4 pts)

DORMIGENE. A urea derivative of a-brome Used as a nervous sedative. isovalerianie wid

DORMIOL. A condensation product of chloral and tertiary amyl alcohol. V. SYNTHETIC

DOSS. Japanese name for a dyewood from an evergreen tree Ilex Mertensii (Maxim), found in the Ogasawara and Okinawa Islands. Contains a dyestuff, dossetin C<sub>15</sub>H<sub>9</sub>O<sub>5</sub>, crystallising in yellow needles; m p. 271 -272° (Ito, J Soc Chem Ind 1908, 440)

DOUBLE SCARLÉT, BRILLIANT SCARLET v Azo- colouring matters

DRACYL. An obsolete synonym for toluene DRAGON'S BLOOD. A resin used as a red pigment (v Pigments; also Resins)

DRIERS (in the United States also termed 'Japans') is the trade term for those metallic oxides, chiefly oxides of lead, manganese (litharge, red lead, manganese dioxide) and cobalt or their inorganic salts, which were formerly used exclusively in the process of oil boiling as 'driers.' More recently acetate, oxalate, and borate of manganese have been employed Since metallic salts of abietic acid have the same property, such salts are also manufactured as 'driers' They salts are also manufactured as 'driers' are obtained either by precipitating rosin soap solutions with solutions of metallic salts, or by fusing together a metallic oxide with colophony Hence these are distinguished in commerce as 'precipitated' driers, and 'fused' driers. The latter are readily discriminated from the former by their not containing any moisture salts of linseed oil fatty acids are now used as driers, they can be prepared either by precipitation or by the dry process, hence both 'precipitated' manganese linoleate, 'precipitated' lead linoleate, and 'fused' manganese linoleate, &c, are obtainable The driers prepared from linseed oil fatty acids, as also from rosin (colophony), are soluble in oil of turpentine, ether, chloroform, and linseed oil Hence they are also termed in commerce ıtself Solutions of these driers in soluble driers' linseed oil or in turpentine oil, or in a mixture of both, are obtainable in commerce under the name 'liquid driers,' 'terebene,' and other fancy names.

The function of these driers appears to be to absorb oxygen from the air and transfer it to the linseed oil, thus accelerating its drying to a flexible film when exposed to the air this assumption, the fact that litharge and manganese dioxide act as driers will be best explained by stating that at the high temperature at which linseed oil is 'boiled,' the oxides saponify the linseed oil, forming therewith lead or manganese soaps, which then act in their turn in the manner indicated above From experiments on the behaviour of cerium salts obtained from various drying oils, Morrell (Chem. Soc Trans 1918, 113, 111) concludes that a change occurs in the intra-molecular linkings when drying oils are polymerised by heat action of cerium a-eleostearate (from tung oil) as a drier may be represented by the following scheme :-

scheme :—
$$\begin{array}{c} \operatorname{CeX}_3 \text{ or } \operatorname{CeOX}_6 \Longrightarrow \operatorname{Ce}_2(\operatorname{XO}_2)_6 \\ \operatorname{CeO}(\operatorname{XO}_3)_6 + \operatorname{drying} \text{ oil} \\ \Longrightarrow & \operatorname{CeX}_3 + \operatorname{OCHY} \\ & | & | & | & | & | \\ \operatorname{OCHZ} \end{array}$$
This representation is also in accord with the

conclusions of Ingle (J Soc Chem Ind, 1917, 318) as to the behaviour of lead linolenate Cerium driers are less effective than cobalt or manganese linoleate in accelerating the drying of a varnish containing a lead drier. The following are the analytical data obtained by the author in the examination of a drier.

	Per cent.	•
Ash	18 6	
Fatty and rosin acids .	. 834	

The more detailed examination gave the following result -

		Per cent
Lead oxide		10.99
Manganous oxide		6 79
lron peroxide	•	0 47
Calcium oxide		0 56
Rosın acıds		$38\ 32$
Oxidised acids		4 24
Other fatty acids		40.86

It should, however, be understood that the results furnished by chemical analysis alone are not sufficient to supply the data on which to base an opinion as to the properties which a drier will impart to the oil. The colour and drying power of the boiled oil to be prepared and other conditions greatly influence the choice and the quantity of a drier to be employed on a large scale.

DROP GUM or GAMBOGE v PIGMENTS

DROSERA WHITTAKERI. Drosera whittakeri is found in Australia, and grows plentifully on the hills near Adelaide. The tuber of this plant consists of an inner solid but soft nucleus full of reddish sap or juice, and an outer series of easily detached thin, and more or less dry, layers of an almost black material. Between these layers are to be found small quantities of a brilliant red colouring matter, the amount varying in tubers of different size and age, but apparently more plentiful in the older plants (Rennie, Chem. Soc. Trans., 1887, 51, 371., 1893, 63, 1083)

63, 1083)

The colouring matter is extracted from the tubers by means of hot alcohol, the solution evaporated, and the residue, containing a little alcohol, is then mixed with water and allowed to stand. The product is dried, sublimed, and the billiant vermilion powder, which contains two substances, is fractionally crystallised from boiling alcohol, or acetic acid.

The more sparingly soluble compound  $C_{11}H_8O_5$  forms red plates, melts at 192°-193°, dissolves in alkaline solutions with a deep red violet colour, and gives a triactifi derivative,  $C_{11}H_5O_5(C_2H_4O)$ ,, melting-point 153°-154°

 $C_{11}H_5O_5(C_2H_3O)$ , melting-point  $153^\circ$ - $154^\circ$  The monosodium compound  $C_{11}H_1O_5Na_2H_2O$ , gives dark reddish-brown needles, the disodium compound  $C_{11}H_6O_5Na_22H_2O$ , brown needles, and the calcium compound  $(C_{11}H_1O_5)_2Ca_3H_2O$ , dark brown crystals

By treatment with stannous chloride or hydrochloric acid, this substance  $C_{11}H_{10}O_{5}$  yields a reduction product  $C_{11}H_{10}O_{5}$ . yellow needles, melting-point  $215^{\circ}-217^{\circ}$ . When dry, this is stable in air, but if left in contact with alcohol or water, it soon becomes red owing to exidation According to Rennie it is probable that the original colouring matter is a trihydroxymethylmaphthogramme, in which case the substance  $C_{11}H_{10}O_{5}$  will consist of the corresponding

hydroquinone derivative. The more readily soluble compound contained in the tubers possesses the formula  $C_{11}H_8O_4$ . It crystallises in red needles, melting-point  $174^\circ-175^\circ$ , dissolves in alkaline solutions with a deep-red colouration, and gives a diacetyl derivative,  $C_{11}H_6O_4(C_2H_7O)_2$ , yellow needles, melting-point  $107^\circ-110^\circ$ .

**DRUMINE.** An alkaloid stated to be present in Euphorbia Drummondii (Ogston, Brit.

Med Jour 1887, 451)

DRY CLEANING. French or chemical cleaning, Nettoyage à Sec A process of removing dirt, grease, &c, from fabrics by solvents other than water, such as petroleum benzine, coal-tar benzene, carbon tetrachloride, &c. The substitution of such liquids for water seems to have been first made in France during the middle of the last century by the application of 'camphene,' a terpene used at that period for illuminating purposes

The process was introduced into the United Kingdom, in 1866, by Messrs Pullar & Sons, who greatly developed it, employing machinery to a large extent in place of the hand labour

formerly used

The solvents mainly used at the present time are 'benzines,' usually obtained from American and Borneo petroleum, of sp gr 0 78; boiling between 80° and 120° The benzols employed are what are known commercially as 90's and They may contain sensible quantities of carbon disulphide, thiophen, and possibly other sulphur products, which are apt to impart a disagreeable smell to the fabric Carbon tetrachloride, on account of its relatively high price, is only used to a limited extent It has the great advantage over the hydrocarbons that it is uninflammable, and can be employed in particular cases where the use either of benzine or of benzol is madmissible (For a method of valuing benzine for the purpose of the dry cleaner, v Trotman, J Soc Chem Ind 1906)

Thiophen in benzol may be detected by the isatin and sulphuric acid reaction (v Thiophen).

(Cf Schwalbe, Chem Zeit 1905)

Carbon disulphide may be detected by the formation of a precipitate with phenylhydrazine, and determined quantitatively by conversion into potassium xanthate by means of alcoholic potash (v Carbon disulphide)

An admixture of paraffin hydrocarbons with the benzol may be recognised by nitrating the sample with nitrosulphuric acid and subsequent

steam distillation.

The carbon tetrachloride employed for degreasing should be free from hydrochloric acid or traces of free chlorine. It should also contain no chloroform or phosgene gas, which are apt to

act deleteriously upon the workpeople

Special soaps, soluble in benzine, are employed in dry cleaning. They not only assist the degreasing, but diminish the risk of fire. The soaps consist of practically anhydrous oleates or stearates of potash or soda made by melting together oleic acid and soap, or by dissolving the alkali in methylated spirit, and adding to the solution the requisite quantity of oleic acid or melted stearic acid. Strong ammonia solution may be employed instead of potash or soda.

(For a description of the methods and

machinery employed in modern processes of dry cleaning, and of the apparatus for the recovery and purification of the solvents used, v. Dyeing and Cleaning, by F J. Farrell, Griffin & Co, Ltd. 1910)

**DULCAMARA.** Buttersweet, Woody night-de. (Douce amère, Fr , Buttersuss, Ger) shade. (Douce amère, Fr, The young branches of the Solanum dulcamara (Linn), were formerly administered as a decoction in rheumatism and skin diseases contain an amorphous glucoside dulcamarin C<sub>22</sub>H<sub>34</sub>O<sub>10</sub> (Geissler, Arch Pharm [iii.] 6, 1010); 

DULCIN. p-Phenetol-carbamide

C2H5O C6H4 NH CO NH2

Used as a substitute for A sweet substance saccharın

DULCITOL v CARBOHYDRATES. DUMASIN. Pentamone v Ketones DUNGING SALT. Sodium Arsenate v ARSENIC

**DUOTAL.** Guaracol carbonate

formed by the action of phosgene gas on an alkaline solution of guaracof Used in the treat-

ment of phthisis

DURA or DARI. A gram imported from India and the Sudan, chiefly as a poultry food position: moisture, 8 45, proteins, 12 98, other ntrogenous substance, 0 08; fat, 3 30, starch, 62 66; other carbohydrates, 9 74, fibre, 1 03, ash, 1 71 pc, food units 113 3, nutrient ratio, 1 · 6 · 1 Contains no alkaloids or cyanogenetic glucosides (Bull Imp Inst. 1913, 11, 33) the use of Sudan Dura as a brewing material, see Briant & Harman, Journ Instit Brewing, 1918, 209

DURENE v. CYMENES

**DURENOL** v PHENOL AND ITS HOMOLOGUES DUTCH PINK or ITALIAN PINK v Pig-

DUTCH ULTRAMARINE. Cobalt blue v COBALT AND PIGMENTS

**DURINE.** A preparation of formaldehyde **DYEING.** The art of dyeing consists in imparting colour to the substance of various

materials, usually textile fabrics, in such a manner that it is not readily removed or altered by those influences to which the dyed material is subsequently exposed Tinting is essentially the same as dyeing, but the term is applied to the production of light shades only. In staining, the colour is not so permanently fixed as in dyeing, but in all these processes the colouring substance is presented to the material in a dissolved condition. In textile printing, the colour is applied to portions of the material only, thus producing a pattern, but in other respects, many printing operations are, in principle, similar to dyeing. In some processes of textile printing, as in painting, on the other hand, a more or less opaque pigment is applied to the surface of the material by means of adhesives, thus partially or entirely obscuring the true surface.

Dyeing is of chief importance in connection

with the textile industries, but many other materials are coloured by dyeing processes, such, for example, as leather, feathers, horn, ivory, bone, straw, and wood.

The various materials above enumerated remarkable differences in behaviour towards any particular dyestuff. Wool, silk, leather, feathers, horn, and animal skin products in general, behave somewhat similarly towards colouring matters, but the dyeing properties of cotton, linen, and other vegetable products are quite different.

Differences quite as marked are to be noted in the behaviour of various colouring matters towards one and the same material, and thus, in dyeing operations, both the nature of the material to be dyed and of the colouring matter

employed, have to be taken into account

The theory of dyeing. The precise cause or causes of the different behaviour of the textile fibres in dyeing is still a matter of uncertainty, but there is little doubt that the chemical and physical properties both of the fibre and of the dyestuff are all factors of the complicated phenomena of dyeing

The earliest theory with regard to the fixation of dyes was that the action is purely mechanical, and it is still urged by many that rarely in dyeing processes is there direct evidence of chemical combination between fibre and dye-Both bodies, as a rule, retain their individual chemical and physical properties unchanged, and the dye can frequently be removed from a fibre by means of alcohol or other solvent

In its earliest form, the mechanical theory was put forward by Hellot and d'Apligny in 1780 (See Hellot's L'Art de la Temture des Lames (1734), and d'Aphgny's L'Art de la Teinture des fils de Coton (1776)) They assumed that the particles of colouring matter became embedded in the minute pores permeating the fibre substance Crum (Chem Soc Trans. 16, 404) strongly supported this theory, while Muller-Jacobs (J Soc Dyers, 1885, 63) expressed much the same idea in attributing the fixation of dyes to membranous diffusion. It has, however, been shown by several investigators that dyeing processes do not obey the general laws of osmosis. v Georgievics (J Soc Chem Ind 1895 149) considers dyeing is chiefly due to surface attraction Witt (Farber-Zeit 1890) put forward a modification of the mechanical theory. He regards the fibres as solid solvents in which the dyestuffs dissolve, the latter being withdrawn from its aqueous solution in the same way as, eg. ether will extract many substances from their aqueous solution The degree to which a dye-bath can be exhausted will then depend upon the relative solubility of the dyestuff in the acid or neutral dye liquor and in the fibre substance under the conditions (concentration and temperature) used Krafft (Ber 1899, 1608) attributes dyeing phenomena to a separation of the dyestuff upon the fibre in a colloidal state, and this view has been supported by Biltz (ibid 1904, 1766), Linder and Picton (Chem. Soc. Trans 1905, 1931), and Dreaper (J Soc. Chem. Ind 1905, 223)

The chemical theory of dyeing was originally put forward by Macquer and Bergman in opposi-tion to the mechanical theory of Hellot and d'Apligny, and has received much experimental

support from Persoz, Berthollet, Chevreul, Kuhlmann, Vignon, Knecht, and many others It is mainly based on the following facts: wool and silk fibres contain amino-acids, and thus possess strongly marked acidic and basic functions Cellulose, the essential constituent of the vegetable fibres, is chemically much less active, and lyes much less readily than wool and silk, but under certain conditions it exhibits weakly acidic All dyestuffs (with few exceptions) possess either acidic or basic properties, or are in the form of salts, which dissociate in the dyebath, and chemical combination between fibre and dyestuff would thus, primâ jacie, be expected Moreover, lanuginic and sericinic acids, which are produced respectively by the hydrolysis of wool and silk, precipitate both acid and basic dyes from their aqueous solution (Knecht and Appleyard, J Soc Dyers, 1889, 72)

A full discussion of the evidence for and against the mechanical and chemical theories of dyeing cannot be here entered upon, but a general review indicates that the attempt to cover all dyeing phenomena by any one simple explanation must fail, and that many factors are concerned, the relative importance of which varies in different cases In the dyeing of wool or silk, with acid or basic dyes, it may be taken as established that chemical combination occurs, but prior to this some hydrolysis of the fibre, and more or less complete ionisation of the dye-Surface concentration of the stuff takes place dye on the fibre probably also occurs, and the coagulation and precipitation of the dissociated colloidal dyestuff also plays a part supported by the observation that the temperatuie of maximum absorption of dye does not (in the cases investigated) correspond with that of maximum fixation (Dreaper, Trans Faraday Soc 1910, pt 1 Haldane-Gee and Harrison, ibid); the former possibly coinciding with the point of maximum coagulation and precipitation, and the latter indicating the point of maximum degree of chemical combination. Whether the superficially precipitated dye penetrates the fibre by osmosis or by 'solid solution' is mainly a question of definition of terms

In the case of the dyeing of vegetable fibres, the same factors probably operate, but their relative importance is not the same, the physical factors—precipitation, surface attraction, and osmosis—predominating over the effect of direct chemical action

Classification of colouring matters. It has already been pointed out that dyestuffs vary in their behaviour If, e g separate pieces of wool are boiled in solutions containing,  $(\hat{a})$  logwood, (b) indigo, (c) magenta, and (d) pieric acid, the following results are obtained: (a) the wool acquires a brown stain; (b) the wool remains undyed, the indigo floating about in an insoluble condition (c) the wool is dyed a magenta colour; and (d) the wool is dyed a full yellow If the experiment is repeated on wool which has been previously boiled with a solution of potassium dichromate, the result will be that in (a) the wool is dyed black, whereas in (b), (c), and (d), similar results are obtained to those of the Logwood, therefore, previous experiments must be applied in conjunction with potassium dichromate or some similar substance, i.e. with a mordant; whereas magenta and picric

acid dye without such aid, and indigo requires a special process, the nature of which is not indicated by the results of the above-described experiments.

Colouring matters may be divided into three groups on the lines indicated above, viz. (1) those which must be applied in conjunction with a mordant, (2) those which dye direct from their solutions; and (3) those requiring special processes

As regards groups (1) and (2) this classifica-tion is practically identical with that of Bancroft (Philosophy of Permanent Colours, 1 118, London. 1794), who, upwards of a hundred years ago, divided colouring matters into substantive dyes and adjective dyes, the former class comprising those which dye directly from their solutions, and the latter being composed of those dyes which require mordants Nietski (Farber-Zeit 1889-90, 8) uses the terms direct dyes and mordant dyes in exactly the same sense, and both Bancroft's and Nietski's terms are still in use. Hummel (The Dyeing of Textile Fabrics, 1st ed. 1885, 147) also classifies dyestuffs into two groups. monogenetic dyes being such as are capable of producing one colour only, while polygenetic dyes produce distinct colours with different mordants The members of the first group are coloured bodies, and these merely require to be fixed upon the fibre, whereas the colouring principles comprising the second group as a rule only develop a colour when chemically combined with a mordant

It will be seen that these three systems of classification closely correspond, the terms substantive dyes, direct dyes, and monogenetic dyes being almost synonymous, as are adjective dyes, mordant dyes, and polygenetic dyes. It should, however, be pointed out that none of these classifications is absolute, magenta, for example, is a direct dye as applied to wool, but requires a mordant when used for cotton.

A distinct method of grouping dyestuffs is that of Benedikt (The Chemistry of the Coal-tar Colours, 1886, 29), who distinguishes between colour acids, colour bases, and neutral colouring Both colour acids and colour bases are fixed on the fibre in the form of salts, but whereas in the colour acids the essential colourproducing group resides in the acid portion of the compound, in the colour bases the reverse is Alizarin ied, dyed on wool, is a salt of which alumina forms the basic and alizarin the acidic constituent Alizarin is therefore a colour acid. Similarly, the red pigment formed on dyeing tannin-moidanted cotton with magenta is a combination of rosaniline base with tannic acid Magenta is therefore a colour base. Indigo and other vat dyes are neutral colouring

All the above schemes of classification are defective, either from the point of view of the practical dyer, or the chemist As regards practical application, the large number of dyestuffs now on the market may, however, be satisfactorily classified into the following groups: (1) mordant dyes, (2) acid-mordant dyes; (3) acid dyes; (4) direct dyes; (5) basic dyes; (6) sulphide dyes; (7) vat dyes; (8) ingrain dyes.

The classification of dyes according to chemical constitution is based upon the existence,

of a certain group of atoms, in all members of each class, upon the integrity of which their colouring power depends. This is termed the chromophorous group, the theory being first put forward by O N Witt, in 1876 The introduction of such a chromophor into a suitable colourless compound is accompanied by the development of colour, eq benzene  $(C_6H_6)$  is colourless, but azobenzene  $(C_8H_5-N=N-C_6H_5)$ is orange-coloured Many substances, however, such as azobenzene, although strongly coloured, are devoid of dyeing power Such bodies are termed chromogens, and in order to convert them into useful dyes, the introduction of a saltforming group is necessary, such groups, in this connection, being termed auxochromous groups Thus, whilst azobenzene is not a dyestuff, both amino azobenzene and hydroxy-azobenzene are useful dyes In these compounds, the azo- group (-N=N-) is the chromophor, and the amino- or hydroxyl- group is the auxochrome

The presence of both a chromophor and an auxochrome does not, however, necessarily indicate that a substance is a useful dye For example, trinitroaniline,  $C_0H_2(NO_2)_3NH_2$ , contains the chromophor  $(NO_2)$  and the salt-forming auxochrome group  $(NH_2)$ , but it is not a useful dyestuff, because the basic character of the auxochrome is neutralised by the strongly acidic chromophor. On the other hand, trimtrophenol  $C_0H_2(NO_2)_3OH$  is the useful dyestuff pieric acid.

Those dyestuffs which do not contain a salt-forming auxochrome cannot be applied by ordinary dyeing processes, and the fact that a special process is required,  $e \, q$  for indigo, is not primarily due to its insolubility, but to its neutral character.

The scheme of classification, based on the chromophor, distinguishes the following groups—

Typical dyes

Acridine yellow, orange, and

Meldola's blue, gallocyanin,

mine, hydhron blue

scarlet

Capri blue.

Methylene blue,

Chromophor

Group

7. Acridine

8 Oxazine

9. Thiazine

1. 180-Nitroso->C-O Resorein green, dioxine, gambine or oxime >с-й он 2 Nitro--NO. Pieric acid, naphthol vellow 3 Azo--N-N- (a) Basic-chrysoidine, Bismarck brown, Janus red Acid-azo- oranges and scarlets, naphthol black, diamond black
(c) Direct—benzopurpurin, diamine colours, direct blacks (d) Mordant—'acid mordant' or 'chrome colours methyl violet, blue, patent 4. Triphenyl Magenta, Victoria methane blue, malachite green 5. Oxyketone CO The alizarm dyes, indan-threne and algol vat dyes 6. Pyrone Rhodamine, violamine, cosins.

12. Quinoline —CH—CH Quinoline yellow, cyanine

N<sub>2</sub>CH

13. Flavone —O—C— Fustic, weld, logwood
Brazil wood

15. Miscellaneous

This classification of dyestuffs into chemical groups is of importance in the identification and testing of dyestuffs, as the constitution of the compound determines its behaviour towards reagents. It is not, however, a convenient one as regards the practical application of the dyestuffs.

Essential properties of a dyestuff. In order to rank as a useful dye, a substance must possess certain essential features A direct dye must not only possess a colour of considerable intensity and permanence, but must also be capable of easy fixation on textile fibres without mjury to fibre or dyestuff For example, both vermilion and azo-scarlet possess a brilliant colour, but the former is not a dyestuff, because it cannot be applied or produced by dyeing processes. A mordant dye, on the other hand. need not possess a strong colour, and may, in fact, be colourless. It must, however, be capable of producing strongly coloured and insoluble colour-lakes in combination with mordants. Vat dyes-e g indigo-are, as a rule, quite insoluble in water, and therefore cannot be applied by an ordinary process The essential feature of this group is that their reduction products are soluble in alkaline solutions, and on this fact the method of application depends

In order to constitute a dyestuff a body must, therefore, either be strongly coloured in itself or be capable of producing coloured lakes. It must also be of such a character that it can be fixed upon textile material from an aqueous solution, and it must produce a colour which possesses sufficient permanence and resistance to washing and other influences. (In so-called 'dry dying' alcohol or other volatile solvent is used)

Operations preliminary to dyeing processes. In any dyeing operation, the chief bodies concerned are the fibre to be dyed and the dyestuff. In their natural condition, all textile fibres contain impurities which not only detract from their appearance, but prevent their being satisfactorily dyed. The removal of these impurities is the object of the washing, scouring, and bleaching processes which the 'raw' fibres undergo.

A great bulk of manufactured textile material chiefly cotton and linen, is used in the undyed condition, but wool and silk are usually dyed

Dyeing processes. Dyeing may take place at various stages of the manufacture of textiles 'Piece dyeing' is generally employed when a plain-coloured material is required, being the most economical and convenient process. If a coloured pattern is required, the material must either be printed, or dyed before weaving, and

he process may then be carried out on the raw rool or cotton ('wool dyeing,' cotton dyeing'), t an intermediate process ('sliver' or 'slubbing yeing') or upon the spun yarn ('yarn dyeing,' hank dyeing,' 'cop dyeing,' 'cheese dyeing,' 'u arf

Wool is dyed in all these forms; silk, almost xclusively in the form of yain, and cotton,

hiefly as warps, cops, or pieces.

The dyeing of wool. Wool may be dyed 71th a greater range of colouring matters than ny other textile fibre. All the various classes f colouring matters enumerated in the classifiation based on practical application (see p. 575) re used in the dyeing of wool, the most apportant groups being the mordant, the acid wordant, the acid and the direct dyes, as well as he *i at dye*, indigo. In boiling solutions, the ffinity of wool for dyestuffs is generally greater han that of silk, but at temperatures below 60° he relative affinity is frequently reversed. By suitable selection of dyestuffs and control of the rocess, it is therefore possible to dye the wool and ilk in a woven fabric either to different shades, r to produce a uniform colour on the two fibres s a general rule, dyeing processes which produce ood results on wool are entirely unsuited for otton, and it is an easy matter to dye one of the wo fibres when present in a mixed fabric and It both ave the other practically unstained bres are to be dyed the same or different colours, wo distinct processes are, as a rule, required

Different varieties of wool behave differently 1 dyeing, the finer wools, such as merino, 1euning a larger amount of dyestuff to produce a iven shade than the coarser qualities,  $e[\hat{q}]$  Lincoln r 'crossbred' wool The absorption of dye is, owever, by no means entirely governed by the nameter of the fibre, eg fine silky mohair is ery difficult to dye, probably on account of the lose texture of the external scales, Fibres aken from different portions of the same fleece lso vary somewhat in dyeing properties, and ool taken from the skin of animals slaughtered or meat, or from diseased animals, also differs om 'fleece' wool as regards its affinity for dyes Il these differences are shown to a much greater stent with some dyestuffs than with others

The affinity of wool for certain dyestuffs may e increased by suitable treatment with acids, lkahs, or oxidising agents (bleaching powder r hydrogen poroxide solution), or may be immished by a treatment with tannic acid olution and in other ways. Various colours may us be produced by a 'piece-dyeing' process y weaving together untreated wool and wool eated with one or other of the above-mentioned

The behaviour of wool towards solutions of netallic salts governs the use of these bodies as nordants. When boiled in solutions of salts f the heavy metals having an acid character g. sodium dichromate), a decomposition of ne salt takes place, and an insoluble basic salt r a hydrated oxide is fixed upon the fibre.

The dyeing of silk. In general behaviour wards colouring matters, silk resembles wool, ut, as a rule, dyes are fixed on silk in a less ermanent manner than on wool. The dyeing f silk appears to depend, to a greater extent, n absorption, and to a less extent on chemical ction. The acid, basic, and direct dyes are TTOT TT

chiefly used on silk, but other classes, eg. the mordant and vat dyes, are also employed, though they tend to detract somewhat from the brilliant lustre and suppleness on which the beauty of silk depends. The maximum absorption of dyes by silk usually takes place at 60°-80°, and

not at 100° as in the case of wool.

The so-called 'dry dyeing' process sometimes employed for silk consists in the use of benzine or other volatile liquid as a solvent for the colouring matter in place of water. The usual dyestuff salts are not soluble in benzine, but the fatty acid compounds or resinates of basic dyes are readily soluble.

The mordanting of silk with metallic salts is usually carried out in cold concentrated solutions of basic salts, chiefly basic aluminium sulphate for colours, and basic ferric sulphate or

ferrous acetate for blacks

The production of black upon silk is a matter of great importance, and is almost a trade to itself. The silk is first immersed in a solution of chestnut extract or other tannin matter, and then in a solution of basic ferious acetate (pyrolignite of non) or ferrie sulphate ('nitrate of iron') By successive alternate treatments in these liquors, the silk may be weighted to the extent of 400 p.c. It is then dyed with logwood extract There are a number of alternative processes for the black dyeing of

Wild silks, of which Tussur is the chief variety, are much more difficult to dve than ordinary silk, the fine fibrillæ of which the fibre is composed being very impervious. The bandlike flat surface of the fibre also reflects the light in such a manner that a white shiny appearance is liable to be produced instead of a rich, intense colour or black

Artificial silks, consisting essentially of cellulose, are more allied to cotton than to silk as regards dyeing properties. Those varieties produced from collodion (Chardonnet silk) dye more reaculy with the basic dyes than do cotton or the artificial silks obtained from a cuprammonium solution of cellulose (Pauly, Lehne, Thicle or 'glanzstofle' silk) or from cellulose xanthate ('viscose silk') There is, however, much individuality in the dycing properties of the various forms of artificial silk

The dyeing of cotton. Cotton is died chiefly in the form of hanks or warps, less frequently as loose cotten or piece goods The dyeing of cotton on the spool, cop or beam is, however, now of

great importance

The dyeing properties of cotton differ greatly from those of wool, and therefore the processes and materials used also differ. Until the discovery of Congo red in 1881, there were three dyestuffs only which were capable of dyeing cotton directly from their aqueous solution turmeric, safflower, and annatto-but there are now a large number of synthetic substantive dyes by means of which any desired shade can be obtained. The very numerous class of acid dyes which are of such great importance in wool and silk dyeing do not produce reasonably fast colours on cotton.

Cotton has no affinity for metallic salts, and therefore does not become mordanted by boiling in their solutions. Indirect methods have therefore to be adopted in applying the

basic mordants required for the fixation of the mordant dvestuffs on cotton This fibre exhibits a remarkable affinity for tannic acid, which is withdrawn from dilute solution and concentrated on the fibre to a considerable extent The tannic acid may then serve for the fixation either of metallic mordants or of basic dyestuffs upon the fibre

The vat dues and sulphide dyes are now of first importance, since, by their use, extremely

fast colours are obtained upon cotton

The so-called *ingrain dyes* are used almost exclusively on cotton The term is applied to designate a class of dyes which are produced in

situ on the fibre by precipitation

Mercerised cotton is viitually a distinct textile fibre, and whilst its dyeing properties are essentially the same as those of ordinary cotton, its treatment requires special precautions on account of its great affinity for the direct dyes

The dyeing of linen. The behaviour of linen towards the respective classes of dyestuffs is similar to that of cotton, but it is more difficult to dye. The inferior dyeing affinity of linen is probably due to the different physical structure of the fibre

The dyeing of jute. Although a cellulose derivative, jute differs from cotton and linen, both in its chemical composition and properties As regards dyeing processes, it behaves, to a certain extent, like tannin mordanted cotton. and may be dyed not only with direct, but with basic colouring matters without the aid of a mordant

The dyeing of leather. Leather resembles wool in dyeing properties, but the difficulty of dyeing it in fast colours is increased by the necessity of using lukewarm liquors only, since boiling water destroys ordinary leather Certain of the newer methods of producing leather, eq chrome tannage, yield a product which will withstand boiling solutions. It is very important as regards the permanence of the leather (when used eq for bookbinding or furniture covering) that no mineral acid should be employed in the dyeing process. As a substitute, formic acid may be recommended The acid and basic colouring matters are chiefly employed, the mordant dyes being used to a less extent

The dyeing of feathers. The chief cause of difficulty in the dyeing of feathers is their natural water-resisting property. This is due partially to their ultimate physical structure, and partially to the natural grease present The midrib is specially difficult to dye Apart from these special difficulties, feathers behave in dyeing, like wool, and may be dyed with acid, basic, direct, or mordant dyes Reds, browns, blues, and other colours are usually dyed with acid dves, but the most satisfactory blacks are obtained by successive and repeated treatments with logwood extract, and ferrous sulphate with potassium binoxalate Dyeing must proceed for a considerable period at the boil, in order to ensure penetration

The dyeing of horn. Horn is dyed in strong solutions of acid dyes by steeping for a long period in the cold or slightly warm baths, which are acidified with acetic or formic acid it contains sulphur, horn may be dyed brown

by steeping in a solution of sodium plumbite, which results in the formation of lead sulphide Black is also produced by alternate steepings in solutions of mercuric nitrate and potassium sulphide

The dveing of ivory. Both acid and basic dyes may be employed in the dyeing of ivory, the former being used with addition of acetic or formic acid in hot solutions. Alcoholic solutions of colouring matters may also be em-Red is frequently dyed by the cochineal ployed single-bath process used for wool (see p. 500)

Black is produced either by the logwood and iron or aniline black process, as used in cotton

dyeing, or by means of silver nitrate

The dveing of wood. The method of dveing wood in bulk is to boil or steam it to remove the resins, and then treat it in an apparatus similar to that employed in creosoting. The wood is placed in a closed vessel which is exhausted by a vacuum pump, and when the air has been sucked out from the pores as completely as possible, the hot dye liquor is admitted into the vessel and put under pressure by a force The direct and basic dyes may be employed Wood is usually dyed black by alternate treatments with logwood extract and sodium dichiomate solution

The dyeing of straw. Straw is largely dyed for use in the manufacture of hats and baskets There are two commercial varieties, distinguished as chip and straw Chip is readily dyed by means of the direct cotton dies or the sulphide dies, and many basic and acid dyes in neutral solution may also be employed. The process must be continued for 2 to 3 hours at the boil

'Straw' is more difficult to dye than 'chip,' on account of the highly silicious character of the surface The straw should be steeped for several hours in warm water before dyeing Blacks are usually obtained with logwood and iron, but for colours the direct cotton and the basic dyes are chiefly used

MATERIALS USED IN DYEING OPERATIONS.

A great variety of chemical substances are used in dveing processes- acids, alkalis, oxidising and reducing agents, metallic salts, and a wide range of organic bodies Two groups only need be here considered, viz (a) mordants, and b) colouring matters

Mordants. The function of a mordant is to chemically combine with the dyestuff employed. and fix the latter upon the fibre It has already been mentioned that many colouring matters do not require the aid of mordants, but dye direct from their solutions This applies to the acid, direct, basic, and vat dyes when used for wool and silk, and the direct, sulphide, and vat dyes On the other hand, the when used for cotton mordant and acid-mordant dyes (as their name implies) demand the use of mordants on wool, as do the mordant and the basic dyes on cotton

It is evident that the chemical properties of the dyestuff will govern the character of the mordant required for its fixation, and two types of mordants may thus be distinguished: (1) the metallic or basic mordants, used for the mordant' dyes; and (2) the acid mordants, required in applying the basic dyes to cotton or other vegetable fibre.

Basic mordants are employed for fixing on

animal and vegetable fibres those colouring matters which are of an acidic character, ie either true acids or bodies containing hydroxylgroups, of which the hydrogen is replaceable by metals They are metallic salts, of which the most important are certain compounds of chromium, aluminium, iron, and tin

Wool is mordanted by boiling it with dilute solutions of these metallic salts, usually with the addition of certain acids or acid salts, e.g. cream of tartar, sulphune acid, formic acid, oxalic acid, &c. During this operation, the wool assists in the dissociation of the metallic salt; an insoluble basic salt being fixed within and upon the fibre, while a more acid salt remains in solution acids or acid salts, which are added along with the metallic salt, possibly assist the dissociation of the latter by forming more sensitive salts (tartrates, exalates, &c), and are conveniently termed assistants

As a rule, the mordanting operation precedes that of dyeing, not infrequently, however, the two operations take place simultaneously, and the dyeing is thus effected by the so-called 'single-bath' or 'one-dip' method, as, for example, in producing cochineal scarlet on wool In such cases, the wool is boiled in a solution containing colouring matter, metallic salt, and assistant, in certain accurately determined relative proportions Combination takes place between the colouring matter and the mordant, but the pigment which would otherwise be precipitated, is retained in solution by the liberated acid of the mordant or the assistant present, and is only gradually withdrawn from solution by the fibre, which thus becomes dyed

In the case of the 'acid-mordant' dyestuffs, the application of the mordant follows

that of the colouring matter

Silk may be mordanted in the same manner as wool, by boiling with metallic salt solutions, but, more usually, it is steeped for about 12 hours in cold solutions of the mordants previously rendered more or less basic and sensitive subsequently washing with (calcareous) water, the metallic salt absorbed by the silk is dissociated, and an insoluble basic salt is precipitated within the fibre

Cotton is mordanted with metallic salts by processes very different from those adopted for wool and silk, since it has not the property of causing the dissociation of the metallic salts when boiled or steeped in their solutions The methods adopted also vary according to the form of the material, ie whether it be cotton-wool, yarn,

or cloth.

With calico, it is usual to impregnate the fabric with solutions of metallic acetates, then to dry it, and expose for several hours to a moist During this so-called 'agewarm atmosphere ing' process, dissociation of the salt takes place, acetic acid escapes, and a more or less basic salt remains fixed upon the fibre

Sometimes the 'ageing' is replaced by a 'steaming process,' in which the fabric is exposed from 1 to 1 hour to the action of high or low pressure steam in a closed chamber This method is adopted by the printer of textile fabrics in the so-called 'steam style,' in which a mixture of colouring matter and metallic salt (usually acetate) is printed upon the fabric, which is then dried and steamed. The metallic salt is decom-

posed, a basic salt is precipitated upon the fibre, and at the same time the mordant combines with the colouring matter and produces the coloured pigment. This method is analogous to the single-bath process of the woollen dyer

Cotton may also be mordanted by imprognating it with a metallic salt solution, drying, and then passing through a solution of an alkaline substance,  $e\,g\,$  ammonia, chalk, sodium carbonate, &c , to neutralise the acid and precipitate upon the fibre a basic salt or metallic oxide. Solutions of an alkali salt, the acid of which forms an insoluble compound with the base of the mordant,  $e\,g$  sodium phosphate, silicate, arsenate, &c, are also employed. The substances used in this manner for fixing the mordant upon the fibre are termed fixing agents

In some cases of mordanting, the cotton is first impregnated with the fixing agent (eg tannic acid, sulphated oil, &c ), then dried and afterwards passed through the mordant solution,

as for example, in Turkey-red dyeing

Special methods of mordanting cotton are occasionally adopted, in accordance with the properties of the metallic salt employed Stannic oxide, for example, may be fixed on cotton by passing the fabric through a solution of sodium stannate, and then through dilute sulphuric acid, and finally washing Alumina is precipitated on the fibre by the mere exposure to an of cotton impregnated with sodium aluminate, in which case atmospheric carbonic acid is the active fixing agent

The chromium mordants are of the first importance, since, with different colouring matters, they yield a considerable range of colours which are remarkable for their permanence The most important chromium mordants are potassium and sodium dichromates, and they are more largely used as mordants in wool dyeing than all other metallic In cotton dyeing they are not used as mordants, but are frequently employed as oxidising agents, eg to develop the colour in dyeing catechu brown or aniline black Dichromates are also used in producing 'chrome yellow' and 'orange' (chromates of lead) on cotton Chromrum fluoride (CrF<sub>3</sub>,4H<sub>2</sub>O) is also used to a limited extent as a mordant for wool Other chromium salts employed as mordants, chiefly for cotton, are chrome alum, chromium acetate, and various chromates of chromium. Chrome alum is obtained in large quantities as a byproduct during the manufacture of alizarin; chromium acetate is prepared by mixing together solutions of chrome alum and lead acetate

Wool is mordanted with chromium by boiling for  $1-1\frac{1}{2}$  hours in a solution containing 3 p c potassium (or sodium) dichromate calculated on the weight of wool. In many cases it is beneficial to add 1 p c sulphuric acid (sp gr. 184) or 15 pc hydrochloric acid (sp gr 116), or 6 pc tartaric acid, or 1 to 2 pc. formic acid, whereby fuller or brighter colours are ultimately obtained When potassium dichromate alone is employed, partial decomposition of the salt occurs, accompanied by some reduction, the wool absorbing some chromic acid and chromium chromate as well as potassium di-When an addition of sulphuric chromate. acid is made, the wool absorbs only chromic acid. In both cases the mordant d wool

has a yellow or buff colour, and only in the dye-bath does the reduction to the real mordanting body, chromic oxide, take place within the fibre through the interaction of the colouring When formie or tartarie acid is the assistant employed, the mordanted wool has a green colour, reduction of the chromic acid taking place already in the mordanting bath, and chromic hydrate being deposited within the wool substance If large amounts of potassium dichromate are employed in mordanting, especially if sulphuric acid is also added to the bath, bad results may be obtained through the colouring matter being oxidised and destroyed by the excess of chromic acid present in the wool This In certain defect is known as 'over-chroming.' cases, potassium dichromate may be usefully applied in the same bath with the colouring matter (eg with alizarin), or by an after process (eg in the case of camwood)

Silk is not usually mordanted with chromium It becomes mordanted, however, if boiled with a solution of potassium dichromate and tarterie acid after the manner of wool, but not if potassium dichromate alone, or with the addition of sulphune acid, is employed. The fibre is

somewhat injured by this process

Cotton may be mordanted with chromium by impregnating it with a moderately concentrated solution of chrome alum, drying, and passing through a boiling solution of sodium carbonate or caustic soda. Better results are obtained by the use of 'chromium mordant G A I' or 'G A II' (M L & B), which are solutions of chromium chromates. Other satisfactory mordants for cotton are chromium bisulphite (Knecht) and sodium chromite (Koechlin). A method in vogue with the printer is to implicance (pad) or print the cotton with a mixture of colouring matter and chromium acetate, dry and steam ('steam colours')

The chief aluminium mordants employed for wool are aluminium sulphate and alum. The best results are usually obtained by boiling the wool for 1-1½ hours with a solution of 8-10 p c of the aluminium salt calculated on the weight of the wool, with the addition of cream of tartar, in the proportion of 3 mols to 1 mol aluminium sulphate. Aluminium tartiate also gives excellent results, but it is cheaper to allow this salt to be formed in the mordanting bath itself in the manner described. With some colouring matters, exalts acid or potassium binoxalate are

to be preferred as assistants to cream of tartar Silk is mordanted by boiling with dilute aluminum sulphate solution, or by steeping 24 hours in a concentrated solution, and then washing in water, preferably calcarcous

washing in water, preferably calcareous

The methods of mordanting cotton with aluminium are various. It may be impregnated with a moderately concentrated solution of alum or aluminium sulphate, dried, and then passed for a few minutes into a hot solution of sodium phosphate, arsenate, or silicate, or ammonium carbonate. Or the cotton may be impregnated with a cold solution of neutralised sulphated oil, or of tannic acid, then dried, and steeped in a solution of aluminium acetate or basic aluminium sulphate. Another method consists in impregnating the cotton with a solution of sodium aluminate, drying and exposing to the a.r, or passing through a solution of am-

monium chloride. Normal and basic aluminium acetates are also employed, the cotton being impregnated with their solutions, then dried, and exposed to a moist warm atmosphere ('ageing'), and finally passed through a hot solution of, eg, sodium arsenate. The aluminium acetates employed are prepared by dissolving aluminium hydrate in acetic acid, or by decomposing a solution of aluminium sulphate with calcium or lead acetate ('red liquor'). The more basic the aluminium salts are, the more readily do their solutions precipitate by heating or on dilution with water, and the larger the quantity of alumina they yield to the fibre when used as mordants

The chief iron mordants employed are ferrous sulphate and acetate, and ferric sulphate, ace-

tate, and nitrate

The ferrous acetate employed is made by dissolving scrap iron in pyroligneous acid. The 'pyrolignite of non' or 'black iron liquor' thus obtained is preferable to that made by the mutual decomposition of ferrous sulphate and lead acetate solutions, being less liable to oxidise because of the presence of tarry matter and reducing agents, eg pyrocatechol, &c. Ferric sulphate is prepared by heating a mixture of definite proportions of ferrous sulphate, sulphuric acid, and nitric acid. Although the last-mentioned substance is only used as an oxidising agent, the final preparation is generally termed by dyers 'nitrate of iron' The socalled 'pure intrate of iron' (ferric intrate) is prepared by dissolving scrap iron in intricacid

Wool may be mordanted by boiling in a solution of 7 p c ferric sulphate [Fe<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>] and 4 p c potassium binoxalate, calculated on the weight of the wool, but iron is now very little used as a mordant for wool

Silk may be mordanted by steeping for about 12 hours in a cold solution of pyrolignite of non (sp gr 1 025), and afterwards washing well. The most usual mordant, employed largely in black-silk dyeing, is the basic ferric sulphate  ${\rm Fe_4(SO_4)_5(OH)_2}$ . The silk is steeped in a somewhat concentrated solution of this salt (sp gr 1 25) for about 1 hour, then squeezed and well washed. The operations are repeated seven of eight times, after which the silk is boiled in an old soap-bath and finally washed. In this process, the very sensitive basic ferric sulphate absorbed by the fibre is decomposed during the several washings, a still more basic and insoluble salt being precipitated within the fibre.

Cotton is usually mordanted with iron by first impregnating it with a cold solution of tanin matter ( $e\,g$  myrobolans), and then, with or without drying, passing into a dilute solution of ferrous acetate, ferric sulphate, or nitrate (sp.gr  $1\,01-1\,02$ ) The cotton may alternatively be impregnated with ferrous or ferric acetate, dried, and exposed to a moist warm atmosphere ('ageing'), finally fixing the mordant by passing through a hot solution of,  $e\,g$ , sodium phosphate

Tin mordants are now rarely used, but if required the stannous salts are employed for wood and the stannous salts for cotton. The chief stannous mordant is stannous chloride or 'tin crystals' It is frequently sold as a somewhat acid solution under the names of 'single muriate of tin' (sp.gr. 1.3), or 'double muriate of

(sp.gr. 1.6). The general name 'tin spirits' en to a variety of solutions of tin in mixtures adrochloric, sulphuric and nitric acids.

Vool is mordanted by boiling in a solution aning 6 pc of the weight of wool of stanchloride, and 6-10 pc cream of tartar e proportions of stannous chloride should rictly avoided, otherwise the dyed wool feels 1 and its general properties are impaired

ilk may be mordanted in the same manner ool, or steeped several hours in a more is concentrated solution of stannic chloride,

hen well washed

otton is mordanted with stannic oxide by egnating the cotton with cold tannin solution sumach) for 3 hours, then squeezing, and, or usually without drying, steeping 1 in a dilute solution of stannic chloride · 102), and washing Cotton thus mored was much used formerly for fixing ant dyes-peachwood, logwood, &c, in i case the stannic oxide constituted the ant, the tannic acid acting as fixing agent that the basic coal-tar dyes are so much yed in dyeing cotton, although the same od of mordanting is employed, the roles of n salt and tannic acid are reveised, for it lly the tannic acid which is the essential ant, while the stannic chloride serves only · fixing agent

zid mordants are employed for fixing basic ing matters on cotton They are not used of or silk dyeing, since these fibres themact as acid mordants, and are dyed ly without the aid of any external mordant innic acid is the mordant par excellence ring the basic coal-tar colours on cotton ibre absorbs tannic acid from its cold soluand, when thus prepared, if immersed solution of a basic colouring matter, the · acid combines with the colour base, a ed lake is formed within the substance of ore, and the cotton becomes dyed yeing process, the acid originally in comon with the colour base is displaced, and in the free state it interferes with the comformation of the colour lake, it is found cial to add a small proportion of sodium nate or acctate to the dye-bath Since the tes of colour bases are somewhat soluble, in

of tannic acid or in alkaline (or soap) ons, it is necessary in practice to fix the acid on the fibre, previous to dyeing, in the of an insoluble metallic tannate This is by passing the tannic-acid-prepared cotton ha hot or cold solution of tartar emetic or c chloride, tannate of antimony or tin is ived on the fibre, and these salts have as in affinity for the colour base as free tannic as. Cotton thus mordanted and dyed is oly permeated with a very insoluble double e of antimony or tin and colour base; a compound which is insoluble in alkaline ns, and therefore fast to washing with ζC.

tannin mordants, chiefly used by the regall nuts and sumach, for light colours; nyrobolans, divi-divi, quebracho, and it extract, for dark colours and blacks already pointed out, tannic acid is somelised as a fixing agent for tin, iron, or ium mordants, and since its iron com-

pound is of a bluish-black colour, it sometimes serves as a useful colouring matter.

Tannic acid is not used as a mordant for wool, but for silk it is largely employed, partly for adding weight to the silk, and partly for the purpose of dyeing it black in conjunction with iron mordants. Its utility as a weighting agent is owing to the fact that under suitable conditions silk absorbs as much as 25 p c of its own weight of tannic acid from a hot solution, whereby the fibre gains not only in weight, but also in volume

Fatty acids are used as mordants, very much in the same way as tannic acid, for fixing the basic coal-tar colours on cotton. They are, however, not attracted by cotton from their solutions, but they form insoluble lakes with the colour bases. The method of procedure is to impregnate the cotton with a more or less concentrated solution of their alkali salts, viz soap, after drying, the fabric is passed through, or preferably steeped in, a cold solution of aluminium sulphate, and washed. Although the fatty acid is thus fixed on the fibre as an insoluble aluminium soap, it is still capable of combining with the colour base when the cotton is passed into a solution of a basic colouring matter Colours dyed by this method are brighter than, but not so fast to washing as those fixed by means of tannic acid.

Instead of ordinary soap, a more convenient fatty-acid mordant to employ is the so-called 'sulphated oil' This is prepared by mixing two parts by weight of castor oil with about one part of sulphuric acid (sp gr 184), and allowing the mixture to stand for 24 hours is then washed with a strong solution of common salt to free it from excess of acid, and is finally neutralised with ammonia or caustic soda product of the action of the acid upon the oil is somewhat complex, but its practical utility depends upon the fact that it behaves like a fafty acid and gives an extremely soluble soap with alkalis Ordinary easter-oil soap produced by boiling the oil with caustic soda is also very soluble, and may be used as a mordant Sulphated oil is frequently called 'Turkey-red oil,' 'alizarin oil,' and 'soluble oil,' since it is largely used in the modern short process of Turkey-red dyeing as the fixing agent for the necessary aluminium mordant, and it gives the characteristic brilliancy of colour obtained by the use of the oil preparations formerly employed Monopole oil is a similar product

Phenohe colouring principles, eg alizarin, and certain acid colouring matters derived from benzidene, stilbene, fluorene, &c (the direct cotton dyes) can also be fixed on cotton, and serve as mordants for basic colouring matters, but they are only occasionally used for this purpose for the production of compound shades

COLOURING MATTERS

(For Classification, see p 575)

GENERAL METHODS OF APPLICATION

CLASS I — Mordant dyes. This group of dyestuffs includes the majority of the natural dyes, also the namerous class of alizarin dyes, a number of derivatives of pyrogallol and other phenols, and certain azoquinone-oximes and nitroso-dyes. The following are the chief members of the group

Mordant dyes of natural origin—Logwood; limawood and other camwood and other cochineal; soluble redwoods' 'insoluble redwoods'; fustic; weld; querci-

tron bark and flavin.

Synthetic mordant dyes -Alizarin; anthrapurpurin, flavopurpurin, and purpurin; nitroalizarin (alizarin orange), alizarin maroon (aminoalizarin), alizarin bordeaux, anthracene brown, alizarin blues, alizarin cyanins, anthracene blues; alizarin greens, corulein, anthracene yellow; mordant yellow, galloflavin, &c; alizarın black; gallem; gallocyanın, cœlestine blue, naphthol green, dark green, gambine, &c

The mordant dyes are largely used in the dyeing of wool and in the dyeing and printing of cotton and silk Generally speaking, they yield colours which are of great fastness to light, soaping, acids, alkalis, bleaching agents, &c As regards cotton dyeing, they have largely been replaced by 'after-treated' direct dyes and the very fast vat and sulphide dyes recently intro-

duced

When a fabric, previously mordanted with a metallic salt, is dyed with a mordant dye, a chemical reaction takes place, resulting in the formation of a new compound which is the actual colouring substance, this compound being, at the same time, fixed upon the fibre, which thus becomes dyed. These colour lakes or pigments may, for the most part, be equally well produced by bringing together the mordant and the dyestuff under suitable conditions in the absence of the fibre Many colour lakes of this class are indeed manufactured in bulk for use in printing or painting.

There are three methods by which mordant dyes may be applied. (1) the mordant may be first applied to the fibre, and afterwards the dyestuff in a separate operation ('mordanting and dyeing method'), (2) the order of applica-tion of mordant and dye may be reversed ('dyeing and saddening method'), or (3) the mordant and dyestuff may be applied simultaneously ('single-bath method') All three methods are in use If equally good results can bo obtained, the 'single-bath' process is to be preferred, but usually the 'mordanting and

dyeing' process gives the maximum fastness

The dying of wool with mordant dies —

When potassium dichromate is comployed as mordant, the 'mordanting and dyeing' process is usually preferable, because, in the 'single-bath' process, the acid necessarily added to dissolve the colour lake formed in the dye-bath, converts the mordant into such a powerful oxidising agent by liberating chromic acid, that most dyestuffs are injuriously affected. In the two-bath process, the wool should be thoroughly washed after mordanting If this operation is omitted, a portion of the mordant passes from the wool into the dye-bath, and, combining with an equivalent amount of colouring matter, forms a precipitate which floats about in an insoluble condition. This not only leads to loss of dyestuff, but also tends, by superficial absorption, to cause the dye to 'rub off' The dyeing process consists in boiling the wool for from 1 to 2 hours with the colouring matter in aqueous The final washing, after dyeing, has for its object the removal of any colouring

matter which is imperfectly fixed; but, however thorough the washing may be, there is usually a small amount of colouring matter present on the dyed wool, which has merely been absorbed without combining with the mordant. This free colouring matter will dissolve out or 'bleed' during milling or scouring, because most mordant dves are readily soluble in alkaline solutions A slight supplementary mordanting ('afterchroming') is frequently used to fix this small amount of free dyestuff

The 'dyeing and saddening' method is not much used with the true mordant dyestuffs, but is the chief process used in applying the acid mor-Since the ultimate intensity of colour depends chiefly upon the amount of dyestuff which is fixed in the first bath, whilst the colour is only developed by the mordant in the second bath, it is obviously more difficult to exactly match a pattern when this process is used than

when the mordant is applied first

In the 'single-bath' method of dyeing, the bath is prepared with the necessary amount of mordant and dyestuff with an addition of sufficient acid (usually oxalic or acetic acid) to dissolve the colour lake formed. This process is therefore only suitable when the colour lake is soluble in dilute acid, and this is particularly characteristic of the tin and aluminium colour lakes, although the single-bath process is also frequently applicable in the case of ferrous sulphate or chromium fluoride mordants

The dyeing of silk with mordant dyes -The mordant dyes are not very largely used on silk, chiefly because they detract somewhat from the lustre of the fibre The degree of fastness of the acid dyes, if suitably selected, is, moreover, usually sufficient for the requirements of the silk dyer. The methods of mordanting have already been dealt with. Dyeing must take place at the boil for about an hour, and the silk is subsequently brightened by working in a warm dilute solution of acetic or tartaric acid,

and dried without washing

Dueing of cotton with mordant dies -The 'mordanting and dyeing' method is generally used in cotton dyeing Aluminium is the chief mordant employed, and, in conjunction with alizarin, it yields the important Turkey-ied dye This is a complex compound containing an alizarin-aluminium-calcium lake in combination with polymerised fat acids In the production of this colour, the cotton is first prepared with sulphated oil, then dued, mordanted with alum, and 'subsequently dyed with alizain complete process involves a large number of operations, for the details of which manuals of dyeing must be consulted. The dyeing of Turkey red is a very ancient process, having been carried out centuries ago in India by the aid of milk (as fatty matter) and munjeet, the Indian madder plant No alum was employed. but the plant with its earthy incrustations contained sufficient alumina to produce the alumina The art spread from the East through Persia and Turkey, and was first introduced into this country (Glasgow) in 1790 Madder has now been entirely replaced by synthetic alizarin in Turkey-red dyeing, and the substitution of 'sulphated oil' for olive oil or soap has also greatly simplified the process.

Another important mordant dye on cotton

is logwood black, which is obtained by dyeing with logwood on tannin-iron mordant. It is largely used in hank and warp dyeing. Catechu brown is produced by the 'dyeing and saddening' method, the cotton being first steeped in a strong solution of catechu, and subsequently treated with potassium dichromate. The latter acts essentially as an oxidising agent, producing the so-called paponic acid on the fibre

A great variety of permanent and bright colours may be obtained with mordant dyes on cotton mordanted with chromium or aluminium, but they are not very largely used, fast colours being chiefly produced by means of the sulphide, but, and margin dyes (Classes VI, VIII, VIII)

vat, and ingrain dyes (Classes VI, VII, VIII)

CLASS II—Acid-mordant dyes. These dyes are suitable only for wool and silk, and are largely used on wool They are now very numerous, and are chiefly azo-compounds containing salicylic acid or orthoaminophenol components. Some, however, are anthracene or pyrogallol derivatives. Unlike the mordant dyes, they usually contain sulphonic acid groups

Many of the acid-mordant dyes may be employed as ordinary acid dyes, since a mordant is not essential to the development of their colour. Usually, however, such dyestuffs produce faster colours when applied in conjunction

with a mordant

An exhaustive list of acid-mordant dyestuffs cannot be given, but the following will serve as They include dyes giving a complete examples acıd-alızarın, anthrarange of colours Redscene, chrome, cloth and milling reds and yellows cloth oranges and yellows, alizarin yellows (certain marks are mordant dyes), chrome yellows, milling orange and yellow, diamond, and mordant yellow Greens alizarin, chrome, diamond and milling greens Blues . acid-alizarin, brilliant-alizarin chrome blues, chromotropes (chromotropes dye red as acid dyes). Violets alizarin heliotropes, anthraquinone and chrome violets, fast violet Blacks and greys: acid-alizarin, acid-chrome, anthracene-acid, anthracene chrome, chrome, chromotrope and diamond blacks acid-alizarin, acid-anthracene, chrome, cloth, diamond, and palatine browns; chromogen

Dyeing of wool with acid-mordant dyestuffs The usual method of application of this class of dyestuffs is to prepare a bath containing the necessary amount of dyestuff along with 1 to 3 p.c sulphuric or 3 to 5 p.c acetic acid, and 5 to 10 pc sodium sulphate (Glauber's salt) (all calculated on the weight of wool) the solution has been brought gradually to the boiling-point, the boiling is continued until the colouring matter is entirely fixed, an additional amount of acid being added if necessary of acid, however, frequently retains the dyestuff The wool is then removed from in solution the bath, the metallic salt is added, and the wool is again boiled in the bath for half an hour. In most cases, petassium dichromate, to the extent of one-half the weight of the dyestuff, is the mordant employed

A modification of this process is used with the metachrome series of dyestuffs The metachrome mordant is a mixture of yellow potassium chromate and ammonium sulphate. The bath is prepared with 3 to 5 p c of this mordant together with the dyestuff. A prolonged boiling is necessary, during which the ammonium salt is decomposed, ammonia being evolved and the liberated sulphuric acid gradually converts the chromate into dichromate and as ists in the fixation of the dye and the formation of the colour lake.

The anthracene chromate dyes are applied in a bath containing colouring matter, dichromate, and sodium sulphate, this being another modification of the usual process.

The mercerol and autochrome dyes are also

applied by special processes

The acid-mordant dyes are of increasing importance in wool dyeing, but are little used on

silk, and not at all on cotton

CLASS III -Acid dyestuffs. The acid dyes constitute the most numerous class of synthetic The chief chemical groups colouring matters of acid dyes are azo- dyes, nitro- compounds, sulphonated basic dies, and phthaleins, greatest number being azo-dyes The azo-dyes contain few greens, blues, and violets, but the reds, oranges, yellows, browns, and blacks are very numerous The nitro- compounds are few in number and are all yellows, the chief being The greater pieric acid and naphthol yellow number of the sulphonated basic dyes, on the other hand, are greens, blues, and violets, but acid magenta belongs to this group. The resorcin or phthalein dyes are chiefly brilliant pinks or reds, the cosins being the most important group

So extremely numerous are the acid dyes, and their names are so varied, that no useful purpose would be served by the enumeration of

a small number of the products

The dyeing of wool with acid dyes—The commercial acid dyes are, in most cases, the sodium salts of sulphonic acids The introcompounds, however, owe their acid character to the presence of intro-groups in the molecule

The simplest and most usual method of dyeing acid colours on wool is to prepare a bath containing the requisite amount of dyestuff with 1 to 4 p c of sulphuric acid and 10 to 20 p c of sodium sulphate (Glauber's salt) wool is dyed in this solution for about an hour, boiling for the greater part of the time dyes require smaller, and some larger percentages The sodium sulphate has a levelling influence on the fixation of the dyestuff, probably by reason of the solvent action which it exerts, since acid dyes are readily removed ('stripped') from wool by boiling in a solution of sodium sulphate Another reason is to be found in the retarding action which it exerts in the liberation of the free colour acid by the sulphuric acid The equation, which may be expressed as

Colour salt + H<sub>2</sub>SO<sub>4</sub> ≥ colour acid + Na<sub>2</sub>SO<sub>4</sub>, is reversible (See also M Firt, Theory of the Acid Dyebath, J Soc Dyers, 1910, p. 33.)

Sodium bisulphate or 'nitre cake' may be used in place of the sulphuric acid and sodium sulphate, or the sulphuric acid may be entirely replaced by acetic or formic acid.

In the case of dyestuffs which do not readily produce level colours on the fabric, the acid may be added to the bath gradually in small portions at a time, or may be substituted by ar. ammonium salt, eg ammonium sulphate

The acid is then liberated very gradually as the ammonium salt is decomposed in the boiling bath, and the colouring matter is therefore converted very slowly from the condition of the soluble sodium salt to that of the more or less insoluble free sulphonic acid

The dyeing of silk with acid dyes —Acid dyes are very largely used in silk dyeing. They are applied in a bath containing a large quantity (up to 25 p.c. in volume) of 'boiled-off' liquor, the soapy solution of the silk gum obtained in the 'ungumming' of raw silk. The bath is rendered slightly and by means of sulphure, acetic, or formic and Dyeing usually takes place at about 80°, at which temperature silk has more affinity for acid dyes than in a boiling bath

After dyeing, the silk is washed and is then 'brightened' by working in a dilute (sulphuric, acetic, or tartaric) acid solution and dried

without washing

The dyeing of cotton with acid dies -There is no satisfactory method of fixing the acid dyes upon vegetable fibres, but for cheap work they are often applied by a 'padding' process
(LASS IV —The direct dyes. The direct dyes

have the special property of dyeing cotton without the aid of a mordant. Lake the acid dyes, they are for the most part azo- compounds, but contain the chromophor -N=N- twice in the molecule, and thus belong to the tetrazogroup of azo- dyes They further resemble the acid dyes in that they are alkali salts of sulphonic acids, but as regards dyeing properties, they differ greatly from the simpler azo-acid dyes Since the introduction, in 1884, of the first of the direct dyes, a very large number have been placed on the market, and the group contains dyes of all colours-reds, oranges, yellows, greens, blues, violets, browns, and blacks. As examples of these dyes may be mentioned the congo, benzo, diamine, mikado, titan, and chlorazol dyes

Primuline is an example of a direct dye which is not an azo- compound. It contains the

chromophor<S>C-

Turmeric is the only important natural direct dye for cotton

The direct dyes are largely used upon all No satisfactory explanation has been given of the fact that they readily dye cotton, whereas the acid dyes are entirely devoid of this property The further fact that the direct dyes dye wool in a neutral, or even slightly alkaline bath, whereas the acid dyes require the addition of acid, also requires explanation.

The dyeing of wool with direct dyes -I hese colouring matters are applied to wool in a neutral or feebly acid bath with or without the addition of sodium sulphate (Glauber's salt). In some cases, an after-treatment with a metallic salt gives improved fastness, the dyes then practically falling into the class of acid-mordant dyes.

As a class, the direct dyes on wool are much faster to milling than the ordinary acid dyes When fixed on wool, they are also faster to acids, washing, and light than when dyed on cotton In some cases, the colour produced by a direct dye on wool and cotton is identical, in other cases it differs considerably

The dyeing of silk with direct dyes -These

and also on account of their property of pe manent reduction to colourless bodies, in the s called 'foulard' style of printing. They a applied in a similar manner to the acid dybut smaller amounts of 'boiled-off liquor' a acid are required

The dying of cotton with direct dyes .- T application of the direct dyes to cotton is ve They are dyed either from a cold, lul warm, or boiling solution, to which is added small amount (up to 2 p c ) of sodium carbona and an amount of a neutral salt (up to 20 gra NaCl or Na SO<sub>4</sub> per litre), which is governed the solubility of the dyestuff employed At l temperatures more concentrated solutions colouring matter are required. Brighter color are obtained by passing the dyed mater through a solution of Turkey-red oil before d ing or by adding this substance to the dye-ba Mercerised cotton has a much grea affinity than ordinary cotton for the direct dy and in dyeing this fibre special precautions.

Many of the direct dyes produce on cott colours of great brilliance and richness, I they bleed out into hot water, and are s more readily attacked by soap. Most of th are also comparatively fugitive to light.

There are three processes by means of wh the fastness-properties of certain of the dir

dyes can be improved

1 The diazotising and developing process This process is applicable only in the case those dyestuffs in which amino- groups c stitute the auxochrome. The cotton is f dyed by the general process above describ then washed, and afterwards worked for a moments in a cold dilute solution of nitrous a (2 p.c. NaNO, and 4 p.c. HCl cn the wei of the cotton) This treatment diazotises The cotton is then well was amino- groups in cold water and entered immediately into developing bath, which contains a solution of phenolic or amino- compound; sufficient al or acid respectively being added to bring 'developer' into solution. The chief develop used are phenol, resorcinol, a- or \(\beta\)-napht aminonaphthol, aminodiphenvlamine, m-to enediamine, and certain of the napht sulphonic acids

Primuline is applied by the process described

A modification of this method is known the 'coupling process'. The cotton is first d with the direct dye, and, after rinsing, is wor in a diazotised solution of the 'coupler' chief body used as coupler is p-nitranil Ready prepared stable diazo- compounds of base are sold under the names 'nıtra: azophor red,' and 'nıtrosamine red'

2. After-treatment with metallic salts this process for improving the fastness of direct dyes on cotton, the fibre is first dyes the ordinary way, and then, after rinsing treated for half an hour in a warm slightly ac fied solution of a suitable metallic salt chiefly employed are copper sulphate, potass dichromate, and chromium quoride; from 1 pc with addition of 2 to 4 pc. of acetic: being used. A mixture of the two first-n tioned salts is frequently used. This process dyes are largely used on silk both for self shad a mainly employed in the case of blues, gre

browns, and blacks. The chromium salts exert the greatest influence in increasing the fastness to washing, and the copper salt in regard to the fastness to light

This process is only applicable to the case of dyestuffs containing free OH or COOH groups

3 After-treatment with formaldehyde -This process, which is chiefly used for navy blues and blacks, is carried out by working the dyed cotton for half an hour at 70° in a solution containing about 3 p c formaldehyde (on the weight of the The fastness against washing is concotton) siderably improved thereby.

Cotton dyed with direct dyes may be 'topped' with basic dyes, for which the direct

dyes act as mordants

CLASS V—The basic dyes. This group of colouring matters is numerically smaller than the mordant, acid, or direct dye groups includes, however, the most brilliant dyes known, rhodamine pink, auramine yellow, malachite green, methylene blue, and methyl violet being well-known examples.

The basic dyes are very diverse in their chemical constitution Many are derivatives of di- or tri-phenylmethane (magenta, malachite green, methyl violet), others are azo- compounds (chrysoidine, Bismarck brown), some are azinederivatives (safranine and induline), whilst the thia/ines contain sulphur as an essential constituent (methylene blue)

The commercial basic dyes are usually hydrochlorides of the colour bases, but some are acetates, oxalates, sulphates, or zinc chloride double salts. They are characterised by great tinetorial power, 1 p c being usually sufficient to produce a full rich colour As a class, they are not fast to light

Janus dyes. These occupy an intermediate position between the basic and the direct dyes. They are azo- dyes of strongly basic character, and possess an extraordinary affinity for both animal and vegetable fibres. They are only of

slight importance

The dyeing of wool with basic dyes -These dyes are little used on wool, the acid dyes being applied in an equally simple manner and producing faster colours Wool has a strong affinity for basic dyes, and readily becomes dyed if A small amount immersed in their solutions of sulphuric or acetic acid may usually be added with advantage Certain of the basic dyes, e g Victoria blue, are very fast to milling when dyed on wool, and such are still employed

The dyeing of silk with basic dyes—Members of this group of colouring matters are largely used by the silk dyer when fastness to light is not important. The silk yarn is dyed at 60°-80° in a slightly acidified solution of the dyestuff, to which is added some 'boiled-off liquor' After rinsing, the dyed silk is brightened in a dilute solution of acetic or tartaile acid

The dyeing of cotton with basic dyes.—Cotton has little affinity for basic dyes, which, on this fibre, must therefore be applied in conjunction with a mordant, and since the dyestuffs of this class are basic in character, the mordant must be of an acidic nature

Three types of acid mordants are available, tannin matters, fat acids, and colour acids

Tannın morde ils -As mentioned dealing with the mordants, cotton has a natu al

affinity for tannic acid, which it withdraws from solution Since tannic acid forms lakes with the basic colouring matters, cotton thus prepared may be dyed with basic dyes, but the colours produced in this manner are not fast against As previously explained, however, if after mordanting with tannin the cotton is treated with a salt of antimony, tin, or iron, an insoluble metallic tannate is formed on the fibre, which compound has the same affinity for basic dyes as tannic acid has The fastness properties of the double tannate of colour base and metal are much superior to those of the simpler tannates of the colour bases

For the production of pale or bright colours, commercial tannic acid, gall nuts, or sumac are the tannins employed. The cotton is saturated ('padded') with a solution of 2° to 3°Tw strength, or steeped for 2 to 3 hours in a solution containing 2 to 5 p c tannic acid or 10 to 30 p c ground gall nuts or sumac, the amount being calculated on the weight of cotton. The varn or fabric after squeezing is then at once passed into a 'fixing bath' containing I to 25 pe of tartar emetic or an equivalent amount of antimony oxalate or fluoride After well washing, the cotton is dyed with 0.1 to 1 p.c. of basic dye

Additions of acetic acid to the tannin bath, ammonium chloride and chalk to the fixing bath, and a little acid to the dye-bath, are frequently The mordanting and fixing baths may be used for several successive lots of material

If the brightest possible shades are not required, some cheaper tannin matter (myrobolans or quebracho) may be employed, and stannic chloride may replace tartar emetic as fixing agent For dark shades, ferrous acetate ('black iron liquor') is used as fixing agent, the blueblack tannate of non producing a composite colour with the basic dyestuff

2 Fatty mordants -If Turkey-red oil or soap is used as mordant, the colour produced with a given basic colour is brighter, but less fast, than when tannin mordant is used

The fat acids are fixed on the cotton as metallic salts, usually aluminium salts; the fibre being first saturated with a solution containing I lb of sulphated oil or soap per litre, then died and passed into a solution of aluminium acctate ('red liquor') of about 8°Tw.

3 Colour acids as mordants - Mention has already been made of the fact that the direct dyes act as mordants towards the basic dyes, compound shades being thus obtained Mordant, sulphide, and acid dyes also possess the property of attracting basic dyes, but these combinations

are not much employed.

The dyeing of jute with basic dyes -Although jute is a vegetable fibre, it does not consist of free cellulose, but of a bastose-cellulose compound, corchorobastose (Cross and Bevan) The dyeing properties of this substance differ materially from those of cellulose, jute, in fact, behaving in a very similar manner to tanninmordanted cotton, and dyeing direct with basic

CLASS VI -The sulphide dyes. The dyes of this group are numerous and important are chi fly employed for the production of fast blacks, blues, greens, browns, and yellows on cotton and other vegetable fibres. They come into the market in the form of powders, which

are usually insoluble in water, but soluble in solutions of sodium sulphide. They are also rendered soluble by other reducing agents, e g. sodium hydrosulphite or glucose in alkaline solution

The sulphide dyes are commercially known under such names as. sulphur, sulphurol, thiogene, thionol, thional, thioxine, cross-dye immedial, katigen, kryogen, pyrogen, pyrol rexoll, eclipse, vidal, or auronal dyes. They are produced by melting together under suitable conditions aromatic nitro-, amino-, or iminocompounds with sulphur and sodium sulphide. The constitution of the sulphide dyes is not known, but the commercial products are probably mixtures of polymers or homologues and not homogeneous bodies. They contain free sulphur in amount varying from I to as much as 12 p.c (Gardner and Hodgson, J Soc Chem Ind 1910, 672), and also contain very variable amounts of sodium sulphide and polysulphides

The dyeing of wool with sulphide dyes -Wool is rarely dyed with sulphide dyes, on account of the injurious action of sodium sulphide on the The wool may be protected to some extent against this action by the addition to the dye-bath of glucose, tannic acid, or ammonium salts. A previous treatment of the fibre with formaldehyde has also been recommended (R Kann, Eng Pat 3492, 1903), but this reduces the affinity of the wool for these dyestuffs

Greater success in the application of sulphide dyes to wool or union material has been attained by E Lodge (Eng Pats 19852, 1913, and 23386, 1914) by the use of sodium hydrosulphite and sodium sulphite as reducing agents, and ammonia as solvent for the reduced dyestuff (J. Soc. Dyers, 1915, 252)

The dyeing of silk with sulphide dyes -Owing to the harmful effect of their alkaline solutions upon the silk fibre, the sulphide dyes are not used

The dyerng of cotton with sulphide dyes -This group of dyestuffs is now of the greatest importance in cotton dyeing for the production of fast blacks, and to a less extent, fast blues, greens, browns, &c The necessary ingredients in the dye-bath, in addition to the dyestuff, are sodium sulphide, sodium sulphate, or other neutral salt, and an alkalı, usually sodium car-The actual and relative amounts of these substances vary considerably with different dyestuffs, depending upon the ease with which the colouring matter employed undergoes reduction and the solubility of the reduced product An addition of Turkey-red oil, monopole oil, or glue, tends to produce more level and brighter shades, glucose aids the reduction Dyeing takes place just below the boil, and it is usually desirable to keep the cotton completely immersed during the operation, in order to prevent irregular oxidation. The baths are used in a very concentrated condition, with the result that only a small proportion of the dyestuff is withdrawn, and for reasons of economy 'standing baths' are employed, that is, the baths are used continuously and replenished with dyestuff, &c. as required. The dyeing takes place in iron, wooden, or lead-lined wooden vessels, copper being nigidly excluded, on account of the ease with which it is attacked by sodium sulphide

Some of the sulphide dyes do not require any treatment subsequent to dyeing other than exposure to air, but with many dyes of the group an after-treatment greatly improves their fastness to light and other influences. object of the after-treatment is to ensure the complete oxidation of the colouring matter, and, as oxidising agents, chromium salts, or hydrogen peroxide are used Steaming the material in presence of air is employed in the case of immedial blue

After-treatment with potassium dichromate and acetic acid is now usually practised, since the use of copper salts has proved to be dangerous in regard to the subsequent tendering of the fibre on account of their catalytic action in inducing acid formation (Gardner and Hodgson,

J Soc Chem Ind 1910, 676)

The tendering of cotton dyed with sulphide blacks has greatly retarded the use of these dye-The defect does not usually develop for some weeks or months after the material has been dyed, and it has been shown to be due to the slow formation of sulphuric acid on the The development of acid arises mainly from the oxidation of free sulphur present on the dyed fibre, but even in the absence of free sulphur, acid development may take place by decomposition and oxidation of the colour molecule.

The risk of tendering, during storage, of cotton dyed with sulphide blacks is entirely eliminated by a final treatment of the material, before drying, with sodium acetate, but, of course, the protective influence of this salt disappears if the material is washed

CLASS VII—The vat dyes. Until comparatively recently, indigo was the sole representative of this group of dyestuffs synthetic production of indigo was, however, followed by that of a number of derivatives of this body Other series of vat dyes have also been produced, and by their use it is now possible to produce on cotton, and in a more limited degree on wool, many bright colours of a degree of fastness hitherto unattainable.

The essential features of a vat dye are its easy reduction to a form soluble in dilute alkali, its attraction in this condition by the fibre, and its ready reoxidation on the fibre to the original insoluble condition. The reduced solution is termed the 'vat,' the composition of the vats differing for wool and for cotton

The vat dyes are mainly derivatives of indigotin or anthracene They include the following . indigo (indigotin), mono-, di-, tri-, and tetra-halogen derivatives of indigotin, methyl and tolyl indigotins, ciba dyes (brom indigotins), thioindigo red and scarlet, helindone dyes (sulphur derivatives), algole dyes, cibanone dyes, and the indanthrene dyes The group includes dyes of all hues

Indigo is the oldest and by far the most important of the vat dyes It has been employed in India for at least 3500 years, by methods not differing essentially from some of those which are still employed. It is used upon all fibres,

but least commonly upon silk.

Various reducing agents are employed for the conversion of the insoluble indigo blue into indigo white, and they are always associated with an alkali to dissolve the latter substance. After saturating the fibre with the alkaline sclution, the indigo white is reconverted into

indigo blue upon the fibre by atmospheric oxidation

The dyeing of wool with indigo vat blue The two chief indigo vats used for wool are the fermentation vat and the hydrosulphite vat, the former being chiefly used for dark shades on woollen cloth (Army and Navy cloth, &c ), and the latter for light colours on slubbing, varn, &c There are several forms of fermentation vat, of which the woad vat is the most important in The precise composition and this country 'setting' of the vat is largely a matter of experience, the ingredients being somewhat as follows. finely ground indigo, 12 to 18 lbs woad, 300 lbs; bran, 10 lbs; madder, 12 to 18 lbs, lime, 12 lbs, water, 600 gallons After warming the liquor, various fermentations set in, the essential bacterium being the Desmobacterium hydrogeniferum The starchy matters present are first converted into glucose, which then splits up into lactic acid, the latter undergoing butyric fermentation with evolution of hydrogen.

Excessive fermentation is prevented by making suitable additions of lime, or it is accelerated, if necessary, by adding further quantities

of bran

In the so-called German or soda vat and the Indian or potash vat, the woad is replaced by bran with the addition of molasses, and the lime

by sodium or potassium carbonate

To avoid the derangements to which all fermentation vats are liable, Collin and Benoist (J Soc Chem Ind 1885, 493) proposed to employ a pure ferment and a completely fermentable food material, but this process has not come into practical use.

Natural indigo is still largely used in the fermentation vats, but in the hydrosulphite vat the synthetic product is generally employed The hydrosulphite vat may be prepared as lbs, caustic soda solution 42°Tw, 3 pints, sodium hydrosulphite solution 20°Tw, 7½ gallons, water, 50 gallons. The liquid is heated to 50° until it becomes clear greenish-yellow solution of indigo white thus obtained is diluted to 250 gallons with water at 50°, to which a little hydrosulphite has been previously added.

The hydrosulphite solution may be replaced by 3 lbs of solid sodium hydrosulphite, in which case the amount of caustic soda must be in-3½ lbs of natural indigo (60 pc indigotin) are about equivalent to 10 lbs synthetic indigo (20 p c paste) After working in the vat, the goods are well squeezed, and then exposed to the atmosphere to oxidise the indigo

white they are then well washed

Dark shades of indigo blue on wool are usually 'bottomed' with a red dye (alizarin, diamine fast red, or barwood) before vatting, and are frequently 'topped' after vatting with orchil or other dyestuff in order to add ' bloom ' to the colour.

In most respects indigo vat blue is an extremely fast colour, but has the defect of being liable to 'rub off'

The dycing of silk with vat indigo —This dye If used, the is not much employed on silk most suitable vat to employ is the hydrosulphite vat set with synthet c indigo.

The dyeing of cotton with vat indigo - . 'he

hydrosulphite vat is largely employed in cotton dyeing as well as in the dyeing of wool, but it is used cold in the case of cotton. Other cotton vats are the 'copperas vat' and the 'zincpowder 'vat.

A strong copperas vat contains, indigo (60 pc), 10 lbs ferrous sulphate, 30 lbs, and quicklime 35 lbs, in about 200 gallons of water The lime decomposes the ferrous sulphate, with the production of ferrous hydroxide, and this in the presence of lime and indigo decomposes water, yielding ferric hydroxide and hydrogen; the latter at once combines with the indigotin and produces indigo white, which dissolves by reason of the excess of lime present The normal colour of the vat liquor thus obtained is brownish-amber

The zinc-powder vat is made up with water, indigo, zinc-powder, and lime In the presence of the lime and indigo, the zinc decomposes water even at the ordinary temperature, and the liberated hydrogen effects the reduction of

the indigo

Vat dyes other than indigo -These may be divided into two groups (a) the indigoids, which may be considered as derivatives of indigotin, and (b) the anthracene derivatives all possess the common property of forming vats when treated with hydrosulphite of soda, but whereas the indigoids may be applied to the animal as well as to the vegetable fibres, the anthracene derivatives are only suitable for cotton on account of the large amount of caustic alkali required for their solution reduced anthracene vat dyes also do not appear to possess much affinity for wool

(a) Indigoids—Examples of dyes of this series are throundigo red and scarlet, helindone red and scarlet, ciba red and scarlet, vat red, helindone orange and yellow, ciba green; brom indigos, ciba blues and greys; ciba violets and heliotropes, helindone brown Some of the algole reds also probably belong to this group

(b) Anthracene vat dyes —Indanthrene red, algole reds, indanthrene yellows and oranges, algole yellows and oranges, cibanone yellows and oranges, indanthrene greens, algole greens; indanthrene, chloranthrene, duranthrene and caledon blues, algole blues, indanthrene grey, indanthrene blowns, algole browns, cibanone

The application of the indigoid vat dies to wool and silk —Vats suitable for the dyeing of wool and silk are obtained by reduction with sodium hydrosulphite in alkaline (caustic soda) Bright reds of excellent fastness are solution obtained with these dyes, and they are used in conjunction with indigo for producing bright purplish blues Separate vats are preferable for the indigo blue and the red dye, since the best conditions of dyeing do not coincide in the two

The application of indigoid vat dyes to cotton -Vegetable fibres are readily dyed with these dyes from a hydrosulphite vat, or alternatively they may be dyed from a sodium sulphide bath, as used for the sulphide dvestuffs. They may as used for the sulphide dyestuffs be mixed with most sulphide dyes and dyed from the same vat.

The application of the anthracene vat dyes -This group of dyes is also applied in the hydrosulphite vat, but usually requires such a strongly

caustic solution that its application to wool and silk is not practicable

By use of the vat dyes, it is now possible to produce a great variety of colours on cotton which are extremely fast to light, washing, bleaching, and other influences. The high cost of these dyes at present somewhat limits their application

Indophenol is a vat due of different type, being obtained by the action of nitroso-dimethyl amline on a-naphthol. On reduction, it is converted into indophenol white, a body sufficiently stable to be sold commercially. On oxidation with chromic acid, indophenol white is reconverted into indophenol blue

Indophenol has been employed in conjunction with, or as a substitute for indigo, but has the defect of being readily affected by acids.

Tyrian purple—As the result of an extremely interesting investigation, P Friedlander has shown (Ber. 1909, 765) that this ancient dyestuff is identical with one of the dibromingotins Tyrian purple, as described by Pliny, was obtained from various species of the mollusc mirrer, and from 12,000 of the species M brandaris Frieldander obtained 14 grams of the colouring matter

Under the law of the ancient Romans the use of Tyrian purple was confined to royalty, and the method of its production was a lost art for many centuries. At the beginning of the 18th century, it was investigated by Reaumur (Mem de l'Acad Royale des Sciences, 1711), but it never again assumed any importance as a dive stuff.

dye stuff
CLASS VIII—The ingrain dyes. The term
ingrain dye may be used to designate those
colouring matters other than mordant dyes
which are produced from their constituent
bodies upon the fibre. Three sub-groups may be
distinguished (a) aniline black, (b) the insoluble

azo- dyes, and (c) the metallic dyes.

(a) Anthre black—This term is applied to the insoluble black pigment which is produced by the oxidation of aniline in acid solution. In 1834 Runge noticed that a blue-black compound was produced by oxidising aniline with chromic acid, but Wilm, in 1860, first described the formation of the product on the cotton fibre. The first successful process for producing aniline black on the fibre was patented by Lightfoot, in 1863. He used potassium chlorate and cupric chloride as oxidising agents, and the black proved to be of such excellent fastness that innumerable modifications of the process have since been proposed, and the literature of the subject is very extensive.

Numerous oxidising agents have been employed in the production of aniline black, eq. chlorates, chromates, salts of copper, vanadium, tungsten, cerium, &c. Aniline black may also be produced by electrolytic (Goppelsroder, Jahresber 1876, 702) or by atmospheric (Green,

J. Soc Dyers, 1908, 231) oxidation

Pure aniline on oxidation yields a blue-black, the homologues and allied compounds giving somewhat different shades; e g orthotoluidine produces a brownish-black, paraphenylene diamine a deep black, and para-aminodiphenylamine (diphenyl base) a greenish black. The oxidation of a-naphthylamine on the fibre results in the formation of naphthylamine violet, but this colour is not of much importance.

The composition of aniline black has been the subject of much investigation (see Armand Muller, Chem. Zentr. 1871, 288; Goppelsroder, Jahresber. 1876, 702; Nietski, Ber. 1878, 1094; Caro Verhandt d Ges Deutscher Naturforchen u Arzte, 1896, 119; Willstatter and Moore, Ber. 1907, 2665; Marsden, J Soc. Dyers, 1908, 9; Willstatter and Moore, Ber 1909, 4118; Green and Woodhead, Chem. Soc. Trans. 1910, 223).

Three chief and successive products of the oxidation of aniline may be distinguished-emeraldine, nigraniline, and ungreenable black

Emerabline is produced by the limited oxidation of amline, several residues linking up to form a chain of 4 molecules (Willstatter and Moore) or a chain or ring of 8 molecules (A.G. Green). The aniline residues are in all probability linked by nitrogen atoms in para-position, since para-compounds only have been obtained on oxidation or reduction.

Emeraldine is a greyish-green body which is always formed during the early stages of the production of aniline black. By the action of oxidising agents, it is converted into nigraniline and by the action of reducing agents it forms first proto emeraldine by addition of two atoms, and then leuco-emeraldine by addition of four atoms of hydrogen (Green, Chem. Soc. Trans 1910, 223). More powerful reducing agents split the molecule and produce amino- and para diamino-compounds or quinone

Nigraniline, the second oxidation product of aniline, appears to contain eight aniline residues, and is said to be produced either by removal of two hydrogen atoms from the emeraldine molecule (Green) or by polymerisation with loss of hydrogen (Willstatter and Moore)

Nigraniline or aniline black is a violet-black body which is unaffected by alkalis, but which is turned green by mild reducing agents (sulphurous acid), owing to its reconversion into emeraldine. Dilute mineral acid also brings about indirectly the production of emeraldine from nigraniline. Oxidation of a portion of the latter substance probably takes place at the expense of the reduction of other molecules, since both quinone and emeraldine are formed in the reaction.

Ungreenable black, the final product in the technical formation of aniline black, is produced by the further oxidation of nigraniline in presence of aniline. It is a very stable body, and is not readily attacked by acids, alkalis, oxidising or reducing agents. Powerful oxidising agents, however, convert it almost quantitatively into quinone, and powerful reducing agents into a brown leuco-compound which is reoxidised to the original black on exposure to air. The presence of free aniline is essential to the production of ungreenable black, which suggests that it is a condensation product of emeraldine and aniline.

Technical production of aniline black—Of the large number of modifications of Lightfoot's original process which have been suggested, very few have withstood the test of practical experience The following technical methods may be distinguished (1) dyed blacks; (2) aged blacks; (3) steam blacks.

(1) Dyed blacks, also known as one-bath blacks, are used for the dyeing of cotton yarn, which is often previously dyed in a sulph de black

vat. The ingredients used are  $(e\ g)$  aniline (12 lbs.), hydrochloric acid (18 lbs.), sulphuric acid (24 lbs.), and sodium dichromate (24 lbs.), in 40 gallons of water. The use of hydrochloric acid alone produces a blush-black, and sulphuric acid alone a reddish-black, the mixed acids yielding a jet black

The yarn is worked in the solution whilst slowly heating to the boiling-point, and the bronzy black thus produced is steamed to convert it into a good black which is ungreenable. This process does not tender the cotton Many

modifications of the process are used

(2) Aged blacks are obtained by padding (saturating) the cotton with a solution containing aniline salt, some oxidising agent (sodium chlorate) and an oxygen carrier (copper sulphate, or vanadium chloride), after which the fabric is dried and 'aged'; the colour being developed by passing through an ageng chamber in which a moist atmosphere and a temperature of about 45° are maintained. The padded cotton has a green colour (emeraldine) which turns black (nigraniline) during ageng. The final change to ungreenable black is brought about by working the fabric in a solution of sodium dichromate

Copper aged blacks are very largely produced on cotton piece goods by a process which differs only in detail from Lightfoot's original process

Varadium blacks are chiefly used in calico printing Lightfoot (Bull Soc Ind de Mulhouse, 1871, 285) first noticed the special efficiency of vanadium salts as oxygen carriers in the production of aniline black, and Witz (Bull Soc Ind de Rouen, 1876, 310) showed that 1 part of vanadium chloride was sufficient to convert 40,000 parts aniline into aniline black

The production of aged blacks is liable to tender the cotton unless the process is very

carefully controlled

Air oxidation black—A G Green (J Soc Dyers, 1909, 191) has introduced an amline black process in which atmospheric oxygen is the oxidising agent, some carrier such as cuprous chloride being present. The mineral acid usually employed is also replaced by formic acid, and by these modifications the hability to tendering of the fibre is greatly reduced.

The process is based on the fact that amline oxidises much more rapidly in presence of a small quantity of a paradiamine or a para-amino-

phenol

After padding, the goods are dried, aged, and chromed as in the case of the ordinary aged black

(3) Steam blacks are more costly than aged blacks, but safer as regards tendering of the fibre. They are produced by padding with an aniline solution which contains sodium chlorate as oxidising agent, and sodium ferrocyanide as oxygen carrier. After padding, the goods are dried, steamed, and chromed Lightfoot's original steam black only differed from this in that it contained potassium ferricyanide in place of the less expensive sodium chlorate and ferrocyanide.

Anilne black, when successfully dyed on cotton, is the fastest and most intense black which it is possible to produce. Its chief drawbacks are the tendency to turn green and the liability to tendering of the fibre, unless great experience and close supervision are exercised

Application of an line black to wool and silk.

—The animal fibres are rarely dyed aniline black.

The black may be produced on silk by the same methods as on cotton, but the process is chiefly confined to the silk-cotton union material used as umbrella cloth.

Aniline black does not readily develop on wool, but if the wool is previously chlorinated (Lightfoot) or boiled with a mineral acid (Bethmann, Eng Pat 21236, 1901), the aged black may be produced—It is railly met with

(b) The insoluble azo-dyis—The sulphonated

(b) The insoluble azo- dyes—The sulphonated azo- dyes which are so largely used in the dyeing of wool and silk have little or no affinity for cotton. Azo- compounds, however, may be produced on cotton by impregnating the fibre with one of the constituents, and then passing it into a solution of the necessary combining body. The non-sulphonated compounds are used in this process, and the insoluble dyes thus obtained are, in many cases, useful colours

The constituent parts of an azo-dye are (a) a phenol or amine, and (b) a diazotised base, and the process (introduced by Holliday, in 1880) consists in saturating the cotton with an alkaline solution of a phenol, drying, and then developing

in a solution of the diazotised base

In calico printing (rarely in dyeing) the fibre may be impregnated with a paste containing the whole of the colour ingredients, and the colour developed by steaming

Substances employed in the process · Phenols—The only phenols practically employed in the padding operation are β-naphthol and β-oxynaphthoic acid anilid (naphthol AS)—Amines are not used

Aromatic bases—The following list gives the principal bases employed and the colours they produce when combined with \(\beta\)-naphthol.—

Colour produced with Base β-naphthol Orange yellow Anılıne Paranitraniline Red (with copper, brown) Paratoluidine Yellow orange Nitroparatoluidine Orange Xylıdıne Red Cumidine Bluish-scarlet Scarlet Phenetidine Blue shade of claret red a-Naphthylamine Ammoazobenzene Cumson Benzidine Purplish-brown Tolidine Reddish-brown Dianisidine Blue (with copper) Azo-black base Black

Padding or grounding —The intensity of the ultimate colour depends entirely upon the concentration of the padding solution, which, for a shade of medium strength may contain, per litre, 25 grams  $\beta$ -naphthol, 45 grams sodium hydroxide solution 36°Tw, and 75 grams Turkey-red oil. Cotton is evenly saturated in the cold with this solution, then dried at a low temperature and developed

The developing process—The most important dye produced by this process is paranitraniline red, which is dyed on a very large scale and used as a substitute for Turkey-red. The diazosolution is prepared as follows. stir to a paste 10 lbs. paranitraniline with 2 gallons water; then redd 2 gallons hydrochloric acid (35°Tw) and 6 gallons water. After mixing well, add 5½ lbs sodium intrite dissolved in 2 gallons water. Dilute to 200 gallons, and add sodium

acctate to neutralise the excess of hydrochloric acid. The colour is produced instantaneously on passing the padded cotton into this solution, the cotton being subsequently washed, dried, and soaped.

Various stable forms of the diazotised base appear under the names azophor red, nitrazol,

and nitrosamine red.

Azo-oranges, yellows, crimsons, claret-reds, &c, are obtained in a similar manner by the use of the various bases given in the above table

Azo-browns are produced by adding an alkaline solution of copper to the  $\beta$ -naphthol prepare, and developing with paranitraniline or  $\beta$ -naphthylamine

Azo-blues are obtained by padding with β-naphthol, developing with dianisidine, and subsequently treating with cupric chloride and

chromic acid

(c) The metallic dyes. Iron buff and Nankin uellow.-These two colours, which differ only in depth of shade, are produced by precipitating They are hydrated ferric oxide on the fibre dyed chiefly on cotton, and produce yellow to light-brown colours which are fast to light, washing, or alkalis, but are readily attacked by The cotton is padded in a solution of a ferrous or ferric salt, and then passed into an alkaline solution. If the cotton is previously mordanted with tannin, more even shades are produced, the black tannate of iron being subsequently destroyed by a treatment with sodium peroxide (Knecht) or other oxidising agent Iron buff is only produced on wool or silk as an intermediate product in dyeing Prussian blue

Manganese bronze—This colour is produced on cotton by precipitating manganous hydroxide on the fibre, and oxidising this to the brown

manganic peroxide.

Chrome yellow is produced on cotton by impregnating with a solution of lead acctate, drying, and passing into a solution of potassium dichromate

Chrome orange —This compound is a mixture of normal lead chromate (chrome yellow) and basic chromate It is obtained by treating cotton dyed chrome yellow with lime water

Prussian blue is produced on cotton by first dyeing an iron buff, and then passing into an acid solution of potassium ferrocyamide. This colour is extremely fast to light, but is sensitive to alkalis. It is produced on silk in a similar manner as a constituent in dyeing that fibre black. Prussian blue is little used on wool, but is readily produced either by a process analogous to the above, or by making use of the fact that when heated in acid solution with wool, the prussiates undergo decomposition with formation of Prussian blue.

COLOURING MATTERS USED IN DYEING

Natural dyestuffs · Indigo —There are two distinct methods of applying indigo in dyeing, and the most important, the vat method, has already been described (v. Vat dyes, p 586) Indigo may also be used as an acid dyestuff by converting it into the disulphonic acid. This sulphonic acid is known as indigo carmine or indigo extract. It dies a brighter blue shade than vat indigo, but it is much less fast to light and washing. It is largely used on wool and silk, but is inapplicable to cotton.

Logwood is a mordant dyestuff and is used for all fibres On wool it is largely employed other dyestuffs for browns, olives, greys, &c

Potassium or sodium dichromate is the usual mordant, but a single-bath logwood black may be produced by using a mixture of logwood, ferrous sulphate, and oxalic acid Logwood is now chiefly employed in the form of unoxidised (for cotton) or oxidised (for wool) extract On silk and cotton, logwood is only used for black dyeing

Brazilwood, peachwood, and Limawood constitute the class of soluble red woods. They are now very little employed, the colours they produce being fugitive to light. They produce reds with aluminum, reddish-browns with chromium, and purples with iron mordants.

Cumwood, barwood, and sanderswood are known as the insoluble red woods. If wool is boiled with the ground wood, it acquires a leddish-brown colour, which is still employed as a 'bottom' for indigo vat blue. They are, however, mordant dyestuffs, producing brownish reds with chrome mordant.

Madder—This dyestuff, once of great importance, has now been almost entirely superseded by synthetic alizarin. It is still used to a limited extent in the dyeing of wool felt, since it penetrates through the material more readily than alizarin. With chromium mordant it gives a reddish-brown, with aluminium a red, and with non a purple shade. The use of madder in the wood vat has already been referred to (v. Vat dyes). This dyestuff is not now

used on cotton or silk

Cochineal —The use of this valuable dyestuff in dyeing is now confined to the production of scarlet shades on woollen cloth, for which purpose considerable quantities are still imported, Stannous chloride is the mordant employed, and the colour is produced by the one-bath method. The aluminium and tin lakes are used as pigments in painting and printing.

Lac-dye may be employed in the same way as cochineal, and gives similar shades. Since the commercial product contains a considerable amount of mineral matter, the ground dyestuff is treated with dilute hydrochloric acid before use

Cudbear and orchit (or archit) —These products are essentially the same dyestuff, the former in the solid and the latter in the condition of paste. It is one of the few natural colouring matters which die wool without a mordant. Wool and silk may be died in a neutral, slightly acid, or slightly alkaline, solution. It produces full, rich crimsons, and is still used to a considerable extent. The colouring matter is not applicable to cotton.

Safflower was formerly much used in dyeing bright pinks upon cotton, but is now obsolete as

a dyestuff

Annatto produces orange shades on silk without a mordant, it is applied in a hot soapbath. Annatto is still used in colouring cheese and butter.

Weld—This mordant dyestuff produces pure yellows in conjunction with aluminium mordant. When used as a 'topping' for indigo vat blue, a special shade of green, known as 'carriage green,' is obtained.

Fusic is the most important of the natural yellow dyewoods, and is still employed in

conjunction with chrome mordant on wool as the yellow constituent in compound browns, olives, greys, &c It is chiefly used in the form of fustic extract

Osage orange is closely alhed to fustic utilisation as a dyestuff is only of recent date and is confined to America.

Quercitron bark and flavin are also natural yellow mordant dyes, producing somewhat brighter shades than fustic, but are now obsolete

Persian berries -This mordant dyestuff is employed by the calico printer for producing

bright orange shades with tin mordant

Turmeric is the only representative of the natural colouring matters which acts as a direct dye towards cotton, silk, and wool It produces yellow shades, which are brighter if dyed in a bath acidified with alum.

Catechu, cutch, and gambier —These dvestuffs are still largely employed in cotton dyeing cotton is steeped in a strong solution of the dye, and attracts the colouring matter The cotton is then passed into a boiling solution of potassium dichromate, which, by oxidation, produces insoluble brown japonic acid upon the A little copper sulphate is frequently added to the catechu bath, whereby fuller and faster shades are produced

Silk is dyed in a similar manner for the production of weighted blacks, the chromium, fixed during the oxidising process, subsequently acting as a mordant for logwood

Other natural dyestuffs —Colouring matters are very widely distributed in the vegetable kingdom, and all countries where the textile arts have attained any great development (notably India, Persia, and Japan) had until recently each its own series of indigenous products which were used as dyestuffs In Donegal, Sutherland, and other localities, the homespun tweeds, &c , are still largely dyed by means of the various lichens, 100ts, barks, nuts, &c , found in the neighbourhood

Synthetic dyestuffs. The number of individual synethetic dyestuffs now at the dyer's disposal is very large, and includes dyes yielding all possible hues. There is infinite variety in the cháracteristics of the various artificial dyes, and in most cases it is not difficult to match a given shade or to comply with given conditions as regards properties Being, therefore, more convenient to use, and in very many cases cheaper and faster than the natural dyestuffs, the latter have been largely superseded

The dyeing properties and the characteristics of the dyed colours depend chiefly upon the chemical groups to which the dyes belong, but, as indicated in the section dealing with the classification of dyes, the chemical groups do not coincide with groups based on the method of application (v. Classification of colouring

matters, p 575).

(1) Nitroso colouring matters — This class includes fast-green or dark-green and naphtholgreen. In conjunction with iron mordants, both these dyestuffs give olive-green colours re-

markable for their fastness to light

Cotton is dyed with dark-green by mordanting with iron, and dyeing in a neutral bath. Wool is mordanted with ferrous sulphate and oxalic acid, and dyed in a neutral bath. Silk is dyed in a similar manner.

Naphthol green which contains non as a constituent is only applicable to the animal fibres. Wool is dyed with the addition of suiphuric acid.

(2) Nitro-colouring matters -- The most important members of this class are pieric acid, Victoria orange, aurantia, palatine orange, naphthalme yellow, naphthol yellow These colouring matters are only applicable to the animal fibres, which are dyed in a bath slightly acidified with sulphuric acid. The colours they yield are not fast to light Pierre acid yellow is remarkable for its extreme freedom from any trace of orange On exposure to light it rapidly becomes brownish, owing to reduction of the nitro groups

(3) The azo- group —These may be divided into monazo, disazo, trisazo, and tetrakisazodyes, according to the number of azo-groups they contain They may be further classified into aminoazo- (NH<sub>2</sub>) and oxyazo- (OH) dyes, and into other subdivisions based upon the state of combination of the azo- group A few of the azo- colours are basic dyes, a large number of the acid dyes, and acid-mordant dyes, belong to this group, and practically the whole of the direct dyes. The azo-dyes vary greatly in their degree of fastness to light, and in other properties.

(4) The di- and tri-phenylmethane group -This group includes most of the basic dyes, such as auramine, magenta, methyl violet, malachite These bodies, when sulphonated, green, &c form acid dyes, and therefore such dyestuffs as acid magenta, guinea green, &c., and also patent blue, Victoria blue, fall under this group As a class, these dyestuffs are fugitive to light

(5) The oxyketone group - This group includes most of the artificial mordant dyes, eg the alizarin fieds, yellows, blues, browns, &c Most of the dyestuffs of this group are extremely fast to light and other influences, which renders them of the greatest importance

(6) The pyronine group —This includes (1) the pyronines proper, a small class of basic dyes, of which pyronine red is an example, (2) the much more important group of the phthaleins latter includes both basic and acid rhodamines, the cosms and the mordant dyestuffs gallem and cœrulem

(7) The acridine group —This includes a few basic dyestuffs of which the chief is acridine

(8 and 9) The oxazine and thiazine group -This group also includes both acid (thiocarmine), basic (methylene blue), and mordant dyes (gallocyanın)

(10) The azine group includes the valuable basic dyes the safranines, and the basic and

acid indulines and nigrosines

The dyeing of mixed fabrics of cotton and wool may be carried out by selecting those colouring matters for which both fibres have an attraction, eg the benzidine dyes, &c., or by first dyeing the wool in the usual manner with an acid colouring matter, then mordanting the cotton with tannic acid and tartar emetic or stannic chloride, and dyeing in the cold with a basic colouring matter The so-called 'crossdyeing' process is carried out by first dyeing the cotton (usually with sulphide or other 'ast' dyes) in warp form, then weaving with a

wool weft and dyeing the latter by means of acid dyes. Logwood blacks on union are dyed by first mordanting the cotton in the cold with myrobolans and nitrate of iron, then the wool with a boiling solution of potassium dichromate, and finally dyeing both fibres thus mordanted in a hot decoction of logwood Mixtures of cotton and silk are dyed by methods similar to those employed for cotton and wool. Fabrics of wool and silk are dyed by methods similar to those usual for wool or silk alone

When dyeing compound shades, the main principle to be observed is to apply simultaneously only such colouring matters as are applicable, when used singly, by methods as similar as possible to each other Basic and acid colouring matters ought not to be applied together in the same dye-bath, since the one requires a neutral, the other an acid, bath in order to yield their full colouring power Basic colouring matters, however, may be quite well applied to mordanted cloth along with phenolic colouring matters, since both these classes usually dye in a neutral bath.

It is very irrational to produce compound shades by the application of a mixture of fast and fugitive colours, the best results are obtained by employing colouring matters of about equal fastness to various influences observed, the dyed colour is hable to change materially in shade under those influences to which it is subsequently submitted, eq. light,

For further information on the subject of dyeing, the following works may be consulted 1813 The Philosophy of Permanent Colours E Bancroft

1835-1861 Recherches chimique sur la Tanture E Chevreul.

1869 Dictionary of Dyeng and Calico-C O'Neill. punting

A Practical Handbook of Dyeing and Calico-printing W Crookes.

1876 Dyeing and Calico-printing. F C Calvert

1885. The Dyeing of Textile Fabrics. J J Hummel

1893 La Pratique du Temturier Vol. 1, 1893, Vol. 2, 1894, Vol. 3, 1897. J Garcon

1901. Dictionary of Dyes, Mordants, and other Compounds used in Dyeing and Calico-printing C Rawson, W M Gardner, and W F Laycock (' Rawson,

1903. Theorie und Praxis der Modernen Farberei A Ganswindt

1904 et seq. Systematic Survey of the Organic Colouring Matters G Schultz and P Julius, transl by A G Green

1907 Farbereichem Untersuchungen Anleitung zur Untersuchung und Bewertung der wichtigsten Faiberei, Bleicherei, Druckerei, und Appretur-Materialien 2nd edit. P Heermann

Étude sur les Industries du Blanchiment, de la Teinture, de l'Impression. O Picquet

1910. A Manual of Dyeing. 2nd edit. E. Knecht, C. Rawson, and R. Lowenthal.

Identification of the Commercial Dye-

stuffs. Mulliken.

1912 Bleaching and Dyeing of Vegetable Fibrous Materials. J. Hubner. Farbestofftabellen G Schultz.

Mercerisation und Appretur P. Gard-

L'Industrie des Matières Colorantes Organiques A Wahl

1913. Synthetic Dyestuffs and Intermediate Products from which they are derived 2nd edit. J. C Cain and J F Thorpe

1915. The British Coal-tar Colour Industry. W M Gardner

The Analysis of Dyestuffs

A G Green DYESTUFFS, IDENTIFICATION OF, ON FABRICS. The recognition of the particular dyestuff which has been employed in dyeing a fabric is frequently a master of considerable difficulty. If one dyestuff only has been used, it may be possible to identify it with little trouble, but when, as is often the case, two or more dyes have been employed to produce a given colour, their separation and recognition

may be a very complex problem

In the identification of dyestuffs, the ordinary basis of qualitative analysis is as far as possible made use of, i e the group to which the colouring matter belongs is first determined, and then tests are applied to identify it as an individual substance. There are, however, such a large number of closely related compounds amongst the dyestuffs, many being isomeric, and differing only (eg) in the position of a sulpho-group, that with the small amount of substance frequently at disposal it is quite impossible to definitely identify a specific commercial pro-

duct This is, however, of minor importance
The recognition of a dyestuff is much facilitated by a knowledge of the dyeing properties of the various groups of colouring matters. In a given case this may greatly limit the possibilities For example, no one with a knowledge of dyeing processes would search for a cochineal scarlet on cotton or a paranitraniline red on wool - The presence or absence of certain metallic bases in the ash of the incinerated material is also a useful indication of the nature of the dyestuff under investigation, and an examination of this point should be made

What may be termed the 'dyeing groups' of colouring matters do not coincide with the 'chemical groups'. For example, the dyeing group of 'basic' dyes, the members of which are fixed on cotton by means of a tannin mordant, includes members of the triphenyl methane, azine, azo-, acridine, and other chemical groups

To determine the dyeing group to which a colouring matter present on a fabric belongs, acid and alkaline solvents are used as 'stripping agents. The following are the general reactions:

ACTION OF 'STRIPPING AGENTS.'

Dyeing group Mordant dyes Basic dyes on wool " on cotton

Acid dyes Direct cotton dyes

Stripping solution 50 c c HCl (conc ) per litre 5 pc acetic acid 3 p.c. sodium hydroxide

3 p.c. ammonia 0 880 s.g. A small piece of mer-cerised cotton is placed in the solution. In each case the fabric is boiled for 5 minutes in the solution.

With direct cotton dyes the small piece of white cotton placed in the solution becomes dyed. With acid dyes this does not occur

The action of the above stripping agents can only be considered as general in regard to a particular group, and many exceptions will be found in dealing with particular members of the group, e.g. some acid dyes are only slightly stripped by boiling in dilute ammonia, and a few of them stain mercerised cotton. Too much reliance must not, therefore, be placed on these stripping reactions.

The most satisfactory group reagents for classifying dyestuffs into the chemical groups are acid or alkaline reducing agents, on the use of which several schemes of identification have been based. Those reagents were first proposed by Witt, in 1886, whose scheme was extended by Weingaertner, in 1887; but the systematic development of the scheme of analysis is mainly due to A. G. Green.

The reactions of dyestuffs are considerably

influenced by the nature of the fibre upon which they have been fixed, which also determines the particular dyeing process necessary. For example, the basic colours dye wool and silk direct, whereas their fixation on cotton involves the use of a tannin mordant. It is convenient, therefore, to treat separately the identification of dyestuffs on animal and on vegetable fibres.

Identification of dyestuffs on wool—Tables for the identification of dyestuffs on wool have been published in Hummel's Dyeing of Textile Fabries; in Knecht, Rawson, and Loewenthal's Manual of Dyeing, in Lehne's Farber Zeitung, and by many other authors; but the most systematic and reliable scheme is that of Green (Analysis of Dyestuffs, 2nd edit 1915) In this scheme, which is now to be described, the dyeing groups are first distinguished by the action of suitable solvents, and then further differentiation is carried out by the use of reducing agents with subsequent reoxidation. The various groups of dyestuffs as dyed on animal fibres, show the following reactions—

Decol	lourised by hydrosul	phito		Not decolourised, but changed in	
Colour restored on exposure to air	Colour not restored by air, but re- stored on oxidation with persulphate	restored other by	Not altered by hydrosulphite	shade Original colour restored by air or persulphate	
Azines, oxazines, thiazines, indi- goids	ines, indi-		Pyrone, acridine, quinoline, and thiazol groups. Some members of the oxyketone group	Most dyestuffs of the oxyketone group	

In Green's scheme the following reagents are employed, and must be used of the strength indicated.—

Dilute ammonia (1 100) —1 cc conc ammonia; 100 cc distilled water

Aqueous-alcoholic ammonia—1 cc conc ammonia; 50 cc strong alcohol or methylated spirit; 50 cc water

Dilute acetic acid (5 p c ) —5 c c glacial acetic acid; 95 c c. water

Dilute alcohol (1:1)—50 cc strong alcohol or methylated spirit; 50 cc water

Dilute hydrochloric acid (1 · 10) —10 cc conc, hydrochloric acid; 90 cc. water.

Caustic soda (10 p c) —10 grams solid caustic soda in 100 c c of water.

Hydrosulphite AX —Dissolve 50 grams hydrosulphite NF cone or rongalite 6, ie the formaldehyde compound of sodium sulphoxylate, in 150 c.c. of hot water. Add to the solution at 80°-90° 0 25 gram of precipitated anthraquimone previously ground to a paste with a little of the solution, and dilute the whole to 500 c c with cold water. The solution should be slightly alkaline. It may become acid on keeping.

Hydrosulphite B.—Dissolve 50 grams hydrosulphite NF conc. or rongalite C in 500 c.c. water, and acidify with 2 c.c. of acetic acid. This is used only for yellows and oranges.

Vol. II.—T.

Persulphate —A cold saturated solution of potassium persulphate, or a 1 pc solution of ammonium persulphate. If alkaline, neutralise carefully with acetic acid.

Sodium acetate (5 p c)—5 grams crystallised

sodum acetate, 100 c c distilled water

Procedure—The reactions are carried out in test-tubes upon pieces of the material about 3-1 inch square, which are covered with about 1-11 inches of the reagent The tests should be carried out exactly as described. Except when specially stated, the reagents are to be applied in succession to fresh portions of the In making 'stripping tests,' the degree of stripping is judged by comparing the remaining depth of shade with that of the original pattern The colour of the stripping solution cannot be relied upon as a guide boiling with dilute acetic acid and dilute ammonia, it is found advantageous to repeat the extraction, as a better stripping is thereby obtained, and with acid dyestuffs any staining of the cotton by the first strong extract is removed during the second boil In testing removed during the second boil with dilute ammonia or sodium acetate, the material is placed in a test-tube with a piece of white mercensed cotton cloth, somewhat smaller in size, and boiled for the time preser bed. If the shade is a pale one, the size of the sample should be increased and that of the cotton

diminished The dilute ammonia is replaced by aqueous alcoholic ammonia in the case of the violet and black dyestuffs (Tables III. and VII), as in these cases the acid dyestuffs are less easily extracted, and the cotton is more liable to be stained by them In making reduction tests, the sample is boiled for from onequarter to one minute with the hydrosulphite, then rinsed well under the tap, and allowed to he on white paper With most dyestuffs which form air-oxidisable leuco- compounds, the colour ieturns immediately or in a few minutes, but with others a longer time is required. The reaction is accelerated by exposing the pattern to ammonia If the colour does not return, the pattern is boiled with a little water in a testtube, and potassium persulphate is added drop by drop, carefully avoiding an excess also fails to cause any return of colour, the dyestuff is to be regarded as a nitro-, nitroso-, or azo- compound The depth of the restored colour varies greatly in different cases, with some dyestuffs the colour reappears with nearly its original depth, but with others (probably on account of the greater solubility of their leucocompounds) only a pale shade may return Safranine and its azo- derivatives are peculiar in yielding a violet colour on reoxidation of the leuco- compound, this being due to the condensation of the leuco-safranine with the formaldehyde present in the hydrosulphite NF

The tables (I to VII, pp. 596-602) give the

scheme of analysis

These tables do not always differentiate between closely related individual dyes, for which purpose further tests must be made stage of the investigation it is very desirable to procure small pieces of material dyed with such known colouring matters as the group tests, combined with the colour of the sample, have indicated as probable Parallel individual tests should then be applied with these and with the sample under examination

The following series of reagents have been found by W M Gardner to be convenient for this purpose They should be applied in the

order given, and used as follows -

Application of cold reagents - Steep a piece of the material 1 by 1 inch in (say) 5 cc of reagent for 5 minutes in a small porcelain

Application of reagents by boiling —In a testtube place 10 cc of reagent and a piece of material 1 by 1 inch Boil gently for 3 minutes

Application of nitric acid test - On a white glazed porcelain plate, spot the material by dipping a glass rod into the acid and then pressing it on to the cloth Note the changes in colour which occur during the formation of the spot, and re-examine after several hours, noting any further change

Reagents for identifying individual dyestuffs

1. Boiling water Use small pieces 2. 5 p c. acetic acid, boiling of white wool,

3. 3 p c. ammonia, boiling. silk, and cotton 4 3 pc soap, boiling along with the 5 Boiling alcohol dyed sample

Concentrated sulphuric acid, cold 3 p c ammonium persulphate, cold.

8 Bleaching powder solution, 5°Tw., cold

9. 50 grams SnCl<sub>2</sub>,2H<sub>2</sub>O+100 cc HC

 $32^{\circ}$ Tw per litre, cold 10. 5 c c. NaNO<sub>2</sub> (5 grams per litre), an

5 c.c. H<sub>2</sub>SO<sub>4</sub> (3 5 grams per litre) boiling.

11. Apply test 10 cold, then wash an steep the material in a solution of sodiu β-naphtholate 5 grams per litre

12 Concentrated nitric acid (spot test)

If the sample under examination shows tl same reactions as any one of the control sample with each of the above reagents, its identif may be considered as clearly established

Knecht (J Soc Dyers, 1904, 98) states the titanous chloride possesses advantages ov both stannous chloride and sodium hydrosulphi as a test for dyestuffs The solution is prepare by boiling together 25 c c commercial titanoi chloride solution (20 pc) and 25 cc HC 32°Tw, and diluting to 1000 cc with water.

IDENTIFICATION OF DYESTUFFS ON COTTON.

Green (Analysis of Dyestuffs, 1915) he worked out an analytical scheme for the detetion of dyestuffs on cotton on the same lines a that previously described for wool

The reagents are as follows .-

1 Dilute ammonia -1 c c strong ammon to 100 c c distilled water

2 Dilute caustic soda -10 grams causti soda in 100 c c water

3 Saline vaustic soda -10 c c caustic sod solution (35-40 p c NaOH) to 100 c c saturate solution of salt

4 Formic acid 90 pc -- Concentrated con mercial acid

5 Dilute formic acid—1 ee formic aci

(90 pc) to 100 cc distilled water 6 Dilute hydrochloric acid -- 5 cc hydro

chloric acid (30 pc) to 100 cc of water 7 Soap solution -10 grams soap in 300 c.

of water 8 Tannin solution -10 grams tannic aci

and 10 grams sodium acetate in 100 cc o

9 Bleaching powder solution —Fresh solutio at 5°Tw

10 Hydrosulphite AX (same as for woo see p 593)

11 Hydrosulphite B (same as for wool, so p 593)

12 Hydrosulphite BX —Dissolve 50 gran of rongalite or hydrosulphite NF cone in 125 c of hot water. Grind I gram of anthraquinor (precipitated, not sublimed) to a fine powde and reduce to a smooth paste with a little of th rongalite solution. Add this paste to the remaining solution, and heat the whole for 1 or 2 minutes at about 90°. Then dilute wit cold water to 500 cc, and, after standing unt cold, add 1½ c c of glacial acetic acid a well-stoppered bottle with greased stopper The reagent should be tested occasionally k trying its effect on cotton dyed with a-naphthy amine bordeaux, which should be fully di charged after boiling for 1 or 2 minutes anthraquinone may be replaced by  $\beta$ -oxyanthr quinone, which can be employed in the san way or previously dissolved in a little alcoh before it is added to the hydrosulphite. the advantage of giving a clearer solution, by is not so generally available

13 Persulphate solution —A cold saturate

solution of potassium persulphate, or a 1 pc solution of ammonium persulphate.

14. Acid stannous chloride —100 grams stannous chloride to 100 c c. hydrochloric acid (30 pc) and 50 cc water. The hydrochloric acid must be free from arsenic. In testing for sulphide colours, this reagent may be replaced with advantage by a strong solution of titanous

15 Chromium fluoride reagent -A solution of 10 grams chromium fluoride and 5 grams sodium acetate in 100 c c water

Procedure: General -All the reactions are performed in test-tubes, with pieces of material about  $\frac{1}{3}$ —\frac{3}{2} of an inch square, which are covered to the depth of from  $1\frac{1}{2}$  to 2 inches with the reagent. The degree of stripping is judged by comparing the remaining shade with that of the original pattern, the colour of the stripping solution cannot be relied upon as a guide testing the colours of calico prints, the separate colours should be cut out and the reactions tried upon them separately With cotton and wool, or cotton and silk unions, the weft is separated from the cotton warp, and each submitted to Doubt may frequently exist as examination to how a particular shade should be classified, whether, for instance, a reddish-blue is to be regarded as a blue or a violet In such cases the tables referring to both colours should be employed, but since the same general scheme is adopted throughout the tables, this can scarcely give rise to confusion. The same remark applies to possible mixtures, thus in examining a certain green shade, both the yellow and the blue tables may have to be used

To distinguish between individual dyestuffs of the same group the same method should be adopted as in the case of wool (see p. 594)

It is obvious that in many instances the analytical procedure may be greatly simplified by exclusion of dyestuffs the presence of which is rendered improbable or impossible by the special circumstances of the case, eg by the material, shade, or mode of application of the colour

Stripping test -A few of the direct dyestuffs are partially stripped by dilute ammonia, and may thus give rise to the impression that they are acid colours To avoid this error, it is advisable to add a small piece of white cotton when carrying out the test With an acid dyestuff the cotton is either not tinted or becomes white on boiling a second time with dilute ammonia

Transference of basic dyes to wool -The tannin mordant is first removed, as in testing for a basic colour, by boiling the pattern for half a minute with saline caustic soda It is then well washed to remove all alkalı, and is boiled with a piece of white wool (half the size of the cotton or less) in a little water for 1 or 2 minutes. In most cases the dye base will leave the cotton almost entirely, and dye the wool a If the colour does not develop on full shade the wool, one or two drops of dilute formic acid (1:100) may be added. In the case of a few dyestuffs which are more difficult to strip (e.g. basic greys), it is necessary to extract the colour with dilute hydrochloric acid (1.20), carefully neutralising the extract with ammonia before adding the wool.

Tannin test for basic dyes -Add a few drops of tannin solution to the formic acid extract. Shake well, and if the precipitate does not form at once, allow to stand a few minutes Some colouring matters such as the rhodamines, gallocvanines, and chrome colours of the rosaniline series (which contain carboxylor hydroxyl- groups in addition to basic groups) only precipitate slowly, and the precipitate, being more finely divided, is sometimes difficult to see

Bleeding test for direct dyestuffs -In testing for direct dyestuffs by the bleeding test, the sample is placed in a test-tube together with a smaller piece of white mercensed cotton cloth, and boiled with soap solution for about a minute The soap solution may be replaced by a 5 p c.

solution of sodium carbonate

Lead acetate test for sulphide dyes—The sample is first boiled for two initutes with a 10 pc solution of sodium hydroxide to remove any adventitious sulphur compounds. It is then well washed and just covered with acid stannous chloride solution The mouth of the test tube is closed by a cap of filter paper closely wrapped round it, into the centre of which is placed, by means of a glass rod, one drop of lead acetate solution. The contents of the test-tube are slowly heated to the boiling-point, when a blackish-brown stain of lead sulphide appears if The brown spot a sulphide colour is present will again disappear on continued boiling on account of the decomposition of the lead sulphide by the hydrochloric acid evolved It is important to pay special attention to the cleanliness of the test-tubes employed for this test, as tubes which have been previously used for hydrosulphite reductions acquire a thin invisible deposit of sulphur upon their walls, which, on boiling with stannous chloride, gives rise to hydrogen sulphide, and thus may lead to error

Reduction and reoxidation tests —The reduction with hydrosulphite BX is carried out by boiling the sample with the reagent for from a The azmes, minute to a minute and a half thiazines, oxazines, &c., and most of the azo-dyestuffs are fully reduced in about half a minute, but the insoluble azo- colours and some direct dyestuffs require longer boiling to complete their reduction In testing the reoxidisability by air, the reduced sample may be exposed to the fumes of ammonia, which in many cases accelerates oxidation

Tests for vat dyestuffs—(See Green and Frank, J Soc Dyers, 1910, 83; and Grand-mougin, Tabellarische Uebersicht der Kupenfarbstoffe)

The analytical tables worked out by Green for the identification of dyestuffs on vegetable fibres, are given on pp 603-609, Tables VIII. to XÍV

ST Mulliken (Identification of the Commercial Dyestuffs, 1910) has published a systematic scheme of analysis, primarily intended for the examination of the dyestuffs in substance. The tables give the reactions of 1475 distinct dyestuffs, and are of great service in the identification of dyes on the fibre

Spectroscopic examination of dyestuffs.—The absorption spectra of dyestuffs may be used as an aid to their identification. In the second

## RECOGNITION OF DYES ON WOOL OR SILK. TABLE I—YELLOW AND ORANGE COLOURS.

		£	or brown	hanged to yellow-brow —Alizarin class. hodant is present	n   c	Alizarın red S, &c., on Sn, alizarin orange SW on Cr, Al. Sn.
	1001 (1:100)	or Vat dyestuff	by persul- lest for a	Mordant absent —Azc Salt dyestuff. Confir: by boiling with 5 p c. sodium acetate and a piece of white cotton stained		Chrysophenine, chrysamine, diamine fast yellow 3G, diamine yellow N, benzo fast yellow and orange, congo orange, cotton yellow, carbazol yellow, pyramine orange, diamil orange, triazol yellow G, sulphon yellow, &c.
	nute amn	Iordant oulphite A	d Colour by air or ordant (Cordant Cordant Cordan Corda Corda Corda Corda Corda Corda Corda Corda Corda Cord	Cotton is stained.— Salt-mordant dye- stuff	15	Autochrome orange, mercerol yellow 2R, mercerol orange 2R, salicine orange G, salicine yellow G, metachrome yellow, &c.
	dyestuff. Boil twice for one minute with dilute ammonia (1:100) Keep the ammoniacal extract	remains white (Mordant or Boil with hydrosulphite AX	Decolourised Colour stored either by air or phate.—Azo class.  Mordant present	totton in stained Salt-morradant dye- salt-morradant dye- salt-morradant dye- stuff  Cotton in stained Salt-morradant dye- stuff  Cotton remai dye- stu		Acid alizarin yellow, eriochrome yellow, alizarin yellow G, 2G, 3G, R, flavazol, diamond flavine, anthracene yellow, chrome fast yellow, fast mordant yellow, salicine yellow D, diamond yellow, chrome yellow, chrome orange, &c.
	acal extract		s on exposure to more quickly on ion with persul-	Cotton is stained — Sali dyestuff Stilbene class Cotton remain white — Indigol Vat dyestuff.	-	Curcumine S, Mikado yellow, golden yellow and orange, direct yellow, stilbene yellow, diamine fast yellow A, AR, diamine orange D, dianil direct yellow S, chloramine orange, sun yellow, &c.
	uff. Boil tw	tripped The (Direct dyes	Decolourised. Colour returns on exposure to air, or more quickly on oxidation with persulphate. Boil with 5 p c.	Cotton is stained—Salt dyestuff Still-bene class white—Indigol Vat dyestuff. Confirm by extended to the confirm of the confirmation of the confirm		Helindone yellow and orange, ciba yellow and orange, ciba indigo yellow 3G, thioindigo yellow, and orange, &c
c. acetic acid	, or Mordant dyest white cotton Keep	Little or no colour is stripped The cotton or is tinted (Direct dyestuff).	changed changed lest for a	Mordant absent —Thi- azol Salt dyestuff. Confirm by boiling with 5 pc sodium acetate and a plece of white cot on The latter is staine	=	Thioflavine S, diamine tast yellow B, FF, M, sulphine, primuline, chloramine yellow, brilliant pure yellow, chlorazol fast yellow, thiazol yellow, Clay ton yellow, chlorophenine, dianil pure yellow HS, naphthamine pure yellow G, mimosa yellow, &c
D	ect, or M	Little or	Colour un only sligh in shade	Mordant present — Flavone or Ketone class		Fustic, querentron bark, weld, flavine, alizarin yellow A and C, galloflavine, &c
Boil twice for one minute with 5	-Acid, Vat, Direct, and a piece of	e AX or B	and colour does not return o air or upon oxidation with —Azo-group or Nitro- Add cone HCl to the moniacal extract	Solution becomes violet or violet-red	6	Metanil yellow, orange IV, Victoria yellow
twice for		ton reman	and colour does not rear or upon oxidation —Azo-group or Niti Add cone HCl to the	Solution becomes red		Fast yellow, Indian yellow, azoflavine, curcumeine, orange II, wool fast yellow.
Boil	Little or no colour is stripped	colour is stripped, but cotton remains white dyestuff. Boil with hydrosulphite AX or		Solution becomes colourless or pale yellow	7	Naphthol yellow S. Martius yellow, citronine.
	or no colo		Decolourised, on exposure to persulphate group.	No change of colour		Tartrazine, oranges G, 2G, &c, R, fast light yellow, xylene yellow, flavazine.
	Little	Much cold Acid dye		olour is not affected — noline or Pyrone group		Quinoline yellow, uranine, eosin orange, turmenc.
	dyestuff.	Deco by	colourised Colour is not restored y air or by persulphate —Azo- group			Chrysoidine, tannin orange, new phosphine, Janus yellow.
	Much colour is stripped.—Basic dyestuff Boll with hydrosulphite AX or B	only very	olourless solution. Boil with dilute HCl (1 · 10)	Fibre and solution pale yellow		Thioflavine T, rhoduline yellow.
	ur is strippe with hydros	Not decolourised or only very slightly. Treat fibre with conc. H.SO.	Colourless Boil with HCl (1 Thiazol	Completely decolourised	61	Auramine O and G.
	Much color Boil	Not deco	Green fluorescent solution :— Acridine group			Phosphine, benzoflavale, rheonine, patent phosphine, acridine yellow, acridine orange, &c.

1 Weld is partly discharged by acetic acid though the extract remains colourless.

minute with 5 p.c. acetic acid

Boil twice for one

## RECOGNITION OF DYES ON WOOL OR SILK TABLE III—PURPLE AND VIOLET COLOURS.

	100) and	Boil with	Colour changed to coange-brown or brown or brown or brown or list present as the change of the chang	18	Alizarın bordeaux, alizarin claret, alizarin maroon.											
	- 1	stuff.	Colour Crange Class or ange or	17	Alizarin red S, &c., on Fe.											
	oholic ammoi	or Salt dyestuff.	Mordant absent Salt dyestuff Confirm by boiling with 5 pc sodium acetate and a specific of which were of white certain	16	Diamine, benzo, dianil, columbia, chlorazol, naphthamme, chlorantme, &c., violets.											
	queous alc	Vat, Mordant, hvdrosulphite AX	solum acctate and a price of white cotton Stained  Stained  Mordant present Acid mordant dyestuff.	15	Acid alizarin violet, acid chiome violet, eriochiome violet, Palatine chiome violet.											
	ute with a	ed Vat,	Decolourised Colour does not return on exposure to air, but is restored by persulphate Triphenylmethane class. A chrome mordan is present	, <b>#</b>	Chromogene violet, metachrome violet, chrome violet											
	or one min	ur is stripį	Occasion of the control of the control occasion of the control occasion of the control occasion occasi	60	Helindone violet, ciba violet, ciba heliotrope, thiomdigo violet, &c.											
c acid	onl twice for te cotton	Little or no colour is stripped	by extraction with anime and sublimation  By extraction with anime and sublimation  Worldant present Azine, Oxazine, or Thiazine class.	12	Gallocyanine, prune, celestine blue, correine.											
c acetı	uff Be of whit		Colour unaffected Pyrone class. A chrome mordant is present	11	Galleme											
te with 5 I	Bul twice for one minute with 5 p c acetic acid  Acid, Vat, Mordant, or Salt dyestuff Bul twice for one minute with aqueous alcoholic ammonia (1 a piece of white cotton		Colour is changed to orange-yellow, restored by persulphate to a violet- brown Anthraquinone class	10	Anthraquinoue violet, alizain duect violet, alizain insol, alizain cyanol violet											
one minut			Decolourised Colour is not restored either by an or by persulphate Azo class	6	Lanacyl violet, azo acid violet, sulphon violet, Victoria violet erio fast purple, wool violet											
il twice for	Vat, Mor	nams white	does not return it is restored by yinethane and Soil with 5 p c cone mmute Luttle colour is stripped Spot right all hire with our hydrochlore acid he un. Yellow hauged spot	œ	Acid violets, alkali violet, guinea violet, formyl violet, kitone violet, &c											
B	1	pped, but cotton rem Boil with hydrosulp	pped, but cotton rem Boil with hydrosulp	pped, but cotton rem Boil with hydrosulp	our is stripped, but cotton remains white Boil with hydrosulphite AX	pped, but cotton rem Boil with hydrosulp	pped, but cotton rem Boil with hydrosulp	pped, but cotton rem Boil with hydrosulp	pped, but cotton rem Boil with hydrosuly	pped, but cotton rem Boil with hydrosulp	pped, but cotton rem Boil with hydrosuly	pped, but cotton rem Boil with hydrosuli	pped, but cotton rem Boil with hydrosuli	en pur en	l-	Pyrone colours, $ey$ , fast acid violets, violamines, and violet $4R$ , &c.
	no colour is stripped													pped, but Boil with	pped, but Boil with	pped, but Boil with
	5		Decolourised One exposite Persulphate Pyrone cli Pyrone cli Pyrone cli Sodum a Much colou Stripped, but tion remans cli fibre with colou sulphurte Colour Colour Colour to fibre to fibre to colour to fibre to colour colo	5	Certain red-shade acid violets, e.g., acid violet 4RS and 5RS, red violets 4RS, &c.											
	Little	Much co	Decolourised Colour returns on exposure to air Azine, Oxazine, or Thiazine class	4	Fast blue R, red shade indulines, &c											
	Soluble	boil twice for a with dilute	Unaffected Al or Cr is piesent in ash On boiling with drifte ammonia, the colour becomes much blue	en   1	Soluble red woods, e g, Brazilwood, Limawood, sapanwood, peachwood, &c.											
	olour 18 st. yestuff or	unute with cohol (1 1	Decolourised Colour does not return on exposure to air, but is restored by per sulphate Triphenyl-methane class  Personal Proceedings of Colour Action 1988	1	Methyl violet, ethyl violet, crystal violet, benzyl violet, brilliant violet, &c.											
	Much colour is Basic dyestuff Red Wood Bo		autholise Triphenyl- methane class  Decolourised Colour returns on exposure to all Azine, Oxazine, or Thiazine class		Neutral violet, roduline violet, methylene violet, tannin heliotrope, iri. violet, &c.											

## RECOGNITION OF DYES ON WOOL OR SILK. TABLE IV.—BLUE COLOURS.

1	ΨX	Co agai	n on expost	i to brown, become to air —Aliza	ing blue rin class	26	Alizarın blue (Anthraquinoline class).
	dyestuff. Boil with hydrosulphite	Colour is not r by air or by	Morda Confi sodu whit	ordant present  at absent —Salt  rm by boiling wit  macetate and a percent of the cotton. Cotton	h 5 p c piece of stamed	25	Diamine, benzo, dianil, Chicago, chlorazol, naphthamine, &c, blues and sky blues, sulphon acid blue, sulphoneyanine, sulphonezurine, erio fast blue SWR, tolyl blue GR extra and 5R extra, &c
monia	ıl with	rised.	ant Boil c so-	Cotton is Salt m	stained .— ordant stuff	24	Autochrome blue
lute am	tuff. Bo	Decolourised. restored either	Test for a Mordant Dresent Bo with 5 p c so dium acetate	로요를 white —	remains Acid-mor- lyestuff	23	Chromotropo blue, acid chrome blue, chrome fast blue, anthracene chrome blue, fast mordant blue, salicine chrome blue, &c
ıth di	dyes	on ex	ourised Co	lour does not retur r, but is restored i	y absent	55	Brilliant milling blue B
ute w	Salt		lass. Tes	riphenylmethan t for a mordant	present	21	Chrome blue
min	nt, or	on ex	lant int fibre miline	Colourless solut Pyrone dyes	stuff	20	Fast acid blue R, violamine 3B
Boil twice for one minute with dilute ammonia ammoniacal extract	-Vat, Mordant,	Colour returns on ex Test for a mordant	Mordant absent Treat fibre with aniline	Blue solution gi evaporation a su residue — Indig dyestuif	blimable	13	Indigo, helindone blue $3\mathrm{G}$ and $2\mathrm{B}$ , ciba blue, thromdigo blue, $3\mathrm{C}$
il twic	-Vat	Solour Test	It	Fibre and solu	tion red	18	Errochrome azurol, errochromo cyanine, chromo worsted blue, chromal blue, chromoxane blue, &c
stuff.	Little or no colour is struped -	Decolourised (	Mordant present — Azine. Oxazine, or hiazine class Treat fibre with cold conc. sulphune acid	Fibre and soluti violet		17	Gallocyanme, celestine blue, correme, prune, gallamme blue, delphine blue, bulhant delphine blue, mdahzaum, lanoglaucine, ultra cyanine, ultra violet, &c
Salt dye	olour		Mor Azu Thia fibre	Fibre and solut	ion green	16	Brilliant alizarin blue, mercerol brilliant blue, indochromine, &c
or Salt	r no	d, or	thra one Test Cr)	Mordant al	sent	15	Indanthrene blue WB.
Mordant, or	Little or no co		er —Anthra- quinone class. Test for a mor- dant (Cr)	5 <b>0</b>		1.4	Alizarin cyanine, brilliant alizarin cyanine, anthracene blue, alizarin indigo blue, acid alizarin blue 2B and GR, &c
Vat. Mo	멸	Fibre 1s not de-		bight yellow to orange- brown, returning with persulphate red-violet, returning with persulphate		13	Alizarin astiol, alizarin uranol, cyananthrol, alizarin direct blue B, alizarin cyanol B, alizarin sky blue, anthraquinone blue, &c
Boil twice for one minute stripped Acid Vat. Ma (1 100) and a piec		Fibre 1	—Anthragu none class Colour of re			12	Alizarin saphirol, anthracyanine, alizarin direct blue EB and E3B, alizarin (yanol EF, brilliant anthrazurol, enocyanine LM, fast sky blue, &c
twice upped	ains w	De	ecolourised by air or by	Colour is not rest persulphate —A	tored either zo class	Ţ	Lanacyl blue, azo acid blue, azo merino blue, azo navy blue, chromazone blue, azoc yanine, &c
	ton rema	Colour	r, but bersul- nenyl- ass	Ammoniacal ex colourless, but be on acidify	comes blue	10	Soluble blue, water blue, alkalı blue, chlorazol brilliant blue 14B, ısamıne blue 8B, brilliant dianil blue 6G, brilliant sky blue 8G, &c
Little or no colour	our is stripped, but cotton remains white	T. Boll with hydros Decolourised Colour	does not return on exposure to any but is restored by persul- phate — Triphenyl- methane class	Treat fibre art fibre with color color bind. Fibre red- con Fibre red- coran color c	Fibre and solution colourless to pale yellow  The property of the property of the party of the p		Patent blue, new patent blue, Neptune blue, xylene blue, brilliant acid blue, kitone blue, erioglaueino, eriocyamine, eriochlorine, cyanole, cyanine, fast acid violet 10B, cyanogene, &c.
Lutti	stripi	Dec Dec			brown to ge-brown	· α	Wool blue, cyanole FF, intensive blue, &c
	lour is	ayes	ar and and sses.	The ammoniac colourless F in a	e is present		Prussian blue
	Much co	Decolour	r retured to e., Oxas azine, go cla	The ammonia- cal extract is blue On	heating	9	Thiocarmine, fast blue, induline, acid cyanine, fluorescent blue, lazuline blue, &c
	M.	) de	Colour returns on exposure to air — Azine, Oxazine, Thazine, and Indigo classes.	adding caustic	yellow	2	Indigo extract, indigo carmine
	od.		On spot	ed Alor Crispre ting with hydrocl ue is changed to b	iloric acid	4	Logwood blue,
	stripped or Logwe	Coil with dilute alcohol (1:1) twice for one minute	1	Decolourised Avreturns on exp air —Safranine stuff	violet colour	8	Janus blue, diazine blue, indoine blue, Janus dark blue, naphthindone, &c.
	h volour fa	rith dilute nce for or	1 1 2 2 E	Decolourised Col- return on exposur is restored by per Triphenylmeth	e to air, but sulphate :	64	Night blue, Victoria blue, setocyanine, seto- glaucine, turquoise blue, &c.
	Mucl Basic	Your tr	Much co Basic with h	Decolourised. Co on exposure to air Oxazine, or This	::—Azine,		Methylene blue, Nile blue, Capri blue, cresyl blue, Meldola's blue, new methylene blue, di- phene blue, &c.

## RECOGNITION OF DYES ON WOOL OR SILK. TABLE V.—GREEN COLOURS.

5 p c acetic acid

witn

minute

one

Boil twice for

r changed to e or orange-i. Alizarin class Original colour does not return on exposure to air, Alizarin cyanine green, alizarin viridine, but is restored by perbrilliant alizarin viridine, &c. sulphate A chrome mordant is present and Colour coorange of brown. with Original colour returns on Cœruleine, alizarin green S, alizarin dark, 100) exposure to an green W. mordant is present Bollv J Colour changed to red-violet The ammonia original colour is restored by perdyestuff. Eriochrome veidone, acid alizarin green. A chrome mordant is sulphate present restored —Azo or mordant Mordant absent -Salt Diamine, benzo, dianil, Columbia, chlorazol, dilute Salt dyestuff. Confirm by boiling with 5 p c sodium acetate naphthamine, &c , greens. and a piece of white cotton or with Colour is not re r by persulphate -ss. Test for a m Wordant been of the property o -Vat, Mordant, hydrosulphite AX Diamond green 2S, chrome patent green N, minute mercerol green, &c one Diamond green B and 3G, chromoxal green, either by air or by Nitroso class. Cyprus green, omega chiome green, eriofor chrome green H and L, acid chrome green, &c twice Little or no colour is stripped gambines, dioxine, Naphthol green B. Salt dyestuff. Boil piece of white cotton chrome green G, &c. Colour does not return Decolourised on exposure to air, but is restored by persulphate Triphenylmethane class. A Cr mordant is present Chromoxane green, chrome green, fast chrome green, &c. Decolourised
Colour returns on
exposure to air
Test for a mordant Mordant absent
Indigoid Vat dye
stuff. Confirm by
extraction with amiline Ciba gieen G, helindone green G. Mordant, or and sublimation test Mordant present -Oxazine or Thiazine 6 Alizarin green B and G class -Acid, Vat, Fibre is not decolourised, but colour changes—
Anthraquinone group. Colour of reduced fibre is— Anthraquinone green, alizarin cyanine green, Orange, returning but cotton remains white with hydrosulphite AX with persulphate to alizarin brilliant green, fast acid green RH, grey-green anthraquinone blue-green. Red-violet returning colour is stripped with persulphate to bluish-green Alizarin emeraldol, alizarin direct green. De lourised Colour is not restored Sulphon acid green, mixtures of an azo either by air or by persulphate .-Azo class blue and yellow. Much colour is stripped. Acid dyestuff Boil ou Acid green, light green, guinea green, wool green, Noptune green, alkali fast green, Decoloursed Colour does not return on exposure to air, but is restored by persulphate —Triphenylö cyanole fast green, erio green, brilliant methane class milling green B, night green, &c. Decolourised Colour returns on exposure to air —Azine, Oxazine, Azıne green S. or Thiazine class Decolourised. Colour does not re Much colour is stripped — Basic dyestuff Boli with hydrosulphite AX Malachite green, brilliant green, fast green, turn on exposure to air, but is restored by persulphate —Triphenyl setoglaucine, solid green, China green, new fast green, &c. methane class Decolourised Original colour re-Methylene green, Capri green, azine green, turns on exposure to air -Azine, fast green M. &c. Oxazine, or Thiazine class Decolourised A dark violet colour returns on exposure to air - Safranine Azo dyestuff Janus green, diazine green.

## RECOGNITION OF DYES ON WOOL OR SILK TABLE VI-BROWN COLOURS

2		
Š		
7		
ממון		
-		

	(1 100)	M	Colour of fibre unaltered Boil with dilute hydrochloric acid (1:10)	Colour 1s not stripped Cr and Cu present	10	Cutch.						
	Ith dilute ammonia	Boil with hydrosulphite AX	Colour of fibi Boil with dilut	Colour is stripped Cr present	6	Anthragallol, anthracene brown, alizarin brown.						
	Boil twice for one minute with dilute ammonia (1		Colour is not restored a by persulphate .— Boll with 5 p c and a piece of white or two minutes	Cotton 1s stained Cr may be present Salt dyestuff	8	Diamine, benzo, dianil, Columbia, Congo, naphthamine, toluylene, Hessian, &c, browns.						
ic acid		-Vat, Mordant, or Salt dyestuff	Decolourised Colour is not restored either by air or by persulphate — Azo group Boil with 5 p.c. sodium acetate and a piece (white cotton for two minutes	Cotton remains white Cr is present Acid- mordant dyestuff	2	Anthracene acid brown, acid alizaim biown, anthracene chromate brown, diamond brown, palatine chrome brown, salicine brown, metachrome brown, mercerol brown, &c Also manganese brown (Mn in ash)						
Boil twice for one minute with 5 p c acetic acid	Little or no colour is stripped -Acid, Vat, Mordant, or Salt dyestuff.	is stripped -Vat,	Decolourised Colour does not return on ex-	Decolourised Colour does not return on ex- posure to an, but is restored by persulphate		Chromogen, oxychromine.						
	ed —Acid, Vat, M	Little or no colour is stripped	Decolourised Colour returns on exposure to air Boil with 5 p.c. sodium acetate and a piece of white cotton for two mnuttes	Cotton 1s stamed —Salt dyestuff stilbene class	ō	Mıkado browns.						
Boul	no colour is strippe		н	<b>-</b>	H	<b>-</b>					Cotton remains white — Indigold vat dyestuff	77
	Lattle or		Much colour is tripped —Acid dyestuff On boiling with hydrosulphite AX, decolourised Colour is not	restored either by air or by persulphate — Azo class	es	Acid brown, fast brown, resorcin brown, guinea brown, solid brown, sulphon acid brown, Neptune brown, wool brown, &c.						
	s stripped .—	yestun rosulphite AX	Decolourised. Decolourised. To voice colour sed. Capturns on restored ether by air or		63	Bismarck brown, Janus brown, tannin brown.						
	Much colour i	Boil with hydrosulphite AX			1	Diazine brown.						

## RECOGNITION OF DYES ON WOOL OR SILK. TABLE VII.—BLACK AND GREY COLOURS.

	of white cotton			Colour changed to brown Original colour returns slowly on exposure to air			Naphthazarıne, alızarın blue black SW, W, brilliant alızarın black, alızarın black S, WR, WX, &c.
	100) and a piece of white cotton		Boll with hydrosulphite AX	narged Boll for a 5 p c sodium e of white cotton	Cotton 1s stained	10	Diamond black FB, FR, NG, GA, 2B, Palatine chrome black, anthracene acid black, alizadine black, monochrome grey, anthracene chromate grey, &c.
	oholic ammonia (1	minute with dilute		The colour is discharged Boll for one minute with 5 p c sodium acetate and a piece of white cotton	Cotton remains white	6	Diamond black PV, PVB, P2B, F, diamond blue black EB, cricefrome black, enochrome grey, chrome fast black, anthracene chrome black, and alizarin black, fast chrome black, and chrome black, &c
	th aqueous alc	Boil for half a	Fibre unaffected	Treat original cold conc ric acid	Colourless	œ	Aniline black.
h c seeme sen	ff. Boll twice for one minute with aqueous alcoholic ammonia (1	Little or no colour is stripped —Mordant dyestuff.		Unaffected Treat original fibre with cold concault sulphure acid	Blue solution	L.	Alızarın cyanıne black.
d o man command on o man more			Millioner	Fibre blue or blue-vrolet, solution crimson Test for indigo by extraction, evaporation, and sublimation			A 'vatted black' (indigo and logwood)
	or Salt dyestu		•	Fibre and solution orange to pale brown			Logwood on Fe, Bonsor's black
•	ripped -Acid, Mordant, or Salt dyestuff.	Little o		Fibre and solution crimson			Logwood on Cr.
	r is stripped —Ac	ped —Acid or stuff		The cotton is stained —Salt dyestuff. On bolling with hydrosulphite AX, the colour is permanently discharged			Union black, ('olumbia, diamine, diamil, benzo, chlorazol, naphthamine, &c, blacks and fast blacks, direct deep black, direct blue black, &c.
The second secon	Little or no colour is st	The colour is stripped Salt dyestuff		The cotton remains white— Acad dyestuff. On boiling with hydrosulpinte AX, the colour is permanently discharged			Naphthol black, naphthylamine black, nerol, azo acid black, Palatine black, sulphocyanine black, sulphon black, Biebrich patent black, ercsol black, &c.
				Much colour is stripped .— Basic dyestuff			Janus black, diazine black, jute black, methylene grey, new fast grey, &c.

# RECOGNITION OF DYES ON VEGETABLE FIBRES TABLE VIII—YELLOW AND ORANGE COLOURS

			DY.	ESI	OFFS,	IDENTIȚICAT	101	N OF, ON FABRICS 603
	wo drops	Reduce original	The colour su unffected or changed in shade, becoming lighter, browner, or bluer. If altered in shade, exposure to air restores the original colour. Apply the lead acctate test.	No H <sub>2</sub> S is evolved Boil with soap solution and white mercensed cotton	The white cotton is not stained.—Mordant or vat dyestuff Boll with 90 pc formic acid Also test ash for me-	Colour not stripped by formic acid No mordant in ash —Vat dyestuff —Colour	18	Helindone orange R, helindone yellow 3G, anthraflavone G, algole yellow 3G and R, algole orange R, indanthrene yellow, indanthrene orange RT and golden orange indanthrene copper, erbanone yellow R and orange R.
	l one or ty	d <b>R</b> educ	affected or cha ter, browner, le, exposure to colour Apply acetate test	svolved ]	The white stained.— vat dyesti 90 p c f	stripped by formic acid Alor Cr in ash — Mordant dyestuff	124	Persian berries on Al'or Cr_mordants, alizarin orange on Al mordant
	h water an	tannin is not obtained lrosulphite BX	uris unaffe ing lighter, I in shade, original co	No H <sub>2</sub> S is e	The w	white cotton is Thiazol salt lyestuff	16	Chlorophenme, chloramme yellow, diamine fast yellow B, FF, and C, Clayton yellow, thuazol yellow, thioflavine S, &c Primuline developed with hypochlorite
	boil wit	nnin is r sulphite	The colo becom altered the	н	2S 1s evolv dy	18 evolved —Sulphide dyestuff		Yellows and oranges of the immedial, katigene, cross-dye, pyrogene, thiogene, sulphur, thional, &c, series
	water, and	able by ta		- I	er gett	Colour is not pped Cr present the ash —Azo rdant dyestuff	1,4	Chrome orange, ahzann yellow R, GG, &c diamond flavine, flavazol, &c
	pool q	precipi	hight y tored b (including soap	corto	white not sta with	lour is stripped -Insoluble azo colour	13	Metanitianiline orange, mtrotoluidine orange, orange from naphthol A('
na (1:100)	da, rinse witl i (1 100)	icid extract ]	or becomes ur is not rest zo dyestuff (3) Boil with	te mercerised		Fibre after re- action is colour- ess —Salt azo dyestuff	12	Chrysophemne, chrysamine, carbazol yellow, toluylene yellow and orange, stilbene yellows and oranges, benzo, Congo, diamino, and diaml yellows and oranges, pyramine orange, Pluto orange, &c
	salır wea	A coloured acid extract precipitable by tannin is not c fibre with hy Irosulphite BX	Decolourised or becomes light yellow. Original colour is not restored by air oxidation —Azo dyestuff (including the Stilbene class) Boil with soap solution	and whi	Salt d	thre after reduc- on is light yellow and can be diazo- tised and de- eloped red with etanaphthol— Primuline azo- colour	11	Primuline developed with phenol or with resortinol, cotton yellow G and R, oriol yellow, dianil yellow, &c
ak ammo	ute with	ly or	ng a huch 18 solu- uff il with		dec	s permanently olourised .— azo class	10	Chrysoidine, tannin orange, Janus yellow, azo phosphine, new phosphine, &c
Boil with weak ammonia (1:100)	Boil for a quarter of a minute with of	The colour is completely or largely stripped, giving a coloured acid extract, which is precliptated by tannin solution.—Basic dyestuff Transfer to wool and boll with		ydrosulphite B	Woolis not decolourised Boil cotton with hydro- chloric acid (1 20)	Not decolourised The cone H <sub>2</sub> SO <sub>4</sub> and alcoholic solutions are fluorescent — Acridine class	6	Phosphine, benzoflavine, actidine yellow, rheonine, orange flavophosphine, &c
	l for a q	The co	The collargel; largel; coloured precipit tion fransfer		Yoolis n Soil cott chloric	Decolourised	œ	Auramme
							1-	Flavinduline
	is not stripped	pletely dest	ns and fibre being Treat original fibre nmonium sulphide	1s not blackened h hydrosulphite BX	olourised Al in ash	A1 absent	9	Throflavine T, methylene yellow, rhoduline yellow
	The colour 18	our is com	both solutions colourless Tre	Fibre 18 Boil with	Not decolourised Test for Al in ask	Al present	5	Alızatın yellow A.
	Ŧ	The col	colou with		Cı prese	lackened nt in ash	7	Chrome yellow or chrome orange (lead chromate).
	ripped d water		Colour is transferred to the wool —Acid	lydro- ite B	P	ermanently colourised .— Azo class	က	Indian yellow, orange IV , G, &c.
	colour is straight	and small piece of white	Colour is to	with h	Not qui	decolourised .— Pyrone or Inoline class	63	Quinoline yellow, eosine orange
	The co		- 1	nr not transferred to the wool. Sn is present in ash			1	Persian berries on tin mordant.

## TABLE IX—RED COLOURS.

Boll with weak ammonia (1:100)

e or two	ed or changed rown. Original d on exposure racene class. c forme acid. for mordant	Cr absent — Anthracene Vat dyestuff	16	Algole red B, indanthrene red, indanthrene claret, ciba scarlet G.
ite with water and one or tw fibre with hydrosulphite BX	Colour unaffected or change o maroon or brown. Origin colour restored on exposure to ar. — Anthracene class Boll with 90 p. c forms and Also test ash for mordant	Cr absent Anthracene Vat dyestuff Cr in ash Anthracene Mordant dyestuff	15	Alizarin, purpurines or alizarin maroon on Cr mordant .
ite with with dibre with	Colour to mare colour to aur Beil w	Colour stripped by formic acid Al in ash —Anthracene Mordant dyestuff	#1	Turkey red, alızarın red, alızarın pınk, alızarın maroon.
or one minuce original	tholPrin	ed to greemsh yellow d to original colour ch can be diazotised dred with betanaph-nuline Azo colour	13	Primuline developed with betanaphthol or with R-salt
Boul for a quarter of a minute with saline caustic soda, rinse with cold water, and boul for one minute with water and one or two drops of weak formic acid (1 100)  A coloured acid extract precipitable by tannin is not obtained Reduce original fibre with hydrosulphite BX	1254 +	No H <sub>2</sub> S is evolved Red vapours on heating —Indigoid vat dyestuff	12	Algole red 5G and scarlet G, algole pink R, ciba red G and scarlet G, ciba bordeaux B, helindone red B and 3B, helindone scarlet S, helindone fast scarlet R, thioindigo red B, thioindigo scarlet, vat red
arth cold 100) nm is not	Become less exposu or Indi	H <sub>2</sub> S is evolved — Sulphide dyestuff	11	Thiogene rubine, &c
oda, rinse v nic acid (1 ble by tani	tored by air or Boll with soap sed cotton te cotton is not de Boll with pyridine	The colour is not stripped Cr in ash —Mordant Azo dyestuff	10	Chrome red, brilliant chrome red, chrome bordeaux, &c.
f a mmute with saline caustro soda, rinse with cold water, and drops of weak formic acid (1 100) coloured acid extract precipitable by tannin is not obtained	Decolourised and colour not restored by air or by persulphate —Azo class. Boll with soap solution and white mercerased cotton. The white cotton is 'The white cotton is 'The white cotton is not stande —Salf dye- stande Boll with stuff Test ash for pyridine	The colour is stripped — Inscoluble Azo colour	os.	Paranitramiline red, alphanaphthylamine bordeaux, chloramisidine pink, introanisidine pink, scarlet and reds from naphthol AC.
te with sa drop	rised and color liphate —Azo tion and white nite cotton is —Salt dye- Test ash for	Cr or Cu present —Azo Salt dyestuff after- treated	×	Diamine fast red F, &c
ter of a minu A coloure	Decolourised by persulphic solution The white stained —Stained —Stuff Tes	No Cr or Cu present — Azo Salt dyestuff	r-	Benzopurpurine, diamine scarlets, diamine reds, benzo fast scarlets, diazo brilliant scarlet, rosanthienes, Zambesi red, erica, diamine rose, geranine, rosophenine, &c.
for a qua	rgely extract, olution. r other nd boil	Wool decolourised Colour not restored by air or by persul- phate —Azo class	9	Janus red.
it stripped Boil	s completely or largely a coloured acid extract, tated by tannin solution tuff (on tannin or other ransfer to wool and boil ydrosulphite AX	Wool decolourised. Colour does not return on exposure to arr, but is restored by persulphate.— Triphenylmethane class	o	Magenta, fuchsine, new fuchsine, isorubine, cerise, grenadine, &c
5	lour is grying precipi dyes t) T vith h	Wool decolourised Colour returns on exposure to air — Azine class	4	Safranines, rhoduline reds and pink, azine scarlet, induline scarlet, neutral red, &c
The col	The extripped, which is I—Basic mordan	Wool not decolourised — Pyrone class	8	Rhodamines, rhodines, irisamine, anisoline, rosazein, rhoduline pink, acridine reds, &c.
hour 18	Trans- rool and h hydro- te AX	Decolourised. Colour not restored by air or persul- phate .—Azo class	61	Crocein scarlets, brilliant croceines, Palatine scarlet, fast reds, &c
The colcus	dyestuff fer to w boll wit sulphi	Not decolourised — Pyrone class	1	Eosines, phloxine, erythrosine, 10se bengal, &c.

## RECOGNITION OF DYES ON VEGETABLE FIBRES. TABLE X.—PURPLE AND VIOLET COLOURS.

		Authrace:  Apply tate feet to the control of the co	ne	20	Indanthrene violet or violanthrene.
	mth	would be a colour of the colou	ie	19	Thiogene dark red, &c
	minute	The colour is unaffected or of charged in stands being a stands of the colour is numbered or of charged in stands of the colour is numbered in stands of the colour or Application of the charged in the colour or of the	or	18	Alizarın cyanıne 3R, alizarın bordeaux
	l for one	and the control of the color of	is	17	Gallem, alızarın violet, alızarın elaret, alızarın eyelamine, alızarın red on Cr mordant.
	i), and bor	Authracein by Colour is unaffected or of the colour or holds of the colour of the colour is unchanged by the colour of	d on Indi-	16	Ciba violet, ciba heliotrope, thioindigo violet, helindone violet
	(or salt solution), and boil for one minute with	ourised ourised ourised ourised ourised ourised ourised our castored o	red on Mor- zine 'not	15	Gallocyanine, modern violet, ultra violet, coreine, &c
	water (	Sulphide dyes	tuff ur is	17	Thiogene violet, katigene violet, hydrone violet, &c
	with cold c acid (1	restored by an or castored by a castored	nt in Ior-	13	Chrome bordeaux, chrome prune, &c
100)	da, rinse eak formi	or Alizable  or Alizable  Doll verised of Alizable  Boll verised of Alizable  acid (11  The collapse of Alizable  with with the collapse of Alizable  with with the collapse of Alizable	our ed ble our	11	Benzidine puce
Boil with weak ammonia (1:100)	e caustic so drops of w	per of the contract preceptiable by tall again and described by a day of the colon and a da	olu- ent rin	11	Alizarın purple
weak ar	ith salm	Croured acide decoloured acide decoloured acide decoloured a personal part of the colour of the colo	lzo uff	0	The undermentioned (col 9) coppered or chromed
3oil with	minute w	dyestuff	Cu t	ை வ	Violets of the diamine, benzo, Congo, Hessian, Columbia, chlorazol, chlorantine, diamil, oxamine, and rosanthrene series
	Boll for a quarter of a minute with salme caustic sods, rinse with cold water and one or two drops of weak forme and (1	ぬかって しゅってきる nhenvimeth	Col- re- ut ls per-	00	Chrome violet
		to larged, striked, s	ns to		Gallocyanine, modern violet, ultra violet, coreine, prune, &c
	ot stripped.	Decolourised Colis not restored half or persulphate —Azo	b <b>y</b>	۰ .	Janus claret red, &c
		Decolourised Col does not return air, but is restor	in ed		Methyl violets, ethyl violet, benzyl violet, erystal violet, &c
	TI.	Hoge (Oxazine or Thiazine) clas	s s	" :	Methylene violet, rhoduline violet, iris violet, neutral violet, tannin heliotrope, &c.
	<del></del>	Pyrone class	-		Anisoline.
	Ir is	The ammonia solution is vio	cal		Acid violets, formyl violets, alkali violet, guinea violet, &c.
	S'ripped —A	Colon trans- tourn	our-	٦	Red shades of soluble and alkalı blues.

# RECOGNITION OF DYES ON VEGETABLE FIBRES TABLE XI.—BLUE COLOURS.

Colour unaffected or becomes darker, browner, &c. the original colour being restored on exposure to air —Anthracene mordant or anthracene vat dyestuff (also Ultramarine). Boil with 90 p.c. formo acid The colour is not much affected Test ash for Cr Cr absent :-Indanthrene blues, algole blues. Anthracene vat dvestuff for one minute with Cr present in ash —Alizarin dyestuff on Cr Alizarın blue, alizarın cyanınes or anthracene blues on Cr mordant. The colour is stripped Al in ash Apply lead acetate test No H<sub>2</sub>S evolved: Alizarin dye-stuff on Al Alizarin cyanines or anthracene blues on Al (or salt solution), and boil mordant, erganone blues. precipitable by tannin is not obtained. Reduce original HoS is evolved Ultramarine. Colour changed to greenish-yellow, which can be diazotised and developed Primuline developed with naphthylamine ether. with betanaphthol -Primuline azo colour Decolourised or yellowish Cholour is restored on exposure to air —Azine, Oxazine, Thazine, or Indigo class, Apply lead acctate test No H<sub>2</sub>S 1s evolved Heat fibre carefully in dry test-tube No coloured vapour Cr in ash —Mor-dant thiazine or oxazine (not falling Brilliant alizarin blue, gallocyanine, gallamine blue, prune, celestine blue, delphine blue, &c. in group (1) soda rinse with cold f weak formic acid (1 Indigo (natural and synthetic), indigo MLB Violet vapours 2B, 4B, 5B, 6B, and T, ciba blue, bromindigo evolved -Indigoid dyestuff Blues of the immedial, katigene, thiogene. H<sub>2</sub>S is evolved :— Sulphide dyestuff pyrogene, sulphur, &c , series, hydrone blues Decolourised and colour not restored by air or persulpate.—Azo elass. Boil with soap solution and white mercensed cotton The white cotton is not for a quarter of a minute with saline caustic s water and one or two drops of stained Colour is stripped by boiling pyridine — Insoluble azo colour Boll with weak aminonia (1 Dianisidine blue, blue from naphthol AC A coloured acid extract The white cotton is stained .—Salt dyestuff. Test ash for Cr and Cu Cr or Cu present The undermentioned (col. 8) coppered or Azo salt dyestuff chromed after-treated No Cr or Cu present —Azo salt dyestuff Blues of the diamine, benzo, Congo, Columbia, chlorazol, dianil, oxamine, Chicago, &c , series Precipitated
Cr present in ash.
Basic mordant
dyestuff. Boil
cotton with
hydrosulphite BX Decolourised The colour is completely or largely stripped, giving a coloured and extract, which is precipitated by tannin solution.—Basic dyeatuif (on tannin or other mortaint) or Basic mortaint dyeatuff. Add chromium intoride reagent to the extract slowly Colour restored only by Chrome blue. persulphate :— Triphenyl-methane class Decolourised Colour returns on Gallocyanine, gallamine blue, celestine blue, exposure to air prune, coreine, ultracyanine, modern cyanine, &c. Boul f Oxazine class -Basic dyestuff I and boil with onte AX Decolourised Colour does not return on ex-Victoria blue, night blue, turquoise blue, colour is not stripped. posure to air, but is restored by per-sulphate —Triphenyl setocyanine, &c. methane class Colour changes to red precipitated —Bas ransfer to wool and hydrosulphite just before being de-colourised Colour returns violet or blue Safranine azo Indoine blue, Janus blue, naphthindone blue, diazine blue, &c. dyestuff Methylene blue, Nile blue, Capri blue, indazine, Basle blue, Meldola's blue, fast blue, Decolourised returns on exposure to air .—Azine, Oxazine, or Thiazine class cresyl blue, rhoduline blue, nitroso blue, &c Not 1 The colour is stripped .—Acid dyestuff (or Prussian blue) plue) The extract is colourless and Prussian blue. remains so on acidification. FeCl<sub>3</sub> gives a blue precipitate The extract is colourless, but becomes blue on acidification. Transferred to wool, the blue is decolourlese by hydrosulphite A, and restored by persulphate:—

Triphenylmethane class Alkali blues, soluble blues, chlorazol brilliant blue, isamine blue, betamine blue, brilliant dianil blue, brilliant sky blue.

# RECOGNITION OF DYES ON VEGETABLE FIBRES. TABLE XII.—GREEN COLOURS.

Boil with weak ammonia (1, 100)

		Reduced oxida	to yellow. Restored by air tion—Indigoid dyestuff	20	Ciba green, helindone green.
	e BX	wn, blue, present in hracene	Reduced colour is blue. Green shade restored by air	19	Algole green.
Boil for half a minute with saline caustic soda, rinse, and boil twice with weak formic acid	osulphit	ord Nut	Reduced colour 14 dark maroon. Green shade restored by air	80	Indanthrene green or viridanthrene, leucole dark green B.
	rth hydr	changed to red, I for Cr and Ni  No Cr or  stuff ash —A	Reduced colour is brownish- olive $H_2S$ evolved on apply ing lead acetate test	17	Indanthrene olive or olivanthrene
	Reduce original fibre with hydrosulphite BX	st ash for Chang st ash for C r N1 — dyestuff	Colour of fibre rather paler, solution brownish-yellow  Colour of fibre becomes grey, solution red Ni mash  Colour unaffected Solution colourless  Cr mash	16	Cæruleine, anthracene green
	se origina	The colour is unaffected &c Test The ash contains Cr or in thracene Mordant difference of the colour	The most of the mo	15	Alizarin green S on Ni-Mg mordant.
twice w		&c &	Colour unaffected Solution colourless Cr in ash	#1	Alizarın green S on Cr mordant.
e, and boil	obtained	The colour is The ash con Anthracene	Reduced colour is brownish- red Green restored by per- sulphate, but not by air Boiling HCl (1 20) gives bright green solution	13	Alizarın viridine, billiant alizarın viridine, alizarın eyanine green
da, rınse	in is not	Decolourised Colour returns on exposure to air —Azine, Oxazine, or	No H <sub>2</sub> S is evolved — Mordant Oxazine (or Thiazine)	ᄗ	Gallanılıc green, ındalızarıne, &c
austic soc	coloured acid extract precipitable by tannin is not obtained.	Decolourised Colour returns on exposure to air Azine, Oxazine, or	No II S is evolved — This is evolved — This is evolved — Sulphide dyestuff Sulphide dyestuff	11	Greens of the immedial, katigene, thiogene, pyrogene, cross-dye, sulphur, and thionol series
saline c		not restor	r is changed to greenish vellow, red to green by air, which can tised and developed red with thol:—Primuline Azo class	10	Primuline developed with amidodiphenylamine
nute with	tract pre	not hate Boil hite	Not decolourised Cr is present in the ash —Mordant Azo dyestuff	6	Diamond green, &c
half a mu	ed acid ex		on the colour read of the colour destroyed read of the colour read of the col	no	Russian green, fast green O, steam green, Alsace green, gambines, dioxine, &c
Boil for	A colour	Decolourised and colour estored by air or persulp Azo or Nitroso class. with soap solution and with soap solution and with mercerised cotton	Cr or Cu present	2	The undermentaoned (col. 6) coppered or chromed.
tripped		Decol restored —Azo o with si	Honor of the control	9	Diamine green, benzo green, Columbia green, chloramine green, &c
e colour is not stripped	rstripped,	0 0	recipitated Cr in ash Fibre lecolourised by hydrosulphite BX, the colour not returning in air, but restored by persulphate —Triphenylmethane class	ıo	Chrome green.
The	ely or largely	Inthorporther estuf	Decolourised Colour does not return on exposure to air, but is restored by persulphate —Tri- phenylmethane class	#	Brilliant green, malachite green, methyl green, Victoria green, setoglaucine, new fast green, &c
	our is completel	prespitated by tanning dyestuff (on tannin or or Basic Mordant dy chromium fluoude to Not precentated) —Basic	by persulphate —Triphenylmethane class closed by the phenylmethane class closed by the phenylmethane class colour returns violet or green on exposure to class  Decolourised Colour returns on exposure to	က	Janus green, diazine green, &c.
	The colour	dyest dyest or Bs chr	Decolourised Colour returns on exposure to air:—Azine, Oxazine, or Thiazine class	ଦୀ	Fast green M, methylene green, azine green, Capri green, &c.
Tì	The	colour is st	ripped:—Acid dyestuff. erred to wool is decolourised ite A and restored by per- riphenylmethane class	-	Acid greens, guinea green, Neptune green, naphthalene green, agalma green.

## TABLE XIII.—BROWN COLOURS.

-	100)		r, paler,	nd Cu	Cr and Cu absent:— Anthracene vat dyestuff, &c.			15	Indanthrene maroon, algole brown, leucole browns, cibanone brown, paramine brown (p-phenylene diamine oxidised on fibre).
		3X	becoming darker ad acetate test	Test ash for Cr and	t .—Mordant HCl (1 20)	r only slightly. e caustic soda	Solution brown, fibre unaffected	14	Cutch.
	ormic acid (1 · 1	hydrosulphite I	Unaltered or changed in shade, becoming darker, paler, yellower, &c. Apply lead acetate test	No H2S is evolved. T	or Cu (or both) present .—Mordant dyestuff. Boil with HCl (1 20)	Not stripped or only shg Boll with dilute caustic: (10 p.c.)	Fibre and solution dull violet	13	Alizarın orange on Cr mordant, alizarin red or purpurın on Cr.
	Boil for half a minute with saline caustic soda, rinse, and boil twice with weak formic acid (1 · 100)	fibre with	ered or cha yellower	No H2	Cr or Cu dyest	Com	pletely ipped	12	Anthragallol, anthracene brown, alizarin brown
		Reduce original fibre with hydrosulphite BX	Unalt	5	II <sub>2</sub> S is evolved — Sulphide dyestuff			11	Immedial cutch, cross-dye brown, katigene browns, pyrogene browns, thiogene browns, &c
				Decolourised and original colour stored by air —Indigoid class				10	Helindone brown G, cıba br <del>ö</del> wn, thıoındıgo brown.
		ot obtained.	yellov and	The shade is changed to greenish yellow, which can be diazotised and developed red with beta-aphthol:—Primuline azo-colour					Primuline developed with metaphenylene diamine, terra-cotta, &c
Boil with weak ammonia (1		coloured acid extract precipitable by tannin is not obtained.	or by per- ith soap	Boll with	color st		Not	00	Iron buff, khakı (oxides of Cr and Fe).
	lf a minute w	st precipitable	ed and colour not restored on exposure to air or by per- :—Azo class and Mineral colours Boil with soap solution and white mercensed cotton	The white cotton is not stained.		Ine colour is n —Mineral Treat with bisulphite in	Decolourised	-1	Manganese bronze
	Boil for ha	acid extra	tored on ex neral col	te cotton 1	lour 18	The colour is stripped —In-soluble azo-colour Test ash for Cu	Cu	9	Paranitraniline cutch (para red coppered).
	tripped.	A coloured	ur not rest LSS and Mi n and whil	The whi	The col	stripped soluble a colour Te for Cu	Cu	5	Para brown (chrysoidine and paranitraniline), benzidine or toludine brown
	colour 18 not stripped.	¥	1 76 40	cotton is Salt dye-	est ash for	presen salt d	Cr or Cu present:—Azo salt dyestuff after-treated		The undermontioned (col. 3) coppered or chromed
	The cc		Decolouris sulphate	The white stained — Stuff Tee		Cr and Cu absent —Azo salt dyestuff		8	Browns of the diamine, benzo, Congo, dianil, chlorazol, Columbia, Hessian, oxamine, and toluylene series.
		ext	The colour is stripped, giving acid extract which is precipitated by tannin solution —Basic dyestuff The dyestuff transferred to wool is permanently decolourised by hydrosulphite AX.—Azo class					63	Bismarck brown, Janus brown, tannin brown, &c.
		Fran	sferred t	o wo by hy	ripped —Acid dyestuff wool it is permanently y hydrosulphite AX.—				Fast brown, naphthylamine brown, acid brown, &c.

# RECOGNITION OF DYES ON VEGETABLE FIBRES TABLE XIV.—BLACK AND GREY COLOURS

		ıte	phite BX g brown,	Colour not changed by reduction (or very slightly) —Anthra-	cene class Test ash for Cr	r absent — inthracene at dyestuff	15	Indanthrene grey, indanthrene black, cibanone black, helindone black
100)		er and dult	h hydrosu e (becomin	Colour no reducta slightly)	Test a	r present — nthracene mordant dyestuff	14	Alizarın cyanınc black, alizarın blue black B.
		boil with wat	ngec n, &	Only sl restore at once presen lene	owly and d to blace by per- it in ash mordar	lour is brown d imperiectly ek by air, but sulphate Cr -Naptha- it dyestuff	13	Naphthazarıne S, alızarın black S, alızarın bluc black SW, naphthomelane
	5)	da	4	ur is brown, but red to black on it —Azine,	NoH Fib red on bleac solut An	2Sisevolved the becomes dish-brown boiling with hing powder ion 5°Tw— oxidation black	21	Aged-amline black, Prussiate black, one-bath amline black, steam amline black, diphenyl black
	Boil with dilute hydrochic acid (1	Boil for one minute with hydrochloric acid (1	precipitable by tannin is not obtained Redu  The colour is unaffected of re- + + + + +	The reduced colour is brown, but is rapidly restored to black on exposure to air — Azine, or the color of the	dolor but bow 50° 7 phic	o is evolved the becomes these or pale to n boiling h bleaching der solution Tw —Sul- de dyestuff	11	Blacks of the immedial, katigene, cross-dye, pyrogene, thiogene, thionol, pyrol, sulphur, &c , series
	dilute hv		table by	ocolour ored by vapours	ised and rair G on heat ndigoid	l colour 1e- lives violet ing in dry l dyestuff	10	Ciba grey G and B
Boil with weak ammonia (1	Boil with		tract precip	stored by air or persulphate — Azo class Boil with soap solution and white mercerised cotton	not stai 15 strip ing p Inso	ned Colour ped by boil- yridine — luble azo colour	6	Azophor black, &c
ıl with wea	ot stripped		A coloured acid extract	air or pers Boil wit thite merce	cotton is Test ash and Cu	struction and the control of the con		The undermentioned (col. 7) coppered or chromed
β	The colour is not stripped	pped (or on	, ,		-3 444		-1	Blacks of the diamine, oxydiamine, benzo, Columbia, diamil, Pluto, &c, series Also diaminogen, diazo blacks, &c (developed or coupled)
	The	The colour is not stripped (or only slightly)	ne colour is largely stripped in the acid extract is pre- trated by famin solution	asic dyestuff. Transfer to wool and boil with hydrosulphite AX	Colour just bef colouri or viole return Safra	Colour becomes red just before being decolourised. Violet or violet-blue colour returns in air—Safranine azo class.  Decolourised Colour returns on exposure to air—Azine, Oxazine, or Thiazine class.		Janus black, Janus grey, diazine grey.
		The	The colour is land the acid equipment	Basic dyestuf wool and bon sulphi	Deco Colour exposu Azine or T			Methylene grey, new methylene grey, new fast grey, nigrisine, &c
		tripped		Bolution	red C	r in ash	4	Logwood black on chrome mordant, noir reduit
		The colour is stripped	Sol	lution o	range	Fe in ash	60	Logwood black on iron mordant.
		The α	Fib	re and	solution	colourless	61	Tannate of iron
	The	dvest	inff tra	nsferred	to woo	dyestuff l and boiled manently ass.	1	Naphthol blacks, naphthylamine blacks, Palatine black, &c.

edition of Vogel's 'Practische Spectralanalyse irdischer Stoffe' (1889), charts were given of the absorption bands of various dyestuff solutions. This work was greatly extended in 'Formânek's Spectralanalytischer Nachweis Kunstlicher organischen Farbstoffe' (1900), which contains a systematic scheme for the spectroscopic identification of upward of 350 distinct dyestuffs In a later publication (1905) entitled 'Qualitative Spectralanalyse anorganischen und organischen Korper,' Formânek has extended his data to include the natural colouring matters, and, in conjunction with E Grandmougin, has republished his tables as 'Untersuchung und Nachweis organischen Farbstoffe auf spectroscopischen Wege'

The value of spectroscopic tests for the identification of dyestuffs on the fibre is limited to those dyestuffs which can be removed without

change by suitable solvents

Spectroscopic examination of dyed fibres has not yielded useful results W = M = G

Note.—Green's scheme of identification of dyes and the annexed Tables are printed by permission of Prof A G Green and Messis Chas. Griffin & Co., Ltd

DYER'S BROOM. Genista tinctoria (Linn) (Dyer's broom, Dyer's greenweed, Genet, Genestrole, Trentanel, Fr ; Ginster, Ger ) is found in the pastures, thickets, and waste places throughout Central and Southern Europe, across Russian Asia to the Baikal, and northward to Southern It is frequent in the greater part of England, but rare in Ireland and Scotland The fact that it contains a yellow colouring matter is recorded by numerous writers, and the following embody the principal references to the dycing and general properties of the plant Bancroft (Philosophy of Permanent Colours, 1813, 2, 108), Gmelin (Handbook of Chemistry, 16, 517); Berthollet (On Dyeing, 1824, 2, 242), Gonfreville (L'Ait de la Teinture des Laines, 501), Leuchs (Farben u Farbekunde, 1846, 2, 309), and Schutzenberger (Traité des Matières Colorantes, 1867, 4, 422)

To isolate the colouring matters, a hot aqueous extract of the plant is treated with lead acetate solution, and the pale yellow viscous precipitate is collected and decomposed by means of boiling dilute sulphuric acid. The clear liquid decanted from the lead sulphate deposits on cooling a dull yellow powder, this the solution poured into a large volume of ether, causing the separation of a dark-coloured resinous impurity. The clear liquid is evaporated, yielding a yellow crystalline residue, which consists of two substances. To separate is filtered off, dissolved in a little alcohol, and these, advantage is taken of the fact that, with sulphuric acid in the presence of acetic acid, one only of these compounds gives an insoluble This is collected and decomposed with water and the product crystallised from It is obtained as yellow needles, dilute alcohol and was found to be identical with the luteolin of weld (Resela luteola) (Perkin and Newbury, Chem Soc Trans 1899, 75, 830).

Genistein C<sub>14</sub>H<sub>10</sub>O<sub>5</sub>, the second colouring matter of dyer's broom, is present in the mother liquors obtained during the purification of the luteolin, and also in considerable quantity in the filtrate from the lead precipitate, from which

it is most readily isolated. To the boiling liquid ammonia is added, causing the separation of a lemon-yellow precipitate, which is collected and decomposed with boiling dilute sulphuric acid. The clear liquid is extracted with ether, and the extract evaporated, leaving a brownish crystalline mass. It is purified by crystallisation from acetic acid, and by conversion into the acetyl derivative.

Ğenistein crystallises in long colourless needles, mp 291°–293° (Perkin and Horsfall, Chem Soc Trans. 1900, 77, 1312), soluble in alkalıs with a pale yellow colouration. Alcoholic ferric chloride gives a dull-red violet colouration, and alcoholic basic lead acetate a lemon-yellow precipitate

Tracetylgenistein C<sub>14</sub>H<sub>7</sub>O<sub>5</sub>(C<sub>2</sub>H<sub>3</sub>O)<sub>3</sub>, colourless needles, m p 197°-201°, and tetrabromgenistein C<sub>14</sub>H<sub>6</sub>Br<sub>4</sub>O<sub>5</sub>, colourless needles, m.p. above 290°, have been described

On digestion with boiling 50 p c potassium hydroxide, genistein gives phloroglucinol and p-hydroxyphenylacetic acid

By methylation with methyl iodide in the usual manner, genistein dimethyl ether and methylgenistein dimethyl ether are produced

Genistein dimethyl ether  $C_{14}H_1O_3(OCH_3)$ , forms colourless leaflets, melts at  $137^{\circ}$ – $139^{\circ}$ , and gives the monacetyl compound

$$C_{14}H_7O_3(C_2H_3O)(OCH_3)_2$$

minute colourless needles, mp 202°-204°. When decomposed with alcoholic potash, it forms methoxyphenylacetic acid and phloroglucinol-monomethyl ether (identified by means of its disazobenzene derivative)

Methylgenistein dimethyl ether

melts at 202°, and the acetyl derivative

forms colourless needles, mp 212°-214°. With alcoholic potash it gives methoxyphenylactic acid and probably methylphloroglucinol-monomethyl ether

Genustein diethyl ether  $C_{14}H_8O_3(OC_2H_5)_2$  forms colourless needles, m p  $132^{\circ}-134^{\circ}$ , whereas acetylgenistein diethyl ether

$$C_{14}H_7O_3(C_2H_3O)(OC_2H_5)_2$$

melts at 168°-170°. Alcoholic potash gives p-ethoxyphenylacetic acid

According to Perkin and Horsfall, genistein may be a trihydroxyphenylketocumaran—

Genistein is a feeble colouring matter, and upon woollen cloth gives, with chromium mordant, a pale greenish-yellow, with aluminium mordant, a very pale yellow; and with iron mordant, a chocolate-brown shade.

Dyeing Properties of Dyer's Broom—In this respect there is a close resemblance between dyer's broom and weld. The dyeing power of the former is distinctly the weaker of the two; otherwise the only point of difference worthy of mention is that shown by the iron mordant, which, in the case of dyer's broom, gives a duller and more drab-coloured shade.

ECGONINE 611

Luteolin is also present in the Digitalis purpurea | Chem Soc 1918, 40, 598) (Urbain and Demeni (digito-flavone) (Fleischer and Fromm, Ber 1899, 32, 1184; v Kostanecki and Diller, ibid. 1901, 34, 3577), and in the flowers of Antirrhinum majus (Wheldale and Bassett, Bio-Chem J lc)

A G P DYMAL. Trade name for 'didymium' salicylate [Nd Pr] [C<sub>6</sub>H<sub>4</sub>(OH) CO<sub>2</sub>]<sub>3</sub>. A reddishwhite substance, used as a dusting powder, having non-irritant and antiseptic properties

DYNAMAGNITE, DYNAMITE v. EXPLOSIVES **DYSANALYTE.** A cubic or pseudo-cubic mineral consisting of titano-columbate of calcium with some ferrous oxide, cerium earths (up to 5 pc), and sodium The amount of Cb<sub>2</sub>O<sub>5</sub> given in various analyses ranges from nil to 26 p c, and the mineral has sometimes been asserted to be identical with perofskite (CaTiO<sub>3</sub>), the difficulties presented in the analysis of this mineral are, however, implied by its name, from δυσανάλυτος, 'hard to undo' It occurs as small black cubes in contact-metamorphic limestone at Sasbach in the Kaiserstuhl Mountains, Baden, and in metamorphic sandstone at Magnet Cove, Arkansas L. J S

**DYSCRASITE** (antimonial silver) Antımonide of silver Ag<sub>6</sub>Sb (Ag 84 3 p c) to Ag<sub>3</sub>Sb (Ag 72 9 pc), occurring as silver-white masses with metallic lustre, but often with a dull yellowish or black tainish The material is  $H = 3\frac{1}{2} - 4$ Orthorhombic crystals, twinned to simulate hexagonal forms, are rare and not distinctly developed. According to T. Liebisch (1910), two compounds are represented, namely, cubic AgoSb, with sp gr 10 05, and orthorhombic Ag<sub>3</sub>Sb, with sp gr 9 63-9 82 The mineral is found in silver mines at Wolfach in Baden (where it is the chief ore), Andreasberg m the Harz, Allemont in France, and the Cobalt district in Ontario. L J S Cobalt district in Ontario.

DYSODILE. A soft laminated carbonaceous substance of a yellowish-grey or green colour burning with a luminous flame, and a smell resembling that of burning caoutchouc in Sicily and in Bavaria (Fricklinger, Jahrb Min 1875, 760; also Church, Chem News, 34,

155, Potonić, Chem Zentr 1907, n 555). **DYSPROSIUM.** Sym Dy At wt 162 5 (162 52 Kremers, Hopkins, and Engle, J. Amer.

troux, Compt rend 1906, 143, 598, Hinrichs, *ibul* 1143) From erbia—one of the fractions obtained from ytterbite—Cleve separated an earth which he called 'holmia', and from this material Lecoq de Boisbaudran isolated a simpler constituent, dysprosia, characterised by a well-marked absorption spectrum showing bands in the ultra-violet as well as in the visible region of the spectrum (Compt. rend. 1886, 102, This oxide, one of the most basic constituents of holmia, was concentrated by precipitations with potassium sulphate and ammonia. Further fractionation of the nitrates and double nitrates gave a product rich in terbium and dysprosium (Urbain, Compt acnd 1904, 139, 736, 1905, 141, 521) The latter is separated from terbium by fractional crystalli-ation of the ethyl sulphates, terbium and yttrium being concentrated in the least soluble fraction, whilst dysprosium accumulates in the middle fraction (Urbain, Ann Chim Phys 1900, [7] 19, 184. Compt rend 1906, 142, 785)
Dysprosium material of very high purity

may be obtained by fractional crystallisation of the rare-earth bromates, followed by fractional crystallisation of the ethyl sulphates (Engle and Balke)

Dysprosium is separated from yttiium by crystallisation of the intrate with the addition of bismuth nitrate

Dysprosium forms a white oxide Dy,O3, but no perovide Its salts have a faint greenish-yellow tint chloride DyCl<sub>3</sub>.6H<sub>2</sub>O, nitrate Dy(NO<sub>3</sub>)<sub>3</sub>,511<sub>2</sub>O, melting at 88° in its water of crystallisation, very similar to bismuth nitrate (Urbain and Jantsch, Compt. rend. 1908, 146, 127), sulphate  $\operatorname{Dy}_2(\mathrm{SO}_4)_2, \mathrm{8H}_2\mathrm{O}$ , builtant yellowish crystals, stable at 110°, and completely dehydrated at 360°. For other salts, vJantsch and Ohl, Ber 1911, 44, 1274

Dysprosium stands first in the list of paramagnetic elements, its oxide Dy<sub>2</sub>O<sub>3</sub> being 12 8 times more magnetic than ferric oxide. high values for the magnetic susceptibility of the rare earths (yttria, erbia, &c) fractionated by Cleve and Nilson are probably due to the presence of dysprosium (Urbain, Compt. rend. 1908, 146, 922) G., T. M

 $\mathbf{E}$ 

EARTHNUT OIL v ARACHIS OIL EAU DE COLOGNE v PERFUMES

EAU DE JAVELLE. Solution of sodium hypochlorite (v Bleach liquor, ait Chlorine)

EAU DE LUCE. Aqua luciæ A milky liquid containing oil of amber, alcohol, and ammonia (v. Amber, Oil of).

EBONITE. Sulphurised caoutchouc RUBBER

EBULLIOSCOPE. An instrument for determining the strength of spirits of wine by means of its boiling-point ECBOLINE v. ERGOT.

N-benzoylhydroxypropylnovec-ECCAINE,

A compound allied to cocaine gonidene ester and of the structure

A crystalline base, more active than cocaine in its an esthetic properties, is non-toxic and can be readily sterilised (Braun and Muller, Ber 1918, 235, Chem Werke Grenzbach, D. R. P. 301139).

ECGONINE v COCAINE AND THE COCA ALKA-LOIDS.

ECHITAMINE, ECHITENINE v DITA BARK. ECHUJIIN v. DIGITALIS

E. C. POWDER v EXPLOSIVES

EDESTIN. A crystalline proteid substance found in the seeds of hemp, sunflower, and easter-bean, capable of uniting with acids to form salts (Osborne, J. Amer. Chem. Soc. 24, 39)

EDIBLE OILS AND FATS. Under this term are compused those oils and fats which serve as food. Most of the edible oils and fats, such as olive oil, cod liver oil, suct, laid, butter-fat, cacao butter, goose fat, &c, are immediately fit for consumption, if 'rendered' with sufficient care Hence they rarely need undergo a process of technical punication, except simple filtration, before they are placed on the market This applies, however, only to the freshly ('first') rendered products, as contact with putrescible organic matter rapidly causes hydrolysis of the oils and fats, so that, e g, lard or olive oil which has been allowed to remain in contact for some little time with the organic tissues from which they have been derived, readily become rancid, and therefore unfit for food in their original state. This applies with especial force to cod liver oil

The large demand made on the market by the rapidly growing consumption of oils and fats has, however, led to the employment of second and even third 'renderings,' in the case of animal fats, and second or third 'expressions' in the case of vegetable oils and fats. Such products contain, as a rule, notable amounts of free fatty acids, and small quantities of extractive substances, all of which must be removed before the oils and fats become suitable for food purposes. The processes applied to the purification and decolourisation are treated as deep secrets. It may, however, be stated that most of the processes are based on physical methods of refining (see OILS AND FATS)

Chemicals involving the use of mineral acids must be altogether excluded from the purview of the refiner, since the mineral acids, however carefully they may be removed by subsequent washing, impart an objectionable flavour to the products, thus rendering them useless for edible purposes Alkalis or alkaline earths are not so objectionable, and are, therefore, used commercially for the removal of fice fatty acids (A notable example of this kind is furnished by the usual method of refining cotton-seed oil) The alkalis and alkaline salts chiefly employed in practice are caustic soda, sodium carbonate, sodium silicate, lime, and magnesia Baryta has also been proposed (and patented), but its poisonous properties would prohibit its use.

Whereas all animal oils and fats, when freshly prepared, are suitable for food, this holds good only in a somewhat restricted sense of vegetable oils and fats. If the vegetable oils and fats are free from objectionable substances they may hold in solution or may have dissolved from the vegetable matter from which they have been obtained by expression or boiling out, they are suitable for edible purposes, and are so used to a very considerable extent.

We know, however, a number of oils which contain non-fatty toxic or injurious sub-tances, such as tung oil, calophyllum oil, cureas oil, croton oil. In the case of castor oil, the fatty acids themselves are injurious to health; hence

castor oil is unusable as a foodstuff. Experience has also shown that the fats belonging to the Chaulmoogra group are injurious to health All these oils must, therefore, be excluded from the list of edible oils

Oils which are regarded as palatable in one part of the world are not used as food in other parts. Thus, whereas rape oil and linseed oil ind a ready market for edible purposes in India, these oils are at present used in exceptional cases only as table oils in Western Europe. They are replaced in richer countries by cotton-seed oil, sesamé oil, arachis oil, poppy-seed oil, and olive oil. The increased consumption of the available edible oils and fats has created, during the last decades, a new and growing industry.

Amongst the most important developments in this industry is the preparation of coco-nut oil and palm-kernel oils for edible purposes (see Coco-nut oil and Palm-kernel oil). Mixtures of several edible oils are now largely prepared as 'table oils' to replace olive oil Amongst these substitutes the most important arachis oil, sesamé oil, cotton-seed oil, and to some extent also soya-bean oil. Mixtures of edible vegetable and animal oils a .d fats are chiefly represented by the 'suet substitutes,' which are prepared by mixing tallow with cotton-seed oil, and by 'lard substitutes,' which represent a mixture of animal fats with cottonseed oil, maize oil, &c Until a few years ago, 'compound lard' contained frequently no lard whatever, but, according to legislation in the United States, 'lard compounds' or 'lard substitutes' must now contain at least 50 pc. of lard (see LARD SUBSTITUTES)

Another development of the industry has been the introduction of hardened or hydrogenated oils, which are now extensively used as constituents of margarine (see Hydrogenated oils)

The manufacture and sale of edible oils and fats has assumed such considerable importance that legislation has stepped in to protect the public against fraudulent practices both as regards the commercial description of a 'blended' fat and the absence of injurious substances (see Margarine)

The changes in the character of fats during cooking have been described by Masters and Smith (Analyst, 1914, 39, 347).

J. L.

EDINOL. Trade name for p-amino-saligenin hydrochloride Used as a photographic developer

**ÉDULCORATION.** The act of washing insoluble matter so as to remove any soluble impurity.

**EFFLORESCENCE.** This term is applied to two or three different phenomena, characterised by the formation of a loose pulverulent deposit on the surface of a solid body

1 Crystallised salts are said to effloresce when they lose their water of crystallisation and fall to powder, e.g. normal sodium carbonate

fall to powder, eg normal sodium carbonate
2 The formation of a pulverulent crystalline deposit on the surface of a porous body, in consequence of the crystallisation of a salt existing in solution within its pores, or formed from materials existing therein, is also called efflorescence; e.g. the crystallisation of potassium nitrate on the surface of the earth or in caves, as

EGG. 613

in India, or on saltpetre plantations; of sodium sulphate and carbonate on walls; of alum on alum-slate; of ferrous sulphate on iron pyrites, &c In all these cases the solution of the salt is drawn by capillary action to the surface of the solid body, and there crystallises in consequence

of the evaporation of the water

3. The term 'efflorescence' is likewise applied to the effect which sometimes takes place when a saline solution is left to evaporate slowly in an open vessel The solid salt separates in dendritic crystals just where the surface of the liquid touches the side of the vessel, and the solution, rising by capillarity between these crystals, forms a fresh crystallisation just above the first layer; and this process is continually repeated till the crystallisation thus formed extends over the edge of the vessel, and sometimes down to the table on which it stands. Sal-ammoniac and other ammonia salts often effloresce in this The effect may be prevented to a certain extent by greasing the edge of the vessel, or better, by quickly boiling the solution, or evaporating it by heat from above EGG. The egg of the common fowl, Callus

EGG. The egg of the common fowl, Callus Bankva, consists of four principal parts the shell, the white, the yolk or vitellum, the white yolk or germinal cell, and the membranes separating these parts. The usual weight of an egg is 40-60 grams, of this the shell and shell membrane form 5-8 grams, the yolk, 12-18 grams: and the white, 23-34 grams.

The shell has the composition

The colouring matter of egg-shells is a bile

pigment (Liebermann, Ber 11, 606)

The shell membrane or membrana putaninis separating the white from the shell is double, the two parts being in contact over the greater part of the inner surface of the shell, but separated by an air-space at the broad end of the egg. These membranes are composed of a proteid substance allied to keratin (Liebermann)

Abderhalden and Strauss have examined the products of hydrolysis of ovokeratin, and find them to be, from 100 grams of substance 39 grams glycine, 35 grams alanine, 11 gram amino-valerie acid, 74 grams leucine, 40 grams proline, 81 grams glutamic acid, and 11 gram aspartic acid (Zeitsch physiol

Chem. 1906, 48, 530)

The white is a faintly yellow albuminous fluid enclosed in large cells of thin membrane. The fluid has sp. gr. 1045, and is really mobile, though it appears viscous because of its confinement by the cell walls. It has an alkaline reaction. Kojo (Zeitsch physiol Chem 1911, 75, 1) gives the following analysis:

Water		87 71 p c
Ash .		04,
Organic solids		11.89 ,,
Total nitrogen		1 75 ,,
Dextrose .		0 55 .,

The organic solids consist mainly of proteins, small amount of dextrose, and traces of fats, ecithin, and cholesterol Salkowski showed that glucosamine was also present (Biochem Zeitsch. 1911, 32, 335).

The protein substances of egg white are all gluco-proteids, and in their solution and precipitation properties are similar to globulins, albumins, or proteoses (v art PROTEINS)

Ovoglobulin separates from egg white partly on dilution with water, and completely if the solution be saturated with magnesium sulphate or ammonium sulphate By repeated solution in water and reprecipitation, part becomes insoluble (Langstein, Hofmeister's Beitrage, 1), and it is probably a mixture of several globulins (Corin and Beiard, Bull Acad Belg [3] 15, 643) The globulins form about 67 p c of the total proteids of egg white (Dillier, Hammarstein's Text-book Physiol Chem 430)

Ovalbumin is the chief proteid constituent of egg-white. It is obtained crystalline by precipitating the globulins with ammonium sulphate, as above, and allowing the filtrate to evaporate (For details of methods, see Hofmeister, Zeitsch. physiol Chem 16, 187, Osborne and Campbell, J. Amer. Chem. Soc. 1900, 22, 422, Rosenkrantz, J. Russ. Phys. Chem. Soc. 1906, 38, 2, and Mayer, Compt. rend. 1906,

143, 515)

Pure ovalbumin crystallises in large, well-formed needles A 2.5 p.c. solution in water becomes turbid at 60°, and at 64° yields a floculent coagulum. The addition of sodium of coagulation (Osborne and Campbell,  $l\,\epsilon$ , Starke, Pfluger's Archiv 12, 18)

The composition of ovalbumin is .

C H N S P () 52 75 7 10 15 51 1 62 0 12 22 90 p c

(Osborne and Campbell, lc 441, in agreement with Hopkins, J. Physiol 25, 306, and Hammarsten, Zeitsch physiol Chem 9, 304)

Many attempts have been made to assign a formula to ovalbumin Latham (Biochem J. 1908, 3, 207 et seq) gives it a molecular weight of 5473, and the formula  $\mathrm{C_{240}H_{397}N_{65}O_{76}S_3}$ , but this must be regarded as an approximation

It has been characterised by determining the various amino-acids yielded on hydrolysis with hydrochloric acid of sp gi 1 19 (v. art Proteins). 100 grams ovalbumin yield alanine, 21 grams, leucine, 200 grams, phenylalanine, 44 grams, tyrosine, 11 gram; cystine, 03 grams, proline, 23 grams; aspartic acid, 15 grams, and glutamic acid, 80 grams (Abderhalden and Piegl, Zeitsch physiol Chem. 1905, 46, 24, Osborne, Jones, and Leavenworth, Amer J Physiol 1909, 24, 252; Levene and Beatty, Biochem Zeitsch 1907, 4, 305)

The specific rotation of pure albumin is given as --31° (Willcock, J. Physiol. 1908, 37, 27), -30 70° (Hopkins, ibid. 25, 306), and

 $-29\,31^\circ$  (Osborne and Campbell,  $l\,c$  )

The precipitation of albumin by acids and salts is a completely reversible reaction (Galeotti, Zeitsch physiol Chem 1905, 44, 461), and follows the law of mass action (Ostwald, Chem Soc. Abstr 1908, 1 375, Gerrini, Zeitsch physiol Chem 1906, 47, 287)

Conalbumin Only about one-half of the proteid of egg white can be obtained as crystal-lised ovalbumin, the other fractions have a higher rotation and sulphur content By heating these fractions at 65°, conalbumin is

614 EGG

obtained as a coagulum, which, when dried at 110°, has the following composition:-

Its specific rotation is about  $-36^{\circ}$  (Osborne and Campbell, lc.; Panormoff, Chem Zentr 1898, 11, 487).

Ovomucoid is found in the residue after all proteids coagulated by heat have been removed from egg white (Neumeister, Zeitsch Biol 1890, 27, 369, Morner, Zeitsch physiol Chem 1893, It forms about 10 p c of the proteids 18, 525) of egg white It is not precipitated from solution by mineral acids or organic acids (except phosphotungstic and tannic acids) or by metallic salts.

The inorganic constituents of egg-white in 1000 parts are: Potash, 276-285 parts, soda, 235-239 parts, lime, 18-29 parts, magnesia, 16-32 parts; iron oxide, 4 4-5 5 parts, chlorine, 238-205 parts, phosphoric acid, 32-48 parts, sulphuric acid, 13-26 parts; silica, 3-20 parts, carbon dioxide, 97-116 parts (Poleck and Weber, Hammarsten's Physiol Chem 1904, 432). Traces of fluorine have also been found (Nickles, Compt rend 43, 885)

The yolk or yellow is contained within a membrane composed of a proteid allied to keratin (Liebermann, Pfluger's Archiv 43) It is a viscous, opaque, yellow, alkaline emulsion having a mild taste, and consists chiefly of water, fats, and colouring matter, together with vitellin, lecithin, cholesterol, &c Muller and Masuyama (Zeitsch f Biol 1900, 39, 542) have found a diastatic enzyme in the yolk, the presence of purme bases has been detected by Mesernitzki (Biochem Zentr 1, 739), and of

creatinin by Salkowski (lc).

1000 parts of the yolk contain approximately. water, 480 parts, fats, 228 4 parts proteid (vitellin), 1563 parts, lecithin, 1072 parts, cholesterol, 175 parts, insoluble salts,

6.12 parts, soluble salts, 3.53 parts

According to Poleck, the ash of yolk of eggs contains, in 1000 parts soda, 51 2-65 7 parts potash, 80 5-89 3 parts, lime, 122 1-132 1 parts, magnesia, 20 7-21 1 parts; iron oxide, 11 9-14 5 parts, phosphoric acid, 638 1-667 0 parts, silicic acid, 5 5-14 0 parts

The fat forms about 30 p c of the yolk, and may be extracted with ether. It is a mixture of solid and liquid fat, and contains 06 pc phosphorus, equivalent to 1504 pc. lecithin, 61 5 p c of neutral fat, and 23 2 p c free acids The neutral fat contains glycerol and acid in such proportions as prove it to consist of triglycerides only (Thorpe, Arch Nedér Sci Exact Nat 1901, 6, 155) On saponification On saponification it yields 40 pc oleic acid; 38 04 pc palmitte acid; and 15 21 p.c stearic acid (Paladino, Biochem Zeitsch 1909, 17, 356)

Barbieri, by extracting yolks with carbon disulphide, obtained tristearin and triolein (Compt rend 1907, 145, 133), besides cholesterol; while MacLean has isolated pure tripalmitin from egg-fat (Biochem. J 1909, 4, 168).

Lecithin (v Lecithin) is present both free and in combination with protein (Manasse, Biochem Zeitsch. 1906, 1, 246) Commercial Commercial lecithms are usually prepared from eggs, a d show a high choline content. Trier identified

B-aminoethyl alcohol in egg lecithin, isolating it as its gold chloride compound (Zeitsch physiol. Chem. 76, 496) Lecithin is best prepared from dried egg yolks by Bergell's method (Ber. 1900, 2584) The extract prepared by boiling with 96 p.c alcohol for 6 hours, is cooled to 0° and precipitated with an alcoholic solution of cadmium chloride The precipitate is filtered off, washed with 96 p c alcohol, dried in air, and extracted with ether The residue is boiled with eight times its volume of 80 p.c. alcohol, and a concentrated solution of ammonium carbonate is added until the solution is free from cadmium. The filtered solution is cooled to -10°, and the precipitate is dissolved in chloroform, precipitated with acetone, and dried in vacuô over sulphuric acid.

After hydrogenation of egg lecithin, Paul and Ochme isolated stearic and palmitic acids, and suggest that linolic and palmitic acids are originally present (Ber. 1913, 1297)

Barbieri (Gazz chim ital 1917, [1] 1) stands alone among a host of investigators in his failure to isolate lecithin from egg yolk affirms that the phosphorus of egg yolk is present as morganic phosphates, that the choline exists only as a degradation product of ovochromin, and that glycerol and fatty acids exist as constituents of fat, and not combined as lecithin

Cholesterol is present in the yolk to the extent of about 1 pc (Juckenack, Zeitsch. Nahr Genussm 1899, 1905; Benecke, Bied Zentr 1881, 568) According to Scrano and Palozzi (Chem Zentr 1911, 11 772), it occurs as its oleic and palmitic esters

Lutein, the colouring matter, was described by Maly (Chem Zenti 1881, 485) as a mixture two substances: vitello-lutein, which is yellow and soluble in petroleum, and vitello-rubrin, a red body soluble in carbon disulphide. The mixture is an orange-red amorphous substance, which is not affected by alkalis, and can therefore be isolated by hydrolysing the fats. C A. Schunk (Proc Roy Soc. 1904, 170), as the result of spectroscopic examination, identified the yellow pigment of the hen's yolk with the L xanthophyll, which he has isolated from The characteristic spectroscopic haviour of the lutein has been suggested as a test of the purity of commercial dried egg yolk (Lewin, Miethe, and Stenger, Pfluger's Archiv Willstatter and Escher (Zeitsch physiol Chem 76, 214) isolated the pigment from the yolks of 6000 eggs, and obtained it in crystalline form, and showed it belonged to the xanthophyll group of pigments; they showed also that a carotin-like pigment was present Palmer (J Biol Chem 23, 261) showed that the pigments of egg yolk were increased by feeding with rations rich in xanthophyll, yellow corn being particularly suitable for this purpose as a winter ration. The point is of importance to the poultry trade, since eggs with highly-coloured yolks are in great request.

Vitellin, the protein constituent of the yolk, is separated by mixing the yolks with brine, in which the protein is soluble, and extracting the fat, &c, with ether. It has the properties of a globulin, and, on dialysis or dilution of the solution, is precipitated as a white granular substance. A solution coagulates at 70°-75°. EGG. 615

Vitellin thus obtained is a mixture of compounds of protein matter with lecithin, containing from 15-30 p.c of the latter, called lecithin-nucleovitellins. The lecithin thus combined is not removed by ether, but is completely removed by alcohol, leaving an insoluble proteid, nucleovitellin, of constant composition, which, on digesting with pepsin, yields paranuclein.

It is probable that all these proteins are

compounds of vitellin with phosphoric acid H<sub>3</sub>PO<sub>4</sub>, or some very simple organo-phosphoric acid. The composition of vitellin, free from

phosphoric acid, is:

(Osborne and Campbell, J Amer Chem Soc 1900, 22, 413)

Plimmer states that another distinct protein 'livetin,' can be obtained from the yolk (Chem Soc Trans 1908, 1500)

Berthelot and André determined the heat of combustion of egg yolk to be 8124 calories per gram (('ompt rend 110, 925)

The germinal vesicle, or white yolk, forms a laver between the yolk and the yolk membrane, with a process extending to the centre of the It is enclosed in a membrane, and contains proteins, lecithin, and a nuclein rich in phosphoius (Meischer Chem Soc Trans 1871,

The space between the shell membranes at the broad end of the egg contains a mixture of oxygen, nitrogen, and carbon dioxide, richei in oxygen than the surrounding air (Hufner, Du Bois Reymond's Archiv de Physiol 1892, Phys Abth 54)

Eggs contain, on the average, about 4.6 mgms of ferric oxide This amount is increased by feeding the hens with foods rich in iron, and 'iron-eggs' thus obtained have been recommended for therapeutic use, but are very costly (Hartung, Zeitsch f Biol 1902, 43, 195, Hofmann, Zeitsch, anal Chem 1901, 40, 450)

Bertrand states that arsenic is normally present in the egg to the extent of  $\frac{1}{200}$  mgm (Compt rend. 1903, 136, 1083), and that 1 mgm boron occurs in 1 kgm dried egg (Compt rend

1913, 156, 2027)

Development —During incubation the egg respires, giving off carbon dioxide and water, and absorbing oxygen If this process be stopped by placing the egg in an ineit gas, or by rendering the shell impervious to gases, the development of the chick ceases The respiration is slight for the first four days of incubation, and then increases until, at the ninth day, the respiration of the egg is equal to that of a fowl, weight for weight. This respiration involves combustion of the fat of the egg, and a steady loss in weight (Pott, Landw Versuchs Stat 23, 203, Chem Soc Abstr. 1879, 474)

Levene states that the quantity of xanthine bases and nuclein compounds increases with the growth of the embryo (Amer J Physiol 1900,

3, x11)

In egg-white the ratio albumin: ovomucoid of egg is constant during incubation (Bywaters and Roue, J. Physiol 46, Proc 33) After 3 days α-lactic acid was found (Anno, Zeitsch. physiol. Chem. 80, 227) In new-laid eggs t'e cholesterol is free, but gradual esterification takes place, until at the time of hatching 40 p c. is in ester form (J. Biol Chem 21, 23)

All the phosphorus in the chick is derived from the yolk and white, and, until the last few days of incubation, no lime is taken from the shell (Voit, Zeitsch f Biol 13, 518, Pott, lc) At the end of incubation, the shell may supply some of the lime required (Carpiaux, Bull Acad Roy Belg 1908, 283, see also Plimmer and Scott, J. Physiol 1909, 38, 247)

The freezing-point of fresh eggs is 0 427°-0 480°, and rises slightly as incubation proceeds, corresponding to a rise of osmotic pressure from 55 atmospheres to 73 atmospheres (Atkins, Biochem J 1909, 480)

The chief uses of eggs are as an article of food and in the tanning industry As the supply of new-laid eggs varies with the season, recourse is had to some means of preserving a proportion of those that come into the market at the time

of greatest production

The following is a list of some processes for preserving whole eggs, which have been patented Saturating with carbon dioxide (Lake, Eng Pat 1653, 1888), treating with formaldehyde vapour (Loft, Eng. Pat 19721, 1915, Corden, Eng Pat 15206, 1914), wrapping air-tight in rubber-coated paper (Jessen, Eng Pat 15580, 1891), coating with gclatin (Mills, Eng. Pat 17717, 1891; Farquhar and North, Eng. Pat 15128, 1892, Coakley, Eng. Pat. 18816, 1899), with gelatinous starch solution (Schultz, Eng Pat 1328, 1902); with drying oils, wax, or varnish, with or without previous treatment with antiseptics (Stukes, U.S. Pat. 649899, Barlow, Eng. Pat. 11054, 1902, Lorne, Eng. Pat. 18439, 1901, and 9898, 1901; Davis and Metz, Eng Pat 105840, 1916), with boric acid (Markham, Eng Pat 3513, 1895), with lime (McArdle, Eng Pat 3592, 1892; Wilson and Wilson, Eng Pat 15461, 1899), with sulphuric acid (Reinhardt, Eng Pat 18130, 1898; Marks, Eng Pat 12867, 1903), with liquid petroleum (Henderson, Eng Pat 101337, 1916), and with water-glass (Stead, Eng Pat. 4910, 1882, Rylander, Eng Pat 23523, 1901)

The following methods (J Soc Arts 1898, 46, 390, J Soc Chem Ind 1898, 261) were found to keep over 90 p c of the eggs in good condition for 8 months. (1) varnishing with vaseline, (2) keeping in milk of lime, and (3) keeping in water-glass solution. Of these the last method is the one least open to objection, and is that chiefly used Eggs preserved in water-glass have a tendency to burst when boiled, but this can be prevented by piercing

the shell with a needle.

A Chinese preparation, 'Pidan,' is made from duck's eggs by coating them with a paste obtained by stirring into an infusion of 13 lbs. strong black tea, 9 lbs lime, 4½ lbs salt, and a bushel of wood ashes After 5 months they are The changes coated further with rice hulls which take place were investigated by Blunt and Wang (J Biol. Chem 28, 125), who deduced that decomposition of both the egg proteins and lipins had taken place. The values for ash, for total and organic phosphorus and especially for ammoniacal nitrogen, are high

The contents of eggs may be preserved by hating at 185°F with a strong solution of a sugar (glucose, maltose, dextrose, &c), until 616 EGG

readily miscible with water, milk, &c, and will keep for some months without deterioration (Fordred, Eng Pat. 15762, 1894, Goodlett, US Pat 692268)

An important method of preserving the whites and yolks of eggs consists in drying the mixture in a current of air at 37°-40° during 18 hours, powdering the product, and again exposing to dry an. The powder is packed in tins, or can be pressed into tablets, and is quite soluble in water (Foschi, Fr Pat 369967, 1906) A similar product is obtained by drying on a hot roller at a temperature above 212°F. (Hatmaker, Eng Pat 28, 1905, see also Eng Pat 8743, 1903, for apparatus)

Substitutes for eggs in baking are made from albumin and cascin obtained from skimmed milk, sometimes mixed with a proportion of dried egg, or from maize meal containing sodium bicarbonate and artificial colouring matter (Beythien and Waters, Zeitsch Nahi Genussm 1906, 11, 272, Bernstein, Eng Pat 14336 and 29189 of 1897) An emulsion of sunflower or other oil with blood serum and lecithin has been suggested (Eng. Pat. 100135, 1915)

The eggs used for the treatment of leather, especially the finer sorts of glove leather, are preserved in the liquid state by the addition of salt and borax (Jean, Mon Sci 1892, 6, 561) Analyses of this material, 'eggio,' show that it contains 2-5 pc more water and 31-6 pc less fatty acid than firsh egg yolk (Boyer, J Soc. Chem Ind 1892, 447) Bein has pointed Bein has pointed out that the addition of salt to yolks materially reduces the amount of fat (Bied Zentr 21, 138) Rotten eggs could well be used for this puipose, as Cech has found that the available oil is not much diminished in amount, and can be extracted by pressure (J pr Chem [2] 22, 338)

Oil of egg, obtained by extracting yolks with ether, or, better, chloroform, forms about 10 p c of the egg, and is used in pharmacy in the preparation of ointments and emulsions, in the leather industry and in tempera painting (Lupus, D. R P 187211) It has no medicinal value (Paladino and Toso, Analyst, 1896, 21, 161) It has the following properties mp 22°-22 5°, sp gr. at 15°, 0 9144, and number, 12, saponification number, 190 2; ether number, 189 0, indine number, 72 1; Hehner's number, 95 16, Reichert-Meissla Chem Zeit 1897, 21, 303). Reichert-Meissl number, 04 (Kitt,

Detection, estimation, and analysis —The presence or absence of egg yolk in pastry can be determined by examination of the fat, there being a great difference in composition and properties between the fat of wheat and of the egg (Spaeth, Zeitsch Nahr Genussm 1896, 10, 171) The phosphorus of the yolk being present chiefly as lecithin, the yolk may be estimated by igniting with sodium carbonate and nitre, determining the phosphoric acid in the residue 1·129 grams phosphoric acid represent 100 grams yolk (Bein, Ber 23, 423) The addition of alkalı is necessary to prevent loss of phosphoric acid during ignition (Juckenack, 7.c) The amount of egg present in pastry, &c, may be determined by the amount of cholesterol present (Cappenberg, Chem Zeit 1909, 33, 985) methods for the determination of the con-

the product contains 40-60 p.c of sugar. It is stituents of egg yolk for commercial purposes, see J Soc (hem Ind 1906, 829)
EGG ALBUMIN v PROTEINS

EGG ALBUMIN v PROTEINS 2 EGG FRUIT. The fruit of Solanum melongena (Linn) A plant of tropical and sub-tropical countries Konig gives, as the mean of three analyses--

Nitrogenous Soluble carbo-matter Fat hydrates Fibre 1 1 0 1 3 9 1 2 Water 93 2

An American analysis (Bull 28, US Dept of Agric ) shows a somewhat larger amount of carbohydrates (5 1 pc), but is similar in other respects

This plant is sometimes called Aubergine. The fruit is used in much the same way as the tomato

EGOLS. A class of antiseptics consisting of the potassium mercury salts of o-mitrophenolsulphonic acid, the individual members being termed 'phenegol,' 'cresegol,' &c, depending on the phenol from which they are derived They are reddish-brown powders, readily soluble in water, and are strongly bactericidal (Gautrelet, Compt rend 129, [2] 113, J Soc

Chem Ind 1899, 854) (v SYNTHETIC DRUGS)
EGYPTIAN BLUE. This is a blue colouring matter which was used by the Romans in the first few centuries of the Christian era, but ceased to be manufactured about the time of the invasion of the barbarians It figures in several beautiful frescoes in the Vatican, and has also been found at Pompen Fouqué discovered the composition to be that of a double silicate of calcium and copper CaO,CuO,4SiO<sub>2</sub>, devoid even of traces of alkali The substance is crystalline, and is hardly attacked by any chemical reagents, which explains the perfect state of its preservation in the paintings for which it was employed 1900 years ago. It may be prepared at a bright-red heat, but this temperature must not be exceeded, as it is then decomposed into cuprous oxide, wollastonite, and a clear light-green glass The higher the temperature, the more of the latter is formed, and, finally, wollastonite disappears entirely By the old method of preparation, described by Vitiuvius, fine sand and soda are intimately mixed with copper filings, then moistened with water, formed into cakes, and the latter dried. The cakes are then fused in an earthen pot until the blue colour is produced Fouqué found this method practicable, but used potassium sulphate in preference to soda as a flux It was not possible to replace chalk or lime by magnesia The ancient Romans employed a large excess of silica, whilst Fouqué employed more basic mixtures, finally purifying the product with hydrochloric acid exquisite colour of this mineral body, the fact that it is absolutely unaffected by moisture, light, and most chemical reagents, as well as the facility and the low price of production, render it desirable that its manufacture should be again resumed (F Fouqué, Compt rend 108, 325-327; and J Soc Chem Ind 8, 291)
The conditions of its production have been studied by Laurie, McLintock, and Miles (Proc. Roy Soc 1914, A, 89, 418) The French ultramarine works of Deschamps Frères reproduced Egyptian blue by fusing in an ultramarine furnace a mixture of quartz (50 parts),

chalk (21 parts), copper oxide (24 4 parts), and sodium carbonate (4 6 parts), all in the finest powder, and completely free from iron. The product, washed with hydrochloric acid and hot water, had the composition corresponding to Fouqué's analysis, and was in all respects similar to the material used by the Romans (Bock, Zeitsch angew Chem 1916, 29, 1 228) EIKONOGEN. Sodium salt of 1-anino-

8-naphthol-6-sulphonic acid Used as a photo-

graphic developer

EKSANTALAL (12H18O Obtained on oxidation by ozone of the sesquiterpene primary alcohol a-santalol (15H24O, which, together with the isomeric \(\beta\)-santalol, constitutes 'santalol,' the oil resulting from the hydrolysis of the hydrogen phthalate made from the oil obtained by distilling sandalwood with steam (Semmler, Ber 43, 1893)

It is dextro-rotatory ( $\alpha_D + 13.5^{\circ}$ ), bp 100°-110°, under 10 mm; spgr 0 9846 at 20°/20, n<sub>D</sub>, 1 48519.

It gives an oxime, which yields the corresponding nitrile and aminc, and gives eksantalol on reduction (Semmler, Ber 41, 1488)

Heated with acetic anhydride and sodium acctate, it forms enol-eksantalal acctate

which, on oxidation with potassium permanganate, yields noreksantahe acid C<sub>11</sub>H<sub>16</sub>O<sub>2</sub> (Semmler, Ber 43, 1722–1725), the methyl ester of which, on reduction by sodium and alcohol, yields noreksantalol<br/>  $\mathrm{C}_{11}\mathrm{H}_{18}\mathrm{O}\,,\,$  and this on oxidation in benzene solution with potassium dichromate and dilute sulphuric acid, gives noreksantalal C  $_{11}\rm{H}_{16}\rm{O},~b~p~92^{\circ}-94^{\circ}/11~mm~,~sp~gr~20^{\circ}/20^{\circ}~0~9964,~n_{_D}~1~48301,~\alpha_{_D}~-30~8^{\circ}$ The enol-noreksantalal acetate gives, on oxidation by potassium permanganate, teresantalic acid  $C_{10}H_{14}O_2$  (Semmler and Zaar, Ber 43,

In all three series, containing 10, 11, and 12 carbon atoms respectively, the same tricyclic system is present as exists in  $\alpha$ -santalol, and Semmler (Ber 43, 1893–1898) assigns to these bodies constitutions based upon the camphene or camphor type, and a santalol and eksantalal are structurally represented as below

$$\begin{array}{c|ccccc} H_2C & CH_2 & CH_2 \\ & \downarrow & CH_2 & \downarrow \\ & \downarrow & CH_2 & \downarrow \\ & \downarrow & CH_3 & CH \\ \hline CH_3 & CH & CH_2 \\ \hline CH_4 & CH_2 & CH_2 \\ \hline \rightarrow & & \downarrow & CH_2 \\ \hline \end{array}$$

EKTOGEN. Trade name for a mixture of zinc dioxide with zinc oxide and carbonate, used as a surgical dressing

ELÆOMARGARIC ACID C18 H32O2 An acid, melting at 48°, obtained by the saponifics tion of elaococoa oil (Japanese wood oil) with

alcoholic potash, and decomposing the trielaomargarin thus formed The glyceride when first formed is liquid, but on exposure to light gradually solidities. Isomeric with elaostearic acid, elaoleic acid, and linoleic acid. On exposing an alcoholic solution to light, plates of claostearic acid, m.p. 71°, are precipitated (Cloez, Bull Soc chim 1876, 26, 286; 1877, 28, 24); traces of iodine or sulphur also bring about this change (Maquenne, Compt. rend. 135, 696) Oxidation with potassium permanganate gives rise to azelaic acid, sativic acid, and dihydroxy-stearic acid (Kametaka, J Soc Chem Ind 1903, 1138, 1908, 1120, Kit, Chem Soc Abstr 1905, 10. Maquenne, lc) Linolic acid also on oxidation yields sativic acid, and on bromination, linolic acid and elaomargatic acid yield the same tetrabromide Elaostearic acid and linoleic acid yield the same diozomde (Majima, Ber 1900, 674) Elaomargane acid and elaosteame acid are converted into claoleic acid when heated at 180° m sealed tubes (Cloez, J. 1878, 38)

ELÆOPTENE. The volatile portion of an essential oil, used in contradistinction to the stearoptene or solid portion

ELAIDIC ACID C18H34O2 A solid stereoisomene modification of oleic acid, obtained by the action of nitrous acid upon olive oil, saponifying the glyceride (claidin) so formed, and decomposing the salt by hydrochloric or sulphuric acid May also be obtained by the action of nitrous acid or of intile acid (sp gr 12) (Edmed, Chem Soc Proc 1899, 190) upon oleic acid, or by decomposing barium oleate with furning nitric acid, m p  $52^{\circ}$  (Saytzeff, Chem Soc Abst 1893, 551), b p  $287.5^{\circ}-288^{\circ}$  at 100 mm (Krafft and Mordlinger, Ber 1889, 819) Soluble in alcohol and other Solutions have an acid reaction. The melted acid gradually absorbs oxygen from the air, and becomes permanently liquid. It may be distilled unchanged under diminished Heated with potassium hydroxide, it is converted into acetate and palmitate of potassium with evolution of hydrogen

Elaidic acid dissolved in benzene, toluene, ligioin, or chloroform, and mixed with caustic soda and shaken, gives a brilliant phosphorescence (Radziszewski, Annalen, 203, 305) Hydriodic acid forms an addition compound (nodostearic acid) with claidic acid (Farb vorm Fried Bayer & Co D R P 180087, Chem Soc Abstr 1907, 1 380), which, on treatment with alcoholic potash, yields oleic and 180-oleic acids (Lebedeff, ibid 1893, 550) On reduction with nickel or copper and hydrogen at 280°-300° elaidic acid is converted into stearic acid (Sabatier and Mailhe, Ann Chim Phys [8] 4, 319) Treated with dilute solution of potassium permanganate in presence of potash, it is converted into a dihydroxy acid, which differs from the corresponding compound obtained from oleic acid by its lower melting-point (995°), and its greater solubility in alcohol and other, and also into pelargonic, azelaic, and oxalic acids (Saytzeff, Ber 1886, 21, Edmed, Chem. Soc. Trans 1898, 627), oxidation with sulphuric acid, however, gives rise to the same dihydioxystearic acid as is obtained from oleic acid under the same conditions (Tscherbakoff and Saytzeff, pr Chem, 1898, 27).

Eluidin, a solid isomeride of clein, is obtained by the action of nitrous acid upon clein,

and is probably a triglyceride C57H104O6 melts at 36°; is nearly insoluble in alcohol. readily soluble in ether The Unquentum oxugenatum and U citrinum of the pharmacopœias, prepared by the action of nitric acid upon hog's lard, contain impure claidin.

ELAIDIN V ELAIDIC ACID

ELALDEHYDE v ALDEHYDE.

**ELARSON.** Trade name for strontium chloroarsenobehenolate

ELATERIN v ELATERIUM

ELATERITE. Elastic bitumen, Mineral caoutchouc A fossil resin, somewhat resembling asphalt, found at Castleton in Derbyshire, in France, Switzerland, &c It is a soft, flexible, elastic, and readily inflammable substance, with a resmous lustre and a dark-brown colour is mainly used as an insulator for electric cables It is rapidly decomposed by strong nitric acid When boiled with alcohol and alcoholic potassium hydroxide it behaves like beesway One sample gave an acid value of 49, and saponification value of 569, and contained 889 pc of unsaponifiable matter.

The sediment from the juice ELATERIUM. of the fruit of the Echallium elaterium (Richard), or squirting cucumber Elaterium is a drastic

purgative

It, and elaterm derived from it, were excluded from the BP in 1914 The latter substance is still official in the United States (Eluterinum, USP) Good specimens of elaterium when extracted with chloroform yield 25 pe of elaterin, which by recrystallisation is obtained in minute shining six-sided plates The dose of elaterin is  $\frac{1}{4}$  to  $\frac{1}{10}$  of a grain, and to it is ascribed the whole of the physiological activity Elaterin is a neutral bitter comof claterium pound, insoluble in hot or cold water leadily soluble in cold alcohol, sparingly in ether. If sulphuric acid be added to a solution of elaterin in melted phenol, a crimson colour is produced which changes to scarlet (Lindo, Chem News, 37, 35, Johannson, Zeitsch anal Chem 24, 154)

On hydrolysis with alcoholic potash, elaterin furnishes acetic acid and an amorphous compound, elateridin Further action of alkali gives clateric acid, also amorphous, and melting at 73°-75° (Berg, Bull Soc chim [m ] 35, 435)

Elaterium of English origin, examined by Power and Moore (Pharm J. [iv | 29, 501), was found to contain 5 3 p c of moisture and 6 7 p c of ash. By a suitable method of extraction, a colourless crystalline product, mp 217°-220°, was obtained, corresponding with the claterin of the pharmacopenas This was found to be non-homogeneous, consisting of at least two substances, one, a-elaterin, melting at 230° with decomposition, and lavo-rotatory, whilst the other, B-elaterin, has a lower melting-point, and is dextro-rotatory Both have the same empirical composition, but the former is physiologically mert, whilst the latter shows marked physiological activity. Examination of commercial elaterin showed that it consists mainly of varying proportions of these two substances, which probably accounts for the marked difference in medicinal value of different samples a-Elaterin has also been found in colocynth, the dried fruit of Citrullus colocynthis (Schraeder) (Power and Moore, Chem. Soc. Trans 97, 99), and has been further investigated by Moore (ibid. 97,

According to Berg (Bull. Soc. chim [ni] 17, 85; [iv ] 7, 385), elaterin does not exist readyformed in the fruit, but is produced after the rupture of the tissues by the action of an enzyme elaterase on a glucoside present. The latter consists of a brittle, amorphous powder, and gives the same leactions as elaterin with sulphuric acid and phenol Power and Moore (Chem Soc Trans 95, 1985), however, on examination of the fresh fruit, failed to confirm this

**ELBON.** Trade name for cinnamoyl-phydroxyphenylurea.

ELDERBERRY. The fruit of Sambucus nigra (Linn) The only analysis quoted by Konig gives, as the composition of the fruit-

> Water 81 9 Crude fibre Pentosans 66

The bernes are used for the preparation of a wine and of 'clderberry syrup,' an old-fashioned From the wine, a brandy is sometimes distilled, which according to analyses by Arnthor and Zink, quoted by Konig, contains 39 2 p c of alcohol (by weight) and has a sp gr of 0 836 According to de Sanctis (Gazz chim ital 1895, 25, 49), elder leaves and stems contain conine Malmejac (J Pharm Chim 1901, 14, 17) attributes the dimetic and purgative action of the bark of the elder to an alkaloid, sambucine, a deliquescent bitter substance, yielding the usual alkaloid reactions He also found elder bark to contain tannin, a resin resembling scammony and a yellowish-red oil possessing a strong odour of elder Elder pith contains both xylan and araban (Browne and Tollens, Ber 1902, 35, 1457) The oil extractable from the juice of red elderberries (S racemosa) by ether, is light yellow in colour, has a sp gr of 0.9072, solidifies at  $-8^{\circ}$ , and melts at  $0^{\circ}$ , contains 6 65 pc of free acid (as oleic acid), 0 66 p c of unsaponifiable matter, has a saponification number 2093, iodine number 814, and Reichert-Meissl value of 154 (Byers Hopkins, J Amer Chem Soc 1902, 24, 771)

The leaves of the elder contain a cyanogenetic glucoside, which, under the influence of emulsin, yields dextrose, benzaldehyde, and hydrogen cyanide; the yield of the last mentioned amounts to 0 126 gram per kilogram of fresh leaves (Guignard, Compt rend 1905, 141, 16; Bourquelot and Danjou, ibid 59) This glucoside, which has been named sambunigrin C14H17O6N, crystallises from ethyl acetate in long, colourless needles, melting at 151°-152°, is readily soluble in water or alcohol, but nearly insoluble in ether (Bourquelot and Danjou, Compt rend.

1905, 141, 598)

According to Ravenna and Tonegutti (Chem Zentr 1910, 1 544) the enzyme which decomposes the glucoside in the plant is not soluble in water The hydrocyanic acid is present in larger quantities than has hitherto been believed, and is most abundant in the stalks

ELECTRARGOL. Trade name for a form of colloidal silver.

ELECTRIRIDOL. Colloidal iridium.

ELECTRO-DEPOSITION and ELECTRO-I LATING. Electrolytic processes are now used to a considerable extent in the precipitation of metals in a reguline form, available for many commercial purposes. Thus the common and more easily corroded metals are coated with thin films of electro-deposited metal of a more permanent and less oxidisable character, in the process of electroplating; for the reproduction of engraved surfaces, type, &c, in electro-deposited copper, electrolysis is applied in electrotyping, and in the deposition of a pure metal from an impure form or a mixture of metals, the principles of electrolysis are applied in the process of electrolytic refining and in chemical analysis

An electric current passed through solutions of metallic salts brings about decomposition of a particular type, accompanied by the deposition or evolution of the products, or others which may have been formed as the result of secondary changes. When copper sulphate solution is electrolysed with a platinum anode, and any cathode, not ordinarily attacked by this salt, copper is deposited at the cathode, and oxygen and sulphune acid are formed at the anode, thus

Other metallic compounds follow a similar The primary products consist of charged atoms or radicals These are termed ions, and in the process of electrolysis migrate to the electrodes, those to the anode being called the anions, and those to the cathode cathions Anions consist of the acidic group of the salt, while the cathions comprise hydrogen, the metals, and basic radicals. At the electrodes, the electrical charges are neutralised, and the substance appears in the nascent form, subsequently being deposited or evolved Elementary amons, such as chlorine, bromine, and iodine may enter into subsequent secondary changes, while others, such as  $NO_3$ ,  $SO_4$ ,  $C_2H_3O_2$ , &c, react with water, giving generally the corresponding acid, and liberating oxygen With soluble anodes—those attacked by the anion—solution of the metal takes place, and thus, in a copper solution, a copper anode supplies metal under the best conditions as rapidly as metal is de-The strength of the solution is thus maintained, a condition necessary for continuous deposition as required for plating and refining

purposes
The deposition of metals follows the usual law of chemical equivalents as enunciated by Faraday in 1835. The ampere-hour is thus associated with a definite amount of metal, which cannot be exceeded, but may not be attained owing to failure in realising the best conditions. The quantities of metals deposited by the coulomb, ampere-hour, and 746 ampere-hours, are shown in the table on next column

These amounts, however, can only be deposited when the cathode is surrounded by a considerable amount of strong electrolyte, and only under this condition can the facts of electrolysis be applied, as in the voltameter, to the determination of the number of coulombs passing through a circuit While for most industrial purposes it is not necessary for the theoretical amount of metal to !e deposited per amperchour, yet the most satisfactory deposits are,

Metal	Chemical	Mgs per coulomb	Grams per ampere-hour	Oza (av ) per 746 ampere-hours
Copper (cupic) ,, (cupious) Gold (aurie) ,, (aurous) Iron Lead Nickel Silver Tin (stannous) ,, (stannie) Zine	31 8 63 6 65 7 197 0 28 0 103 5 29 3 108 0 59 0 29 5 32 5	0 329 0 658 0 680 2 040 0 290 1 070 0 302 1 118 0 610 0 305 0 336	1 182 2 364 2 440 7 320 1 040 3 870 1 085 4 025 2 190 1 095 1 210	31 0 62 0 58 6(Troy) 176 0 ,, 27 3 101 0 28 5 96 5(Troy) 57 5 28 7 31 8

in many cases, obtained when this condition is most nearly attained Failing the full amount of metal per ampere-hour, hydrogen is deposited, and this often leads to a dark powdery deposit unsuitable for most industrial purposes. Such deposits, caused usually by excessive current, are said to be 'burnt'. In some solutions—for example, those of copper, iron, or nickel sulphate this inferior deposit seems to result from the simultaneously deposited hydrogen due to a weak electrolyte In other examples, that of the alkaline or eyanide copper bath, it would seem (from the fact that much hydrogen may be deposited without 'burning') more probably due to the deposition of traces of basic compounds with the metal (Bancroft, Trans Amer Electrochem Soc vi [ii] 27) This view is supported by the fact that the presence of substances in the solution, which are good solvents for such basic compounds, prevents burning to a considerable extent. Thus, in the alkaline copper-bath, 'burning' is reduced, and the quality of the deposit improved by the addition of such salts as sodium bisulphite and potassium cyanide In other examples, notably those of tin and lead, the deposit is much improved by the addition of a trace of a colloid, such as glue or gelatine

Migration of ions. It is well known that ions migrate to their respective plates with definite speeds under unit potential gradient If deposition proceeds more rapidly than this rate of migration, the cathode liquid becomes impoverished of its metal, and this involves a falling off in the ampere-hour yield. The deficiency of metal around the cathode is counteracted to a limited extent by the ordinary process of diffusion, but there is a limit to the rate at which metal may be satisfactorily deposited, unless special steps are taken to make up for this slow migration Methods of circulation and stirring are based on this fact, and also on the further fact that the heavier solution formed around the anode falls to the bottom of the solution, and, owing to its superior conductance, gives rise to the conduction of an undue proportion of the current and consequently thicker deposits on the portion of the cathode in this heavier layer, hence a limit to the rate at which deposition may be safely allowed to proceed is se", and the time required to deposit a given amount of metal varies correspondingly.

E.M.F. for deposition. With an insoluble anode, a definite minimum E M.F is required to maintain electrolysis Thus, when copper sulphate is decomposed with a platinum anode, the total change is as follows:-

$$CuSO_4 + H_2O = Cu + O + H_2SO_4$$

This, as a chemical change, is endothermic, and electrolytically it demands a minimum EMF of about 1 25 volts for its continuance a copper anode is used, no change in the com-Energy changes position of the solution occurs at both electrodes are equal and of opposite character, and hence a much smaller E M F., viz. that required to overcome the ohmic resistance of the solution, is required With efficient anode solution and a good conducting electrolyte, such as is obtained in an acidified copper sulphate solution, a PD of about 03-05 volt will maintain a fair rate of deposition In the case of nickel, where there is usually inefficient anode solution and a poorly conducting electrolyte, a much higher PD is required, partly on account of the greater stability of nickel com-Figures given, therefore, for P D 's at bath terminals are of a very elastic character, depending upon the efficiency of anode solution, strength and temperature of the solution, distance between electrodes, presence or otherwise of conducting substances in the solution, and current density required, while the conditions controlling the allowable current density at the cathode are chiefly the strength of the electrolyte and its motion

The deposition of copper may be accomplished by simple immersion methods, resulting in a chemical exchange with the more positive

metal plated, thus

$$\frac{Zn}{Fe}$$
 +  $CuSO_4 = \frac{Zn}{Fe}$   $SO_4 + Cu$ 

The process is but little used, and then only for very cheap work Deposits on a surface which is undergoing simultaneous corrosion, cannot be sufficiently adherent to withstand the frictional processes of polishing and burnishing

The single-cell process, practised at one time, involves the generation of current and the deposition of metal in the same apparatus The process is illustrated in the Daniell cell, in which, if the copper cylinder is replaced by any conducting substance, copper is deposited

For the deposition of copper on biass and similar electro-negative metals, and on blackleaded or metallised non-conducting surfaces of wax, guttapercha, plaster, earthenware, &c, the following solution is of wide application

The composition is variable. Strong solutions admit of more rapid deposition without 'burning,' while the acid imparts conductance From such a solution deposition can be carried on at the rate of 40 amperes per square foot without stirring, while, with a properly stirred electrolyte, this current density may be increased up to 250 or 300 amperes per square foot

Previous to deposition, brass is cleaned first in potash (5-10 p.c) to remove grease. It is

of oxide, or tarnish, and, after rinsing, is scoured with pumice or ordinary sand or scratch-brushed. After finally rinsing, it is immersed in the solution, and should be covered immediately The deposit is pink, though dull When of sufficient thickness, the work is removed, rinsed, and the dull surface improved by scratchbrushing or polishing

For deposition upon non-conducting substances such as wax, guttapercha, &c, the following methods may be used .—

(1) Wax and guttapercha Thorough 'yet careful brushing with the finest plumbago causes sufficient to be retained to render the surface conductive Such surfaces when suspended in the copper solution, receive a deposit beginning at the metal connection, and gradually spreading over the whole surface, the thickness

of the deposit not being exactly uniform
(2) Plaster of Paris and unglazed earthenware may be rendered conductive by first seaking in linseed oil and allowing to dry They are thus linseed oil and allowing to dry rendered impervious to the solution, and the skin of dried oil will now retain blacklead, when this substance is applied by gentle brushing, sufficient to impart the necessary conductance

(3) Flowers, feathers, and other delicate objects which will not withstand the pressure of even gentle brushing, may be treated to produce a fine layer of some metal or sulphide on the surface by simple chemical means usual method is to paint them or immerse them in a weak solution of phosphorus in carbon disulphide, and, after drying, a brief immersion in silver nitrate solution leads to the precipitation of silver, which imparts the necessary conductance Again, the object to be treated may first be immersed in silver nitrate solution, and subsequently exposed to sulphuretted hydrogen, which produces a layer of silver sulphide, sufficient to cause the immediate deposition of copper at all points

A more successful method for wax and guttapercha involves covering these substances with copper by simple precipitation substance, usually as a mould or impression of a surface to be reproduced in electrotype copper, is first blackleaded and is then laid face upwards in a solution of copper sulphate filings are then spinkled on the surface filings in contact with the blackleaded surface produce innumerable voltaic couples, which bring about the deposition of copper quickly and completely The mould is then rinsed to detach the filings, and placed in the bath, when copper is at once deposited uniformly over the whole surface

On positive metals such as non, zinc, tin, Britannia metal, pewter, &c, copper cannot be successfully deposited from the acid sulphate solution on account of the ease with which these metals turn out the copper by simple immersion, and are, moreover, corroded by the acid more stable and more satisfactory compound for this class of work is the double eyanide of copper and potassium (CuCN KCN) Solutions of this compound are usually prepared by (1) dissolving the carbonate in a solution of potassium cyanide; (2) precipitating the sulphate with ammonia, dissolving the precipitate in creess of ammonia, and decolourising the blue then passed through aquafortis to remove scale solution with potassium cyanide. Many other methods are available and are in use The strength of such solutions may be varied between wide limits up to 4 ozs (or even more) of metal per gallon. An excess of potassium cyanide is essential, owing to the formation of insoluble single copper cyanide (CuCN) at the anode. The course of electrolysis in such double cyanide solutions (including those of zinc, silver, and gold) is very different from that of the salts of the mineral acids. In the case of copper, the reaction may be simply represented as follows.—

Anode CuCN KCN Cathode 
$$\left| \begin{array}{ccc} & & \text{Cathode} \\ & \leftarrow & \text{Cu(CN)}_2 \text{--} \text{K}^+ & \rightarrow \end{array} \right|$$
 then:

 $\begin{array}{ll} {\rm Cu(CN)_2 + Cu = 2CuCN} & {\rm K + CuCN \ KCN = Cu + 2KCN} \\ & {\rm (Insoluble)} & {\rm (deposited)} \end{array}$ 

The copper in the salt thus migrates as a complex ion to the anode, and there forms an insoluble compound This, unless removed, offers considerable resistance, and increases the difficulty of maintaining a normal current Sufficient potassium cyanide for its removal is formed at t'e cathode, and can be best brought to the anode by stirring, which facilitates the process of deposition by keeping the anode The solution of the anode deposit is more usually accomplished by using an excess of potassium cyanide called 'free cyanide' From such a solution copper is not deposited upon iron, tin, Britannia metal, pewter, &c, by simple immersion, and only very slightly on The solution is conveniently used warm, and yields a deposit of copper accompanied by hydrogen, which, as a rule, does not seriously interfere with the physical character of the deposit. The deposit is, under the best qualitative conditions, smoother than that from the sulphate solution, and once the positive metal has been coated in this solution, it may be removed, and, after thorough rinsing, introduced into the sulphate solution for a thicker deposit The cyanide solution, when strong, and used within certain limits of current density, yields twice as much copper per ampere-hour (viz 2 36 grams) as the sulphate, the cyanide compound being the monovalent cuprous salt.

Applications of copper deposition — The metal is most extensively deposited as a covering for more positive and more corrodible metals, and as a preliminary deposit before coating with the more expensive metals Deposits of copper on iron may then be burnished or polished and the surface improved by metal-colouring, when, by chemical processes, the surface is covered with a uniform film of oxide or sulphide by immersion in solutions producing these compounds, such as the salts of copper and soluble sulphides Uniform and pleasing effects are readily obtained, and may be rendered more permanent by subsequent lacquering metal is also deposited on a considerable scale in the electrotyping process, thin, but rigid reproductions of type, engraved plates and blocks being made in electro-deposited copper, and subsequently strengthened by 'backing with a lead-antimony alloy, which is then planed down to the required thickness.

Analytically, copper deposition is applied

The in the separation of this metal from more positive metals in an acid solution with a platinum anode From a solution of brass in nitric acid, the copper may be most completely deposited without interfering with the zinc The copper deposit made on a platinum dish or suitable electrode is then weighed, and the zinc obtained by either a chemical or a further electrolytic method (v Analysis) On a commercial scale, the deposition of copper has been applied by Cowper-Coles to the production of parabolic mirrors for lighthouse work, and to the production of sheet, wire, and tubes The manufacture of deposited tubes of large diameter has been successfully carried on by Elmore who arrange an agate burnisher to travel backwards and forwards along the tube during deposition, thereby producing a compact deposit of good mechanical properties. Tubes up to 16 feet long by 8 feet diameter, and weighing more than 3 tons, have been produced by this method More usually, however, tubes are of smaller dimensions, and are subsequently drawn down or opened longitudinally for the production of sheets Copper is deposited on a most extensive scale in the refining process, possibly as much as three-quarters of the total output of the metal annually being refined in this manner, using a solution of the sulphate with thick anodes of the crude metal The crude metal contains up to 10 p c of impurities deposited metal is almost entirely free from impurity Positive impurities pass into the solution, and are retained there by the application of the principle used in electrolytic analysis Electro-negative impurities, as silver and gold, pass as chloride and metal respectively into the mud at the bottom of the tank, from which they may be recovered Tin, antimony, arsenic, and bismuth are prevented from being deposited by a careful control of the conditions, and the sum-total of impurities in the refined metal is usually less than 0 05 p c

Deposition of silver. Silver, being an electronegative metal, is readily deposited by simple immersion. The process is applied commercially under the name of 'whitening' Solutions for this purpose contain a small proportion of a silver compound, such as chloride or cyanide, dissolved in cream of tartar, sodium chloride, or similar solvent. The deposition on properly prepared work is a case of simple replacement, and only very thin deposits are obtainable by this means. The same materials may also be used in the form of pastes, and rubbed on the prepared work with a wet rag, this method finding application in silvering clock and instrument dials.

Electro-deposition is most frequently practised. For this purpose the nitrate solution is unsuitable on account of its instability, being serviceable only as a voltameter electrolyte where the very electro-negative metal platinum is used. For silvering on more positive metals, those which usually require silver plating, a solution of the double cyanide of silver and potassium is used. This may be prepared from the metal as follows:—

(a) The metal is converted into intrate, and precipitated with potassium cyanide. When precipitation is just complete, the clear liquid is decanted and the precipitate washed and

subsequently dissolved in more cyanide solution, a further supply of cyanide being added for 'free cyanide'

(b) Precipitation may be effected by common salt, and the washed chloride dissolved in excess

of potassium cyanide

(c) A silver plate may be made the anode in a 5 p c solution of potassium cyanide, using a small copper or carbon cathode contained in some of the solution in a porous cell. The silver dissolves electrolytically, and the process can be continued until sufficient silver has passed into the solution and a good deposit is obtainable from it.

The solution is used cold, and yields a dull deposit of the metal, which may subsequently be scratch-brushed, burnished, and polished The addition of a small proportion of carbon disulphide (1 oz in 250 gallons) completely alters the character of the deposit from dull to bright. 'Bright plating' is, therefore, much practised The silver compound in the solution being somewhat unstable, special precautions are necessary before placing work in the solution for deposition Copper, brass, and nickel silver may be plated directly, though previous to deposition it is common practice to pass such work through a solution of mercury cyanide This solution can be prepared by treating a soluble mercuiic compound with excess of potassium cyanide, or by dissolving the oxide in cyanide A usual strength is 1 oz of metal per gallon From such a quicking, solution a bright deposit of mercury is obtained on brass, copper, and nickel silver by simple immersion, with the advantages that the more negative metal mercury is less oxidisable than copper, &c, and prevents simple immersion deposition of silver when the work is put into the bath, and possibly promotes the adhesion of the subsequently electrolytically deposited silver, by amalgamation More positive metals, like iron and zine, are first coppered in the alkaline bath, and subsequently treated as copper With the lead alloys it is usual to pass their straight from the cleansing processes through a separate silver solution, from which a rapid deposit is first made This is termed striking,' after which the work is transferred to the main solution to complete the deposition

The current density permissible is controlled largely by the metal content of the solution, Excessive current produces a light-grey, dark grey, or black deposit, which, being powdery, is unable to withstand the processes of burnishing

and polishing

Silver anodes should be of fine metal, preferably annealed, and, therefore, more susceptible to solution. Their behaviour is an indication of the condition of the solution. If a dull deposit collects upon them, with a falling off in the current, more free cyanide is required to dissolve the insoluble single cyanide which forms upon them. On the other hand, excessive free cyanide unduly promotes the solution of the anode, limits the permissible current density at the cathode, lowers the cathode efficiency, and promotes simple immersion deposition on work which is not 'quicked'. The amount of free cyanide present can easily be determined in this and other cyanide solutions by titration with standard silver intrate.

The deposition of gold is practised considerably on account of the pleasing colour and permanence of the deposit Deposition is easy owing to the strong electro-negative character of the metal, and for electro-gilding, as with electro-silvering, cyanide solutions are used. Gold may be applied to metallic surfaces by three methods, viz · (1) mercurial gilding; (2) simple immersion; and (3) the electrolytic method. Mercurial gilding is used for heavy deposits of gold. An amalgam of gold is first prepared by pouring mercury on to hot gold. Excess of mercury is squeezed through chamois leather, and the pasty amalgam preserved in balls under water—It is applied to the prepared metallic surface with a wire brush, which has first been wetted with a weak solution of mercuric This is rubbed on the amalgam and nitrate then on to the work, depositing a layer of amal-When a uniform coating is obtained, the mercury is expelled in a muffle, and the process repeated for thicker deposits The dull surface is improved by scratch-brushing. This process is still largely employed

Simple immersion methods are followed, for mere films of metal, in solutions of very varied composition. A moderately strong solution of sodium pyrophosphate is prepared (5-10 pc), and gold chloride added a little at a time. The warm solution at once gives up its metal, but the deposits can, in any case, only be very thin, though they may be thickened somewhat by 'quicking' in a mercury solution and re-

immersing in the gold solution

Gilding with the separate current is largely practised, using a solution prepared by the

following or similar methods.

(1) Making a gold sheet the anode in a cyanide solution (4 ozs per gallon), using a cathode contained in a porous cell. The process is continued until sufficient metal is lost from the anode or until the solution yields a good deposit.

(2) Precipitating gold chloride with ammonia, avoiding excess, and dissolving the washed pre-

cipitate in potassium cyanide solution

The metal content may vary from a few pennyweights to a few ounces per gallon solution is used warm with a gold anode, which is electrolytically corroded, or by a less desirable, though common, method of using an insoluble platinum anode and making frequent small The colour of the additions of gold chloride deposit varies considerably with the conditions, chiefly that of current Other conditions, such as temperature, size of anode, distance between electrodes, metal content, also apparently affect the colour; but these varying conditions primarily influence the current. Cold and weak solutions give a pale colour, which may be improved by warming. Excessive current produces darker and foxy deposits, which are, however, improved by scratch-brushing. More satisfactory deposition proceeds when work is kept in motion. The colour of The colour of the deposit is heightened somewhat by a preliminary coating of copper, while a deposit of silver has the reverse effect, though occasionally required. The colour is modified also by the addition of copper and silver cyanides to the solution Thick deposits are most satisfactorily obtained by occasional scratch-brushing and 'quicking.'

The deposition of nickel is now very largely applied on account of the hardness of the deposit and its permanence, together with the exceptional lustre which can be imparted to it by polishing. The solution most commonly employed is .

Nickel ammonium sulphate . 12 ozs Water . . . . 1 gallon

The conditions which affect the electro-deposition from this solution have been studied by Bennett, Kenny, and Dugliss (J. Phys Chem 1914, 18, 373, Bennet, Rose, and Tinkler, *ibid* 1915, 19, 564; Hammond, Trans. Amer. Electrochem Soc 1916, 30, 103) To this solution, additions are frequently made to increase the conductance, sal-ammoniae or common salt being used. Owing to the fact that no free mineral acid is permissible, and the tendency also to the formation of nickel hydroxide at the cathode, small additions of acids, such as citric, boric, or acetic, are made with varying success The only modification in the above solution which adds any appreciable advantage, is that of nickel sulphate, which is so much more soluble than the double salt, and from which, therefore, much stronger and therefore better, conducting solutions can be prepared thick deposits, Langbein recommends the use of 3 lbs of nickel sulphate per gallon, with the addition of ethyl sulphate or magnesium ethyl On account of the greater stability of nickel compounds and the poor solution of the anodes, and also the absence of free mineral acid, a distinctly higher EMF is required than for copper deposition, viz 5-6 volts for 'striking,' that is, rapidly coating the work with nickel, and then 2-3 volts for depositing the bulk of the metal The absence of free acid necessitates especial care in the preliminary cleansing processes, most work being finally passed through a weak solution of cyanide (5 pc), and quickly and thoroughly rinsed before being put into the vat. The nickelplating solution is usually worked cold, and is conveniently contained in lead-lined Anodes may be of east or rolled metal, the east being much superior, as its porosity assists its corrosion under the influence of the current Rolled metal is but little attacked by the amon SO<sub>4</sub>, and, failing the solution of the anode, sulphuric acid is produced and oxygen evolved; thus

## $SO_4 + H_2O = H_2SO_4 + O$

This acidity is to be avoided, and with it there is a diminution in the metal content of the solution which is similarly undesirable. Cast anodes are therefore much more largely used, more uniform conditions being maintained with them. On account of the high E.M.F. used, some hydrogen is evolved at the cathode, though the current efficiency should not ordinarily fall below 90 p.c. Occasional motion of the work is desirable in order to disengage any bubbles which may collect. Once deposition has commenced, the work should not be exposed more than momentarily, otherwise an imperceptibly thin film of oxide will form and almost entirely prevent the adhesion of further deposited metal. In fact, it is Toonsidered impossible to successfully deposit nickel on such

a surface or on a previous deposit. On this account, previously nickelled work should be entirely freed from its nickel, and, while this may be done by acid solutions, common practice follows the method of polishing, at the same time preparing the surface for the new deposit Copper and brass work is mckelled directly after the usual cleaning operations, whilst positive metals like zinc and Britannia metal require a previous deposit of copper from the cyanide According to Canac and Tasilly (Compt 1914, 158, 119), alummium is readily plated with a durable coating of nickel by first plunging the metal in a boiling potassium hydroxide solution, scrubbing it with milk of lime, followed by steeping in a 0.2 p.c solution of potassium cyanide for some minutes, after which it is transferred to a bath of 500 grams of hydrochlone acid and 1 gram of iron in 500 e.e. of water Between each bath it is well washed with water, and then plated electrolytically in a solution of nickel chloride The success of the plating apparently depends on the use of the ferruginous bath Iron and steel may be nickelled without coppering, the polished surface being first rendered somewhat dull by scouring with finest pumice. The deposit should be kept bright, and thus is attained by a current density not exceeding 5 ampeies per square Higher current densities produce dull deposits, tending to become powdery on the edges. The nickel deposit is very hard, and the final lustre is obtained by 'finishing' with Sheffield lime, which produces a highly lustrous surface—In addition to the extensive cycle and motor industry, nickel-plating is applied con siderably for small machine parts and instru-ment work and for 'facing' stereotypes, the deposit being much more durable than the stereotype metal

The deposition of iron is carried out on lines similar to those for nickel, solutions of ferrous sulphate or ferrous ammonium sulphate being used These must be entirely free from mineral acid, more so than in the case of nickel, and water should be well boiled before being used, to expel air and prevent the oxidation of ferrous to ferric compounds Ferric compounds under electrolysis yield little metal, being first reduced to ferrous at the cathode The process is not much used beyond the application in steel-The iron deposit facing? engraved plates obtained is free from carbon, but is very hard Anodes of the softest wrought iron containing a minimum of carbon are desirable, and require occasional scouring to remove the layer of insoluble carbon. The deposition of iron has also been used for thicker deposits, which have subsequently been removed and examined to investigate the properties of pure iron, and to more successfully determine the effect of impurities. For 'steel facing,' the engraved plates are cleaned, any scouring being done with only a light brush. The plate is then hung in the bath and covered quickly by 'striking. Deposition is then continued more slowly for 10-15 minutes. The deposit is very hard and withstands the printing process much better than copper. When the deposit shows signs of wear by the exposure of copper, the metal is simply and readily removed by immersion for a short time in a 5-10 pc sulphuric acid. By voltaic

action the iron is soon dissolved from the plate, and, after a light scouring with a fibre brush and subsequent rinsing, the plate is again immersed in the solution to receive a further coating of iron, thus considerably lengthening For a review of the work done in the its life commercial electro-deposition of cadmium, see Mathers and Marble, Trans Amer Electrochem Soc 1914, 25, 297 et seq

Of several cobalt solutions the following vield the best results. (1) crystallised cobalt ammonium sulphate, 200 grams per litre of water; (2) cobalt sulphate, 312 grams; sodium chloride, 196 grams; boric acid, nearly to saturation, water, 1000 cc. From these solutions, cobalt plates on brass, iron, steel, copper, tin, nickel-silver, lead, and Britannia metal, the coating being firm, adherent, hard, and uniform When polished, the surface, although brilliantly white, has a slight bluish tint Compared with nickel-plating, the plating with solution (1) is four times, and with solution (2) fifteen times, more rapid, but the voltage required is greater than that used for most nickel-plating baths Since cobalt is harder than nickel, a much thinner coating of cobalt will offer the same protective coating as a greater weight of nickel deposit, thick deposits of cobalt are superior in appearance and character to those of nickel (Kalmus, Harper, and Savell, J Ind Eng Chem 1915, 7, 379, Chem Soc Abst. 1915, 107, 11 465)

Deposition of alloys. (1) Brass. From mixed solutions of zinc and copper sulphates with the addition of acid, copper can be completely deposited free from zinc, this principle being applied in electrolytic analysis and also in With salts more alike in stability, this condition being attained in the double evanide olutions, the two metals are deposited simultaneously in proportions varying with conditions The cyanides are invariably used for this purpose, the zine and copper compounds being used in about the same quantity The mixed solution may be prepared in many ways, of which the following are in common use

(a) Treating solutions of the mixed sulphates with excess of ammonia and adding about 30 p c more cyanide (KCN) than is required to decolour-

ise the blue solution

(b) Preparing a solution of potassium cyanide (1-1½ lbs per gallon) and saturating about three-quarters of it with the mixed carbonates of zinc and copper, subsequently adding the remaining solution for 'free cyanide'

The solution may be used warm or cold If cold, a higher concentration is desirable, and also a smaller proportion of zinc compound If warm, the zine compound should be increased and the solution may be safely diluted Brass anodes are used, which, in a warm solution with a fair proportion of free cyanide, dissolve under the influence of the current. Lack of free evanide gives rise to the formation of a deposit of the two single compounds on the anode, adding resistance and reducing the current, and thus altering the colour of the deposit The colour and composition of the deposit vary with several conditions In general, the conditions which demand a higher E M.F. give rise to a lighter deposit containing more zinc, whilst those which reduce the resistance of the solution

tend to the production of redder deposits co taining more copper. Thus more yellow depos are produced in cold dilute solutions with lit free cyanide, whilst the copper in the depo is increased when the solution is strengthened warmed, or more free KCN added. The co position of the deposit is controlled also by t proportion of the two metals in the solution and the fact that from cyanide solutions copi and zinc are ordinarily deposited in the prop-tion of 63 6 of copper 32 5 of zinc accoun approximately for the 2 1 proportion in deposit yellow brass The colour of the deposit lightened by the addition of ammonia ammonium carbonate, these substances assisti the solution of the zinc compound, which is l soluble than the copper compound, and p venting the formation of basic compounds

The methods of deposition of brass foll on the same lines as those of copper from t

cyanide solution

(2) Other alloys are deposited to only The choice of solution very limited extent depends upon the ease of mixture of solution from which the constituents can be separate deposited Thus for bronze, the sulpha chloride and cyanide solutions, are unsuital A satisfactory mixture can be made from t double oxalates of the two metals with amn nium oxalate, the solution being worked wa with a bronze anode Similarly, a solut from which nickel silver may be deposit may be prepared by mixing the separate dou cyanides of copper, zinc, and nickel with pot sum cyanide, or by dissolving the alloy nitric acid, precipitating the mixed carbona and dissolving the mixture in evanide deposition of these alloys is subject to the sa conditions as for brass, attention being paid those conditions which facilitate the deposit of the more positive constituents, viz tin, 71 and nickel

The electrodeposition of zine has of late ve been extensively carried on As a protect coating for iron and steel an electro deposit ] great advantages over the ordinary process galvanising The sulphate solution is commo used In some cases zinc anodes are employ and these to a large extent maintain a consti composition of solution Other processes emp insoluble anodes, resulting in the formation sulphuric acid equivalent in quantity to the z The deposition of zinc from si deposited acid solutions demands special conditions, a for general plating purposes the free acid tl formed is neutralised by filtering the liq through zinc dust From neutral solution there is the tendency for the formation of ba compounds leading to the growth of mo deposits Such addition agents as gluc beta-naphthol, &c, lead to a considera improvement in the character of the deposit

Another extending application of z deposition is found in the recovery of z from liquors produced by treating certain z ores with weak sulphuric acid. The zinc ores with weak sulphuric acid usually present to the extent of 6-10 pc. electrolysis this is reduced considerably, the acid solution formed by the use of insolu anodes is then used for extracting the z compounds from a further quantity of ore process has been applied largely at Anaconda reguline deposits may be obtained from solutions of the stannite, stannate, or double exalate with ammonium, with ease. They are prepared as

(1) The hydrate is precipitated from stannous chloride with alkali, and the precipitate dissolved in excess of alkali

(2) By converting the metal to hydrated oxide with nitric acid, and dissolving the washed oxide in alkalı

(3) By adding a solution of ammonium ovalate to tin chloride A white precipitate first forms, and this is soluble in excess.

Solutions containing from 1 to 2 ozs of metal per gallon are of convenient strength these solutions tin may be obtained as a reguline deposit within certain limits of current density Excessive current produces a grey spongy deposit, which may, however, be considerably improved by the addition of a small amount of glue or gelatine, thereby extending the limit of allowable current Solutions of moderate strength yield metal quantitatively, and the anode suffer normal corrosion, maintaining the strength of the electrolyte The dull white deposit is improved by gentle scratch-brushing with a soft brush Tin may also be deposited without the current by using an oxalic acid solution to which tin chloride is added metal is deposited by simple immersion, and the process may be improved by the addition of grain tin with the work, which facilitates deposition by local voltage action, while tin The process is hastened passes into the solution by immersing the small work in zinc trays, the zine passing into the solution as an anode this case the solution must be replemshed with tin chloride, and rejected when any large amount of zinc has accumulated in it

The deposition of lead may be similarly offected from alkaline solutions, which yield reguline deposits, whilst the acetate produces the well-known crystalline deposit. The process has but little application, but more recently the deposition of the metal has been successfully applied by Betts to the refining of the metal The electrolyte is a solution of lead fluosilicide, which may be made by dissolving the oxide or carbonate in the acid With unrefined anodes the lead passes into solution, whilst silver is retained in the anode mud. Good reguline deposits are obtained by the addition of glue or gelatine in small amount

More recently, lead perchlorate has been successfully used Perchloric acid is prepared from the sodium salt with a slight excess of hydrochloric acid Insoluble sodium chloride is filtered off and the acid liquid evaporated down to 135° C until excess of hydrochloric acid is expelled The perchloric acid solution is diluted and neutralised with litharge, a small proportion of the acid being set aside to add to the lead perchlorate solution, so that lead and free acid are present to the extent of about 5 pc. each. Very satisfactory deposits are obtained from this solution, especially with the addition of peptone in the proportion of 0.05 p c, and deposits of considerable thickness may thus be produced. With moderate current Vol. II. -T.

Deposition of tin may be successfully carried densities deposition proceeds almost quantitaout from a number of solutions. The chloride | tively, and this with the high chemical equivalent yields a beautiful crystalline deposit, but of lead has suggested its use for voltmeter purposes

A feature of lead deposition is the production of lead peroxide at the anode. This has long been known, and the production of coloured rings and films was first observed by Nobili, in Thus in a solution of lead acetate, or of 1826litharge in alkali, and a polished anode of mickel or iron, coloured rings are obtained if a wire The colours cathode is held close to the anode are due to the gradually varying thickness of the film, which brings about interference in the light falling upon them With anodes of other shapes the colours follow the shape of the anode, though they pass off into a more or less rounded form These colours are called 'metallochromes' With excessive current a dull-brown deposit of lead peroxide is at once formed

Anodic deposition of lead peroxide is also applied in the estimation of small amounts of lead in certain alloys from solutions containing nitric acid

The deposition of platinum is carried out from a solution of the chloride in sodium phosphate A platinum anode is insoluble and the solution needs the addition of the platinum compound. A brown colour sometimes occurs owing to some of the metal passing into the colloidal form

Palladium may be successfully deposited from a solution of its double chloride, and the process has been applied by Cowper-Coles for producing a non-corrodible covering for electro lytically deposited parabolic mirrors in copper

Cobalt deposition follows on lines similar to those for nickel, mercury, which may be deposited from the nitrate solution, has been applied to the construction of electrolytic meters, in which a small and definite fraction of the total current is shunted through a cell in which mercury is deposited, the weight of the mercury deposit automatically recording the current passing through the meter, and at constant potential difference the energy absorbed in the S F. circuit

ELECTRUM v NICKEL **ELEMI** v Olfo-resins ELEMI, ESSENTIAL OILS OF, v OLYO-

**ELEMICIN** (3 4 5-trimethoxyallylbenzene) A constituent of the essential oil of elem has been synthesised by Mauthner (Annalen, 1917, 414, 250) by condensing allyl bromide and pyrogallol-2 6 dimethyl ether, which yields 2 6 dimethoxyphenyl allyl ether (I), which, by heating at 220° is converted into 2 6-dimethoxy-4-allyxphenol (II), this, on methylation, yields elemicin (III.).

626 ELEMOL.

**ELEMOL**, C<sub>15</sub>H<sub>26</sub>O, a naturally-occurring monocycle sesquiteipene alcohol found in Manilla elemi oil; b.p  $152^{\circ}-156^{\circ}/17$  mm; D<sup>20</sup> 0 9411;  $n_{\rm D}$  1 5030;  $\alpha_{\rm D}-5^{\circ}$ . Forms a benzoate; b p  $214^{\circ}-218^{\circ}/10$  mm, D<sup>20</sup> 1 0287;  $n_{\rm D}$  1 5378;  $\alpha_{\rm D}-6^{\circ}$  Elemol, when reduced by means of formic acid, sodium hydrogen sulphate, or zinc-dust, yields Elemene C15H24; bp 115°- $117^{\circ}/10 \text{ mm}$ ,  $D^{20} 0 8797$ ,  $n_{D} 1 4971$ .

Elemol is the first monocyclic sesquiterpene alcohol to be found in Nature or to be obtained artificially (Semmler and Liao, Ber 1916, 49, 794).

ELIQUATION or LIQUATION. A metallurgical term to denote the operation of separating substances by taking advantage of their different fusibilities

ELLAGIC ACID. Ellagic acid, C14H6O8, is obtained from numerous tannin matters which contain an ellagitannin by boiling the aqueous extract with dilute sulphuric acid. In many cases it is found in the free state, due probably to the hydrolysis, by fermentation, of the ellagitannin or ellagic acid glucoside originally present, and such a fermentation frequently occurs when aqueous extracts of the tannin matters are allowed to stand for some time

Divi-divi (Casalpinia coriaria [Willd]) (Lowe, Zeitsch anal Chem 1876, 14, 40), myrobalans (Terminalia chebula [Ratz]) (Lowe, ibid), algarobilla (Casalpinia brevifolia) (Zolffel, Beilstein, 2, 1085), and valonia (Quercus aegilops [Linn]) yield considerable quantities of ellagic acid, and are the best natural sources of this material It has also been isolated from oak galls (Chevreul, Ann Chim Phys. 1828, [2] 9, 329), oak bark (Etti, Montash 1880, 1, 226), the bark of Picca excelsa (Link) (Strohmer, ibid 1881, 2, 539), pomegianate ind (Rembold, Annalen, 1867, 143, 288), Quebracho colorado (Perkin and Gunnel, Chem Soc Trans 1896, 69, 1307), Arctostaphylos uva-ursi (Spreng) and Contaria myrtifolia (Linn) (Perkin, abid 1900, 77, 424), Hamatoxylon campeachranum (Linn) (Perkin, ibid 1897, 71, 1137), the fruit of Casalpinia digyna (Rottl) (Nierenstein, Chem Zeit 1909, 87), and the bank of Terminalia catappa (Linn) (ibid) In fact, ellagic acid probably always occurs, if only in minute quantity, in all plant products which contain gallotannin Most interesting are the Oriental bezoar stones, concretions which are found in the stomachs of goats and other animals, and consist largely of ellagic acid These originate, without doubt, from the fact that the animal has fed upon plants in which some quantity of an ellagitannin is present

Ellagic acid was first discovered by Chevreul (Ann. Chim. Phys l.c.) in oak galls, and more closely examined by Braconnot (ibid 9, 187) According to Pélouze, it possessed the composition C<sub>7</sub>H<sub>4</sub>O<sub>4</sub> (*ibid*. 54, 356), whereas Merklein and Wohler (Ann Chem. Pharm whereas 55, 129), who isolated it from bezoar stones and termed it 'bezoaric acid,' showed that its true formula is C<sub>14</sub>H<sub>6</sub>O<sub>8</sub>

The precipitate of crude ellagic acid, which is produced by boiling the tannin extract with dilute mineral acid, is purified by wasning with alcohol and subsequent recrystallisation. A (Barth and Goldschmiedt, Ber. 1878, 11, 846; similar procedure is also satisfactory with the Schiff, ibid. 1879, 12, 1534; Zolftel, Arch.

deposits so frequently produced by fermentation and alluded to above. Ellagic acid is most readily crystallised from pyridine (Perkin and Nierenstein, Chem Soc. Trans 87, 1416), and the product, which is either a pyridine compound or contains pyridine of crystallisation, is washed with alcohol.

Lowe (Zeitsch Chem 1868, 4, 653) was the first to prepare ellagic acid synthetically by oxidising gallic acid with arsenic acid, and this was subsequently accomplished by Ernst and Zwenger (Annalen, 1871, 159, 32) by heating ethyl gallate with sodium carbonate solution in the presence of air, and by Griessmayer (ibid 1871, 160, 55) by heating gallic acid with water and iodine. It is readily produced by oxidising gallic acid dissolved in acetic acid by means of potassium persulphate and sulphuric acid (Perkin and Nierenstein, lc), and together with flavellagic acid by the addition of potassium persulphate to a solution of gallic acid in sulphuric acid (Perkin, ibid 1906, 89, 251) Under similar conditions, gallotannin produces ellagic acid (Nierenstein, Ber 1908, 41, 3015, and 1909, 42, 353), and it can also be prepared by boiling the tannin with hydrogen peroxide solution (ibid 1907, 40, 917) Herzig and Bronneck obtained good results by passing air through an ammoniacal solution of gallie acid ethyl ester, and also of gallotannin (Monatsh 1908, 29), whereas Sisley (Bull Soc chim 1909, [4] 5, 727) prepared ellagic acid from tannin by means of alcoholic potassium hydroxide solution According to Rupe, a 60 p c yield is produced when gallie acid in sulphuric acid solution is oxidised with sodium nitrite (Chemie d naturl Farbstoffe, 1909, 2 162)

Ellagic acid crystallised from pyridine forms prismatic needles, which are converted by alcohol into a pale yellow crystalline powder, but when purified by means of its acetyl derivative, the product is almost colourless (Perkin and Nieren-It is very sparingly soluble in all the usual solvents, dissolves in alkaline solutions with a pale yellow colour, and from these liquids, when diluted with hot alcohol, is deposited on acidifying in a crystalline con-When heated it does not melt below 360°, and sublines at higher temperatures with considerable carbonisation. With nitrie acid containing nitious acid and subsequent dilution, it gives a blood-red colouration (Griessmayer's reaction), and this property, considered at one time to be characteristic, is known to be also possessed by flavellagic acid (Perkin)

According to Merklein and Wohler (lc), the alkaline solution of ellagic acid becomes reddishvellow on exposure to air, and deposits black crystals of potassium glaucomelanate

When boiled with phenylhydrazine in alcoholic solution, or when carefully heated with phenylhydrazine, ellagic acid forms the product  $C_6H_5NH\ NH_2,C_{14}H_6O_8$ , which crystallises in yellow needles, and is readily reconverted into ellagic acid Ellagic acid gives similar compounds with aniline and quinoline (Goldschmiedt, Monatsh 1905, 26, 1139)

Pharm. 229, 123), colourless needles, mp 343°–346° (Perkin and Nierenstein); tetrabenzoylellagic acid  $\rm C_{14}H_2O_8(C_7H_5O)_4$ , colourless needles (Goldschmiedt and Jahoda, Monatsh 1892, 13, 151), m p 332°-333° (Perkin), cllagic acid monomethyl ether (14H<sub>5</sub>O<sub>7</sub>(OCH<sub>3</sub>), yellow crystalline powder (Goldschmiedt, Monatsh 1905, 26, 1139); diacetylellagic acid monomethyl ether C<sub>14</sub>H<sub>3</sub>O<sub>7</sub>(OCH<sub>3</sub>)(C<sub>2</sub>H<sub>3</sub>O)<sub>2</sub>, crystalline powder (Goldschmiedt); ellagic acid dimethyl ether C<sub>14</sub>H<sub>4</sub>O<sub>6</sub>(OCH<sub>3</sub>)<sub>2</sub>, crystalline powder (Goldschmedt), and ellagic acid tetramethyl other C<sub>14</sub>H<sub>2</sub>O<sub>4</sub>(OCH<sub>3</sub>)<sub>4</sub>, colourless microscopic needles (Goldschmiedt, Herzig and Pollak, Monatsh 1908, 29, 263), have been described

Ellagic acid gives the following salts  $KC_{14}H_5O_8$ , minute yellow needles (Perkin and Wilson, Chem Soc Trans 1903, 83, 134); K<sub>2</sub>C<sub>14</sub>H<sub>4</sub>O<sub>8</sub>, prisms (Merklem and Wohler, Perkin and Wilson), K<sub>2</sub>C<sub>14</sub>H<sub>4</sub>O<sub>8</sub>, KOH, microscopic prisms (Merklem and Wohler),

crystalline powder, \*NaC<sub>14</sub>H<sub>5</sub>O<sub>8</sub>,H<sub>2</sub>O, yellow needles (Ernst and Zwenger, Annalen, 1871, 159, 32)

The tinctorial properties of ellagic acid are somewhat feeble, but in 1887 it was placed on the market by Messrs Meister, Lucius, and Bruning, under the name of 'Alizarin Yellow in paste,' and was recommended as yielding, with chromium mordants, greenish-yellow shades of considerable permanence

Rembold (Annalen, 1867, 143, 288) states that when ellagic acid is treated with sodium amalgam, it gives \gamma-hexahydroxydiphenyl

three distinct sufohydrocllagic acids C14H10O6.  $C_{14}H_{10}O_7$ ,  $C_{14}H_{10}O_9$ , and glaucohydrodluqic acid, but, according to Nierenstein (Ber 1908, 41, 1649), this latter compound is in reality the pentahydroxydiphenylmethylolid referred to later By the action of boiling potassium hydroxide solution (Barth and Goldschmiedt, Ber 1879, 12, 1242), ellagic acid gives herahydroxydiphenylencketone (\*) C<sub>1</sub>, H<sub>2</sub>O<sub>7</sub>, by fusion with potassium hydroxide *B-hexahydroxydiphenyl*, and by fusion with sodium hydroxide \gamma-hexahydroxydrphenyl, together with some of the  $\beta$ - compound

When ellagic acid is distilled with zinc-dust fluorene is produced (Rembold, Ber 8, 1494. Barth and Goldschmiedt, ibid 11, 846, cf also Graebe, *ibid* 1903, 36, 212)

As a result of their investigation, Barth and Goldschmiedt (ibid 1879, 12, 153) suggested that ellagic acid could be represented by one of the two following formulæ:

and of these the first was preferable

Whereas such a constitution requires the presence of five hydroxyl groups, Schiff (Ber 1879, 12, 1534) considered that acetylellagic acid was  $C_{14}H_{2}(C_{2}H_{3}O)_{4}O_{8}$ , and this was subsequently corroborated by Zolffel (Arch. ellagic acid.

Pharm, 229, 123) Schiff, at the same time, proposed two formulæ for ellagic acid

the latter of which is identical with that preferred by Graebe ( $l \epsilon$ ) In a later investigation, Goldschmiedt and Jahoda (Monatsh 13, 49), as the result of their preparation of tetrabenzoylellagic acid, were satisfied of the existence in ellagic acid of but four hydroxyls, and con-

Graebe (Ber 1903, 36, 214), among other criticisms, pointed out that a substance having this constitution, and which may be more clearly written as follows (1) -

$$\begin{array}{c|c} COOH & COOH \\ \hline (1) & OH & OH \\ \hline \end{array}$$

would yield, on distillation with zinc-dust, not fluorene, but an isomeric hydrocarbon (2), if this, indeed, were capable of existence. In the same paper this author announced the interesting fact that diphenylmethylolid

on distillation with zinc dust, gives not only diphenyl and methyldiphenyl, but also fluorene; and in a theoretical discussion of the subject, suggested that Schiff's second formula (given above) most probably represents the true constitution of this substance

The subject was remvestigated somewhat later by Perkin and Niercustein (le), who, on digesting ellagic acid with boiling potassium hydroxide solution, obtained Barth and Goldschmiedt's so-called hexahydroxydiphenyleneketone ('13H<sub>8</sub>O<sub>7</sub>) From this substance a penta-acetyl derivative ('13H<sub>1</sub>O<sub>7</sub>(('2H<sub>3</sub>O)<sub>5</sub>, piismatic needles, mp. 224°-226°, and a pentabenzoyl derivative, ('13H<sub>3</sub>O<sub>7</sub>(('7H<sub>5</sub>O)<sub>5</sub>, plates, mp. 257°-259°, were prepared. As, moreover, zinc-dust distillation gave fluorene, there could be little doubt that this compound was in reality pentahydroxydiphenylmethylolid, as, indeed, was surmised by Graebe.

From their results, together with a study of other similar compounds, Perkin and Nierenstein considered that Schiff's formula, and which can be more clearly written thus .

Finally, Herzig and Pollak (Monatsh 1908, 29, 263), by a study of the methylation products of ellagic acid, obtained results further corroborating this constitution.

When Goldschmiedt's tetramethylellagic acid is digested with alcoholic potash and methyl iodide, diphenyl-1 2.3:6.7 8-hexamethoxy-5 10-di-carboxylic methyl ester (mp. 109°-111°)

is obtained, together with a small quantity of diphenylmethyloid-2 3 6 7.8-pentamethoxy-5-carboxylic acid methyl ester (2), m.p. 187°-189°.

By means of alcoholic potash, these compounds (1) and (2) yield respectively diphenyl-1 2 3 6 7 8-hexamethoxy-5 10-dicarboxylic acid (3), mp 238°-240°, and diphenyl-2 3 6.7 8-pentamethoxy-1-hydroxy-5 10-dicarboxylic acid (4), m.p. 200°-203°:

(4) 
$$CH_3O$$
 $OCH_3$ 
 $COOH$ 
 $OCH_3$ 
 $OCH_3$ 

Ellagic acid is probably not a tanning material in the true sense of the term, but is of considerable importance nevertheless in the tanning process. Thus the ellagitannic acid absorbed by the hide is at least in part converted into ellagic acid, and the deposit thus formed adds weight and bloom to the finished article.

Other members of this group have been prepared from m-hydroxybenzoic acid, p-hydroxybenzoic acid, proto-catechuic acid, sym. dihydroxybenzoic acid, gallic acid, and from ellagic acid itself—An account of these is given below

## FLAVELLAGIC ACID

Flavellagic acid  $(^{\circ}_{14}\text{H}_{0}^{\circ})_{9}$  is obtained in conjunction with ellagic acid when gallic acid is oxidised with potassium persulphate and strong sulphuric acid. By the employment of a dilute acid, flavellagic acid almost free from cliagic acid can be produced, and for this purpose the following process is recommended (A. G. Perkin, Chem. Soc. Trans. 1906, 89, 251). 20 grams of gallic acid are treated with 160 c. c. of 96 p. c. sulphuric acid, 66 c. c. of water added, and the hot solution thus obtained is cooled to 50°, and maintained at this temperature during the gradual addition of 40 grams of potassium persulphate. A. G. Perkin and F. M. Perkin (Chem. Soc. Trans. 1908, 93, 1194) have obtained the same compound by the electrolytic oxidation of gallic acid in the presence of sulphuric acid,

and more recently Bleuler and Perkin (bid. 1916, 109, 530) have prepared it by heating gallic acid at 110°-120° with arsenic and sulphuric acids.

Flavellagic acid crystallises from pyridine in small yellow prismatic needles which contain pyridine, and do not melt below 360°. With nitric acid containing nitrous acid and subsequent dilution, it gives the blood-red colouration (Griessmayer reaction), which is also produced by ellagic acid. Dilute alkalis dissolve it with a yellowish-green colouration, and by distillation with zinc-dust fluorene is produced. Acetyl-flavellagic acid.  $C_{14}HO_{9}(C_{2}H_{4}O)_{5}$ , colourless needles, melts at  $317^{\circ}-319^{\circ}$ , and benzoyl-flavellagic acid.  $C_{14}HO_{9}(C_{14}H_{2}O)_{5}$ , prismatic needles, melts at  $287^{\circ}-289^{\circ}$ . Flavellagic acid dyes mordanted woollen cloth shades somewhat resembling but stronger than those produced by ellagic acid.

Chromium Aluminium Tin Iron.
Yellowish- Pale greenish- Pale Dark oliveolive yellow. yellow brown.

By the action of boiling 50 p.c. potassium hydroxide solution, flavellagic acid gives hexahydroxydiphenylmethylolid

colourless needles (mp above 300°). This compound dissolves in solutions of the alkali hydroxides with an orange-yellow tint, and this, on dilution with water and exposure to air, develops a strong bluish-violet colouration. The acetyl derivative  $C_{13}H_2O_8(C_2H_3O)_8$ , colourless prismatic needles, melts at  $232^\circ-234^\circ$ . Flavellagic acid is hydroxyellagic acid, and possesses the following constitution.

(cf Herzig and Tscherne, Monatsh 1908, 29, 281)

Flavellagic acid gives with diazomethane the methyl ether C<sub>14</sub>HO<sub>4</sub>(OMc)<sub>5</sub>, mp 245°, and this is converted by methyl iodide and potassium hydroxide into methyl-3 4 5 6 2'.3'.4'-heptamethoxy/liphenyl-2.6'-dicarboxy/late

 $C_{12}H(OMe)_7(CO_2Me)_2$ 

m p 83°-87°. The latter, on hydrolysis with potassium hydroxide, yields the acid

 $C_{12}H(OMe)_7(CO_2H)_2$ 

m p  $163^{\circ}$ - $167^{\circ}$ , in the anhydrous condition. From water it crystallises with  $1\rm{H}_2\rm{O}$  and then melts at  $95^{\circ}$ - $100^{\circ}$  (decomp ) (M. von Bronneck, Monatsh 1908, 29, 281)

When gallic acid 3 4-dimethylether, or gallic acid trimethylether is oxidised by means of potassium persulphate and sulphuric acid at 45° (Herzig and Schmidinger, Monatsh. 1910, 31, 918), flai ellagic acid tetramethylether

yellow needles, m.p. 270°-271°, is produced, the acetyl derivative of which melts at 270°-271° By means of methyl sulphate and potassium hydroxide, the heptamethoxy compound is obtained

## CERULEOEILAGIC ACID

Cœruleoellagic acid is prepared by heating ellagic or flavellagic acid (1 part) with sulphuric acid (monohydrate 10 parts) to 200° and allowing the temperature to fall to  $185^{\circ}-190^{\circ}$  and remain there for 30 minutes. The product isolated by pouring into water is purified by conversion into its acctyl derivative, and this is subsequently hydrolysed by means of sulphuric acid in the presence of acctic acid. It may also be obtained by the oxidation of ellagic or flavellagic acids (1 part) in sulphuric acid solution (10 parts) with arsenic acid (1 part) at  $100^{\circ}-130^{\circ}$  (Perkin, Chem. Soc. Trans. 1916, 109, 529)

Cæruleocllague acud, C<sub>14</sub>H<sub>6</sub>O<sub>10</sub>, separates from hot pyridine in small pale yellow prismatic needles which melt above 360° and are very sparingly soluble in the usual solvents. Concentrated solutions of the alkali hydroxides dissolve it with a greenish-yellow colour, which on dilution become green and finally of a blue colour, and these changes appear to arise from oxidation. Sodium and potassium carbonate solutions also give blue liquids. On distillation with zinc-dust fluorene is produced.

Acetylcoruleoellagic acid,  $C_{14}O_{10}(C_2H_3O)_6$ , colourless needles, mp 330°-332°, and benzoylcoruleoellagic acid  $C_{14}H_{10}(C_7H_5O)_6$ , mp 343°-345°, have been prepared. The latter is more sparingly soluble in benzoic anhydride than the corresponding derivative of flavellagic acid, and this property provides a means for the separation of the two substances.

By digestion with 50 p.c. potassium hydroxide solution diluted with its own volume of alcohol, coeruleoellagic acid gives a compound  $\rm C_{12}H_{10}O_8$ , evidently octohydroxydiphenyl. This consists of an almost colourless crystalline powder, soluble in dilute potassium hydroxide solution, with a reddish-violet colouration which becomes brown on exposure to air. The acetyl derivative  $\rm C_{12}H_2O_8(\rm C_2H_3O)_8$  forms colourless needles melting at  $\rm 177^\circ{-}178^\circ$ 

The constitution assigned to corruleoellagic acid is as follows —

a formula which represents it as either a dihydroxy-cllagic or hydroxy flavellagic acid

Coruleoellagic acid possesses well-marked dyeing properties of a similar character, though more strongly developed than those of ellagic and flavellagic acids, and may have practical utility. Owing to its sparing solubility it gives the best results in the paste form. The following shades are obtained by the employment of mordanted woollen cloth —

Chromium Aluminium Tin Iron.
Deep olive- Greenish- Dull Greenishyellow yellow black

In the case of the aluminium mordant the

employment of chalk in the dye-bath exerts a detrimental effect

## CATELLAGIC ACID.

Schiff (Ber. 12, 2590) on heating protocatechuic acid with arsenic acid, obtained a substance which he named catellagic acid, and to this he assigned the formula  $C_{14}H_{10}O_7$  or  $C_{14}H_8O_7$  Perkin and Nierenstein (Chem. Soc. Tians 1905, 87, 1417) obtained a similar substance by the oxidation of protocatechuic acid, and also parahydroxybenzoic acid with potassium persulphate and sulphuric acid to the latter authors, catellagic acid C14H6O6 crystallises from pyridine in colourless needles, which melt above 360°, and sublime with but moderate carbonisation at higher temperatures Solutions of the alkali hydroxides dissolve it with a pale yellow colour, and with nitric acid it gives a magenta coloured liquid Dracetylcatellagic acid, colourless prismatic needles, melts at 322°-324°

Catellagic acid, by distillation with zincdust, gives fluorene, and is closely related to ellagic acid. Its constitution (Perkin and Nierenstein) may be expressed as follows—

From the oxidation products of parahydroxybenzoic acid, Perkin and Nicrenstein isolated, in addition to catellagic acid, a second compound  $\rm C_{14}H_8O_6$ , colourless needles, mp above  $\rm 360^\circ$  This, which gives the diacetyl derivative  $\rm C_{14}H_6O_6(\rm C_2H_3O)_2$ , colourless needles, mp  $\rm 267^\circ - 268^\circ$ , and by distillation with zinc-dust fluorene, possesses the formula

and evidently originates from the interaction of one molecule of protocatechuic acid and one of parahydroxybenzoic acid

## METELLAGIC ACID

Metellagic acid (Peikin and Nierenstein, Chom Soc Trans 1905, 87, 1425) is obtained in small amount together with other substances by oxidising metahydroxybenzoic acid in sulphuric acid solution with potassium persulphate at about 30°. The product is fractionally crystallised from acetic acid, the second crop acetylated, the acetyl derivative thus obtained purified by repeated crystallisation from acetic anhydride and hydrolysed with sulphuric acid in the usual manner.

Metallagic acid  $\mathrm{C_{14}H_6O_5}$  crystallises from acetic acid in colourless needles which readily sublime at high temperatures. With caustic alkaline solutions it behaves similarly to catellagic acid giving a yellow colour which is discharged on heating. Distilled with zine-dust a small quantity of a crystalline hydrocarbon is produced which appears to be identical with fluorene. Acetylmetelluque acid  $\mathrm{C_{14}H_6O_5(C_3H_3O)}$  crystallises in colourless leaflets, in p. 269°-271°

The reactions of metellagic acid are in (Burm.) (nat ord. Myrsinaceæ) harmony with those required by a member of the ellagic acid series, and the following formula assigned to it is probably correct .-

In the mother liquors obtained during the purification of acetyl metellagic acid, a second acetyl compound, in p 237°-243°, was isolated This on hydrolysis gave a substance crystallising in glistening yellow needles, the nature of which has not been ascertained

By the oxidation of suitable hydroxybenzoic acids, a series of hydroxydiphenyldimethyloid derivatives have thus been prepared, viz

Metellagic acid  $C_{15}H_5O_4(OH)$ Catellagic acid  $C_{15}H_4O_4(OH)_2$ Resoflavine  $C_{14}H_3O_4(OH)_3$ Ellagic acid  $C_{14}H_2O_4(OH)_4$ Flavellagic acid  $C_{14}HO_4(OH)_4$ Coruleoellagic acid C14O4(OH)a

AGP

ELLAGITANNIC ACID. A tanning principle present in divi-divi (Lowe, Frdl 14, 35), in algarobilla (Zolffel, Arch Pharm 229, 123), in myrobalans (Lowe, lc 44, Zolffel, lc), and probably to some extent in all plant products which contain a gallotannin (v ELLAGIC ACID)

According to Lowe, the molecule of ellagitannic acid contains 5 hydroxyl groups replaced by acetyl It gives precipitates with gelatin, albumen, alkaloids, tartar emetic, and a nearly black precipitate with feiric acetate Its alcoholic solution gives, with lead acetate, a vellow precipitate

Zolffel (lc) considered it a sugar-free tannic acid of the formula

## $C_{6}H_{2}(OH)_{2}COO_{6}H_{2}(OH)_{2}COOH_{2}$

whereas Perkin and Nierenstein (lc) suggested it to be a glucoside or a condensation product of two molecules of gallotannin

Nierenstein (Ber 1910, 43, 1267) has obtained it in a pure state by repeated treatment of the crude substance from myrobalans with alkalı and ethyl chloroformate, and then decomposing with pyridine according to Fischer's method; and gives its formula as

$$C_{26}H_{28}O_{19},3H_2O$$

It crystallises from a mixture of pyridine and acetic acid in pale-yellow plates, which, after sintering at 300°-306°, melt at 329°-336°  $[a]_{p}^{17} + 1802^{\circ}$  So obtained, it readily yields ellagic acid with dilute sulphuric acid, but is not decomposed by 10 pc sodium carbonate solu-As it is hydrolysed by emulsin to luteoic acid. Nierenstein considers it to be a glucoside of this compound (v Tannins) A G P ELUTRIATION. The separation of the

lighter from the heavier particles of a powder by washing and decantation

EMANIUM v ACTINIUM

EMBARIN. Trade name for sodium mercurisalicylsulphonate

EMBELIC ACID. The active constituent of the fruit of the Indian plant Embelia riles

Employed as an anthelmintic

EMBOLITE. Native silver chlorobromide containing chlorine and bromine in indefinite proportions It is the commonest of the hornsilvers or cerargyrites  $(q \ v)$ , occurring as compact horn-like masses or as small, indistinct cubic crystals of a greenish-grey colour Frequently a small amount of rodine is also present, and the formula is Ag(Cl,Br,I); the colour of this iodembolite variety is usually greenish-yellow to orange-yellow Large quantities have been mined with other ores of silver in Chile, Mexico, and Broken Hill in New South Wales Prior and Spencer, Min Mag 1902, 13, 174)

EMERALD v BERYL, and GLUCINUM EMERALD GREEN. Hydrated chromium sesquioxide (v Chromium), also applied to Schweinfurth green (v art Arsenic, also Pia-MENTS)

EMERALD-NICKEL v ZARATITE EMERALD, ORIENTAL, v CORUNDUM

EMERY. (Émeri, Fi , Smirgel, Schmirgel, Ger) An impure variety of coundum. It is, however, more of the nature of a rock, rather than a simple mineral, consisting e-sentially of a fine-grained aggregate of corundum (50-60 pc) and magnetite (30-40 pc) Several other minerals are also present in smaller and variable amounts Hamatite and limonite are present as alteration products of the magnetite, and others, such as diaspore, margarite (emerylite), muscovite, kyanite, have probably been derived from the corundum In addition, there may also be tourmaline, chloritoid, biotite, staurolite, rutile, idociase, pyrites, iron spinel (hercynite), &c The material is dull black, dark grey, or bluish-black, and has much the appearance of a fine-grained iron-ore; sp gr 3 75-4 31. It is massive and opaque, breaking with a more or less regular fracture, and is magnetic abrading power (measured by the amount of glass abraded by a given weight of emery in a certain time) ranges from 65 to 32, as compaied with sapphire at 100. These values are only roughly proportional to the amount of alumina (corundum) present in the mixture, depending also to some degree on the texture of the material.

The analyses on next page are of material from Naxos, J-IV (I and II, the extremes as regards the amount of alumina of seven analyses by Oser, quoted by S. A. Papavasihou, 1913; III and IV, by E. Ludwig, quoted by G. Tschermak, 1894), Asia Minor, V, by R. Kramer, 1907, VI, Peekskill, New York, by G. H. Williams, 1887. Several older analyses have been published by J. L. Smith (Amer J. Sci. 1850, 1851, 1866), and by R. Jagnaux (Bull Soc. Min. France, 1884)

Analysis III corresponds with the mineral composition corundum 52 4, magnetite 32 1, tourmaline 115, muscovite 2, margarite 2 p.c.; and IV with corundum 50, magnetite 33, tourmaline 9, chloritoid 4, muscovite 3, calcité lpc.

The only sources of supply are Greece, Asiatic Turkey, and the United States deposits in the island of Naxos in the Grecian archipelago have been known since remote Here, on the north-east side of the times

	Ι¹	II	III	IV.	v	VI 2
Al <sub>2</sub> O <sub>3</sub>	66 16	$60\ 05$	57 67	$56\ 52$	60 10	46 53
Fe <sub>2</sub> O <sub>3</sub> FeO		24·03 ( 4 02		34.65	33.20	41 29
MnO	0.25	0 85			_	
CaO	. 162	1 51	0.43	0 90	0.48	
MgO	. 0 38	1 05	0.83	0.43		9.43
K,O			0.31	0 40		-
Na,O	. —		n d	0.60		*******
SiO,	. 3 23	509	564	545	1 80	242
TiO,		-	n.d.	n d		0.51
$B_2O_3$		****	1 15	0.88	-	
$H_2O$		3 67	0 70	0.42	562	_
	99 76	100 27	100 09	100 25	101 20	100 19
Sp gr.	4 06	384	371	3.98	*******	

island, the emery occurs as irregular beds and lenticular masses in crystalline limestone (marble), associated with crystalline schists and intiuded by granito. Large isolated blocks also occur weathered out on the surface. The methods of mining are very primitive, and resort is made to the old method of fire-setting, the surface of the rock being heated by fires and suddenly cocled by water to develop cracks. The production is the monopoly of the Greek government, and amounts to about 10,000 tons per annum. Emery also occurs on the neighbouring islands of Irakhia, Sikinos, Paros, and Amurgos, but these deposits are not worked (On the Naxos deposits, v. W. H. Cottrell, Foreign Office Rep., Miscell. Ser. No. 368, London, 1895. G. Tscheimak, Min. Petr. Mitt. 1894, anv. 311. S. A. Papavasiliou, Zeits Deutsch Geol. Gesell. 1913, lxv. 1–123.)

The important deposits in western Asia Minor, around Smyrns, were discovered by J Lawrence Smith in 1847 Here also the emery occurs as lenticular bodies in bluish crystalline limestone, associated with mica-schist, gneiss, and granite; and as loose blocks weathered out in the reddish surface soil The principal mines are situated on the Gume Dagh or Messogis Mountain in the Aidin district (E of Ephesus and S E of Smyrna) Others are near Manisa (the ancient Magnesia), near Kula and Adula (N. of Ala-Shehr=Philadelphia), and on the neighbouring islands of Samos and Nikaria The output of 20,000 to 30,000 tons per annum is shipped from Smyrna. (On the Asia Minor deposits, v J L Smith, Amer J Sci 1850, 1851, and Original Researches in Mineralogy and Chemistry, 1884, R. Kramer, Klemasiatische Schmirgelvorkommnisse Inaug -Diss Leipzig, Beilin, 1907)

In the United States, deposits have been worked at Peekskill in Westchester Co, New York, occurring here as segregations in gabbro, and in amphibolite-schist at thester in Hampden Co, Massachusetts The annual output amounts to only about 1000 tons, and that from Chester has now fallen off. (On the United States occurrence, n. G. P. Merrill, The Non-Metallic Minerals, 2nd edit New York, 1910)

An abrasive material mined at Wildenreuth and other places in Bavaria under the name of emery (Schmirgel, Ger) is really a mixture of garnet and quartz. In the trade the name emery has also been applied to the refuse from cutting gems and from gem-washings.

' Also Cr<sub>2</sub>O<sub>5</sub> trace <sup>2</sup> Also P<sub>2</sub>O<sub>5</sub> trace, S 0 01

In the preparation of the material for the market, after being carefully hand-picked at the mine, it is crushed under stamps and sorted into various grades of fineness through wire sieves. The finest 'flour of emery' is prepared by washing and elutriation Emery powder is used by lapidaries and stone workers, for polishing plate glass and grinding lenses In the form of emery paper, emery cloth, and emery sticks, the powder is fixed by means of glue For emery wheels the powder is mixed with a cementing material, such as shellac, sodium silicate, or vulcanite, or it is baked with loam or kaolin and thus embedded in terra-cotta or porcelain Such wheels are much used by lapidaries and dentists, and for grinding and polishing metals, &c Attempts have been made to use pieces of emery, in place of diamond, in the crowns of rock-drills (On the technical uses of emery, v A Haenig, Der Schmirgel und seine Industrie, Wien, 1910) L J S.

EMETAMINE v IPECACUANHA

EMETINE v IPECACUANHA. EMIN RED v PRIMULINE

EMODIN v GLUCOSIDES, RHUBARB; SYN THETIC DRUGS

EMPRESSITE. A silver-tellurum mineral AgTe, discovered in the Empress-Josephine mine, Kerber Creek, Colorado, associated with galena and native tellurum. Lustre, metallic, streak, greyish-black, fracture, conchoidal, hardness, 3–3 5, sp gr. 7 51 (Bradley, Amer J. Sci. 1914, 38, 163)

EMPYROFORM v SYNTHETIC DRUGS.

EMULSIN. The term 'emulsin' applied originally to the enzyme or mixture of enzymes, present in bitter and sweet almonds, which hydrolyses amygdalm to glucose, benzaldehyde, and hydrogen cyanide. The same enzyme was shown subsequently to hydrolyse many natural glucosides as well as the synthetically prepared  $\beta$ -glucosides, though it was entirely without action on the stereoisomeric  $\alpha$ -alkylglucosides (Fischer, Zeitsch physiol Chem 1898, 26, 60) The name is now used to comprise all enzymes which hydrotyse  $\beta$ glucosides Such enzymes are widely distubuted in plants besides the kernels of fruits, they also occur in the leaves and young twigs of the higher plants, in most mould fungi, especially in Polypoius species parasitic on trees, and in bacteria Henry and Auld (Proc Roy Soc 1905, B, 76, 568) have identified emulsin as present in yeast.

β-Glucosides are also hydrolysed by certain animal extracts, e.q intestinal mucous membrane, a fact which is taken to indicate the presence of emulsin

There are, however, well-marked differences in the number of substances which these various emulsin preparations are able to hydrolyse, and the investigations of Armstrong and others (Chem Soc Proc 1910, 334) indicate, that whilst the leaf enzyme (emulsin) is, as a rule, a specific for the glucoside normally occurring in the particular plant, the seed enzyme, especially that of the almond, is able to attack a number of glucosides. For this reason almond emulsin is considered to be a mixture of several closely related enzymes. According to H. E and E F. Armstrong and Horton (Proc Roy So. 1908, 80, 321), it contains a \$\textit{\textit{Boundary of Soc. 1908}} \text{ 1908}.

632 EMULSIN

proper, which attacks  $\beta$ -glucosides; a cyanase, which eliminates hydrogen cyanide from amygdalin, an amygdalase, splitting off glucose from amygdalin; and a luclase, acting on milk sugar (cf. Rosenthaler, Arch. Pharm 1910, 240, 105)

Almond emulsin hydrolyses  $\beta$ -glucosides and also the disaccharides, lactose, gentiobiose, cellose, and melibiose It further hydrolyses the synthetical  $\beta$ -methylgalactoside The emulsin of Aspergillus niger, on the contrary, has no netion on lactose or on  $\beta$ -galactosides In referring to emulsin, it is therefore advisable

to specify the source of the enzyme

Preparation —Ground sweet almonds from which the fat has been expressed are macerated with water and a little toluene or chloroform at the ordinary temperature for 12-24 hours The liquid is expressed through a cloth, protein precipitated with a few drops of acetic acid, and the clear filtrate precipitated with alcohol added 50 c c. at a time till no further precipitate The liquid is decanted and the precipitate rapidly washed with absolute alcohol and ether to dehydrate it, and dried in a vacuum Alternatively the original precipitate may be collected on a filter paper, dued roughly on bibulous paper, and dissolved in water in a closed bottle, to which a little toluene is added The clear solution so obtained is very active, and retains almost constant activity for many It is essential to carry out this preparation as rapidly as possible

The dried enzyme is a soft, white powder, giving most of the reactions of proteins, but losing these on purification. Its chemical nature has not been the subject of any detailed investigation. Active preparations are to be

obtained in commerce

Emulsin exhibits maximum activity in neutral solution Alkali stops the action, small quantities of acids usually increase the activity, owing to their effect in neutralising

alkaline impurity

Experiments with emulsin are usually carried out at about 37°; the optimum temperature is 45°. Emulsin is less resistant to destruction by heat in presence of alcohol below 50 p.c. but above this point the reverse is the case, owing to the fact that in stronger alcohols the enzyme is precipitated, and in this condition is more resistant to heat. In sterilising plants containing enzymes it is best to use alcohol of such strength as to produce a liquid containing about 60 p.c. of alcohol (Bourquelot and Bridel, J. Pharm ('him 1913, 7, 65)

Emulsin is detected qualitatively by its action on amygdalin, liberating hydrogen eyanide. The activity is tested quantitatively either against amygdalin or saliein, by measuring the amount of sugar formed in a given time.

Emulsin is also able to bring about synthetic change (Armstrong, Proc. Roy. Soc. 1905, B, 76, 592), van't Hoff, Sitzungsber K. Akad. Wiss Berlin, 1910, 48, 963). Emulsin exhibits hydrolytic activity in alcoholic solutions. Although unable to hydrolyse salicin in presence of 95 p.c. alcohol, it becomes active in 90 p.c. alcohol, hydrolysis continuing slowly until 37 p.c. of the glucoside has been decomposed. As the strength of the alcohol is diminished equilibrium is more rapidly attained, and a greater proportion of the salicin undergoes hydrolysis (Bour-

quelot and Bridel, Compt rend 1912, 154, 944). Similarly, emulsin is inactive when suspended in solutions of glucosides in acetone or ethyl acetate, but becomes able to effect hydrolysis when 10 pc of water is added to the acetone or the wet ester is used When emulsin acts on a solution of glucose in 85 p c alcohol, the corresponding  $\beta$ -methyl or  $\beta$ -ethyl glucosides are formed, the same equilibrium point being reached in solutions containing originally glucose or alkyl The more dilute the alcohol the greater is the hydrolysis and the less the synthesis (Bourquelot and Bridel, Compt rend 1912, 155, 319) The position of the equilibrium is independent of the quantity of emulsin added and of the temperature, but it varies with the strength of the alcohol and with the amount of glucose present. The property of emulsin to facilitate synthesis has been utilised by Bourquelot to prepare a large number of glucosides of the alcohols, including such substances as geraniol and glycol, and also the corresponding galactosides

The emulsin contained in a number of plant preparations is able to hydrolyse dl-benzaldehyde cyanohydrin, and also to synthesise this compound from benzaldehyde and lydrocyanic acid In some cases the synthesis is asymmetric, and an optically active cyanohydrin is formed (cf. Rosenthaler, Arch. Pharm. 1913,

251, 56)

(For a list of the glucosides attacked by emulsin, v Giucosides) E F A

ENAMEL BLUE. Cobalt blue (v PIGMENTS) **ENAMELS** are vitreous compositions capable of adhering by fusion to the surface of metal or of pottery The base of the enamel is generally a colourless glass, in which are suspended particles of an opaque metallic oxide or salt, generally stannic oxide Up to about 900°, tin dioxide exists finely suspended in the enamel, but at a higher temperature combination or solution may occur and opacity is diminished Titanic oxide has been used in place of stannic oxide, and antimony compounds have been employed. Opacity may also be obtained by the use of arsenious oxide, calcium phosphate, cryolite, or fluorspar. A preparation known as 'artificial cryolite' sometimes replaces the mineral When bone ash is used, a little nitie may be added to remove the grey colour; the nitre also prevents reduc-tion of the lead The vitreous base of most The vitreous base of most enamels is a glass containing much lead, but when the presence of lead is objectionable, as in enamelled vessels for culmary purposes, soda-and potash-glasses should be used. If great fusibility be required, it may be secured by the association of borates with the silicates Borate enamels are very hard, but are deficient in elasticity, and may be affected by moisture. Colour is imparted to the enamels by the introduction of various compounds, generally metallic oxides, the use of which, in many cases, has been known from remote ages Percy's analyses of the enamelled bricks of Babylonia (circa 700 B c) showed that a dense white was obtained by the use of stannic oxide, a nich yellow by means of antimony, apparently in the form of lead anti-moniate ('Naples yellow'); and a deep blue by copper, probably introduced as oxide, but existing in the enamel as a copper sodium silicate. The ancient Egyptian blue enamel, or glaze, ENAMELS 633

contains a definite copper calcium silicate, forming crystals remarkable for their intense pleochroism, to which Fouqué gave the name of vestorien (Bull Soc Fr Min [12] 1889, 36) (v. EGYPTIAN BLUE). The ancient Egyptians seem to have been ignorant of the art of enamelling on metal.

The enamelling of pottery is a simple operation, since it consists in the mere fusion of an opaque glaze on to the surface of the ware. But the application of a vitreous body to a metallic surface is less easy, and it is to this process that the term 'enamelling' is sometimes restricted, while the works of art produced by enamelling on metal are themselves known as 'enamels'

The most ancient enamels found in North-Western Europe are referable to the prehistoric Iron Age, sometimes distinguished as Late Celtic In these ancient enamels, brilliantly coloured opaque glass is embedded in cells of shallow cavities in bronze. It is probable that the vitreous material was applied as a powder, mixed with a vehicle so as to form a paste, and then heated sufficiently to cause its fusion, and thus secure firm attachment to the metal

A similar method, known as champ levé enamelli g, was extensively employed at a much later date in Germany and at Limoges in France—a town which for centuries was the chief centre of artistic enamelling. In this process the design is traced on a plate of copper or bronze, and all the parts to be enamelled are excavated by the graver, these hollows are then filled with enamel, which is afterwards fused, and the entire surface is finally ground smooth. The design thus appears in permanent colour, each tint being defined by ridges, or separated by areas, of metal which are the clevated parts of the original ground. The exposed metal is sometimes gilt.

Another and probably older method of inlaid enamelling is that known as cloisonné Here the design is outlined by strips of metal fixed to the metallic base, and the spaces between these bands are filled with enamel, which is then fused and the surface ground Each coloured area is, therefore, sharply separated from its neighbour by the edge of the metal fillet Of this character were the early Byzantine enamels, which were generally executed on plates of gold. The Chinese have for many centuries (at least since the 13th) been extremely skilful in this work, and the Japanese have applied a similar method to The Japanese the ornamentation of porcelain attach the cloisons to the base with a gummy cement, and even when the ground is a metal, they generally use no solder In some cloisonné work the surface is not ground down, so that the fillets stand in relief above the general surface of the enamel

The use of transparent enamels for the ornamentation of metalwork was introduced by the Italian goldsmiths of the 13th century, and afterwards became popular in France. The design was generally worked in low relief, or chased on gold or silver, and brilliant translucent enamels were applied in such a way that the varying thickness produced differences of tint. This style is known as basse-taille. In certain rare enamels, probably of the 15th century, the pattern is outlined by strips of metal without any foundation, and the spaces in this grating

are filled in with coloured transparent enamels; such cloisonné work is termed plique-à-jour

Artistic enamelling has been largely used, not only for inlaid and encrusted work, but for the production of paintings on metal. The painted enamels of Limoges, introduced in the 15th century, were usually executed on slightly convex plates of copper, the subjects being generally in white and grey on a deep-violet ground, but the process was extended to the decoration of dishes, ewers, and other objects, and various colours were employed, sometimes heightened with gold In this country, enamel painting was largely practised, by the family of Bone, during the latter part of the 18th and the early years of the following century While their smaller works, or miniatures, were executed on gold, the larger examples were on plates of The copper plate, having been annealed copper and cleaned with dilute sulphuric acid, was coated on both sides with white enamel powder it was then fired, and the enamelled face ground This white enamel, which was rendered so opaque by means of stannic oxide as to conceal the metal ground, was imported from Venice in the form of round cakes. The surface Venice in the form of round cakes to be painted was further prepared by the application of a second coat of cnamel, more fusible and transparent than the first, this material, known to the enameller as 'flux,' was used in the form of rods and beads of white Venetian Upon the fluxed surface, carefully ground smooth, the painting was executed in enamel colours, or powdered glasses mixed with an appropriate vehicle, and as it was necessary to vitrify each colour separately, a complicated subject would require several successive firings in a muffle, each liable to injure or spoil the work

Recent years have witnessed a great revival in artistic enamelling, with the introduction of various improvements in technical details

In addition to art enamels, there is a growing demand for coarse work in enamelled sheet-iron for street plates, railway notices, and other advertisements, which are generally executed in blue and white In applying enamel to metal it is necessary to use a material which will adhere firmly to the base, at the same time, it must fuse at a temperature lower than that which softens the metal, or melts the solder if any be present, and the enamel should have a coefficient of expansion by heat as nearly as possible that of the metal ground It is obviously important that, when enamel is used as a lining for saucepans, stew-pans, and other vessels for culinary purposes, it should be free from all ingredients which are poisonous, or may become so by contact with food An important industry has been developed in recent years, especially in Germany, Austria-Hungary, and France, in the manufacture of vessels for domestic use, in cast or sheet metal, soft sheet steel is commonly used for hollow ware

The object to be enamelled, having been carefully cleaned by picking in an acid bath, is coated with a thin layer of ground enamel. This primary coating need not be opaque, and generally contains a little cobalt or nickel oxide. A suitable material is obtained, according to P. Randau, by fusing a mixture of finit or quartz 30 parts, borax 25, and felspar 30 This product is then ground with kaolin 10 75 parts, felspar 6,

634 ENAMELS.

and magnesia, 1.25; the clay tends to lessen the fusibility. The finely powdered ground enamel 18 mixed with water to a creamy consistency and applied to the ware, slightly heated, after the coating has dried, the object is fired in a muffle furnace at a temperature rather under 1000°. After this stage of enamelling, the ware is scrubbed to remove scale, and a second coating is applied, this time opaque white enamel may be made, according to the same authority, from a mixture of flint 375 parts, borax 275, tin dioxide 30, soda 15, nitie 10, ammonium carbonate 75, and magnesia 7, this mixture is fritted and then ground with flint 6 12 parts, tin dioxide 3 66, soda 0 7, and magnesia 0 7. The cover enamel is suspended in water and spread over the ground enamel, and the ware is then carefully fired in a muffle at a lower temperature than that of the first firing In some cases a third coat of enamel is applied, while in other cases the ground layer is dispensed with, and only a single white coating Finally the surface may be decorated by painting or transfer printing in enamel colours, or by photographs In some enamels, like granite ware. the colour is applied by spraying

(On enamels, v Sir H H Cunynghame, On the Theory and Practice of Art Enamelling upon Metals, 3rd ed London, 1906, Lewis F Day, Enamelling, London, 1907, Alexander Fisher, The Art of Enamelling upon Metal, London, 1906; Philip Eyer, Die Eisenemaillierung, Leipsig, 1907, Paul Randau, Die Fabrikation des Emails und das Emailheren, Vienna, 4th ed 1909, Enamels and Enamelling, English translation of P Randau by Charles Salter, London, 1900; Julius Grunwald, translated by Herbert H Hodgson, The Theory and Practice of Enamelling  $\mathbf{F} \mathbf{W} \mathbf{R}$ on fron and Steel, London, 1909)

ENARGITE. Copper sulphaisenate Cu<sub>3</sub>AsS<sub>4</sub> or 3Cu<sub>2</sub>S As<sub>2</sub>S<sub>5</sub>, occurring as orthorhombic crystals with perfect prismatic cleavages and usually a prismatic habit The colour is ironblack with a metallic lustre, sp gr 4 44 It is not a common mineral, but in a few coppermining districts, for example, at Butte in Montana, Morococha in Peru, and in the Sieria de Famatina in Argentina, it is of some importance as an ore of copper

ENDIVE. Cichorum endima (Linn), the blanched leaves are used as a salad or garnish

According to Konig, their average composition is-

Other carbo-Water Protein

Fat Sugar hydrates Fibre 0.1 0.8 1.8 0.6 Ash 94 1 18

There are two chief varieties—with crinkled leaves and with smooth leaves. The former is apparently richer in nitrogen and ash, but poorer in carbohydrates (Dahlen, Landw Jahrb 1874, 3, 312, and 4, 614)

An analysis of the ash of endives by Richardson (Annalen, 67, 377) gave the following

figures -

 $MgO SO_3 P_2O_5$ K<sub>2</sub>O  $SiO_2$   $Fe_2O_3$ Na<sub>2</sub>O CaO 379 12 1 120 18 52 30 24 6 3 4 H. I.

ENDLICHITE v VANADIUM ENESOL v Synthetic Drugs ENGLISH BROWN v Azo- COLOURING MATTERS.

ENGLISH GREEN, or CHROME GREEN, v. CHROMIUM; PIGMENTS.

ENGLISH PINK, or ITALIAN PINK, v.

ENGLISH RED, or VENETIAN RED, v. PIGMENTS

ENHYDROS v. AGATE.

ENSILAGE is a process of preserving green fodder for cattle The term silo is applied to the pit or other depository in which the fodder (silage) is contained, and appears to be of Basque derivation, and originally meant a pit for storing This method of preserving grass and other succulent forage crops has been in use for many years in Germany, whence the practice was carried to France, and afterwards to America. Although the German system had been fully described by Johnston, in 1843, in the Transaction of the Highland Agricultural Society (new series, 9), no general notice of it was taken in these islands until 1882, when a report of the British Legation at Washington on ensilage, as practised on Mr Havemever's estate in New Jersey, was published as a blue-book, since which time many silos, of varying forms and construction, have been made in this country The object of all these is to exclude the air from the mass of fodder, allowing only a greater or less fermentation according to the practice of different authorities, and to this end silos have been formed (1) wholly or partly underground, (2) in the form of enclosed buildings. Silage has also been largely made by building up the green fodder in the form of an ordinary stack, of which only the top is covered Whether it be silo or stack, the principal factor is the superposition of a sufficient pressure, either by dead weight or by some mechanical system of screws, levers, or similar appliances The changes which forage undergoes in the silo are due to fermentation, and, as may be supposed, this action varies considerably with the materials stored and with the condition of storage, as, for example, green maize undergoes puncipally lactic fermentation, whilst in English fodder fermentation produces alcohol which is almost immediately converted into aldehyde If the air has not been properly expelled from the silo, or if the fodder has been pitted in a very damp condition, the aldehyde becomes quicky converted into acetic acid

Silage is of two kinds-viz. 'sweet' and 'sour.' Sweet silage is the result of allowing the grass to remain without pressure, and so permitting air to mingle with it until such time as a temperature of 125° to 150° F. is reached by the fermentation process, sour silage, on the other hand, results when pressure is at once applied, whereupon an acetic or lactic fermentation takes place It appears that a temperature of about 125° is sufficient to kill the bacteria which produce acid fermentation; 'if the bacteria are killed and the silo is covered and weighted, the enclosed mass of green fodder will remain sweet, and be practically preserved under the same conditions as fruits, vegetables, or meats are preserved when canned be the case, it will be at once intelligible that by less packing of the fodder when put into the silo, and extending the time of filling until the temperature rises to a point fatal to the bacteria, the resulting silage will be sweet and free from

acidity; whilst sour silage is produced by at once consolidating, covering up, and weighting the grass fodder so as to prevent the temperature rising to the point fatal to the bacteria (A Voelcker, J Roy Agric Soc 20).

Two samples of sweet silage showed only a trace of acid, not exceeding in quantity that contained in ordinary fresh meadow grass —

A Annual Street	_ 1					
	Silage made from					
THE SECOND FLOW	Clover and rye grass mixed	Meadow grass				
Moisture	75 80	74 40				
*Albuminoids .	2 53	2 56				
Sugar and other carbo-	200	2017				
hydrates soluble in water	1 43	2 99				
Crude vegetable fibre	18 31	17 90				
Mineral matter (ash)	1 93	2 15				
(1000)	- 00	0				
	100 00	100 00				
*Containing nitrogen	0 40	0 41				
Volatile acids (calculated acetic acid)	0 01	0.01				
Non-vole tile acids (calculat	ed					
as lactic acid)	0.01	0.02				
-	1					

A sample of sour silage made from grass placed in a heavily weighted silo showed the following composition —

Water	76 01
Soluble albummoids	0.91
Insoluble albummoids	1.78
Soluble carbohydrates	2.87
Crude fibre	15.82
Soluble mineral (ash)	1.63
Insoluble mineral matter (ash)	0.98
	100 00
Total nitrogen	0.43
Volatile acids (calculated as acetic acid)	0 19
Non-volatile acids (calculated as lactic acid)	1 24

This silo was provided with a tap at the bottom from which juice was drawn off from time to time. This drainage had a sp gr of 1.022, and contained, per imperial gallon.

	Grains
Albuminoids	1,008 56
Fixed acid (lactic acid) .	516 42
Volatile acids (butyric and	
acetic acids)	476 70
*Carbohydrates and amides	1,528 42
Mineral matters (ash) .	1,768 20
Water .	64,701 70
	70,000 00
*Containing non-albuminous	•
nitrogen	35.84

The dry matter (dried at 100°) which was left on evaporation of its drainage contained.

*Albuminoids . Lactic acid (Carbohydrates and amides	20 90 10 71				
Mineral matter (ash)		$\frac{31}{36} \frac{73}{66}$			
		100 00			
*Containing nitrogen †Containing nitrogen	•	$\frac{3}{0} \frac{34}{51}$			

The subject of changes during ensilage has been very carefully investigated both at Rotham sted by Lawes and Gilbert, and at Woburn by Voelcker The results obtained present a striking similarity, and it will suffice if one set of tables (those of Dr Voelcker) is referred to here, although the Experiments on Ensilage conducted at Rothamsted (Lawes and Gilbert) should also The sile to which these figures relate was one made at Woburn in 1886, and was of the character generally termed 'sour' It was weighted with stones in strong clm boxes to 112 lbs per square foot, and was unprovided with any drain The silo was closed on July 3 and opened on December 16, the contents being very fair,' although 'not really fine,' meadow grass The numbers, in a condensed form, are given in the following table

QUALITY AND COMPOSITION OF THE GRASS PUT INTO SILO, AND OF THE SILAGE PRODUCED

	Total weight	Water	Volatile acids	Non-volatile acids	Soluble albuminoids	Insoluble	Digestible fibre	Woody fibre	Soluble carbo- hy drates, &c	Soluble mineral matter	In-oluble mmeral matter	Total nıtrogen	Albuminoid	Non-albuminoid nitrogen
Grass put in silo Silage removed Loss Loss per 100 lbs of fresh grass	Ibq 32,780 30, *89 2,391 7 29	lbs 23,108 22,035 1,073 3 27	109 +109 +0 33		lbs 47 197 +150 +0 46	lbs 733 252 481 1 47	lbs 3278 2326 952 2 90	lbs 2670 2409 261 0 79	lbs 2265 2254 11 0 03	1b4 430 471 + 11 +0 12	lbs 249 230 19 0 06	lbs 151 155 +4 +0 01	lbs 123 73 50 0 15	lbs 28 82 + 54 +0 16

The main features shown in the table are as follows. The total loss due to formentation, evaporation, &c, in making the silage was 7.29 pc on the fresh grass. Of this, 3\{\} pc consisted of water, so that the loss in dry matter was only about 4 p.c. The loss of total introgen when, as here, no drainage was allowed to flow away, is very slight; but the introgenous bodies have undergone considerable change from the albu-

minoid to the non-albuminoid condition, the fibre has been diminished, insoluble albuminoids have been lessened, and soluble albuminoids increased (Voelcker, J. Roy. Agric. Soc 23, 410)

Since the nutritive value of food depends to a considerable extent on the albuminoid compounds contained in it, it becomes of importance from the economic point of view, to determine the quantity of such compounds in fresh green 636 ENSILAGE

food and in silage Kinch found that 55 p c of the total nitrogen present in the silage of grass was of a non-albuminoid nature, whilst in the original grass only 9 p c was of this form. In the case of mangold leaves, Kellner found that of the total nitrogen present, 27 8 p c. in the original leaves, 45 5 p c in the silage, and as much as 59 7 p c in silage preserved in stoppered jars, was non-albuminoid. Chifford Richardson, examining maize silage, found

Original stalks
Silage No 1
, No 2
Dried fodder
. Per cent total introgen as non-albuminoid
. 21 2
. 44 6
. 44 6
. 15 6

Silage from young maize . 53 3 , , , older , . 47 1

These results with grass, mangolds, and maize show that in ensilage a large portion of the albuminoids is converted into non-albuminoid introgenous substances, whilst in the ordinary drying of fodder no such change seems to take place. This is made clear by a reference to Dr. Voelcker's table given above under the heading 'Quality and composition,' &c, where the woody fibre, as indeed the whole of the fibre, has been diminished, insoluble albuminoids are lessened, and the soluble albuminoids increased

Ensilage is largely practised in Canada and in America for the sake of having a supply of green' fodder during the winter months, especially in Canada where the winter is long In England, since the interest attaching to novelty has waned, it has gradually died out of practice, except as a resource for adoption in unusually wet seasons, when haymaking is impossible or is so long delayed as to threaten loss of the crops In such circumstances those who possess the simple tackle necessary for putting up silage stacks in the open field have a material advantage over their neighbours Otherwise, in a country in which roots are easily grown and preserved for winter use, and where the actual period of the year in which vegetation is suspended is comparatively short, it would seem pieferable, having regard to the deterioration which occurs in ensilage, especially as regards albuminoids, to adhere to the more ordinary practice of hay-making

.The application of silage as fodder and its practical effects with stock have been carefully investigated both at Rothamsted and Woburn

(See Report on the Practice of Ensilage at Home and Abroad, Jenkins in J. Roy. Agric. Soc., new series, 20., On the Chemistry of Ensilage, Voelcker, ibid., Experiments on Ensilage conducted at Crawley Mill Farm, Woburn, 1884-5 and 1885-6, Voelcker, ibid. 22., Experiments on Ensilage conducted at Crawley Mill Farm, Woburn, 1886-7, Voelcker, ibid. 23., articles in the Agricultural Gazette, April. 27, 1855 et. seq., by Sir. J. B. Lawes and Sir. Henry. Gilbert, reprinted in pamphlet form under the title of Experiments on Ensilage conducted at Rothamsted.)

ENSTATITE. Metasilicate of magnesium, usually with some ferrous oxide isomorphously replacing magnesia, (Mg, Fe)SiO<sub>3</sub>, and with increasing iron passing into bronzite and hypersthene lt belongs to the orthorhombic

series of the pyroxene group, and is of importance as a rock-forming mineral and as a constituent of meteoric stones. Rarely it is clear and transparent and of a rich green colour, when it is used as a gem-stone, such material is found in the 'blue ground' of the South African diamond mines.

L. J. S.

ENTEROL v Synthetic drugs. ENZYMES and ENZYME ACTION v. Fermentation, Barley, Chemical Affinity

EOSAMINE v Azo- colouring matters
EOSIN v Triphenyl methane colouring

EOSOTE v Synthetic drugs

EPHEDRINE, C<sub>10</sub>H<sub>15</sub>ON, crystals, m p 39°, b.p 225°, occurs in Ephedra vulgaris. It is a secondary base, furnishes crystalline salts, and on 'exhaustive methylation' yields eventually trimethylamine and a-phenylallyl alcohol  $({}^{\circ}_{9}H_{10}O)$ . On distillation the hydrochloride furnishes methylamine and propiophenone When heated with hydrochloric acid, ephedrine undergoes with hydrochloric acid, ephedrine undergoes isomerisation to  $\psi$ -ephedrine (isoephedrine), crystals, in p 117°, which also occurs with it in E vulgaris. Both bases are poisonous and ephedrine hydrochloride has been used as a mydratic (Ngai, Chem Zentr 1894, 1 470, Merck's Bericht, 1893, 13; Ladenburg and Oelschlagel, Ber 1889, 22, 1823; Miller, Arch Pharm 1902, 240, 481; Flaecher, ibid 1904, 242, 380, Schmidt and collaborators, ibid 1905, 243, 73, 1906, 244, 239, 241, 269; 1908, 246, 210, 1909, 247, 141, 1911, 249, 305. 246, 210, 1909, 247, 141, 1911, 249, 305, 1913, 251, 320, Apoth Zeit 1911, No 37; Fourneau, J. Pharm. Chim. 1904, [v1.] 20, 481, Emde, Arch Pharm 1907, 245, 662, Gadamer, ibid 1908, 246, 566; Calliess, Apoth Zeit 1910, 25, 677) According to Rabe (Ber 1911, 44, 824) the ammonium base of ephedrine decomposes into water, trimethylamine and  $\alpha$ -phenylpropylene-  $\alpha\beta$ -oxide, and he regards ephedrine and  $\psi$ -ephedrine as optically isomeric β-methylamino-a-OH CHPh CHMe NHMo phenyl-propane-a-ols (cf Emde, lc.) This view is shared by Schmidt (Arch Pharm 1914, 252, 89, 1915, 253, 52), (cf Emde,  $\bar{l}$  c.) who considers the isomerism due to the difference in the spatial arrangement of the —CHOH group Uf also Eberhard (Arch Pharm 1915, 253, 62). a-Ephedrine has been synthesised by Nagai (Eng Pat 120936) and shown to differ from Fourneau's products and to be the racemic form of the natural variety (see Chem Soc Abst 1919,

The name ephedrine has also been applied by Spehr to a crystalline alkaloid C<sub>13</sub>H<sub>19</sub>ON, m p 112°, obtained from Ephedra monostachya, a drug used as a remedy for syphilis and gout in Bessarabia and Wallachia Ordinary ephedrine is known commercially as 'ephedrine (Merck)' and the second species as 'ephedrine (Spehr)'

EPICARIN,
HO C<sub>10</sub>H<sub>6</sub> CH<sub>2</sub>—OH COOH

A derivative of β-naphthol used in the treatment of skin diseases V Synthetic drugs

EPICHLORHYDRINS α-epichlorhydrin

EPICHLORHYDRINS α-epichlorhydrin (γ-chlorpropylene oxide) CH<sub>2</sub> CH CH<sub>2</sub>Cl Prepared by treating glycerol with phosphorus pentachloride or by heating the dichlorhydrins with

EPRONAL. 637

hydrochloric acid (Berthelot, Ann Chim Phys [3] 41, 299); by treating  $\alpha$ - or  $\beta$ -dichlorhydrin with caustic potash (Reboul, Annalen, Spl. 1, 221, Tollens and Munder, Zeitsch Chem 1871, 252; D. R. P. 239077), or with alkali carbonates or alkaline earth hydroxides (D R P 246242) Glycerol is treated with hydrogen chloride at 120°-130°, and the product distilled in vacuô. The fraction boiling at 50°-120°. which contains the dichlorhydrins, is treated with strong caustic potash solution, and the The portion boiling product distilled in vacuô below 75° is washed with water and distilled. The fraction boiling at 118°-119° is crude epichlorhydrin, and may be purified by redistillation in vacuo (Fauconnier, Bull Soc chim ii 50, Mobile liquid with odour resembling that of chloroform, bp 1166° (corr), sp gr 1 2031 at 0°/4° (Thorpe, Chem Soc Trans 1880, 206), insoluble in water With sodium in ethereal solution epichlorhydrin yields allyl alcohol and the diallyl ether of glycerol (Tomor, Ber 1888, 1290, Kijner, J Russ Phys Chem Soc 24, 31), by heating with alkyl iodides, chloriodohydrin alkyl ethers are produced (Paal, Ber 1888, 2971), ammonia gas converts it into tri hlorhydroxypropylamine (Fauconnier, Compt rend. 107, 115), with potash in the presence of an alcohol, it yields the corresponding dialkyl ether of glycerol (Zunino, Atti Real Accad Lincei, 1897, ii 348, 1907, 9, i 309), with phenols and sodium ethoxide, glyceryl diaryl others are formed, whilst with sodium hydroxide, glycide aryl ethers are produced (Boyd and Marle, Chem Soc Trans 1908, 838, 1909, 1803; Zunino, Atti Real Accad Lincei, 18, i 254, Cohn and Plohn, Ber 1907, 2597, Lindemann, *ibid* 1891, 2145); concentrated nitric acid oxidises it to  $\beta$ -chloriactic acid, alcoholic ammonia and ethylsodiomalonate convert it into chlorhydroxypropylmalonamide (Traube and Lehmann, Ber 1899, 720) combines with acid chlorides to form dichlor-Haller and March have synhydrin esters thesised ketolactoric acids and other unsaturated compounds by condensing epichlorhydrin with the sodium derivatives of benzoyl acetic esters of acetone dicar boxylic ester and of acetylacetone (Compt rend 132, 1459, 136, 434, 137, 11, 1203) It has also been condensed with aniline (Fauconnier, *ibid* 106, 605, 107, 250), with *p*-toluidine (Cohn and Fiiedlander, Ber 1904, 3037), with prussic acid (Lespieau, Compt rend 127, 965), and with salicylic acid (Lange, D R P 184382; Chem. Soc Abstr 1906, i 930) Organomagnesium compounds react with it to form substituted chlorhydrins of the type

# R·CH CH2OH

(Riedel, Chem Zentr 1907, 1. 1607, D. R. P. 183361; Fourneau and Tiffeneau, Bull Soc chim. 1, [4] 1227) Epichlorhydrin has been used as a solvent for resins for photographic purposes (Phot Korr 1899, 2)
β-Epichlorhydrin (β - chlorpropylene oxide)

CH<sub>2</sub> CHCl CH<sub>2</sub>. Obtained together with the a- compound and other products by treating the allyl chloriodohydrins formed by treating alcohol with iodine monochloride, with dry sodium hydroxide (B.got, Ann. Chim Phys. 22,

[6] 468) Liquid, bp 132°-134° Phosphorus pentachloride converts it into By-dichloropropylene, and sodium amalgam into allyl alcohol

Polymerules (CaHbOCl). Obtained in the preparation of pyrazine bases by the distillation of glycerol with ammonium phosphate and ammonium chloride, by treating the high-boiling fractions with hydrochloric acid, it is obtained crystalline, mp 112°-113°, bp 232°-233°. Readily soluble in hot alcohol, sparingly soluble in water

(C<sub>3</sub>H<sub>5</sub>O(1)<sub>5</sub> Obtained by treating α-epichlorhydrin with hydrofluoric acid in a platinum dish. Yellow oil, soluble in alcohol, benzene, ether, or acetic acid. Dilute sulphune acid at 200° converts it into a substance resembling cellulose (Paterno and Olivieri, Gazz chim ital. 24, 1.

305; n 541)

Chlorbutylene oxide CAH,OCL Obtained by treating a mixture of butenylglycerol and glacial acetic acid with hydrogen chloride, b p. 125 5° (corr ) at 738 mm , sp gr 1 098 at 15°/0 (Zikes, Monatsh 6, 352)

**EPIDIORITE.** A rock consisting essentially of plagioclase-felspar and hornblende, and with a more or less marked schistose structure It has been formed by the action of pressuremetamorphism on dolerite (q v), the augite of been thereby transformed into Rocks of this class are of wide which has hornblende distribution, and being tough and hard they are much quarried for road-mending, for example, in Cornwall Auriferous quartz-veins are often associated with epidioritis LJS

EPIDOTE. Basic silicate of calcium, aluminium, and ferric iron, HCa2(Al, Fe)3S13O13, the alumina and ferric oxide replacing one another isomorphously, Fe<sub>2</sub>O<sub>3</sub> ranging from 1 6 pc (in the elinozoisite variety) to 17 pc. The monoclinic crystals are usually much elongated in the direction perpendicular to the plane of symmetry, the habit thus being prismatic. The colour ranges from green to brown, and is usually a characteristic yellowish-green resembling that of a pistachio-nut, hence the name pistacite sometimes applied to this mineral Sp gi 33 -35, H 6-7Good crystals are found at many localities, and very fine ones at the Knappenwand in the Untersulzbachthal in Salzburg These are of a deep green colour and are sometimes cut as gem stones Epidote is strongly pleochroic, and when manganese (Mn<sub>2</sub>O<sub>3</sub>) enters isomorphously into the composition this character becomes still more pronounced This is seen in the withamite variety from Glencoe, Scotland, and in the cherry-red manganese-epidote (piedmontite) from San Marcel, Piedmont. As a rock-forming mineral epidote is abundant in crystalline schists and metamorphic rocks; and it is often present in weathered igneous rocks A rock composed of epidote and quartz is known as epidosite or epidote-schist, and sometimes carries gold Banded epidosites have been used as ornamental stones. L. J. S.

EPINEPHRINE v ADRENALINE; also SYN-

EPININE. Trade name for 3 4-Dihydroxyphenylethylmethylamine See Adrenaline EPIOSINE. Trade name for 1.methyl 4 5-

diphenylene imidazole EPIRENAN & SYNTHETIC DRUGS

EPRONAL. Ethylpropylacetylurethane

EPSOM SALTS. Magnesium sulphate (v. MAGNESIUM

EPSOMITE or Epsom Salts, MgSO<sub>4</sub>.7H<sub>2</sub>O<sub>5</sub> as a mineral usually occurs as silky white fibres forming efflorescences in limestone caves, mine galleries, &c ('rystals are orthorhombic with spheroidal hemihedrism and a prismatic habit, the prism angle being nearly a right angle (89° 26') Although good crystals are readily obtained by artificial means, in nature they are of rare occurrence They have, however, been found in the salt deposits at Stassfurt in Prussia; where also a massive variety ('reichardtite') forms thin layers in carnallite. This epsomite forms thin layers in carnallite is no doubt derived by the hydration of the more abundant kieserite, MgSO<sub>4</sub>,H<sub>2</sub>O Albany Co, Wyoming, and on the border of British Columbia and Washington, there are several small lakes containing fich deposits of compact white epsomite—At Epsom in Surrey, from which place the name is derived, the salt occurs only in solution in mineral waters

LJS

ERBIUM or Neo-Erbium Sym Er At wt 167 7 A dark-grey powder, sp gr 4 77 at 15° (S Meyer, Monatsh 1899, 20, 793)

Sources - Ytterbite (gadolinite), yttrotant-

alite, euxenite, venotime

The oxide of this metal belongs to the group of yttrium earths, forming with dysprosium and thulium the erbium sub-group, the salts of which are distinguished by their colour and by their characteristic absorption spectra in solution

Separation and purification of the yttrum earths. The following methods have been employed in separating the closely related

members of the yttrium earths

1 Fractional decomposition of nitrates (Bahr and Bunsen, Annalen, 1866, 137, 1) nitrates, heated until nitrous fumes are evolved, leave a vitieous residue of basic mitrates, the solutions of which, in hot water, slowly deposit, on cooling, small pink needles containing the greater proportion of the coloured erbium earths. The crystals are extracted with 3 p.c. intric acid, the solution evaporated, the residue again heated, and the whole series of operations repeated many times. The erbium earths accumulate in the less soluble basic salts, whilst the remaining yttiium earths, and any cerum earths which may be present, accumulate in the final mother liquors, and are not readily freed completely from the erbium earths Further separation is effected by heating these , nitrates from the final mother liquors so strongly that the residue consists of oxides mixed with insoluble basic nitrates. On extraction with water, the erbium compounds remain undissolved, and the operation is repeated till the more soluble yttrium salts no longer show absorption bands

According to Wichers, Hopkins and Balke (J Amer. Chem Soc 1918, 40, 1615) the nitrate fusion method is very efficient for separating erbium and yttrium and is much better than any of the later methods, although the cobalticyamde and the nitiite precipitation methods (v supra) are both good, the latter being the more practicable and efficient

2 Differences in the solubility of the oxides. -The oxides are mixed with water and treated

with an amount of nitric acid insufficient to convert the oxides into normal nitrates The mixture becomes hot, and, after cooling, concentrated nitric acid is added until the product assumes a reddish tint. The precipitate is then digested with alcohol, which dissolves the normal nitrate, to the exclusion of the basic nitrates The latter consists chiefly of erbium salts mixed with some ytterbium and scandium compounds, the more basic yttiium and cerite metals passing into solution The final separation of erbium from ytterbium and scandium is effected by fractionally precipitating the basic nitrates by adding oxides to the solution of normal nitrates In this operation ytterbium and scandium are precipitated as basic nitrates, while cibium remains dissolved (v. Drossbach, Ber. 1902, 35,

3 Fractional precipitation by alkalis or aniline—The addition of ammonia to hot solutions of the salts of the yttnum group occasions the precipitation of basic salts which

are easily filtered

The neutral chloride dissolved in 50 p.c. alcohol is decomposed at 90° with alcoholic aniline, or the hydroxide is treated with aniline hydrochloride in aqueous solution (Prossbach, Ber 1896, 29, 2452, Mullmann and Baur, *ibid* 1900, 33, 1748; 1902, 35, 2382, Urbam, Comptrend 1901, 132, 136, Kruss, Annalen, 1891, 265, 1; Zertsch anorg ('hem 1893, 3, 108, 353, 407).

4 Fractionation with ammonium carbonate and acetic acid —The hydroxides of the yttrium group are moderately soluble in concentrated aqueous ammonium carbonate, and from this solution the earths may be fractionally precipitated by acetic acid (Dennis and Dales, J Amer Chem Soc 1902, 24, 400)

5 Fractionation of oxalates, acetales, acetylacctonates, ethylsulphates (Urbain, Ann Chim. Phys 1900, [7] 19, 184, Maic, Ber 1902, 35, 2370), formates (K A Hofmann, ibid 1908, 41, 308), ferrocyanides (Bohm, Die Darstellung der seltenen Eiden), bromates (James, J Amei Chem Soc 1908, 30, 182), chromates (Kruss and Zoose, Zeitsch anoig Chem 1893,

Various methods of separation of erbium from yttrium have been examined by Willand and James (J. Amer. Chem. Soc. 1916, 38, 1198), who find that fractional precipitation with (1) sodium nitrite, (2) with potassium cobalticyanide from a hydrochloric acid solution, (3) with sodium phosphate from a dilute nitric acid solution of the oxides on the whole yield the best results. In all three methods the last fraction was obtained by the addition of oxalic acid to the final liquois

Erbia (erbium ouide) Er<sub>2</sub>O<sub>3</sub> A pink powder obtained by heating the oxalate at 575°-845°; sp gr. 864, sp ht 0065 So prepared is apt to retain carbon dioxide

The salts of erbium have a pink or reddish colour, a sweet, astringent taste, and their solutions exhibit characteristic absorption bands in the red, green, and blue regions of the spectrum The wave lengths corresponding with the two most prominent bands are as follows: λ 522 5-523 5 and 647 5-651 5

Unless ionic dissociation is repressed, dilute solutions of the various orbium salts exhibit

dentical absorption spectra (Zeitsch. physikal. hem 1906, 56, 624; v Purvis, Proc Camb.
Phil Soc. 1903, 12, 202, 206) The absorption ands undergo displacement when the solutions re placed in a magnetic field (J Becquerel, lompt rend. 1907, 145, 1412)

(For accounts of the flame, spark, and arc mission spectra, v Thalen, ibid 1880, 91, 326, Exner and Haschek, Wien Akad Ber 1899, 108, Ia, 1071, 1123, Eder, Chem Zentr 1916, 1 242; ibid ii 455) According to Eder erbium is

robably complex.

Erbium salts exhibit signs of radio-activity

Strong, Amer Chem J. 1909, 42, 147). **Erbium sulphate** E<sub>12</sub>(SO<sub>4</sub>)<sub>3</sub>,8H<sub>2</sub>O nonoclinic crystals, separating from aqueous olution at  $20^{\circ}-25^{\circ}$ , isomorphous with the orresponding yttrium, praseodymium, and leodymium salts (Kraus, Zeitsch Cryst 1901, 4, 307)

The double sulphates Er<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>, K<sub>2</sub>SO<sub>4</sub>, 4H<sub>2</sub>O and  $\text{Er}_{2}(SO_{4})_{3}(NH_{4})_{3}SO_{4}H_{2}O_{4}$  are readily

oluble in cold water

 $\text{Et}(\text{NO}_3)_3,5\text{H}_2\text{O}$ Erbium nitrate Large eddish crystals stable in air. When heated on platinum wire, the nitrate imparts an intense reenish colour to the flame, which, when viewed hrough the spectroscope, shows bright lines in he yellow-green and orange-blue regions of the pectrum. These bright lines correspond in position with the dark bands of the absorption pectrum.

Erbium formate E<sub>1</sub>(CHO<sub>2</sub>)<sub>3</sub>,2H<sub>2</sub>O Crystal-

ises from cold aqueous solution

Erbium oxalate Er<sub>2</sub>(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub>,10H<sub>2</sub>O (K A fofmann, Ber 1910, 43, 2631) is a reddish, paringly soluble microcrystalline powder

Erbium platinocyanide E12[Pt(CN)4]3,21H2O, ike the corresponding yttiium salt, crystallises n red prisms with a green metallic reflex other salts, see Morgan and James, J Amer them Soc 1914, 36, 10, Katz and James, ibid G T M 913, 35, 872

ERECTHIDIS OIL consists of a terpene 110 II 16, which boils at 175°, and at 185° has a p gr of 0 838 This body absorbs a molecule of ydrochloric acid without the separation of a rystalline compound. The portion of the oil thich boils above 200° also consists of a terpene Beilstein and Wiegand, Beil 15, 2854)

EREPSIN v Enzymes, art FERMENTATION

ERGAMINE, ERGOSTEROL, ERGATHI-NEINE, ERGOTIN, ERGOTININE, ERGO-'OXINE v ERGOT

ERGOT OF RYE. Spurred 14e, Ergota, 3 P , USP Secale cornutum (Ergot de seigle, er; Mutterkorn, Ger) Ergot of 1ye consists f the dried sclerotium (resting stage) of the ungus Claviceps purpurca (Tulasne) developed rom the ovary of the ryo (Secale cereale) The rains are dark violet-black, 1-4 centimetres ong, slightly curved and tapering at both ends The same or a closely related fungus occurs on number of other grasses, wild and cultivated, ind on some Cyperaceae The first definite nention of ergot which the writer has been able o trace, is in the 1582 edition of Adam Lonicer's Kreuterbuch, cap ccclxx. p 285, where its ise by midwives is also referred to From an ccount by the medical faculty of Marburg, sublished in 1597, down to the middle of the

19th century, there is a voluminous literature in German and in French, on ergotism, there seems little doubt, however, that the epidemic ignis sacer of the Middle Ages was also due to the presence of ergot in flour (cf Ehlers, L'ergotisme, Paris, 1896) Only one outbreak of ergotism in England (due to wheat) has been described (Phil Trans Roy Soc 1762, 52, 530) The use of eigot by midwives was forbidden in several countries about 1770-1780, but in the second decade of the last century it was definitely introduced into scientific medicine by the American physicians Prescott and Stearns, and now ergot is only important as a drug, being chiefly employed as a stimulant of uterine muscle In the United States ergot growing on wild grasses (especially on species of Elipmus, cf. also Amberg, Schweiz Woch Chem Pharm 1911, 49, 489), has occasionally given rise to extensive poisoning of horses and cattle Further botanical and historical details are given by Fluckiger and Hanbury, p. 740. Cooke, Brit Fungi, 772, Pharm J [m ] 1, 702, Paisons, Rep US Commissioner Agriculture, 1880, 136, Holmes, Pharm J 1886, [m ] 16, 684; Kobert, Historische Studien, Halle, 1889, Carruthers, Rep. Roy. Ague. Soc. 1874, [n.] 10, 443 (with illustrations of eigot on other grasses)

The chief commercial varieties of ergot of rye come from Russia and from Spain eigot with large grains, is probably of no special value, because several writers agree that the larger grams have a smaller percentage of alkaloid than the smaller ones Although they are not official, other eigots, growing on wild grasses, deserve attention as a source of alkaloid Ey that on Mondia cardea has been found to contain as much as 0.81 p.c. Festuca ergot contains 0.34 p.c. (Pharm J. 1913, 130). Perhaps 'Diss' ergot on Ampelodesmos tenar (Link), occurring along the coast of Algeria, might thus be utilised (if Germaix et Chassang, Rev. méd et phaim de l'Afrique du Nord, Alger, 1898, 1, 45, 78, 113, 146, 184, also Holmes, Pharm J 1886, [m ] 16, 684) Diss ergot has better keeping properties than eigot of rye, about which the Brussels Conference agreed that it should not be more than one year old and should be preserved entire. The rapid deterioration of eigot seems to be due to oxidation, which is favoured by the presence of the oil. Hence removal of the latter by light petroleum has been recommended Aymonier (J Pharm Chim 1897, [vi] 6, 359) suggests coating the ergot with a dilute ethereal solution of tolu balsam

The literature on the chemistry of ergot is as voluminous as that on ergotism first investigation is recorded by J G Model in his Chymische Nebenstunden (Wittenberg, 1771), seven years after von Munchhausen had first recognised the drug as a fungus Among older writers we need, however, only mention here Tanret (Compt rend 1875, 81, 896), who first obtained the crystalline alkaloid ergotinine, and Buchheim (Aich. exp. Path Pharm 1874, 3, 1) who, albeit without experimental evidence, expressed the view that the activity of ergot is due to decomposition products of protein. The active principles of ergot have only been obtained pure in recent years. They are a complex alkaloid and some simpler bases

Ergotoxine  $\mathrm{C}_{35}\mathrm{H}_{41}\mathrm{O}_6\mathrm{N}_8$ , mp 162°-164°, an amorphous alkaloid yielding crystalline salts, of which the phosphate  $\mathrm{C}_{35}\mathrm{H}_{41}\mathrm{O}_6\mathrm{N}_5$ ,  $\mathrm{H}_3\mathrm{PO}_4\mathrm{H}_2\mathrm{O}$ , m.p. 187°-188°, is the most readily obtainable (Barger and Carr, Chem. News, 1906, 94, 89, Trans Chem. Soc. 1907, 91, 337, Barger and Ewins, ibid 1910, 97, 284; 1918, 113, 235). This alkaloid had been obtained in an impure form as 'amorphous ergotimne' by Tanret, and was termed hydroergotimne by Kraft (Arch. Pharm 1906, 244, 336, 1907, 245, 644), who, without analyses, first recognised it as a hydrate of Tanret's ergotimne

The alkaloids of ergot are best obtained on a small scale by extracting the powdered drug with ether and shaking the ethereal solution with a weak acid (citie, tartaric) Excess of strongly ionised mineral acids precipitates the salts of the ergot alkaloids, which, on account of their high molecular weight, are semi-colloids. This property led many of the earlier investigators into error, but may be put to account when working on the large scale. The residue left on evaporation of an alcoholic tincture of ergot is extracted with light petroleum, to remove the fat. It is then dissolved in ethyl acetate and shaken with citric acid, sodium bromide solution then precipitates the mixed hydrobromides almost quantitatively. The latter are dissolved in dilute caustic soda, from which ether removes only ergotinine The caustic soda solution is then neutralised, made alkaline with sodium carbonate and extracted again with ether The crude ergotoxine left on evaporation, together with that from the mother liquois of the crystallisation of ergotinine (see below), is dissolved in 80 p.c. alcohol and a slight excess of syrupy phosphoric acid in alcohol is added On standing for a few days the ergotoxine phosphate crystallises in thin needles, which on recrystallisation from 50 parts of 90 pc alcohol may be transformed into broad plates They are soluble in 313 parts of cold and 14 parts of boiling 90 p c alcohol By shaking with cold distilled water a 1 pc solution is obtainable, which is strongly opalescent and sets to a thick jelly on the addition of electrolytes. For physiological experiments ergotoxine is best dissolved in dilute sodium hydroxide

Ergotinine, although not physiologically active, may be considered here. The alkaloid left on evaporation of its ethereal solution (above) readily crystallises from absolute alcohol in needles, mp  $229^{\circ}$  (corr.), and having the composition  $C_{3b}H_{39}O_5N_5$ . It is extremely soluble in cold chloroform, and dissolves in 292 parts of absolute alcohol at  $18^{\circ}$ , in alcohol  $[\alpha]_D = +338^{\circ}$ ; the solutions are strongly fluorescent. Prolonged boiling with alcohol lowers the rotation and produces an amorphous decomposition product. Heating in alcoholic solution with slightly more than one equivalent of (phosphoric) acid transforms ergotinine into ergotoxine phosphate which may be crystallised. The same change takes place slowly in cold dilute acetic acid. Conversely ergotoxine is dehydrated to ergotinine by boiling with methyl alcohol or by warming with acetic anhydride.

Both alkaloids give the following colour

reaction (Tanret). A solution of 1 mg. of the alkaloid in ether or in ethyl acetate is allowed to evaporate on the surface of concentrated sulphuric acid (containing oxides of nitrogen). A transitory orange colour is developed changing to violet-blue Keller (Schweiz Woch Chem. Pharm. 1894, 32, 121) has improved this by dissolving 2-3 mg of the solid alkaloid in sulphuric acid on a watch glass and stirring in a trace of anhydrous ferric chloride, when an orange colour is produced, changing to deep red and then becoming blue or green at the edges Alternatively the solution of the alkaloid in glacial acetic acid may be poured on sulphuric acid, without mixing The test yields a positive result with the crude alkaloid from I gram of ergot

Both alkaloids are sensitive to general alkaloidal precipitants, potassium mercuric iodide giving a faint opalescence with 1 part of alkaloid in 1-2 millions of water. On dry distillation both alkaloids yield a crystalline sublimate in isobutyryl formamide CH(CH<sub>3</sub>)<sub>2</sub>CO CO NH<sub>2</sub> (Barger and Ewins, Ic) Ergotoxine probably contains a carboxyl group and ergotinine is the corresponding lactone. Both contain one N-methyl group, but no O-methyl group (Barger and Ewins, Chem. Soc. Trans. 1918, 113, 235). Ergotimine does not yield any crystalline salts.

The total alkaloids of ergot (i.e. ergotoxino +ergotinine) may be estimated by exhausting the finely powdered drug with light petroleum, mixing it with magnesia, extracting with ether and shaking the ether with dilute acid. After making alkaline with ammonia the alkaloids are again shaken into other, and estimated gravimetrically (Keller, l.e.). For details and results of analysis, sie Allen's Comm. Organic Analysis, 1913, vol. vi. p. 21. The alkaloidal content varies roughly from 0.1 to 0.4 p.c., and in Russian and Spanish ergot is generally between 0.2 and 0.3 p.c. Konig (Apot Zeitsch 1912, 27, 879) found 10 samples of German ergot to contain 0.032-0.14 p.c. An approximate idea of the ergotoxine content may be obtained by physiological means (Barger and Dale, Bio-Chem. J. 1907, 2, 240)

In addition to a specific alkaloid, the active principles of ergot comprise a number of simpler bases, derived from the amino-acids of protein by loss of carbon dioxide. Of these \$\beta\$-iminazo-lylethylamine (from histidine) and \$p\$-hydroxy-phenylethylamine (from tyrosine) are therapeutically the most important. They can be prepared synthetically and are also formed by putrefaction. Their physiological action has been much studied of late and they have been applied practically.

Histamine (β-iminazolylethylamine 'β-1,' 4-β-aminoethylglyoxaline, ergamine)

$$\begin{array}{c} \text{CH=N} \\ | \\ \text{NH\cdotCH} \end{array} \hspace{-0.5cm} \text{CCH}_2 \hspace{0.5cm} \text{CH}_2 \hspace{0.5cm} \text{NH}_2$$

was obtained first synthetically by Windaus and Vogt (Ber 1907, 70, 3691) and then by Ackermann (Zeitsch. physiol. Chem. 1910, 65, 504) by the putrefaction of histidine Its intense physiological activity was not recognised until it was isolated from a commercial dialysed ergot preparation (Barger and Daie, Chem. Soc. Trans.

1910, 97, 2592). Kutscher (Zentralb Physiol 1910, 24, 163) isolated it simultaneously from ergot, but did not correctly identify it substance is certainly present in eigot as such, but its amount may be increased by such processes as prolonged extraction and dialysis. The base can be obtained directly from histidine by putrefaction (D R. PP 2528724, 256116, Hoffmann, La Roche & Co, D. R P 250110, F. Bayer & Co; the yield is inconstant but may reach 60 pc or more) It may also be made from histidine by heating with acids (Ewins and Pyman, Trans Chem Soc 1911, 99, 339, yield 25 p.c ) It is best obtained by reduction of the nitrile of iminazolylpropionic acid (Pyman, Trans Chem Soc. 1911, 99, 668), which nitrile in its turn is obtained most conveniently from histidine by the action of chloramine-T (Dakin, Biochem J 1916, 10, 322), although it may also be synthesised, according to Pyman (lc; cf also Koessler and Hanke, J Amer ('hem Soc 1918, 40, 1706) The most important salts are the *phosphate* C<sub>5</sub>H<sub>9</sub>N<sub>3</sub> 2H<sub>3</sub>PO<sub>4</sub> forming large crystals, and the dipicrate

### $C_5H_9N_3(C_6H_3O_7N_3)_9$

yellow rhombic leaflets, mp  $238^{\circ}-242^{\circ}$  (corr) The free base is readily soluble in water, its solutions give Pauly's histoline reaction with sodium p-diazobenzene sulphonate; a very distinct rose-pink colouration is still obtainable at  $1\ 10,000$ 

**Tyramine** (p-Hydroxyphenylethylamine) OH C<sub>8</sub>H<sub>4</sub> ('H<sub>2</sub> ('H<sub>2</sub>NH<sub>2</sub> is similarly derived from tyrosine by loss of carbon dioxide. On a small scale it can be obtained by heating tyrosine to 260°-270° under reduced pressure (Schmitt and Nasse, Annalen, 1865, 133, 211, Ehrlich and Pistschimuka, Ber 1912, 45, 1006, thus obtained a 50 pc yield) On a larger scale it is best got by reduction of p-hydroxybenzylcyanide according to Barger (Trans Chem Soc 1909, 95, 1123), who first isolated It also occurs as a product of it from ergot putrefaction (Barger and Walpole, J. Physiol 1909, 38, 343). The free base melts at 161° and boils at 175°-180° and 8 mm. It dissolves in 95 parts of water at 15° and can be extracted from aqueous solutions by shaking many times with ether, more readily by amyl alcohol gives Millon's reaction, the most characteristic derivative is the dibenzoyl compound C<sub>22</sub>H<sub>19</sub>O<sub>3</sub>N, m p 170°

### Acetylcholine

#### OH(CH<sub>3</sub>)<sub>3</sub>N CH<sub>2</sub> CH<sub>2</sub>O OC CH<sub>3</sub>

occurs in ergot and ergot extracts in varying small amounts. It was isolated by Ewins (Biochem J 1914, 8, 44) from a BP liquid extract. It is hydrolysed almost at once by rendering its solution alkaline with caustic soda.

Physiological action of the active principles— Ergotinine is inert, but ergotoxine causes, on intravenous or intramuscular injection, a prolonged rise of blood pressure and contraction of the pregnant uterus. Repeated administration leads to dry gangrene, eg of the cock's comb, which is coloured purplish black by small doses, a reaction used at one time for the testing of ergot preparations (for criticism v. Cronyn and Henderson, J Pharm exp. Therap 1909 1, 231). Ergotoxine is the poisonous principle Vol. II.—T.

responsible for epidemic ergotism (Kobert attributed the two types of this disease to the alkaloid cornutine and the resin sphacelinic acid, but both these substances seem to have been ergotoxine in an impure or modified form: cf Barger and Dale, Blochem. J., 1907, 2, 240). \$\beta\$-Immazolylethylamine is one of the most potent of drugs, causing marked contraction of the isolated guinea-pig's uterus in a bath containing 1 25,000,000 p-Hydroxyphenylethylamine resembles adrenaline in action, as in structure, but is much less potent; it causes a rather transitory rise of blood pressure Acetylcholine is not a therapeutically desirable constituent It has a muscarine-like action, inhibiting the heart, and stimulates intestinal muscle (Hunt and Taveau, Hygienic Bulletin, Washington, No 73, 1911) In most ergot pieparations the amount is negligible For a detailed account of the last three and most of the following bases, see Barger, the Simpler Natural Bases, Long. mans, 1914

In addition to  $\beta$ -immazolylethylamine, phydroxyphenylethylamine and acetylcholine, there have been found in ergot quite a number of other sulstances of bio-chemical interest, which are not peculiar to the drug. They are which are not peculiar to the drug mert, or have only a slight physiological action Putrescine and Cadaverine (Rielander, Sitzber. Ges Naturw Marburg, No 7, 1908), isoamyl-Physiol Soc amine (Barger and Dale, Proc May 15, 1909), and agmatine (Engeland and Kutscher, Zentralbl. Physiol 1910, 24, 479) are all amines, derived respectively from ornithine, lysine, leucine and arginine, by decarboxylation Agmatine has an action on the uterus like that of B-immazolylethylamine, but very much feebler It is guanidylbutylamine

# NH, C(NH) NH CH, CH, CH, CH, NH,

and was first discovered in herring roe by Kossel (Zeitsch physiol Chem 1910, 66, 257). A few amino-acids have also been found as such leucine (Buchheim, lc), asparite acid (Bargei and Dale, Biochem J 1907, 2, 240), leucine, isoleucine, valine (van Slyke, J Pharmac exp Therap 1909, 1, 265). A mixture of these was described as an active principle, 'clavin' by Vahlen (Acrh exp Pharm Path 1906, 55, 131). There are further present choline which gives rise to the odour of trimethylamine, when ergot is treated with sodium hydroxide (Brieger, Zeitsch physiol Chem 1887, 11, 184), (for acetylcholine, see above), betaine and uracid (Rielander, lc), vernine C<sub>16</sub>H<sub>20</sub>O<sub>8</sub>N<sub>8</sub>,3H<sub>2</sub>O, which also occurs in leguminous seedlings (Schulze and Bosshard, Zeitsch physiol Chem 1887, 10, 80), lactic acid, mannitol, trehalose, phytosterols and fats (see below)

Inert constituents which so far have not been found elsewhere are secale ammonulphonic acid  $C_{16}H_{24}O_{16}(NH_2)SO_3H$  (Kraft, lc), prisms, mp 200°, soluble in water giving an acid solution; ergothnoneine  $C_9H_{16}O_2N_3S_2H_2O$ , isolated by Tanret (J Pharm Chim. 1909, [vi] 30, 145) in an amount of 0 l pc and shown by Barger and Ewins (Trans Chem Soc 1911, 99, 2336) to be the trimethylbetaine of thiolistidine; secalonic acid  $C_{14}H_{14}O_6$ , a crystalline yellow colouring matter, mp. 244°, present to the extent of about 2 pc (Kraft, lc) This substance is probably the lactone of an hydroxy

acid, and possibly identical with sclerocrystallin Path. Pharm 1877, 6, 174). Another crystalline yellow colouring matter  $C_{15}H_{14}O_{7},H_{1}O_{7},m_{1}O_{1}$ . 350°, after regeneration of a crystalline acetyl derivative, has been described by Freeborn (Pharm. J. 1912, [iv.] 34, 568) The most important colouring matter of ergot, sclererythrin, is not known in a pure condition 18 the cause of the deep purplish red colour of the outer cells of the grain and is valuable for the recognition of ergot in flour, &c. substance is acidic, and apparently present as a salt, for it is not extracted by other until the ergot has been moistened with water containing It is soluble in aqueous alkalis tartaric acid with a red colour. Clavicepsin C18H34O16, m p 198° when anhydrous,  $[\alpha]_D^{20} = +142 27^\circ$ , was extracted from ergot by Marino-Zuco and Pasquero (Gazz chim ital 1910, 41, 368) It is hydrolysed by acids to two molecules of glucose and one of mannitol The oil is quantitatively the most important constituent (mostly 30-33 p.c.) and has been examined by Rathje (Arch Pharm 1908, 246, 692), who found D. = 0.9250,  $n_D = 1.4685$ saponification number 179 3, acid number 11 38, 10dine number 740, Hehner number 96 25, Reichert-Meissl number 0 63 and true acetyl number 27 44 The percentage composition is oleic acid 68, hydroxyoleic acid 22, palmitic acid 5, unsaponifiable matter 0 35, morganic matter 0 2, alkaloid 0 6, and glycerol 7 5

Ergosterol C<sub>27</sub>H<sub>42</sub>O,H<sub>2</sub>O, m p 165°, and fungsterol C<sub>28</sub>H<sub>40</sub>O,H<sub>2</sub>O, m p 144°, have been separated by Tanret (Compt rend 1908, 147, 75). The second at least seems to exist in other

fungi.

The detection of ergot in flour is based on the presence of alkaloids and of sclererythrin, and on the fact that the ergot particles float when the flour is shaken with an alcoholchloroform mixture of density 1 435 After pouring off, alcohol is added, when the ergot sinks and can be examined microscopically For further details, see Allen's Commercial Organic Analysis, vol vii, 1913, p. 23, and Marino-Zuco and Duccini (Gazz. chim. ital. 1914, 44, 437)

For a critical discussion of the ergot preparations of the British Pharmacopæia, see Carr and Dale (Pharm J 1913, [iv.] 37, 130) It should be noted that 'ergotin' is not a pure substance, but a purified soft extract The varying activity of different extracts should be controlled by physiological test

ERGÖTINE v. ERGOT OF RYE. ERGOTININE v ERGOT OF RYE.

ERICA B v PRIMULINE

ERICIN. Trade name for salicylic methoxy-

methyl ester C<sub>6</sub>H<sub>4</sub><0H OCH<sub>2</sub> OCH<sub>3</sub>

ERICOL. Guaracol acetate. ERICOLIN is found in several plants belonging to the order Ericacea, especially in the herb of the marsh wild rosemary, Ledum palustre (Linn) (Rochleder and Schwarz, Annalen, 84, 368), less abundantly in the common heath or ling, Calluna vulgaris (Salisb) (Rochleder, ibid 354), in Rhododendron ferrugineum (Linn ) (Schwarz, ibid. 361), and in the red bearberry, Arctostaphylos Uva-ursi (Spreng) (Kawalier, Sitz. W. 9, 29).

To prepare it from L palustre, the chopped (Dragendorff and Podwyssotzki, Arch. exp. leaves are thrown into boiling water and boiled for several hours. The filtrate obtained after adding basic lead acetate to the liquid is evaporated in a retort to one-third of its bulk Thal's modification of this process (Ber. 16, 1502), the liquid is concentrated in vacuo, the separated lead salt filtered off, and the liquid freed from lead by hydrogen sulphide. The filtrate is then evaporated to a syrup, and the ericolin dissolved out by anhydrous ether alcohol. The substance left on evaporating this solvent is repeatedly dissolved in a mixture of ether and alcohol till it no longer leaves any residue (Rochleder and Schwarz).

On heating the mother liquor obtained in the preparation of arbutin from the leaves of red bearberry with dilute hydrochloric or sulphuric acid, ericolin is deposited as a resinous precipitate, which may be purified by solution in alcohol and precipitation by water (Kawalier)

Rochleder and Schwarz, after deducting 10 6 p.c ash, found erreolin to contain 51.71 p c carbon and 7 19 hydrogen, whence they deduced its formula to be  $C_{34}H_{56}O_{31}$  (51 00 C and 7 00 H).

Thal (lc), after separating an admixed substance (containing 57.5 pc carbor and 7.29 hydrogen), which was insoluble in ether alcohol, from ericolin, assigned to the latter body the formula C<sub>28</sub>H<sub>24</sub>O<sub>3</sub>, but subsequently altered it to C<sub>26</sub>H<sub>30</sub>O<sub>3</sub> to explain the decomposition equations of the substance. Ericolin, as prepared by Thal, is a brown-yellow, odouiless, hygroscopic, bitter substance decomposing even below 100°. It is decomposed by water. By extracting successively with benzene, chloroform, and a mixture of ether and alcohol, different substances go into solution; it appears that the cricolin is thus broken up into a sugar and ericinol With dilute sulphuric acid warmed to 100°, a sugar and a resinous brown yellow viscid mass, hydroericinol, are produced. An analysis of the latter body gave numbers corresponding with the formula  $x(C_6H_{12}O_2)$ , subsequently modified to  $x(C_5H_{10}O_2)$ .

The equations representing these decompositions are as follows :-

$$\begin{array}{c} {\rm C_{28}H_{30}O_3 + 4H_2O = C_8H_{12}O_8 + C_{20}H_{28}O} \\ {\rm Ericolm} & {\rm Ericinol.} \\ {\rm C_{20}H_{26}O + 7H_2O = 2(C_{10}H_{20}O_4)} \\ {\rm Hydroericinol.} \end{array}$$

Power and Tutin, however, could not isolate ericolin by Thal's process, and they found that ericinol was identical with furfural (Chem. Zentr 1907, 1 916)

Thal also isolated ericolin from Calluna vulgaris (Salisb), and recognised it by the smell of ericinol in many Ericacea and Vacciniea.

Kanger extracted ericolin from the leaves of the cowberry (Vaccinium vitis idea) by means of ether alcohol. It was then purified by precipi-tating foreign substances with lead acetate, and removing excess of the latter with sulphuretted hydrogen The substance so obtained was of a resinous nature, completely soluble in ether alcohol, and corresponded with the composition  $C_{16}H_{25}O_{10}$  (Chem. Zeit. 1903, 27, 794). Ericolin is probably not a chemical individual, but consists of a mixture of glucoside-like substances.

Ericolin is poisonous towards many microorganisms, particularly hose belonging to the colon group, but has no action on acid-fast bacilli. Twort (Proc. Roy. Soc. 1909, B. 81, 248) has used it as a medium for the isolation and culture of human tuberculous bacilli from sputum.

ERIGERON CANADENSE is an oil consisting of a terpene boiling at 176°, which, when pure, has sp.gr. 0 850-0 870, and an optical rotation +52°. It is soluble in an equal volume of alcohol, and rapidly resimfies The commercial oil has sp.gr. 0 8549-0 8963, optical rotation, +28° 48′ to 84 20°; and b.p. 172°-178° (Kepler and Pancoast, Amer. J. Pharm. 75, 216).

According to Rabak (Pharm Review, 1905, 23, 81; 1906, 326), the oil obtained from the fresh herb (0 66 p.c.) is of a bright-yellow colour with a peculiar odour like that of caraway. On exposure to air, it deposits crystals, and a brownish resinous mass is formed. The oil from the dried herb (0 26 p.c.) was darker in colour, had a powerful aromatic odour, and gave no crystals when exposed to air. It also differed from the fresh herb oil in sp. gr., optical rotation and other characteristics. In all probability, the terpeneol found in erigeron oil is a decomposition product. The oil absorbs two molecules of h, drochloric acid to form a solid dihydrochloride C<sub>10</sub>H<sub>16</sub>.2HCl, melting at 47°-48° (Beilstein and Wiegand, Ber. 15, 2854, Wallach, Annalen, 227, 292)

ERINOID. See GALALITH

ERIOCOCCUS CORIACEUS. This is a scale insect consisting of a small pinkish white sac occurring in clusters, and which encase the stems of young shoots of the Eucalypti, more especially those on old or burnt stumps. These insects contain a colouring matter which dyes wool mordanted with chromium, aluminium, tin, and iron, brown, light amber, pale orange-brown and sage-green shades respectively. The colours thus obtained are not specially brilliant, and are not fast to alkalis (Gurney, Report of the Australasian Association for the Advancement of Science, 1898).

A. G. P.

ERIOGLAUCINE. The trade name of the dyestuff obtained by condensing benzaldehydeosulphonic acid with ethylbenzylanihnesulphonic acid (J Soc Chem Ind of Basle, Eng Pat 25128; Gnehm and Schule, Annalen, 299, 347)

ERODIAMINE C<sub>10</sub>H<sub>17</sub>ON<sub>3</sub>, light yellow plates, m p 287°, and rutaecarpine C<sub>18</sub>H<sub>17</sub>ON<sub>3</sub>, light yellow needles, m p 258°, occur in the fruit of Erodia rutaecarpa (Asahina and Kashiwati J. Pharm. Soc Japan, 1915, No 405, 1293; Asahina and Mayeda, ibid. 1916, No 416, 871; Amer Chem Soc. Abstr. 1916, 607, 1917, 332).

The annexed constitutional formulæ have been assigned. Perhaps they are not fully established, but they are of considerable interest as showing that the alkaloids are derivatives of

indole; they also show a relationship to harmaline and harmine. On boiling with alcoholic hydrogen chloride erodiamine (I) is converted into isoerodiamine  $C_{19}H_{17}ON_3,H_2O$ , in p 155°–156°, and this is decomposed by alcoholic potash into carbon dioxide, N-methylanthianilic acid and a base  $C_{10}H_{12}N_2$ , in p 120°, which has been identified as  $2-\beta$ -aminoethyl indole. Isoerodiamine is regarded as

$$\begin{array}{c|c} \operatorname{HN} \operatorname{C}_6\operatorname{H}_4\operatorname{CH} : \operatorname{C} \operatorname{CH}_2\operatorname{CH}_2 \operatorname{N} \operatorname{CO} \operatorname{C}_6\operatorname{H}_4\operatorname{N(Me)} \operatorname{CHOH} \\ & & \\ \end{array}$$

Rutaecarpine (II) is similar to erodiamine, but more soluble in alcohol, hot potassium hydroxide in ammonium hydroxide yields anthramlic acid, and an unidentified product, which on fusion with potassium hydroxide, like the base  $C_{10}H_{12}N_2$  from erodiamine, yields 2-indole carboxylic acid G. B.

ERUBESCITE v. BORNITE

ERUCIC ACID C22H42O2 A homologue of acrylic acid obtained, together with oleic acid, by the saponification of the fixed oil of white mustard. It was found together with beheme acid by Goldschmiedt (Sitz W 70, 451) in the oil expressed from black mustard, and by Fitz (Ber 1871, 442) in the fat oil of grape seeds — It is best obtained by saponifying rape oil (from Brassica campestris, Linn ) with alcoholic potash distilling off the alcohol and dissolving the acid liberated on the addition of sulphuric acid in three times its volume of 95 pc alcohol, on cooling to 0°, crystals of erucic acid separate out in an almost pure condition (Reimer and Will, Ber 1886, 3320). It may be obtained by saponifying colza oil, freeing the erucic acid from arachidic acid by treatment with glacial acetic acid, in which the latter is insoluble (Ponzio, Gazz. chim ital 1904, ii 50) It is also present in cod liver oil (Bull Ber 1906, 3570)

Erucic acid crystallises in colourless needles, mp 34°; bp 281° at 30 mm Heated with phosphorus oxychloride, it forms an anhydride  $\rm C_{44}H_{82}O_3$ , an oil which may be crystallised in a freezing mixture By treatment with nitrous acid, or better, by warming erucic acid with dilute nitric acid to the melting-point and then adding sodium nitrite, brassidic acid (brassic acid) is produced. This acid stands to crucic acid in the same relation as elaidic to oleic acid Cf Mascarelli, Atti R Accad Inneel, 1917, [v] 26, 1 71 Fused with potash, erucic acid yields aradic (arachidic) and acetic acids Concentrated nitric acid converts erucic acid into brassilic acid (C<sub>13</sub>H<sub>24</sub>O<sub>4</sub>), pelargonic acid, arachidic acid, and a small quantity of dinitiononane (Fileti and Ponzio, J. pr. Chem [2] 48, 323), and concentrated sulphuric acid into behenolactone (Shukow and Schestakow, Chem. Zentr 1908, With hydriodic acid and amorphous phosphorus, it is converted into behenic acid (Goldschmiedt, J. 1876, 579) On treatment with phosphorus triodide, iodobehenic acid is obtained, which on treatment with alcoholic potash, yields iso-erucic acid, m p 54° (Ponzio, lc) Erucic acid is converted into behenic acid by heating with iodine in a sealed tube for 4 hours at 270° (Reychler, Bull Soc chim [3] 1, 296). When bromine is added to erucic acid under water, a dibromide C22H42Br2O2 is produced. It crystallises from alcohol in small white nodular crystals, mp 42°-43°, and is reduced to erucic acid by sodium amalgam in alcoholic solution. Its barium and lead salts (C<sub>22</sub>H<sub>41</sub>Br<sub>2</sub>O<sub>2</sub>)<sub>2</sub>Ba and (C<sub>22</sub>H<sub>41</sub>Br<sub>2</sub>O<sub>2</sub>)<sub>2</sub>Pb are white, viscid, easily decomposable precipitates (Otto, Annalen, 127, 182, 135, 226). Erucic acid dibromide heated in scaled tubes to about 145° for 7–8 hours, with 4 or 5 mols potassium hydroxide in alcoholic solution, yields the potassium salts of behenolic acid

$$C_{22}H_{40}O_2(=C_{22}H_{42}Br_2O_2-2HBr)$$

(Holt, Ber 1892, 961). Behenolic acid, on treatment with zinc-dust and acetic acid, yields brassidic acid and not crucic acid. At ordinary temperatures, only 1 mol of hydrobromic acid is abstracted by potash in alcoholic solution, monobromerucic acid  $\rm C_{22}H_{41}BrO_2$  is formed, with rise of temperature and separation of potassium bromide. On diluting the solution with water and adding hydrochloric acid, the acid separates as an oil which soon solidifies. It melts at 33°-34° and remains liquid for a long time. It is heavier than water, and insoluble therein, but dissolves readily in alcohol and ether. It unites with bromine, forming a dibromide  $\rm C_{22}H_{41}BrO_2^{\circ}Br_2$ , which melts at 31°-32°, and forms amorphous viscid salts

When erucic acid dibromide is triturated with water and an excess of silver oxide, and heated untilit turns brown and evolves hydrogen bromide, the mass, on boiling with water, yields a yellow oil which is a mixture of liquid hydroxyerucic acid  $C_{22}H_{42}O_3$  and dihydroxybehenic acid  $C_{22}H_{44}O_4$ The latter partially crystallises out on long standing For complete separation, the washed oil is saponified with baryta water; the precipitate is exhausted with ether and the dissolved barium hydroxyerucate is decomposed by hydrochloric acid. The hydroxyerucic acid, which separates as a colourless oil, is purified from alcohol It is viscid, lighter than water, insoluble therein, but miscible in all proportions with alcohol and other. Its salts  $C_{22}H_{41}O_{4}M$ are amorphous All excepting those of the alkalı metals and barıum are insoluble in water. Hydroxyerucic acid may also be formed from monobromerucic acid by the action of silver oxide Boiled with potash lye, it is converted into dihydroxybeheme and (Haussknecht, Annalen, 163, 40) Dihydroxybeheme and, mp 1322-133°, is also obtained by the oxidation of crucic acid with potassium permanganate (Hazura and Grussner, Monatsh 9, 947) Sodium erucate C<sub>22</sub>H<sub>41</sub>NaO<sub>2</sub> is soluble in alcohol, and gives precipitates of lead and barium erucates on the addition of an alcoholic solution of the acetates of these metals The silver salt is a curdy precipitate, which becomes coloured

nexposure to light.

Ethyl erucate  $C_{12}H_{41}(C_{2}H_{5})O_{2}$  is a colourless oil boiling above 360° without decomposition. The amide  $C_{22}H_{41}O(NH_{2})$ , mp 84° (Reimer and Will, Ic), mp. 78°-79° (Krafft and Tritschier, Ber. 1900, 3580), and the annide, mp. 55°, are crystalline bodies readily soluble in ether or

benzene, sparingly soluble in alcohol.

ERUCIN. A crystalline body said to exist in white mustard seed. It is insoluble in water, sparingly soluble in alcohol, easily in ether or in oils (Simon, Pogg. Ann. 64, 593)

n ols (Simon, Pogg Ann 64, 593)

Dierucin C<sub>3</sub>H<sub>5</sub>OH(C<sub>22</sub>H<sub>41</sub>O<sub>2)2</sub> When rape oll is allowed to stand for a long time, a yellowish

tallow-like deposit is found frequently in the casks. This, by repeated solution in ether and subsequent addition of alcohol, may be obtained in silky needles. Dierucin melts at 47° and is readily soluble in ether and light petroleum, insoluble in cold but soluble in hot alcohol. A trierucin could not be separated from rape oil, but was obtained by heating dierucin and erucic acid at 300°, mp. 31° (Reimer and Will, lc) (v. Oils and Fats)

ERVASIN. Trade name for acetyl p-creso-

ERYTHRENE. See BUTINENE.

**ERYTHRIN**, Erythric acid, Erythrinic acid, Erythrolecanoric acid  $C_{20}H_{22}O_{10}, H_2O$ , is a constituent of most lichens from which archil is prepared 1t was discovered by Heeren (Schweigger's Jour f Chem. 59, 513) in Rocella tinctoria (D C), from which lichen and several others of the same genus it may be extracted by boiling water, or better with milk of lime (cf. also Kane, Stenhouse, and Hesse (l c); Schunck, Annalen, 61, 69; De Luynes, Annales de Chim [4] 2, 385; Menschutkin, Bull Soc chim. [2] 2, 424)

The method adopted by Stenhouse to prepare this substance from the Rocella fucifirms is as follows (Annalen, 68, 72, and 149, 290). Three pounds of the lichen are mace lated for 20 minutes in a milk of lime made by shaking ½ lb of lime in 3 gallons of water, and the product filtered by means of a bag filter. The clear liquid, as it passes through, is immediately precipitated with hydrochloric acid, as prolonged contact with the lime decomposes part of the crythrin. The crude crythrin collected on bag filters is freed from acid and calcium chloride by stirring it up once or twice with a considerable quantity of water and again collecting

Erythrm forms small colourless needles, melting at 148° (Hesse, Ber. 1904, 37, 4693), and not at 164°, as stated by Ronceray (Chem. Zentr 1904, n 1504), and is sparingly soluble in water, easily soluble in alcohol, ether, and alkalis Alcoholic ferric chloride gives a purple-violet colouration, which passes to brownish-red on addition of excess of the reagent

When boiled with water, crythrin gives (a) orsellinic acid and (b) picro-erythrin (Schunck,

Annalen, 61, 65)

 $\mathrm{C_{20}H_{22}O_{10}+H_2O}=(a)~\mathrm{C_8H_8O_4}+(b)~\mathrm{C_{12}H_{16}O_7}$  whereas by boiling with alcohol, picroerythrin and orsellinic acid ethyl ester (c) are produced (Heeren, l~c; Kane, Annalen, 39, 25)

 $C_{20}H_{22}O_{10} + C_2H_5OH = (c) C_8H_7(C_2H_5)O_4 + C_{12}H_{16}O_7$ 

With amyl alcohol, a similar reaction occurs with formation of (d) amylorsellinate and picroerythrin (Hesse, Annalen, 139, 22)

 $C_{20}H_{22}O_{10}+C_5H_{12}O=(d)$   $C_8H_7O_4$   $C_5H_{11}+C_{12}H_{16}O_7$ On the other hand, excess of lime water (Lamparter, Annalen, 134, 255) gives orcin,  $CO_2$ , and erythrite

$$C_{20}H_{22}O_{10} + 2H_2O = 2C_7H_8O_2 + CO_2 + C_4H_{10}O_4$$

Tribromerythrin  $2\mathrm{C}_{20}\mathrm{H}_{19}\mathrm{Br}_3\mathrm{O}_{10},3\mathrm{H}_2\mathrm{O}$  (Hesse, Annalen, 117, 310; and *ibid* 139, 32) consists of white crystalline spherules which cake together a little above  $100^\circ$  and melt at  $139^\circ$ . Boiled with alcohol, it is resolved into *ethyldibromorsellinate* and *brompicrocrythrin*.

Picroerythrin or erythro-orsellinic acid C<sub>19</sub>H<sub>14</sub>O<sub>7,3</sub>H<sub>9</sub>O

forms colourless prisms, melts at 158°, is readily soluble in hot water, and gives with ferric chloride a purple-violet colouration. It appears to be most readily obtained pure by Hesse's method of boiling erythrin with amyl alcohol (*l c*)

Digestion with hot baryta water converts pieroerythrin into CO<sub>2</sub>, orcin, and erythrite

$$C_{12}H_{16}O_7 + H_2O = CO_2 + C_7H_8O_2 + C_4H_{10}O_1$$

The experiments of de Luynes, Menschutkin, and Hesse have confirmed the view first put forward by Berthelot, that erythrin is the monolecanoric ester (1), and picroerythrin the monoorsellinic ester of erythrite (2).

According to Hesse (Ber. 1904, 37, 4693), erythrin contains a free carboxyl group, and must therefore be represented as follows—

$$\begin{smallmatrix} \mathrm{C_4H_6(OH)_3 \ O \ C_6H_2(OH)CO \ O \ C_6H_2(OH)COOH} \\ \mathrm{CH} \end{smallmatrix}$$

whereas picroerythrin has the formula

On the other hand, Zerner (Monatsch , 1914, 35, 1021) considers that the structure of erythem is best represented thus

 $\beta$ -Erythrin  $\mathrm{C}_{21}\mathrm{H}_{24}\mathrm{O}_{10}$  is prepared by the action of lime water on some varieties of the R fuciformis (Menschutkin,  $l\,c$ ). It is a white crystalline powder; in p. 115°–116° (Lamparter,  $l\,c$ ), nearly insoluble in water, soluble in alcohol and ether with violent evolution of  $\mathrm{CO}_2$ . Boiled with water, orsellinic acid, and  $\beta$ -picrocrythrin are produced, with alcohol, the latter body together with othylorsellinate

β-Picroerythrin  $C_{13}H_{16}O_{10}$  crystallises in needles, very soluble in water and alcohol Boiled with baryta water, it is resolved into  $CO_2$ , erythrite, and β-orcinol (Menschutkin)

According to Hesse (lc), it is probable that  $\beta$ -erythrin is not a chemical individual, but consists mainly of erythin, the properties of which are modified by the accompanying impurities.

A G. P

ERYTHRITE or Cobalt-bloom. A mineral consisting of hydrated cobalt arsenate,  $Co_3(AsO_4)$ ,  $SH_2O$ , crystallising in the monoclinic system and isomorphous with the iron salt vivianite Sometimes, as at Schneeberg in Saxony, it occurs as beautiful radially-arranged groups of blade-shaped crystals with a bright crimson colour and brilliant lustre. More usually, however, it forms an earthy, peach-blossom red encrustation on other cobalt minerals, of which it is an alteration product—It has been found in the Dolcoath mine in Cornwall, Thuringia, Allemont in France, Cobalt in Ontario, Chile, &c. The name was given by F. S. Beuda.t,

in 1832, from  $\epsilon\rho\nu\theta\rho\delta s$ , red — The same name was also used by T Thomson in 1843 for a flesh-red variety of orthoclase-felspar

L. J S.

**ERYTHRITOL.** Erythrol, Erythromannite, Erythrite (v. Carbohydrates).

ERYTHRODEXTRIN v DEXTRIN

ERYTHROL NITRATE v. SYNTHETIC DRUGS. ERYTHROPHLEINE C<sub>28</sub>H<sub>43</sub>O<sub>7</sub>N (?) is an amorphous alkaloid from Sassy bark of Erythrophlœum guineense, G. Don, a large leguminous tree from the West Coast of Africa The bark is used as an arrow poison and in ordeals, and is also known as Casca, Red water, or Mancona bark. The alkaloid yields only amorphous salts, and is the only known one having a digitalis-like action on the heat Heated with concentrated hydrochloric acid it is hydrolysed to methylamine and an amorphous acid Erythrophleine hydrochloride is used to a limited extent in medicine (Gallois and Hardy, Bull Soe chim 1876, 26, 39, Harnack, Chem Zentr 1897, 1 301, Power and Salway, Amer J. Pharm 1912, 84, 337)

# ERYTHROSIN (Tetravodofluorescein)

C20H8O5I4

is prepared by electrolysing a mixture of fluorescein (1 part) and finely powdered iodine (1.5 parts) in soda solution. The vessel is provided with a diaphragm, and the fluorescein mixture occupies the anode space, whilst a dilute solution of soda or sodium hydroxide forms the kathode liquid (D. R. P. 108838, 1899), Chem. Zentr. 1900, i. 1176, cf. also Frdl. 1877–87, 2151.

The commercial product is a red-brick powder dissolving in dilute alkalis without any marked fluorescence It can be purified by dissolving in aqueous ether, filtering and shaking with dilute sodium hydroxide. The sodium salt 14 precipitated by the addition of strong alkali, and, after filtering, washing, and recrystallisation from alcohol, it is dissolved in water, and the free erythiosin is liberated by treatment with hydrochlone acid (Mylius and Foerster, Ber When pure, it is of a lighter 1891, 24, 1482) colour than the impure product, almost insoluble in absolute ether, benzene, or chloroform, but more readily so in acetone, alcohol, and aqueous ether The colour of erythrosin fades under the influence of light and of hydrogen peroxide, particularly in the presence of potassium hydroxide (Zeitsch Farbenindustr 7, 299). Light also causes it to act hamolytically in photodynamic reactions (Sacharow and Sachs, Chem Zentr 1905, 1 1420)

Erythrosin considerably enhances the action of light in its influence on bacteria (Mettler, Chem. Zentr 1905, ii 1037, Huber, ibid 1906, ii 148, Sachaiow and Sachs, lc) Erythrosin is employed in the cotton and paper industries, and Thouert (Compt rend. 1909, 148, 36) has attempted to apply it to colour photography, but with little success so far (cf also Stark, Physikal Zeitsch 1907, 8, 248), it has also been used as a sensitiser in photo-electric experiments (Schlenderberg, J. Phys. Chem. 1908, 12, 574) as an indicator in analysis (Mylius and Foerster, lc.; J. Amer. Chem. Soc. 1899, 21, 359).

Erythrosin is often mixed with rose Bengal (tetralodochlorofluorescein) to suppress its fluorescence, and with common salt to increase its solubility. The former mixture is often used in France to colour foodstuffs, but the presence of rose Bengal in erythrosin is forbidden in America. Jean (J. Soc Chem. Ind. 1908, 70) gives a method for testing and estimating rose Bengal in erythrosin.

Silver erythrosin, which is of an intense blue colour, is prepared by dissolving 1 gram of erythrosin in 400 c c water, adding 7 1 c c. of a 10 p c solution of potassium bromide, and then 12 c c. of a 10 p c silver intrate solution. It forms a colloidal solution which becomes 'ripened' on waiming or standing for a few days, depositing a very fine sediment which has a pure blue colour in transmitted light (Luppo-Cramer, Zeitsch Chem Ind Kolloide, 1907, 1 227; 1908, 2, 325)

The sodium or potassium salt is sometimes known as Erythrosin B, Pyrosin B, Iodeosin

Dianthine B, Eosin J, &c

ERYTHROXYANTHRAQUINONE  $C_{14}H_sO_3$  is produced, along with an isometic hydroxyanthraquinone, by the action of phthalic anhydride on phenol, in presence of sulphuric acid,  $C_8H_4O_3+C_6H_6O=H_2O+C_{14}H_8O_3$  Salicylic acid, anisole, and even anisic acid behave in the same way as phenol. The two hydroxyanthraquinones are separated by boiling with water and barium carbonate, crythroxyanthraquinone is unaltered by this treatment, whereas in the case of the isomeride a soluble barium compound is formed

Érythioxyanthraquinone may also be prepared from aminoanthraquinone, dissolved in acetic acid to which a little sulphuric acid has been added, by adding potassium intrite till the colution becomes yellow. After a short time, on adding water and boiling, yellow flakes of crythroxyanthraquinone separate, and increase as the acetic acid evaporates (D. R. P. 97688, Rower, Ber. 1882, 15, 1793, Perger, J. pr. Chem.

[2] 18, 147).

It is also formed by oxidising the amino-salt with fuming sulphuric acid (D R P 67063, Frdl. in 203), or by heating crythroxyanthraquinone acid (C<sub>15</sub>H<sub>8</sub>O<sub>5</sub> at 270° (Birnkow, Ber 1887, 20, 2438), and by heating anthraquinone-a-monosulphonic acid with a mixture of alkali hydroxide and a salt of an alkaline carth metal under pressure (Fr. Pat 336867, J. Soc. Chem.

Ind 1904, 438, ibid 1905, 841).

Erythroxyanthraquinone crystallises from alcohol in groups of pomegranate-yellow needles, more soluble in hot than in cold alcohol melts at  $173^{\circ}$ – $180^{\circ}$  (Baeyer and Caro), at  $190^{\circ}$ (Liebermann), at 191° (Romer) It begins to sublime at 150°, and condenses to form fine needles, reddish-yellow in colour, like alizarin It is almost insoluble in dilute ammonia, and only slightly soluble in strong ammonia With lime or baryta water it gives a dark-red, almost insoluble lake, decomposable by carbon dioxide The absorption spectrum of a sulphuric acid solution of erythroxyanthraquinone differs from that of its isomeride, hydroxyanthraquinone; the latter body is also more easily converted into alizarin by fusion with caustic otash than the former (Baeyer and Caro, Ber. 7, 986).

Nitric acid oxidises ethythroxyanthraquinone to phthalic acid.

When sodium erythroxyanthraquinone sulphonate is heated with bromine and water at 120°-123°, 2.4-dibromo-1-hydroxyanthraquinone is formed, crystallising in reddish-yellow needles, m.p. 233°. When heated with p-toluidine at 80°-100°, this bromine derivative loses the bromine atom in the -4- position, and is converted into bromoquinizarin blue

which crystallises in blue needles, giving a green colouration with sulphuric acid (D. R. P. 127532, 1902).

Erythroxyanthraquinone forms an acetate  $C_{16}H_{10}O_4$ , yellow needles, m >  $176^\circ$ - $179^\circ$  (Liebermann and Hagen, Ber 1882, 5, 1804)

The constitution of erythroxyanthraquinone has been established by Liebermann (Ber 10, 611, 11, 1611), who synthesised it from a reduction product of quinizarin, hydroxyhydro- $C_6H_4$   $C_{CH(OH)}$   $C_6H_3(OH)$ , anthranol which the hydroxyl-group in the benzene nucleus occupies the ortho-position Frand (Ber 12, 237) was also able to show that in methylhydroxyanthraquinone, a derivative of ordinary hydroxyanthraquinone, the hydroxyl-group was in the para-position, and hence, in the isomeric erythroxyanthraquinone, the hydroxyl- group must be in the ortho-position with regard to the ketone group. Erythroxyanthraquinone is therefore o-hydroxyanthraquinone (v Alizarin AND ALLIED COLOURING MATTERS)

ESCALIN. Trade name for a paste of

aluminium powder and glycerol

ESERAMINE, ESERIDINE, ESEROLINE v ORDEAL BEAN

#### ESERINE or PHYSOSTIGMINE

# NHMe CO·NHC<sub>13</sub>H<sub>15</sub>N OH

(Herzig and Meyer, Monatsh 1897, 18, 389, Heubner, Chem Zentr 1905, n 1111, Straus, Ann 1913, 401, 350; 1914, 406, 332, the most recent work is by Polonovski and callaborators, e q part viii Bull Soc. chim 1918, 23, 356) is the most important of the alkaloids occurring in Calabar bean, in the seed of which it was first found by Jobst and Hesse, in 1864 (Annalen, 1864, 129, 115, Hesse, ibid 1867, 141, 82).

It is present in variable amount (0 04–0 27 pc Carr and Reynolds, Pharm J 1908, 542), and may be extracted by shaking the powdered seed with warm alcohol containing about 1 gram of tartaric acid per litre. The alcohol of the extract is distilled off and the residue is taken up with water, filtered to remove resinous matter and repeatedly shaken with ether until the latter ceases to be coloured. The aqueous extract is then made alkaline with sodium bicarbonate and finally repeatedly shaken out with ether. On evaporation, the alkaloid separates in scaly crystals which may be purified by recrystallisation from ether.

Eserine forms thin rhomboidal, colourless lamellæ, m p 105°-106°, which gradually assume a rose tint and even a yellow colour when exposed to air. It is sparingly soluble in water,

but readily so in alcohol, ether, chloroform, benzene, and carbon disulphide. Eserine is lævo-rotatory, its specific rotation varying between  $-82^{\circ}$  and  $-120^{\circ}$ , depending on the solvent employed (Petit and Polonovski, Bull. Soc. chim. 1893, 9, 1008, cf Orloff, Chem. Zentr. 1897, 1. 1214). Polonovski and Nitzberg, Bull Soc chim 1915, 17, 235 et seq , idem 1916,

When eserine is isolated in the form of its salicylate, the base liberated with sodium carbonate, extracted with ether, and finally allowed to crystallise from a mixture of benzene and petroleum, it separates in stout prisms, mp.  $86^{\circ}-87^{\circ}$ ,  $[\alpha]_{D}$  -75  $8^{\circ}$  By recrystallising in the presence of a crystal of Petit and Polonovski's modification, mp 105°-106°, it is completely converted into the latter, and has  $[\alpha]_D$ -75 8°; the alkaloid is therefore dimorphous (Salway, Chem. Soc. Trans 1911, 2151)

When eserine or its salts are distilled with caustic potash in a current of hydrogen, methylamine and carbon dioxide are amongst the products of decomposition, whilst the residual liquid contains eseroline, a colourless crystalline body, which turns red by the action of moisture Eseroline is also formed, together with methyl carbamide, when eserine is heated in a scaled tube with alcoholic ammonia at 150°. When heated with aqueous potash in presence of air, yields rubreserine C13H16N2O2, red needles, these, in alkaline solution, are gradually converted into eserine blue, which, according to Ehrenberg (Chem Zentr 1894, n 439), is a true dye, colouring wool and silk without mordants. The constitution of cserine blue is not known, of Salway, Chem Soc Trans 1912, 986 not poisonous, but it has a paralysing effect on the heart and central nervous system of the frog (Heubner,  $l \epsilon$ ).

Salts of eserine. Eserine salicylate

$$C_6H_4(OH)CO_2H,C_{15}H_{21}O_2N_3$$

forms slightly bitter, colourless, stout prisms, m p.  $180^\circ-181^\circ$  (Salway), soluble in 150 parts of cold and 30 parts of boiling water, in which it forms an acid solution, it is more readily soluble in alcohol and in chloroform

Eserine metacresotate (metahydroxytoluate)

$$\mathrm{C_{15}H_{21}O_2N_3~C_6H_3(OH)MeCO_2H}$$

m.p. 156°-167°, is prepared by mixing ethereal solutions of metacresotic acid and of eserine Escrine benzoate, m p. 115°-116°, soluble in four parts of cold water, and eserine hydrogen citrate, are prepared similarly The two former are not deliquescent, and yield neutral aqueous solutions, whereas the latter, and also normal eserune tartrate, prepared by dissolving eserine in tartaric acid and crystallising over sulphuric acid, are very deliquescent (Petit and Polonowsky, ic). The methodide forms deliquescent yellow prisms, m p 100° (decomp), and the sparingly soluble picrate crystallises in feathery yellow needles, mp. 114°. Eserme yields no definite aurichloride or platinichloride

Eserine sulphate  $(C_{15}H_{21}O_2N_1)_2H_2SO_4$  is prepared by adding a 10 p c. sulphuric acid, drop by drop, to an ethereal solution of eserine until the newly formed salt no longer separates. After filtration, it is carefully dried at about 40° It

is a white or yellowish-white micro-crystalline powder, odourless, and having a bitter taste It is readily soluble in water, alcohol, and in chloroform. Gold chloride solution, when added to the aqueous solution of the salt, gives a beautiful purple colour, whilst platinic chloride precipitates a yellowish white salt When evaporated with ammonia, the sulphate, like the salicylate, yields a blush residue. If a little eserine sul-phate is placed on a glass slip and moistened with water, and then a drop of sulphuric acid added, characteristic crystals of a spaningly soluble sulphate separate This characteristic appearance is maintained for some time, but eventually the crystals become only and then resinous (Reichard, Pharm Zeit 1909, 50, 375) Eserine and its salts turn red on treatment with alkali or on standing in solution, but the sulphite C<sub>15</sub>H<sub>21</sub>O<sub>2</sub>N<sub>3</sub>,H<sub>2</sub>SO<sub>3</sub>, the aqueous solution of which does not turn red, but remains colourless, has been prepared by Merck by treating a solution of eserine with sulphurous acid, and evaporating the mixture (D R P 166310; J Soc Chem Ind 1906, 443) It is a white powder, readily soluble in water

A derivative, the intensity of the fluorescence of which exceeds that of all known substances. is obtained by allowing an aqueous solution of physostigmine to stand for a few months until it has become deep blue, then adding phthalic and. The substance forms dark-blue crystals with a blood-red fluorescence, and colours cotton, silk, alcohol, ether, &c , dark blue with no fluorescence The aqueous solution, on the other hand, is colourless by transmitted light, but shows a beautiful red fluorescence by reflected light (Goubert, Compt. rend. 1909, 149, 852).

Escrine and its salts should be kept in amber-

coloured, well-stoppered phials

Test for eserine or its salts -A small fragment (the size of a grain of sand) is placed in a porcelain capsule and dissolved in a drop or two of fuming mitric acid The dark-yellow solution formed on warming is evaporated with stirring, the residue, which is of a pure green colour, dissolves in water, strong alcohol, and in sulphuric acid to a clear green solution. 0 005 gram of the alka-loid can thus be detected. The green colouring matter has been termed chloreserine (Ferreira da Silva, Compt rend 1893, 117, 330, Formanck, Chem Zentr 1895, 1.1148, Reichard, *lc*) When exposed to nitric acid fumes, the blue residue becomes violet-blue, and when dissolved in a drop of nitric acid it forms a beautiful reddishviolet solution, which soon changes to blood-red, and on standing or on dilution becomes greenishyellow Escrine reduces molybdic acid very slowly in the cold, but rapidly gives a blue colouration on warming (Reichard, lc) ammonium selenite in concentrated sulphuric acid, eserine gives a lemon-yellow colouration, which becomes orange, and then paler after standing for 3 hours (da Silva, Compt rend 1891, 112, 1266) Eserine can also be detected by means of the polarisation microscope (Kley, Rec trav chim 1903, 22, 367) (For the estimation of eserine or physostigmine, see Heikel, Chem Zentr 1909, 1 949)

Eserine and its salts are of considerable therapeutic value, being employed chiefly as a myotic to contract the pupil of the eye and to decrease intraocular pressure in glaucoma.

According to Pal (Chem. Zentr 1900, in. 1031), esemine forms an antidote to the poisonous effect of curare.

Bilateral antagonism exists between eserine and curare in their effect on muscle. Those muscles which, like the diaphragm, are last paralysed by curare are first set free again by the injection of eserine. The latter also stimulates the respiratory centre (Rothberger, Pfluger's Archiv. 1901, 87, 117)

(For other investigations on the physiological effect and use of eserine or physostigmine, see Matthews and Brown, Amer. J. Physiol. 1904, 12, 173, Edmund and Roth, ibid. 1908, 23, 28, 46, Joseph, ibid. 1909, 23, 215, Anderson, J. Physiol. 1905, 33, 414; Magnus, Pfluger's Archiv. 1908, 123, 99, Kress, ibid. 1905, 109, 608; Modrakowski, ibid. 1907, 118, 52, Unger, ibid. 1907, 119, 373; Harnack, Chem. Zentr. 1908, ii. 338, Winterberg, ibid. i. 750, Loewi and Mansfeld, ibid. 1910, i. 941; Powsner, Biochem. Zeitsch. 1907, 2, 339, Fuhner, Arch. expt. Path. Pharm., 1917, 82, 205.) (V. Ordeal Bean.)

ESPARTO, Stipa maicrochloa (tenacissima) (Linn.) This grass is a native of Spain, Portugal, Greece, and North Africa It is largely used as a raw material for paper making, having been introduced into this country by Routledge. For this purpose it possesses many advantages, as it grows abundantly without cultivation, is easily harvested, yields a large percentage of cellulose of high quality with comparatively simple treatment.

Of the various kinds known to commerce, such as Spanish, Tripoli, Arzen, Oran, Sfaz, Gabes, &c, the first-named is preferred by paper makers, as it yields more and better fibre. It

therefore commands a higher price

To give some idea of the importance of esparto as a paper-making material, it may be mentioned that the annual imports into the United Kingdom are between 200,000 and 300,000 tons

It is also used for the purpose of making coarse mats and topes (v Spon's Enc of the

Industrial Arts)

Esparto arrives in this country firmly pressed The first operation which it underinto bales goes in the paper mill is that of 'dry-picking,' in contradistinction to a subsequent process known as 'wet picking'. The grass is spread out on tables covered with coarse wire cloth, and any root ends, weeds, &c, which may have found their way into the bales through im-Any sand or perfect gathering are picked out The redirt escapes through the wire cloth moval of dut can be more completely effected by machinery. For this purpose the grass is fed into a mechanical duster or willow, where it receives a violent shaking, the dirt being loosened and removed The cleaned esparto is then fed into 'vomiting' boilers, where it is treated with caustic soda solution at pressures varying from 10 to 40 lbs per sq inch for from 3 to 5 hours The amount of soda required is about 17 lbs of 60 pc caustic soda per cwt, but it varies with the quality of the grass, the form of boiler, the pressure at which it is boiled, and the time allowed. When the boiling is completed, the liquor is run away, evaporated to dryness, and ignited, and the alkali recovered in the form of

carbonate This is dissolved in water, converted into caustic by means of lime, and is again available for use.

The boiled grass is washed once or twice in the boiler, and then subjected to the 'wetpicking' process, whereby any unboiled portions are removed. The process is being gradually abandoned in favour of what is known as the 'presse-pâte' system, which consists in purifying the pulp by passing it when bleached through a series of strainers and knotters. It is then made into a coarse web of pulp on the 'presse-pâte,' which may be described as a paper machine without the drying cylinders.

The subsequent treatment of esparto does not differ essentially from that of other fibres, and

need not be particularly described

The haulms (really leaves) of esparto vary in length from 0.3 to 0.5 in, the mean thickness being about 1.5 mm, the individual fibres are 100-400 mm. long, and 0.009-0.5 mm thick

The fibres possess, owing to their fineness and their tendency to curl, considerable 'felting' properties. Paper made from esparto is soft to the touch, 'bulks' well, and at the same time is

strong.

Esparto fibre can be recognised in a paper by the fact that when boiled with a solution of aniline sulphate a rose-pink colour is developed. A similar colour is produced with straw cellulose, but of a paler shade. With rodine and sulphuric acid the wall of the actual bast-fibre assumes a rusty colour, showing that it is not composed of cellulose, with aniline sulphate (cold) it is stained yellow, and with phloroglucinol and hydrochloric acid a crimson colour is assumed, thus demonstrating the presence of bodies characteristic of lignified walls. They can be distinguished by the size and shape of the peculiar serrated epidermal cells, which are to be seen when the commercial fibre is examined under the microscope.

Moreover, espaito pulp always has a number of the short conical, often hooked, hairs which line the inner (upper) surface of the 'iolled' leaf, and which serve to distinguish it from straw, the outer (lower) surface of the leaf is

glabrous

The chemical characteristics of espaito are those of the ligno-celluloses and the pecto-celluloses, the latter predominating

Its composition is shown by the following analyses (Hugo Muller).—

				Spanish.	African.
Cellulos	8e			$48\ 25$	45 80
Fat and	d wa	х.		2 07	2 62
Aqueou	ıs ex	tract		10 19	9.81
Lignin	and	pecto	us		
subst				$26\ 39$	29 30
$\mathbf{A}\mathbf{s}\mathbf{h}$				372	3 67
Water		•		$9\ 38$	8 80
					•
				100.00	100.00

The names 'esparto,' 'sparto,' and 'alfa' ('halfa'), are also given to the leaves of another grass, Lygeum spartum (Loefl.), which likewise grows in Spain and North Africa. These leaves are used as an inefficient substitute for those of Stipa. Microscopically they differ, in that the hars have a considerable lumen, and are

usually blunt-ended; the epidermis has peculiar little rounded silica cells; and the ribs of bast fibres are isolated (not linked together to form a continuous band, as in Stipa)

ESTERIFICATION. The reaction between an acid and a monohydric alcohol appears, at first sight, to be exactly analogous to the neutralisation of an acid by a monoacid base: the equations are:

 $CH_3 CO OH + C_2H_5 OH = CH_3 CO OC_2H_5 + H_2O$ 

The product formed in the second reaction is a metallic salt, and that in the first reaction is termed an alkyl salt or more commonly an ester The esters can be regarded as metallic salts in which the metallic radicals have become replaced by alkyl- groups Corresponding with mono-, di-, and trivalent-metallic radicals are the monodi-, and trivalent hydrocarbon radicals, methyl CH<sub>3</sub>—, ethyl C<sub>2</sub>H<sub>5</sub>—, propyl C<sub>3</sub>H<sub>7</sub>—, &c, methylene CH<sub>2</sub>=, ethylidene C<sub>2</sub>H<sub>4</sub>=, &c, glyceryl C<sub>3</sub>H $\equiv_5$ , &c The process of the corversion of an acid into its alkyl-salts is usually termed esterification, and is the common method by means of which esters are prepared Although this reaction appears to be similar to the process of neutralising an acid by means of an alkali, there are two important points of difference

1. The reaction between an acid and an alkalı in aqueous solution is instantaneous, whereas the reaction between an acid and an alcohol is usually slow, and lends itself readily to study as a time reaction

2. In the process of neutralisation between strong acids and strong alkalis, the reaction proceeds to completion when equivalent quantities of acid and alkali are used, but when equivalent amounts of acid and alcohol are brought together, complete conversion into ester and water never takes place The reaction begins slowly and proceeds until a state of equilibrium is established. The rate at which the reaction proceeds, and the final equilibrium depend upon the specific acid and alcohol employed and upon the temperature process of esterification is a typical balanced or reversible reaction ·

$$R CO_2H + R' OH R CO_2R' + H_2O$$

as the water formed in the process of esterification can hydrolyse the ester to a certain extent This reaction is one of the best to study in order to illustrate the effects of mass action

The reaction between acetic acid and ethyl alcohol was studied by Berthelot and Pean de St. Gilles, in 1862, they showed that working with equivalent quantities of alcohol and acid, equilibrium is reached when 66 pc. of the reacting compounds have been transformed into ester and water. They were able to show that, by working with an excess of alcohol, a higher percentage of the acid can be transformed, as illustrated by the following numbers.

Equivalents of alcoholfor lequiva- 0.205 1015204012 50 lent of acid

p.c. of acid trans-formed into ester 19 42 66.5 78 83 88 93 10^

According to Guldberg and Waage's law of mass action formulated in 1867, the rate of chemical reaction is proportional to the concentrations (molecular) of the reacting substances. and to a constant which is characteristic of the reaction and retains the same value provided the temperature is constant. In the balanced reaction

$$CH_3 CO_2H + EtOH \ge CH_3 CO_2Et + H_2O$$

the direct reaction proceeds at a rate which can be denoted by K a b, where K is the specific constant for this reaction, and a and b are the concentrations of the and and alcohol at the time of measuring the rate Similarly, the reverse reaction proceeds at a rate K' c d, where c and d are the concentrations of the ester and water at the particular time. When the state of equilibrium is reached, we have:

$$K a b = K' c d$$

Using one equivalent of acid and one equivalent of alcohol, Berthelot and St Gilles showed that equilibrium is reached when 0.66 mol of acid and of alcohol have been transformed, leaving 0 33 mol in the free state, we thus have

$$K \times 0.33 \times 0.33 = K' \times 0.66 \times 0.66$$
  
or  $K = \begin{bmatrix} 0.66 \\ 0.33 \end{bmatrix}^2 = 4$ 

The ratio K/K' is denoted by K, and is usually termed the equilibrium constant ratio of the constants for the direct and reverse reactions, and is independent of concentration, but may vary with temperature

This equilibrium constant enables us to calculate the amount of acid transformed when any known quantities of the given acid and alcohol are mixed at the given temperature and kept until equilibrium is attained

With a mixture of a, b, c, d equivalents of acid, alcohol, ester, and water at the given temperature, a-x and b-x equivalents of acid and alcohol will be present when equilibrium is attained, and c+x and d+x equivalents of ester and water will also be present, as each equivalent of acid reacts with an equivalent of alcohol and produces one equivalent of both ester and water

$$K_{K'} = \frac{(c+x)(d+x)}{(a-x)(b-x)}, \text{ but } K_{K'} = 4$$

$$\cdot \cdot \cdot \frac{(c+x)(d+x)}{(a-x)(b-x)} = 4$$

and x can be calculated when a, b, c, d are

It is necessary to distinguish between esterification by means of the acid and alcohol alone, and esterification of an acid by an alcohol in the presence of a catalytic agent The former process is generally termed 'direct esterification,' or sometimes 'autocatalytic esterification,' as it has been suggested that the organic acid itself can act as a catalyst to a certain extent second process is termed the catalytic method of est rification, and is the method most commonly used in the laboratory for the prep ration of esters.

Catalytic esterification. The modern laboratory method is due to E Fischer and Speier (Ber. 1895, 28, 3252). It consists in heating the acid with excess of alcohol, usually about five times the theoretical amount, in the presence of a catalyst. The most efficient catalysts are strong acids, and hydrogen chloride (about 3 p c. of the alcohol) or concentrated sulphuric acid (about 10 p.c. of the alcohol) is generally When hydrogen chloride is used it is advisable to prepare a 3 p c solution of this in the alcohol before adding the organic acid The 3 pc solution is best prepared by weighing the alcohol and passing in the dry gas by means of a delivery tube which just touches the surface of the liquid, and weighing from time to time until the requisite increase in weight is attained The organic acid is then added, and the whole boiled in a reflux apparatus for 3 hours. The greater part of the alcohol is distilled off, the residue poured into about 5 times its volume of cold water, and the solution neutralised with sodium carbonate It is then extracted with ether, the ethereal solution dued and distilled, when a 90 p c yield of the ester can usually be obtained Esters which are readily hydrolysed by water, or esters which are extremely soluble in water, can sometimes be isolated by subjecting the reaction mixture to distillation under reduced pressure J. and M Phelps (Amer J. Sci 1907, [iv | 24, 194) advise heating the acid. eg succinic or benzoic, with about its own weight of 1 p c alcoholic hydrogen chloride, and a small amount of solid zinc chloride at 100°-110°, and afterwards passing in a current of alcohol (compare abid 1908, 26, 281) For constants and properties of common esters, see Mathews and Faville (J. Physical Chem. 1918, 22, 1)

The function of the catalyst is to accelerate the rate of esterification; many acids which react very slowly with alcohol alone,  $e\,g.$  acctic acid at the ordinary temperature, can be transformed rapidly into esters in the presence of a small amount of catalyst Hydrogen chloride is a more efficient agent than sulphuric acid, the ratio being three to one (Kailan, Monatsh 1909, 30, 21), but sulphuric acid is frequently used, as it is more convenient to work with Phelps (Amer J. Sci 1908, [iv ] 260, 290, 296) has examined theaction of various acid sulphates, eg NaHSO<sub>4</sub>, KHSO<sub>4</sub>, piperidine and aniline hydrogen sulphates, but finds that they are not so efficient as sulphuric acid, and similarly he has studied the action of metallic chlorides on the catalytic action of hydrogen chloride Charcoal (Freundlich, Zeitsch physikal Chem 1906, 57, 412) and finely divided platinum (Sudborough and Turner) also act as feeble catalysts. It is usually stated that the presence of the catalyst does not affect the equilibrium, and that provided the ratio of alcohol is kept the same, the percentage of ester formed when equilibrium has been obtained, is the same in both catalytic and direct processes; but Phelps does not agree with this statement Ultra violet does not agree with this statement light brings about esterification of solutions of benzoic and other cyclic acids, especially in the presence of a trace of hydrochloric acid (Stoermer and Ladewig, Ber 1914, 47, 1803).

The amount of ester formed by the catalytic process depends upon the specific acid and alcohol used, and tends to become less when the

ester is readily hydrolysable. In the case of esters extremely soluble in water, or very readily hydrolysed, it is not advisable to pour into water, but to distil immediately under reduced pressure. Phelps and Eddy (Am. J. Sci. 1908, 26, 253) recommend adding solid potassium carbonate to the reaction mixture and distilling under reduced pressure. Certain finely divided metallic oxides have a catalytic action on the formation of esters.

A method of catalytic esterification consists in passing a mixture of the vapours of the acid and alcohol through a tube containing finely divided titanium dioxide at 290°-300° (Sabatiei and Maihle, Compt iend 1911, 152, 494; A. Maihle, Chem Zeit 1913, 37, 777 and 806)

According to Hauser and Klotz (Chem Zeit 1913, 37, 146), glueinum oxide is more efficient than titanium oxide, and even tertiary alcohols yield esters when this oxide is used

Senderens and Aboulenc (Compt rend 1911, 152, 1671, 1855, 1912, 155, 168, 1012) state that by the addition of 5 p c of aluminum sulphate or of potassium hydrogen sulphate to a mixture of equivalent amounts of acetic acid and ethyl alcohol an 82 p c yield of ester is obtained, and that 1 p c of sulphuric acid by voluine gives an 86 5 p c yield. The actual catalyst is stated to be ethyl hydrogen sulphate.

Various esters derived from cyclic alcohols, e.g., cyclohexanol, have been prepared by the method. The temperature, however, must not be too high and the method does not work with aromatic acids containing a carboxylic group directly attached to the benzene nucleus.

For use of (a) anhydrous calcium chloride, see Chem Soc Abstr 1911, 1 601, (b) nickel sulphate Kurtenacker and Habermann, J Pr Chem 1911, (u) 83, 541, and (c) dilute sulphuric acid Bodioux, Compt rend 1913, 156, 1079, 157, 938, 1428 A convenient method for certain esters appears to be a saturated aqueous solution of sodium chloride as solvent and a little sulphuric acid as catalyst

Relationship between constitution and esterification using the catalytic method. V Meyer and Sudborough (Ber 1894, 27, 510, 1580, 3146) were the first to point out any clear relationship between the constitution of an acid and its conversion into an ester by the catalytic method. Their original method was to saturate the alcoholic solution of the acid with hydrogen chloride, to leave at the ordinary temperature for several hours, and then to isolate and weigh the ester formed. Subsequently the Fischer-Speier method was used. They were able to show that practically any substituted benzoic acid, in which the two ortho- positions with respect to the carboxylic group are substituted, does not yield appreciable amounts of ester when its alcoholic solution is saturated with hydrogen chloride and kept for several hours at the ordinary temperature, or when the acid is boiled with a 4 p c. solution of hydrogen chloride m alcohol. All substituents react in much the same manner, eg CH<sub>3</sub>, Cl, Br, F, I, CO<sub>2</sub>H, NO<sub>2</sub>, &c, but the substituents F, OH, CH<sub>3</sub> have not so marked an effect as nitro-, carboxylic, and

halide radicals; eg. acids of the type  $C_gH_2Me_3CO_2H(CO_2H . Me_3=1:2:4:6)$  are esterified to an appreciable extent when hydrogen chloride is passed through their

boiling alcoholic solutions for several hours (Meyer and Sudborough, Ber 1894, 27, 1580, Meyer, *ibid.* 1895, 28, 182, 1254; compare also Kahn, *ibid.* 1902, 35, 3857).

The presence of a second benzene nucleus, as in naphthalene- and anthracene-carboxylic acids, has much the same effect as nitro- or halogen substituents; thus 2-chloro-1-naphthoic acid, anthracene-7-carboxylic acid and 1-chloroanthracene-9-carboxylic acid do not yield esters by the catalytic method, whereas the isomeric 3-chloro-2-naphthoic acid, anthracene-1-carboxylic acid, and 3-chloroanthracene-2-carboxylic acid yield 90 p.c. of ester under the same conditions.

It has been shown by V. Meyer (Ber. 1895, 28, 3197, and Rupp, *ibid* 1896, 29, 1625) that the dibasic acids—tetrachloro, tetrabromo, and tetraiodo-terephthalic acids, tetrachloro, tetrabromo, and tetraiodo-isophthalic acids, i.e. acids in which both carboxylic groups are diortho- substituted, do not yield esters when heated with a 3 p c solution of hydrogen chloride in alcohol An acid in which one of An acid in which one of the carboxylic groups is dioitho-substituted and the other is not, yields an acid ester under similar conditions, e g 2 6-dimethylterephthalic

acid, 
$$C_6H_2Me_2(CO_2H)_2$$
 gives  $CO_2Me \underbrace{\hspace{1cm}}_{Me} CO_2H$ 

(Jannash and Weiler, Ber 1895, 28, 531) Similarly, hemipinic acid, 3.4-dimethoxyphthalic

(Wegscheider, Monatsh 1895, 16, 135) further examples, see Wegscheider, ibid. 1900, 21, 621, 638, 1902, 23, 405, McKenzie, Chem Soc Trans 1901, 79, 1135) An appaient exception is met with in the case of 3 6-dichlorophthalic acid, which, according to Graebe (Ber. 1900, 33, 2026), yields the acid ester

$$Cl$$
 $CO_2Et$ 
 $Cl$ 
 $CO_2H$ 

when its alcoholic solution is saturated with hydrogen chloride and kept at the ordinary Tetrachloro-, tetrabromo-, and temperature tetraiodo-phthalic acids also yield acid esters (Rupp, *ibid* 1896, 29, 1625), o-nitrophthalic acid and papaverine acid (Miller, Annalen, 1881, 208, 243, Wegscheider and Lipschitz, Monatsh. 1900, 21, 790, Wegscheider, *ibid* 1002, 22, 260), belanger 1902, 23, 369) behave in a similar manner, yielding neutral esters, and it is probable that the esters are formed by the conversion of the acids into their anhydrides under the influence of the hydrogen chloride or sulphuric acid, and the subsequent addition of alcohol to the anhydrides. 3 6-Dichlorobenzoylbenzoic acid and tetrachlorobenzoylbenzoic acid are also readily esterified, and it is possible that these react with alcohol as the tautomeric hydroxyphthalides; eg.:

$$\begin{array}{l} C_6 \text{Cl}_4 \swarrow \begin{array}{c} \text{CPh(OH)} \\ \text{CO} \\ \text{\longrightarrow} \text{CO}_2 \text{Et} \cdot C_6 \text{Cl}_4 \cdot \text{CPh(OH)}_2 \\ \text{\longrightarrow} \text{CO}_2 \text{Et} \cdot C_6 \text{Cl}_4 \cdot \text{CPh} \\ \end{array}$$

The results obtained with tri- or even (Graebe). hexa-carboxylic acids show the same inhibiting effect of ortho-substituents. Thus trimesic acid benzene 1 3:5-tricarboxylic acid, yields a neutral ester, hemimellitic acid, the 1 2 3-tricarboxylic acid, yields a dimethyl ester , pyromellitic acid, benzene-1 2 4 · 5 · tetracarboxylic acid, yields a normal ester, prehnitic acid, 1 · 2 : 3 · 4 · tetracarboxylic acid, gives a dimethyl ester in the cold, and the neutral ester on heating, the formation of the latter is probably due to the fact that an anhydride is first formed, and this, with methyl alcohol, gives the dimethyl ester

which is further esterified to the normal ester Dinitro-pyromellitic acid, mellitic acid, and hexahydromellitic acid do not yield esters (Meyer, Ber 1894, 27, 1590, 28, 182, 1896, 29, 840) Isohydromellitic acid, on the other hand, yields a monomethyl ester (Van Loon, ibid 1895, 28, 1272)

Substituted benzoic acids in which the substituents are not in ortho-positions, give good yields of esters by the Fischer-Speici method, and this difference in behaviour has been used for the following purposes 1 Separation of an ortho-disubstituted benzoic acid from isomerides (compare Martz, Ber. 27, 3147, for separation of dinitro-benzoic acids; Jannasch and Weiler, ibid. 3445, for separation of isomeric trimethyl-benzoic acids) Rosanoff and Prager (J Amer Chem Soc 1908, 30, 1912) show that a mixture of benzoic acid with an ortho-substituted benzoic acid can be separated by partial esterification in the cold with a 3 p c solution of hydrogen chloride in ethyl alcohol 2 Determination of the constitution of particular substituted benzoic acids. (Compare V. Meyer, Ber 1895, 28, 187, for 2-methyl - 5 - bromobenzoic acid, Zincke and Francke, Annalen, 1896, 293, 123, for 4-acetyl-5-bromo-iso-phthalic acid, and Matthews, Chem. Soc Pioc 1900, 16, 187, for 2 3 5-trichlorobenzoic acid. See also Blaise, Compt rend. 1898, 126, 733, for substituted succinic acids)

When the carboxylic group is not directly attached to the benzene nucleus, but is united to a side chain, the presence of ortho-substituents does not prevent the formation of esters, eg. 2 4 6-tribromo-3-aminophenylpropionic acid and 2 4 6-tribromophenylpropionic acid, mesitylacetic and mesitylglyoxylic acid all gave good yields of esters when the alcoholic solutions are saturated with hydrogen chloride. Kellas (Zeitsch-physikal. Chem 1897, 24, 221) and Goldschmidt (Zeitsch. lect ('hem 1909, 15, 4) have shown that the introduction of one ortho-substituent into the benzoic acid molecule retards esterification to an appreciable extent, the nitro- group having a greater inhibiting effect than Me, Cl. Br. or I Kailan also shows that o-aminobenzoic acid hydrochloride is esterified more slowly than its isomerides (Monatsh 1906, 27, 1038), but he finds that o-nitrocinnamic acid is esterified more readily than the p- and m-nitro acids (ibid 1907, 28, 1163). Goldschmidt was able to show that using a dilute alcoholic solution of the acid with hydrogen chloride as catalyst, the velocity of the reaction can be determined by means of the equation for a unimolecular reaction, K=

 $1/t(\log a/a - x)$ ; since under these conditions the mass of the alcohol can be regarded as constant, and the reaction is practically non-reversible in the presence of the large excess of alcohol. The concentration of the acid at any given moment can be determined by direct titration with standard barium hydroxide solution, using phenolphthalein as indicator. When, however, the ester formed is readily hydrolysed, as in the case of ethyl formate, ethyl trichloracetate, and ethyl pyruvate, it is necessary to titrate with ammonium hydroxide solution, using litmus as indicator, as the end point is not definite when baryta is used, owing to the instantaneous hydrolysis of the ester by the baryta. The constant K is directly proportional to the concentration of the hydrogen chloride in the absence of water, and falls somewhat as t (time) increases, owing to the fact that the water produced during the esterification has a retard-The influence of small amounts of ing effect. water has been studied by Kallan (Monatsh 1906, 27, 543, 997, 1907, 28, 115, 571, 673, 705, 965, 1069, 1137, 1187, 1908, 29, 799; 1909, 30, 1), who has shown that the relationship between the rate of esterification and the amount of water can be represented by means of an equation of the type, eg in the case of cinnamic acid

$${\begin{aligned} &\overset{1}{K} &= 0.665 + \frac{4876}{c} - \frac{1.678}{c_2} + \left(2.911 - \frac{3.212}{c} + \frac{4.358}{c_2}\right) & \\ & + \left(-27.54 + \frac{31.78}{c} - \frac{2.284}{c_2}\right) & \\ & \end{aligned}} w^2$$

for concentrations of water w=0 3 to 1 3, and for concentrations of hydrogen chloude c=0 15 to 0 66 (Compare also Fitzgerald and Lapworth, Chem. Soc Trans 1908, 93, 2168.)

The inhibiting action of water on a mixture of acid and alcohol is readily shown by comparing the rates of esterification of the same mixture of acid and alcohol. (a) in presence of water; (b) in presence of a corresponding quantity of benzene (Kailan, Zeitsch physikal Chem 1915, 89, 676) When 25 or 50 volume p.c. of alcohol is used the relative effects of benzene and water are as 1.100 or 150 The effect of water on the esterification of dibasic acids has also been examined (Kailan, ibid. 1913, 85, 706; 1914, 87, 619)

Sirks (Rec. trav. chim. 1908, 27, 237) has

Sirks (Rec. trav. chim. 1908, 27, 237) has obtained the following values for the esterification constants of the isomeric dintrobenzoic acids at 25°, using ethyl alcohol containing a little water with hydrogen chloride as catalyst . 3:4-dintro-acid, 0 0086, 3.5-acid, 0 0050, 2:3-acid, 0 0005, 2.5-acid, 0 0003, 2.4-acid, 0 0002, and 2.6-acid, nil. The values for  $\mathbf{E}_{\mathrm{ktoh}}^{25\circ}$  obtained for benzoic, m-, and o-nitrobenzoic acids were 0 0132, 0 0071, 0 001.

These numbers indicate the retarding effect of one or two nitro-groups in the ortho-position; even one m-group retaids, but the introduction of a second intro-group into the meta-position of the m-nitro acid molecule, increases the rate.

From a comparison of the rates of esterification of these acids with their dissociation constants, Sirks comes to the conclusion that acids with small dissociation constants are esterified more readily by the catalytic method than acids with high dissociation constants; but an examination of certain alkyl-derivatives of aliphatic acids shows that the statement does not hold good generally.

Sudborough and Lloyd, by using Goldschmidt's method of determining esterification constants, have shown that the introduction of one substituent into the acetic acid molecule has a retarding effect, whatever the nature of the substituent The introduction of two substituents usually has a more marked effect than one, and three a greater influence than two, when the two or three substituents are alike. A comparison of the esterification constants for the normal fatty acids gave the following values for  $E_{MeOH}^{15^{\circ}}$ . formic 1124, acetic 104, propionic 92, and from butyric to stearic values between 50 and 54 The introduction of a carbonyl-group in place of the α-methylene group of a normal fatty acid produces a retarda on (Kailan, Monatsh 1907, 28, 1187, Sudborough, Chem Soc Trans 1912, 101, 1227) The introduction of methyl- groups into the molecule of methyl hydrogen succinate also produces a lowering of the esterification constant (Bone, Sudborough, and Sprankling, Chem Soc Trans. 1904, 85, 534)

In the case of unsaturated acids, the following generalisations have been drawn (compare Sudborough and Lloyd, Chem. Soc. Trans 1898, 73, 81, Sudborough and Roberts, ibid 1905, 87,1840, Sudborough and Thomas, ibid. 1907, 91, 1033, Sudborough and Gittins, ibid. 1909, 95, 315, Sudborough and Davies, ibid. 975). 1 An abunsaturated acid is esterified much less readily than its saturated analogue, in many cases the ratio is 1 40. 2 An ab-unsaturated acid is esterified far less readily than the isomeric acids, in which the double linking is further removed from the carboxyl-group As a rule, βγ- unsaturated acids are esterified more readily than their saturated analogues,  $e g \beta \gamma$ -phenylcrotonic acid has  $E_{MeOH}^{15^{\circ}}$  86 6, and the saturated  $\gamma$ -phenyl-n-butyric acid  $E_{NeOH}^{15^{\circ}}$  56 25 γδ-Acids and other unsaturated acids, in which the double linking is further removed from the carboxyl- group, are esterified at much the same rate as the corresponding saturated acids, e gE for oleic acid is 543, and for palmitic acid 497 3 In most cases the introduction of a substituent into the a- position in the acrylic acid molecule has a greater inhibiting effect than when introduced into the \$\beta\$- position 4 The effect of a substituent in the cis- position is more marked than in the trans-, thus an acid of the

The difference in the rates of esterification of an  $\alpha\beta$ - unsaturated acid and the saturated analogue, a difference which has been confirmed by Kailan (Monatsh. 1907, 28, 1137) in the case

<sup>&#</sup>x27;The letter E is used to denote the ester'fication constant of an acid, using normal hydrogen chloride as catalyst, and the temperature and specific alcohol are also given, as the constant varies with these.

of cinnamic and hydrocinnamic acids, affords a very convenient method for separating mixtures of the two. This is well shown in the preparation of hydrocinnamic acid. The acid is obtained by the reduction of cinnamic acid, and is liable to contain small amounts of the latter, which are difficult to remove by crystallisation, but a complete separation can be effected by the action of a dilute solution of hydrogen chloride in ethyl alcohol. The saturated acid is completely transformed into ester, and, under suitable conditions, the unsaturated acid iemains A similar method can also be unesterified. used with advantage for the separation of a mixture of an  $\alpha\beta$ - unsaturated acid and the isomeric  $\beta\gamma$ -acid, this method is preferable to the older method used by Fittig, in which the  $\theta\gamma$ -acid was transformed into the  $\gamma$ -lactone (Sudborough and Thomas, Chem Soc Trans. 1911, 99, 2307)

Gyr (Ber 1908, 41, 4308) has determined the esterification constants of a number of mono-, di-, and tri- substituted acetic acids, and confirms the results obtained by Sudborough and Lloyd The values, for example, for acetic, phenyl acetic, dipnenyl acetic, and triphenyl acetic, at 25° are respectively 7.76, 3.57, 0.20, and 0.001 worked with absolute methyl alcohol, and recommends boiling some three or four times with small amounts of metallic calcium in order

to remove the last traces of water

Kistiakowsky (Zeitsch physikal Chem 1898, 27, 250) has determined the velocity of reaction of an acid and alcohol in mixtures of the alcohol and water containing hydrogen chloride this case the esterification and hydrolysis proceed side by side, and the velocity constants for the two are denoted by K and K'. Then:

$$K = \frac{1}{t} \frac{\xi}{a} \log_e \frac{\xi}{\xi - x} \text{ and } K' = \frac{1}{t} \frac{a - \xi}{a} \log_e \frac{\xi}{\xi - x}$$
$$K + K' = \frac{1}{t} \log_e \frac{\xi}{\xi - x}$$

when the amounts of water and alcohol are large compared with the amount of acid present Edenotes the gram-molecules of acid transformed into ester at equilibrium, and x the number of gram-molecules transformed at end of time t K can also be calculated from K+K' by

means of equation ·

$$K = (K + K')^{\xi}_{\alpha}$$

The values of K and K' and also the value  $\frac{\xi}{z}$ are independent of the original concentration of

the organic acid

The same values for K and K' are obtained when a mixture of acid, alcohol, water, and catalyst, or ester, alcohol, water, and catalyst are used, provided the proportion of water and alcohol and the concentration of the catalyst are kept the same

The temperature coefficient between  $t = 25^{\circ}$ and  $t = 30^{\circ}$  is about 10 pc per degree, using mixtures of alcohol and water containing

13-70 p.c. alcohol

The values of both K and K' fall on passing from formic to acetic, and again from acetic to monochloracetic

The effect of the proportion of alcohol and water present is marked The value of K in the

case of formic acid rises from 0 to 300 as the percentage of alcohol increases from 0 to 914. whereas the value for K' falls from 198 to 60. as the percentage of alcohol is increased from 0 to 914 In the absence of a catalyst, the velocity of esterification falls as the organic acid is used up, whereas the velocity of hydrolysis is first small, and gradually increases as fresh acid is formed, and both K and K' diminish with an increase in the amount of alcohol present

Direct method of esterislication. The reaction between alcohol and a mineral acid is frequently used as a method for the preparation of ethyl esters of inorganic acids, e y ethyl chloride bromide, nitrite, sulphate (for ethyl bromide, cf Weston, Chem Soc Trans 1915, 107, 1489,

Holt, ibid. 1916, 109, 1) Norris, Watt, and Thomas (J. Amer Chem Soc 1916, 38, 1071) have studied the reaction between different alcohols and dilute solutions of hydrochloric and hydrobromic acids Thev find that the rate of formation of esters in the case of primary alcohols tends to decrease as the molecular weight of the alcohol in-Of the three groups of alcohols, the tertiary are the most, and the primary least, reactive

Menschutkin, working with the fatty acids (J pr Chem 1881, [n] 24, 49; *ibid* 1882, [n] 25, 193, Ber 1897, 30, 2783), was one of the first to draw attention to the relationship between the rate of esterification and the constitution of the acid and of the alcohol, and found that the amount of acid converted into ester in a given time diminished with the complexity of the acid molecule, and also with the complexity of the alcohol molecule. By heating a given amount of acetic acid with equivalent amounts of different alcohols for one hour at 135°, Menschutkin was able to show that, on the whole, a primary alcohol is esterified more rapidly than a secondary, and a secondary than a tertiary. Some of the results obtained by Menschutkin for primary alcohols are not true measures of velocity, as equilibrium had already been obtained within the hour It is also pointed out that the introduction of side chains or of negative substituents into a primary alcohol also tends to lower the rate of esterification, and the greater the number of such groups introduced the more pronounced is the effect. The introduction of an olefine linking into the molecule of the alcohol in the  $\alpha\beta$ - position also diminishes the rate of esterification. Subsequent experiments made by heating the alcohol with given weights of acetic anhydride gave similar results Still later work with hydroxy-polymethylenes (Chem. Soc. Trans. 1906, 89, 1533), using the acetic anhydride method, indicates that these compounds are esterified more readily than analogous open-chain compounds, eg the value for methyl-propyl-carbinol and cyclopentanol are respectively 98 and 189, and that cyclic tertiary alcohols are esterified least readily and primary most readily. The results also indicate that methyl- groups in the o- position with respect to the hydroxyl- group also produce retardation, but not to the same extent as whon attached to the carbon atom to which the OH is attached Michael and Wohlgast (Ber. 1909, 42, 3157) have determined the rates

of esterification of trichloracetic acid with following numbers, using the equation for a various alcohols at 25° and 50°, and give the bimolecular reaction:—

Primary alcohols	$K \times 10^s$	Secondary	K × 10° 25°	Tertiary	K×10' 25°
Methyl Ethyl	3690 24,000 650 4,210 725 1200 740	Dimethyl-carbinol Methyl-ethyl ,, Methyl-propyl ,, Methyl-hexyl ,,	. 98 . 90 . 76 . 109	Ter-butyl Ter-amyl	. 118
Heptyl Octyl Cetyl	1160 1860 2500 16,300				

According to Michael, factors which are of importance are. 1 Degree of association of the alcohol; as this is greater in the case of the simpler alcohols, the rate of exterification would be reduced to a greater extent 2 Readiness with which the alcohol decomposes into H and RO in order to react with the carbonyl- group 3 The influence of alkylgroups on the affinity of the RO and H for the unsaturated atoms of the carbonyl- group Michael claims that his results do not afford the slightest evidence for the view that atoms or groups can produce effects by mere filling of Space

As regards the influence of the constitution of the acid on the velocity of esterification, Menschutkin drew the following conclusions as the result of experiments made by heating the respective acids for one hour with isobutyl The introduction of an alkyl- group alcohol into the formic acid molecule invariably produces a retardation in the esterification. The introduction of alkyl- groups into the acetic acid molecule also produces a retardation, and this is greatest when three alkyl- groups replace the three hydrogen, eg trimethylacetic acid, and is least when only one alkyl- group is introduced Unsaturated acids of the type of crotonic and cinnamic are esterified less readily than their saturated analogues

Lichty (Amer Chem J 1895, 17, 27; 1896, 18, 590) found, however, that in the case of the chlorinated acetic acids the introduction of a chlorine atom into the acetic acid molecule tends to facilitate the formation of ester when the direct method is used. The presence of an  $\alpha\beta$ - ethylene linking has not the same marked effect in direct esterification as in catalytic esterification (Thomas and Sudborough, Chem Soc Trans 1912, 101, 317) It is obvious that the results obtained by the direct and catalytic methods of esterification are not comparable, and Sudborough and Feilmann (Chem. Soc Proc. 1897, 13, 241) have suggested that at least two factors are concerned in the process of esterification when using the same alcohol, viz (1) strength of the acid esterified, (2) the presence of substituting groups in close proximity to the carboxyl- group When the direct method of esterification is used, the former appears to be the determining factor, whereas with the catalytic method the latter is of primary importance; but in either case both factors are involved Compare also Flurscheim, Chem Soc. Trans. 1909, 95, 718

Rosanoff and Prager (J. Amer. Chem. Soc. 1908, 30, 1895) have also shown that durthosubstituted benzoic acids can be esterified by the direct method, provided they are heated with

excess of the alcohol at a comparatively high temperature (183°) for about 100 hours. The reaction proceeds even at the boiling-point of alcohol, but requires prolonged heating. Rosanoff and Prager determined both the velocity constant K and the reciprocal of the equilibrium constant, viz  $\mathbf{K}'=\mathbf{K}'/K$ . They used different weights of alcohol and acid in each experiment, and by taking the volume of the solution into account, they calculated K by means of the equation

$$K = \frac{2\ 3026v}{t\sqrt{(a-b)^2 + 4ab}\mathbf{K'}} \times \frac{2ab - x(a+b-\sqrt{(a-b)^2 + 4ab}\mathbf{K'})}{\log_{10}\ \frac{2ab - x(a+b+\sqrt{(a-b)^2 + 4ab}\mathbf{K'})}{2ab - x(a+b+\sqrt{(a-b)^2 + 4ab}\mathbf{K'})}}$$

where a and b are the molar concentrations of the acid and alcohol, v is the volume of the mixture in litres, x the number of gram-molecules of the ester formed after time t-(in days). The values of K obtained for a few acids are.

2 3 5-Tribromobenzoic . 0 370 3 4 5 0 900 2 4 6 0 088 2 . 4 . 6-Trichlorobenzoic . 0 031 Acetic . 1 450 Propionic . 0 950 Isobutyric . 1 070 Trimethylacetic . 0 180 Chloracetic 2 460 Dichloracetic 6 380 at 80° Trichloracetic 12 000	Benzoic .	0 630
2 . 4 6-       "0 088         2 . 4 . 6-Trichlorobenzoic       0 031         Acetie       1 450         Propionie       0 950         Isobutyrie       1 070         Trimethylacetic       0 180         Chloracetic       2 460         Dichloracetic       6 380 at 80°	2 3 5-Tribromobenzoic	0 370
2 . 4 . 6-Trichlorobenzoic       . 0 031         Acetic       . 1 450         Propionic       . 0 950         Isobutyric       . 1 070         Trimethylacetic       . 0 180         Chloracetic       . 2 460         Dichloracetic       . 6 380 at 80°	3 4 5- ,,	0 900
Acetie       .1 450         Propionie       0 950         Isobutyrie       .1 070         Trimethylacetie       0 180         Chloracetie       .2 460         Dichloracetie       .6 380 at 80°		0 088
Propionic	2.4.6-Trichlorobenzoic	0 031
Isobutyric       .       1 070         Trimethylacetic       .       0 180         Chloracetic       .       2 460         Dichloracetic       .       6 380         at 80°	Acetic	1 450
Trimethylacetic         0 180           Chloracetic         2 460           Dichloracetic         6 380		0 950
Chloracetic 2 460 at 80°		1 070
Dichloracetic 6 380 at 80°		0 180
Trichloracetic 12 0001		
· · · · · · · · · · · · · · · · · · ·	Trichloracetic	12 000)

and it is pointed out that in many of these cases an increase in velocity of esterification runs parallel with an increase in the electrolytic dissociation constants of the acids, although the two sets of constants are not proportional

the two sets of constants are not proportional Michael and Oechslin (Ber 1909, 42, 317) have determined the amounts of ester formed when various mono- substituted benzoic acids are heated with methyl alcohol in sealed tubes at 135° without a catalyst, but they did not determine the actual constants. They found that many m- and p- substituted benzoic acids containing negative substituents are esterified somewhat more readily than benzoic acid, but that an ortho- substituent, with the exception of Cl, tends to retard esterification, and that the effect is the greatest in the case of the methylgroup.

The following are some of the values obtained after 7 hours. benzoic 11·1, o-chlorobenzoic 11·9, o-bromobenzoic 10·4, o-iodobenzoic 9·7, o-nitrobenzoic 10·0, salicylic 7·7, and o-toluc 6·2

The results obtained by Rosanoff and Prager and by Michael and Oesc'.elin are compatible

with the view that at least two factors are concerned in esterification either by the catalytic or the direct method; that in the former case the chief factor is the presence of substituents in close proximity to the carboxyl- group (steric hindrance) whereas the strength of the acid, as measured by its dissociation constant, is of but secondary importance; in the direct method the relative values of the two factors are frequently (Note The dissociation the exact opposite constant for s-trichlorobenzoic acid is not known) The results do not justify the conclusion drawn by both Rosanoff and Michael, that substituents do not retard esterification by mere space filling

For direct esterification of dibasic acids, see Kailan (Zeitsch physikal Chem 1915, 89, 676)

The reaction between a mercaptan and an acid is analogous to that between an alcohol and an acid, and the product is the ester of a thio-acid. The reaction has been studied in detail by L. S. Pratt and E. E. Reed (J. Amer. Chem. Soc. 1915, 37, 1934) at a temperature of 243°, and using different mercaptans. The limiting values are much less than when the corresponding alcohols are used, and tend to dimmish as the molecular weight of the alcohols increases.

Other methods for preparing esters. which cannot be obtained by direct esterification, eg. the esters of diortho-substituted benzoic acids, can be prepared by the following methods 1 Action of the alcohol on the acid chloride or acid anhydride. The latter action is accelerated by the presence of small amounts of a strong acid (Reychler, Bull Soc chim Belg. 1907, 21, 428) 2 Action of the alkyl iodide on the dry silver salt, or, in the case of aromatic acids, by the action of alkyl chlorides on dry potassium salts in the presence of a trace of a base such as triethylamine Benzyl benzoate is readily prepared by heating together at 95°-100° potassium benzoate (100), benzyl chloride (120), and triethylamine (1) 3 Action chloride (120), and triethylamine (1) of methyl sulphate on an aqueous solution of the sodium salt of the acid (Werner and Seybold, Ber 1904, 37, 3658) 4 Action of an ethereal solution of diazomethane on the acid

Sulphonic acids. Sulphonic acids do not give good yields of esters when warmed with alcohol alone or with alcohol and a mineral acid. This is probably due to the fact that the ester formed immediately reacts with the alcohol, giving the free acid and an ether:

$$C_6H_5SO_2\cdot OMe + MeOH = C_6H_5SO_2OH + Me_2O$$

(comp Krafft and Roos, Ber 1893, 26, 2823, Wegscheider and Furcht, Monatsh 1902, 23, 1094) The esters are not formed when the sodium salt of the acid is warmed with alkyl iodide and alcohol (Wegscheider and Furcht); but can be obtained by the action of methyl sulphate on the acid or by the action of the alcohol on the acid chloride at low temperatures (Krafft, Ber. 1892, 25, 2256)

Esters of dibasic acids. A dibasic acid can give rise to both an acid and a normal ester, e.g. CO<sub>2</sub>H CH<sub>2</sub> CO<sub>2</sub>H—>CO<sub>2</sub>H CH<sub>2</sub> CO<sub>2</sub>Me and CO<sub>2</sub>Me CH<sub>2</sub> CO<sub>2</sub>Me. With the exception of diortho-substituted aromatic dibasic acids, the normal esters can be obtained by the Fischer-Speier method of esterification. For the preparation of the acid es'ers, the following methods

are resorted to: 1. Addition of alcohol or of sodium alkyl oxide to the acid anhydride. 2. Partial hydrolysis of the normal ester 3. Partial esterification of the acid. 4 The action of alkyl iodides on the acid silver salts, if these can be prepared, or on the acid potassium salts. In the case of a symmetrical dibasic acid, the same acid ester is formed by all these methods, although, in most cases, except those coming under the first heading, the yields are not very good, as free acid and normal esters are also formed, and the separation is not always easy

With an unsymmetrical dibasic acid, two isomeric acid esters (the a and the b)  $^1$  can exist. and which of these is formed depends largely on which method of preparation is used Wegscheider has carried out numerous experiments (comp. Monatch. 1895, 16, 75, 1897, 18, 418, 629, 1899, 20, 685, 1900, 21, 621, 638, 787, 1902, 23, 317, 357, 369, 393, 405, 1903, 24, 413, 915, 1905, 26, 1039, 1906, 27, 487, 777; 1907, 28, 819, 1908, 29, 525, 531, 535, 541, 557, sulphonated benzoic acids, *ibid* 1902, 23, 1093, tribasic acids, *ibid* 1910, 31, 1253) on the following acids hemipinic (3 4dimethoxyphthalic), nitrohemipinic, camphoric, mitro-, bromo-, amino-, and hydroxy-terephthalic, 3- and 4-nitrophthalic, 4-hydroxyphthalic, phthalonic, homophthalic, aspartic, phenylsuccinic, and papaverinic, and has drawn the following conclusions under the assumption that the two carboxylic groups differ appreciably as regards their relative strength 1 By the addition of alcohol to the anhydride and by the action of alkyl iodides on the acid potassium or acid silver salts, the stronger carboxyl- group is esterified, so that under these conditions a-acid esters are formed. If there is but little difference in the relative strengths of the two carboxylgroups, then a mixture of the two acid esters is obtained, but the a- compound preponderates;  $e q = C_6 H_2(CO_2 H)_2(OMe)_2(1 \cdot 2 \cdot 3 \cdot 4)$  gives, by these methods, the acid ester

Exception phenyl succinic anhydride Diazomethane and the acid give mainly neutral ester, but also small amounts of acid ester derived from the stronger carboxyl- group 2. By the process of catalytic esterification the carboxylic acid group, which is freer from adjacent substituents, is esterified first, thus opianic acid gives the b- ester C<sub>8</sub>H<sub>2</sub>(CO<sub>2</sub>Et)(CO<sub>2</sub>H)(OMe)<sub>2</sub>,

1 2 3 4

but neutral ester and small amounts of the aisomeride may also be formed, especially if the
conditions are such that anhydride formation
can occur. The same rule holds good for
direct esterification if the possibility of anhydride
formation is excluded. 3 The addition of
sodium alkylate in the absence of alcohol and
water to the anhydride yields the b- acid ester,
with appreciable amounts of the isometide
4. The partial hydrolysis of the neutral ester
gives the acid ester, which is difficult to prepare

<sup>&</sup>lt;sup>1</sup> The term b-ester is given to the acid ester with the higher dissociation constant, and the isomeride is termed  $t_{-6}$  a- (Ber 1902, 35, 4329, 1903, 36, 304).

by the method of catalytic esterification; eg dimethylhemipinate gives the a-acid ester. Exceptions the hydrolysis of methyl hydroxyterephthalate, 4-hydroxyphthalate, 4-nitroph-

thalate, aminoterephthalate, phthalonate.
For formation of esters of phthalic acid in presence of hydrochloric acid and hydrolysis of the esters, see Wegscheider and Amann (Monatsh 1915, 36, 549 and 633)

In the case of the majority of symmetrical dibasic acids the two carboxylic groups are esterified at the same rate; in other words, the replacement of OH by OEt does not affect the rate of esterification In the case of both maleic and oxalic acids, however, there is a retardation produced by the conversion of one of the CO<sub>2</sub>H groups into a CO<sub>2</sub>Et This is attributed by Kailan (Zeitsch physikal Chem 1914, 87, 619) to the proximity of the two carboxylic groups to one another, and as the retardation is greater in the case of maleic acid, it is argued that in this acid the two carboxylic groups are closer than in oxalic acid

Theories of esterification.—Direct method The reaction between the acid and alcohol is not a simple ionic reaction, since the acid with the highest dissociation constant is not necessarily esterified most readily, and in the case of a dibasic acid it is not always the stronger CO2H group which is esterified first. The view usually accepted at the present time is that an additive compound of alcohol and acid of the type ∕OEt

R C OEt, ie a dialkyl-derivative of an ortho-OH

carboxylic acid, is first formed by the addition of a molecule of alcohol to the carbonyl- group of the acid (Wegscheider, Monatsh 1895, 16, 137, Henry, Ber 1877, 10, 2041) Michael suggests that before this additive compound is formed an unstable 'Kekulé double molecule' is formed, perhaps by the coming into play of the two latent valencies of the oxygen atoms

$$R < \stackrel{C=O}{<} \stackrel{OH}{<} Et$$

it is well known that many esters can form additive compounds with sodium ethoxide, and, therefore, presumably with ethyl alcohol, although compounds of the latter type have not been isolated Pechmann (Ber 1898, 31, 503) has been able to show that the esters of diorthosubstituted benzoic acids cannot yield additive compounds with sodium ethoxide, probably owing to the space filling (steric hindrance) of the group in the ortho-position; by analogy, the diortho-substituted acids would not yield the orthocarboxylic acid derivative at all readily, and this would account for the fact that diorthosubstituted benzoic acids are not readily esterified

Catalytic esterification According to Michael, the processes of direct and catalytic esterification are essentially different, the former is a reaction between the organic acid and alcohol, and the latter a reaction between the organic acid and an additive compound of the alcohol and catalyst (Ber. 1909, 42, 312). This reaction does not consist of addition to the carbonylgroup of the acid, but in union of the acid with the alcohol-catalyst complex to form a large 'double molecule,' which decomposes immediately into ester, water, and catalyst. The fact

that esterification is facilitated by the presence of the catalyst is accounted for by the assumption that the alcohol-catalyst complex contains a greater store of energy, is therefore more reactive than the alcohol itself, and can more quickly develop the maximum entropy of the alcohol-acid system It is questionable whether these views are in harmony with the facts established by Wegscheider, viz that the same acid ester is obtained from an unsymmetrical dibasic acid, whether the direct or catalytic method of esterification is adopted, provided that anhydride formation is excluded. Goldschmidt also comes to the conclusion that the two processes are essentially different (Zeitsch Elektrochem 1909, 15, 4) He claims to have shown that the presence of sodium trichloracetate does not depress the velocity of esterification of trichloracetic acid by the direct method, and, therefore, the direct esterification cannot be a simple case of hydrion catalysis Later experiments of Sudborough and Turner show that the presence of salts of trichloracetic acid does diminish the rate of esterification of the acid, and hence Goldschmidt's conclusion is not justified. The fact that in direct esterification the strength of the acid is an important factor is compatible with the view that the organic acid itself plays the same part in direct esterification that the catalyst does in catalytic esterification agrees again with the fact that the reaction in certain cases of direct esterification is bimolecular (Goldschmidt, Ber 1896, 29, 2208), provided the reaction is not between the ions of acid and alcohol.

Kaılan (Monatsh 1908, 29, 799) is also of opinion that the processes of direct and catalytic esterification are not analogous, since the addition of a small amount of water increases the rate of esterification of trichloracetic acid by the direct method, but diminishes the activity of trichloracetic acid when it is used as a catalyst.

Various views are held with regard to the function of the catalyst Goldschmidt and Uddy (Zeitsch physikal Chem. 1907, 60, 728; comp also Ber 1906, 39, 711, and Zeitsch. physikal Chem. 1910, 70, 627) conclude that the hydrions of the catalyst are the active agents, since the activity of different acids is in the order of their relative strengths, eg. the relative values of the acids hydrochloric, pieric, trichloracetic, and dichloracetic as catalysts are indicated in the order given, and the addition of salts of the acid used as catalyst diminishes the activity of the catalyst They further conclude that the hydrions unite with the alcohol, and

that the complex ion (C2H6O,H) is the active agent. They suggest the following modified formula for calculating the velocity constant:-

$$K.c = \frac{1}{t}(n+r+a)\log_{\theta}\frac{a}{a-x}-x$$

where c =concentration of catalyst, n =concentration of water, and r =hydrolytic constant of the complex alcohol ion for

$$(C_2H_6^+O,H)$$
  $r=1.5.$ 

The constants obtained by means of this equation agree better, and do not show the tendency to decrease as t increases, a tendency which is marked when the constants calculated

by means of the ordinary equation for a unimolecular reaction are used.

A marked exception to the depressing effects of the addition of salts is met with in the case of hydrogen chloride, where the addition of chlorides such as calcium chloride, tetramethyl-ammonium chloride, and aniline hydrochloride, have no appreciable effect on the catalytic activity of hydrogen chloride (Goldschmidt, Sudborough)

Many chemists are of opinion that in catalytic esterification, as in catalytic hydrolysis of esters by means of mineral acids (see art on Hydro-LYSIS), a complex is formed between the substance transformed (acid in esterification and ester in hydrolysis) and the catalyst This compound is probably an oxonium salt containing quadra-

valent oxygen, e.g. R C The salt. only small amounts of which may be present,

is ionised into (R CO<sub>2</sub>H,H) and Cl, and the esterification proceeds by the reaction of the complex cation with the alcohol (Acree and Jol nson, Amer Chem J. 1907, 38, 334), eg..

$${\scriptstyle (R~CO_2^+H,H)+C_2H_5OH=R~CO_2Et+H_2O+\overset{+}{H}}$$

and not with the ions of the alcohol as suggested This reaction is more rapid than by Lapworth the simple reaction between the acid and alcohol, and in this way the accelerating effect of the catalyst is accounted for

The retarding effects produced by small amounts of water are, according to Lapworth (Chem Soc Trans 1908, 93, 2167, 2196), due to the water competing with the organic acid for the catalyst, and giving use to the com-

plex ion (H2O,H), which is not catalytically active The fact that in alcohol containing water, the velocity of esterification at constant volume decreases more rapidly than the concentration of the catalyst (Goldschmidt and Uddy, Zeitsch physikal Chem 1907, 60, 735, Kailan, Monatsh 1906, 27, 543, 997, Annalen, 1907, 351, 186), is also held to be opposed to the view that a complex hydrion is formed with the alcohol

Work on the catalytic activity of strong acids tends to show that this activity is exerted in two ways · (a) by means of hydrions, (b) by the non-ionised acid, and that the effect due to the non-ionised acid is greater for highly ionised than for slightly ionised catalysts. methyl and ethyl alcohol the retarding influences of water on esterification is of much the same magnitude for different acids, and this is held to support the view that an intermediate compound of alcohol and catalyst plays a most important part in the change (Goldschmidt and Thuesen, Zeitsch. physikal Chem. 1912, 81,

Compare also Snethlage, ibid. 1913, 85, 211; 1915, 90, 91 and 139, Acree, Amer Chem J 1912, 48, 352; Dawson and others, Chem Soc. Trans. 1913, 103, 2135; 1914, 105, 1093, 1915, 107, 1426; 1916, 109, 1262; McBain and Coleman, ibid 1914, 105, 1517; H S Taylor Zottoch Electro Chem 1014, 203 Taylor, Zeitsch. Electro Chem. 1914, 20, 201.

(For the conversion of acid amides into esters, and for the transform tion of methyl into ethyl Vol. II .- T.

esters, both processes of alcoholysis, see ait HYDROLYSIS.) JJS

Trade name for basic aluminium ESTON. acetate. Used as an antiseptic powder in dressing wounds

ESTORAL v Synthetic drugs

ESTRAMADURITE v FERTILISERS

ETA (E) ACID. 1-Naphthylamine-3.8-disulphonic acid v NAPHIHALENE

ETHELEN. Trigallic acetal
ETHAL v WAYES
ETHYL C2H5 or CH3—CH2 is a univalent
basylous radical not known in the free state, the body  $C_4H_{10}$  or  $(C_2H_5)_2$  which was named othyl (afterwards diethyl) by its discoverer, Frankland (Chem. Soc. Trans. 2, 263), being now regarded as butane

Ethyl acetamide v Ethylamine (infra)

Ethyl acetate v ACETIC ACID.

Ethyl acetoacetate v Ketones

Ethyl alcohol v ALCOHOL

Ethyl antimonite (C<sub>2</sub>H<sub>5</sub>O)<sub>3</sub>Sb, bp 115°-120°, from action of C<sub>2</sub>H<sub>5</sub>OH upon Sb<sub>2</sub>O, in presence of anhydrous CuSO<sub>4</sub>, colourless liquid (McKey, Chem Soc Trans 1909, 95, 604)

Ethyl arsenite (C<sub>1</sub>H<sub>5</sub>O)<sub>3</sub>As. A liquid formed by action of C<sub>2</sub>H<sub>5</sub>OH upon As<sub>4</sub>O<sub>6</sub> in presence of anhydrous CuSO<sub>4</sub>, the yield being 4 5 p c (Lang, McKey, and Gortner, Chem Soc Trans 1908, 1908). 93, 1368) According to Crafts (Bull Soc ching 1870, ii 14, 99), it is obtained in very small yields by

(1) Action of Et<sub>3</sub>S<sub>1</sub>O<sub>3</sub> upon As<sub>4</sub>O<sub>6</sub> at 200°,

(2) Action of Ag<sub>3</sub>AsO<sub>3</sub> upon Etl at 150°,

(3) Action of AsBr, upon NaOEt Auger (Compt rend 1906, 143, 907) obtained a yield of 12 p c by heating EtOH and As4O6 in a sealed tube far above the boiling-point of the alcohol

Ethyl arsenate. A liquid boiling above 200° with partial decomposition. Decomposed by water immediately. Formed by heating together

water immediately. Formed by heating together in sealed tubes at  $100^{\circ}$  Ag<sub>3</sub>AsO<sub>4</sub> and Et1 (Crafts, Bull Soc chim 1870, ii 14, 99)

Ethyl boride (C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>B Tricthyl boride, Borethyl, b p 95°, sp gr 0 696 at 23°, V D at  $132^{\circ} = 3598$ ; at  $101.6^{\circ} = 3.76$ , calc = 3.399From action of  $Zn(C_2H_5)$ , upon  $(C_1H_5)_3BO$ ,, or from  $BCl_3$  and  $Zn(C_2H_5)_2$  (Frankland, Trans Roy Soc 1862, 167, Proc Roy Soc 1876, 25, 165)

A spontaneously inflammable, colourless liquid burning with green flame; decomposed by HCl, evolving  $\mathrm{C}_2\mathrm{H}_6$ , and forming  $\mathrm{B}(\mathrm{C}_2\mathrm{H}_6)$ , c'l is slowly decomposed by water, and destroyed by HNO3, with formation of boric acid Absorbs NH3, forming an only liquid NH3B(C2H5)3 of an aromatic odour, which is fairly stable even in contact with O  $B(C_2H_5)_3$  absorbs O and forms  $B(C_2H_5)_3O_2$ , bp 125° This, by the action of water, yields ethylboric acid  $B(C_2H_5)(OH)_2$ , a crystalline solid resembling naphthalene in appearance, can be sublimed at 40°, has a sweet taste, and acid action, but does not combine sweet taste, and action, but does not combine with bases By action of  $Zn(C_2H_5)_2$  upon this body we get  $B(C_2H_5)(CC_2H_5)_2$  and  $B(CC_2H_5)_3$ , bp  $112^\circ$ , V D. (dissoc) 2.78. By further action of  $Zn(C_2H_5)_2$  upon ethylboric acid we get  $B(C_2H_5)_2CC_2H_5$ ; bp.  $102^\circ$ , V D. = 3.914 Water converts this into  $B(C_2H_5)_2CH$ , a spontaneously inflammable liquid, which absorbs c:ygen, yielding B(C<sub>2</sub>H<sub>5</sub>O)C<sub>2</sub>H<sub>5</sub>OH; m.p. 8°.

Ethyl bromide  $C_2H_6Br$  Hydrobromic or Bromhydric ether; mp  $-116~0^\circ$  (Ladenburg and Krugel, Ber 32, 1818),  $-125~5^\circ$  (Schneider, Zeitsch. physikal Chem. 22, 233); bp. 38 22° (corr) (Thoipe and Rodger, Phil Trans. 1894, A, 11, 472), sp gr 1 4189 at 15° (Mendéleeff), l 4733 at 0° (Pierre), l 4685 at 13 5°; crit temp 236° (Pawlewski, Ber. 16, 2633)

Preparation — 1 By action of P and Br upon alcohol (Serullas, Ann Chim Phys [2] 34, 99) 40 parts red P and 160 parts absolute alcohol are slowly mixed with 100 parts Br, kept cool until all action is over, the mixture is then distilled, and the distillate purified by washing with water and rectification (Personne, Compt rend 52, 468) —2 Absolute alcohol is heated with Br and distilled, and the distillate is freed from excess of Br by washing with KOH and redistillation (Lowig, Annalen, 3, 288)—3 4 parts KBr and 5 parts of a mixture of 2 parts H<sub>2</sub>SO<sub>4</sub> and 1 part alcohol of 96 p c are distilled, and distillate purified as above (De Vrij, J. 1857, 441, Tabourg, Bull Soc chim 1911, [iv] 9, 124) Weston (Chem. Soc. Trans. 1915, 108, 1489) has given details of a modification of De Viij's method, using sodium bromide in place of potassium bromide and paying attention to the rate of heating, whereby yields of 80–90 p c of the theoretical amount of pure ethyl bromide can be obtained (cf. Holt, Chem. Soc. Trans. 1916, 109, 1.)—4. ('2H<sub>4</sub> in presence of AlBr<sub>3</sub> at 0° combines with HBr to form ('2H<sub>5</sub>Br. (Gustavson, J Russ Phys Chem Soc 16, 95) To prepare pure  $\rm C_2H_5Br$ , the commercially pure bromide is shaken with  $\rm H_2SO_4$  and allowed to separate It is then shaken with dilute alkali and distilled with 10 p c of sweet almond oil (Adrian, Chem Zentr 1895, 1 20)

Ethyl bromide is a colourless liquid with a sweet taste, and burning after-taste. Its vapour has an esthetic properties (Robin, Compt. 1 end 34, 649.) Webster, Bio. Chem. J. 1906, 1.328., Cole, J. Physiol. 29.) It is slightly soluble in water, and miscible with alcohol and ether. The vapour is decomposed by heat, yielding C<sub>2</sub>H<sub>4</sub> and HBr, or, if the temperature be higher, C. is deposited (Lowig). Burns in air with a green flame, evolving HBr. By introduction of Cl. two isomeric bodies of the composition C<sub>2</sub>H<sub>4</sub>(IBr, b. p. 84.5° and 104°-108°, and 2 isomerides of composition C<sub>2</sub>H<sub>3</sub>Cl<sub>2</sub>Br, b. p. 137°, 151° (Lescœur, Bull. Soc. chim. 1878, 29, 483) are obtained. At 100° Br. reacts, forming CH<sub>2</sub>Br CH<sub>2</sub>Br, whilst at 200° CH<sub>3</sub> (HBr<sub>2</sub>. is the product (Meyer and Muller, Ber. 24, 4247, J. pr. Chem. [2] 46, 161). By action of Br. and P. upon alcohol, we get, further, C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub>, b. p. 160°; C<sub>2</sub>H<sub>3</sub>Br<sub>3</sub>, b. p. 180°; C<sub>2</sub>H<sub>2</sub>Br<sub>4</sub>, b. p. 196°-200° (Bertrand, Finot, Bull. Soc. chim. 34, 28). (v. Ethylene bromide). Heated with AlBr<sub>1</sub>, yields successively C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub>, C<sub>2</sub>H<sub>2</sub>Br<sub>4</sub>, and C<sub>2</sub>Br<sub>6</sub> (Mouneyrat, Compt. rend. 127, 109). With ammonia it yields NH<sub>2</sub>(C<sub>2</sub>H<sub>5</sub>), HBr., and with ethylamine, diethylammonium bromide NH(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>HBr; similarly, with diethylamine, it yields N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>HBr.

Ethyl chloride C<sub>2</sub>H<sub>5</sub>Cl Chlorhydric or Hydrochloric ether; mp. -142 5° (Schneider, Zeitsch. physikal Chem 22, 233); b.p. 12 5° (Regnault, J. 1863, 67), crit. temp. 187 2; crit. press. 51·7 atm (Berthoud); sp.gr. 0 9214 at 0°; 0 9176 at 8°, 0 920 at 0° (Pierre), 0 £51

at b p. (Ramsay, Chem. Soc. Trans. 35, 470); S.V 75 8 (Kopp). (Robiquet and Colin, Ann Chim. Phys [2] 1, 343; Regnault, *ibid.* 71, 355, Kuhlman, Annalen, 33, 108; Lowig, Pogg Ann 14, 346, Gm 7, 367; Gerh. 11, 308) For a table of its vapour pressure between  $-30^\circ$  and  $+100^\circ$ , see Henning, J. Soc. Chem. Ind. 1920, 1. T.

Known as far back as 1500, and its preparation described by Glauter in 1648. From alcohol by the action of  $S_2Cl_2$  or metallic chlorides (Rouelle [1759]), from alcohol and HCl (Basse [1801]). Composition first ascertained by Robiquet and Colin—Also formed by action of Cl upon  $C_2H_6$  (Darling, Annalen, 150, 216), and of HCl upon ether in scaled tubes (Berthelot)

The action of HCl upon alcohol gives a very poor yield of C<sub>2</sub>H<sub>5</sub>Cl, unless ZnCl<sub>2</sub> is added to the alcohol before passing in HCl (droves, Chem Soc Trans 1874, 636) Kiuger (J pr Chem [2] 14, 193) recommends that a mixture of 1 part of ZnCl<sub>2</sub> in 82 parts of alcohol should be saturated with HCl gas in the cold, and then heated to boiling, HCl being passed in during distillation. A reflux condenser stops the alcohol vapour, and the C<sub>2</sub>H<sub>5</sub>Cl passes on and is collected in a well-cooled receiver.

The formation of  $\mathrm{C_2H_5Cl}$  in this process is partly due to the action of HCl upon alcohol, and partly to the umon of the nascent  $\mathrm{C_2H_4}$  with HCl (Schorlemmer, Chem Soc Trans 1875, 308), this view is sustained by the fact that if amyl alcohol be thus treated, both primary and secondary amyl alcohol are produced, and the latter can only be formed from the radical  $\mathrm{C_5H_{10}}$  Ethyl (bloride is also formed by the action of PCl<sub>5</sub> upon alcohol, and also, to some extent, by the action of Cl upon alcohol. Hence the formation of ethyl chloride in the manufacture of chloral

It is a colourless mobile liquid, has a pungent ethereal smell, and a sweetish taste, is inflammable and burns with a green-edged flame, evolving HCl. It is spaingly soluble in water (1 pirt in 3000), but mixes readily with alcohol and ether. It dissolves phosphorus, sulphui, fats, oils, and many resins. It combines with many metallic chlorides—eg SbCl<sub>5</sub>, Fe<sub>2</sub>Cl<sub>6</sub>—to form crystalline compounds. Mainly used for Ethylation in the manufacture of dye stuffs and drugs, and as a refingerant and solvent

It is used in medicine like ether, but, on account of its volatility, it is mixed with an equal volume of alcohol, and is then known as 'alcoholised muriatic ether, B.P.' It is decomposed at a red heat into  $C_2H_4$  and HCl, at higher temperatures carbon separates, and ('H<sub>4</sub>, H, and HCl are set free Strong HNO<sub>3</sub>, when boiling, liberates HCl and forms  $C_2H_5NO_3$  in small quantity (Thénard, Boullay) SO<sub>3</sub> absorbs it, forming SO<sub>3</sub>  $C_2H_5Cl$ , a fuming liquid, bp. 130°—probably the three isomerides  $C_2H_5O$  SO<sub>2</sub> Cl, CH<sub>2</sub> ClCH<sub>2</sub> SO<sub>2</sub> OH, and CH<sub>2</sub>(OH) CH<sub>2</sub>SO<sub>2</sub> Cl (Purgold, Zeitsch Chem 1868, 669; Ber. 6, 502). When vapour of  $C_2H_5Cl$  is passed through boiling  $H_2SO_4$ , HCl is evolved (Thénard), also  $C_2H_4$  and SO<sub>2</sub> (Boullay). When the vapour is led through a tube containing heated soda-lime, a mixture of CH<sub>4</sub> and H is evolved, and NaC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> and Na<sub>2</sub>CO<sub>3</sub> are formed (L. Meyer, Annalen, 139, 282), while, according to Dumas and Stas,  $C_2H_4$  and Macl are formed. On

AgNO<sub>3</sub>,C<sub>2</sub>H<sub>5</sub>Cl has no action until after prolonged contact, but even after some months only a very small quantity of AgCl is formed (Thénard, Boullay); but AgCl is rapidly thrown down when C2H5Cl is heated with a solution of AgNO, in sealed tubes (G. C. Foster)

Heated with bromine and non wire at 100° C<sub>2</sub>H<sub>5</sub>Rr and ethylene dibromide are formed (V Meyer and Petrenko-Kutschenko, Ber 25, 3307

NH, has only a slight action at ordinary temperatures, but on heating a mixture of NH3 and C2H5Cl, ethylene and a deposit of NH4Cl are formed (Dumas and Stas, Ann Chim Phys 73, When a mixture of ethereal NH, and C2H5Cl is exposed to sunlight, crystals of ethylamine hydrochloride separate (Stas) When heated with alcoholic  ${\rm NH_3}$  to  $100^\circ$  in sealed tubes,  $\overline{NH_2(C_2H_5)}$ ,  $\overline{HCl}$  is also produced, together with traces of  $\overline{NH(C_2H_5)}_2$ ,  $\overline{HCl}$  and  $\overline{N(C_2H_5)}_3$ ,  $\overline{C_2H_5}$  (Groves, Chem Soc Trans. 1861, 13, 331) KOH slowly removes Cl from C<sub>2</sub>H<sub>5</sub>Cl, yielding (in aqueous sol ) alcohol and KCl Alcoholic KOH and C<sub>2</sub>H<sub>5</sub>Cl heated to 100° in sealed tube deposits KCl and forms ( $C_2H_5$ )<sub>2</sub>O (Balard, Ann Chim Phys [3] 12, 302). Passed into heated sol of  $K_2$ °, ethyl sulphide is formed, with alcoholic KHS, mercaptan is formed (Reg-

Chlorine in the sunlight yields di-, tri-, tetra-, penta-, and hexa- chloro- derivatives of ethane (Regnault), of the dichloro- derivatives CH3 CHCl2 is the main product. If, however, ethyl chloride is heated with a halogen carrier, CH2CI CH2CI is the chief product (D Ahns and Kautzsch, J pr Chem [2] 1909, 80, 305, Staedel ibid 303; Ber 6, 1403, Meyer and Muller, J pr Chem [2] 46, 161)

Trichlorethane C<sub>2</sub>H<sub>3</sub>Cl<sub>3</sub> 1 Chlorethylene chloride CH<sub>2</sub>Cl CHCl<sub>2</sub>, b p 115° (Pierre, Annalen, 80, 127), 113 7° at 760 (Stadel, Ber 15, 2563), 113 5°-114° at 753 2 (Schiff, Annalen, 220 97), sp gr 1 4223 at 0° (P), 1 4577 at 9 4°/4°, 1 2945 at 113 5°/4° (Schiff, 1 4406 at 25 6°,  $\mu_{\rm A} = 1$  4719 at 22°  $R_{\infty}=42.26$  (Kanonnikow), from  $C_2H_3Cl$  (Regnault, Ann Chim Phys [2] 69, 151), or CHCl, CH, (Meyer and Muller, J pr Chem [2] 46, 174), and SbCl.

2 Ethenyl trublorude ('H<sub>3</sub> CCl<sub>3</sub>; bp 74 5° (Geuther, J 1870, 435), 74 1° at 760 mm (Stadel, Ber 15, 2563), spgr 1 3465 at 0° (Pierre, Annalon, 80, 127), 1 372 at 16° (Regnault, Annalen, 33, 317), 1 3657 at 0°, 1 3249 at 26°;  $\mu_{\rm A} = 14199$  at 21°,  $R_{\rm C} = 42$ 0 (Kanon-level),  $R_{\rm C} = 42$ 0 (Kanon-le nikow); by the chlorination of ethyl chloride (Regnault, ibid 33, 317) Decomposed by sodium ethoxide, giving C2H2Cl·O C2H5, C2H4O2,

and C<sub>2</sub>H<sub>3</sub>(OEt)<sub>3</sub>.

Tetrachlorethane C<sub>2</sub>H<sub>2</sub>Cl<sub>4</sub>.

 $1~as\text{-}\mathrm{CH_2Cl}~\mathrm{CCl_3}$ , b.p.  $135^\circ$  (Regnault, Ann Chim Phys [2] 69, 162), 138 6° (Pierre, Annalen, 80, 130), 135 1° (corr) (Geuther, Brockhoff, J. 1873, 317); sp gr 1 6116 at 0° (P), 1 576 at 19° (R), 1 5825 at 0°, 1 5424 at 26°,  $\mu_{\rm A}$ ab  $t^{\circ} = 14772 - 0000437t$ ;  $R_{\infty} = 5072$  (Kanonnikow). Formed by chlorination of ethyl chloride (Laurent, Annalen, 22, 292), or action of Cl upon CH<sub>2</sub>Cl CHCl<sub>2</sub> (Regnault), by heating at 110° acetylene tetrachloride and AlCl<sub>3</sub> (Mouneyrat, Bull. Soc. chim. 1898, 19, 499).

Decomposed by sodium ethoxide

Decomposed by somular convaries  $C_2HCl_2 O C_2H_5$  and  $CH_2(OC_2H_5)CO_2Na$ . 2. s-CHCl<sub>2</sub> CHCl<sub>2</sub>; b p 147 2°, sp gr. 4°/4° =1 6208, 17°/17° =1 6013,  $\mu_D$  at 17° = 1 495587 $\pm$ 0 000006 (Veley, Proc Roy. Soc. 1910, B 82, 217), R $\propto \pm$ 50 6 (Kanoninkow). From C2H2 and SbCl5 (Berthelot, Jungfleisch, Annalen, Suppl 7, 254), from dichloraldehyde and PCl<sub>5</sub> (Paterno, Pisati), from ethylene chloride and AlCl<sub>5</sub> at 75° if Cl and C<sub>2</sub>H<sub>5</sub> be led into the mixture (Mouncyrat), by long heating

yields HCl and C<sub>6</sub>Cl<sub>6</sub>

If S<sub>2</sub>Cl<sub>2</sub> and I p c of its weight of reduced iron be saturated with (I, and if the excess (I be removed with CO<sub>2</sub>, and then dry acetylene be passed in at a temperature of 40°, and the cycle repeated until no more C<sub>2</sub>H<sub>2</sub> is absorbed, sym-tetrachloroethane, pentachloroethane, and hexachloroethane are produced, the two latter m small quantity (Michel, Zeitsch angew Chem 1906, 19, 1095) The sym-tetrachlorethane is distilled off from the mixture, either alone or ın steam (D. R. P. 174068), Chem. Soc. Absti 1906, 1 169)

Pentachlorethane CHCl<sub>2</sub> CCl<sub>3</sub>, b p 159 t° (corr), 161 7° at 760 mm (Stadel), mp. 18, sp.gr. 1 7089 at 0°, 1 6926 at 10 15 '/4" (Therpe, Chem Soe Trans 37, 192),  $\mu_{\Lambda}=1.4871$  at 251°,  $R_{\infty}=59.05$  (Kanonnikow) By action of chlorine upon C2H3Cl also from chloral and PCI, (Paternò, Annalen, 151, 117), or AlCI, (Mouneyrat, Bull Soc chim 1898, 19, 260)

Hexachlorethane CCLCCL trichloride, art CARBON)

(For mixed halogen derivatives of ethane, see Mouneyrat, Bull Soc chim 1898, 19, 499, Swartz, Rec trav chim 17, 234 and 235, Bull Acad Roy Belg 36, 519, ibid 1909, 60 and 728, ibid 1899, n 281, Elbs and Newmann, J. pt. Chem 1898, 58, 245, Biltz, Bci 1902, 35, 3524, Lemoult, Compt. 1end. 1903, 136, 1333. Besson, Bull. Soc. chim. [3] 11, 920. Nef, Annalen, 298, 353, 354, 358, and 360, Thiele and Peter, dud 369, 149 Lawrie, Amer Chem J 1906, 36, 487)

Ethyl isocyanate. F theyl  $C_2H_5N$  C O, bp 60°, sp gr 0.8981, heat of formation (Lemoult, Compt rend 1898, 126, 43); absorption spectrum (Hartley, Dobbie and Walker, Chem Soc Trans 1901, 856). By the action of dry KCNO upon  $\mathrm{KC_2H_5SO_4}$  at 180°-250° (Wurtz, Ann Chim Phys 1854 [3] 42, 43), distillate is a mixture of isocyanate and isocyanurate, which can easily be separated by By passing CNBr into absolute distillation alcohol and ether (Ponomarew, Bei 1882, 15, 515, Mulder, Rec trav chim 1, 210, 2, 133) by passing CNBi (Mulder, ibid 3, 306) or CNC1 (Cloez, Compt rend 44, 428, Annalen, 102, 355) into C<sub>2</sub>H<sub>5</sub>ONa in absolute alcohol, by the action of diazomethane on sym-ethylmtrousea (Degner and Pechmann, Ber 1897, 30, 653), by heating the mercuic chloride derivative of ethyl thiolcarbamate (Anschutz, Annalen, 1908, 359, 202)

In 1895 Nef (ibid 287, 310) showed that the compound hitherto known as ethyl cyanate was ethyl isocyanate. Ethyl isocyanate is a mobile It is decomliquid having a suffocating smell posed by water into diethylcarbamide, and by ammonia into ethylcarbamide

Ethyl cyanide. Propionitrile  $C_2H_3C:N$ ,

m.p -103 5° (Schneider, Zertsch. physikal. Chem 22, 233), bp. 98 1° (corr.) (Engler, Annalen, 133, 153), 97 08° (corr.) (Thorpe, Chem Soc. Trans 37, 205), sp. gr. 0 8010 at 0°, 0 7938 at 7 36°/4° (Thorpe), 0 7015 at 97°/4° (Schiff, Ber 19, 567), H. C. 440 7 Cals (Berthelot and Petit. Compt. rend. 108, 1217). Latent lot and Petit, Compt rend 108, 1217), latent heat of vapoursation (Louguinine, Chem Zentr 1900, 1 451), molecular refraction 25 10 (Bruhl, Zeitsch physikal Chem 16, 214; Eykman, Rec trav chim 12, 172, Costa, J. 1891, 336); magnetic rotation (Perkin, Chem Soc. Trans 55, 701), cut temp 2912°, crit. density 0 2401 (Ter-Gazaman, J. Chim. Phys. 1906, 4, 140)

From Ba(C<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>)<sub>3</sub> and KCN (Pelouze, Annalen, 10, 240), from cyanogen chloride and Zn(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> (Gal, Zeitsch Chem 1868, 252), from Zn(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> and (CN)<sub>2</sub> (Frankland and Graham, Chem. Soc. Trans. 37, 740), from proponamide and P<sub>2</sub>O<sub>5</sub> (Dumas, Malaguti and Leblanc, Annalen, 74, 329) By heating ethyl socyanide to 240° (Nef, ibid 280, 296)

Preparation—By heating C<sub>2</sub>H<sub>5</sub>I with pure KCN and alcohol to 180°, C<sub>2</sub>H<sub>5</sub>Cl dissolved in three times its weight of alcohol is heated with KCN to 100° or 150° (Rossi, Annalen, 159, 79) C2H5I is heated with KCN in closed tubes to 180°, and the product distilled, and the distillate is freed from excess of C2H51 by means of a solution of CaCl2, upon which the C2H5CN swims, while the heavier  $C_2H_5I$  sinks to the bottom (Williamson, Phil Mag [4] 2, 205, Gauthier, Ann Chim Phys [4] 17, 180)

It is a mobile colourless liquid with an agreeable odour, soluble in water, from which CaCl, or NaCl causes it to separate, miscibility with water (Rothmund, Zeitsch physikal Chem 26, 453), V D 1 928 It combines with the hydracids, e q HCl, the product  $C_3H_5NHCl$ forms monoclinic prisms soluble in water and melting at 121° (Gauthier) HCl gas passed into a well-cooled solution of the cyanide in acetic acid gives acetyl chloride and C<sub>2</sub>H<sub>5</sub>CONH<sub>2</sub>,HCl (Colson, Bull Soc chim [3] 17, 57) Chlorine combines with C<sub>3</sub>H<sub>5</sub>N to form dichlorpropionitile CH<sub>3</sub> CCl<sub>2</sub> CN, a liquid body, b p 103°-107°, which polymerises on standing to a solid  $(C_3H_3Cl_2N)_3$  Propionitrile also forms compounds with metallic chlorides and with cyanogen chloride C<sub>3</sub>H<sub>5</sub>N CNCl is a liquid decomposed by water, b p 60°-68° (Hencke, Annalen, 106, 280) Compounds with AlCl<sub>3</sub> (v Perrier, Compt rend 1895, 120, 1423) Propionitrile is hydrolysed by acids or alkalis yielding propionic acid, electrolytic reduction yields n-propylamine (Ahiens, Zeitsch Elektrochem 1896, 3, By the action of metallic sodium, polymerisation ensues, the products being sodium cyanide, ethane, and α-iminopropionylethyl cyanide (Meyer, J. pr. Chem. [2] 38, 326, 45, 63)

Ethyl isocyanide, Ethylcarbylamıne C<sub>2</sub>H<sub>5</sub>N C; b p 79°; sp gr. 0 759 at 4°, 0 7150 at 44 5° (Gauthier, Ann Chim Phys. [4] 17, 233); molecular refraction 16.26 (Costa, Gazz. chim. ital 22, i 104) By the interaction of C<sub>2</sub>H<sub>5</sub>I and AgCN (Gauthier, Annalen, 151, 239), by distilling KCN with KC2H5SO4, propionitrile also being formed; by digesting ethylamine with chloroform and alcoholic potash (Hofmann, ibid. 147, 107).

It is a colouriess liquid having a very repulsive smell. It is transformed into propionitiile by heating at 240°, by heating with alcoholic H.S ethylthioformamide is produced; with sulphur and CS, ethylthiocarbimide; combines with CNCl, with C<sub>2</sub>H<sub>5</sub>Cl at -10° yields C<sub>2</sub>H<sub>5</sub>N(Cl)O C<sub>2</sub>H<sub>5</sub> (Nef, *ibid*. 280, 296). HgO oxidises it to C<sub>2</sub>H<sub>5</sub>N . C: O. Yields a chloride C<sub>2</sub>H<sub>5</sub>N (Cl<sub>2</sub>) by treatment with SO<sub>2</sub>Cl in ether at -15°, an oil, bp 102° (Nef, *lc*) Yields a bromide, C<sub>2</sub>H<sub>5</sub>N (Br<sub>2</sub>, with bromine in This gives ethylamine dibromide with water, ethyl isocyanide dibromide hydrobromide and ethylthicarbimide with H,S, ethyl isocyanide dibromide hydrochloride with dry HCl, an only ethnodide with ethyl rodide, and diphenylethylguanidine with aniline (Guillimand, Bull Soc chim 1904, in 31, 605, 1905, in 33, 652) Compounds with cobaltous, ferious, and ferric chlorides (v. Hofmann and Bugge, Ber. 1907, 40, 3759) (For further reactions, v. Nef, l.c)

Ethyl derivatives of cyanuric acid. Although cyanuric acid exists in only one modification, probably the 180-(11) esters derived from the tautomeric forms (I. and II.) are known.

Ethyl cyanurate (C<sub>2</sub>H<sub>5</sub>OCN)<sub>3</sub>, mp 29°, b p 275° (Claesson, J. pr Chem [2] 33, 131) By action of CNCl on C<sub>2</sub>H<sub>5</sub>ONa (Hofmann. By action of CNCI on C<sub>2</sub>H<sub>5</sub>ONa (Hofmann. Ber 19, 2061), also from C<sub>2</sub>H<sub>5</sub>ONa (3 mols) and trimethyl cyanurate (H), from CNI of CNBr and C<sub>2</sub>H<sub>5</sub>I (Ponomarew, Bull Soc chim [n] 41, 315), by heating the lead salt of the diethyl ester (P.), together with ethyl isocyanurate by treating silver cyanurate with C<sub>2</sub>H<sub>5</sub>I (C, P); from di-potassium cyanurate and C<sub>2</sub>H<sub>5</sub>I (P)

Oily liquid readily soluble in alcohol, ether, CHCl3 and CS2, sparingly so in water Converted into ethyl isocyanurate by long heating at  $180^{\circ}-200^{\circ}$ , yields cyanuric acid with HCl, cyanuric chloride with PCl<sub>5</sub>, melamine with strong ammonia and the diethyl ester with

baryta water

Diethyl cyanurate, Diethylcyanuric acid HO(CN)<sub>3</sub>(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> (Ponomarew, Ber 18, 3267, Hofmann, *ibid* 19, 2077) By warming ethyl cyanurate with concentrated baryta water, by treating methyl cyanulate or mono-potassium cyanurate with C2H5I

Thick plates from water, insoluble in ether, sparingly soluble in alcohol Decomposes on heating into diethyl isocyanurate, yields cyanuric acid on warming with HCl. Pb salt and  $C_2H_5I$  yield triethyl cyanurate, but if C<sub>2</sub>H<sub>5</sub>I contains free I, triethyl isocyanurate also formed.

Ethyl isocyanurate, Triethylcarbimide (C<sub>2</sub>H<sub>5</sub>NCO)<sub>3</sub>, mp 95°, bp 276°, absorption spectrum (Hartley, Dobbie and Walker, Chem. Soc Trans. 1901, 849), heat of formation (Lemoult, Compt rend. 125, 869) By polymerisation of  $C_2H_5NCO$  (H. D and W.); also from potassium cyanate or cyanurate, or from

Pb salt of the diethyl ester and KC<sub>2</sub>H<sub>5</sub>SO; parts of H<sub>2</sub>SO<sub>4</sub>, 5 parts H<sub>2</sub>O, and 15 parts of by heating ethyl cyanurate at 180°-200°; alcohol (85 pc) (Stinde, Dingl. poly. J 181, together with ethyl cyanurate from silver cyanurate and C2H I (Habich and Limpricht, Annalen, 109, 102)

Rhombic prisms (Rammelsberg, J. 1857, 273; Fock, Ber 19, 2076) moderately soluble in hot water, readily so in alcohol. KOH yields N(C<sub>2</sub>H<sub>5</sub>)H<sub>2</sub> and K<sub>2</sub>CO<sub>3</sub>.

Diethyl isocyanurate, Diethylcyanuric acid H(NCO)<sub>3</sub>(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>, mp 173° Obtained in preparing the triethyl ester (Habich and Limpricht, Lc); together with cyanuric acid by heating anhydrous mono-potassium cyanurate with  $C_2H_5I$  at  $170^\circ-180^\circ$  (Ponomarew, Ber 18, 3270); by heating diethyl cyanurate (Hofmann, ıbıd. 19, 2078)

Hexagonal columns, readily soluble in hot water, alcohol, and ether, sparingly so in cold water. Yields  ${\rm CO_2}$  and  ${\rm N(C_2H_5)H_2}$  by heating in a scaled tube at 100°. Pb salt and  ${\rm C_2H_5I}$ yield ethyl isocyanurate

Ethyl diketobutyrate v Ketones
Ethyl ferrocyanide (C<sub>2</sub>H<sub>7</sub>)<sub>4</sub>Fe(CN)<sub>6</sub> From
Ag<sub>s</sub>Fe(CN)<sub>6</sub> and C<sub>2</sub>H<sub>5</sub>I (Freund, Ber 21, 935)
Rhombic crystals from CHCl<sub>2</sub>, easily soluble in water and alcohol, insoluble in ether, ligroin, and  $\mathrm{CS}_2$  Decomposed at  $212^\circ-214^\circ$ , with formation of ethyl isocyanide. The aqueous solution gives a deep violet tint with FeCl3, and a white precipitate with HgCl<sub>2</sub> Molecular weight by eryoscopic method indicates the monomolecular formula (Buchbock, Zeitsch physikal Chem

Ethyl fluoride C,H,F Obtained impure by action of CaF, and H,SO, upon C, H,OH (Reinsch, J pr Chem 1840, 19, 314); by heating KC<sub>2</sub>H<sub>5</sub>S<sub>4</sub> and KFHF (Fremy, Compt rend 1854, 38, 393) Prepared pure from AgF and C,H, I (Moissan, Ann Chim Phys 1890, [6] 19, 272) Colourless gas with ethereal odour, liquefying at  $-32^{\circ}$  under atmospheric pressure Burns in air with a blue flame Readily soluble in  $C_2H_5Br$ ,  $C_2H_5I$ ,  $C_2H_5OH$ , and other Decomposed at  $100^\circ$  by heating with KOH in a scaled tube into KF, C, H, ()H, and ether

Ethyl formamide H CONHC, H<sub>5</sub>; bp 199°, sp gr 0 952 at 21° By distilling an aqueous solution of H COONH<sub>3</sub>(C<sub>2</sub>H<sub>5</sub>); separates from distillate upon addition of KOH (Linnemann). By action of NH<sub>2</sub>(C<sub>2</sub>H<sub>5</sub>) upon CCl<sub>3</sub> CHO (Hofmann, Ber. 5, 247); a thick colourless

Ethyl formate H CO<sub>2</sub>C<sub>2</sub>H<sub>5</sub>; b p 54 4° at 760 mm., sp.gr 0 9376 at 0°/4°; b p 53 4°-753.6° at 754.4 mm, sp.gr. 0.8731 at  $53.4^{\circ}/4^{\circ}$  (R. Schiff, Annalen, 220, 1063), b p at 760 mm 54.3°, sp.gr. 0.9480 at  $0^{\circ}/4^{\circ}$ , critical temperature, 235.3°; critical pressure, 35,550 mm (Young and Thomas, Chem. Soc. Trans. 1893, 1003) 1193). Capillarity constant at boiling-point, a2 = 4 528 (Schiff, Annalen, 223, 75); H C p 388 cals. (Berthelot, Ogier), 391 7 (Delépine)

By distilling a mixture of 6 parts alcohol (90 o c), 7 parts sodium formate and 10 parts (conc) H<sub>2</sub>SO<sub>4</sub> (Liebig), by distilling a mixture of ethyl oxalate with oxalic acid (Lowig, J 1861, 599), by heating a mixture of glycerol, oxalic acid, and alcohol in a reflux condenser, and then distilling (Lorin, Bull. Soc chim [ii ] 5, 12) On the large scale a mixture of 9 parts of starch and 29 parts of MnO<sub>2</sub> (of 85 pc) is heated with 20

- A mobile colourless liquid with a pleasant aromatic smell; is used for flavouring arrack and rum; is decomposed by water, for which it has strong affinity Sodium in absolute ether decomposes it into CO, H and C<sub>2</sub>H<sub>5</sub>ONa (Freer and Sherman, Amer Chem. J 18, 581) Cl forms with it in sunlight dichlorethyl formate H CO2C2H3Cl2, a colourless only liquid, decomposed on heating, and which by action of alkalis is decomposed into acetic, formic, and hydrochloric acids—By further action of (I in sunlight, perchlorethyl formate Cl CO2C2Cl5, b p 200°, sp gr 1 705 at 18°, is formed (Clove, Ann Chim Phys [3] 17, 297)

Ethyl hydrate v ALCOHOL

Ethyl hydride, Ethane, Dimethyl  $C_2H_6$  (Frankland and Kolbe, Chem Soc Trans 1, 60), m p. -171 4°; b p at 749 mm -85 4° (Ladenburg and Krugel, Ber 32, 1821), -172 4° and -84·1° (Cardoso and Bell, J Chim Phys 1912, 10, 497), -88 5° (Mass and McIntosh, J. Amer Chem. Soc. 1914, 36, 737), -89 3° (Burrell and Robertson). Gas: sp gr 1 036, condensed at 4° and 46 atmospheres pressure (Calletet, J Annalen, 282, 245; Olszewski, Ber 1894, 27, 3306, and Keunen, Chem Zentr 1897, 11 540; Cardoso and Bell, lc), heat of vaporisation 146+1010 ergs per mol; density at the bp 0 5490 (Maass and McIntosh, lc); sol in water at  $t^{\circ}$ 

(1) 
$$\frac{94556-035324t+0006278t^2}{100}$$
  
(2)  $8710-033242t+000603t^2$ 

(1) Schickendants (Annalen, 109, 116), (2) Bunsen (Gasometry) Sol in alcohol=1.5 (Berthelot, J 1867, 344); H (' p 373 3 ('als. (Th)=372 3 (Berthelot, Ann (him Phys [6] 30, 559); H.F (from amorphous C)=11.7Cals (B); occurs in crude petroleum

Pittsburg gas consists of ethane mixed with a little CH<sub>4</sub> and CO<sub>2</sub>, and is used for smelting

and lighting purposes.

By action of K upon C<sub>2</sub>H<sub>5</sub>(N, also by action of water upon Zn(C<sub>2</sub>H<sub>3</sub>), (Frankland and Kolbe, Chem Soc Trans 3, 338); by action of Zn upon C<sub>2</sub>H<sub>5</sub>I in presence of water; by electrolysis of acetic acid or acetates (Kolbe, Annalen, 69, 279), by heating 9 parts of  $C_2H_5I$  with 2.6 parts of AlCl<sub>3</sub> to 140° or 150° (Kohnlein, Ber 16 562); by action of  $Hg(C_2H_5)_2$  upon  $H_2SO_4$  (Schorlemmer, Annalen, 132, 234); by heating an excess of BaO2 with acetic anhydride and sand (Schutzenberger, Zeitsch Chem 1865, 703; Darling, Annalen, 150, 216), by the action of nascent H (from ZnCu couple or Zn dust) upon a mixture of C<sub>2</sub>H<sub>5</sub>I and C<sub>2</sub>H<sub>5</sub>OH (Sabanejeff, Ber 9, 810). By combining C<sub>2</sub>H<sub>4</sub> or C<sub>2</sub>H<sub>2</sub> with H in presence of reduced Ni (Sabatier and Senderens, Compt. rend 124, 1360; 128, 1173) Ethane is now produced on an industrial scale by means of this reaction carried out under pressure details of the process, see Sprent (J Soc Chem Ind 1913, 171) From EtI and potassium hydride in sealed tubes at 180°-200° (Moissan, ibid 134, 389) Reduced Ni at 325° breaks ethane up into C and CH4 (Berthelot)

Ethyl iodide  $\rm C_2H_5I$  (Gay-Lussac, Ann. Chim. Phys [1] 91, 89, Serullas, *ibid.* [2] 25, 323 and 42, 119; Frankland, Chem Soc Trans 2, 263; 3, 32), mp -118° (Schneider); bp 72·34° (Linnemann, 70.770), bp. 71·3° (Andrews), 72 57° (corr ) (Thorpe and Rodger), 71 9°-72° (Lanebarger); sp gr 1 9444 at 14·5° (L), 1 9755 at 0° (Pierre), 1 9309 at 15° (Mendeléeff), sp gr 1 1810 at 72 2°/4° (Schiff), sp gr 1 9653 at 4°, 1 9433 at 15°, 1 9243 at 25° (Perkin), V D 5 417 (Marchand)

From alcohol and HI (Gay-Lussae); from alcohol, iodine, and phosphorus (Sciullas). Personne (Compt rend 42, 468) suggested the use of red P, and, according to Beilstein (Annalen, 126, 250), I part 1cd P and 5 parts of alcohol (90 pc) are placed in a retort, and 10 parts of iodine slowly added After 24 hours the C2H5I is distilled off, washed with NaOH solution and with water, and dried over CaCl, Cusmer (Ber 1884, 649) dissolves I in liquid paraffin adds yellow P and then alcohol Walker (Chem Soc. Trans 1892, 717; cf ibid 1905, 1596) obtained a 93 pc yield from alcohol, iodine, and equal parts of ied and yellow phosphorus (cf Adams and Voorhees, J. Amer Chem Prepared in quantitative Soc 1919, 41, 789) yield by adding Et<sub>2</sub>SO<sub>4</sub> slowly to a warm con-centrated aqueous solution of KI (Weinland and Schmid, D. R. P. 175209). Also from action of HI upon ether (Cottrel Rogers), and together with iodoform and NaI by electrolysis of a solution of 20 grams C<sub>1</sub>H<sub>5</sub>COONa, and 20 grams KI in 20  $\epsilon \epsilon$  of water (Miller, Hofer, and Reindel)

C2H5I is a colourless, pleasantly smelling liquid, almost insoluble in water, but miscible with alcohol and ether. It precipitates silver from its solutions in the cold, easily forms double compounds with silver salts, metals, and ammonia Br displaces the I, and forms C, H, Br, Cl forms C<sub>2</sub>H<sub>3</sub>Cl (Dumas, Stas, Annalen, 35, 162) ICl forms C<sub>2</sub>H<sub>5</sub>Cl and I<sub>2</sub> HCl has no action, but, on the contrary, C<sub>2</sub>H<sub>5</sub>Cl is decomposed by HI into C<sub>2</sub>H<sub>5</sub>I and HCl (Lieben, Zeitsch ('hem 1868, 712) By action of HI at 150°, ethane and iodine are formed HNO3 also causes the separation of I (Marchand, J pr Chem 1884, 33, 186) Metallic Ag abstracts I, and leaves butane (C<sub>4</sub>H<sub>10</sub>) By heating with water to 150°, we get alcohol (Reynoso, J 1856, 567), with excess of alcohol yields ether and some C, H<sub>4</sub> (Lieben, Rossi, Annalen, 158, 166). Is gradually turned brown by exposure to light, owing to formation of free I and liberation of butane (Frankland, Chem Soc Trans 3, 322) In presence of mercury the reaction is much more rapid, owing to combination of the liberated I with Hg is used largely for the preparation of other ethyl compounds

Ethyl nitrate C<sub>2</sub>H<sub>5</sub>O NO<sub>2</sub>, Nitric ether; mp -112° (Schneider), bp 86 3° at 728°4 mm; sp gi 1 1322 at 0°, 1°1123 at 15 5° (Kopp, analen, 98, 367), bp 87 6°; sp gi. 1 1305 at 4°, 1 1159 at 15°, 1 1044 at 25° (Fekin), H F p = 40,870, H F v = 38,750, H C = 325, 690 (Th 4, 214) Obtained from alcohol by the action of nitric acid in presence of some substance which will destroy nitrous acid, eg. urea (Millon, Ann Chim. Phys [3] 8, 233).

Preparation —400 grams of pure mitric acid (sp gr 1 40, previously heated with 1 pc of urea in order to destroy lower exides of nitrogen)

are mixed with 300 grams absolute alcohol and 100 grams of urea nitrate, and distilled; when half the liquid has distilled off, a mixture of 400 grams of nitric acid and 300 grams of alcohol is allowed to drop into the retort and the distillation continued (Lossen, Annalen, Suppl. 6,

Silver nitrate, heated with ethyl iodide (or bromide) and alcohol, yields, not ethyl nitrate, but the nitrite, together with aldehyde (Bertrand, Bull Soc chim 33, [n ] 566) If, however, EtI be added gradually to powdered AgNO<sub>3</sub>, kept cool, a theoretical yield of EtNO<sub>3</sub> is formed (Von Biron, J Russ Phys Chem. Soc 30, 667).

Ethyl nitrate is a colourless liquid with pleasant odour and taste; almost insoluble in water, but miscible with alcohol and ether Ammonia forms ethylamine nitrate; nascent hydrogen yields hydroxylamine and other bases (Lossen). Sulphuietted hydrogen yields mercaptan (Kopp, Annalen, 64, 320). Alcoholic potash decomposes it with separation of crystals of potassium nitrate

Ethyl nitrite C<sub>2</sub>H<sub>5</sub> O NO, Nitrous ether, b p 16 4° (Liebig), 17° (Mohr, J 1854, 561, Brown, The Harm J 15, 400), 17 5° (Dunstan and Dymond, *ibid* 18, 861), sp gr. 0.90° at 15.5° (Br); 0.917 at 0°/0° (D and D), H.F.p. = 30,610, H.F.v. = 28,870 (Th. 4, 217).

This body was discovered in 1681 by Kunckel Obtained from alcohol by action of nitric acid, especially in presence of some oxidisable matter,

eg copper

Preparation —500 grams of potassium nitrite and 1 litre alcohol (45 pc) are mixed and cooled by means of ice, a mixture of sulphuric acid (500 grams), water (500 grams), and alcohol (500 grams) is then slowly dropped in, the heat produced volatilises the ether, which is con-densed in an efficient condenser (Feldhaus, Annalen, 126, 71) Dunstan and Dymond Dunstan and Dymond (lc.) recommend that 34 5 grams sodium nitrite in 120 grams water be slowly mixed with a mixture of 13 5 grams of sulphuric acid with 32 grams alcohol and about 75 c c water kept cool by ice The ethyl nitrite separates out as a layer floating on the surface As thus prepared, the liquid contains only traces of alcohol, from which it is freed by shaking with water Alkali ethyl sulphates react with alkali nitrites, yielding nitrous ester and nitroethane (Ray and Neogi, Also formed by Chem Soc Trans 1906, 1900) the interaction of MgSO<sub>4</sub>, NaNO<sub>2</sub>, and C<sub>2</sub>H<sub>5</sub>OH (Matuschek, Chem Zeit 1905, 29, 115)

Ethyl nitrite is a mobile liquid with a pleasant ethereal smell and taste, hable to undergo decomposition, especially in presence of water It can be preserved if mixed with alcohol and a small quantity of glycerol. It is decomposed by sulphides in accordance with the following

equation:

 $C_2H_5NO_2+3H_2S=NH_3+C_2H_6O+H_2O+3S$ 

(Kopp, Annalen, 64, 321)

It is slightly soluble in water and miscible with alcohol Mixed with alcohol it constitutes the 'sweet spirit of nitre,' spiritus ætheris nitrosi, of the British Pharmacopæia, which is thus prepared. To 1000 millilitres of alcohol (90 p.c.) add gradually 100 millilitres of sulphuric acid, and then 125 millilitres of nitric acid, stirring Transfer the mixture to a retort or constantly

flask, in which 100 grams of copper in fragments have been placed, and to which is attached an efficient condenser with a receiver containing 1000 millultres of 90 p c alcohol Surround the receiver with ice-cold water and distil gently at a temperature which is at first about 77°, and rises to 80°, but does not exceed 82°, until the volume of liquid in the receiver has been increased to 1600 millilitres Then allow the contents of the retort to cool, add 25 millilitres of nitric acid, and resume the distillation until the liquid in the receiver is increased to 1700 millilitres. Mix this liquid with 1000 millilitres of the alcohol so as to make the whole contain 2 66 p c. of ethyl nitrite.

Ethyl oxalylsuccinate v Ketones

Ethyl oxide  $(C_2H_5)_2O$ , Ether, Ethyl ether, Vinic ether, formerly called Sulphuric ether, Phosphone ether, &c., according to the acid used in its preparation, mp = 117.6° (Archibald and Melntosh), mp = -112.6° (Ladenburg, and Krugel), bp 34.97° (Regnault), 34.6° at 762 mm (Schiff, Annalen, 220, 332), 34.9° (Kopp), sp gr 0.7360 at  $0^\circ$ , 0.7185 at 17.5° (Hager), 0.6950 at 34.6°/4° (Schift), 0.7024 at 15°, 0.7099 at 25°/25°, 0.7201 at 15°/15° (Perkin, Chem S.c. Trans 45, 474), sp gr 0.71982 at 15°/15° (Wade and Finnemore), sp gr 0.7183 at 17.1° (Bruhl), sp gr 0.70942 at 25° (Linebarger), MM = 4.777 at 20°, freezes at -129° to a crystalline mass, which melts at -117.4° (Olzewski, Monatsh 5, 128), capillarity constant at bp  $a^2 = 4.521$  (Schiff), S at 17.5° = 8, critical temperature, 194°, critical pressure, 35.61 atmospheres (Ramsay and Young) Vapour tension 67.49 mm at  $-20^\circ$ ; 182.34 mm at  $0^\circ$ , 286.4 mm at  $10^\circ$ ; 433.2 at 20°, 636.3 at 30°.

The preparation of ether from alcohol and sulphuric acid was first described by Valcrus Cordus (1540), and the method of making this 'oleum vitroli dulce' was published in 1552 by Conrad Gessner Frobenius, 1730, named it sulphuric ether Val Rose (Scher. J. 4, 253) showed that it contained no sulphur, but its composition was first made out by Saussure and also by Dumas and Boullay (Ann Chim Phys [2] 36, 294) Its constitution was first studied by Liebig (Annalen, 60, 31, 23, 39) and Regnault (Ann Chim Phys [2] 71, 352) Williamson, in 1850-51 (B. A. 1850, 65, and Phil Mag. [3] 37, 350), established the views now held as to its constitution and formation.

Formation — From alcohol, by action of phosphoric or arsenic acid (Boullay), by action of boron fluoride (Desfosses, Ann. Chim Phys. [2] 36, 294), of ZnCl<sub>2</sub> (Masson, Annalen, 31, 63), of SnCl<sub>4</sub>, and other metallic chlorides (Kuhlmann, *ibid* 33, 97, 192) From alcohol by heating with dry NH<sub>4</sub>Cl to  $260^{\circ}$ –400° (Berthelot, Annalen, 83, 110) From C<sub>2</sub>H<sub>5</sub>NaO and C<sub>2</sub>H<sub>5</sub>I (Williamson) From alcohol and HCl, HBr, or HI at 200° or 240°, or by heating alcohol with ZnCl<sub>2</sub>, SnCl<sub>4</sub>, HgCl<sub>2</sub>, CaCl<sub>2</sub>, MgSO<sub>4</sub>, or chrome alum to 300° (Reynoso, Ann Chim Phys [3] 48, 385) Also from C<sub>2</sub>H<sub>5</sub>I and Na<sub>2</sub>O (Green, Bull. Soc. chim [1,] 29, 458)

The continuous method now adopted is due to Boullay (J. Pharm Chim. 1, 97), and the true explanation of the action was first given by Williamson (l.c.). The alcohol and sulphuric acid

first form  $\rm C_2H_5HSO_4$  and  $\rm H_2O$ , and then, by the action of  $\rm C_2H_5HSO_4$  upon more alcohol,  $\rm (C_2H_5)_2O$  and  $\rm H_2SO_4$  are formed. That this is true Williamson proved by using amyl sulphunic acid and ethyl alcohol, when amyl-ethyl ether was produced (cf. Prunier, J. Pharm. Chim. 1897, 5, 513). Addition of  $\rm Al_2(SO_4)_3$  increases rate of production of ether owing to catalytic action (Senderens, Compt. rend. 1910, 151, 392).

Preparation -Boullay's method is usually A mixture of 9 parts H<sub>2</sub>SO<sub>4</sub> with 5 employed parts of alcohol (90 pc) is placed in a retort, and alcohol is slowly dropped in and the temperature is maintained between 130° and 140°. An iron retort lined with lead is usually employed, and the ordinary mercurial thermometer, which is, of course, liable to fracture, is replaced by a bulbed tube containing some non-drying oil. The tube is then either exhausted by the airpump, and hermetically scaled, or its open end is closed by a plug of cotton wool, and two marks are made on the tube corresponding to positions of the surface of the oil at 130° and 140° respec-In order to avoid risk of explosion by contact of the vapour with direct flame, coils of lead tubing, conveying superheated steam or the vapour of some liquid of high boiling-point (e g coal-tar phenol), are used as a source of heat The distillate is treated with milk of lime or solution of soda in order to remove the SO, which is always simultaneously formed. It is then rectified in an apparatus devised by Mohr, which consists of a still from which the vapour is led through a vessel kept at 38° by warm water Here the alcohol and water vapours are condensed, and the ether passes over, and is condensed in an efficient condenser

By means of an apparatus devised by Soubeiran, ether can be prepared in a fairly pure state by one operation. A mixture of about 30 lbs of H<sub>2</sub>SO<sub>4</sub> with 20 lbs of alcohol of 85 p c is heated rapidly to 130° in a copper still, to the bottom of which two tubes convey alcohol from a tinned copper reservoir. The neck of the retort (which is of lead) is connected with the first condenser, which is a copper vessel kept warm by allowing the waste water from the worm tub (where the ether is condensed) to flow over the outside of it Here the steam and vapour of alcohol are condensed, and the liquid formed can be drawn off from below by means of a stopcock. The ether vapour then passes on to the purifier, which consists of a vessel filled with wood charcoal saturated with a solution of caustic soda, which not only removes the SO<sub>2</sub>, but also absorbs the oil of wine The pure ether passes on, and is condensed in a well-cooled worm greatest yield of other is obtained at 140°-145°; below this temperature the yield is less, and above this much SO, is evolved (Prescott and Norton, Amer Chem J 6, 243) Other apparatus have been described by Annaratone (Fr Pat 408089; J Soc Chem 1910, 592), Fritzsche (Zeitsch anal Chem 1897, 36, 298), Barbet et fils et Cie (Fr. Pat 479435, 1914). Sulphune and has been largely replaced by benzenesulphonic acid in this preparation (Krafft, Ber 26, 2831; D. R 69115, Frdl. m. 11).

Commercial ether contains water, alcohol, aldehyde, methyl compounds, ketones, and vinyl Llcohol. A. J. Jones (Pharm. J. 1919, 103, 71)

found that in commercial samples, methyl compounds range from 0.02 to 0.06 p.c., sp.gr. 0.7182-0.7211, and ketones (as acctone) from 0.09 to 5.0 pts per 10,000 Vinyl alcohol may be removed by shaking with water or treatment with bromine, phenylhydrazine, or potassium hydroxide (Poleck and Thummel, Ber. 22, 2863), and the other impurities by treatment with dry Fe(OH)<sub>2</sub>, prepared from crystallised FeSO<sub>4</sub>, and an equimolecular weight of powdered lime (Gaborini, Bull Assoc Chim Sucr Dist 1909, 26, 1165); by treatment with potassium hydroxide and powdered potassium permanganate, and finally drying with calcium chloride (Fierz; Wade and Finnemore, Chem. Soc Trans 1909, 1842, François, Chem Zent 1897, n 144; Fritsch, Chem Zeit 1909, 33, 759) Phosphorus pentoxide may be substituted for calcium chloride (Lieben, Annalen, Suppl 7, 218) (For other methods, v Guigues, J. Pharm. Chim 1906, 24, 204, and Ekenberg, Zeitsch anal Chom 36, 245)

According to the BP, 1914, ordinary ether should have a sp gr of 0.726, 15°/15°, and boil between 34° and 36°. Purified ether (Æether purificatus of the BP) should give no yellow colouration with solid potassium hydroxide in the dark, or with a solution of freshly prepared solution of potassium iodide To test for presence of methyl compounds shake 2 vols of the ether with I vol of 20 pc alcohol and I vol. of water, allow the mixture to separate, and draw off the lower layer Mix 5 millilitres of this lower layer with 2.5 millimetres of an aqueous solution (1 in 50) of potassium permanganate and 0.2 millilitre of sulphuric acid At the end of 3 minutes add 0.5 millilitre of an aqueous solution (96 in 100) of oxalic acid, followed by I millilitre of sulphuric acid, and then by 5 millilitres of decolourised solution of fuchsin, and mix thoroughly. No violet colour should be produced within 20 minutes

Wade and Finnemore (lc) find that ether and water form a binary mixture, b p 34 15°, containing 1 3 p c water, and that slight traces of alcohol greatly increase the solubility of water in other 1 gram ether dissolves 0 023 gram water at 15°-20° and 0 019 at 10° (cf. Schunke, Zeitsch physikal Chem 14, 331; Napier, Bull Soc chim. 29 [11] 129); 1 gram water absorbs at 0°, 0 12465, at 10°, 0 09599, at 20°, 0 07487, at 30°, 0 06370 gram ether (Schunke, lc; Tolloczko, Ber 28, 808; Klobbie, Zeitsch physikal Chem 24, 615; Osaka, J. Soc Chem Ind 1909, 625). If ether be shaken with an equal volume of CS<sub>2</sub>, a turbidity is produced if water be present. A better test is with anhydrous copper sulphate or amalgamated alumınıum.

Decompositions—Pt black, CrO<sub>3</sub> or HNO<sub>3</sub> oxidise ether to acetic acid; ozone produces ethyl peroxide Ether vapour and air at 100°, in the presence of a copper catalyser and platinum-black, yield acetaldehyde and formaldehyde (Orlow, J. Russ. Phys. Chem. Soc. 1908, 40, 799; cf. Legler, Annalen, 217, 392). When ozonised oxygen is passed into ether, a solution is obtained containing  $C_2H_2O_4$ ,  $C_2H_4O_2$ , and  $H_2O_2$ , also a small quantity of  $CH_2O_2$  (A. W. Wright). Ether combines with  $H_2SO_4$  to form  $C_2H_5LSO_4$ ; by action of SO<sub>3</sub>, we first get (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>SO<sub>4</sub>, but by excess of SO<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>HS<sub>2</sub>O<sub>7</sub> and a small quantity

of, a body of the composition  $CH_2(SO_3C_2H_5)_2$ are formed. When ether is heated with water and a small quantity of H<sub>2</sub>SO<sub>4</sub> to 150°-180°, alcohol is formed (Erlenmeyer, Zeitsch. Chem. 1868, 343) According to Lieben (Annalen, 165, 136), the reaction goes on, though very slowly, without H2SO4, and at the ordinary temperature Ether combines easily with HI and yields CoH, I (Silva, Ber 8, 903), and is decomposed by heating with zinc-dust into C2H4 and water (Jahn, Monatsh 1, 675), bromine and aqueous ether yield bromacetaldehyde (Maugin, Compt rend 1908, 147, 747).

Instances of the spontaneous explosion of ether on heating are not unknown, due probably to the presence of ethyl peroxide produced by autoxidation Ether occasionally contains hydrogen peroxide and acctaldelyde and a relatively large amount of vinyl alcohol (Kassner, Arch. Pharm 1912, 250, 436). Hydrogen peroxide may be detected by the immediate red colouration produced when 5 c.c is treated with 1 c c of N/10 potassium thiocyanate solution and 2 drops of a 5 pc freshly prepared acidified solution of ferrous ammonium sulphe to (Dietze, Apoth Zeit 1915, 30, 165) There should be no liberation of iodine when 2 cc. of a 10 pc solution of cadmium potassium iodide are shaken with 10 cc of the sample. Per-oxides or peroxidised compounds may be formed in ether when improperly stored. The degree of oxidation is dependent upon the nature of the container, the purity of the ether, the amount of air present, and, in the case of glass containers, the intensity of the light which accelerates the oxidation Peroxides may be removed by means of ferrous hydroxide or by shaking with mercury

Mixtures of ether vapour and air explode when heated to about 190° if the amount of ether in the gaseous mixture reaches 1 gram

A hydrate  $(C_2H_5)_2O$ ,  $2H_2O$  is known; it is a solid body, which only exists below  $-3.5^{\circ}$ , and is obtained by the rapid evaporation of aqueous obtained by the rapid evaporation of aqueous ether on filter paper (Tanret, Bull Soc chim 30, 505; Compt rend 87, 765) Dry ether and chlorine at  $-80^{\circ}$  yield a white crystalline chloride  $(C_2H_5)_2OCl_2$ , mp  $-51^{\circ}$ ; bromine similarly yields a reddish-yellow bromide  $(C_2H_5)_2OBr_2$ , m.p.  $-40^{\circ}$  (McIntosh, Chem. Soc. Trans 1905, 789) A bromide  $(C_2H_5)_2OBr_3$ , ied plates, mp 22°, is also known (Schutzenberger, Annalen, 167, 86).

berger, Annalen, 167, 86).

Ether combines with various metallic salts and forms crystalline bodies, eg BeCl<sub>2</sub>, 2(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O, large prisms (Atterberg, Ber 9, 856); HgBr<sub>2</sub>,3(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O (Nicklès, J. 1861, 200); AlBr<sub>3</sub>,(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O (N.). TiCl<sub>4</sub>,(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O; crystalline, mp 42°-45°; bp 118°-120°, decomposes into TiCl<sub>3</sub>(Oc<sub>2</sub>H<sub>5</sub>) (Bedson, Annalen, 180, 236; Chem Soc. Trans 1876, 1, 311)

Procentees—A colourless mobile bound with

Properties —A colourless mobile liquid with asant smell Volatilises very quickly, and pleasant smell by so doing produces great cold "It is extremely inflammable, and its vapour, which is very heavy, forms an explosive mixture with air Ether is readily soluble in concentrated HCl solution (Draper, Chem. News, 35, 87), and in concentrated H<sub>2</sub>SO<sub>4</sub>, from which it is separated by addition of ice (Riedel, D R P. 52982, Frdl n 551) It is miscible with almost all

organic liquids, and with liquid  ${\rm CO}_2$ . It is a good solvent for resnns, fats, alkaloids, &c., and sulphur and phosphorus dissolve in it to a small extent. It also dissolves I and many salts, e q Fe<sub>2</sub>Cl<sub>6</sub>, HgCl<sub>2</sub>, PtCl<sub>4</sub>, and many gases, not-

ably ammonia

In laboratories ether should always be kept over mercury or sodium in completely filled, well-stoppered bottles, preferably made of blue, green, or brown glass, in cool places excluded from the light (cf Baskerville and Hamor, J Ind Eng Chem 1911, 3, 301 et seq; J Soc Chem Ind 1911, 830)

Its vapour when inhaled produces insensibility (Faraday, 1818), and it is used as an anæsthetic in surgery It is also occasionally used in surgery for producing local anæsthesia, this is done by directing a spray of ether upon the skin, when the extreme cold produced by evaporation produces complete insensibility.

It is also used as a solvent in the preparation of collodion, in the extraction of tannic acids, of

several alkaloids, &c

Chlorine acts violently upon ether, setting fire to it, and liberating carbon (Cruikshank) ether be kept cool by icc, and in the dark, substitution, roducts are obtained. The Clreplaces the H atom by atom from one ethyl group first, CHHHCHHOC<sub>2</sub>H<sub>5</sub> in the following order.

(Lieben, Annalen, 111, 121; 146, 180; Abeljanz, 164, 197, Jacobsen, Ber 4, 215, also Liebig, Annalen, 1, 220; Malaguti, Ann Chim Phys [2] 70, 338; [3] 16, 5; Fritsch and Schumacher, Annalen, 279, 302)

# Monochlorether, Monochlorethyl oxide

CH, CHCl O C, H,;

bp 97°-98°, by the chlorination of ether (Lieben); by saturating a mixture of aldehyde and alcohol with HCl (Frapolli, Wurtz, Annalen, 108, 226); by the action of PCl<sub>5</sub> upon acetal (Bachmann, ibid 218, 39) It is decomposed by H<sub>2</sub>SO<sub>4</sub>into HCl<sub>7</sub>C<sub>2</sub>H<sub>4</sub>O, and C<sub>2</sub>H<sub>5</sub> OH C<sub>2</sub>H<sub>5</sub>ONa forms NaCl and acetal; water decomposes it into HCl. aldehyde, and alcohol (Laatsch, Annalen, 218, 36)

#### Dichlorether, Dichlorethyl oxide

#### CH,Cl CHCl O C,H,;

b p 140°-145°, sp gr 1 174 at 23°; by action of Clupon ether (Malaguti, Annalen, 32, 15; Wildman and Gray, J Amer Chem Soc 1919, 41, 1122); by action of HCl upon a mixture of chloraldehyde and C<sub>2</sub>H<sub>5</sub> OH (Natterer, Monatsh 5,496), by action of HCl upon chloracetal (N)

Dichlorether is decomposed on heating in a tube into  $C_2H_5Cl$  and other products  $C_2H_5ONa$  forms first *chloracetal*  $CH_2Cl$   $CH(OC_2H_5)_2$ . By heating with excess of C2H5ONa in sealed tubes, ethyl glycol acetal CH<sub>2</sub>(OC<sub>2</sub>H<sub>5</sub>)CH(OC<sub>2</sub>H<sub>5</sub>)<sub>2</sub> is formed In ethereal solution, Zn(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> yields ethyl chlorether CH<sub>2</sub>Cl CH(C<sub>2</sub>H<sub>4</sub>)OC<sub>2</sub>H<sub>5</sub>; b p 141°, sp gr 0 9735 at 0° With excess of 141°, sp gr 0.9735 at 0° With excess of Zn(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> we get CH<sub>2</sub>(C<sub>2</sub>H<sub>5</sub>)CH(C<sub>2</sub>H<sub>5</sub>)OC<sub>2</sub>H<sub>5</sub>; bp 131 4° at 749 6 mm; sp gr 0.7865 at 0° (Lieben, Annalen, 178, 14) Strong caustic potash forms alcohol, a body C<sub>8</sub>H<sub>16</sub>Cl<sub>2</sub>O<sub>3</sub>, a little glycollic acid, and two isomeric bodies of CHCCO (Abdustry 31, 164) the composition C<sub>4</sub>H<sub>9</sub>ClO<sub>2</sub> (Abeljanz, ibid. 164, 218) · α-C<sub>4</sub>H<sub>9</sub>ClO<sub>2</sub>, chloraldehyde alcoholate CH<sub>2</sub>Cl CH(OH)O·C<sub>2</sub>H<sub>5</sub>, bp 93°-95°; and β-C<sub>4</sub>H<sub>4</sub>ClO<sub>-</sub> oxychlorater CH<sub>2</sub>OH·CHCl·O·C<sub>2</sub>H<sub>5</sub>,

also formed by chlorination of ether (Jacobsen, Ber 4,217), bp 151°-155°. With NH<sub>2</sub> OH yields glyoxime (Hantzsch and Wild, Annalen, 289, 293). It is decomposed by water, and in presence of marble yields monochloracetaldehyde alcoholate, which, on distillation, gives monochloraldehyde and monochloracetal (Fritsch

Trichlorether CHCl<sub>2</sub> CHCl O C<sub>2</sub>H<sub>5</sub>; b p 167°-168°, 155°-158° (Copaux), 170°-175° (Oddo and Mameli); sp gr. 1 3303 at 14° (O and M). By the action of PCl<sub>5</sub> upon dichloracetal CHCl<sub>2</sub> CH(OC<sub>2</sub>H<sub>5</sub>)<sub>2</sub> (Krey, J 1876, 475), by hydrolysis of the compound formed by action of chlorine on triethyl boric ester (Copaux, Compt. rend 127, 721), by chlorinating ab-dichloroethyl ether with I molecule of Cl at temperature of boiling saturated salt solution At 230°-240° in a scaled tube, yields dichloroaldehyde and  $\rm C_2H_5Cl$  HCl at 110° for 6 hours, and distillation with concentrated H<sub>2</sub>SO<sub>4</sub>, yield the same products Conversely, dichloroaldehyde in alcohol with HCl yields trichloroether Boiled with water, trichloroether yields acctaldehyde, di-chloroacetaldehyde hydrate (m p 55°-56°, bp 965°-975°), dichloroacetal, and other very complex compounds (Oddo and Mameli, Gazz chim ital 33, ii 373; also ibid 36, 1 480), decomposed by heating with C.H.ONa, yielding dichloracetal

Tetrachlorether CCl<sub>2</sub> ('HCl O C<sub>2</sub>H<sub>5</sub>; b p. 189 4° at 749 1 mm, 79° at 16 mm, 189 7° at 758 7 mm.; sp gr 1 4225 at  $18^{\circ}/4^{\circ}$  (Weber and Foster, J Amer Chem. Soc 31, 1410), sp gr 1 4379 at 0°, 1.4182 at 15 2°. Formed by action of Cl upon monochlorether in contact with I (Wurtz, Vogt, Zeitsch Chem. 1871, 689), or of chloral alcoholate CCl<sub>3</sub> CH(OH)O C<sub>2</sub>H<sub>5</sub> upon PCl<sub>5</sub> (Henry, Ber 5, 101, 435, Paternò, Pisati, J 1872, 303) From trichlorvinyl ethyl ether and chlorine (Godefroy) A liquid with a camphorlike smell, decomposed by heating with water into C<sub>2</sub>H<sub>5</sub>OH, HCl, and chloral Alcoholic KOH forms with it trichlorunal ethal ether CCl<sub>2</sub> CCl O C<sub>2</sub>H<sub>5</sub> Heated with alcohol, forms trichloracetal (Pergami).

Pentachlorether. Two isomerides:

(1) CCl, CCl, O C<sub>2</sub>H<sub>5</sub>, bp 190°-210° with decomposition; spgr 1 645 (Jacobsen, Ber 4, 217); by action of Cl upon trichlorvinyl ethyl ether (r supra) (Busch, Ber 11, 445)

(2) CCl<sub>1</sub> CHCl O CH<sub>2</sub> CH<sub>2</sub>Cl; b p 235°; sp gr 1 577 at 80° By action of PCl<sub>5</sub> on a compound of chloral and glycol chlorohydrin

#### CCl<sub>3</sub> CH(OH)O CH<sub>2</sub> CH<sub>2</sub>Cl

(Henry, Ber 7, 763)

Hexachlorether  $CHCl_2$  CHCl O CHCl  $CHCl_2$ ; b p  $250^{\circ}$ , by action of PCl<sub>5</sub> upon dichloral dehyde saturated with HCl (Paterno, Pisati, J 1871, 508)

Octochlorether C4H2Cl8O, from ethylidene oxychloride and Cl in sunlight (Roth, Ber 8, Camphor-like smelling, volatile crystals

Perchlorether (C<sub>2</sub>Cl<sub>5</sub>)<sub>2</sub>O (Regnault, Annalen, 34, 27); sp gr 1 900 at 14 5° (Malaguti, Ann Chim Phys [3] 16, 4); the ultimate product of the action of CI upon ether in sunlight, is a solid, forming orthorhombic scales, m p. 69°, which possesses a smell like that of camphor Decomposed by heating at 300° into C2Cl6 and CCl3 COCl; with K2S yields perchlorvinyl ether

(C<sub>2</sub>Cl<sub>3</sub>)<sub>2</sub>O, with liberation of S (Malaguti, Ann. Chim Phys [3] 16, 19) This body is a liquid, b p 210°, sp gr 1 652 at 21°, which combines with Cl in sunlight to form (C<sub>2</sub>Cl<sub>2</sub>)<sub>2</sub>O, or with Br

vielding (C2Cl3Br2)2O

Ethyl peroxide (C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>O<sub>3</sub>, produced by leading dry ozone into absolute ether (Berthelot, Bull Soc chim ii 36, 72), is a thick syrup which does not solidify at - 40°, is partially volatile on heating, but easily explodes, is miscible with water, by which it is decomposed into C2H5OH and H<sub>2</sub>O<sub>2</sub>

Ethyl phosphates.

1 . Normal ethyl phosphate (C2H5)3PO4, b p  $215^{\circ}$  , sp gr 1 072 at 12°. By action of heat on lead diethyl phosphate Pb(C<sub>2</sub>H<sub>2</sub>)<sub>4</sub>(PO<sub>4</sub>)<sub>2</sub> (Vogeli, Annalen, 69, 190), from C<sub>2</sub>H<sub>3</sub>I and Ag<sub>3</sub>PO<sub>4</sub> (Clermont, Ann Chum Phys [3] 42, 330), from PO(I<sub>3</sub> and C<sub>2</sub>H<sub>2</sub>ONa (Limpitcht, Annalen, 124, 213), from PO(I<sub>3</sub> and C<sub>4</sub>H<sub>2</sub>ONa (Limpitcht, Annalen, 124, 213), from PO(I 134, 317), from  $\stackrel{\circ}{1}{1}{20}_{5}$  and absolute elcohol (Carius, ibid 137, 121) Colourless liquid, miscible with and slowly decomposed by water, forming diethyl phosphoric acid

2 Diethyl phosphoric acid (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>HPO<sub>4</sub>, a syrupy liquid obtained by action of P<sub>2</sub>O<sub>5</sub> upon It is a monobasic acid, and forms wellmarked salts, eg  $\text{Ca}[(\text{C}_2\text{H}_5)_2\text{PO}_4]_2$ , crystalline needles . Pb[(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>PO<sub>4</sub>]<sub>2</sub>, needles , m p 180°,

soluble in water

3 Ethyl phosphoric acid H<sub>2</sub>(C<sub>2</sub>H<sub>5</sub>)PO<sub>4</sub>, an oily liquid decomposed by heating, evolving (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O, C<sub>2</sub>H<sub>5</sub>OH, and C<sub>2</sub>H<sub>4</sub> Equal parts of alcohol (95 p c) and syrupy H<sub>3</sub>PO<sub>4</sub> are heated to 60° or 80°, diluted with water and neutralised with BaCO3, filtered, the Ba salt allowed to crystallise, and then decomposed by H2SO4 (Pelouze, Ann Chim Phys [2] 52, 37)

Also produced by action of P2O5 upon ether or alcohol or of syrupy  $\rm H_3PO_4$  upon ether (Vogeli, J 1847-48, 694) Forms crystalline salts with the metals, of these the lead salt is least soluble

Ethyl phosphites.

Triethyl phosphite (C2H5)3PO3; bp 191° (or 188° in stream of H); sp gr 1 075 at 15 5° (Railton, Chem Soc Trans 1855, 7, 216), from

PCl<sub>3</sub> and alcohol or sodium ethoxide

An ethereal-smelling liquid, which absorbs O and is converted by PCl<sub>5</sub> into PO(OC<sub>2</sub>H<sub>5</sub>)Cl<sub>3</sub>, PCl<sub>3</sub>, and C<sub>2</sub>H<sub>5</sub>Cl (Geuther, Hertz J 1876, 207) By heating with NaOC<sub>2</sub>H<sub>5</sub> yields Na(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>PO<sub>3</sub>; with (C<sub>2</sub>H<sub>5</sub>O)<sub>2</sub>O, gives

H<sub>2</sub>(C<sub>2</sub>H<sub>5</sub>)PO<sub>3</sub>

(Jaehne, Annalen, 256, 269).

2 Duthyl-phosphorous acid H(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>PO<sub>3</sub>; b p 184°-185°, sp gr 1 0749 (15 5°/4°) From P<sub>4</sub>O<sub>6</sub> and C<sub>2</sub>H<sub>3</sub>OH (Thorpe and North, Chem Soc Trans 1890, 634); from PCl<sub>3</sub> and C<sub>2</sub>H<sub>5</sub>OH (Levitsky, J Russ Phys Chem Soc 1903, 35, 211, Milobendzi and Sachnowski, Chem Soc Abst 1918, 1 477), from  ${\rm C_2H}_5{\rm I}$  and  ${\rm Pb_3(PO_3)_2}$  (Michaelis and Becker, Ber 1897, 1005). Colourless mobile liquid with an alliaceous odour

3 Ethyl-phosphorous acid  $H_2(C_2H_5)PO_3$ , very unstable syrupy liquid; obtained by action of PCl<sub>3</sub> upon weak alcohol (Wurtz, Ann Chim Phys [3] 16, 218). The salts do not crystallise well, but are more stable than the acid

Chlorule of ethyl-phosphorous acid C<sub>2</sub>H<sub>5</sub>POCl.; b p 117 5° (corr); sp gr 1·30526 at 0°/4° (Thorpe, Chem Soc. Trans. 37, 346); from PCl<sub>3</sub>

and C<sub>2</sub>H<sub>5</sub>·OH (Menschutkin, Annalen, 139, 343); decomposed by water into HCl, C<sub>2</sub>H<sub>5</sub> OH, and H<sub>3</sub>PO<sub>3</sub>; also by heating to 165° into C<sub>2</sub>H<sub>5</sub>Cl, P, PCl<sub>3</sub>, and P<sub>2</sub>O<sub>5</sub> (Chambon, J 1876, 205)

Diethyl pyro-phosphorous acid is not known, but the zinc salt  $(C_2H_5)_2P_2O_5Zn$  is not known, but the zinc salt  $(C_2H_5)_2P_2O_5Zn$  is produced when  $Zn(C_2H_5)_2$  is heated with  $P_2O_5$  to 140° (Dilling, Zeitsch Chem 1876, 266)

Ethyl selenic acid C<sub>2</sub>H<sub>5</sub>HSeO<sub>4</sub>, very unstable, produced from H<sub>2</sub>SeO<sub>4</sub> and C<sub>2</sub>H<sub>5</sub>OH (Fabian, Annalen, Suppl 1, 244). The salts are also unstable, the most permanent being the strontium salt, they can be crystallised with the ethyl-sulphuric acid salts

Ethyl selenhydrate, Ethyl selenomercaptan  ${
m C_2H_5SeH}$  , b p 53 5°; sp gr 24°/4° 1 3954 (Tschugaeff, Ber 1909, 42, 49) From ethyl iodide and sodium hydroselenide Nauscous-

smelling liquid

Ethyl selenide, Selenethyl  $(C_2H_5)_2Se$ : b p 108°, discovered by Lowig (Pogg Ann 37, 552) By action of  $K_2Se$  upon  $(C_2H_5)_2C_2O_4$ (L), by action of K('2H5SO4 upon K2Se (Joy, Annalen, 86, 35; Rathke, ibid 152, 210), crude product is then distilled again with KOH, C<sub>2</sub>H<sub>5</sub>HSO<sub>4</sub>, and a small piece of P or S (Pieverling, ibid 185, 331, Ber 9, 1469) A colourless mobile liquid with a not unpleasant smell, is insoluble in water, acts as a base . H(I precipitates from a solution in dilute  ${\rm HNO_3}$  an oily liquid (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>SeCl<sub>2</sub>, which gives with aqueous  $NH_3$  a crystalline oxychloride  $[(C_2H_5)_2Se]_2OCl_2$ . Combines with  $C_2H_5I$  to form crystalline  $(C_2H_5)_3Sel$  (Pieverling) This is a white substance resembling Epsom salts in appearance. The (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>SeCl<sub>2</sub> forms double salts with several metallic chlorides, e g

## [(C,H5)aSeCl],ZnCl2

Ethyl diselenide  $(C_2H_5)_2Se_2$ ; b p  $186^\circ$ ; formed as a by-product in the preparation of (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>Se, was first obtained by Wohler and Dean (Annalen, 97, 1) from KC<sub>2</sub>H<sub>5</sub>SO<sub>4</sub> and K<sub>2</sub>Se<sub>2</sub> (Rathke, *ibid* 152, 212), has an intensely disagreeable smell, and acts as a poison (Pieverling) A solution in dilute HNO<sub>3</sub> gives with HCl monoclinic crystals of  $(C_2H_5)_2Sc_2O_2(OH)_2Cl_2$ , which are soluble in water, and are reduced by SO, to ethyl diselenide

Ethoxy-selenyl chloride C<sub>2</sub>H<sub>5</sub>O SeOCl, m p. 10°; bp 175°, not without decomposition. From SeOCl<sub>2</sub> and absolute alcohol Heated in a tube to 200° yields Se, C<sub>2</sub>H<sub>5</sub>Cl, HCl, and H<sub>2</sub>O (Michaelis and Landmann, Annalen, 241,

156Diethylselenite SeO(OEt)<sub>2</sub>; bp 183°-185°, not without decomposition, sp gr 149 at 165°. From ScoCl<sub>2</sub> and NaOEt in dry ether at 180°-190°, also from Ag<sub>2</sub>SeO<sub>3</sub>, and C<sub>2</sub>H<sub>5</sub>I at 85° m sealed tube A thick liquid, decomposed by water into C<sub>2</sub>H<sub>5</sub>OH and H<sub>2</sub>SeO<sub>3</sub> Decomposed by heating in a tube at 200° with separation of Se (M and L.)

Ammonium monoethylselenite

#### C<sub>2</sub>H<sub>5</sub>O SeO ONH<sub>4</sub>;

a crystalline, very hygroscopic acid, formed by action of alcohol and ammonia upon SeO2 at temperature of melting ice. It is soluble in alcohol, insoluble in ether (Divers and Hada, Chem Soc. Trans. 75, 538).

Ethyl sulphates.

Mono-ethyl sulphate  $C_2H_5HSO_4$  (Ethyl-sulphu) ic acid; Sulphethylic acid; Sulphovinic

acid); sp gr 1 316 at 16°

First noticed by Dabit (Crell's Ann [1802] 1, 394); and again by Serturner (Gilb Ann 60, 53; 64, 67, [1818-1820]; see also Heeren, Pogg. Ann. 7, 193; Hennel, Trans. Roy Soc 1826; 240, 1828, 365, Serullas, Ann Chim Phys [2] 39, 153; Liebig and Wohler, Pogg Ann 22, 487, Liebig, Annalen, 9, 17, 32, 13, 32, 38; Dumas and Boullay, Ann Chim Phys [2] 36, 300)

Formed by action of sulphuric acid upon alcohol (Dabit, Serturner, Serullas), also by absorption of ethylene in sulphuric acid (Hennel, Berthelot, Ann Chim, Phys [3] 43, 385), by heating ether with strong H<sub>2</sub>SO<sub>4</sub> (Hennel, Magnus, Pogg Ann 27, 386), from S<sub>2</sub>Cl<sub>2</sub> and alcohol (Heusser, Annalen, 151, 249), from HO SO, Cl and alcohol in the cold (on heating, this gives C<sub>2</sub>H<sub>5</sub>O SO<sub>2</sub>Cl, Claesson, J pr Chem.

[2] 19, 245)

Preparation -A mixture of equal parts of absolute alcohol and strong sulphure acid is heated for 4 hours on the water-bath, by this treatment 59 p.c. of the H2SO4 used is converted into mono-ethyl sulphate (Berthelot, Bull Soc chim ii 19, 295) By using 3 molecules alcohol and 1 molecule H<sub>2</sub>SO<sub>4</sub> (ie 3 to 2 by weight), 77 p c of the sulphuric acid is utilised (Claesson, J pr Chem [2] 19, 246) The mixture is diluted with water and neutralised with lead or barium carbonate, filtered, evaporated to crystallisation, the crystals dissolved in water, and the acid set free by means of H<sub>2</sub>SO<sub>4</sub>, or, when lead carbonate is used, by means of H<sub>2</sub>S. The piecipitated BaSO4, PbSO4, or PbS is removed by filtration, and the liquid evaporated in vacuo over sulphuric acid Mono-ethyl sulphate is a syrupy liquid, very soluble in water, soluble also in alcohol, but insoluble in ether

Heated with water, yields alcohol and sulphuric acid; on heating with alcohol to 140°, ether and H<sub>2</sub>SO<sub>4</sub> are formed Oxidising agents, e q MnO<sub>2</sub> or K<sub>2</sub>CrO<sub>4</sub>, give aldehyde (Jacquemin and Lies-Bodard, J. 1857, 345). The salts are very soluble in water, and can be heated in dilute solutions without decomposition, though in strong solution, or by prolonged heating, they are decomposed with formation of alcohol, a sulphate, and free H<sub>2</sub>SO<sub>4</sub> This decomposition may be diminished by the addition of an alkali hydroxide or carbonate By dry distillation, the salts yield 'heavy oil of wine,' a liquid containing diethyl sulphate and olefines (c) Kremann, Monatsh. 1917, 38, 53) HCl gas yields C<sub>2</sub>H<sub>5</sub>Cl and a metallic sulphate (Kohler, Ber 19, 11,

1929)

By distilling sodium ethyl sulphate in a vacuum, Lilienfeld (Austrian Pat 63526) obtains

Vacuum, Imenied (Austrian Fat 65526) obtains ethyl sulphate in 83-90 p c yield.

Salts (v. Marchand, Pogg Ann 32, 456, 41, 595). LiC<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>, H<sub>2</sub>O; NaC<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>, H<sub>2</sub>O, KC<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>, crystallises in anhydrous monoclinic plates, soluble in 0.8 part H<sub>2</sub>O at 17°, (Schabus, J. 1854, 560; Marignac, J. 1855, 608)

 $Mg(C_2H_5SO_4)_2,4H_2O$ ,  $Ca(C_2H_5SO_4)_2,2H$ ,O

forms monoclinic laminæ, soluble in 0.8 part water at 17° (M). Ba( $\rm C_2H_5SO_4$ )<sub>2</sub>,2H<sub>2</sub>O, monoclinic plates, sp gr. 2.080 at 21.7°, soluble in 0.32

parts of water at  $17^{\circ}$  (M.); Pb(C<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>)<sub>2</sub>,2H<sub>2</sub>O; Harge rhombic tables —  $Pb(C_2H_5SO_4)_2$ ,  $PbO_5$ amorphous; soluble in 0 54 part H,O at 17°

amorphous; soluble in 0.54 part  $H_2O$  at 17° Diethyl sulphate  $(C_2H_5)_2SO_4$ , sp gr. 1.1837 at 19° (Claesson, J. pr. Chem. [2] 19, 257), b.p. 208° (with decomposition), 113.5° at 31 mm, 118° at 40 mm, 120.5° at 45 mm. (Villers, Bull. Soc. chim. ii. 34, 26); m.p. about  $-24.5^\circ$  (V), occurs in 'heavy oil of wine' Formed by leading  $SO_3$  vapour into absolute ether (Wetherill. Annalen. 66, 117), ii. small ether (Wetherill, Annalen, 66, 117). in small quantity by distillation of C2H5HSO4; by dropping alcohol into chloride of ethyl sulphuric acid C<sub>2</sub>H<sub>5</sub>O SO<sub>2</sub>Cl, or by action of H<sub>2</sub>SO<sub>4</sub> upon alcohol (Claesson), from Ag<sub>2</sub>SO<sub>4</sub> and C<sub>2</sub>H<sub>5</sub>I (Stempnewsky, Ber 11, 514) A nuxture of 1 vol absolute alcohol and 2 vols H<sub>2</sub>SO<sub>4</sub> ıs dıstılled slowly *in racuó* (Vilhers) Absolute alcohol is mixed with Nordhausen sulphuric acid, the ether separated by means of CHCl, and the product distilled in vacuo (Claesson and Lundvall, Ber 13, 1699)

Is an oily liquid, with an odour like that of peppermint, insoluble in water, and slowly decomposed by that liquid even in the cold, when heated with water, yields alcohol,  $\rm H_2SO_4$  and  $\rm C_2H_5HSO_4$  Heated with alcohol, forms ether  $\mathrm{C_2H_5HSO_4}$ and C<sub>1</sub>H<sub>5</sub>HSO<sub>4</sub>, warm baryta water forms alcohol and Ba(C<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>)<sub>2</sub> (Vilhers)

Unsaturated hydrocarbons are present to

the extent of about 5 pc in 'heavy oil of

wine

Chloride of ethyl sulphuric acid (1, H<sub>3</sub>O SO, C1; b p  $151^{\circ}$ - $154^{\circ}$  (corr.) (Claesson, J. pr. Chem. [2] 19, 248), with slight decomposition.  $93^{\circ}$ - $95^{\circ}$  at 100 mm (Muller, Ber 6, 227) From C<sub>2</sub>H<sub>5</sub>Cl and SO<sub>3</sub>, together with isomerides (v Ethyl chloride), by dropping alcohol into SO<sub>2</sub>Cl<sub>2</sub> (not ruce versa, diethyl sulphate is then formed) (Behrend, J. pr. Chem. [2] 15, 28), from a salt of ethyl sulphure acid and PCl<sub>5</sub> (Purgold, Ber 6, 505), from ethyl chloroformate Cl CO<sub>2</sub>C<sub>2</sub>H<sub>5</sub> and fuming sulphuric acid (Wilm, Ber 6, 505), by passing C<sub>2</sub>H<sub>4</sub> into HSO<sub>3</sub>Cl, SO<sub>4</sub>H C<sub>2</sub>H<sub>4</sub> SO<sub>2</sub>Cl is also produced (Muller, *ibid*. 6, 227)

A liquid with an irritating smell, exciting tears, and decomposed by contact with water into HCl, H2SO4 and alcohol When the chloride is added to methyl alcohol or ethyl alcohol, ethyl sulphuric acid and CH3Cl or C2HaCl are produced, with amyl alcohol, amyl sulphuric acid and ethyl chloride are formed If, on the other hand, absolute al-(Muller) cohol is dropped into the chloride, diethyl sulphate is produced (Claesson)

Bromethyl sulphuric acid

 $CH_2Br\ CH_2\ OSO_2OH(?)$ ;

from ethylene bromide and SO<sub>3</sub> (Wroblewski, Zeitsch Chem 1868, 563) The barium salt is anhydrous, and is decomposed by continued heating with water, with separation of BaSO4 and formation of glycol (Beilstein and Wiegand, Ber 15, 1370) An isomeric body

CH<sub>3</sub> CHBr OSO, OH(?)

is formed by acting upon ethylene bromide with Ag<sub>2</sub>SO<sub>4</sub> and water (B and W), the barium salt is very soluble in water, and its solution decompose on warming into BaSO<sub>4</sub>, HBr, and glycol **Dibromo-diethyl sulphate** (CH<sub>2</sub>Br CH<sub>2</sub>)<sub>2</sub>SO<sub>4</sub>,

by action of ethylene bromide upon Ag<sub>2</sub>SO<sub>4</sub> in

presence of benzene (Beilstein and Wiegand, Annalen, 15, 1369); a heavy oil insoluble in water, soluble in benzene and ether, decomposed on heating. Heated with water, yields bromide of ethyl sulphuric acid, and finally HBr, H<sub>2</sub>SO<sub>4</sub>, and glycol.

Ethyl sulphites.

Mono-cthyl sulphite C<sub>2</sub>H<sub>5</sub>O SO OH, Ethyl sulphirous acid, Hyposulphethylic acid. The potassium salt KC<sub>2</sub>H<sub>5</sub>SO<sub>3</sub> is obtained by passing SO, into a well-cooled solution of NaOC, H5 in absolute alcohol (Rosenheim and Liebknecht, Ber 1898, 408) The ammonium salt is prepared by passing NH<sub>3</sub> followed by SO<sub>2</sub> into absolute alcohol (Divers and Ogawa, Chem Soc Trans 1899, 535) The alkali salts are very unstable in the solid state decomposing with evolution of SO2. Dilute mineral acids bring about this change in solution; ethyl iodide and alcohol at  $150^{\circ}$  yield ethyl sulphonate (q v)By treatment with PCI, diethyl sulphite yields the chloride of ethyl sulphurous acid C2H5O SOCI which boils at 122°, is decomposed by water into HCl, SO<sub>2</sub>, and alcohol, and gives, by further action of PCl<sub>5</sub> at 180°, thionyl chloride,  $\rm C_2H_5Cl$ , and POCI, (Michaelis and Wagner, Ber 7, 1074) Ethyl sulphurous acid is isomeric with ethyl

Delivity supintious with is sense in the single sulphone and C<sub>2</sub>H<sub>5</sub> SO<sub>2</sub> OH (q v ).

Diethyl sulphite C<sub>2</sub>H<sub>5</sub>O SO OC<sub>2</sub>H<sub>5</sub>; sp gr
1 1063 at 0° (Carius, J pr Chem [2] 2, 279); b p 161 3° From alcohol and S<sub>2</sub>Cl<sub>2</sub>; 18 parts of absolute alcohol are mixed with 50 parts of S<sub>2</sub>Cl<sub>2</sub>, previously warmed to 60°, the mixture is kept at this temperature for an hour, and then distilled, the portion boiling above 150° is redistilled (Warlitz, Annalen, 143, 74)

A colourless liquid, with mint-like odour, miscible with alcohol and ether, insoluble in water Decomposed by water; with PCl<sub>5</sub>, yields a chloride C<sub>2</sub>H<sub>5</sub>OSOCl, which, on distillation, splits up into C<sub>2</sub>H<sub>5</sub>Cl and SO<sub>2</sub> (Geuther, Annalen, 224, 223; Michaelis and Wagner, Ber 7, 1074). Decomposed by KOH into potassium ethyl sulphonate (Rosenheim and Liebknecht, ibid. 31, 405) Decomposes at 200° into SO<sub>2</sub> and ether (Prinz, Annalen, 223, 374), or by heating with SOCl<sub>2</sub>, to 120°, into SO<sub>2</sub> and C<sub>2</sub>H<sub>5</sub>Cl (Geuther).

Ethyl sulphydrate C<sub>2</sub>H<sub>5</sub>SH. Mercaptan, Ethyl mercaptan; b p. 36 2° (Nasım, Ber 15, 2882), sp gr 0 83907 at 20°/4° (N), 0 835 at 21°. By distilling Ba(C<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>)<sub>2</sub> with Ba(HS)<sub>2</sub> (Zeise, Annalen, 11, 1), by action of KHS upon C<sub>2</sub>H<sub>5</sub>Cl (Regnault, Ann. Chim Phys [2] 71, 390); by heating absolute alcohol saturated with SO<sub>2</sub> in a scaled tube, there are also formed C<sub>2</sub>H<sub>5</sub>HSO<sub>4</sub>, (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O, and H<sub>2</sub>SO<sub>4</sub> (Endemann, Annalen, 140, 336), from P<sub>2</sub>S<sub>5</sub> and alcohol (Kekulé, Annalen, 90, 310) It is best prepared by saturating a solution of KOH, sp gr 1 3, with H<sub>2</sub>S, mixing with a solution of Ca(C<sub>2</sub>H<sub>5</sub>SO<sub>4</sub>)<sub>2</sub> (also of sp gr. 13), and distilling on water-bath (Liebig, Annalen, 11, 14); according to Claesson (J pr. Chem [2] 15, 193), (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>S is also formed; NaC<sub>2</sub>H<sub>5</sub>S is prepared from the product of this action, the alcoholic solution evaporated to dryness, the (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>S removed by means of benzone, and the C<sub>2</sub>H<sub>5</sub>NaS decomposed by dilute H<sub>2</sub>SO<sub>4</sub>

It is a colourless liquid with a penetr ting garlic-like smell, slightly soluble in water, from which it crystallises on evaporation with 18H<sub>2</sub>O,

m.p. 12° (Claesson). It gives with bromine  $C_2H_5Br$ , HBr, and BrS (Friedel, Ladenburg, Annalen, 145, 189) HNO<sub>3</sub> forms  $C_2H_5SO_2$  SC<sub>2</sub>H<sub>5</sub>, or, if the acid be stronger, ethyl sulphonic acid Treated with Cu powder, loses its disagreeable odour, but is considerably decomposed thereby (Finish)

It forms crystalline compounds with S<sup>Cl</sup><sub>4</sub> (Demarcay, Bull. Soc chim [2] 20, 127). It absorbs NO, forming a dark-red solution

The hydrogen of mercaptan is replaceable by metals forming mercaptides; e.g.  $C_2H_5KS$  and  $Hg(C_2H_5S)_2$ , transparent plates, m.p.  $76^\circ$ – $77^\circ$  (Otto, Ber 13, 1290, 15, 125), very slightly soluble in cold alcohol, soluble in 12–15 parts hot alcohol (Liebig, Annalen, 11, 17). gives with HNO<sub>3</sub> the mitrate  $Hg(C_2H_7S)NO_3$  (Hofmann and Rabe, Zeitsch anorg Chem 17, 26),  $Pb(C_2H_5S)_2$ , yellow precipitate, formed when  $Pb(C_2H_5O_{22})_2$  and  $C_2H_5HS$  are mixed in alcoholic solutions  $As(C_2H_5S)_3$ , a foul-smelling oil formed from  $As(C_3H_3C_2)_2$  and  $C_2H_5NaS$ , decomposed by heat into As and  $(C_2H_5)_2S_2$   $Pt(C_2H_5S)_2$  pale-yellow precipitate (Zeise).

precipitate (Zeise). Ethyl sulphide ( $\rm C_2H_5)_2S$ ; b p 91° (Pierre, J 1851, 51), 91 9° at 754'7 mm (Beckmann, J. pr. Chem [2] 17, 451), 92·2°-93° ( $^{\circ}$  corr.) at 754 mm. (Nasim, Ber 15, 2282), 92·1° (corr.) (Thorpe and Rodger); sp gr 0·8367 at 0° (Pierre), 0 83676 at 20°/4° (N) From K $_2$ S and C $_2$ H $_5$ Cl (Regnault, Ann Chim Phys [2] 71, 387); from Zn(C $_2$ H $_5$ l $_2$  and SOCl $_2$  (Gauhe, Annalen, 143, 266), from KHS and KC $_2$ H $_5$ SQ $_4$  in alcohol (Hobson, Chem Soc Trans 10, 56), also from P $_2$ S $_5$  and (C $_2$ H $_5$ ) $_2$ O (Beckmann) To prepare the pure sulphide, heat the crude sulphide with Cu powder to 260°-280° (Finckh, Ber 1894, 1239)

It is a colourless liquid with strong garlic-like smell, insoluble in water. Bruhl states that in the pure state it has an ethereal odour. Chlorine acts upon it in the dark, forming substitution products: (1) (C<sub>2</sub>H<sub>3</sub>(I<sub>2</sub>)<sub>2</sub>S, vellow liquid, b.p. 167°-172°, sp.gr. 1.547 at 12°, (2) (C<sub>2</sub>H<sub>3</sub>(I<sub>3</sub>)<sub>2</sub>S, b.p. 189°-192°, sp.gr. 1.219 at 13.5°; (3) (C<sub>2</sub>H(I<sub>4</sub>)<sub>2</sub>S, b.p. 217°-222° By prolonged action of chlorine, C<sub>2</sub>Cl<sub>4</sub> is formed, and possibly (C<sub>2</sub>Cl<sub>5</sub>)<sub>2</sub>S (Riche, Ann Chim. Phys. [3] 43, 283; Annalen, 92, 358) (C<sub>2</sub>H<sub>2</sub>Cl<sub>3</sub>)<sub>2</sub>S also results from action of chlorine upon (C<sub>2</sub>H<sub>3</sub>Cl<sub>2</sub>)<sub>2</sub>S, which can be obtained from C<sub>2</sub>H<sub>4</sub> and Cl<sub>3</sub>S<sub>2</sub> (Guthrie, Annalen, 116, 241) Bromine forms (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>SBr<sub>2</sub>, a yellowish-red unstable body, which, by the action of KI, yields (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>SI<sub>2</sub>, a black liquid (Rathke, Annalen, 152, 214)

Ethyl sulphide combines with many metallic chlorides to form crystalline compounds, e.g.  $(C_2H_5)_2S$  HgCl<sub>2</sub>, white crystals, soluble in alcohol and ether, mp 119°,  $(C_2H_5)_2S$  HgI<sub>2</sub>, yellow needles, mp 110°;  $2(C_2H_5)_2S$  TiCl<sub>3</sub>, dark-red crystals (Demarcay) (For addition compounds, see Abel, Zeitsch physiol Chem 20, 268; Werner, Zeitsch anorg Chem. 15, 14; 17, 101; Smiles, Chem Soc Trans 1900, 164.)

Nitric acid (of sp gr 1 2) forms ethyl sulphoxide or diethyl sulphine oxide  $(C_2H_5)_2SO$ , a soluble oily liquid, solid at low temperatures, and decomposed by heating. Fuming intric acid converts this into diethyl sulphone  $(C_2H_5)_2SO_2$ ; bp. 248°; mp. 70°-72° (Oefele, Annalen, 127, 370; 132, 88);  $R\infty = 47.53$  (Kanonikow, J Russ. Ph.s Chem. Soc. 15

451); also formed from  $Pb(C_2H_b)_4$  and  $SO_2$ (Frankland and Lawrence, Chem. Soc. Trans 35, 244), and from (C2H5)2S, and a solution of KMnO<sub>4</sub> (O) Rhombic plates, soluble in 6.4 parts water at 16°

Triethyl sulphine (C2H5)3S forms compounds

C,H,HS (Cahours, ibid 135, 352, 136, 151), forms rhombic plates of disagreeable odour, and soluble in water By action of Ag<sub>2</sub>O, yields triethy, sulphine hydroxide (C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>SOH, a strong base, displaces NH4, and has caustic action on the skin, it forms a crystalline deliquescent mass, which unites with acids to form deliquescent salts yielding double compounds with many metallic salts;  $e\ g$ :

## PtCl<sub>4</sub>·2(C,H<sub>5</sub>)<sub>3</sub>SCl

dark-reddish prisms, soluble at 20  $7^{\circ}$  in 30 parts of water (Dehn, Annalen, Suppl 4, 90; Jorgensen, J pr Chem [2] 6, 82, Kraut, Annalen,

210, 321)
When  $(C_2H_5)_2S$  is heated with  $CH_3I$ , diethylmethyl sulphine rodide (C2H5)2CH3SI, an unstable non-crystalline body, is formed, from which other salts can be obtained; the chloride (C2H5)2CH3SCl forms many double salts (Kruger, J pr Chem The isomeride, ethyl-methylethyl [2] 14, 195) sulphine iodide, prepared from C<sub>2</sub>H<sub>5</sub>CH<sub>3</sub>S and C<sub>2</sub>H<sub>5</sub>I, is crystalline (Kruger)

Ethyl disulphide  $(C_2H_2)_2S_2$ , bp 152 8°-153.4° (corr) at 730 mm (Nasm, Ber 15, 2882), sp gr 0 99267 at 20°/4° (N) By distillation of KC<sub>2</sub>H<sub>5</sub>SO<sub>4</sub> with K<sub>2</sub>S<sub>2</sub> and water (Zoise, Annalen, 11, 1, Morin, ibid 32, 267, Lowig, Kopp, ibid. 35, 345, ('ahours, ibid 61, 98), by action of I upon a slight excess of a solution of C2H5NaS (Kekulė; Linnemann, *ibid* 123, 279), by action of S upon C<sub>2</sub>H<sub>5</sub>SNa (Bottger, *ibid* 223, 348); by distilling liquid thioacetaldehyde (Klinger,

Ber 32, 2195)

A colourless liquid, with garlic-like smell, very slightly soluble in water, with dilute HNO3, gives  $(C_2H_5)_2S_2O_2$ , an only liquid, which may be regarded as the ethyl salt of ethyl thiosulphonic acid C2H5SO3SC2H5.

Ethyl trisulphide  $(C_2H_5)_2S_3$ , from  $K_2S_3$  and  $KC_2H_5SO_4$  (Cahours, Bull. Soc chun. [2] 25, 184); also from (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>S<sub>2</sub> and S (H Muller); a colourless liquid, with unpleasant smell.

Ethyl tetrasulphide  $(C_2H_5)_2S_4$ , from  $S_2Cl_2$  and  $C_2H_5SH$  (Claesson, J pr Chem [2] 15, 214), a heavy oily liquid, with disagreeable odour

Ethyl pentasulphide  $(C_2H_5)_2S_5$ ; by heating  $(C_3H_5)_2S_4$  and S at 150°; a tough elastic solid

Ethyl sulphinic acid  $C_2H_5HSO_2$ , by action of Zn(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> upon SO<sub>2</sub> (Wischin, Annalen, 139, 364; Hobson, Chem Soc Trans 1858, 10, 58), the sodium salt is produced by oxidation of sodium mercaptide by dry oxygen (Claesson, J. pr. Chem. [2] 15, 199) The free acid is a syrupy liquid with a sweet taste, by action of HNO3, yields ethyl sulphonic acid and (C<sub>1</sub>H<sub>5</sub>SO<sub>2</sub>)<sub>3</sub>NO, crystals, m p 81 5°, very slightly soluble in water (Zuckschwerdt, Annalen, 174, 308) Forms salts, of which NaC<sub>2</sub>H<sub>6</sub>SO<sub>2</sub> crystallises from alcohol in anhydrous crystals (C.); the lead salt was prepared from Pb(C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>

and SO<sub>2</sub> by Frankland and Lawrence (Chem. Soc Trans 35, 244)

Ethyl chlorosulphinate Cl SO, C<sub>2</sub>H<sub>5</sub>. A faintly yellow hquid, b p 29°/13 mm, which decomposes at its bp, 122°, under ordinary pressure, into sulphui dioxide and ethyl chloride, formed by treating well-cooled ethyl alcohol with thionyl chloride (Stahler and Schiim, Bei 1911, 44, 319)

Ethyl sulphonic acid C<sub>2</sub>H<sub>5</sub> SO<sub>2</sub> OH by the oxidation of C2H5SH (Lowig and Weidmann, Pogg Ann 49, 329, Kopp, Annalen, 35, 346), C<sub>2</sub>H<sub>3</sub>CNS or of (C<sub>2</sub>H<sub>3</sub>)<sub>2</sub>S<sub>2</sub> (Muspratt, Chem. Soc Trans 1849, 1, 45) by HNO<sub>3</sub>, from C<sub>2</sub>H<sub>3</sub>I and K<sub>2</sub>SO<sub>3</sub> (Graebe, Annalen, 146, 37) By heating ethylene sulphonic acid with P and HI at 170° (Kohler, Amer Chem J 20, 688), by oxidation of C<sub>2</sub>H<sub>5</sub>CNS with hypochlorites (De Coninck, Compt. rend 126, 838), by action of cone HNO3 upon mercury mercaptide nitrate (Hofmann and Rabe, Zeitsch anorg Chem 17, 26) The acid is a crystalline deliquescent body; by action of ICI, at 150°, we get dichlorethyl sulphonic acid, excess of ICl<sub>3</sub> gives C<sub>2</sub>Cl<sub>6</sub> (Spring and Wessinger, Ber 15, 445), unites with bases to form salts (For halogen derivatives, see Kohler, Amer Chem J 19, 737, 20, 690, 21, 361 For salt Rosenheim and Liebknecht, Ber 31, 412) For salts, see

Ethyl sulphonate  $C_2H_5$  SO<sub>2</sub> OC<sub>2</sub>H<sub>5</sub>, bp 213 4° (corr), sp gr 1 1712 at 0°/0° 1 1452 at 20°/4° (Nasm, Ber 15, 2884), by action of ethyl sulphonic chloride  $C_2H_5$ SO<sub>2</sub>Cl and  $C_2H_5$ NaO (Carius, J 1870, 726, J pi Chem [2] 2, 262), from  $C_2H_5$ I and silver sulphite (archive sulphite (archive))

sodium ethyl sulphite (q v)

Ethyl sulphonic chloride  ${\rm C_2H_{2}SO_2Cl}$ , b p 177 5° (corr ) (Carius), 171° (Otto, Ber 15, 122), sp gr 1 357 at 22 5° (C), by action of PCL on NaC2H5SO3 (Gerhardt and Chancel, Compt rend 35, 690), by action of Cl upon ethyl oxysulphide (S and W lc) Colourless liquid,

with unpleasant smell

Ethyl thiocyanate C<sub>2</sub>H<sub>3</sub>SCN, bp 143° (Berthelot, Compt rend 130, 444), 146° (corr) (Buff, Ber 1, 206), sp gr 1 033 at 0°, 1 0126 at 19°, 0 8698 at 146° (B), by action of KCNS upon KC<sub>2</sub>H<sub>5</sub>SO<sub>4</sub> (Cahours, Ann Chim Phys [3] 18, 264, Annalen, 6<sub>5</sub>1, 95), or C<sub>2</sub>H<sub>5</sub>Cl (Lowig, Pogg Ann 67, 101), or C<sub>2</sub>H<sub>5</sub>I (Johnson and Levy Amer Chem J 1907, 38, 456), from Et<sub>2</sub>SO<sub>4</sub> and KCNS, yield 87 pc (Walden, Ber. 1907,

A mobile liquid with unpleasant odour, insoluble in water, miscible with alcohol or ether, by action of sodium forms NaCN and (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>S (Hofmann, Ber 1, 184), by heating with solution of KOH, yields (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>S<sub>2</sub>, KCN, and KCNO (Bruning, Annalen, 104, 198), with alcoholic KOH, forms (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>S<sub>2</sub>, CO<sub>2</sub>, NH<sub>3</sub>, but no KCNS (Lowig), but with alcoholic KHS yields C<sub>2</sub>H<sub>5</sub>HS and KCNS (L) Decomposed by hot solution of bleaching powder, yielding CO2, hot, yield (EtO)<sub>2</sub>SO<sub>2</sub> and NaCN Further action gives Et<sub>2</sub>S<sub>2</sub>, NaCN, and NaCNO, if all the hypochlorite is used up By action of aqueous  $NH_3$  at  $100^\circ$ , it yields  $(C_2H_5)_2S_2$ ,  $NH_4CN$ , urea, oxalic acid, and other products (Kremer, J. 1858, 401; Jeanjean, J. 1862, 364) Py passing Cl into it, there results (CNCl)<sub>3</sub> and a liquid C<sub>2</sub>H<sub>3</sub>SCl<sub>3</sub>, bp. 134°-135° (James, J. pr.

Chem [2] 30, 316) A chlorine substitution product  $C_2H_4\text{ClSCN}$  was obtained by James (viul [2] 20, 352) by heating  $C_2H_4\text{ClBr}$  with KCNS and alcohol, an oil heavier than water, with unpleasant smell, bp. 202°-203°; mp below --20°

Ethyl thiocyanate forms compounds with HCl, HBr, and HI (Henry, J 1868, 652; Glutz, Annalen, 153, 312) (For many other compounds, see Wheeler and Johnson, Amer Chem. J 26, 345, J Amer Chem Soc 24, 680, Wheeler and Jamieson, *ibid* 24, 743, Wheeler

and Mernam, ibid 24, 439)

Ethyl isothiocyanate C<sub>2</sub>H<sub>5</sub>NCS (L), Ethylthrocarbinide, mustard oil, b p 134°, sp gr 1 0192 at 0°, 0 9972 at 22°, 0 8763 at 133 2° (Buft, Ber I, 206), by action of C<sub>2</sub>H<sub>5</sub>NCO upon P<sub>2</sub>S<sub>5</sub> (Michael and Palmer, Amer Chem J 6, 259), also in very small quantity when Hg(CNS)<sub>2</sub> is heated with C<sub>2</sub>H<sub>5</sub>I (Michael, ibid. 1, 417), also by heating CSCl<sub>2</sub> and (C<sub>2</sub>H<sub>5</sub>)NH<sub>2</sub> (Rathke, Annalen, 167, 211) A solution of ethylamine in alcohol is treated with CS2, when crystals of ethyl ammonium ethyl dithiocarbamate separate out,  $2(C_2H_5)NH_2+CS_2=(C_3H_5)NHCSSNH_3(C_2H_5)$ A solution of this body is then treated with HgCl2, when the mercury compound is precipitated.

 $\begin{array}{l} (\mathrm{C_2H_5})\mathrm{NH} \times \mathrm{S} \times \mathrm{NH_3}(\mathrm{C_2H_5}) + \mathrm{Hg}(\mathrm{T_2} \\ = 2(\mathrm{C_2H_5})\mathrm{H_3}\mathrm{N}\mathrm{C1} + [\mathrm{NH}(\mathrm{C_2H_5})\mathrm{CS_2}]_2\mathrm{Hg} \end{array}$ 

On distillation, this body yields the mustard oil, HgS and H<sub>2</sub>S (Hofmann, Ber 1, 171, 2, 452) Delépine (Compt. 1end. 144, 1125) improves the yield of dithiocarbamate by using I molecule each of CS,, amine, and soda, and then distilling with basic lead acetate in place of HgCl2. Mustard oil is a liquid with very disagreeable smell, and in contact with the skin produces a burning sensation (Hofmann, Ber 1, 26), with absolute alcohol forms monoethyl throurethane (C<sub>2</sub>H<sub>5</sub>)NH CS OC<sub>2</sub>H<sub>5</sub>, a liquid, bp 204°-208°, with aldehyde ammonia yields the compound C<sub>2</sub>H<sub>15</sub>N<sub>5</sub>S, mp 135°-136° (Dixon, Chem. Soc. Trans. 1888, 414)

Thiocarbimides react with phenylhydrazine, yielding thiosemical bazides (Marckwald) Br in chloroform a ring compound, C2OS2(NEt)2, melting at 45°, is formed (Freund and Bachrach, Annalen, 285, 184)

Ethyl thiophens  $C_4SH_3(C_2H_5)$ 2-Ethyl theophen, bp 132°-134° (corr), sp gr 0 990 at 24° By action of  $C_2H_5Br$  and Na upon C4SH3I (Meyer and Kieis, Ber 17, 1560), also by action of Br upon thiophen, C<sub>2</sub>H<sub>5</sub>Br and Na (Schleicher, *ibid* 18, 3015, *ibid* 19, 671).

3-Ethyl throphen, bp. 135°-136°; spg1 1 0012 at 16°. By heating ethyl succinic acid with P<sub>2</sub>S<sub>3</sub> (Damsky, Ber 19, 3284, Gerlach, Annalen, 267, 146)

Di-ethylthiophen, bp 181°, sp gr 0 962 at 14°/14° An oil obtained by action of Na upon (',H<sub>5</sub>I and 10doethylth10phen (Muhlert, Ber

19, 633).

Ethyl telluride (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>Te, Tellurethyl; b p (°-138° (Marquardt and Michaelis, Ber 137°-138° 21, 2045; Heeren, J 1861, 565), from K,Te and KC,H,SO<sub>4</sub> (Wohler, Annalen, 84, 69; 35, 111; Mallet, Chem Soc. Trans 1853, 5, 71) A heavy, viscid, reddish-yellow liquid, with a very unpleasant smell, burns in air with blue flame, evolving fumes of TeO,; hardly soluble in water, and oxidised by exposure to air, poisonous  $\frac{\mathrm{HNO_3}}{\mathrm{(monoclinic\ prisms)}}$  forms  $\frac{(\mathrm{C_2H_5})_2\mathrm{Te},\mathrm{HNO_5}}{\mathrm{Strong\ HCl\ precipitate_3}}$  the oily compound  $(\mathrm{C_2H_5})_2\mathrm{TeCl_2}$  from a solution of the nitrate, ammonia converts this into [(('2H<sub>5</sub>)<sub>2</sub>Tc]<sub>2</sub>('l<sub>2</sub>O (hexagonal pusms), which, when treated with Ag<sub>2</sub>SO<sub>4</sub>, yields

 $[(C_2H_5)_2Te]_2,H_2SO_4$ 

(colourless pusms) By action of Ag,O upon a solution of the chloride, (C2H5)2TeO is produced,

which decomposes on concentration

When  $(C_2H_5)_2$ Te is acted upon by  $C_2H_5I$ , triethyl tellurium rodide (C, $H_5$ )<sub>3</sub>Te1 is formed, m p 90°-92° (Cahours, Bull Soc. chim. [2] 4, 40), which crystallises in yellow prisms

Ethyl ditelluride  $(({}^{\iota}_{2}H_{5})_{2}Te_{2}^{-1}$  is said to be formed in the preparation of  $({}^{\iota}_{2}H_{5})_{2}Te_{3}$  and to be a dark-red liquid, with high boiling-point

#### ETHY LAMINES

Compounds in which one or more atoms of

H in NH<sub>3</sub> are replaced by C<sub>2</sub>H<sub>5</sub>

Ethylamine C<sub>2</sub>H<sub>5</sub> NH<sub>2</sub>, b p 18 7°, sp gr
0 6964 at 8°, 0 708 at 2° (Hofmann, Ber 22, 699); mp -852°, bp 19° to 20° (Ladenburg and Krugel), sp gr 0 7013 at 4°, 0 6892 at 15° (Perkin), cut. temp 1852° (Schmidt), 1832° and cut piess 554 atm. (Berthoud), H C=4085 Cals (Berthelot, Ann Chim. Phys [5] 23, 244) From ethyl carbinide and KOH (Wurtz, Annalen, 71, 330, also Compt 1end 28, 223), from C<sub>2</sub>H<sub>5</sub>Br and NH<sub>3</sub> (Hofmann, Annalen, 74, 159), by the putrefaction of yeast and of flour (Hesse, J. 1857, 403, Sullivan, J. 1858, 231), by heating alcohol with NH<sub>4</sub>Cl at 300° (Berthelot, Ann Chim Phys [3] 38, 63), by action of nascent H upon acetomtrile (Mendius, Annalen, 121, 142) It is also produced in the dry distillation of the residues from beet-root in sugar manufacture (Duvillier and Buisine, Ann Chim Phys [5] 23, 317), by action of KC<sub>2</sub>H SO<sub>4</sub> upon alcoholic NH<sub>3</sub> at 120° (Eilenmeyer and Cail, J. 1875, 617), by action of Na upon CH<sub>3</sub> CONH<sub>2</sub> in boiling amyl alcohol (Gucibet, Compt 1end 129, 62), from EtOH and phospham at 225°-250° (Vidal, D. R. P. 64346), by action of Zn filings and HCl in the cold on aldehyde ammonia (Trillat and Fayollat) (Jean), by electrolytic reduction of nitroethane in H<sub>2</sub>SO<sub>4</sub> at 70°-75° (Pieron, Bull Soc chim [3], 21,784)

Preparation -(1) 1 volume of ethyl nitrate C<sub>2</sub>H<sub>5</sub>NO<sub>3</sub> is treated with 3 volumes of alcoholic ammonia for 12 hours at 100°, the free bases are separated from the NH<sub>3</sub> by neutralising with H<sub>2</sub>SO<sub>4</sub> or HCl and extracting with alcohol; the sulphates or chlorides are then decomposed by NaOH, and the free bases combined with pierre acid and crystallised The pierate of (C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>N separates first, in yellow needles, then the C2H5NH2 salt in blown prisms, the picrates are then separated and decomposed by HCl, and the bases liberated by KOH (Carey Lea, Chem. News, 5, 118).

(2) Crude & C, H, Cl, from manufacture of chloral, is heated for an hour with 3 times its volume of alcohol (95 p c ) previously saturated with ammonia, in an iron vessel heated by a On cooling, the NH<sub>4</sub>Cl is filtered water-bath off, and the remainder distilled until free from NH<sub>3</sub> and alcohol, the residue (hydrochlorides of

mon-, di-, and tri-ethylamine, mixed with a little NH<sub>4</sub>(I) is treated with a strong solution of NaOH, and the liquid drawn off and dried by means of solid NaOH; it consists of a mixture of the three bases, ethyl oxalate is then added, and the  $({\rm C_2H_5})_3{\rm N}$ , which is unacted upon, is distilled off, the residue consists of a mixture of solid diethyl oxamide  ${\rm C_2O_2(NHC_2H_5)_2}$  and liquid ethyl diethyloxamate  ${\rm C_2O_2N(C_2H_5)_2}$  over which are separated and the diethyl oxamide, after recrystallising from hot water, decomposed with KOH (Hofmann, Ber 3, 109, 776, also Duvillier and Buisine, Compt rend 88, 31)

According to Garner and Tyrer (Chem Soc Trans 1916, 109, 174), a mixture of mono-, di-, and tri-ethylamine can be separated by fraction-

ating with a 10-bulb column.

(3) By the interaction of ethyl bromide and excess of ammonia in alcoholic solution. By regulating the addition of the ethyl bromide to the ammoniacal solution it is possible to obtain the primary and secondary bases without any sensible admixture of the tertiary base (see Werner, Chem Soc. Trans 1918, 113, £)9)

Properties—Is a mobile liquid, with strong ammoniacal odour, is strongly caustic, and has an alkaline reaction, burns with yellow flame, is miscible in all proportions with water, displaces NH<sub>3</sub> from its salts, and, like ammonia, precipitates metallic hydroxides and oxides from salts It. however, dissolves the hydroxides of Al (Wurz), Au, and Ru (Carey Lea), and does not dissolve those of Cd, Ni, and Co. The chloride is decomposed on distillation into C<sub>2</sub>H<sub>3</sub>/NH, NH<sub>3</sub>, C<sub>2</sub>H<sub>4</sub>, and C<sub>2</sub>H<sub>5</sub>(T (Fileti and Piccini, Ber. 12, 1508). CrO<sub>4</sub> oxidises ethylamine to aldehyde and N (Carstanjen, J. 1862, 330).

Caro's acid oxidises EtNH, yielding CII,5OOH, CH<sub>2</sub>CN, CH<sub>3</sub>CH NOH, nitroethane, and CH<sub>4</sub>C(OH)NOH (Bamberger, Ber 35, 4293) It forms a compound C<sub>2</sub>II<sub>5</sub>NHK (Titherley)

Salts (v Wurtz, Annalen, 76, 329)—(C.H.)NH, HCl, deliquescent plates, mp 76°-80°, bp 315°-320° (with decomposition), sp gr 1 2045 at 21°/4° (Schiff and Monsacchi), molecular refract 35 97 (calc 35 95) (Kanonnikow, J pr Chem [2] 31, 347), soluble m alcohol, soluble in 0 42 part of water at 17° (S and M Zeitsch physikal Chem 24, 513) Prepared by heating 1 volume of C.H.(1 with 3 volumes alcoholic ammonia to 100° (Groves, Chem Soc Trans 1861, 13, 331) The hydrochloride and hydroiodide exhibit isodimorphism, both are monoclinic at ordinary temperatures, changing at higher temperatures to uniaxial modifications C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>, HgCl<sub>2</sub>, by action of alcoholic HgCl<sub>2</sub> on alcoholic ('\_1H<sub>5</sub>NH<sub>2</sub>, a crystalline precipitate, soluble in hot HCl solution. With excess of HgCl<sub>2</sub> in aqueous solution a white precipitate

# $2\mathrm{C}_2\mathrm{H}_5\mathrm{NH}_2,\mathrm{HgCl}_2+2\mathrm{HgO}$

is thrown down; by heating  $C_2H_5NH_2$  with excess of  $HgCl_2$ , a yellow precipitate

#### ClHg·OHg·NH C2H,

separates; whilst a crystallisable compound  $ClHg\cdot NH\cdot C_2H_5$  remains in solution (Köhler, Ber.

12, 2308)  $2C_2H_5NH_2$ , Pt(12, yellow powder, combines with  $(C_2H_5)NH_2$ , and forms

4C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>,PtCl<sub>2</sub>,2H<sub>2</sub>O

colourless crystals, soluble in water

(C<sub>2</sub>H<sub>5</sub> NH<sub>2</sub>,HCl)<sub>2</sub>,PtCl<sub>4</sub>, orange-yellow flat rhombohedrons (Weltzien, Annalen, 93, 272), hexagonal rhombohedrons (Topsoe, J. 1883, 619), sp.gr. 2 255 at 19°/4° (Clarke, Bei 12, 1399). C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>,HCl,AuCl<sub>3</sub>, golden-yellow monoclinic prisms, soluble in water (T)

 $(C_2H_5NH_1)_2$ ,  $H_2SO_4$ , deliquescent body, soluble in alcohol, forms double salts with other sulphates, eg ethyl-ammonium alum

 $(C_2H_5NH_2)_2$ ,  $H_2SO_4$ ,  $AI_2(SO_4)_3$ ,  $24H_2O_4$ 

octahedral crystals, soluble at 25° in 689 parts of water.

Ethylchloramine C<sub>2</sub>H<sub>5</sub>NHCl. A very pungent-smelling oil; sp gr 1 067 at 0°, formed by action of NaClO upon C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>,HCl. Heated with HCl, yields ethylamine (Berg, Ann. Chim.

Phys [7] 3, 319)

Ethylduchloramine, ethylmitrogenchloride

C<sub>2</sub>H<sub>5</sub> NCl<sub>2</sub>; b p 88°-89° at 762 mm.,
sp gr 1 230 at 15°, by passing Cl through
aqueous C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub> (Wurtz, Compt rend 11,
810), prepared by distilling a mixture of

C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>, HCl with 2! times its weight of bleaching powder and a little water (Tschermak, Ber.
9, 146), a yellow only liquid with a penetrating
smell, insoluble in water and acids, decomposes
on long standing into HCl, NH<sub>4</sub>Cl,C<sub>2</sub>H<sub>5</sub>NH<sub>3</sub>, HCl,
CHCl<sub>3</sub>, CH<sub>3</sub>CN, and CH<sub>3</sub>COCl (Kohler, Ber 12,
1870) The pure substance keeps unchanged
for long periods under water (Tschermak, ibid
12, 2129) Alkalis decompose it into acetic
acid and NH<sub>4</sub>, behaves towards many bothes as

12, 2129) Alkans decompose to mee accea and and NH<sub>1</sub>, behaves towards many bodies as free (1 Zn(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub> forms (C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>N and Zn(1<sub>2</sub> Ethyl di-iodoamine (2<sub>1</sub>H<sub>3</sub>NI<sub>2</sub>, and Ethyl dibromanine (2<sub>1</sub>H-NB<sub>12</sub>, are formed similarly; the former is a dark-blue liquid (Wurtz)

Ethyl nitramine C<sub>2</sub>H<sub>5</sub>NH NO<sub>2</sub>, m p + 6°, sp gr 1 1675 at 15° A colourless liquid, with an acid reaction, prepared by the action of NH<sub>3</sub> upon diethyl dimitro-oxamide. It gives two classes of alkyl derivatives (Umbgrove and Franchimont, Rec trav chim 16, 388). It is also formed from methyl ethylcarbamate and cone. HNO<sub>3</sub> (Franchimont and Klobbie, ibid. 7, 356).

Nitroethylisontramine ('H<sub>3</sub> ('HNO<sub>2</sub> N(OH)NO is obtained by the action of NO upon the Na derivative of introethane. It forms metallic salts, which are hydrolysed by mineral acids, giving NO and ethylintrolic acid (Traube, Annalen, 300, 106)

Annaich, 300, 100)

Diethylamine ( $({}^{\circ}_{2}\Pi_{5})_{2}NH$ , mp – 40° (Hofmann, Ber 22, 699); bp 57 5° (Hofmann, Annalen, 73, 91), 55 5° at 759 mm (Oudemans, Rec trav chim 1, 59); crit temp 223 3, crit press 36 6 atm (Berthoud), sp gi 0 7262 at 0°, 0 7107 at 15° (Hofmann, Ber. 22, 669), 0 7226 at 4°, 0 7116 at 15°, 0 7028 at 25°. From NH<sub>3</sub> and  $C_{2}H_{5}II$  (Hofmann), or NH<sub>3</sub> and  $C_{2}H_{5}NO_{3}$  (Lea, J 1861, 493, cf Gainei and Tyrer, lc) From  $C_{2}H_{5}Br$  (Werner, Chem Soc. Trans 1918, 113, 899) Prepared by the action of KOH upon the  $C_{2}O_{2}N(C_{2}H_{5})_{2}OC_{2}H_{5}$  obtained in manufacture of ethylamine (qv)

Also by action of very dilute NaOH solution upon  $[(C_2H_5)_2N(NO)]_2$ ,  $H_2SO_4$  (Kopp, Ber 8, 622). Also by action of EtOH upon phospham

at 225°-250° (Vidal, D R P 64346). By heating p-toluenesulphodiethylamide with chlorsulphonic acid at 130°-150° (Marckwald and v Droste-Hulshoff, Ber 31, 3263) By dissolving 50 grams diethylaniline in 148 grams HCl (sp gr 11), diluting with 75 cc H<sub>2</sub>O, adding 32 grams NaNO<sub>2</sub> in 50 cc H<sub>2</sub>O and pouring slowly through a reflux condenser into a boiling solution of 85 grams NaOH in 2 litres of H<sub>2</sub>O (Norms and Kimberley, Amer (hem J. 20, 60)

Price, Brazier, and Wood (J Soc Chem. Ind 1916, 35, 147) have described a convenient mode of preparing large quantities of diethylamine in the laboratory by a simple modification

of this method

NO C6H4 NEt2 + NaOH =NHEt2+NOC6H4 ONa

It a colourless, inflammable liquid with strong ammoniacal smell, very soluble in water, is a strong base, and acts like C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>, except that it does not dissolve Zn(OH), (Carey H<sub>2</sub>O<sub>2</sub> oxidises it to diethylhydroxylamine (Dunstan and Golding) The nitrate decomposes suddenly at 170°, and yields nitrosodiethylamine (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>NNO (Franchimont, Rec trav chim. 2, 95); the hydrochloride (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>NH,HCl forms non-deliquescent plates, mp 215°-217° (Wallach, Annalen, 214, 275), bp 320°-330° (W), sp.gr 1 0475 at 21°/4° (Schiff and Monster), and combines with many many and stately sacchi), and combines with many metallic chlorides,  $eg[(C_2H_6)_2NH,HC]_2PtCl_4$ , orangeyellow monoclinic crystals (Topose, J 1883, 619)

Diethylchlorumine (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>NCl A pungent-smelling oil, bp 91°, sp gr. 0 943 at 0°, formed by action of NaClO upon (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>NH It decomposes on keeping with formation of (('2Ho)2NH,HCl (Berg, Ann Chim Phys [7] 3,

320)

Diethylnitramine Et<sub>2</sub>N NO<sub>2</sub>, bp 206 5° at 757 mm, 93° at 16 mm, spgr 1 057  $15^{\circ}$ Formed by action of Eti upon EtNK·NO<sub>2</sub>; also by action of highest concentrated HNO<sub>3</sub> upon as-diethylurea It is decomposed by KOH at 150°-160° into EtNH<sub>2</sub>, HNO, and CH<sub>4</sub>CHO (Franchimont, Rec trav chim 6, 149, F and Umbgrove, *ibid* 16, 396)

Diethylisonitramine An oil, bp 46°-50° at 18~mm , sp gr  $1.000~at~15^{\circ}$  , formed by interaction of Etl and  $C_2H_5NAgNO_2$  KOH at 100° decomposes it into alcohol, N, and CH<sub>3</sub>CHO

(U and F. lc 399).

Nutrosodiethylamine  $(C_2H_5)_2$ NNO, bp 176.9° (corr); sp.gr. 0.951 at 17.5°, VD = 3.36 (calc = 3.53) (Knecht, Ber 10, 979) b p V D by distilling neutral solution of (C2H5)2NH,HCl with a concentrated solution of KNO<sub>2</sub> (Geuther, Annalen, 128, 251, J. 1871, 695), also from (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>NH and NOCl in ether solution at -15° to -20° (Solonina, J. Russ. Phys. Chem. Soc 30, 431), a yellow oil with aromatic smell, decomposed by HCl into (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>NH and HNO<sub>2</sub> Alcoholic KOH at 140° forms NH<sub>3</sub> HNO<sub>2</sub> Alcoholic KUII at 170 and (C<sub>2</sub>H<sub>5</sub>)NH<sub>2</sub>. Na amalgam forms N<sub>2</sub>O and  $(C_2H_5)_2NH$ . Triethylamine  $(C_2H_5)_3N$ ; b p 89°-89 5° at

736 5 mm. (Bruhl, Annalen, 200, 186), 89°-90 5° at 767 8 mm (Hofmann); spgr 0.735 at 15° (Hofmann, Ber. 22, 700), 0.7426 at 4°, 0.7331 at 15°, 0.7527 at 25° (Perkin), crit temp. 267 1° (Pawlewski, Ber 16, 2633), remains liquid at 75° or 15° or 15 -75° under 10 mm. pressure (H.) By the hours at 140°, β-ethyl-α-benzylhydroxylamine

action of NH<sub>3</sub> upon C<sub>2</sub>H<sub>5</sub>I (Hofmann, Annalen,

73, 91) or upon C<sub>2</sub>H<sub>5</sub>NO<sub>7</sub> (Carey Lea).

\*\*Preparation\*\*—(1) Crude ethylamine (from C<sub>2</sub>H<sub>5</sub>Cl and NH<sub>3</sub>) in alcoholic solution is treated with C, H, Cl (Duvillier and Buisine), (2) by dis-

tillation of (C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>N OH.

Properties—An only liquid with strong ammoniacal smell, slightly soluble in water; strong base. On strongly heating its salts,  $(C_2H_5)_3N$  is destroyed, with formation of a little  $(C_2H_5)_2N$  NO (Geuther, Zeitsch Chem 1886, 513)

The hydrochloride forms feathery non-deliquescent scales, and combines with many metallic salts to form double compounds N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>,HCl, sp gr 1 06885 at  $21^{\circ}/4^{\circ}$  1 part of water dissolves 15 parts salt (Schiff and Monsacchi) (C<sub>2</sub>H<sub>s</sub>)<sub>3</sub>N,HNO<sub>3</sub>, mp 98°-99° (Franchimont, Rec trav chim 2, 339).

Tetra-ethylammonium hydroxide

(C,H<sub>5</sub>)<sub>4</sub>N OH

formed from the iodide (C2H5)4NI by the action of Ag<sub>2</sub>O. It forms very deliquescent needles, is a strong base, which eagerly absorbs CO<sub>2</sub>, displaces NH3, precipitates metallic hydroxides, and gives a blue solution with sugar and CuSO, is decomposed by dry distillation, forming

tructhylamine, ethylene, and  $H_2O$ , when heated with  $C_2H_5I$ , gives  $(C_2H_5)_4I$  and alcohol Salts  $-(C_2H_5)_4NCI$ , forms deliquescent crystals, sp gr 1 0801 at 21°/4° (Schiff and Monsach), and yields many double salts,  $eg[(C_2H_5)_4NCl]_2HgCl_2$ , tetragonal crystals;  $(C_2H_5)_4NCl_1HgCl_2$ , triclimic crystals (Topsoc, J. 1883, 620),  $[(C_2H_5)_4NCl]_2PtCl_4$ , orange-yellow, crystalline (monoclinic (T)), difficultly soluble in water, almost insoluble in alcohol, (C2H5)4NBr, light-orange needles, mp. 78°, combines with Br to form an unstable pentablomide (Marquart, Ber 3, 284), gives with alcoholic I solution a precipitate of (('.2H<sub>5</sub>)<sub>4</sub>NI<sub>3</sub>, quadratic piisms (Weltzien, Annalen, 91, 33), which is also formed along with iodoform when the monoiodide is electrolysed between platinum electrodes (Goecke, Zeitsch Elektrochem 10, 249), the monoiodide (C2H5)4NI forms large crystals soluble in cold water and in alcohol, insoluble in ether, sp gi. 1 559 (Schroder, Ber 12, 562)

a-Ethyl hydroxylamine C  $_2$ H  $_5$ O NH  $_2$ . b p 68°, sp gr 0 8827 at 7 5° (Gurke, Annalen, 205, 277) , by the action of HCl upon benzoyl ethylhydroxylamine ethyl ether  $\dot{C}_6H_5CO$   $\dot{N}\dot{C}_2H_5$   $\dot{O}\dot{C}_2^{\prime}H_5$ , or upon the ethyl ether of benzaldoxime

C<sub>6</sub>H<sub>5</sub> CH NOC<sub>2</sub>H<sub>5</sub>

(Lossen and Zanni, ibid 182, 223)

A strongly smelling inflammable liquid, miscible with water, alcohol, or ether; gives with AgNO<sub>3</sub> a white precipitate, which, on heating, is changed to metallic silver, with acids, forms salts. eg C<sub>2</sub>H<sub>2</sub>ON,HCl, scaly plates, mp 128°, decomposed by heating with HCl to 150° into C<sub>2</sub>H<sub>5</sub>Cl, NH<sub>3</sub>O, HCl, and NH<sub>4</sub>Cl; unites with PtCl<sub>4</sub> to form (C<sub>2</sub>H<sub>7</sub>ON,HCl)<sub>2</sub>PtCl<sub>4</sub>; prismatic crystals, soluble in water and in absolute alcohol (Lossen and Zanni) The sulphate (C2H7ON), H2SO4 crystallises with difficulty, and is soluble in water and in alcohol.

 $\beta$ -Ethyl hydroxylamine  $C_2H_5NHOH$ ; m p. 59°-60°, with decomposition; sp gr 0 9079 at 63 9° (Bruhl). Formed by heating for 8

with conc HCl (Behrend and Leuchs, Annalen, 257, 239); from hydroxylamine and EtI (Lobry de Bruyn, Rec. trav. chim 13, 48; Hantzeh and Hilland, Ber. 31, 2065; Dunstan and Golding, Chem. Soc Trans. 75, 807); by heating 1 part m-nitrobenzylaldoxime-N-ethyl ether with 7 parts cone HCl (Kjellin, Ber 26, 2378); by electrolytic reduction of nitroethane in H<sub>2</sub>SO<sub>4</sub> solution at 15°-20° (Pierron, Bull. Soc. chim. [3] 21, 784).

It is very easily soluble in water or alcohol,

less so in ether, benzene, or cold ligroin reduces Barreswil's solution, and is reduced by

HI to  $C_2H_5NH_2$ .

\$\textit{\beta}\$B\$-Diethyl hydroxylamine (\$C\_2H\_5\$)\_N OH, bp 130°-134°. with decomposition (Dunstan and Golding), bp. at 15 mm 47°-49° (Lachman), \$\text{COMP}\$ (D. and G.), 0.8784 sp gr 0 8771 at 15°/15° (D. and G.), 0 8784 (L). An oil formed by action of Etl upon NH<sub>2</sub>OH, and also from  $(C_2H_5)_2$ NH and H<sub>2</sub>O<sub>2</sub> (D. and G. Chem. Soc. Trans. 75, 800). From addition product of diphenylnitrosamine and Zn(Et)<sub>2</sub> by decomposition with water, along with diphenylamine and Zn(OH), (Lachman, Ber 33, 1022)

Is a feeble base, soluble in water, alcohol, or ether. Reduces AgNO<sub>3</sub> and HgCl<sub>2</sub>, and, on heating, CuSO<sub>4</sub> Reduced with HI, yields heating,  $C_1$  ( $C_2H_5$ )<sub>2</sub>NH.

αβ-Diethyl hydroxylamine C<sub>2</sub>H<sub>5</sub>NH OC<sub>2</sub>H<sub>5</sub>; b.p. 83°; sp gr 0 829 at 0°; formed by interaction of a-ethylhydroxylamine and EtBr. It is easily soluble in water, alcohol, or ether Heated

with cone. HCl at 200°, yields  $HN_2(C_2H_5)$  and  $C_2H_5(l)$  (Lossen, Annalen, 252, 230).

The hydrochloride  $C_4H_{11}NOHCl$  of an isomeric compound is obtained by action of Sn and HCl upon C<sub>2</sub>H<sub>5</sub>NO<sub>3</sub> (Lossen, *ibid* Suppl 6, 238) The salts are more soluble in water than those of hydroxylamine. The free base is a syrup, easily soluble in water, and has strong basic properties. The platinum salt is in the form of orange-yellow plates, soluble in warm alcohol and in water

Triethyloxamine (C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>NO is a colourless crystalline mass obtained from ββ-diethylhydroxylamine and EtI. Also from NEt, and

H<sub>2</sub>O<sub>2</sub> (Dunstan and Golding; Lachman)
It is very easily soluble in water, less so in CHCl<sub>3</sub> and insoluble in ether Decomposes on heating. Reduces Barreswil's solution on heating It gives a precipitate with CuSO<sub>4</sub> and AgNO<sub>3</sub> On reduction, yields triethylamine

Triethyl methyl ammonium iodide

# $N(C_2H_5)_3CH_3I$

obtained by action of  $N(C_2H_5)_3$  upon  $CH_3I$  (Hofmann, Annalen, 78, 277) (for other Salts, v Topsoe, J. 1883, 620) The picrate melts at v Topsoe, J. 1883, 620) The picrate 267°-268° (Lossen, Annalen, 181, 374)

Diethyl dimethyl ammonium iodide, obtained from N(CH<sub>3</sub>),H and C<sub>2</sub>H<sub>5</sub>I; or from N(C<sub>2</sub>H<sub>5</sub>),H and CH<sub>3</sub>I (V. Moyer and Lecco, Annalen, 180, 177) (for other Salts, v Topsoe, lc) The picrate melts at 285°-287° (Lossen, Annalen, 181, 374)

Ethyl trimethyl ammonium iodide is formed from  $(CH_3)_3N$  and  $C_2H_5I$  (Müller, *ibid*. 108, 1) (for other *Salts*, v. Topsoe). The picrate melts at 299°-300° (Lossen)

Triethyl propyl ammonium iodide  $N(C_2H_5)_3(C_3H_7)I$ 

crystallises in large needles (Mendius, Annalen, **M**1, 136).

Diethyl \*\*soamylamine  $N(U_1H_5)_2(C_5H_{11})$ ; b p 154°; from the distillation of  $N(U_2H_5)_3(C_5H_{11})OH$  (Hofmann),  $[N(U_2H_5)_2C_5H_{11},HCl]_2PtCl_4$ , yellow

### Triethyl isoamyl ammonium iodide

$$N(C_2H_5)_3(C_5H_{11})I$$

formed by the action of  $N(C_2H_5)$ , upon  $C_5H_{11}I$  (Hofmann, Chem. Soc. Trans. 1852, 4, 313, Annalen, 78, 279), the hydroxide decomposes on heating as follows -

C<sub>5</sub>H<sub>11</sub>(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>N OH

 $= N(C_5H_{11})(C_2H_5)_2 + C_2H_4 + H_2O$ 

# Diethyl methyl isoamyl ammonium iodide

 $N(C_2H_5)_2CH_3(C_5H_{11})I$ 

is formed from N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>(C<sub>5</sub>H<sub>11</sub>) and CH<sub>3</sub>I (Hofmann)

### Ethyl-methyl-150amylamine

NC2H2(CH3)(C5H11)

b p  $135^{\circ}$ ; is obtained by distillation of the hydroxide of the radical of the preceding compound (Hofmann).

Acetethylamide CH<sub>3</sub> CONHC<sub>2</sub>H<sub>5</sub>; b p 205°, sp gr 0 942 at 4 5° (Wurtz); b p. 204°-205° (Titherley), from ethyl acetate and ethylamine (Wurtz, Ann Chim Phys [3] 30, 491, Compt. rend 36, 180; Annalen, 76, 334) or from C<sub>2</sub>H<sub>5</sub>NCO and HC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> (Wurtz, *ibid* 88, 315), from CH<sub>3</sub>CONHNa and Et1 at 150° The hydrochloride forms deliquescent needles, melting at 60° With NaNH, yields CH3CONNaC2H5, which, with KMeSO4, gives CH3CONMeEt, an oil boiling at 180°, and soluble in water (Titherley, Chem Soc Trans 1901, 393) By action of Cl yields ethyl acetyl nitrogen chloride NClEt(C2H3O) (Norton, Tehermak, Bull Soc chim. [ii ] 30, 106, Compt rend 86, 1409), an oily unstable liquid, slightly soluble in water

N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>C<sub>2</sub>H<sub>3</sub>O; bp 0 9248 at 8 5° (Wallach, Acetdiethylamide 185°-186°; sp gr Annalen, 214, 235)

Ethyl hydrazines; Ethylamide.

1 Ethyl hydrazine C<sub>2</sub>H<sub>5</sub>HN NH<sub>2</sub>; b p 99 5° at 709 mm, is prepared by reducing nitrosodiethyl urea N(C<sub>2</sub>H<sub>5</sub>)H·CO N(NO)C<sub>2</sub>H<sub>5</sub> by means of zinc-dust and acetic acid; this yields ethyl hydrazine urea (semi-carbazide)  $N(C_2H_5)H$  CO  $N(NH_2)C_2H_5$ , and this, by action of HCl, forms  $C_2H_5HN$   $NH_2$ ,  $CO_2$ , and  $C_2H_5NH_2$ . by passing in HCl the chloride separates out, which is then decomposed by strong KOH solution and solid KOH, and the resulting oily liquid distilled over BaO (Fischer, Annalen, 199, 287); also by reducing ethylnitramine with Zn and HCl (Thiele and Meyer, Ber 29, 963)

Is a colourless liquid, with faintly ammoniacal odour; has great affinity for water, in which it is soluble, as also in alcohol, ether, chloroform, and benzene; fumes in air and destroys cork and caoutchoue; reduces Barreswil's solution in the cold. Br water liberates N. It precipitates metallic oxides, and gives the isonitrile reaction with KOH and CHCl<sub>3</sub>. Forms two chlorides: (1) C<sub>2</sub>H<sub>8</sub>N<sub>2</sub>,2HCl, white needles, very

soluble in water and alcohol; on heating to 110° yields (2) C<sub>2</sub>H<sub>8</sub>N<sub>2</sub>,HCl, an amorphous horn, deliquescent body. The sulphate crystallises from alcohol in thin plates, and is very soluble in water By heating C<sub>2</sub>H<sub>5</sub>NH NH<sub>2</sub> with K<sub>2</sub>S<sub>2</sub>O<sub>7</sub> at 80°-100°, potassium ethyl hydrazine sulphonate C<sub>2</sub>H<sub>5</sub>NH NHSO<sub>3</sub>K is obtained; this forms fine needles, very soluble in water, much less

soluble in alcohol or ether (Fischer)

2 as-Diethylhydrazine (C2H5)2N NH2; bp. 96°-99°; prepared by action of zinc-dust and acetic acid upon a solution of mitrosodiethylamine (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>(NO)N; 30 grams (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>(NO)N dissolved in 300 grams water are mixed with 150 grams Zn dust and 150 grams C2H4O (of 50 p c) kept at 20°-30°, and, towards the end, at 40°-50° Excess of Zn is then dissolved by HCl, an excess of NaOH added, and the mixture distilled, the distillate is neutralised with HCl and NH<sub>4</sub>Cl separated by crystallisation the mother liquor  $(C_2H_5)_2N_2H_2$  mixed with  $(C_1H_5)_2HN$  is freed by solid KOH. By the action of HCNO diethylhydrazine urea (semicarbazide) NH<sub>2</sub>CO NH N(C<sub>2</sub>H<sub>5</sub>), is formed 4 grams of this compound are heated with 15 grams of HCl in a sealed tube at 100°, diluted and the NH4Cl separated; on treatment with KOH, pure  $(C_2\Pi_5)_2N_2H_2$  is obtained (Fischer, Annalen, 199, 308).

Is a colourless liquid with a faintly ammoniacal smell, very hygroscopic and soluble in water, alcohol, ether, chloroform, and benzene; reduces Barreswil's solution only on warming; is monobasic; the salts are soluble in water and alcohol, and difficult to crystallise  $[(C_2H_5)_2N_2H_2$   $HCl]PtCl_4$  forms fine yellow needles, soluble in water With SOCl<sub>2</sub> in icecold ether, yields (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>N NSO, an aromatic oil, b p 73° at 20 mm It is slowly hydrolysed by water (Michaelis and Storbeck, Ber 26, 310)

3 sym-Diethylhydrazine C<sub>2</sub>H<sub>5</sub> HN NH C<sub>2</sub>H<sub>5</sub>, b p at 758 mm 84°-86°. Formed by heating at 100° for 3 hours 1 part sym-diethyl-diformylhydrazide and 2 parts fuming HCl, an oil; heated with cone HCl, at 155°, yields NH<sub>4</sub>Cl and  $\rm C_2H_5Cl$  With yellow HgO forms HgEt<sub>2</sub>. With HNO, yields ethyl nitrite The hydrochloride melts at 160° with decomposition (Harries, Ber 27, 2279)

Triethyl-azonium iodide (C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>NNH<sub>2</sub>I 18 formed by action of diethylhydrazine upon 11 times its weight of C2H5I (Fischer, Annalen, 199, 316). It forms needles very soluble in water and in hot alcohol; insoluble in ether and strong alkalis By action of Ag<sub>2</sub>O, gives the strongly alkaline free base, which, on heating, yields  $(C_2H_5)_2N$   $NH_2$  and  $C_2H_4$ . By the action of nascent H (Zn and  $H_2SO_4$ ), the iodide yields HI, NH<sub>3</sub>, and  $(C_2H_5)_3N$ 

Tetra-ethyl tetrazone  $(C_2H_5)_2N$  N.N  $N(C_2H_5)_2$ ; by the action of yellow HgO upon a cold aqueous solution of (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>N NH<sub>2</sub> (Fischer, Annalen, 199, . 319), is an only liquid with a disagreeable smell, not volatile without decomposition, and which does not freeze at  $-20^{\circ}$ . When decomposed by heat, yields N and  $(C_2H_5)H_2N$ . Forms an explosive oily compound with I. Is a strong base, though its salts are very unstable. It reduces silver solutions with formation of a mirror The solution decomposes on heating, yielding N,  $N(C_2H,)_2H$ , and aldehyde The Pt doubl salt [(C2H5)4N4,2HCl]2PtCl4 forms yellow prisms soluble in water.

Ethylene C<sub>2</sub>H<sub>4</sub>, Olefiant gas, Heavy car-etted hydrogen, Bicarburetted hydrogen, buretted hydrogen, Elayl, Ethene, Etherin, discovered by Deiman, Van Trootswyk, Bondt, and Lauwerenburgh (Crell's Ann. 1795, [2] 195, 310, 430) Obtained by action of H2SO4 upon alcohol; the addition of Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> increases the rate of production (Senderens, Compt rend 1910, 151, 392). The sulphuric acid can be replaced by syrupy phosphoric acid, sp gr 1.75 (Newth, Chem Soc. Trans 1901, 915); or alcohol vapour may be passed over heated phosphoric acid, or over thin layers of aluminium sulphate at 350°-400°; by the dry distillation of many organic bodies (hence occurs to the extent of 4 or 5 pc in coal gas). Also formed when CS<sub>2</sub> and H<sub>2</sub>S (or PH<sub>3</sub>) are led over heated copper (Perthelot, Annalen, 108, 194) To prepare the gas, a mixture of 25 parts (by weight) of alcohol and 150 parts sulphuric acid is heated to 170°, and a mixture of 1 part of alcohol and 2 of H2SO4 is allowed to drop slowly into the vessel by means of a funnel. The gas is passed through H2SO4, NaOH solution, and then again through H2SO4. For the preparation of small quantities of pure gas, an alcoholic solution of C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> is warmed with granulated Zn (Sabanejeff, J. Russ. Phys. Chem. Soc 9, 33) Equal volumes of CO and H led over coke impregnated with reduced Ni and Pd heated at  $95^{\circ}-100^{\circ}$ , yield  $8.3~{\rm p.c.}$  C<sub>2</sub>H<sub>4</sub> (Orloff, 1bid. 40, 1588)

Ethylene is produced on an industrial scale by the method of Maihle by passing the vapour of alcohol over amorphous alumina heated to 360°. For details, see Sprent, J. Soc. Chem.

Ind 1913, 171

Properties —A colourless gas, liquefied by a pressure of 42½ atmospheres at -11°, b p -105° (Calletet, Compt rend 94, 1224), -102°-103° (Wroblewsky, Olszewsky, Monatsh 4, 338), -103 9° (Burrell and Robertson); mp -169°, bp at 757 mm. -104 3°, sp gr 0 6095 at -102 7° (Ladenburg and Krugel, Ber 32, 49, 1821); -102 5°, sp gr 0 565 (Maass and McIntosh, J. Amer Chem. Soc 1014 36 737). crit. temp. 9 5° becomes 1914, 36, 737); crit. temp. 95°; becomes liquid at 1° and 45 atmospheres, at 4° and 50 atmospheres, at 8° and 56 atmospheres, and at  $10^{\circ}$  and 60 atmospheres (C);  $L_{\rm D} = 1.26031$ (Batuecas, Helv Chim Acta, 1918, 1, 136) I volume of water absorbs at  $t^{\circ}$ ,

#### $0.25629 - 0.009136t + 0.0001881t^{2}$

volumes C<sub>2</sub>H<sub>4</sub> (Bunsen) Alcohol (of sp gr 0 792 at 20°) absorbs  $3.59498 - 0.05772t + 0.000681t^2$ volumes (Carius, Annalen, 94, 133); H.C.p. =341.4 Cals; H.F. (from amorphous C) = -9.4 Cals. (Berthelot, Ann. Chim. Phys. [5] 23, 180).

Ethylene forms an explosive mixture with oxygen, and explodes, even in the cold, in contact with ozonised oxygen (Houzeau and Renard, J 1873, 319) Led through a hot tube, it splits up into C, H, CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, and C<sub>2</sub>H<sub>2</sub> (Berthelot, Annalen, 139, 277; Bone and Coward, Chem. Soc. Trans. 1908, 1197). When heated in an iron tube under a pressure of about 70 atmospheres it undergoes rapid polymerisation at 380°-400°. The liquid polymerisation products consist of mixtures of saturated, e-hylenic, and polymethylene hydrocarbons; viz.

iso-pentane, hexane, heptane, octane, nonane, amylene, hexylene, nonanaphthene, and  $\alpha\beta$  decanaphthene, hen-, deca-, dodeca-, tetradeca-, and pentadeca-naphthenes At atmospheric pressure ethylene does not polymerise, even at 600° (Ipatieff, Ber. 1911, 44, 2978) Decomposed 600° (Ipatieff, Ber. 1911, 44, 2978) Decomposed by leading over reduced Ni at 300° into C, C<sub>2</sub>H<sub>6</sub>, CH<sub>4</sub>, and H (Sabatier and Senderens, Compt rend. 124, 616, 1358). By the action of the electric spark it forms C<sub>2</sub>H<sub>2</sub> and H, and then C and H (Wilde, Zeitsch. Chem 1866, 735) It combines with H to form C<sub>2</sub>H<sub>6</sub>, when the mixture is passed through a hot tube (Ann. Chim. Phys [4] 3, 431; Bull Soc chim [ii.] 39, 145) or when in contact with Pt black in the Chim. Phys [4] 9, 431; Bull Soc chim [ii.] 39, 145), or when in contact with Pt black in the cold (Wilde, Ber 7, 354) By heating a mixture of C<sub>2</sub>H<sub>4</sub> and CO<sub>2</sub> at 400°, aldehydo is formed (Schutzenberger, Bull. Soc. chim. [ii.] 31, 482) CrO<sub>3</sub> converts it into aldehydo at 120° (Annalen, 150, 373). Fuming sulphuric acid absorbs readily large quantities of C<sub>2</sub>H<sub>4</sub> at ordinary temperatures, as does also common H SO<sub>2</sub> at 160°-170° (Butlerow Ber 6, 193) H<sub>2</sub>SO<sub>4</sub> at 160°-170° (Butlerow, Ber. 6, 193), forming C<sub>2</sub>H<sub>5</sub>HSO<sub>4</sub> (Faraday). C<sub>2</sub>H<sub>4</sub> combines at 100° with strong HI or HBr solution, but not with HCl (Berthelot, Annalen, 104, 184; 115, With bromine water forms a mixture of othylene bromohydrin and ethylene dibromide (Read and Williams, Chem Soc Trans 1917, 111, 240). N<sub>2</sub>O<sub>4</sub> reacts with ethylene to form cthylene nitrosite, and  $N_2O_5$  to form ethylene nitrate (Demjanow, Chem. Zentr. 1899, 1–1064) In presence of AlBr<sub>3</sub>, HBr unites at 0° with C<sub>2</sub>H<sub>4</sub>, forming C<sub>2</sub>H<sub>5</sub>Br With Cl<sub>2</sub>O, chlorethyl chloracetate CH<sub>2</sub>Cl CO<sub>2</sub>H<sub>4</sub>Cl is formed, whilst ClO<sub>2</sub> yields CH<sub>2</sub>Cl CO<sub>2</sub>H. C<sub>2</sub>H<sub>4</sub> combines with BF<sub>3</sub> ir sunlight to form C<sub>2</sub>H<sub>3</sub>BF<sub>2</sub> (Landolph, Ber. 12, 1586), a liquid, b p. 124°-125°, sp gr 1 0478 at 23°. Ethylene acts as a divalent radical, and forms salts with most of the acids.

radical, and forms salts with most of the acids.

Ethylene acetates (Wurtz, Compt rend. 43, 199; Atkinson, Phil. Mag [4] 16, 433).

1 Ethylene monoacetate C<sub>2</sub>H<sub>4</sub>(C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>)OH, monoacetate glycol; bp. 190°-190 5° (corr) (Perkin, Chem Soc Trans. 45, 505); sp gr. 1·1108 at 15°/15°, 1 1018 at 25°/25° (P.); produced by action of KC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> upon C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> in presence of alcohol (Atkinson, 1c; cf. Seelig, D. R. P. 41507). A liquid not misciple with water: yields

41507) A liquid not miscible with water; yields glycol C<sub>2</sub>H<sub>4</sub>(OH), by heating with C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> and alcohol (Demole, Annalen, 177, 45); decomposed by heating with C2H3OCl into acetochlorhydrin

by neating with  $C_2H_3OC1$  into accoording the and ethylene diacetate (Lourenço, thid 114, 127) 2 Ethylene diacetate  $C_2H_4(C_2H_3O_2)_2$ ; b p  $186^\circ-187^\circ$ ; sp gr  $1^\circ-128$  at  $0^\circ$ ; prepared from  $C_2H_4Br_2$  or  $C_2H_4I_2$  and  $AgC_2H_3O_2$  (Wurtz, Ann Chim. Phys. [3] 55, 433), or from  $C_2H_4Br_2$  and anhydrous  $KC_2H_3O_2$  at  $150^\circ-200^\circ$  (Denoted in the second of th mole).

A liquid soluble in 7 parts of water, isomeric with ethylidene diacetate CH<sub>3</sub> CH(C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>)<sub>2</sub>; b p 108 8°; sp gr. 1 061 at 12°; prepared from aldohyde and (C<sub>2</sub>H<sub>3</sub>O)<sub>2</sub>O. Dichlorethylene diacetate C<sub>2</sub>H<sub>2</sub>Cl<sub>2</sub>(C<sub>2</sub>H<sub>3</sub>O)<sub>2</sub>C; b.p. 120° at 20 mm; is produced by action of chloracetic acid upon C<sub>2</sub>H<sub>2</sub> (Prudhomme, Zeitsch. Chem. 1870, 397).

Ethylene acetochlorhydrin

# CICH<sub>2</sub>·CH<sub>2</sub>·O·C<sub>2</sub>H<sub>3</sub>O

b p. 145°; sp gr. 1 1783 at 0°; by heating a mixture of glycol and acetic acid at 100° with HCl (Simpson, Annalen, 112, 147); from C<sub>2</sub>H<sub>4</sub>

and CH2Cl CO2H (Schutzenberger and Lippman, i id. 138, 325); insoluble in water; yields ethylene oxide by action of KOH.

Ethylene acetoiodohydrin ICH, CH, O C, H<sub>3</sub>O,

heavy oil; and

Ethylene acetobromohydrin

BrCH, CH, O C, H,O

b p. 161°-163°; are prepared from

C2H4(C2H4O2)OH

by action of HI and HBr respectively, the former in the cold in presence of  $C_2H_4(OH)_2$  and  $CH_3$   $CO_2H$  (Simpson, Annalen, 113, 123), the latter at 100' (Demole, ibid 173, 120)

Ethylene glycols.

Ethylene glycols.

1 Ethylene glycol CH\_OH CH\_OH, glycol, mp —17.4°, bp 197.37° (Ladenburg and Krugel, Ber 32, 1821); bp 197°-197.5° at 764.5 mm (Wurtz), 196°-199° (corr) (Perkin, Chem Soc Trans. 45, 504), 195° at 751.3 mm (Pribram and Handl, Monatsh 2, 673); sp.gr. 1 125 at 0° (W), 1·1279 at 0° (P and H.), 1 1168 at 15°/15°, 1 112 at 25°/25° (Perkin).

Discovered by Wurtz (Corput, 1974.43, 190.

Discovered by Wurtz (Compt 1end 43, 199; Ann Chim Phys [3] 55, 400). Formed by action of ethylene diacetate upon KOH (W), from  $C_2H_4\mathrm{CH}$  and moist silver oxide at  $160^\circ-200^\circ$ (Simpson, Annalen, Suppl 6, 253); by heating ethylene monacetate with Ba(OH), or with water in sealed tubes (Debus, Annalen, 110, 316); by heating a strong aqueous solution of neurine, when glycol and N(CH<sub>3</sub>)<sub>3</sub> are formed (Wurtz, *ibid* Suppl. 6, 200); by heating ethylene chloride or bromide with PbO and 15-20 times its volume of water at 140°-170° (Eltekow, Ber. 6, 558); or with ferric oxide, or with an alkali formate and an alcohol under pressure; from ethylene bromide and water at 140°-150° (Niederist, Annalen, 186, 393), together with polyethylene glycol, by action of C<sub>2</sub>H<sub>4</sub>O (oxide of ethylene) upon water in scaled tubes (Wurtz, 1bid. 113, 255)

A mixture of glycols, in which ethylene glycol predominates, may be prepared by decomposing petroleum vapours in an iron tube heated at about 650°, converting the product into dichlorethane, &c., purifying this compound by distillation and treating the distillates with alkali carbonates in a closed vessel (Hibbert,

Met & Chem Eng. 1918, 19, 571)

Preparation -1 600 grams ethylene bromide, 700 grams fused potassium acetate, and 1500 grams alcohol (sp gr 0 83) are heated in a flask with reflux condenser until all action ceases, cooled, filtered from KBr and filtrate distilled, the portion distilled above 140° is decomposed by slight excess of Ba(OH)<sub>2</sub> and heated for 12 hours on water-bath The excess of baryta is removed by CO<sub>2</sub>, the liquid evaporated to onethird its volume, the residue treated with a mixture of alcohol and ether, the liquid removed from the crystals and distilled, first on water-bath, to separate alcohol and ether, and then over the flame; the portion distilling above 185° is then redistilled (Erlenmeyer, Annalen, 192, 244).

2. A mixture of 188 grams ethylene bromide, 138 grams potassium carbonate and 1 litre of water [or 32 parts C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub>, 25 parts K<sub>2</sub>CO<sub>3</sub>, 350 parts H<sub>2</sub>O (Groscheintz, Bull Soc. chim. [ii] 31, 293)], is heated with reflux condenser for

10 hours, is then evaporated to small volume on water-bath, the liquid portion poured off th crystals of KBr, the latter are washed with absolute alcohol. The filtrate and alcoholic washings are then distilled as in previous method; a loss owing to formation of C2H3Br is experienced by this process (Zeller and Huffner, J pr. Chem [2] 11, 229) Haworth and Perkin (Chem. Soc Trans. 1896, 175) obtained a 60 p c yield by digesting 138 grams K<sub>2</sub>CO<sub>3</sub> in 1000 c c. H<sub>2</sub>O with 188 grams C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> under a reflux condenser. The solution is evaporated very slowly to a syrup, and then extracted with a mixture of alcohol and ether.

3 1 part of ethylene bromide and 26 parts water are heated to 100° in a sealed vessel for 130 hours. This method gives a yield of over 60 p.c. of the theoretical amount (Niederist,

Annalen, 196, 354)

4. Glycol diacetate is decomposed with powdered Ca(OH)2, and the mixture distilled under reduced pressure. The yield of glycol is 93 p.c. of theory (Henry, Rec. trav. chim. 18. 221).

5 It may also be prepared by heating ethylene diacetate with methyl alcohol and hydrochloric acid (Henry, Bull. Acad. Roy

Belg. 1896, 732).

1 mol of ethylene dichloride, 5 vols. of methyl alcohol and 3 mols, of sodium formate are heated together at 165°-170° in an autoclave for 7 hours After cooling, the methyl alcohol is separated and distilled at ordinary prossure, the glycol under reduced pressure (Brooks and Humphrey, J. Ind. Eng Chem 1917, 9, 750)

Glycol is a colourless, odourless, syrupy liquid with sweet taste, miscible with alcohol and water, but almost insoluble in ether parts ether dissolve 1 1 part of glycol (Knorr, Ber. 30, 912). It dissolves NaCl, ZnCl<sub>2</sub>, and many other chlorides; also K<sub>2</sub>CO<sub>3</sub>, KOH, and

Ca(OH),

By action of oxidising agents, e.g. platinum black, it forms glycollic acid. Nitric acid converts it into glycollic acid, glyoxylic acid, and oxalic acid (Debus, Annalen, 110, 316) Heated with solid KOH to 250°, glycol yields potassium oxalate and hydrogen (Wurtz) Heated with large quantity of water at 200°-210° yields aldehyde (Nevole, Bull Soc. chim. [11 ] 25, 289); ZnCl, at 250° decomposes glycol into aldehyde, crotonaldehyde, and water. By electrolysis of an acid (H2SO1) solution of glycol, formic acid, formaldehyde, CO,, traces of a sugar and a non-volatile acid, not glycollic, are (Lob and Pulvermacher, Zeitsch. formed Elektrochem 1910, 16, 1) Chlorine forms a crystallisable halogen compound, m.p. 39°, b.p. 200°; and an only body,  $C_8H_{12}O_4$ , b.p. 240° (Mitscheilich, J. 1863, 485)  $H_2O_2$  in presence of ferrous compounds yields glycollic aldehyde (Fenton and Jackson, Chem. Soc. Trans. 1899, 75). Heated with aldehydes, glycol yields ethylene oxide derivatives, e.g glycol and crianthaldehyde, heated at 125°-130° for some days yield ethylene cenanthylidene oxide

C<sub>6</sub>H<sub>13</sub> CH CCH<sub>2</sub>

b p 180°; similarly, valeraldehyde and propaldehyde yield C<sub>4</sub>H<sub>9</sub> CH COCH<sub>2</sub>, b p 142°;

and C<sub>2</sub>H<sub>5</sub> CH OCH<sub>2</sub>, b p. 105°-107°, respectively (Lochert, Bull. Soc. chim. [ii.] 48, 337, 716). With Na and EtI, yields glycol monoethyl ether, a liquid boiling under 748 mm at 134 8°, and having a density of 0 93535

at 15°/15° (Palomaa, Ber 42, 3873).

Glycol is converted into glycollic acid by the action of Bacterium accti, provided the acid be neutralised (e.g. by CaCO<sub>3</sub>) as fast as formed, otherwise after a short time the glycollic acid formed stops the further action (A. J. Brown, Chem. Soc Trans, 51, 638). NH<sub>4</sub>Cl and glycol at 300°-400° form collidine C<sub>8</sub>II<sub>11</sub>N HCl (Hofmann, Ber. 17, 1905). Phosgene gas at ordinary temperatures converts glycol into glycol carbonate CO(OCH<sub>2</sub>)<sub>2</sub> (with liberation of HCl), a stable compound, soluble in water, alcohol, and ether, and crystallising in colourless needles (Nemirowsky, J pr Chem [2] 28, 439) Heated with strong formic acid for some time (Henninger, Ber 7, 263), or with ovalic acid (Lorin, Bull. Soc chim. [n ] 22, 104), gives glycol diformin  $({}^{\circ}_{2}H_{4}(\mathrm{CH}_{2}\mathrm{O})_{2}, \mathrm{\ b\ p.\ 174}^{\circ}; \mathrm{\ m\ the\ latter\ case}$  formic acid is also produced. With mineral acid or dehydrating agents yields acetaldehyde, ethylidene ethylene ether and chiefly diethyno other  $O < \begin{array}{c} CH_2 CH_2 \\ CH_2 \end{array} = \begin{array}{c} CH_2 \\ CH_2 \end{array} = \begin{array}{c} O \\ O \end{array}$  (Faworsky, Russ Phys. Chem Soc 1906, 38, 741), By action of sodium on glycol, we get sodium ethylenate C<sub>2</sub>H<sub>4</sub>(OH)ONa, a white crystalline deliquescent body, which, on heating to 190° with more sodium yields disodium ethylenate, a deliquescent mass The former,

with C<sub>2</sub>H<sub>2</sub>Br in a sealed tube, yields ethylene, glycol, NaBr, and probably NaCHO<sub>2</sub> (Wurtz).

De Coninck (Bull Acad Roy Belg. 1905, 360) distinguishes between glycol and glycerol as follows Uranyl sulphate is dissolved in a small quantity of water, excess of ethylene glycol added, the whole placed in a flask and exposed to sunlight In 2 hours the hquid is quite green, and in 2½ hours uranous sulphate is precipitated With glycerol no such colouration or precipitation takes place

Glycol forms a number of cyclic esters and cyclic compounds analogous to the metallic saccharates (Bischoff, Ber. 40, 2803; Grun and Bockisch, ibid 41, 3465).

### 2. Diethylene glycol

# HO CH, CH, O CH, CH, OH

b p. 250°, sp gr. 1 132 at 0°; V.D. = 3 78 at 311° (calc. 3.67) (Lourenço, Ann. Chim. Phys. [3] 67, 275), from ethylene oxide and water, or better, ethylene oxide and glycol at 100° (Wurtz); from glycol and ethylene bromide or bromhydrin at 115°-120° (Lourenço); by action of glycol monoacetate upon sodium glycol at 130°-140° (Mohs, Zeitsch. Chem. 1866,

An only liquid, soluble in water, alcohol, and ether; concentrated HI at 100° forms C2H4I2; oxidised by HNO<sub>3</sub> to glycollic acid, oxalic acid, and diglycollic acid C<sub>4</sub>H<sub>6</sub>O<sub>5</sub>. The chlorhydrin C<sub>4</sub>H<sub>9</sub>O<sub>2</sub>Cl, b p. 180°–185° (L.), 190°–200° (W.), is formed by leading HCl into ethylene oxide, or from ethylene oxide and ethylene chlorhydrin at 140° (Wurtz, Ann. Chim. Phys [3] 69, 338). I molecule ethylene chlorhydrin and 2 molecules glycol are heated at 140° saturated with HCl

and again heated at 400° (Lourenço, ibid. [3] 67, 290); a liquid soluble in water.

The bromhydrin, b p. 205°, is prepared from glycol and ethylene bromide at 160° (L)

#### 3. Tri-ethylene glycol

# C2H4(OH) O C2H4 O C2H4OH

b p  $290^{\circ}$ , sp gr 1.138; a liquid miscible with alcohol and water, and oxidised by HNO3 to diglycol-ethylenic acid CaH10O6 (Louienco, Wutrz).

The chlorhydrin  $C_6H_{13}O_3C1$ , bp  $222^{\circ}-232^{\circ}$ ;

is a liquid soluble in water (L)

The bromhydrin C6H13O3Br; bp. (with decomposition) 250° (L)

4 Tetra-ethylene glycol C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>; b p. 230°,

at 25 mm (L. ; W ) The chlorhydrin  $C_8H_{17}O_4CI$  , b p. 262°–272° , is soluble in water (L)

5. Penta-ethylene glycol  $C_{10}H_{22}O_6$ ; b p 281° at 25 mm.; a viscid liquid soluble in water, alcohol, and ether (L).

6 Hexa-ethylene glycol  $C_{12}H_{26}O_7$ , b p 325° at 25 mm; a very viscid liquid (L).

Ethylene bromide C2H4Br2 or CH2Br CH2Br, m p 9 975° (Moles, Anal Fis Quim 1912, 10, 131), 10 0.2° (Biron, Zeitsch physikal Chem 1913, 81, 590), b p 131 6° (R), 129 5° at 745 mm (Anschutz, Annalen, 221, 137), 130 39° (corr.) (Thorpe and Rodger); sp gr 2 1804 at 20° (Bron, 1c), 2 2132 at 0°/4°, 2 1901 at 10 9°/4° (Thorpe Chem Soc Trans 37, 177); by action of C<sub>2</sub>H<sub>4</sub> and Br (Balard, Ann Chim Phys. [2] 32, 375); formed together with ethyldene bromide CH<sub>4</sub>CHBr, by heating CL H Br, with Br to 180° (Chandlaron, Appellar C<sub>2</sub>H<sub>5</sub>Br with Br to 180° (Tawildarow, Annalen, 176, 14) From C<sub>2</sub>H<sub>5</sub>Cl, Br, and iron wire at 176, 14) From C<sub>1</sub>H<sub>5</sub>Cl, Br, and iron wire at 100° (V. Meyer and Petienko, Ber 25, 3307), also from C<sub>2</sub>H<sub>2</sub>Br, Br, and AlBr<sub>3</sub> at 65°-70° (Mouneyrat, Bull. Soc chim. [m] 19, 497)

Preparation.—1 Ethylene, purified by passing through H.SO4 and NaOH solution, is led into bromine covered with water, contained in a wide tube, until the colour of the Br disappears; the product is washed with NaOH and water, dried by CaCl, and distilled (Erlenmeyer and Bunte, Annalen, 168, 64; Erlenmeyer, ibid 192, 244; Prideaux, Chem News, 1916, 113, 277) 2 By heating together at 150° in glacial acetic acid glycol diacetate and HBr, a quantitative yield of C2H4Br, is obtained (Perkin and Simonsen,

Chem. Soc. Trans. 1905, 856).

Is a colourless mobile liquid with a pleasant smell and sweet taste; VD 6.485 (Regnault). When heated with water to 150°-160°, it yields, first glycol, and, on longer heating, HBr and aldehyde (Carius, Annalen, 131, 173) Zn acts upon an alcoholic solution of C2H4Br2 with formation of C<sub>2</sub>H<sub>4</sub> and ZnBr<sub>2</sub> (Gladstone, Ber. 7, 364), by heating with KI and water, ethane is produced. Alcoholic KOH yields C,H<sub>3</sub>Br, and in excess C<sub>2</sub>H<sub>2</sub>. Heated with AlBr<sub>3</sub> at 100°-110°, yields C,H, and C<sub>2</sub>H<sub>2</sub>Br<sub>4</sub>; with Cl and AlCl<sub>3</sub>, yields C<sub>2</sub>Cl<sub>6</sub> (Mouneyrat)

Aqueous KOH produces, after long contact, C<sub>2</sub>H<sub>3</sub>Br (Stempnovsky, Annalen, 192, 240) By heating with strong NaOH and KOH solution, glycol and C<sub>2</sub>H<sub>3</sub>Br are produced. If C<sub>2</sub>H<sub>3</sub>Br be heated with excess of H2O and PbO to 220° aldehyde is formed (due to decomposition of C<sub>2</sub>H<sub>4</sub>(OH)<sub>2</sub>, first formed, into C<sub>2</sub>H<sub>4</sub>O and H<sub>2</sub>O) (Eltekow, Ber. 6, 558, and Nevole, *ibid* 9, 447)

Heated with strong HI solution, yields C2H4I2 (Sororkin, Zeitsch. Chem 1870, 519) Heated with Ag<sub>2</sub>O and water, aldehyde is produced, and when heated with Ag<sub>2</sub>CO<sub>3</sub> and water, yields glycol (Beilstein and Wiegand, Bei 15, 1368). Treated with Ag<sub>2</sub>SO<sub>4</sub> and benzene, yields dibromethyl sulphate (C<sub>2</sub>H<sub>4</sub>Br)<sub>2</sub>SO<sub>4</sub>; with Ag SO<sub>4</sub> and water, bromethyl sulphune acid (C<sub>2</sub>H<sub>4</sub>Br)HSO<sub>4</sub> (B and W)

Bromethylene bromide ('H\_Br ('HBr\_, bp. 186 5° (Wurtz, Annalen, 104, 243), 187°-188° at 721 mm (Denzel, ibid. 195, 202), 191° (Tawiklarow, ibid. 176, 22), 73° at 11 5 mm, 83° at 18 mm, 187°-188° at 751 5 mm (Anschutz, ibid. 221, 138), spgr 2 624 at 16° (T), 2 620 at 23° (W), 2 6189 at 17 5°/4°, 2 6107

at 21 5°/4° (A ).

Prepared from monobromethylene and Br (Wuttz), from C<sub>2</sub>H<sub>3</sub>I and Br (Simpson, J. 1857, 461), from C<sub>2</sub>H<sub>3</sub>Br and Br at 170° (Hofmann, J. 1860, 364), or from C2H4B1, and Br at 200° (Tawildarow), decomposed by KC2H3O2 at 150° or by H<sub>2</sub>O and PbO into dibromethylene and HBr C<sub>2</sub>H<sub>5</sub>NaO yields ethylidene dibromide (Gray, Chem Soc Trans 1897, 1021) Alcoholic KOH gives C<sub>2</sub>H<sub>2</sub>Br<sub>2</sub>, C<sub>2</sub>H<sub>3</sub>, and bromacetyleno C<sub>2</sub>HBr When heated with excess of absolute alcohol, C<sub>2</sub>H<sub>5</sub>Br and probably C<sub>2</sub>H<sub>3</sub>BrO bromaldehyde, are formed (Glockner, Annalen, Suppl. 7, 110)

The isomeric body CH<sub>3</sub>CBr<sub>3</sub> is not known. Dibromethylene dibromide ('.H.Br. isomerides are known

1 Acetylidine tetrabromide; CH,Br CBr, bols with decomposition at 200°; bp 103 5° at 13 5 mm; sp gi. 2 9292 at 17 5°/4°, 2 9216 at 21 5°/4° (Anschutz, Annalen, 221, 140) Formed from as-dibromethylene CH<sub>2</sub> CBr, and bromine (Lennox, *ibid* 122, 124; Chem Soc Trans 1862, 206, Reboul, Annalen, 124, 270) From C<sub>2</sub>H<sub>5</sub>Br and Br at 180° a compound C<sub>2</sub>H<sub>2</sub>B<sub>14</sub> is produced, which boils with decomposition at 208°-211° (Tawildarow) By heating a mixture of pyrotaitanc acid and biomine with water at 150° (Bourgoin, Ann Chim. Phys. [5] 12, 427)

2 Acctylene tetrabromide CHBr, CHB12, b p 137°-137 2° at 36 mm, 114° at 12 mm, at 15 mm 124°-126°, at 54 mm 151°, sp gr. 2 9710 at 17·5°/4°, 2 9629 at 21 5°/4° (Anschütz, Ber. 12, 2974, and Annalen, 221, 139); decomposes at 190° into HBr, Br, and C, HBr,

By leading C.H. into bromine covered with water (Reboul, Annalen, 124, 269, Berthelot, Bull Soc chim [11] 5, 97); the crude C<sub>2</sub>H<sub>2</sub>Bi<sub>4</sub> is then reduced by means of alcohol and Zn dust to C<sub>2</sub>H<sub>2</sub>Br<sub>2</sub>, when the admixed C<sub>2</sub>H<sub>3</sub>Br Br<sub>2</sub> is converted into C2H3Br, the C2H2Br2 is then purified by distillation and treated with bromine (Anschutz, Annalen, 221, 139), by action of Br and AlBr<sub>3</sub> at 105°-110° upon C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> (Mouneyrat, Bull Soc chim [III] 19, 498) A liquid smelling of camphor and chloroform, A liquid smelling of campnor and chlorotorm, decomposed by heating or by action of Na, by heating with alcoholic KOH, C<sub>2</sub>H<sub>2</sub> and C<sub>2</sub>HBr are produced—Cl produces C<sub>2</sub>Cl<sub>4</sub>Br, (Bourgoin, Bull Soc chim [n] 23, 4); when heated with Br and water to 180°, yields C<sub>2</sub>Br<sub>4</sub> and C<sub>2</sub>Br<sub>6</sub> (Anschutz); with C<sub>4</sub>H<sub>6</sub> and AlCl<sub>3</sub>, produces anthracene; heated with annline, dimethylanline, or alcoholic NH<sub>3</sub>, yields CHBr. CBr<sub>2</sub>.

Pentabromethane CHBr<sub>2</sub> CBr<sub>3</sub>, m.p. 56°-57°

(Bourgoin), 54° (Denzel, Ber 12, 2208), 48°-50° (Reboul); bolls with decomposition at 210° (at 300 mm.) (Denzel). From C<sub>2</sub>HBr<sub>3</sub> (Elbs and Newmann, J. pr. Chem [2] 58, 254; Lennox, Annalen, 122, 125) or C<sub>2</sub>HBr (Reboul, *ibid.* 124, 268) and bromme By slow oxidation (by exposure to air) of C<sub>2</sub>HBr<sub>3</sub> (Demole, Bull. Soc chim. [ii] 34, 204); from C<sub>2</sub>H<sub>2</sub>Br<sub>4</sub> and Br at 165° (Bourgoin, *ibid.* 23, 173). Forms prismatic needles soluble in alcohol and other.

Partmether (\* Br. \* (APRON WEIRROMER\*)

Perbromethane C<sub>2</sub>Br<sub>6</sub>v Carbon Tribromide. Ethylene bromoiodide; mp 28°; bp 163°; sp gr 2 516 at 29°. From C<sub>2</sub>H<sub>3</sub>Br and cone H1 at 100° (Reboul, Annalen, 155, 213) From C<sub>2</sub>H<sub>4</sub> and IBr (Simpson, J 1874, 326) Needles, with clocker KOH and Large 1975. with alcoholic KOH yields, in the cold, C<sub>2</sub>H<sub>3</sub>I and IIBr, and on warming HBr, HI, and C<sub>2</sub>H<sub>2</sub>. Forms iodoform with alcoholic CH<sub>2</sub>COOK at 100°

Ethylene bromhydrin C<sub>2</sub>H<sub>4</sub>Br·OH, Glycol bromhydrin, b p. 150°-152°, at 18 mm 63°-64° (Henry, J. 1889, 1321), 155° (Lourenço, Ann Chim Phys [3] 67, 287); sp gr 1·66 at 8° (H); sp gr 1·7195 at 18 6° (H.) From glycol and C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> at 115°-120° (Lourenço); from glycol and HBr at 100°, and from CH<sub>2</sub>I CH<sub>2</sub>OH and browner (Hapry); also from 3 mate shoots bromme (Henry); also from 3 mols. glycol and 1 mol PBr<sub>5</sub> (Demole, Ber 9, 48), from cthylene and HBiO (60 p.c.) (Mohiewsky, J Russ Phys. Chem Soc 30, 900); with zincolar the state of the dust and alcohol is reduced to C2H4, together with ethylene dibromide by the action of ethylene on bromine water (Read and Williams,

Chem. Soc Trans. 1917, 111, 240)

Ethylene dichloride, Elwyl chloride, Dutch liquid CH2ClCH2Cl; bp 84 9° (Pierre, J 1847-48, 63), 83 5° (corr ) (Thorpe, Chem Soc. Trans 37, 182), 84 5°-85° at 750°9 mm (Bruhl, Annalen, 203, 10), 83 3° at 749 mm (R Schiff, bbid. 220, 96), 84 1° at 760 mm (Stidel, Ber. 15, 2563); sp gr. 1 2808 at 0°/4° (Thorpe), 1 2803 at 0° (Pierre), 1 2562 at 20° (Haagen, Bull Soc. chum. 10, 355), 1 2566 at 9.8°44° Bull Soc. chim. 110, 355), 1 2656 at 98 /4°, 1 1576 at 83 3°/4° (R. Schiff), 1 2606 at 14 4°, 1.204 at 62 5° (Landolt and John), bp 83 93° (corr ) (Thorpe and Rodger) refractive index; at  $t^{\circ}$   $\mu_{A} = 1 \ 441466 - 0 \ 000446t$ , mol refract = 34 06 (Kanonnikow); capill constant at b p  $a^2 = 4$  198 (R. Schiff, Annalen, 223, 72) By action of  $C_2H_4$  upon Cl (Deiman, Trootswyk, Bondt, and Lauwerenburgh, Crell's Ann 1795, 2000) 2, 200), by leading C<sub>2</sub>H<sub>4</sub> into SbCl<sub>5</sub> or into molten CuCl<sub>2</sub> (Wohler, Pogg Ann. 13, 297), from C<sub>2</sub>H<sub>5</sub>Cl and SbCl<sub>5</sub> at 100° (V Meyer and Muller, J pr Chem [n | 46, 173); also by action of glycol upon PCl<sub>3</sub> (Wutz) or upon HCl at 100° (Schorlemmer, Chem. Soc Trans. 1881, 1); from ethylene diamine and NOCI in carefully cooled m-xylene solution (Solonina, J Russ Phys Chem. Soc 30, 606) Prepared best accord-ing to Limpricht's method (Annalen, 94, 245) Ethylene is passed into a gently heated mixture of 2 parts manganese dioxide, 3 parts common salt, 4 parts water, and 5 parts sulphuric acid, and when the mass turns yellow it is distilled It can be prepared in good yield from ethylene and chlorine if the temperature is kept at 0°; at higher temperatures considerable quantities of trichloroethane are formed It forms an

(Liebig, Annalen, 14, 37). Heated with AlCl<sub>3</sub> to 80°-85° is decomposed into HCl and C<sub>2</sub>H<sub>2</sub> (Mouneyrat, Bull. Soc. chim. [ii.] 19, 446). With NH<sub>3</sub> it forms bases, N<sub>2</sub>H<sub>4</sub>C<sub>2</sub>H<sub>4</sub>, N<sub>2</sub>H<sub>2</sub>(C<sub>2</sub>H<sub>4</sub>)<sub>2</sub>, &c. With ethylamine, it yields collidize (trimethylpyruline C<sub>5</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>N), C<sub>2</sub>H<sub>5</sub>Cl, and also NH<sub>3</sub> and N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub> (Hofmann, Ber 17, 1907).

Chlorethylene CH<sub>2</sub>: CHCl, Vinyl chloride; at ordinary temp a gas; bp -18° to -15°; by action of alcoholic KOH upon ethylene chloride (Regnault, Annalen, 14, 28), or ethylidene chloride (Wurtz and Frapolli, ibid. 108, 224). To prepare it the ethylere dichloride and alcoholic potash are left in contact for 3 or 4 days in the cold, and the mixture is then warmed Combines with Cl to form C2H3Cl3 Polymerises in sunlight, and forms a solid insoluble mass of sp gr. I 406 (Baumann, Annalen, 163, 317)

(Regnault, J pr Chem [n ] 18, 80) Also by action of Zn shavings in alcoholic solution upon the trichlorethyl ester of acetic acia (Jocitsch and Faworsky, J Russ Phys Chem Soc 30, 998) Polymerises easily into a solid amorphous msoluble body

2 Sym-Dichlorethylene (Acetylene dichloride) CHCl. CHCl, bp 55°, by distillation of the compound of C<sub>2</sub>H<sub>2</sub> with SbCl<sub>5</sub> (Berthelot and Jungfleisch, Annalen, Suppl. 7, 253); also by action of Zn upon an alcoholic solution of CHClBr CHClBr, or by leading C<sub>2</sub>H<sub>2</sub> into aqueous solution of ICl (Sabanejeff, Annalen, 216, 263).

216, 262)

Trichlorethylene CHCl CCl<sub>2</sub>; bp 87 4°–
87 55°; sp gr. 1 4904 at 4°/4°, 1 4598 at 25 5°/25 5° (Veley, Proc Roy Soc. 1910, B 82, 217), 85°8′–86 0°/741 6 mm, m.p. —83°, D<sub>40</sub> 1·4649; D<sub>45</sub> 1 4695 (Erdmann, J pr. Chem. 1912, 11 85, 78) From isomeric compounds C<sub>2</sub>H<sub>2</sub>Cl<sub>4</sub>, and alcoholic KOH (Berthelot and Jungfleisch), also by action of nascent H (Zn and H<sub>2</sub>SO<sub>4</sub>) upon C<sub>2</sub>Cl<sub>8</sub> (Fischer, J. 1864, 481): from anhydrous chloral and P<sub>2</sub>S<sub>5</sub> at 160°-170° (Paternò and Oglaloro, Ber 7, 81); combines with Cl to form C<sub>2</sub>HCl<sub>5</sub> By action of ammonia in aqueous solution on tetrachloroethane (Comp. prod. chem d'Alais; Eng. Pat, 132755)

For observations on its autoxidation, see

Erdmann (l c)

Tetra-chlorethylene C<sub>2</sub>Cl<sub>4</sub> v Carbon dichloride, art Carbon Cf Weiser and Wightman, J. Phys Chem. 1919, 23, 415.

Hexachlorethane C2Cl6 v Carbon trichlor-

ide, ait Carbon.

Ethylene chlorhydrin HO CH2 CH2Cl Chlorethyle alcohol, allycol chlorhydrin; b p. 128° (Wurtz, Annalen, 110, 125), 132°, 51°-52° at 22 mm.; sp gr. 1 2005 at 18 6° (Henry, J. 1889, 1321), 1 24 at 8°. From glycol and HCl (gas) at 100° (W.) If excess of HCl be used, C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> is produced (Schorlemmer, Chem. Soc. Trans. 39, 143); from glycol and Cl<sub>2</sub>S<sub>2</sub> (Carius, Annalen, 124, 257); from C<sub>2</sub>H<sub>4</sub> and ClOH (Carius, thid. 126, 197. Butlerow, thid. ClOH (Carius, ibid. 126, 197; Butlerow, ibid. oily liquid, insoluble in water When heated 144, 40) Dry HCl is led into glycol kept at with alcoholic KOH, it yields KCl and C<sub>2</sub>H<sub>3</sub>Cl, 148°; the C<sub>2</sub>H<sub>4</sub>Cl(OH) distils off (Ladenburg, and with K it forms H, KCl, and C<sub>2</sub>H<sub>3</sub>Cl Ber. 16, 1408) A liquid miscible with water;

yields alcohol with Na amalgam and H<sub>2</sub>O Chourenço, Annalen, 120, 92). Oxidised by CrO<sub>3</sub> to chloracetic acid (Kriwaxin, Zeitsch. Chem. 1871, 265); with KOH, yields ethylene oxide Cf Gomberg, J. Amer Chem. Soc. 1919, 41, 1414.

Ethylene chlorobromide CH<sub>2</sub>Cl CH<sub>2</sub>Br; bp 107-108° (Lesceur, Bull Soc. chim [ii ] 29, 484), sp gr. 1 79 at 0° (L), 1 705 at 11° (Montgolfier and Giraud, ibid 33, 12), 1 689 at 19°. By action of Br upon CH<sub>2</sub>Cl CH<sub>2</sub>I (Henry, Annalen, 156, 14); from C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> and SbCl<sub>5</sub> (Lossner, J. pr. Chem. [2] 13, 421) or  $C_2H_4Br_2$  and  $HgCl_2$  at  $150^\circ-180^\circ$  (M and G.); by action of Br upon  $C_2H_4Cl(OH)$  at  $130^\circ$  (Demole, Ber 9, 556). To prepare it, first chlorine and then ethylene is passed into Br, kept under a mixture of equal vols of strong HCl and water (James, J pr. Chem [2] 26, 380, also Simpson, Proc Roy Soc 27, 118), decomposed by alcoholic KOH, with formation of C<sub>2</sub>H<sub>3</sub>Cl

Ethylene chloroiodide CH,Cl CH,I; m.p. -15 6° (Schneider, Zeitsch physikal Chem 19, -15 6° (Schneider, Zeitsch physikal Chem 19, 157), bp. 140° (Sorokin, Zeitsch Chem 1870, 519), 140 1° (corr) (Thorpe, Chem Soc Trans 31, 189), 137°-138° (Meyer and Wurster, Ber. 6, 964); srgr 2 151 at 0° (S), 2 16439 at 0°, 2·13363° at 15·3°/0° (T) From CH<sub>2</sub>l CH<sub>2</sub>l and ICl; from C<sub>2</sub>H<sub>4</sub> and an aqueous solution of ICl (Simpson, Proc Roy. Soc 12, 278) Decomposed by alcoholic KOH and HI and Cl; H (1), pascent H wedle H(1) HI and (1) H C<sub>2</sub>H<sub>3</sub>Cl; nascent H yields HCl, HI, and C<sub>2</sub>H<sub>4</sub>, moist Ag<sub>2</sub>O at 160° produces glycol (Simpson) Finely divided silver at 160° yields C<sub>2</sub>H<sub>4</sub> and C<sub>2</sub>HCl CH<sub>2</sub>Cl (Friedel and Silva, Bull Soc. chim. [n.] 17, 242) Heated with HI solution, yields C<sub>2</sub>H<sub>4</sub> and C<sub>2</sub>H<sub>4</sub> [Sorokin]

Ethylene evaphydrin HO CH, CH CN: h.p.

Ethylene cyanhydrin HO CH<sub>2</sub> CH<sub>2</sub>CN; b p 220°-222° at 723.5 mm; sp gr 1 0588 at 0°. By long digestion of C<sub>2</sub>H<sub>4</sub>O (ethylene oxide) with anhydrous HCN at 50°-60° (Etlenmeyer, Annalen, 191, 273); miscible with alcohol and water; 100 parts ether dissolve 2.3 parts  $C_2H_4(OH)CN$  at 15°; insoluble in Cs. Forms a chloride with PCl<sub>5</sub>, which boils at 174°–176° under 752 mm, and has sp gr 1 443 at 18 5° (Henry, Bull Acad. Roy Belg. in. 35,

360)

Ethylene cyanide C<sub>2</sub>H<sub>4</sub>(CN)<sub>2</sub>, Succinonitrile; m.p. 54 5° (Nevole and Tscherniak, Bull Soc. chim. [n.] 30, 101), 51°-52° (Henry, ibid. 43, 618); bp 158°-160° at 20 mm (Pinner, Ber 16, 360), 265°-267° (H); sp gr 1 023 at 45° By heating on a water-bath CH<sub>2</sub>Br CH<sub>2</sub>Br with KCN and sleebel (sp. gr. 0.84) (Suprep. Proc. KCN and alcohol (sp gr. 0 84) (Simpson, Proc. Roy. Soc. 10, 574), the liquid is filtered from the KBr, and distilled, when the alcohol is first removed, then the distillation is conducted under diminished pressure; also is produced by electrolysis of potassium cyanacetate (Moore, Ber. 4, 520); is a white amorphous mass soluble in water, chloroform, and alcohol; slightly in carbon disulphide and ether; decomposed when distilled at ordinary pressure Decomposed by heating with KOH or HCl into NH<sub>3</sub> and succinic acid; with AgNO<sub>3</sub> forms plates of C<sub>2</sub>H<sub>4</sub>(CN)<sub>2</sub>,4AgNO<sub>3</sub>, which is soluble in water and alcohol, and explodes when heated (Simpson).

Ethylene diamine NH<sub>2</sub>  $C_2H_4$ NH<sub>2</sub>; m.p. 85° (Kraut, Annalen, 212, 251); b.p. 116·5°; sp.gr. 0·902 at 15° (Kraut). By heating  $C_2H_4CI_2$  or  $C_2H_4Rr_2$  with alcoholic NH<sub>3</sub> to

100° (Cloez, J. 1853, 468); by distillation of 18-diaminopropionic acid (Neuberg and Neimann, Zeitsch. physikal Chem 45, 110), by treatment of C<sub>2</sub>H<sub>4</sub>(CN), with Sn and HCl (Fairley, Annalen, Suppl 3, 372). 42 grams of CH<sub>2</sub>Cl CH<sub>2</sub>Cl and 510 cc aqueous NH<sub>3</sub> are heated for 5 hours to 115°-120° in a sealed tube; the product is then evaporated until it begins to crystallise, and is then precipitated with 3 times its volume of absolute alcohol, the precipitated (',H<sub>4</sub>(NH<sub>2</sub>),,2HCl is then washed with alcohol and the base set free by distillation with powdered NaOH (Kraut, lc., also Hofmann, Ber 4, 666) The by-product from the manufacture of chloral, consisting of CH2Cl CH2Cl, CH3 CHCl2, &c, can be used. A thick oily liquid, with faint ammoniaeal smell, soluble in water, not miscible with other or benzene, the aqueous liquid, which contains  $C_2H_4(NH_2)_2,H_2O$ , cannot be freed from water by BaO, but only by distillation over Na Caro's acid oxidises it to glyoxime (Bamberger and Seligman, Ber 36, 3831) With HClO and HBiO, yields tetrachloro and tetrabromo derivatives, eg Bi N C.H4 NBr. (Chattaway, Chem Soc Trans 1905, 381) Decomposed by HNO, into ethylene oxide and oxalic acid (Hofmann, J 1859, 386), is a strong base and forms salts, combines with ('S, to form crystals of

soluble in water, insoluble in alcohol and ether C2H4(NH2)2,2HCl, long silvery needles insoluble ın alcohol C2H4(NH2)2HCl, PtCl4, yellow leaflets, slightly soluble in water (Greiss and Martius, Annalen, 120, 327); C<sub>2</sub>H<sub>4</sub>(NH<sub>2</sub>)<sub>2</sub>(HCNS)<sub>2</sub>, large prisms, very soluble in water, in p 145° (Hofmann, Ber 5, 245) Forms many compounds with metallic salts and a periodido C<sub>2</sub>H<sub>4</sub>(NH<sub>2</sub>)HI I<sub>4</sub>, which melts at 218°.

Diethylene diamine, Preparatine

$$(C_2H_4)_2(NH)_2$$
 or  $C_2H_4 < NH > C_2H_4$ 

m p  $104^{\circ}$  (Berthelot) , b.p  $170^{\circ}$  (Hofmann, J. 1858, 343),  $145^{\circ}$ – $146^{\circ}$  (Berthelot),  $140^{\circ}$ (Rosdalsky) From ethylene chloride and (Rosalsky) From ethylene thiolide and alcoholic NH<sub>3</sub> (Cloez, Hofmann, Natanson, Annalen, 98, 291), from sodium ethylene glycollate and acid derivatives of ethylene diamine by heating at 250°-350° (D R P 67811), by heating the sodium compound of sym-diacyl ethylene diamine with anhydrous glycol or ethylene bromide at 200° (D. R. P. 73354); by reduction of ethylene oxamide with zinc-dust and KOH (D. R. P. 66461) (For other methods of preparation, see D R. P 70055, 70056, 73125, 74628, 79121, 98031, 100232)

Forms a hydrate C<sub>4</sub>H<sub>10</sub>N<sub>2</sub>,6H<sub>2</sub>O, mp 44° (Berthelot) Piperazine is not changed by heating with HCl or H<sub>2</sub>SO<sub>4</sub> at 200°. With carbonyl chloride it forms tricarbonyl piperazine It forms a periodide  $C_4H_{10}N_2$  2H11<sub>4</sub>3H<sub>2</sub>O, m.p 283° The anhydrous periodide melts at 280° (Herz, Ber. 30, 1585). Piperazine reacts with 2 molecules HB<sub>1</sub>O to form an additive compound, which, on treatment with dilute alkalı or piperazine solution, yields n-dibromq-piperazine. This latter can form an additive compound with 2 molecules of HBrO (Chattaway, Chem. Soc. Trans. 1905, 951). Piperazine reacts

with an excess of EtI at temperature of waterbath to form triethyl piperazonium iodide, mp 240°. Colourless needles, insoluble in cold water, ligroin, C<sub>2</sub>Cl<sub>3</sub>, ('<sub>6</sub>H<sub>6</sub>, and CS<sub>2</sub>; soluble in waim alcohol (Van Rijn, Chem Zenti 1898, 1, 727) (v Priperazine)

sym-Diethylene diethyldiamine

 $(C_2H_4)_2(NH\cdot C_2H_5)_2, H_2O$ 

b p 165° (Schmidt and Wichmann, Ber 24, 3247); from (C<sub>2</sub>H<sub>4</sub>)<sub>2</sub>(NH<sub>2</sub>)<sub>2</sub> and C<sub>2</sub>H<sub>5</sub>I, or from C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> and NH<sub>2</sub>C<sub>2</sub>H<sub>5</sub> (Hofmann, J 1859, 389); from piperazine and an excess of KEtSO<sub>1</sub> in aqueous solution (Van Rijn, Chem Zentr 1898, i 727) It combines with salts of Cd, Cr, Zn, and Ni

# as-Diethylene diethyldiamine

 $NH_2 \cdot C_2H_4 \cdot N(C_2H_5)_2$ 

mobile liquid; bp 145°, sp gr. 0827 at 185°/185°; from bromethylphthalimide and diethylamine with subsequent hydrolysis, yields a platinochloride, mp 230°; an airochloride, mp 161°-163°, and a picaute, mp 211° (Ristenpart, Ber 1896, 29. 2526)

Triethylene diamine (C<sub>2</sub>H<sub>4</sub>)<sub>3</sub>N<sub>2</sub>; b p 210°; soluble in water (Hofmann, J 1858, 343; Harries, Annalen, 294, 350)

Diethylene triamine

# NH<sub>2</sub>·C<sub>2</sub>H<sub>4</sub> NH C,H<sub>4</sub>·NH<sub>2</sub>

b p. (with slight decomposition) 208°; formed with other products by action of NH<sub>3</sub> upon C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub>; the double salt with PtCl<sub>4</sub> being less soluble than that of triethylene triamine, it can be separated from that body (Hofmann, J 1861, 514)

A strongly alkaline liquid miscible with water and alcohol. Its salts are soluble in water, less so in alcohol, 2[(C<sub>2</sub>H<sub>4</sub>)<sub>2</sub>N<sub>3</sub>H<sub>5</sub>,3HCl]3PtCl<sub>4</sub> forms golden-yellow needles.

Triethylene triamine

, 
$$C_2H_4 < NH C_2H_4 > NH$$

b p.  $216^{\circ}$  (Hofmann, J. 1861, 514); forms salts;  $[(C_2H_4)_3H_3N_3,3HCl]_2PtCl_4$ , long golden-yellow needles moderately soluble in water; with hydrobromic acid gives  $(C_2H_4)_3N_3H_3,3HBr$  from acid solutions: from weaker acid solution  $(C_2H_4)_3N_3H_3,2HBr$  Tetraethylene triamine  $(C_2H_4)_4N_3H_3$ ; by

Tetraethylene triamine  $(C_2H_4)_4N_3H_3$ ; by action of alcoholic NH<sub>3</sub> upon  $C_2H_4Br_3$  (Hofmann, Ber 3, 762); forms 3 series of salts with 1, 2, or 3 molecules of monobasic acid; the hydrobromides are amorphous bodies, almost insoluble in water, alcohol, and ether.

Triethylene tetramine

m p.  $+12^{\circ}$ , b p  $266^{\circ}$ - $267^{\circ}$ ; sp gr. 0 9817 at 15°; a strongly alkaline syrup from  $C_2H_4(1_2)$  and  $C_2H_4(NH_2)_2$ , or from  $C_2H_4Br_2$  and alcoholic  $NH_3$  (Hofmann, J. 1861, 519); yields

a yellow amorphous powder almost insoluble in water

Ethylene diformin C<sub>2</sub>H<sub>4</sub>(CHO<sub>2</sub>)<sub>2</sub>; bp 174°; by heating glycol with pure formic acid (Heninger, Ber. 7, 263); also by heating oxalic acid with glycol (Lorin, Bull. Soc. chim [ii.] 22, 104) Decomposed when heated at 200° in sealed tube into CO<sub>2</sub>, CO, H<sub>2</sub>O, and C<sub>2</sub>H<sub>4</sub>.

Ethylene ferrous chloride C<sub>2</sub>H<sub>4</sub>FeCl<sub>2,2</sub>H<sub>2</sub>O; by heating FeCl<sub>3</sub> and other to 150° (Kachler, Ber. 2, 510); forms prisms fairly soluble in water. The corresponding compound

is prepared by leading C<sub>2</sub>H<sub>4</sub> into a strong aqueous solution of FeBr<sub>2</sub> in sunlight (Chojnacki, Zeitsch. Chem. 1870, 420); forms light-green deliquescent crystals

Ethylene iodide CH<sub>2</sub>I CH<sub>2</sub>I; mp.  $81^{\circ}-82^{\circ}$  (Anonstein and Kramps, Ber. 13, 489); sp gr. 207; from C<sub>2</sub>H<sub>4</sub> and I in sunlight (Faraday, Gm. 4, 682), also in the shade, especially if warmed (Regnault, Annalen, 15, 67); a mixture of I and alcohol (absolute) is saturated with C<sub>2</sub>H<sub>4</sub> with repeated agitation (Semenow, J. 1864, 483)

Crystalline plates or prism3 decomposed by heating; when warmed with alcohol to 70°, IC<sub>2</sub>H<sub>4</sub> OC<sub>2</sub>H<sub>5</sub> is produced, HgCl<sub>2</sub> gives in the cold C<sub>2</sub>H<sub>4</sub>ICl, and on heating C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> (Maumené, J. 1860, 345); alcoholic KOH converts it into C<sub>2</sub>H<sub>3</sub>I (Regnault, Annalen, 15, 63).

Ethylene iodohydrin CH<sub>2</sub>I CH<sub>2</sub>OH; from C<sub>2</sub>H<sub>4</sub>(OH)<sub>2</sub> and gaseous HI (Simpson, *ibid*. 11., 121); also by heating C<sub>2</sub>H<sub>4</sub>(OH)Cl with KI (Butlerow and Ossokin, *ibid*. 144, 42).

An oil, b p. 85°/25 mm.; sp.gr. 2:1649 at

18 6° (Henry, Ber 24 Ref 75).

Ethylene nitrate NO<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>NO<sub>3</sub>; b p. 114°-116° with explosion (Demjanow, Chem Zentr 1899, 1 1064); sp gr. 1 5099 at 4°, 1 4960 at 15°, 1 4860 at 25° (Perkin). Is formed when glycol is treated with HNO<sub>3</sub> and H<sub>2</sub>SO<sub>4</sub>, also by interaction of C<sub>2</sub>H<sub>4</sub> and N<sub>2</sub>O<sub>5</sub> in the cold (Demjanow); an oily liquid which is inflammable, explodes on percussion, and is poisonous; KOH converts it into alcohol (Henry, Ber. 3, 529, also Champion, Compt. rend. 73, 571) On reduction yields CH<sub>3</sub>CHO, C<sub>2</sub>H<sub>4</sub>(NH<sub>2</sub>)<sub>2</sub>, and H<sub>2</sub>N CH<sub>2</sub>CH<sub>2</sub>OH

Ethylene nitrite

# C<sub>2</sub>H<sub>4</sub>(NO)<sub>2</sub> or NO<sub>2</sub> CH<sub>2</sub>CH<sub>2</sub>·NO

m.p below  $-15^{\circ}$ ; bp.  $96^{\circ}8^{\circ}$ ; sp gr  $1^{\circ}2156$  at  $0^{\circ}$  (Berton) Is prepared by distilling glycerol trinitrite with more than an equivalent amount of glycol Soluble in alcohol to form  $C_2H_5ONO$  (Bertoni, Gazz. chim ital 15, 353), insoluble in and slowly decomposed by water.

Nitroethylene can be prepared in 50 p.c. yield by the dehydration of β-nitroethyl alcohol by phosphoric oxide or sodium hydrogen sulphate; an almost colourless mobile liquid, b.p. 98·5°; D<sup>13</sup> 8·1 073. Vapour violently attacks the eyes and throat Readily undergoes polymerisation; with alkali the change occurs with explosive violence. Combines with bromine in ethereal solution, yielding nitroethylene dibromide, colourless liquid, b.p. 97°/21 mm. With aniline gives N-β-nitroethyl aniline NHPh-CH<sub>2</sub>-CH<sub>2</sub>-NO<sub>2</sub>, colourless leaflets, m.p. 37° (Wieland and Sakellarios, Ber. 1919, 52, B, 898).

Ethylene passed into an ice-cold mixture of nitric acid (D 1 4) and concentrated sulphuric acid containing 20 p.c. of sulphur trioxide yields on steam-distillation and fractionation under diminished pressure ethylene dimitrate, colourless oil, b.p. 105 5°/19 mm. D<sup>17</sup> 1·433, and B-n troethyl nitrate, colourless poisonous oil, b.p.

d20-122°/17 mm. D<sup>18</sup> 1 468. The latter is converted by phosphoric oxide into nitroethylene

β-Nitroethyl alcohol is a colourless odourless liquid, b p. 103°/11 5 mm. D13 3 1 309, miscible with water, alcohol, and ether obtained by the action of silver nitrite on ethylene iodohydiin (Wieland and Sakellarios, Ber. 1920, [B] 201.

CH<sub>2</sub> 0, b p. 125°, sp gr. Ethylene oxide

0.8909 at  $4^{\circ}/4^{\circ}$ ; 0 8824 at 10°/10° (Perkin, Chem. Soc. Trans. 1893, 488). From C<sub>2</sub>H<sub>4</sub>(OH)Cl and KOH (Wurtz, Annalen, 110, 125, Chim. Phys [3] 69, 317), also from  $C_2H_4(OH)Cl$ and C<sub>2</sub>H<sub>4</sub>(OH)ONa (Scheitz, Zeitsch. Chem. 1868, 379), also from  $C_2H_4I_2$  and  $Ag_2O$  at 150°, from  $C_2H_4Br_2$  and  $Ag_2O$  at 250°, or from  $C_2H_4Br_2$  and  $Na_2O$  at 180° (Greene, J. 1877, 522); also by treatment of  $C_2H_4CI(OC_2H_3O)$ with KOH (Demole, Annalen, 173, 125); by dropping chlorethylacetate on to a cooled mixture of powdered KOH and sand (Roithner, Monatsh. 15, 666).

A pleasant-smelling liquid miscible with water; when heated with water in a closed tube, yields glycol, and polyethylene glycol. Has str ng basic properties, although not an alkaline reaction. It combines with acids, and precipitates metallic hydroxides from solutions of Mg, Al, Fe, and Cu salts. With Na amalgam forms C2H6O. Pt black oxidises it to glycollic acid; PCl<sub>5</sub> produces CH<sub>2</sub>Cl CH<sub>2</sub>Cl

(Wurtz, Ann. Chim. Phys. [3] 69, 317, and 355). Equal molecules of  $C_2H_4O$  and Br at  $0^\circ$  form C<sub>2</sub>H<sub>4</sub>(OH)Br, C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> and other bodies (Demole, Ber 9, 47), but 2 molecules C2H4O and 1 molecule Br, on standing, yield large red or small yellow prisms of (C<sub>2</sub>H<sub>4</sub>())<sub>2</sub>Br<sub>2</sub>; m.p. 65°; b p with decomposition at 95°, they are insoluble in water, but soluble in alcohol or contact with Hg separates the Br and leaves dioxyethylene (Wurtz, Ann Chim Phys [3] 69, 321). By long contact with Zn(12 or KHO, a polymeride which melts at 56° and is insoluble in ether, is produced (Wurtz, Bull Soc. chim. [11.] 29, 530).

CH2O CH2 Dioxyethylene, Diethylene dioxide

mp 9°, bp 102°, sp.gr. 1 0482 at 0°. From (C<sub>2</sub>H<sub>4</sub>)<sub>2</sub>O<sub>2</sub>Br<sub>2</sub>, and Hg by contact at ordinary temperature. The mixture is then distilled, and distillate rectified over KOH (Wurtz, Ann. Chim. Phys. [3] 69, 323), has no marked basic properties, but combines slightly with acetic acid on heating to 120°. Lourenço cold. [3] 67, 288) obtained it by heating C<sub>2</sub>H<sub>4</sub>(OH)<sub>2</sub> and C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> to 160°, but the product boiled at 95°. Forms an iodide with 2 atoms of I, m p 84°, which is decomposed by water and air, but soluble in benzene; also a sulphate and a picrate melting respectively at 100° and 66°. It also forms a compound with HgCl<sub>2</sub>, soluble in, but slowly decomposed by, water (Faworsky, J. Russ. Phys. Chem. Soc. 38, 741; Paternò and Spallino, Atti. R. Acad Lincei, [v.] 16, i. 87)

Chlorethylene oxide  $C_2H_3ClO$ ; b.p.  $70^{\circ}-80^{\circ}$ . long contact of acetylene chloriodide CHCl: CHI with 40-50 vols. of water at 200°-220° (Sabanejeff, Annalen, 216, 268); difficultly

soluble in water.

Bromethylene oxide  $C_2H_3B_1$  O, bp  $89^{\circ}-92^{\circ}$ . From biomoglycol bromhydim CHBi, CH2OH and KOH (Demole, Ber 9, 51); soluble in water.

Ethylene phosphides 1. Phosphorus, Organio

COMPOUNDS OF.

Ethylene selenocyanate  $C_2H_4(SeCN)_2$ ; m p. 128° (Proskauer), 138° (Hagelberg, Ber. 23, From CH<sub>2</sub>Br CH<sub>2</sub>Br and 1092). alcoholic KCNSe (Proskauer, ibid 7, 1281) Colourless needles, insoluble in cold water or ether, difficultly soluble in hot water or cold alcohol, dissolves unchanged in strong HNO3, but on heating gives ethylene selemous acid  $C_2H_4(SeO_3H)_2$  With alcoholic alkali, yields

othylene disclende (Hagelberg).

Diethylene disulphide  $S < {\rm CH_2\ CH_2\ CH_2\ S}$ ; m.p.  $111^\circ-112^\circ$ ; b p  $199^\circ-200^\circ$ ; V D = 4 28 (Huselberg). mann, Annalen, 126, 280). From  $C_2H_4B_{12}$  and  $K_2S$  (Crafts, *ibid.* 124, 110; 128, 220); also by heating  $C_2H_4CS_3$  or  $C_2H_4S_2Hg$  with  $C_2H_4Br_2$  at 150° (H).

Crystallises from alcohol in needles or leaflets, from ether in thick prisms which are volatile at ordinary temperatures, combines with Br and certain salts, but not with NH3; oxidised by  $\mathrm{HNO_3}$  to  $(\mathrm{C_2H_4})_2\mathrm{S_2O_2}$  The bromide

$$(C_2H_4)_2S_2Br_4$$

is a lemon-yellow amorphous body, in p 96°, very unstable, the nodide (C<sub>2</sub>H<sub>4</sub>)<sub>2</sub>S<sub>2</sub>I<sub>4</sub>, mp. 132°-133°, forms non-black monochnic needles, insoluble in water The oxide (C2H4)2S2O2 can also be obtained by decomposition of the chloride or bromide; with water, forms crystals which decompose without fusion on heating, and are soluble in water, almost insoluble in alcohol. The peroxide  $(C_2H_4)_2S_2O_4$ , obtained by heating  $(C_2H_4)_2S_2$  with fuming HNO<sub>3</sub> to 150° (Crafts, Annalen, 125, 124), crystallises from strong HNO, in small crystals which are insoluble in

Ethylene ethylidene disulphide is obtained by heating ethylene sulphide almost to dull redness in absence of air and in a current of hydrogen sulphide. After being shaken with aqueous alkalı is obtained as an almost colourless liquid, b p 172-173° (O Matter, D R P 313650)

Ethylene hydroxysulphide C<sub>2</sub>H<sub>4</sub>(OH)SH (Throethylene glycol), from HO CH<sub>2</sub>CH<sub>2</sub>Cl and alcoholic KHS, the KCl is filtered off and the filtrate acidified with dilute HCl and evaporated at 30°-40° (Carius, Annalen, 124, 258). Is a heavy liquid, almost insoluble in water, soluble in alcohol, the salts are fairly soluble in alcohol; the mercury compound Hg(C2H5OS)2 crystallises from alcohol in needles.

Throdiethylene glycol (C2H4OH)2S is formed when the above compound is heated, H2S being evolved. It is a crystalline solid; in p. under 60°, soluble in alcohol, insoluble in water, forms salts and double compounds with metallic salts.

Ethylene mercaptan C<sub>2</sub>H<sub>4</sub>(HS)<sub>2</sub>: b.p. 146°; sp gr. 1-123 at 23 5°. From CH<sub>2</sub>Cl CH<sub>2</sub>Cl (or bromide, Werner, J. 1862, 424) and alcoholic KHS (Lowig and Weidmann, Annalen, 36, A colourless liquid, soluble in alcohol and alkalis, forms insoluble mercaptides with heavy metals;  $Pb(C_2H_4)S_2$  yellow,  $Cu(C_2H_4)S_2$ chreme green. Ethylene dimethyl sulphid. C<sub>2</sub>H<sub>4</sub>(SCH<sub>3</sub>)<sub>2</sub>, b p. 183°, from C<sub>2</sub>H<sub>4</sub>Br<sub>2</sub> and CH<sub>3</sub>SNa (Ewerlof, Ber. 4, 716) Ethylene

diethyl sulphide  $C_2H_4(SC_2H_5)_2$ , b.p.  $210^{\circ}-213^{\circ}$ (E.) gives on oxidation first  $(C_2H_5SO)_2C_2H_4$ , m.p. 170°; and next  $(C_2H_5SO)_2C_2H_4$ , m.p. 136·5° (Beckmann, J. pr. Chem. [2] 17, 469).

Ethylene thiocarbonates.

dithiocarbonateC<sub>2</sub>H<sub>4</sub>COS<sub>2</sub>, 1. Ethylene ethylene oxythrocarbonate; m p. 31°; by action of HNO<sub>3</sub> upon C<sub>2</sub>H<sub>4</sub>CS<sub>3</sub> in the cold (Husemann, Annalen, 126, 269); forms long thin tables, insoluble in water, soluble in alcohol, ether, and

Ethylene trithiocarbonate C2H4CS3; mp 39 5° (Husemann), 33 4° (Konowalow), 39° (Frasetti); sp gr. 1 4768. From  $C_2H_4Br_2$  and Na<sub>2</sub>CS<sub>3</sub> in presence of absolute alcohol (Husemann, Annalen, 123, 83), by action of water upon AlBr, C2H4Br, CS2 (Konowalow); by action of alcoholic KOH upon ethylene xanthate  $\begin{array}{l} \mathrm{C_2H_4(S\ CS\cdot OEt)_2} + \mathrm{KOH} \\ = \mathrm{KS\cdot CO\ OEt} + \mathrm{C_2H_4CS_3} + \mathrm{EtOH} \end{array}$ 

a theoretical yield; it reacts with dil KOH thus:  $C_2H_4CS_3 + EtOH + 3KOH$ 

$$-KSCOOEt+C_2H_4S_2K_2+2H_2O$$

(Frasetti, Ber. 38, 488). Forms golden-yellow, rhombic prisms, slowly soluble in alcohol. easily soluble in ether, very soluble in CS2,

CHCl<sub>3</sub>, and C<sub>6</sub>H<sub>6</sub>.

ββ'-Dichloroethyl sulphide (CH<sub>2</sub>Cl CH<sub>2</sub>)<sub>2</sub>S. so-called 'mustard gas,' was apparently obtained by Guthrie (Chem. Soc. Trans 1860, 12, 116, 1861, 13, 35) and by A Niemann (Ann 1860, 113, 288) by passing ethylene into cooled sulphur dichloride:

$$\cdot 2CH_2 \cdot CH_2 + SCl_2 = (CH_2Cl CH_2)_2S$$

The conditions for its formation by this reaction have been studied by Myers and Stephen (J Soc Chem. Ind 1920, 39, 65, T), and by Gibson and Pope (Chem. Soc Trans 1920, 117, 272) May also be prepared by converting ethylene into ethylene chlorohydrin, thence into thiodiglycol by treatment with sodium sulphide, and thiodiglycol into  $\beta\beta'$ -dichloroethyl sulphide by the action of hydrogen chloride

(1) CH<sub>2</sub> · CH<sub>2</sub>+HClO=CH<sub>2</sub>Cl·CH<sub>2</sub> OH

(2) 2CH<sub>2</sub>Cl CH<sub>2</sub>·OH+Na<sub>2</sub>S

$$=$$
(HO CH<sub>2</sub> CH<sub>2</sub>)<sub>2</sub>S+NaCl

(3)  $(\mathrm{HO}\cdot\mathrm{CH_2}\cdot\mathrm{CH_2})_2\mathrm{S} + 2\mathrm{HCl}$ 

$$=(CH_2CICH_2)_2S+2H_2O$$

(V. Meyer, Ber. 1886, 19, 3260, Clarke, Chem. Soc Trans. 1912, 101, 1583). For the conversion of ethylene chlorohydrin into dichloroethyl sulphide, see Gomberg, J Amer. Chem Soc. 1919, 41, 1414.

 $\beta\beta'$ -Dichloroethyl sulphide is more readily procured by the action of ethylene on sulphur monochloride at about 60°

$$2CH_2: CH_2+S_2Cl_2=(CH_2Cl\cdot CH_2)_2S+S$$

The product has either the constitution

or is a colloidal solution of S in (C2H4Cl)2S. According to Gibson and Pope, Guthrie's compound is a disulphide of the constitution

(Guthrie, lc; Gibson and Pope, l.c., cf. Josephson and 1919, 38, 348, T, et seq.).

technical scale by this reaction were worked out by A. G. Green.

A continuous vacuum still for its manufacture is described by Streeter (J. Ind. Eng. Chem 1919, 11, 292)

 $\beta\beta'$ -Dichloroethyl sulphide, when pure, boils at 120°/34 mm, 122 5°/37 mm., and 217°/760 mm. and readily solidifies in cold weather, forming long, colourless needles or prisms, which melt at 135° to a liquid of sp gr. 15°/4°=1.285. It is almost insoluble in water, and is readily hydrolysed by boiling with an alcoholic solution For other physical properties, see Gibson and Pope; for a number of synthetic reactions, see Davies (Chem. Soc. Trans. 1920, 117, 297). It causes intense inflammation and suppuration when in contact with the skin or when breathed First used on account of its physiological and toxic action as an offensive

agent by the Germans during an attack on Ypres in June, 1917 For a method of quantitative estimation of the substance in air, see J. Amer. Chem. Soc.

1920, 42, 266.

The existence of the so-called chloroethylene dithiodichloride (C<sub>2</sub>H<sub>3</sub>Cl)<sub>2</sub>S<sub>2</sub>Cl<sub>2</sub> (?), said to be obtained by the action of C<sub>2</sub>H<sub>4</sub> upon boiling S<sub>2</sub>Cl<sub>2</sub>, and described as a light yellow liquid with a pleasant smell, is doubtful.

On treating  $\beta\beta'$ -dichloroethyl sulphide with concentrated intric acid it is converted into ββ'-dichloroethyl sulphoxide (CH<sub>2</sub>Cl CH<sub>2</sub>)<sub>2</sub>SO, colourless plates, m.p. 51°, and afterwards the sulphone (CH<sub>2</sub>Cl CH<sub>2</sub>)<sub>2</sub>SO<sub>2</sub>, m.p. 110° (Spica, Gazz. chim. ital. 1919, 49, ii. 299) Treated with bromine the dichloroethyl sulphide yields (CH<sub>2</sub>Cl CH<sub>2</sub>)<sub>2</sub>S,2Br<sub>2</sub>, an orange-coloured substance which rapidly loses bromine, and is converted into a bright-yellow solid

(CH2Cl·CH2)2S,Br2

m.p 43°-44° (Gibson and Pope)
Ethylenethiccyanate C<sub>2</sub>H<sub>4</sub>(SCN)<sub>2</sub>; m p. 90°.
From CH<sub>2</sub>Cl-CH<sub>2</sub>Cl and alcoholic KCNS (Sonnenschem, J. 1855, 609, Buff, Annalen, 100, 229); CH<sub>2</sub>B<sub>1</sub> CH<sub>2</sub>Br may be used with advantage (Glutz, *ibid.* 153, 313); forms rhombic plates slightly soluble in cold water, more so in hot, and in alcohol, decomposed by heat, volatile in steam; its vapour excites tears.

Ethylidene acetamide CH<sub>3</sub>CH(NHC<sub>2</sub>H<sub>3</sub>O)<sub>2</sub>; m p. 169°; formed by acting on acetamide with aldehyde; forms large prisms decomposed by

acids (Tawildarow, Ber. 5, 477).

Ethylidene acetate CH<sub>3</sub>·CH(C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>)<sub>2</sub>; p.p. 169°; sp gr. 1061 at 12°. From aldehyde and  $(C_2H_3O)_2O$  at 180° (Geuther, Annalen, 106, 249), from alcoholic C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>K and CH<sub>3</sub>CHCl·C<sub>2</sub>H<sub>3</sub>O<sub>2</sub> (Schiff, Ber 9, 304); by passing acetylene into a solution of mercuric oxide in glacial acetic acid to which an alkyl sulphate, particularly methylene sulphate, has also been added (Eng Pat. 112766, 1918); or \(\beta\)-naphthalene sulphonic acid dissolved in glacial acetic acid may be used instead (Eng. Pat. 112765); has a smell like that of onions, and is decomposed by water into aldehyde and acetic acid.

Ethylidene azine  $CH_3CH:N\cdot N:CHCH_3$ ; b p.  $95^\circ-96^\circ$  at 760 mm.; sp gr. 0.832 at  $17^\circ$ . Prepared by shaking an ethereal solution of aldehyde with an aqueous solution of hydrazine hydrate. The conditions for its preparation on a At '80° it is unchanged, but at a higher tempera-

ture gives off nearly all its N as gas It is fairly stable to alkalis, but is hydrolysed by acids into aldehyde and hydrazine (Curtius and Linkeisen, J. pr. Chem. 1898, [ii.] 58, 310).

Ethylidene biuret  $NH < {\rm CO \ NH} > {\rm CH \cdot CH_3}$ ,

Trigenic acid, by leading HCNO into well-cooled aldehyde (Liebig and Wohler, Annalen, 59, 296), crystallises in small prisms, slightly soluble in water, nearly insoluble in alcohol, has a slightly When heated, liberates CO2, NH3, acid taste. and traces of collidine C<sub>8</sub>H<sub>11</sub>N (Herzig, Monatsh. 2, 398). Is hardly decomposed by NaBrO. HNO3 decomposes it and liberates cyanuric acid The silver compound AgC4H6N3O2 forms amorphous precipitate, soluble in boiling water.

Ethylidene bromide CH, CHBr, b p. 114°-115° (Tawildarow, *ibid.* 176, 12), 109°-110° at 751 mm. (Anschutz, *ibid.* 195, 202), 110 5° at 760 mm. (Anschutz, *ibid.* 221, 137), 108°-110° (corr.) (Perkin, Chem. Soc. Trans 45, 523), b p. 107°-108° (V. Meyer and Muller, J. pr Chem. [2] 46, 168), sp gr 2 129 at 10° (T). 2 1001 at 17 5°/4°, 2 0891 at 20 5°/4° (Anschutz) 2 0822 at 21 5°/1°, 2 1029 at 15°, 2 0854 at 25° (P) From C2H5Br and Br at 170° (Hofmann, Proc Roy. Soc. 10, 619; Caventou, Compt rend 52, 1330) Is produced by action of PCl<sub>3</sub>Br<sub>2</sub> but not of PBl<sub>5</sub> upon aldehyde (Paterno and Pisati, Ber 5, 289), also by heating CH<sub>2</sub>: CHBr with HBr to 100° (Reboul, Compt. rend. 70, 398). A heavy liquid; alcoholic KHS has no action upon it On heating with water and PbO to 130°, aldehyde and HBr are produced (T) With SbCl<sub>5</sub> gives CH<sub>3</sub> CHCl<sub>2</sub> (Henry, Bull. Soc chim [n ] 42, 262)

Ethylidene bromiodide CH3CHB1I, below -20°; b p 142°-143°, sp g1 250 at 1°, 2 452 at 16°. From C<sub>2</sub>H<sub>3</sub>Br and HI in the cold (Pfaundler, J. 1865, 483; Reboul, Annalen, 155, 212); by shaking CH<sub>3</sub>CHI<sub>2</sub> with IBr solution (Simpson, Bull Soc chim [ii ] 31, 412); decomposed by alcoholic KOH or Ag<sub>2</sub>O into HI and C<sub>2</sub>H<sub>3</sub>Br, Br yields CH<sub>4</sub>CHBr,

Ethylidene chlorobromide CH CHClBr; m p. 16 6° (corr ) (Schneider, Zeitsch. physikal Chem. 19, 157); b p. 81°-84·5° at 750 mm, 82 7° at 760 mm. (Stadel, Ber. 15, 2563); sp gr. 1 667 From C2H, Br and concentrated H(1 at 100° (Reboul, Annalen, 155, 215); from C<sub>2</sub>H<sub>5</sub>Cl and Br in sunlight (Denzel, ibid. 195, 193). By action of Cl upon C<sub>2</sub>H<sub>5</sub>Br (Lescœur, Bull. Soc chim [11] 29, 483) Is decomposed at ordinary temperature by alcoholic KOH on long standing; with Ag<sub>2</sub>O gives aldehyde; and with

AgC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> forms CH<sub>3</sub>CH(C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>)<sub>2</sub> (L.)

Ethylidene chloriodide CH<sub>3</sub>CHCH , b p

117°-119°; sp gr 2 054 at 19° By agitating a

mixture of CH<sub>3</sub>CHI<sub>2</sub> and ICl, formed also from CH3CHCl2 and Al2I6 (Simpson, Bull. Soc chim

[1i.] 31, 411)

Ethylidene or Ethidene chloride CH<sub>3</sub>CH(1<sub>2</sub>, m.p. -101 5° (Schneider, Zeitsch. physikal Chem. 22, 233), b.p. 57·4°-57 6° at 750·9 mm, sp gr. 1·1743 at 20°/4° (Bruhl, Annalen, 203, 11), 1·2124 at 0°, 1·1805 at 22°/0°; respectively. fractive index  $\mu_{\Lambda}$  (at  $24.7^{\circ}$ ) = 1 405724.  $R_{\alpha}$ = 34·1 (Kanonnikow); bp 57°-57 7° at 738 7 mm., sp gr. 1.2013 at 0° (Pribram and Handl, Monatsh 2, 650), b p. 56.7°-56.9° at 749.1 mm., sp gr. 1.1895 at 9.8°/4°, 1.11425 at 56.7°/4° (R. Schiff Annalen, 220, 96), 57.32° (corr.) (Thouge

and Rodger), crit. temp. 254 5° (Pawlewsky, Ber. 16, 2633); H.C.p. = 267 1 cals. (Berthelot and Ogier, Bull. Soc chim. [11] 36, 68). From C<sub>2</sub>H<sub>5</sub>Cl and Cl (Regnault, Annalen, 33, 312); from C<sub>2</sub>H<sub>4</sub>O and PCl<sub>5</sub> (Beilstein). By action from C<sub>2</sub>H<sub>4</sub>O and PCl<sub>5</sub> (Beilstein). By action of HCl upon copper acetylene (Sabanejeff,  $\imath b \imath d$ . 178, 111). Is prepared by passing a mixture of chlorine and  $C_2H_5Cl$  over animal charcoal heated to  $250^\circ{-}400^\circ$  (Damoiseau, Bull. Soc chim. [n ] 27, 113) Is obtained as a by-product in the manufacture of chlora! Is decomposed by alcoholic KOH into HCl and C2H3Cl. Sodium produces at 180°-200° hydrogen, acetylene, cthylene, ethane, and C₄Ĥ₃Cl (Tollens, Annalen, 137, 311) By action of Cl, CH, CCl3 and CH2Cl CHCl2 are produced.

Ethylidene di-150-nitramine is formed by the interaction of  $CO(Et)_2$ , or COEtMe, and NO, in  $C_2H_5ONa$  The lead salt is anhydrous and crystallises in cubes. The dimethyl ether torms rhombic (rystals, and melts at 75°. The 180-nitramine itself is hydrolysed by mmeral acids into acetaldehyde NH<sub>2</sub>OH and oxides of N. With Na amalgam, is reduced to hydrazine (Traube, Annalen, 300, 120).

Ethylidene diethyl ether

CH<sub>3</sub> CH(OC<sub>2</sub>H<sub>5</sub>)<sub>2</sub> (v Acetal)

Ethylidene dimethyl ether CH3CH(OCH3)2, Dimethyl acetal, b.p 64 4° (Dancer, Annalon, 132, 240), 62 7°-63 3° at 757 6 mm (R Schifl, ibid 220, 104), sp gr 0 8787 at 0°, 0 8590 at 14°, 0 8476 at 25° (D), 0 8013 at 62 7°/4° (S); capillarity constant at bp  $a^2 = 4.092$  (Schiff, Annalen, 223, 74), occurs in clude wood spirit (D.). 2 volumes of wood spirit, I volume aldehyde, and I volume of glacial acetic acid are heated to  $100^{\circ}$  (Alsberg, J. 1864, 485), also by oxidation of CH<sub>3</sub>OH and C<sub>2</sub>H<sub>5</sub>OH by MnO<sub>2</sub> and H2SO4 (Wurtz)

Ethylidene methyl propyl ether

 $CH_3$   $CH(OCH_3)(OC_3H_7)$ 

b p $~103^{\circ}\text{--}105^{\circ}$  (Bachmann, Annalen, 218, 46).

Ethylidene ethyl propyl ether

CH<sub>3</sub> CH(OC<sub>2</sub>H<sub>5</sub>)(OC<sub>3</sub>H<sub>7</sub>)

b p 124°-126° (Bachmann, lc).

Ethylidene dipropyl ether CH<sub>3</sub> CH(OC<sub>3</sub>H<sub>2</sub>)<sub>2</sub>, b p 146°-148°, sp gi. 0 825 at 22° (Girard, J. 1880, 695).

Ethylidene methyl isobutyl ether

 $CH_3 CH(OCH_3)(OC_4H_9)$ 

bp 125°-127° (Claus and Tramer, Ber 19, 3005; Bachmann, l.c.).

Ethylidene ethyl isobutyl ether

 $CH_3 \cdot CH(OC_2H_5)(OC_4H_9)$ 

b p  $155^{\circ}$  (Claus and Trainer, lc).

Ethylidene di isobutyl ether CH<sub>3</sub> CH(OC<sub>4</sub>H<sub>9</sub>)<sub>2</sub>, b.p.  $168^{\circ}$ -170°, sp.gr. 0.816 at 22° (Claus and Trainer,  $l\epsilon$ ; Guard,  $l\epsilon$ ).

Ethylidene methyl 180amyl ether

 $CH_3 CH(OCH_3)(OC_5H_{11})$ 

b p 141°-144° (Bachmann, l.c.)

Ethylidene ethyl isoamyl ether  $CH_3 \cdot CH(OC_2H_5)(OC_5H_{11})$ 

b p. 165°-167° (Claus and Trainer, lc.).

Ethylidene di isoamyl ether

 $\mathrm{CH_3CH}(\mathrm{OC_5H_{11}})_2$ 

b p 210.8° (corr.), sp.gr. 0 8347 at 15° (Claus and Trainer, lc).

Ethylidene hydramine CH<sub>3</sub>·CH(OH)NH<sub>2</sub> v. | Aldehyde-ammonia, art. Aldehyde.

Ethylidene imine CH<sub>3</sub>CH: NH; m.p. about 85°; b.p. 123°-124°. If aldehyde ammonia be kept in vacuó over H<sub>2</sub>SO<sub>4</sub> for several days, the imine is formed. It crystallises in brilliant colourless crystals, which have an odour like It is soluble in water, EtOH, acetamide CH<sub>3</sub>COOH, CHCl<sub>3</sub>, C<sub>6</sub>H<sub>6</sub>, and toluene. The vapour density at 260° agrees with the simple formula, but at lower temperatures the V.D is higher. It forms a picrate, and combines with HCN to form ('H<sub>3</sub>·CH(NH<sub>2</sub>)CN. It polymerises to form trimeric ethylidine imine

which forms a trinitroso- derivative, with N<sub>2</sub>O<sub>3</sub> in CHCl<sub>3</sub>; opaque yellow-white needles, melting at 161° (Delépine, Compt. rend. 1897, 125, 951, and 1907, 114, 853).

Ethylidene iodide CH<sub>3</sub>CHI<sub>2</sub>; b p. 177°-179° sp.gr. 284 at 0°. By combination of C2H2 and HI (Berthelot, Annalen, 132, 122, Semenow, Zeitsch. Chem. 1865, 725); from CH<sub>3</sub>CHCl<sub>2</sub> and Al<sub>2</sub>I<sub>6</sub> (Gustavson, J. Russ. Phys. Chem. Soc. 6, 164); from C<sub>2</sub>H<sub>3</sub>Br and HI (Friedel, Ber 7, 823). Decomposed by alcoholic KOH into HI and C.H.I.

Ethylidene malonic acid CH<sub>3</sub>·CH C(COOH)<sub>2</sub>; only known in combination, the ethyl ester CH<sub>3</sub>CH:  $C(COOC_2H_5)_2$ , bp. 220°, 115°-118° at 17 mm.; sp.gr. 1.0435 at 15°. From CH<sub>2</sub>(CO<sub>2</sub>Et)<sub>2</sub> (1 mol.) aldehyde (2 mols), and  $(C_2H_3O)_2O$  ( $(1\frac{1}{2})_2O$ ) and  $((1\frac{1}{2})_2O$ ) and ((Annalen, 218, 157). An ethereal oil smelling of camphor; by long contact with Ba(OH) in the cold yields ethoxy malonic acid C5H8O5, malonic acid C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>, and aldehyde.

Ethylidene oxide CH<sub>3</sub> CHO v. ALDEHYDE.

Ethylidene ethylene oxide

$$CH_3 \cdot CH < \frac{OCH_2}{O\dot{C}H_2}$$

b p. 82° (Verley, Bull. Soc. chim [iii.] 21, 276). 82 5° at 765 8 mm.; sp gr. 1 002 at 0°. aldehyde and excess of glycol at 100° (Wurtz, Compt. rend. 53, 378; Annalen, 120, 328); a hquid with irritating smell, soluble in 1½ vols water, from which CaCl<sub>2</sub> separates it; with acetic acid gives glycol diacetate.

Ethylidene ethoxyacetate CH<sub>3</sub>CH(OEt)OAc; bp. 125°-130°; sp gr 0 941. By heating together acetal and acetic anhydride for 3 hours at 150°. A pleasant-smelling oil, slowly decomposed by cold water into alcohol, aldehyde, and acetic acid, rapidly on boiling.

Ethylidene sulphonic acids.

1. Ethylidene hydroxysulphonic acid

### CH<sub>3</sub>CH(OH)(HSO<sub>3</sub>)

Salts of this acid are formed by action of aldehyde upon alkaline hydrogen sulphites; they are crystalline bodies, soluble in water, and decomposed on heating into water, aldehyde, and sulphites.

2. Ethylidene chlorsulphonic acid

obtained by action of CH3CHCl2 upon neutral alkaline sulphites at 140°; the salts are crystalline, and the acid is fairly stable.

3. Ethylidene disulphonic acid

By oxidation of trithioaldehyde (C<sub>2</sub>H<sub>4</sub>S<sub>3</sub>) or thialdin by KMnO<sub>4</sub> solution (Guareschi, Gazz. chim. ital. 9, 75; Annalen, 222, 302). 10 grams thialdin are treated with 30-35 grams ZnMn<sub>2</sub>O<sub>8</sub> and 300 grams water; after filtering, BaH<sub>2</sub>O<sub>2</sub> is added and excess of Ba precipitated by CO<sub>2</sub>; the Ba salt is then precipitated by alcohol. forms a strongly acid only liquid, very soluble in water and alcohol, and is very stable; forms stable salts Na<sub>2</sub>C<sub>2</sub>H<sub>4</sub>S<sub>2</sub>O<sub>6</sub>,H<sub>2</sub>O, small silky crystals soluble in 6071 parts of alcohol (90 p.c.) at  $24.5^{\circ}$ ;  $K_2C_2H_4S_2O_6, 2H_2O$  forms large transparent crystals soluble in 1.56 parts of water at 17°

ETHYL ACETATE v. Alkyl aretates, art.

# ETHYL-m-AMINOPHENOL

$$\langle \frac{OH}{} \rangle NH \cdot C_2H_5$$

(1) Prepared by adding monoethyl aniline to fuming sulphuric acid, converting the ethyl and and and into the sodium salt, melting this with potassium hydroxide at 220° dissolving the product in water, and acidifying with hydrochloric acid. The filtered solution is neutralised with sodium carbonate, and the mono ethyl-m-amino phenol extracted with benzene (Bad. Anil. und Soda Fabrik, D. R. P. 48151).

# (2) Sodium-m-phenylene oxamato

is mixed with sodium ethyl sulphate, alcohol, and sodium carbonate, and heated in an autoclave fitted with a stirrer to 180°. After distilling off the alcohol, and extracting the mass with water, the residue is boiled with dilute sulphuric acid, and the cooled solution of ethyl-m-phenylenediamine sulphate is mixed with sodium nitrite, warmed, and neutralised with sodium carbonate (idem D R P. 76419).

(3) Aniline-2 · 5-disulphonic acid is neutralised with alkali, and the solution diluted, mixed with sodium ethyl sulphate, and heated in an autoclave to 170°-180°. Sodium carbonate is added to the cooled solution which precipitates ethyl-m-ammophenol (Farberfabrik, vorm F. Bayer & Co, D R. P. 82765); m.p. 62°; b.p 176°/12 mm

### ETHYLANILINE

Is manufactured by heating aniline hydrochloride with alcohol in an autoclave at 180°. The hydrochloride separates out on cooling. Or aniline hydrochloride may be boiled with diethylaniline under a reflux condenser, hydrochloric acid added, and the hydrochloride separated by crystallisation. The base boils at 206°/760 mm.; sp gr. 0.954 at 18°. Yields methyl ethyl aniline on methylation, and benzyl ethyl aniline on benzylation.

ETHYLBUTYLACETIC ACID v. OCTOIC ACIDS. ETHYL-iso-BUTYLACETIC ACID v. OCTOIC ACIDS.

ETHYL BUTYRATE v. BUT'RIC ACID.

MINE

ETHYLKETENE v. KETENES.

ETHYLKETENECARBOXYLATEv. KETENES. ETHYLMALONIC ACID v. GLUTARIC ACID and MALONIC ACID

ETHYLMORPHINE v. OPIUM.

ETHYLPROPYLACETIC ACID v. HEPTOIC ACTOS

#### ETHYL-o-TOLUIDINE

$$\begin{array}{c} \mathrm{CH_3} \\ \end{array}$$

May be prepared by heating a mixture of otoluidine, ethyl alcohol, and hydrochloric acid, or a mixture of o-toluidine, ethyl alcohol, and a small quantity of iodine. Boils at 213°-214°;

sp.gr. 0 9534.

EUBORNYL. Trade name for bornyl-a-bromo-isovalerate, obtained by brominating chloro-isovaleric acid and esterifying borneol with the product. A syrupy aromatic liquid, b.p. 175°-178°. Used as a sedative (v. Syn-THETIC DRUGS).

α-EUCAINÉ. Methyl ester of N-methylbenzoyl-triacetone-alkamine carboxylic acid. First obtained by Merling (Ber Deut pharm Ges. 1897, 173) by acting on acctone with ammonia to form triacetoneamine, treating this with hydrocyanic acid to form the cyanhydrin, of \(\beta\)-eucaine.

ETHYL DIPHENYLAMINE v. DIPHENYLA- | which by hydrolysis yields triacetone-alkaminecarboxylic acid. On benzoylation and methylation it forms a-eucaine. Formerly employed as a substitute for cocaine. Its use is now superseded by  $\beta$ -eucaine, which is less painful and not so toxic

> B-EUCAINE (benzoyl-vinyl-diacetone-alkamine)

$$\begin{array}{c} \text{CH} \\ \text{O\cdot CO C}_{6}\text{H}_{5} \\ \text{CH}_{2} \\ \text{(CH}_{3})_{2}\text{C} \\ \text{NH} \end{array}$$

A local anæsthetic, used chiefly as the lactate or hydrochloride, in 2-3 p.c. solution. Is prepared by condensing acetone with oxalic acid and ammonia to form the hydrogen oxalate of diacetoneamine, condensing this with paraldehyde to produce the cyclic vinyl-diacetoneamine (Harries, Annalen, 1897, 328; 1898, 346), reducing by means of sodium amalgam to a mixture of the cis and trans forms of the trimethyl hydroxy piperidine (D. R. P. 95622), treating with sodium amylate to produce the cis form only of this compound (D. R. P. 95621), and treating with benzoyl chloride (D R P 97672), and finally with hydrochloric or lactic acid to obtain the corresponding salt

EUDOXIN v. SYNTHETIC DRUGS.

**EUCAIRITE.** Selenide of silver and copper (Ag,Cu)2Se, crystallising in the cubic system. It is lead-grey, but usually tarnished and dull; sp gr. 75. It is found massive and granular embedded in calcite in the Skrikerum copper mine in Småland, Sweden, and in the Cacheuta mine in the province of Mendoza, Argentina.

L. J. S. EUCALYPTOL. The volatile oil of the leaves of Eucalyptus globulus, consists mainly of a body C<sub>10</sub>H<sub>18</sub>O, b p. 176°-177°; sp gr. 0 923 at 16°. It is optically mactive, has a camphorlike smell, and is probably identical with cincol and cajeputol (v. OILS, ESSENTIAL).

EUCÂLYPTUS GUM v. Kino.

EUCALYPTUS MACRORHYNCHA. Eucalyplus macrorhyncha (F. v. M.), a fair-sized tree, is the 'red stringy bark' of New South V'ales,

EUCAINE, EUCHININ, EUCODIN, EUCOL, and the ordinary stringy bark tree of Victoria (Smith, Chem. Soc. Trans. 1898, 73, 697).

The leaves yield under favourable conditions a very large amount (10 pc.) of a crystalline glucoside termed by Smith myrticolorin, which can be isolated in the crude condition by mere extraction with boiling water. The solution on cooling became semi-solid owing to the separa-tion of crystals, and these can be purified by extraction with ether to remove chlorophyll and crystallisation first from alcohol and subsequently from water. It formed pale-yellow needles, gave on hydrolysis quercetin and glucose, and at first appeared to be a new glucoside of quercetin. Though very similar to rutin its identity with this glucoside was unsuspected in that rutin by hydrolysis was presumed at that time to give quercetin and 2 molecules of rhamnose (Schunck, *ibid.* 1888, 53, 264). Schmidt in 1908 (Arch. Pharm 246, 214), however, pointed out that rutin in this manner yields not only rhamnose but glucose, and the probability that myrticolorin—as also viola quercitrin and osyritrin—were identical with rutin was subsequently confirmed by Perkin (*ibid.* 1910, 97, 1776)

A. G. P

EUCALYPTUS OIL v OILS, ESSENTIAL EUCALPYTUS RESIN v. RESINS

EUCLASE. Basic silicate of aluminium and glueinum (GlO, 17·3 pc.) HGlAlSiO<sub>5</sub>, crystallising in the monoclinic system. The welldeveloped crystals have a very perfect cleavage parallel to the plane of symmetry, hence the name, from εθ, well, and κλάσις, fracture They are colourless, yellow, green, or blue; sp gr 3.1 (higher than that of beryl), H 71/2 The mineral is not attacked by acids, and the water is expelled only at a high temperature It is found as single isolated crystals in gold washings at Ouro Preto in Brazil, and on the River Sanarka in the southern Urals. Minute crystals have also been found in mica-schist in the Grossglockner district of the Austrian Alps, and in granite in the Fichtelgebirge, Bavaria The mineral is occasionally cut as a gem-stone, but it has the drawback that it readily splits along the perfect cleavage. L J. S.

EUCODAL. Trade name for the hydrochloride of dihydrohydroxycodemone Used as a nageotic

**EUCODEINE.** Syn. for codeine methyl bromide.

EUCOL. Trade name for guarcol acetate

EUCOLITE v EUDIALYTE

EUCRYPTITE. Orthosilicate of aluminium and lithium (Li<sub>2</sub>O, 11 9 pc) Li<sub>2</sub>AlSiO<sub>4</sub>, crystallising in the hexagonal system and belonging to the nephelite group of minerals. It occurs intimately intergrown with albite as an alteration product of spodumene at Branchville, Connecticut. An artificial orthorhombic form of Li<sub>2</sub>AlSiO<sub>4</sub>, dimorphous with eneryptite, has been called pseudo- or β-eucryptite (F. M. Jager and A. Simek, 1914).

L. J. S.

EUCUPIN. Trade name for iso-amylhydro-

cuprein.

EUDESMIN C22H26O6 occurs in the kinos of eucalyptus species, the oils of which contain cincole and pinene, but not phellandiene It may be obtained by extracting a concentrated aqueous solution of the kino with ether, purifying by treatment with chloroform, and crystallising from methyl alcohol Colourless prismatic needles, m p 107°,  $[a]_{\rm D}^{21}$  -64 4° (m chloroform). Contains 4 methoxy- groups, and is unacted upon by hydroxyl or ketone reagents. Red solution with sulphuric acid. Nitric acid converts it into dinitroeudesmin, colourless needles, m p. 214°. Boiling cone. nitric acid forms 4:5-dinitroveratrole (2 mols). Chlorine in acetic acid solution gives dichloroeudesmin; colourless rectangular plates, mp. 163°. Dibromoeudesmin, prepared in like manner, colourless needles, m.p. 172°,  $[a]_{\rm p}^{22}+69.4$ ° (in chloro form); this, on oxidation with permanganate, gives 6-bromoveratric acid, m.p. 184°. Diiodocudesmin, needles, m p. 175° (Robinson and S.nith, J. Roy. Soc. New South Wales, 1914, 48, 449)

**EUDIALYTE.** Silicate and zirconate (ZrO<sub>2</sub>,

14-16 p.c.) of sodium, calcium, and ferrous iron, occurring as large well-developed rhombohedral crystals. These are brownish-red or rose-red in colour; sp.gr. 29-3·1. The mineral occurs in nepheline- and sodalite syemites in the Juliane-haab district in south Greenland, at Magnet Cove in Arkansas, on Sedlovatoi Island in the White Sea, and on the Kola peninsula in Russian Lapland The name cudialyte, from εδ, well, and διαλύειν, to dissolve, refers to the fact that the mineral is readily decomposed by acids.

Eucolite, a closely related variety, differs in being optically negative, instead of positive This is from the nepheline-syemites of the Langesundsfjord in Norway L J S

EUDRENINE. A solution containing cucaine

and adrenaline.

EUFORMOL. Dextrm formaldehyde

**EUGENOL** (HO)(CH<sub>3</sub>O)C<sub>6</sub>H<sub>3</sub>·CH<sub>2</sub>·CH : CH<sub>2</sub> (C<sub>3</sub>H<sub>5</sub>: CH<sub>3</sub>O: HO=1 3.4) occurs in oil of cloves from the leaves of Eugenia caryophyllala (Thunb) (Bonastre and Ettling, Annalen, 9, 68), in the oil from the leaves of the Ceylon cinnamon tree (Stenhouse, Annalen, 95, 103, Schimmel & Co, Chem. Zentr 1902, 11 1486), from Cinnamomum pedatinervium (Meissn.) (Goulding, Chem Soc. Trans 1903, 1101), C Tamala (Nees et Eberm.) (Schimmel & Co., Chem. Zentr. 1910, 1 1720), C cassia (Plume), and C zeylanicum (Nees) (Duyk, Chem. Zentr. 1896, ii. 358, Weber, Arch Pharm. 230, 232); from the fruit of Pimenta officinalis (Lindl.) (Bonastre, lc., Oeser, Annalen, 131, 277); from the sap of Acacia Farnesiana (Willd) (Walbaum, J. pr Chem. 68, [2] 235), in the oil from Ocimum Basilicum (Linn) (Schimmel & Co, Chem Zentr. 1900, 1 906); in calamus oil (Thoms and Beckstroem, Ber. 1902, 3187); in Ylang Ylang oil (Schimmel & Co, Chem. Zentr 1903, 1 1087; n. 272); in olive oil from Pugli (Canzoneri, Gazz chim. ital. 1897, 27, 11. 1); in oil of nutmeg (Power and Salway, Chem. Soc. Trans. 1907, 2041); in sassafras leaves and bark (Pomeranz, Monatsh. 11, 101; Power and Kleber, Chem. Zentr. 1897, u. 42); in the oil from the leaves of the bay tree (Mittmann, Arch. Pharm. 27, [3] 539); in oil from Maltese oranges (Gildemeister and Stephan, Arch. Pharm 235, 583), and in other ethereal The methyl ether of eugenol

# $(CH_3O)_2C_6H_3CH_2\cdot CH:CH_2$

occurs in the oil of Asarum canadense (Linn.) (Power and Lees, Chem Soc. Trans. 1902, 67), of A. arifolium (Michx.) (Miller, Arch. Pharm. 240, 371); in oil from Paracoto bark (Jobst and Hosse, Annalen, 199, 75; Wallach and Rheindorff, ibid. 271, 300), in Ylang Ylang oil; in oil from the bark of Cinnamomum pedatinervium, in oil from Maltese oranges (v. supra); and in oil of citron (Schimmel & Co, Chem Zentr 1899, ii. 879).

Oil of cloves, which contains eugenol and a terpene as the chief constituents, is treated with potash; the potassium salt of eugenol separates, is pressed out, washed with alcohol, and the eugenol set free by the addition of a mineral acid (Wasserman, Annalen, 179, 369). It is obtained by a similar process from oil of cinnamon leaves or from that of Primenta officinalis (Lindl.). Another method consists in separating eugenol from oil of cloves by means of the compound it forms with sodium accetate (Chem.

EUGENOL 687

Werke Byk D. R. P. 100418; Chem. Zentr. 1899, i. 764). In the former process, a residue is left after treating the oil of cloves with potash, and by distilling this under reduced pressure, acetyl eugenol, m p. 29°, is obtained (Erdmann, J. pr Chem. 1897, [2] 56, 146). By reducing coniferin in alkaline solution with sodium amalgam, eugenol is formed (Chiozza, Chem Zentr. 1888, 443). Methyl eugenol (allyl guaiacol) has been prepared synthetically by treating veratrol (dimethylcatechol) with allyl iodide and zinc-dust (Moureau, Compt. rend. 121, 721).

Pure eugenol is a colourless oil; b p. 244 5° (Frankforter and Lando, J. Amer. Chem Soc 27, 641), 247 5° (Wassermann, Annalen, 179, 366), 253'5° (corr ) (Perkin, Chom Soc Trans 1896, 1247); sp.gr. 1 0785 at 4°/4°, 1 0734 at 10°/10°, 1 0663 at 20°/20°; specific magnetic rotation, 2·1974; molecular magnetic rotation at 15°, 18.727 (Perkin, l.c.); dielectric constant, 60 (Mathews, J. Phys. Chem. 9, 641),  $\mu_{\rm D}$  1.54437, sp. heat, 0 5024; mol. heat of combustion, 1304 cals. (F. & L. lc) Eugenol reddens litmus, smells strongly of cloves, and has a burning taste It does not reduce Barreswil's solution but reduces ammoniacal silver solutions on long standing Eugenol may be detected by the following reactions (i) ferric chloride imparts a blue colour to its alcoholic solution; (ii) zinc chloride yields a transitory pale-yellow colour; (iii) sulphuric acid gives first a brown colour, changing quickly to purple and finally to wine-red; (iv) Burgess (Analyst, 1900, 25, 265) makes use of a reagent prepared by dissolving 10 grams of mercuric salt in 25 n c. sulphuric acid, and making the solution up to 100 cc. with the latter. By shaking 2 c.c of eugenol with 5 c.c of the reagent, and allowing the solution to stand, a pale violet colour is developed (For estimation of eugenol in oil of cloves, v. Verley and Bosling, Ber 1901, 3359; Thoms, Chem. Zeit. 1903, 27, 954; Arch. Pharm. 241, 592; Spurge, Pharm J. 1903, [4] 16, 701, 757; Schimmel & Co., Chem. Zentr. 1903, n. 1124; Reich, Zeitsch Nahr. Genussm 1909, 18, 401.)

Eugenol and its derivatives are readily converted into the corresponding derivatives of the isomeric isoeugenol (v. infra), which on oxidation yields vanillin and its derivatives Chromic acid oxidises eugenol to acetic acid, carbon dioxide, and water (Wassermann, lc.); chromyl chloride, however, combines with eugenol, and on decomposing the product with water, vanillin can be extracted (Novarine, US. Pats. 365918, 365919; Ber. Ref. 1896, 812). Nitric acid oxidises eugenol to oxalic acid and a resin, but in ethereal solution, 5-nitro-eugenol is formed (v. infra). Ferric chloride oxidises it to dehydroeugenol, mp. 105°-106° (Cousin and Hérissey, Compt. rend. 1908, 146, 1413). Oxidised with potassium permanganate, eugenol yields homovanillin, vanillin, and vanillic acid (Erlenmeyer, Ber. 1876, 273) Hydrogenisation of either eugenol or isoeugenol in the presence of platinum black, yields dihydroeugenol (propylguaiacol), b.p. 246°-248° (Fournier, Bull. Soc. chim. [4] 7, 23).

5-Nitroeugenol is prepared by acting on eugenol in ethereal solution with nitric acid. It forms large triclinic crystals, m.p. 43°-44°

(Weselsky and Benedikt, Monatsh. 3, 388). With tin and hydrochloric acid, it produces aminochlorhydroeugenol hydrochloride; aminochlorhydroeugenol melts at 97°. Nitroeugenol, heated with acetic anhydride and sodium acetate, forms intro-aceto-eugenol, mp. 61°, which, oxidised by potassium permanganate at 70° (Tiemann, Ber 1876, 418), yields nitrovanilic acid. C<sub>4</sub>H<sub>2</sub>(COOH)(OCH<sub>3</sub>)(OH)NO<sub>2</sub>, decom-

posing at 210° without melting.

Acyl and aryl derivatives of eugenol are obtained by treating eugenol with acyl or aryl anhydrides in the cold, in presence of some condensing agent such as sulphuric acid, zinc chloride, pyridine, &c. The chlorides may be substituted for the anhydrides (Merck, D. R. PP. 103581, 109445, Chem Zenti 1899, 11 927, 1900, 11 407; Riedel, D. R. P 189333; Chem. Zentr 1908, 1. 185, Pond & Beers, J. Amer. Chem. Soc. 19, 825). Acetyl eugenol, which occurs in oil of cloves (Erdmann, J. pr Chem. 1897, [2] 56, 146), melts at 29°, and boils at 281°–282° (752 mm.) (Freyss, Chem Zentr. 1899, 1 835), benzyl eugenol has b p. 235°, and on heating with potash and subsequently with potassium dichromate and sulphuric acid, the benzyl ether of vanillin is obtained (Boehringer and Sons, D. R. P. 65937; Ber Ref. 1893, 211) Eugenol carbonate, m.p. 93°-94°, is obtained by passing phosgene gas into an alkaline solution of eugenol, on treatment with ammonia, the carbamate is formed (F. von Heyden Nachf D. R P 58129, Ber Ref 1892, 186), the car-bonate can also be obtained from eugenol and the diphenyl ester of carbonic acid (D. R. P. 99057, Chem Zentr. 1898, n. 1190) adding eugenol to a cooled mixture of pyridine and chlorsulphonic acid in carbon disulphide, potassium eugenyl sulphate, m.p. 203°, is obtained, the latter is transformed by heating with excess of potash into potassium isoeugenylozone, yields potassium vanillin sulphate (Verley, Bull. Soc. chim. 1901, [3] 25, 46).

### Eugenyl acetic acid

$$C_6H_3(CH_2 CH : CH_2)(OCH_3)(OCH_2 CO_2H)$$

is obtained by heating eugenol with chloracetic acid in the presence of sodium hydroxide for 20 hours, and decomposing the sodium salt thus formed with hydrochloric acid, m.p. 75°. By heating the acid with potash and amyl alcohol at 150°, isoeugenyl acetic acid, m.p. 92°-94°, is produced; and oxidation of the latter with potassium permanganate yields vanillin acetic acid (Gassmann and Krafft, Ber. 1895, 1870).

Eigenyl phosphoric acid is obtained by treating eugenol with phosphorus oxychloride and decomposing the resulting chloride with water; in p. of anhydrous acid, 105° (Boehringer & Sons, D. R. P. 98522; Chem. Zentr. 1898, ii. 950; Oeser, Annalen, 131, 277). On heating with alcoholic potash, isocugenyl phosphoric acid is obtained, and by heating the acid sodium salt of the latter to 150°, isocugenol is obtained quantitatively.

Eugenyl alcohol is prepared by heating eugenol in 20 p c. sodium hydroxide solution with formaldehyde at 60°; m.p. 37°-38° (Monasse, Ber 1902, 3894).

Eugenol glucoside, m.p. 132°, is produced by the action of acetochlorhydrose on the potassium 688 EUGENOL.

salt of eugenol (Michael, Amer. Chem. J. 6,

Eugenol and isoeugenol form compounds with proteins which are stated to be used in the treatment of tuberculosis (Belart, Eng. Pat. 9246, 26075; Fr. Pat. 389889; J. Soc. Chem. Ind. 1908, 998; 1909, 1325).

For compounds with iodine, v F. v. Heyden Nachf. D. R P. 70058; Ber Ref. 1893, 915; Ingle, J. Soc. Chem. Ind. 1904, 422) The Nsubstituted aminoethyl compounds of eugenoland isoeugenol- acetamide are stated to be used as local anæsthetics (Einhorn, D. R. P. 208255; J. Soc. Chem Ind 1909, 491).

Bromine in excess forms with eugenol, dibromeugenol dibromide, m p 118°-119°, which, treated with zinc-dust in alcoholic solution, yields dibromeugenol, m.p. 59° (Chasanowitz and Hell, Ber. 1885, 823, Auwers, ibid 1901, By treating eugenol with bromine in chlorotorm at 0°, bromeugenol is formed (Lando, J. Amer Chem. Soc. 1905, 27, 641) Tubromeugenol and other brom- derivatives (v. Lando, l c., Auwers and Muller, Ber. 1902, 35, 114;Hell and Anwandter, ibid. 1895, 2085)

Eugenol methyl ether occurs in several ethereal oils, and can be obtained from veratrol (v. supra). It is also obtained by the action of methyl iodide and caustic potash on eugenol (Ciamician and Silber, Ber. 1890, 1164); b p 247°-248°. It is readily converted into isoeugenol methyl ether, b p 263°, which yields methyl vanillin and veratric acid on oxidation with potassium dichromate and sulphuric acid

iso Eugenol

 $(HO)(CH_2O) C_6H_3 \cdot CH CH \cdot CH_3(C_3H_5 CH_3O HO = 1 3.4)$ occurs in small quantities in oil of nutmeg (Power and Salway, Chem Soc Trans 1907, 2041), and as the methyl ether in the oil from Asarum arifolium (Michx) (Miller, Arch Pharm. 240, 371). It is prepared by adding 5 parts of eugenol or an equivalent quantity of oil of cloves to 12.5 parts of potash, dissolved in 18 parts of amyl alcohol, and heating for 16-20 hours at 140°. The product is steam distilled, and the residue acidified with sulphuric acid. The isoeugenol separates out, is washed with sodium carbonate solution, and purified by distillation (Tiemann, Ber. 1891, 2870; 1894, 2580, Lairi, Eng Pat 17547, J. Soc. Chem Ind. 1891, 854, Haarmann and Reimer, D. R. P. 57808, Ber. Ref 1892, 94). The amyl alcohol may be replaced by methyl, ethyl, or butyl alcohols (Gassmann, Compt. rend. 124, 38; F. von Heyden Nachf. D. R. P. 70274; J. Soc Chem Ind 1892, 633). Other methods are to heat eugenol with 4 parts of caustie potash and some water at 220° (Einhorn and Frey, D. R P 76982; Frdl m 863; Ber 1894, 2455); to heat the alkalı derivative of eugenol to 200° in the absence of air (Fritzsche, D. R P. 179948; J. Soc. Chem. Ind. 1907, 1216); and to heat the acid sodium salt of isoeugenylphosphoric acid to 150° (v. Eugenylphosphoric acid).

iso Eugenol forms colourless crystals, m p. 33°; b.p. 264°-265° (J. Soc. Chem. Ind. 1892, 633), 193 5° (100 mm.), and 266 5°-268.5° (760 mm ); sp gr. 1.0994 at 4°/4°, 1.0944 at 10°/10°, 1.0872 at 20°/20°; specific magnetic rotation 2.5778; molecular magnetic rotation at 15°, 21·469 (Perkin, Chem Soc Trans. 1896, 1247). On

oxidation, isoeugenol and its derivatives yield the corresponding derivatives of vanillin. isoeugenol, when treated with sodium peroxide, yields vanillin itself (Haarmann and Reimer. Eng. Pat. 11952, 14928; J. Soc. Chem. Ind. 1895, 506; 1897, 633; cf. Majert, D. R. P. 82924; Ber. Ref. 1895, 878). Ozone oxidises isoeugenol in the same way, and vanillin is also produced by the electrolysis of the sodium salt of isoeugenol and subsequent addition of acid (Otto and Varley, U.S. Pat. 553039, 553593, Ber. Ref 1896, 249; cf. Otto, Ann. Chim Phys. 1898, [viii.] 13, 120). Ferric chloride oxidises isoeugenol to dehydroisoeugenol, m.p. 133° (Cousin and Hérrisey, Compt. rend. 1908, 147, 247).

sso Eugenol is readily polymerised by small quantities of mineral acids, acid chlorides, or metallic haloids, forming discougenol, which crystallises in colourless needles, mp. 178° (F. von Heyden Nachf. D. R. P. 70274; J. Soc Chem. Ind 1892, 633), 180° (Puxeddu, Gazz. chim. ital. 39, i. 131) Similarly, the ethyl ether of isoeugenol, when dissolved in alcohol hours, polymerises into diethyldissoeugenos, m.p. 130° (Puxeddu, l.c.; Wassermann, Annalen, 179, 375).

The acyl and aryl derivatives are obtained by methods similar to those employed for the eugenol compounds, but the solutions must be kept cool to prevent polymerisation taking place, acetyl isoeugenol, mp. 79°-80°; benzoyl isoeugenol, mp. 103°-104°; benzyl isoeugenol, mp. 48° (Pond and Beers, J. Amer. Chem. Soc. 19, 825; Merck, D. R. PP. 103581, 109145; Chem. Zentr. 1899, 11. 927; 1900, 11. 407; Haarmann and Reimer, D. R. P. 57568; Ber. Ref 1892, 93) If acetyl chloride is added to isocugenol at 54°, and the temperature is finally raised to 80°, diacetyldisseugenol is obtained, and by hydrolysis with acids, disoeugenol is formed (Tiemann, Ber. 1891, 2870; 1894, 2580). Methylene chloride reacts with an alkaline solution of eugenol forming methylenedissoeugenol, m p. 50°-52°, which on oxidation yields methylenedivanillin (Lesault & Co., D. R. PP. 75264, 76061; Ber Ref. 1894, 812, 928).

5-Nitrossoeugenol is formed by the action of nitric acid on isoeugenol in acetic acid solution, on recrystallisation from isobutyl or amyl alcohols it forms a reddish-vellow mass, which decomposes at about 150° (Puxeddu and Comella, Gazz chim. ital. 1906, 36, ii. 450).

isoEugenol carbonate is prepared in the same way as eugenol carbonate (q.v.) (F. von Heyden Nachf D R P. 61848; Ber. Ref. 1892, 486).

By adding excess of bromine to isoeugenol, bromisoeugenol dibromide, m.p. 138°-139°, 19 formed. If only 1 molecule of bromine is added, an unstable isoeugenol dibromide, m.p. 95°, is produced Di- and tri-soeugenol dibromides have been prepared (Hell and Portman, Ber. 1895, 2088; Hell and Bauer, Ber. 1904, 1128; Zincke and Hahn, Annalen, 329, 1).

iso Eugenol methyl ether occurs in the oil of Asarum arifolium (v. supra). It can be prepared by warming eugenol methyl ether with alcoholic potash (Ciamician and Silber, Ber. 1890, 1165) or by treating isoeugenol with dimethyl sulphate and aqueous potash (Mannich, Aich. Pharm. 248, 127); bp. 263°.

isoEugenol is distinguished from eugenol by the following reactions. (1) ferric chloride gives an olive-green colouration with its alcoholic solution, (11) zinc chloride, a bright rose-pink colour; (m.) sulphune acid, a rose-pink colour changing to light-brown

EUGUFORM, EUMYRDIN, EUPHTHAL-MIN, EUPORPHIN v SYNTHETIC DRUGS.

EUKODAL. Oxydihydrocodemone hydrochloride Used as a narcotic (Freund and Speyer, Munch med Woch 1917, 64, 380).

EULATIN. Trade name for a mixture of p-bromobenzoic acid, o-aminobenzoic acid, and antipyrine

**ÉŬONYMUS** (Euonymi Cortex, B.P.) (deleted from U.S. P. 1916) is the root bank of E. atropurpureus (Jaquin), a shrub indigenous to the United States, and known there as 'wahoo' or burning bush.' It is said to contain an amorphous bitter principle and a glucoside euonymin; the latter, according to Hans Meyer (cf. Schmiedeberg, Aich exp Path. Pharm. 1883, 16, 163), possesses a digitalis-like action, causing systolic arrest of the flog's heart in very small doses Nothing is known about it chemically. For a. account of earlier investigations, v Rogerson (Chem Soc. Trans. 1912, 101, 1040), who isolated dulcitol (2 1 p c of the bark) and other substances, but could not find any glucoside The latter negative result may have been due to hydrolysis by the steam distillation employed The name euonymin is also applied to dried extracts of varying composition, brown euonymm being from the root bark, and green (containing chlorophyll) from the stem back

EUPAD. An antiseptic consisting of equal weights of chloride of lime and bone acid.

**EUPHORBIC** EUPHORBIUM, ACID, EUPHORBONE v Gum resins.

EUPHORINE v SYNTHETIC DRUGS.

**EUPHRASIA TANNIN** v TANNINS.

EUPHYLLIN. Trade name for a compound of theophyllme with ethylene-diamine white crystalline soluble powder, used as a diuretic (v. SYNTHETIC DRUGS).

**EUPITTONIC ACID, Eupittone, Pittacal,** Hexamethoxyaurin C25H26O9

$$O \ \, \underbrace{\begin{array}{c} \mathrm{OMe} \\ \mathrm{OMe} \end{array}} \ \, \mathrm{C[C_6H_2(\mathrm{OMe})_2\mathrm{OH}]_2}$$

Pittacal is a blue colouring matter obtained by Reichenbach in 1835 from the highboiling portions of beechwood tar. Gratzel also from the same products obtained a brown substance which gave the same blue colour reaction with baryta as the pittacal of Reichenbach. The colouring matter, which only exists in small quantity in this brown substance, was extracted and examined by Liebermann (Ber. 9, 334). It is an orange-red crystalline powder, soluble in alcohol and acetic acid, forming brown solutions. It forms with alkalis purple. and with ammonia blue solutions, from which carbon dioxide precipitates blue salts, soluble in pure water. Calcium, magnesium, barium, and tin salts also give blue precipitates with the alkaline solutions. The calcium and magnesium salts are soluble ın water, Hydrochloric and sulphuric acids form red Vol. II.-T.

solutions, and the latter turns pure blue on warming. In acid solutions, it dyes animal fibres orange, in ammoniacal solution especially with tin mordant, blue-violet Liebermann called this colouring matter eupittone, and regarded it as at least closely allied to pittacal (Bei 11, 1145) Hofmann (Ber. 12, 1371, 2216), by the action of hexachlorethane C<sub>2</sub>Cl<sub>6</sub> on dimethyl pyrogallate mixed with alcoholic potash or soda, obtained a substance identical in composition with the cupittone of Liebermann, and strongly resembling it in properties. It is an orangeyellow substance, soluble in acetic acid, and precipitated therefrom by alcohol. It melts at 200° and decomposes partly into a blue sublimate. It forms the same alkali salts and gives the same sulphuric acid reaction as cupittone. This body is hexamethoxy-rosolic acid C<sub>19</sub>H<sub>8</sub>(OCH<sub>3</sub>)<sub>6</sub>O<sub>3</sub> Heated with alcoholic ammonia, it yields hexamethoxy-rosandine

which crystallises in broad colourless needles, and has no tractorial properties. When 2 grams cupittone are heated with 20 cc alcoholic methylamine (10 pc) at 150°-160° for 21/2 hours, it yields hexamethoxydimethylformylparaleucamline

 $CH[C_6H_2(OMe)_2NHCH_3]_2C_6H_2(OMe)_2NHCHO$ m p 187° (Liebermann, Bei 34, 1034) gallol is produced in attempting to replace the methoxyl groups by hydrogen. Its formation is analogous to that of iosolic acid

$$3C_8H_{10}O_3+CO_2=C_{25}H_{26}O_9+2H_2O$$

Hofmann terms this body eupittonic acid, and uses the name as synonymous with pittacal and Eupittonic acid cannot be obtained eupittone by heating the sodium derivative of dimethyl pyrogallate or of dimethyl-methyl-pyrogallate separately, in air, a mixture of the two, how-ever, with excess of soda gives sometimes as much as 10 pc of its own weight of the acid. The action is comparable with that in which rosamline is produced from aniline and toluidine.

$$-2C_8H_{10}O_3 + C_9H_{12}O_4 = C_{25}H_{26}O_9 + 3H_2$$

The oxygen necessary to remove the hydrogen is obtained from the atmosphere, as in a closed vessel no such change as the above takes place. Eupittonic acid appears to be dibasic, the sodium and barium salts,  $C_{25}H_{24}Na_2O_9$  and  $C_{25}H_{24}BaO_9$ , have been prepared. Paper steeped m a solution of the acid, and dried, is a most sensitive test for free alkalis on account of the blue colour of the alkaline salts of cupittonic Hydrochloric acid, however, produces With acetic anhydride, the same blue colour a diacetyl derivative,  $C_{25}H_{24}O_9(C_2H_3O)_2$ , mp. 265°, is produced, which, curiously, is insoluble in alkalis, although its molecule ought to contain four hydroxyl groups

The dibenzoyl derivative  $C_{25}H_{24}O_9(C_7H_7O)_2$  forms golden needles, mp  $232^\circ$ , the dimethyl ether  $C_{25}H_{24}Me_2O_9$  has mp  $242^\circ$ , the diethylother  $C_{25}H_{24}Et_2O_9$  has mp.  $201^\circ-202^\circ$ ; and the 10do compound C25H26O9I4 forms brown shining

Eupittone black, noreupittone, hexaoxyaurin  $C_{19}H_{14}O_7$  or  $[C_6H_2(OH)_3]_2C: C_6H_2(OH)_2: O$ is formed by adding eupittone to sulphuric acid at 140° and keeping the temperature at 125° until the mass is pure blue; it is a dark shining black powder, and dyes mordanted fibres through a dirty violet to a deep black, the hydrochloride  $C_{19}H_{14}O_{7}, HCl, C_{2}H_{6}O$  is deep blue (Liebermann and Widermann, Ber. 34, 1033)

EUPNINE. Trade name for a preparation

of caffeme and rodine

**EUPORPHINE.** Trade name for apomorphine methyl bromide.

EUPYRIN v SYNTHETIC DRUGS.

**EUQUININE.** Trade name for quinine ethyl carbonate

EUREKA v. NICKEL

EURESOL. Syn for resorcinol monoace-

EURHODINES, EURHODOLS v AZINES EUROBIN. Trade name of a pharmaceutical preparation consisting mainly of the

diacetate of chrysarobin (q v)**EUROPHEN** v Synthetic drugs

**EUROPIUM.** Eu = 152 (cf Urbain and Lacombe, Compt rend 1904, 138, 637, and Jantsch, ibid 1908, 146, 173)

Source --The oxide of this extremely late element is present in very small amounts in monazite sand, and is extracted from this mineral in the fraction containing the yttrum earths

Exhaction.—In separating europium from its associates, advantage has been taken of the isomorphism existing between the double intrates of bismuth and those of the metals of the rare earths.

Bismuth nitrate gives rise to a series of double salts having the general formula

$$3M^{II}(NO_3)_2, 2B_1(NO_3)_3, 24H_2O$$

where the bivalent metal may be magnesium, zinc, nickel, cobalt, or manganese (Urbain and Lacombe, Compt. rend. 1903, 137, 568). Magnesium bismuth nitrate

$$3 {\rm Mg} ({\rm NO_3})_2, 2 {\rm Bi} ({\rm NO_3})_3 24 {\rm H}_2 {\rm O}$$

is isomorphous with the corresponding double magnesium nitrates

$$3 {\rm Mg}({
m NO_3})_2, 2 {
m M^{IM}}({
m NO_3})_3, 24 {
m HI}_2 {
m O}$$

where MIII Ce, La, Nd, Pr, Sm, Eu, Gd, and Tb

To a mixture of these metals in the form of then double magnesium intrates is added a large excess of magnesium bismuth intrate When the product is crystallised methodically from nitric acid solution, the double magnesium nitrates of cerium, lanthanum, praseodymium, neodymium, and samanium first separate, followed by the double salt of bismuth, leaving curopium and gadolinium in the mother liquors As the fractional crystallisation proceeds, the rare earths yielding double nitrates less soluble than the bismuth double salt, are gradually climinated until at a certain point the middle fractions contain no other tervalent metal than When this point is reached, all the samarum has been crystallised out, and thus separated from the europium and gadolinium The bismuth still in solution is removed from the mother liquor by hydrogen sulphide, and the europium and gadolinium separated by further fractionation of the filtrate, the europium double nitrate being the first to crystallise from this solution (Demarçay, Compt. rend. 1900, 130 1019, 1469, Urbain and Lacombe, *ibid.* 1904, 138, 627). This crystallisation process is a satisfactory method for obtaining pure europium, samarium, and gadolinium, and affords a means of separating quantitatively europium and samarium (cf. Eberhard, Zeitsch. anorg. Chem. 1905, 45, 374, James and Robinson, J. Amer. Chem. Soc. 1911, 33, 1363).

J. Amer Chem Soc. 1911, 33, 1363).

Europia (Europrum oxide) Eu<sub>2</sub>O<sub>3</sub>, obtained from the purified double salt, shows no samarium spectrum, and only the strongest lines of gadolimum (Demarçay, Compt. 1end 1901, 132, 1482), it has a pale pink tint, but when produced by heating the sulphate at 1600°, the colour is somewhat deeper

Europous chloride EuCl, is a colourless amorphous substance, forming a neutral solution

in water, which oxidises at 100°.

$$12EuCl_2 + 3O_2 = 8EuCl_3 + 2Eu_2O_3$$

Europium sulphate  $\rm Eu_2(SO_4)_3.8H_2O$  forms pale-pink crystals, stable in air and dehydrated completely at 375° (Compt\_rend\_1904, 138, 627)

The europium salts yield pale-pink solutions and in strong nitric acid they exhibit an absorption spectrum consisting of eight somewhat faint bands

The spark spectrum is characteristic, and was first observed by Lecoq de Boisbaudian in that given by crude samaria, three very intense line in the blue being ascribed by him to a new earth  $Z_t$ . The reversion spectrum shows the band  $\lambda 615$  referred by de Boisbaudian to the earth  $Z_\zeta$  (Compt. rend. 1893, 116, 674., 117, 199)

The most intense lines of the arc spectrum are very characteristic, and constitute a delicate test for europium, which can be thus identified in the crude oxides from monazite after icmoving the ceium (Eberhard, Zeitsch anorg Chem 1905, 45, 374). The cathode phos-The cathode phosphorescence spectrum exhibits the anomalous iay referred by ('rookes to the 'meta-element' S<sub>δ</sub> Pure europia, however, give no phosphorescence spectrum unless diluted with some other earth. With lime in the proportion Eu<sub>2</sub>O<sub>3</sub> 99CaO, a red phosphorescence is observed, and similar effects are obtained on mixing europia with alumina or gadolima Mixtures with excess of the latter give a white. and with excess of europia a red, phosphorescence (Urbain, Compt rend 1906, 142, 205, and Seventh Congress of Applied Chemistry, 1909, section 10, 94)

(Ultra-violet and arc spectra, v Exner and Haschek, Sitzungsber Wien Akad 1902, Ha, 3, 42, and Die Wellenlangen der Bogenspektra, 1904)

Europium has been detected spectroscopically in the sun's chromosphere, in α-Bootis and in β-Geninorum (Lunt, Proc Roy Soc 1907, Λ, 79, 118)

EUROSAMARIUM. An element detected by Eder, and occurring associated with europium and gadolimium in monazite (Sitzungsber K K. Akad Wiss Wien 1917, 11a, 126, 473)

**EUSCOPOL.** Syn for mactive scopolamine hydrobromide

EUSOL. An antiseptic, used in surgery, consisting of a solution prepared by shaking up in a litre of water 25 grams of a mixture of equal weights of chloride of lime and boric acid, and filtering off the sed-ment; the solution

contains about 0.54 p.c. of hypochlorous acid, | the maximum pressure which corresponds with and 1.28 p.c. of calcium biborate. Although evaporation occurs

EUSTENIN. Trade name for an addition product of theobromine-sodium and sodium todide

EUSYNCHITE v. VANADIUM.

EUXANTHIC ACID, EUXANTHONE, and EUXANTHONIC ACID v. Indian Yellow

EUXENITE. A rare-earth mineral consisting essentially of columbate and titanate of yttium and erbium, with smaller amounts of uranium, thorium, cerium earths, iron, &c Crystals are orthorhombic, but are very rare and indistinctly developed; the mineral is more often massive, with a blackish-brown colour and a brilliant pitchy lustro; sp gr. 46-51, H 61 In appearance and character, it closely resembles several other rare-earth minerals, particularly polycrase, from which it differs in containing a lower ratio of (Cb,Ta)2O5 TiO, It is occasionally found in the felspar quarries at several places in the neighbourhood of Atendal in Norway (v W C Biogger, Die Mineralien der Sudnorwegischen Gramt-Pegmatitgange, 1906) Also in the pegmatite veins of Madagascar, Cooglegong in Western Australia, South Sherbrooke in Ontario, &c Named from εὔξενος, friendly to strangers, in allusion to the several rare-earths it contains

EVAPORATION. The term 'evaporation' is used in connection with technical or manufacturing operations when a liquid is converted into vapour, but in chemical manufactures it is generally limited to those processes where the object is to separate a liquid from a solid which is dissolved in it, or to separate a more volatile from a less volatile liquid. In cases where the liquid to be evaporated is one of the products of the process, and is collected by condensation, the process is known as Distillation  $(q \ v)$ a solid substance contains water, not as water of crystallisation, and without presenting visible moisture or wetness, the process of removing this moisture under suitable circumstances is known as Desiccation of Drying  $(q \ v)$ . The operation of evaporation, however, is conducted on the greatest scale in the generation of steam for steam engines, and in the production of steam generally for manufacturing purposes. Although more water is evaporated in connection with the steam engine than for any manufacturing processes, yet great quantities of steam are used as a convenient means of applying and distributing heat, especially when regulated and moderate temperatures are required. In such cases the heat is applied to the vessel to be heated by jacket spaces or through coils of pipes immersed in the liquid to be raised in tempera-Such applications are made in sugar and confectionery manufacture, and in distillenes, breweries, and dyeing, &c.

Water and many other liquids evaporate without the application of heat. A vessel of water freely exposed at the ordinary temperature of the atmosphere soon evaporates and disappears. This is due to the fact that water vapour has a sensible pressure at ordinary temperatures, and as gases and vapours freely diffuse throughout each other, any water placed with a free surface in a confined space continues to evolve and diffuse its vapour until the water vapour present in the whole space is . 5

the temperature Although evaporation occurs in this way without any application of increased temperature, yet heat is absorbed by the vapour formed from the liquid, and if the evaporation be rapid and be conducted under such circumstances that heat from surrounding objects is prevented from reaching the liquid, then its temperature will be greatly reduced. This fact temperature will be greatly reduced is illustrated by the well-known lecture experiment of placing a shallow copper cup filled with ether upon a block of wood, and under it a drop of water When an is blown upon the ether it rapidly evaporates and carries off so much heat from the copper that the water drop under it is trozen and the copper adheres to the wood A liquid is said to be volatile when its vapour has a considerable pressure at ordinary atmospheric temperatures At 20°, for example, the pressure of water vapour (or tension as it is somewhat mappropriately called) is 1739 mm mercury, whilst the pressure of ether vapour at the same temperature is 433 3 mm and that of alcohol 44.5 mm If the liquids be placed in atmospheres whose pressures could be kept at these points by artificial means, then these liquids would boil As the evaporation under these circumstances is very rapid, relatively to the flow of heat to the liquid from surrounding bodies, if only atmospheric temperature be available, the heat is abstracted so quickly that many liquids freeze

Advantage is taken of these properties for many purposes in the use of an currents passed over the surfaces of liquids to evaporate them without the application of other than natural heat. Reduced pressures are also applied to evaporate without heating or to reduce the temperature necessary in order to avoid chemical decomposition and also to produce cold mechanically

Evaporation in steam boilers. The evaporative efficiency of the steam boiler has been more closely studied than that of any other type of evaporating apparatus, and the principles of the economical application of heat for evaporating are more completely known in this connection than in any other

The leading types of steam boilers now in use for the purposes of motive power are as follows Lancashire, Cornish, Locomotive multitubular, Marine multitubular, and Water-tube or tubu-

lous types

Obviously in an article of this character detailed description of the various types of boiler is uncalled for, the reader is therefore referred to the numerous text-books existing on the subject, such, for example, as "Steam-Boiler Construction," by W.S. Hutton; "Steam-Boilers, Their History and Development," by H. H. P. Powles, "Steam-Boilers," by C. H. Peabody and E. F. Miller

Evaporation in open vessels. In many chemical manufactures it is necessary to evaporate off water in which the substance, the object of the process, is dissolved, and in such cases, when heat is to be applied, it is desirable to economise fuel to the greatest possible extent. The heat of waste gases is therefore used as much as possible. The method of applying heat varies considerably with the chemical nature of the substance dealt with, and choice of economical apparatus is often limited by the corresive nature.

of concentrating oil of vitriol from the dilute acid, a series of shallow open lead pans receives the diluted acid and heat is applied below hot furnace gases are allowed to impinge upon the bottom of the vessel containing the weakest acid, and therefore requiring the greatest quantity of heat for evaporation, and as the gases cool they impinge upon the succeeding vessels of stronger and stronger acid, requiring less quantity of heat until the acid has become as concentrated as it is safe to trust to leaden Instead of this plan, in some cases vessels the furnace gases are passed over the surface of the acid in order to heat it from above, and the volume of gases in motion removes the vapour as fast as it is formed The advantage of surface evaporation arises not from the fact of its being economical, because liquids are heated with difficulty from above. The hot liquid in contact with the gases expands and has no tendency to cause convection currents, but rather remains stationary above the colder portions, and accordingly it is always advisable to heat from below when possible In many processes, however, during evaporation, a deposit comes down and this, coating the bottom of the vessel, prevents the free contact of the metal with the liquid The resistance to the passage of heat from the metal to the liquid is thereby much mereased, and the bottom suffers rapid destruction from over-heating In boilers using very hard water this is a fruitful source of trouble and danger

When surface evaporation is resorted to, large surfaces are exposed in shallow tanks over which the flame and hot gases from a furnace are allowed to pass The pans are covered in so that the radiant heat from the brickwork assists the action of the flame and hot gases. Arrangements of this kind are used in evaporating weak alum liquors, and the tanks are built up of bricks bound together with a cement composed of lime and alum shale. Sometimes an open pan is placed above the gases to contain the feed liquid and heat it in its progress to the evaporating

Evaporation of brine. Brine is sometimes evaporated by allowing the liquor from the springs to flow over masses of twigs in such a manner as to expose very extensive surfaces to the action of air currents, by pumping over a succession of these surfaces it is gradually con-When artificial heat is applied to the evaporation of brine, large shallow salt pans are used They consist of flat quadrangular vessels of plate non supported upon suitable brick walls, forming flues which distribute the heat from furnaces placed in the centre part under the vessel The flames strike directly upon the bottom of the vessel, and the hot products of combustion are conducted outwards by flues leading to the stack or chimney. In this way they traverse the whole bottom surface of the evaporating pans. The pans are of great area, often 60 feet long by 30 feet wide. The water vapour is collected by a large conical hood and carried off by a separate chimney. The sides of the lower part of the cone are so constructed that they can be opened to admit air from the direction in which the wind is blowing. This assists the evaporation by carrying off the water

of the liquid to be evaporated. In the process | vapour generated and exposing the surface to a continual air current.


If heat enough to cause a liquid to boil be applied, an an current will not increase the evaporation, but will tend to diminish it by cooling the vessel containing the hquid; but when the liquid does not reach the boiling-point, the continuous removal of the atmosphere in contact with the liquid is necessary. If this be not done, then, so soon as the atmosphere has become saturated with the vapour, that is, so soon as the water vapour is present in quantity sufficient to produce the pressure which corresponds with the temperature of the liquid, the evaporation ceases.

It is a necessary condition of evaporation at temperatures less than the boiling-point, that the atmosphere into which the evaporation takes place should be continuously ranewed, and it is advisable that the air into which the water evaporates should not be of lower temperature than the liquid. When the temperature of the air is lower than that of the liquid, the volume of vapour carried off is much reduced, as the air must be heated by the vapour before it is capable of carrying off the volume equivalent '> the vapour pressure. Producer gas is used for evaporating the brine at Ischl salt works, and the operation is conducted in deep rectangular plate-iron vessels grouped together, and fed continuously by a system of distributing pipes.

Steam is much used to supply the heat iequired for evaporation, especially when the substances to be dealt with are of an easily decomposable nature if overheated. The heat is often applied to the vessel by means of a steam jacket, that is, the vessel is made double walled, and the interspace is utilised to receive the hot steam. When the temperature requires to be higher than 100°C, the jacket space is constructed to withstand pressure, and steam under boiler pressure is supplied, but steam is most commonly used at atmospheric pressure and applied to the liquid by passing through rows or coils of pipes namersed in it The arrangement of these pipes varies, and in some cases they are so connected as to be capable of hingeing out of the vessels for ready access and cleaning purposes.

Milk is now rapidly evaporated by spraying into heated air without raising the milk temperature to such a point as would cause chemical change

Evaporation under reduced pressures. In the older processes of sugar manufacture, the sugar liquois were concentrated by the direct application of heat from a fire upon the vessels containing them, but the greatest difficulty was experienced in preventing decomposition when the solution became concentrated. To avoid this difficulty, the vacuum pan was introduced by Howard, and in various modified forms it is still used. The boiling-point of a liquid depends upon the pressure to which it is subjected, and by reducing the pressure the boiling-point is greatly reduced. The vacuum pan consists essentially of a globular copper vessel containing the sugar liquor, and having the lower portion steam jacketed, while steam also passes through a coil of pipe immersed in the liquid. The upper part of the vacuum 1 an carries a circular steam dome or cylinder from which a pipe passes to a vessel intended to collect any liquor which may have primed over From this vessel it is run back into the pan again An air pump exhausts the apparatus and passes the steam to a condenser of usual construction. Fig 1 shows a vacuum pan made by Messrs. Manlove, Alliot, & Co. The vacuum pan A is formed in two por-tions bolted together. It is fed with liquor from the vessel B by suitable cocks, the pipe connections a, b being arranged to equalise the pressure within the vessel, so that the flow may be towards A. The live steam is admitted at the valve c, and passes through coil p into the jacket E, from whence it discharges and drains by the pipe F. The cylindrical dome or cylinder a prevents any great priming, but any liquid which may find its way by the pipe H is caught in the cylinder I and returned to A by the pipe K The pipe L leads to the air pump and condenser



A thermometer placed in the upper part of the pan indicates the temperature of the operation, whilst a vacuum gauge or barometer column is used to show the pressure within. The boilingpoint of the liquid at atmospheric pressure is about 250°F, and this is reduced to 150°F by using the vacuum pan

Fig. 2 shows a vacuum pan of modern construction, made by the Murlees Watson Co., Ltd., of Glasgow. In this vacuum pan, the liquor is heated in the vessel a by coiled steam pipes b, and also by reason of its passing upwards through small tubes c, through a steam chamber, d, the down flow of the liquor taking place through the central large tube e. In other respects, the operation of this vacuum pan is similar to that shown in Fig. 1.


Evaporation under reduced pressure by triple and multiple effect. It is evident that as a liquor becomes more concentrated the boiling-point rises, and in evaporating liquids liable to chemical change by overheating, the pressures under which boiling is effected must be arranged to keep the temperature within safe limits. In the case of sugar liquor this safe limit is 227°F Steam at 5 lbs. per square inch above atmosphere has this temperature, and if it be applied to a pan containing a weak sugar liquor at atmospheric pressure, the liquor will boil.

Neglecting losses by radiation and like bottom of the apparatus, the liquor, after causes, the steam coming from it has the same passing through the 'climbing film' tubes, amount of heat as was imparted to the liquor being conducted thereto by a set of 'falling by the steam under slight pressure, and if it be film' tubes also within the steam jacket B


conducted to the coil and jacket of a second pan in which a slight vacuum is maintained, it will generate almost its own weight of steam at lower pressure and temperature, and so evaporate the If the steam so water from the second pan generated be taken to the coil and jacket of a third pan, its heat can again be utilised to evaporate the liquor there, provided the vacuum is sufficient to allow it to boil. This process may be continued until the original heat of the steam supplied to the first pan is dissipated by conduction and radiation. This is the principle of the 'triple effect' apparatus, by which a great economy is effected in the amount of fuel needed to evaporate a given weight of houor. The steam from one pan at higher pressure proceeds to the coil and jacket of the next, and the steam from the second proceeds to the coil and jacket of the third, the pressures within the pans proceeding in descending order as well as the temperatures

Fig 3 is an illustration of a triple-effect evaporator of the 'Hagemann' type, constructed by the Mirilees Watson Co, Ltd, for concentrating sugar. In this evaporator, the liquor is fed through an internal perforated pipe D at the bottom of the vessel A, and the flow of liquid from vessel to vessel is automatically regulated by traps E, the liquor passing through the vessels A, B, and C in turn, the reduction in pressure being so arranged that it is least in A. more considerable in B, and greatest of all in C, so that the boiling-points are lower in each succeeding vessel. The heating surface is obtained by means of a number of vertical tubes F, through which the liquor passes, these tubes being arranged in a steam or vapour space, a. divided by horizontal baffle plates it steam enters the steam space at the top above the upper baffle plate, and passes to the intermediate and lower spaces through apertures in the baffle plates slightly larger than the tubes which pass through them, the rush of steam through the openings removing the water of condensation from the surface of the tubes, and improving their efficiency. The upper part of each vessel acts as a separator, the vapour escaping, past a baffle J to prevent priming, through an outlet K either to the next effect or to the condenser

In the Kestner 'climbing film' evaporator, one complete section of which is illustrated in Fig 4, vertical tubes are also used for the passage of the liquor, but they are fewer and of greater length than in the apparatus last described Steam supplied through an inlet A to a jacket B causes ebullition of the liquor, supplied to a tank c, releasing a volume of vapour in the tubes D which occupies the central portion of each tube and carries up a film of the liquor on the internal surface. A fixed baffle E, by means of suitably shaped vanes, produces a whirling motion of the liquor and vapour mixture as it emerges into the separator F, whereby the liquor is thrown outwards and drawn off through an outlet g, and the vapour escapes, through a 'save-all' H, by an outlet JIn another form the separator is placed at the bottom of the apparatus, the liquor, after bassing through the 'climbing film' tubes, passing through the 'climbing film'



connected to form a multiple effect apparatus, the liquor and vapour from the outlets a and J of one section being respectively led to the tank c and inlet A of the next



The Yaryan evaporator. This ingenious evaporator is the invention of H T Yaryan, of Ohio, USA, and it is now largely used in this country, the Continent, and America utilises evaporation under reduced pressures as with the triple effect and also multiple effect differs from other inventions of the same class in the method of applying heat to the liquid to be evaporated, whereby the liquid is heated in small quantities at a time, and is brought into very effective contact with the heating surfaces Instead of using coils immersed in a liquid which is also heated by jacketed surfaces, the heat is applied by steam to the exterior surfaces of tubes, and the liquid to be evaporated is pumped within these tubes. The tubes are arranged in straight cylindrical drums, and pass from end to end, terminating in chambers with partitions so placed that when the liquid is pumped in at one end of a series it must pass through the whole series before escaping into the separating chamber. Fig. 5 is a section of

Any number of sections as described may be one drum of the evaporator, showing that the liquid is pumped in at one end, travels along in the direction of the arrows, and finally escapes with the steam which is generated into the separator The tubes are divided into sections termed 'coils,' and each section consists of fine tubes coupled at the ends so as to form a con-The liquid and vapour pass tinuous passage from section to section in series, and so come into effective contact with the whole surface As the liquid is pumped in it evaporates, and the steam produced carries it forward through the tubes, so that a mass of liquid and steam is brought in contact with every part of the whole heating surface, moving at a high velocity because of the steam generated, which has only the one direction of escape, that is, through the whole system of tubes in each towards the separator. The liquor under treatment issuing from the last set of tubes into the separator falls to the bottom, as shown in Fig. 5, and the vapour using up flows in the direction of the arrow to the catch all, where any liquor entrained is caught during its passage through the number of small tubes with which this vessel'is fitted,




Fig 5

and, striking against the end cover, falls to the bottom and is drained off into a receiver This ensures dry vapour being passed on to the evaporating shell of the next vessel, where it in turn boils the liquor which is inside the tubes, and so on through the various effects until the vapour from the last vessel reaches the condenser. The advantages of this apparatus are its economical working and rapid flow of the liquor under treatment. Only a small quantity of liquid is in the apparatus at one time, and so it is exposed to heat for a much shorter period than in the ordinary vacuum pan, which is an important point. The appaiatus avoids entirely the loss of colour which always occurs to some extent in the older vacuum pans because of the long period of time taken to evaporate the large quantity of liquid they contain, amounting in some cases to thousands of gallons The Yaryan evaporator can be started very rapidly, as it contains but a small mass to be heated At the same time, the details are arranged to permit of ready access to the tubes and ready cleaning. This is greatly assisted by the fact that the tubes are Fig 6 shows a quintuple-effect straight evaporator with its vacuum and auxiliary pumps in general use for soda ash recovery in paper mills, caustic soda, caustic potash, meat extract,

distillery refuse, and other liquids readily amenable to concentration.

According to the patentees of the Yaryan apparatus, their double effect will evaporate





Fig 6.

16 lbs. of water for every pound of coal consumed under the boiler, 23½ lbs in a triple effect, 30½ lbs in a quadruple effect, assuming that the boiler is so constructed as to evaporate 8½ lbs of water per pound of coal

The Yaryan apparatus has, since its first introduction in 1888, been greatly improved in design and efficiency, and large sextuple-effect apparatus working on the Yaryan principle have been constructed by the Mirrlees Watson Co, Ltd, for the distillation of fresh water from salt water, producing from 25 to 350 tons per day of 24 hours, and also for use in the Solviy process of soda manufacture in Europe and America Fig 7 indicates the essential features of such an appuatus for the distillation of salt water Salt water, previously warmed by being passed through heaters, is supplied by a pipe A to heating tubes in the lowest evaporator B<sub>6</sub>, and thence is passed in succession by means of pipes c through heating tubes in the remaining evaporators It has then reached boiling-point, and is circulated through the evaporating tubes of the first evaporator B<sub>1</sub>, to which boiler steam is supplied by a pipe D. The mixture of liquor and vapour is conducted from the evaporators  $B_1-B_6$ , to the respective separators  $E_1-E_6$ , by pipes F, the outlets for the liquor and vapour from each separator to the next evaporator stage being by pipes  $\alpha$  and H respectively. ('oncentrated brine is drawn off from the last separator  $F_6$  by a pipe J, and the cordensed fresh water passes from stage to stage by pipes к, and is finally drawn off through an outlet L

In the Lillie type of evaporator the tubes are



also arranged horizontally, but the steam is passed through them and the liquor is allowed to fall over their external surfaces.

Evaporation in mechanical refrigerators. A separate class of mechanical refrigerators exists in which the cooling or refrigerating effect is produced by evaporation. Liquids used air ether, anhydrous ammonia, and anhydrous sulphur dioxide. In all of these the process of the machine is identical. The volatile liquid is during ice

introduced into a chamber at a pressure at which it is liquid, and the vapour is rapidly withdrawn from the vessel by a pump. The rapid evaporation absorbs heat so quickly that the temperature falls and a brine or other liquid is cooled within a coil of pipes immersed in the liquid. By circulating this brine, the low temperature may be utilised for reducing the temperature of a cold store room or for producing ice.

The vapour drawn off by the pump is com- 1. pressed and cooled by passing through tubes surrounded by water, and so becomes liquid again It is then injected into the evaporating chamber again, so that the process is practically continuous, and but slight renewal of the volatile liquid is necessary to make up for small les kages.

The same principle is applied in Carre's freezing machine for freezing water. In this case the vapour is removed from the surface of the water by a pump and condensed by strong sulphuric acid. Ammonia is also utilised in another apparatus in solution in water for the same purpose. By alternate expulsion from the water and absorption, cold is obtained

The rapid evaporation of liquid carbonic acid, air, and other bodies which are gaseous at normal temperatures and pressures, has also been used to produce low temperatures for liquefying the more permanent gases.

EVERNIC ACID and EVERNINIC ACID. Evernic acid, or lecanoric acid monomethyl ether, was first isolated by Stenhouse from the Everma prunasti (Annalen, 68, 83), and has been found also by Hesse (Ber 1897, 30, 366) to exist in the Ranalina pollinaria (cf. also Zopf, Annalen, 1897, 297, 271).

The lichen is extracted with diluted milk of lime, the extract neutralised with acid, the precipitate collected, dried, and digested with a little boiling alcohol The hot alcoholic liquid. treated with its own volume of water, deposits crystals of evernic acid (Stenhouse)

Evernic acid crystallises in small colourless needles, mp 168°-169°, readily soluble in hot alcohol, and when boiled with solutions of the alkalı hydroxides or baryta water, gives (O2, orcinol, and everninic acid

$$C_{17}H_{16}O_7 + H_2O = C_9H_{10}O_4 + C_7H_8O_2 + CO_2$$
  
Ramalic acid suffers a similar decomposition

Fischer (Ber 1913, 46, 3253, and 1914, 47, 505) showed that when evernic acid was methylated by means of diazomethane, in ethercal solution, a neutral crystalline ester identical with trimethyl-lecanonic acid methyl ester was obtained, and he pointed out that, as the methoxy group in everninic acid is in the paraposition to the carboxyl group, the structure of evernic acid must be.

From this it follows that the isomeric ramalic acid (v. Ramalic acid) may be expressed as

For descriptions of the Na, Ba, Ca, Cu, Pb, and Ag. salts of evernic acid, of Hesse (J. pr. Chem 1915, [ii ] 92, 425)

Diacelyl-evernic acid  $C_{17}H_{16}O_7Ac_2$  is a crystalline powder, m p.  $144^\circ$ 

Everninic acid (orsellinic acid monomethyl ether) resembles benzoic acid in appearance, melts at 157°, and yields the ethyl ester

m p. 72°. By digestion with hydriodic acid, everninic acid gives orcinol, ('O\_, and 1 molecule of methyl rodide According to Hemrich (Ber 1904, 37, 1406), two formula are possible for everninic acid:

$$(1) \begin{array}{c} \mathrm{CH_3} \\ \mathrm{OCH_3} \\ \mathrm{OCH_3} \end{array} \\ \mathrm{OH} \\ (2) \begin{array}{c} \mathrm{COOH} \\ \mathrm{OH} \\ \mathrm{OCH_3} \\ \end{array}$$

Fischer and Hoesch (Annalca, 1912, 391, 347) have, however, shown that orsellinic acid amethyl ether, prepared by them, and which has the structure (2), is identical with everning acid.

This conclusion has been further substantiated by the synthesis of everninic aldehude. m p. 65°C (and from it of evernine acid) (Hoesch, Ber 1913, 46, 886), by the careful methylation of orcyl aldehyde with methyl sulphate, and 2N caustic soda in the presence This when treated with an acctone solution of methylchlorocarbonate, yielded methylcarbonato-everninic aldehyde, needles, mp. 77°C, and the latter, when oxidised with potassium permanganate, was converted into methylcarbonato-everninic acid, needles, decomposing at about 100°C, which on hydrolysis with N caustic soda solution yielded everninic acid.

These changes may be represented thus:

Hesse (J. pr Chem. 1915, [n ] 92, 425) has described the following derivatives of everning acid :-

Acetyl everninic acid C9H9O4Ac, colourless prisms, m.p 111°C.

Nutro-everninic acid, orange-red needles,

тр 195°C CaHaOa(NO2)2HaO Dinitro-everninic acid golden-yellow needles, mp 87°C—the potassium salt is very explosive

253) of the isomerum of evernic acid and ramalic acid, and which was based upon the older formula of orsellinic acid, is as follows:

A G P and A. E E

EVERNIIN v. Gums

EVODIA MELIÆFOLIA. This tree, belonging to the Rutacew, is found in China and Japan. where its bark is largely employed in dyeing and It was formerly described by in medicine Loureiro as Pterocarpus flavus, but this error was eventually corrected by P W Squire (Pharm J 1888, [3] 18, 785), who showed it to be really Evodia glanca, which is synonymous with E melia folia By qualitative tests, Martin, Tokio (Arch Pharm 1898, 13, 337) and Squire (le) suspected the presence of berberne, and this colouring matter was subsequently isolated by Perkin and Hummel (Chem Soc Trans 67, A G P

**EVODIN**  $C_{17}H_{20}O_6$  A crystalline substance found in the fruit of *Evodia rutacarpa*, used in China and Japan in medicine and perfumery First isolated by Keimatzu by treatment of the dued fruits with benzeae, mp 285° Insoluble in water, alcohol, ether, and light petroleum, slightly soluble in hot benzene, ethyl acetate, and chloroform, soluble in acetic acid and alkali solutions (cf. Asahina and Ishio, J. Pharm Chim. 1916, [viii.] 13, 53)

EXALGIN. Trade name for methyl acetandide (v. Synthetic Drugs)

EXODIN & SYNTHETIC DRUGS

EXPLOSION, GASEOUS, Within years there has been an enormous increase in the extent to which the internal combustion engine has been utilised, and there are now many millions of h p at work on this principle Gas engines using coal gas, producer gas, or blast-furnace gas, are now working for all powers up to 2000 h p per cylinder, and a large number of heavy oil engines are also in use In addition, there are many thousands of petiol motors at work

In internal combustion engines working on the constant volume cycle, the motive power is produced by the explosion of mixtures of inflammable gas or vapour and air in certain proportions, and the pressure produced by the explosion is utilised to propel the piston of an engine, as is done by the steam pressure in a steam engine

Seeing that a proper understanding of the phenomena associated with gaseous explosions is necessary, not only for persons in charge of internal combustion engines, but also for those responsible for the safety of workers both in coal mines and in factories where explosive mixtures may be encountered or used, it is desirable first to consider generally the proparties of such mixtures and the mode in which flame is propagated through them. At the

Hesse's expression (J. pr Chem. [ii ] 57, outset, it should be recognised (1) that, under given conditions of temperature and pressure, the "range of inflammability" of mixtures of any particular combustible gas and air (or oxygen) may be strictly defined, so that all mixtures outside of certain 'lower' and 'upper' limits are non-inflammable; (2) that the ignition of any given inflammable mixture at a particular pressure is determined by the attainment of a certain definable thermal (or electrical) condition at some point or region in it; (3) that after ignition the speed and mode of propagation of flame through the mixture are usually capable of measurement or analysis by well-established experimental methods, and (4) that when such a mixture is inflamed in a closed vessel, the maximum pressure attained on explosion may be accurately determined. But notwithstanding the truth of this statement, it is needful to warn the reader against errors involved in the loose employment of such terms as 'unition temperature,' speed of inflammation or flame propagation,' pressure of explosion,' etc., unless the concomitant physical cucumstances are simultaneously defined

#### A -IGNITION PHENOMENA

Chemical change may be determined in a gaseous explosive mixture at a much lower temperature than is required to develop flame Thus, for example, if electrolytic gas be heated in an enclosure to a temperature somewhat higher than 400°, the formation of steam can usually be detected after a lapse of a few On gradually raising the temperature the rate of interaction between the hydrogen and oxygen rapidly increases until at length a point would be reached when the rate at which heat is lost from the system (eg by conduction through the walls of the enclosure) is more than counterbalanced by the rate at which it is developed by the interaction. The reaction would then become self-propellant

The temperature at which such autogenous (or self-propellant) reaction begins in a given gaseous mixture is usually termed its temperature' or 'syntion point,' although it is not ordinarily a well defined physical constant, being dependent, for example, upon the manner in which heat is conveyed to the mixture is clear also that the ignition temperature, as thus defined, does not necessarily correspond with the appearance of flame in the mixture. There is, indeed, usually a definite 'pre-flame period, the duration of which can be controlled, and which varies considerably with the different gas mixtures Only when a mixture is fired by adiabatic compression, and the pre-flame period is negligibly short, can its 'ignition temperature' be regarded as a physical constant and determined with precision. In most cases the ignition temperatures assigned to a series of gas mixtures merely give an approximate index of their relative ignitibilities

The Determination of Ignition Temperatures -Several experimental methods (of which three principal ones will be hereafter described) have been used to determine the so-called 'ignition temperatures' of gases The results naturally differ according to the details of the method adopted, and must therefore be carefully differentiated.

(1) A common method is to pass successive samples of a particular inflammable mixture into a vessel whose temperature is gradually raised until the lowest degree at which flame appears has been found. In this way the 'relative ignitibilities' of a series of gas-air mixtures may be determined, and for such purpose the method is a useful one. Thus, for example, one of us (R V W) has recently determined by 'this method the 'relative ignitibilities' of a series of methane-air mixtures, expressed in each case as the lowest temperatures of the particular enclosure required to inflame each mixture, as follows.

Per cent r cent { 6 0 70 80 90 100 120 130 Temperature 695° 698° 701° 707° 713° 726° 732 It may be noted, in passing, that the most ignitable of these mixtures is not the one (9 45 pe methane) which on combustion gives the greatest heat development per unit volume, but rather those containing a considerable The method does not, howexcess of oxygen ever, disclose the lowest temperature at which a given mixture can be ignited, because the walls of the heated vessel, by inducing chemical combination on their surface, retard the actual appearance of flame

(2) A better method (first employed by H B Dixon and H F Coward, Chem Soc Tians 1909, 98, 514), which to a large extent eliminates the effect of heated surfaces, consists in heating the combustible gas and the air (or oxygen) separately to a known temperature and then allowing them to mingle, increasing the temperature of each gas stream, pari pursu, until their mixture results in flame The ignition temperature ranges of some combustible gases, at atmospheric pressure, as determined by this method, are as follows—

Ignition Range Combustible Gas In o**vyč**en In air Hydrogen 580° 590°C. . 580°-590°C Carbon monoxide . 637° 658° 644°--658° Cyanogen 803°-818° 850°-862° Ethylene 542° 547° 500°-519° 416°-440° 406°-440° Acetylene 346°-379° Hydrogen sulphide 220°-235° . 556°-700° 650°-750° Methane . 520°-630° 520°-630° Ethane.

These figures record for a given gas the lowest temperature at which any mixture of it with air or oxygen would inflame. But they afford no information as to the effect of varying the proportions of the gas and air.

(3) The temperature produced by the adiabatic compression of a gaseous explosive mixture within a closed cylinder may be accurately calculated from the fundamental gas laws, and this method has been successfully employed by H. B. Dixon and his co-workers for determining the 'ignition temperatures' of various mixtures of hydrogen and oxygen, as follows.—

Mixture			Igni dete	tion Temperature rmined by Adiabat Compression
$\substack{2 H_2 + O_2 \\ 2 H_2 + 2 O_2 \\ 2 H_2 + 8 O_2 \\ 2 H_2 + 16 O}$	:	:	•	. 526°C. . 511° 478° 472°

According to Dixon and Croft's determinations by this method of the ignition-points of mixtures containing electrolytic gas, whereas successive additions of hydrogen or nitrogen progressively raise the ignition temperature of the undiluted gas by regular increments, as would be supposed, successive additions of oxygen, on the other hand, lower it, as shown in the following table—

THE IGNITION-POINTS OF MIXTURES CONTAINING ELECTROLYTIC GAS BY ADIABATIC COMPRESSION

(By H. B. Dixon and J. M. Crofts, 1914)

The observed raising effects of successive dilutions with hydrogen and nitrogen call for no comment, save that the relative greater effect of hydrogen, as compared with nitrogen, may be attributed to its greater thermal conductivity, but the lowering effect of oxygen is indeed puzzling, and its meaning can only be conjec-Dixon and Crofts have suggested that it may be due either to the formation of some active polymende of oxygen under the experimental conditions (which seems doubtful), or that the concentration of oxygen in some way or other brings about increased ionisation of the This at once raises the larger combustible gas question of whether or not ignition is a purely thermal problem, as until recently has generally been supposed

Ignition by adiabatic compression is utilised in the Diesel oil engine, and in gas engine practice pre-ignition of the charge may take place during the compression stroke it too high a compression is employed. Knowledge of the degree of compression required to cause the ignition of a particular gaseous mixture is, therefore, of considerable industrial value.

Ignition by a Source of Heat of Short Duration This is a very common way of igniting explosive mixtures in everyday practice, although up to the present experimental data regarding it are very meagre. It will be realised that if the inflammation of a large volume of gas is attempted by rapidly heating a small portion of it, the size and duration of the source of heat employed, as well as its temperature, will be of importance, because the effectiveness of any means of ignition depends upon whether or not the volume of mixture first inflamed communicates by its combustion sufficient heat to the adjacent layer of mixture to inflame it in its turn In general, the shorter the duration of the source of heat employed to ignite a given gas mixture, the higher must be its temperature. Unfortunately, until now this aspect of the question has been but little explored, but the point may be illustrated by some recent determinations of the 'relative ignitibilities' of mixtures of various gases with air expressed in the terms of the lowest temperatures at which a hot body of minute area must be raised in order to cause immediate ignition of the particular mixture to which it was momentarily applied. Such temperatures have obviously only an empirical interest.

> Minimum temperature of hot body required for Gas immediate ignition by (Mixtures with air) momentary contact 747°C. Hydrogen 931° Carbon monoxide 1000° Ethylene 878° Coal gas 995° Petrol (Fraction 0-80°C.)  $1062^{\circ}$ Benzene 1033° Ether .

Ignition by Electric Sparks .- It is still a matter of controversy whether ignition by means of an electric spark is a thermal phenomenon, or whether there is simultaneous ionization of the mixture by the spark, the ionization playing a necessary and important part in promoting the combustion of the gases So far as the secondary discharge is concerned, ionization of the mixture between the electrodes undoubtedly precedes the passage of the discharge, and it is possible that this preliminary ionization may render the molecules more reactive than usual towards each other It is known that the exposure of some gaseous mixtures to a source of intense ionization causes their combination at quite low temperatures. The point of controversy would, therefore, appear to be whether in the ignition of gaseous mixtures by an electric spark ionic rather than thermal phenomena predominate; whether, in fact, an electric spark is more effective as an ionizer than is a flame or a heated solid.

Insufficient experimental work has been done to enable a decision to be reached on this matter. The character of the phenomena of spark ignition seems to differ from that of the phenomena of ignition by the prolonged contact of a heated surface Thus, the type of curve that can be constructed connecting 'igniting-current' or 'spark-energy' with percentage composition of mixtures of any given inflammable gas with air differs from the type of curve connecting ignition temperatures, as determined by the first method described, with percentage composition (see Fig. 1). It may be, however, that this difference is due to a large extent, if not entirely, to the short duration and small size of the electric spark; also that the rapidity with which flame can travel in a mixture away from the source of heat becomes of importance equally with the susceptibility of the mixture to ignition when the source of ignition is very small and lasts but a short time. It may be noted that the spark ignition curve in Fig. 1 is nearly the inverse of the speed-percentage curve for flame in methane-air mixtures described in the ensuing section of this article (see Fig. 2).

Limits of Inflammability.—If the whole volume of a gaseous mixture be raised to its ignition-temperature, as it is, for example, in the cylinder of a Diesel engine, it necessarily follows that the whole of it is inflamed. On the other hand, the introduction into a gaseous mixture of a small source of heat at a tempera-

mixtures of various gases with air expressed in ture considerably higher than the ignitionthe terms of the lowest temperatures at which temperature of the mixture need not cause a hot body of minute area must be raised in complete propagation of flame.

It is well known that when a flame is introduced into a mixture of an inflammable gas and air containing but little inflammable gas, a 'cap' or aureole is formed around the flame; and that this cap is larger the greater the proportion of inflammable gas present. The coalminer regularly uses this formation of a cap as an indication of the presence of firedamp in the workings of the mine It has been observed also that in a current of gas and air, poor in combustible gas, passing a lamp flame or similar source of heat, the cap or aureole may become separated from the burning body by which it is originated and be carried along for a limited distance, giving a 'cloud' of flame But unless this cloud were fed by some mixture richer in combustible gas than that in which it originated it would gradually die away a uniform mixture of gas and air the flame could only be carried on automatically throughout the extent of the mixture if the inflammable gas were present in quantity above what is termed the 'lower limit' of inflammability.

When a source of heat is introduced into a stagnant mixture of combustible gas and air, two things are necessary to ensure propagation of flame throughout that mixture, namely, (a) an adequate means of ignition, and (b) a sufficient

proportion of inflammable gas

Supposing that too small a proportion of combustible gas is present, only a small quantity of heat per unit volume of mixture is generated when the layer surrounding the initial source of heat is inflamed, and the products of combustion have to impart heat to a considerable volume of mert gases. The number of collisions between molecules of combustible gas and of oxygen that are chemically effective is therefore small. Such collisions, resulting in combination, will occur only in the neighbourhood of the mitial source of heat around which an aureole will form of a size dependent on the nature and quantity of the combustible gas present.

As the proportion of combustible gas is increased a proportionately greater quantity of heat is evolved per unit volume of mixture, and a smaller quantity of inert gases is present to absorb it. At length a point is reached when the amount of heat contained in the products of combustion of any given layer is just sufficient to raise the adjacent layer to its ignition temperature. Flame is then propagated progressively throughout the mixture without any necessity for the continued presence of the source of heat which started the inflammation, and the mixture is said to 'inflame' or 'explode,' according to the rapidity of the propagation.

To ensure propagation of flame throughout the mixture, therefore, it is necessary for the initial source of heat to be of a volume, intensity, and duration sufficient to raise the layer of gases surrounding it to the ignition temperature of the mixture, and for the heat contained in the products of combustion of this first layer to be sufficient to raise the next layer to its ignition temperature, and so on. An inflammable mixture can be defined as a mixture in which flame can spread to any distance, independently c., and away from, the original source of ignition

under given physical conditions, certain limits of composition within (but not outside of), which self-propagation of flame will take place after ignition has once been effected. These limits, defined in terms of the percentage of the combustible gas present, are usually referred to as the 'lower' and 'higher' limits of inflammability of the gas under the given conditions. For any particular gas they differ slightly with the position of the source of ignition, since the progress of the flame may be assisted or 'retarded' by convection-currents dependent upon whether it has to pass in an unward or downward direction. For example, with mixtures of methane and air and of acetone vapour and air the limits, expressed as percentages by volume of the mixtures, are as follows for upward, downward, and horizontal propagation of flame :-

Lower limit

Methane . Acetone vapour	5 40	Downward 6 00 2 35	Horizontal 5 40 2.20
7		Higher Limit	

Downward Upward Horizontal Methane 13 40 14 30 8 50 9 50 Acetone vapour . 9 70

The limits of inflammability of a number of gases and vapours, mixed with air at atmospheric temperature and pressure, are given in the table that follows. For all but hydrogen and carbon monoxide the criterion of inflammability adopted for the determinations was propagation of flame throughout a 21-litre sphere, ignition being at the centre. The figures for hydrogen and carbon monoxide refer to upward propagation of flame.

Gas		Lower limit	Upper limit
Hydrogen Carbon mond	oxide .	рс. . 41 . 125	71 5 73 0
Methane .		. 56	14 8
Ethane .		3 1	10 7
Propane .		2 2	7 4
Butane .		. 17	57
Pentane		. 1.4	4 5
Ethyleue .		3.0	$22\ 0$
Acctylene .		. 3.3	$52\ 3$

Vapour	Lower limit	Upper limi
Benzene	p.c. 1.5	рс 5 б
Toluene	14	54
Ethyl alcohol .	$4 \cdot 4$	
Methyl alcohol .	. 6.5	24.0
Ethyl ether	. 20	5.0
Acetone	. 225	96
Carbon disulphide	$.~~4\cdot 2$	
Petroleum benzme	. 1.1	38
Gasoline	. 1.5	5.3

The Effect of Temperature.-From what has been said in the preceding paragraph, it is evident that the self-propagation of flame through a mixture is, in principle, dependent on the temperature that the portion of the mixture which is burning can impart to the portion which is about to burn, from which it follows that the higher the initial temperature of the

In the case of mixtures of any particular | mixture the less dependent is the propagation combustible gas with air (or oxygen) there are, | of flame upon the heat generated by combustion Hence the effect of increasing the initial temperature of mixtures of an inflammable gas and air should be to widen the range of inflammability, lowering the lower, and raising the higher

> Thus it has been found experimentally that a progressive increase in the initial temperature of mixtures of methane and air from 20° to 700° caused a corresponding decrease in the amount of methane required to enable self-propagation of flame to take place from 6 00 to 3 25 p c (downward propagation) Similarly, the higher limit was raised gradually from 13 40 p.c at 20° to 16 40 at 600°

The Effect of Pressure —As exemplifying the influence of the initial pressure of gaseous mixtures upon their limits of inflammability, the case of mixtures of methane and an may be cited. A reduction of the initial pressure, at 20°C, from 760 to 380 mm has but little effect on the limits; but below a pressure of 300 mm mercury the effect is marked, the limits narrowing rapidly until, at a pressure of 120 mm no mixture of methane and air is capable of selfpropagation of flame (downwards). On the other hand, an increase of pressure above 760 mm slightly raises both the lower and the higher limit, the figures at 6 atmospheres being 6 40 and 14 05 p c of methane respectively, for downward propagation of flame

The lower limit with mixtures of hydrogen and air, and of carbon monoxide and air, is also raised by increasing the initial pressure above 760 mm, but with these two gases the higher

limit is lowered

Mixtures of Inflammable Gases -In the case of a mixture of inflammable gases such as coal gas, a general formula is available by means of which its limits of inflammability with air may be calculated if its composition, and the corresponding limits of the individual gases are known If a, b, c, . are the relative proportions of its components, and A, B, C, their respective limits of inflammability, then L, the limit for the mixed gases, is given by the equation:-

$$L = \frac{a+b+c+\dots}{a-b-c} \qquad (1)$$

This relationship holds both at the lower and the higher limit. It implies that if a limitmixture with air of one inflammable gas is mixed in any proportion with a limit-mixture with air of another inflammable gas, a limit-mixture results

It has been necessary to emphasize the fact that this formula refers to mixtures of combustible gases with air, for the reason that the presence of any considerable quantity of an uninflammable constituent, such as nitiogen, in the commercial gas would narrow the range of inflammability, just as the range for a pure gas is nairowed if the oxygen-content of the air is reduced.

The effect of Incombustible Gases -The effect of a reduced oxygen-content of the 'air,' or of the presence of inert constituents in the 'combustible gas,' is more marked at the higher than at the lower limit. This is well illustrated

methane with 'atmospheres' containing different proportions of oxygen. These determinations afford an index of the general effect of 'mert' gases on the limits of inflammable gases in general when the actual data are not available. the effect of carbon dioxide is rather more pronounced than that of nitrogen.

Atmosphere		Methane-p c			
Oxygen	Nitrogen	Lower limit	Higher limit		
20 90	79 10 (an)	5 60	14 82		
$19\ 22$	80 78		12.93		
18 30	81 70	-	11.91		
17.00	85 00	5 80	10 55		
15 82	84 18	5.83	8 96		
14 86	85 14	6 15	8 36		
13 90	86 10	6.35	7 26		
13 45	86 55	6 50	6 70		
$13\ 25$	86.75	No mixture	e capable of		
		propagating flame			

Speed of Propagation of Flame —During the propagation of flame in any 'limit' mixture, a balance is struck between heat generated by combustion and heat employed in starting combustion, together with heat lost by conduction Theoretically, therefore, proand radiation vided that the amount of energy imparted to the system by the initial source of ignition is small, so that no appreciable impetus to the propagation of flame occurs near the source of ignition, flame should travel in a limit-mixture at a uniform speed. This is found to be so in practice, the speed of the flame is slower in a limit mixture for downward than in one for upward propagation, but under either condition it is quite uniform throughout the extent of the mixture

It is otherwise when the proportion of inflammable gas lies at some intermediate point between the limits, that is to say, when the mixture is such that one layer on burning is capable of imparting to the uninflamed layer adjacent to it a temperature higher than the ignition-temperature of the mixture. The phenomena of propagation of flame in such mixtures are dependent in a marked degree upon the conditions under which ignition is effected, so that it is necessary when speaking of the propagation of flame through them to specify exactly those conditions

When ignition of a mixture of combustible gas and air has been effected at the open end of a long tube which is closed at the other end, the flame can be propagated in two principal ways, namely, (a) by transmission (by radiation, conduction, and convection) of the heat developed by the burning of one layer of gas to the layer adjacent to it, and (b) by the transmission of a sufficiently high pressure

According to the first mode of propagation, the flame travels along the tube at a uniform speed for a certain distance, and then develops vibrations, swinging backward and forward with oscillations of increasing amplitude. The mean speed of the flame during this vibratory movement, which comprises a series of compression and rarefaction waves, is usually greater than the speed of the uniform movement. According to the nature of the mixture, either the vibrations may continue to the end of the productive of mechanical disturbance of the tube, or the flame may become extinguished mixture. The speed of the uniform movement

by determinations of the limits for mixtures of during a particularly violent vibration (eg. during rarefaction) Or again, a sudden permanently greater enhanced speed of flame may be developed as the result of the second mode of propagation coming into play (during a compression).

During this second mode of propagation cach layer of gas is compressed so suddenly that it is raised beyond its ignition temperature by the heat of compression, and in burning it compresses in turn the unburnt layer in front As already stated, the speed of the flame during this mode of propagation is permanently enhanced, it is, in fact, the maximum speed attainable by the flame and is a physical constant for each mixture of gases The phenomenon is termed the 'detonation wave'

When ignition takes place at the closed end of a tube open at the other end, the flame travels at a rapidly increasing speed towards the open end If the tube is long enough, the flame may eventually assume a vibratory character and the detonation-wave may be developed.

What has been said in this short summary refers only to mixtures of a combustible gas with air, mixtures with pure oxygent behave somewhat differently With regard to the latter, it may be stated that, in general, (a) the preliminary period of uniform movement is shorter, (b) a rapid acceleration follows, and (c) this is immediately succeeded by the detonation-wave without any vibratory movement taking place


Of the different phases in the propagation of flame in mixtures of gases, the two regarding which most information is available are the 'uniform movement' and the 'detonation uave.'

Some further account of these will now be given.

The Uniform Movement—The initial slow propagation of flame that takes place when an inflammable mixture at rest is ignited at a point, is usually regarded as controlled by the transference, by conduction, of the heat developed by the combustion of the mixture immediately surrounding the point of ignition, whereby successive contiguous portions of the mixture are raised in temperature until chemical action becomes rapid The initial speed of propagation of flame in a given mixture away from the point of ignition should mainly depend, according to this view (a) on the conductivity for heat of the unbuint mixture, and (b) on the velocity with which a moderately heated layer begins to react chemically and so to use gradually in temperature, or, in other words, on the rate of change of reaction velocity with temperature

Under certain conditions, with all inflammable mixtures of gases and air at atmospheric temperature and pressure, the initial slow propagation of flame can be maintained at a uniform speed over a considerable distance of travel from the point of ignition. The conditions most favourable, or perhaps even necessary, to initiate and maintain this 'uniform movement' of flame are (a) that the inflammable mixture should be contained in a long straight tube open at one end and closed at the other; and (b) that ignition should be at the open end by a source of heat not greatly exceeding in temperature the ignition temperature of the mixture, and not then depends on the composition of the mixture (presumed to be at atmospheric temperature) for the umform movement of flame in mixtures and pressure), and on the diameter of the tube of some of the common combustible gases with in which it is contained

With a tube of a given diameter the speed



of the uniform movement of flame in a mixture can be regarded as a definite physical constant for that mixture, and its measurement is of considerable value in enabling a judgment to be formed of the behaviour of the mixture under normal conditions of burning

In Fig. 2 are shown speed-percentage curves air over the whole range of inflammability for each gas All the measurements were made in a tube of 25 cm internal diameter, so that they are strictly comparable one with another Apart from the striking differences in the speeds obtainable in different gases displayed by this diagram, a noticeable feature of the curves is the fact that with each gas the maximum speed is obtained not, as might perhaps be expected, with the theoretical mixture for complete combustion, but over a range of mixtures containing an excess of inflammable gas.

The extent to which the speed of the uniform movement is affected by the size of the tube along which the flame travels has been determined for mixtures of methane and air results are recorded in Fig. 3 For all practical purposes it can be assumed that the speed of the uniform movement of flame in mixtures with air of gases other than methane will be affected to a similar degree, so that the absolute speeds of the flames in pipes 30, 60, and 90 cm in diameter will be respectively about 25, 3, and 35 times as great as the speeds recorded in the 25 cm tube (Fig. 2)

It should be noted that the uniform speeds of the flames in tubes of large diameter (greater than 10 cm for methane-air mixtures) are not, strictly speaking, those of the 'uniform morement' as previously defined, since they are not controlled solely by conduction of heat. A




Fig. 2.

second factor, namely, convection, comes into uniform movement of flame for any complex play when the tubes are large, and is responsible gas mixed with an, from a knowledge of its for the enhanced speed of flame.

found that a similar expression to that used for calculating the limits of inflammability of mixed combustible gases can be employed to determine, with fair accuracy, the maximum speed of


composition and the maximum speeds of each Mixtures of Inflammable Gases -It has been of the components. The expression is -

$$S = \frac{a+b+c+}{a+b+c+} . \quad (11)$$

$$S_a + S_b + S_c + . . .$$

the proportions of the different simple consti-tuents in the complex inflammable gas (coal-gas, for example); and S<sub>2</sub>, S<sub>b</sub>, S<sub>c</sub> . . . the the composition of the mixture of the complex

in which S is the speed required; a, b, c . . . maximum speeds of flame in mixtures of the




gas and air in which the speed of the uniform | curves for the uniform movement of flame in movement of flame will be greatest, and mixtures of methane with nitrogen-oxygen formula (11) for determining the speed of flame in the limit-mixtures, and (though with rather less accuracy) in the mixture containing the gases in theoretical proportions for complete combustion.

The value of these formulæ will be appreciated if reference be made to Fig 2, wherein are recorded, for comparison with the simple gas-air mixtures, the speeds of the uniform movement of flame in mixtures of the composite gases CH<sub>4</sub>+H<sub>2</sub> and 3CH<sub>4</sub>+H<sub>2</sub> with air as determined by actual measurement in a tube 25 cm. in diameter. It will be seen that the gas with the lower speed of flame and the smaller range of inflammability (methane) has the predominating influence in deciding the magnitude of these constants for the mixed gas

Effect of Incombustible Gases —In addition to restricting the limits of inflammability, the presence of an incombustible gas mixed with the combustible gas has the effect of reducing the speed of flame in any of the mixtures with air. This effect is illustrated in Fig. 4, which shows the speed-percentage

mixtures containing less oxygen than air, the



curve for air-mixtures being given for comparison. This diagram shows also the effect of nitrogen in reducing the range of inflamma-bility and establishes the fact, theoretically deducible, that the speed of travel of flame is the same in all limit-mixtures that comprise the same constituent gases. The experiments from which these curves are constructed were made in a tube 5 cm in diameter, so that the speeds are not directly comparable with those recorded in Fig. 2.

B .- THE DEVELOPMENT OF 'DETONATION' IN GASEOUS EXPLOSIONS THE EXPLOSIVE WAVE AND 'RETONATION'-RATIO OF EX-PLOSTON

It was in the year 1881 that Berthelot and Vieille, and independently also Mallard and Le Chatelier, announced the discovery of the rapid acceleration of the initial velocity of inflamma-

tion in the progress of gaseous explosions and the final attainment of the enormously higher constant velocity of the 'explosion wave.' Since that date, the phenomena associated with the development of 'detonation' and the propagation of the ex-plosion wave has been extensively studied in this country by H. B Dixon and his co-workers

Ma'lard and Le Chatelier devised a photographic method for investigating the initial phases of gaseous explosion, and the development of detonation, which consisted in recording on a sentitised plate, moving at a known velocity on a vertical plane, the movement of the flame as it travelled along a horizontal glass tube A graph compounded of the two velocities was thus obtained

H B Dixon's experimental method consisted in photographing the explosion flame travelling along a horizontal tube on a highly sensitive film rotated vertically with a constant high velocity (varying, however, between twenty-five and fifty metres per second in different experiments), the explosion tube being placed at such a distance from the camera that the size of the image was about onethirtieth that of the flame In this way it was found possible to analyse the progress of an explosion from its point of origin up to the final attain-

ment of its maximum force and velo-city in 'detonation.' The investigation also included the discovery of the wave of 'retonation,' which is thrown back through the still burning gases from the point where detonation starts (a phenomenon also discovered by Le Chatelier independently in 1900), of the effects of collision between two explosion waves, and of the passage of reflected waves through the hot

products of explosion.

The phenomena associated with the development of an explosion in a gaseous mixture, fired in a closed tube by a spark passed between wires a few inches from the closed end, are very clearly shown in the photograph reproduced in Fig 5 which is analysed in Fig. 6 This photograph was taken during an experiment in which carbon disulphide was exploded with a quantity of oxygen represented by the expression  $CS_2+5O_2$ . It is reinforced by another reflected wave) the The flame in starting at the point o, sends or velocity of the retonation wave is always

mysible compression waves in both directions along the tube, which travel in advance of the tlame with the velocity of sound through the unburnt gases, as represented by the dotted lines om, on in the diagram. The flame itself, travelling at first more slowly than the compression waves, traces the curves on and ob. The compression wave on, on reaching the closed end of the tube, is reflected back again as NC, and, on meeting the flame (which is still travelling in the direction OA), retaids it, and passes thence through the hot and probably still burning gases as the visible wave co instant later it overtakes at D the front of the flame, travelling in the direction on, thereby accelerating it and increasing its luminosity in






FIG. 5.

consequence of the quickened combustion. flame then continues to move forward with rapidly accelerated velocity until 'detonation' is set up at the point E At this point a strongly luminous wave of compression EG (the 'retonation wave') is sent backwards through the still burning gases, which, on reaching the near end of the tube, is reflected back again as GH The detonation 'wave' EF passes onward through the mixture with its characteristic uniform high velocity and intense luminosity Fig. 7 is a similar photograph showing the development of 'detonation,' and the phenomenon of 'retonation' in the case of mixture of cyanogen and oxygen  $(C_2N_2+O_2)$  fired near the closed end of a tube.

Except in special circumstances (e.g. when it is reinforced by another reflected wave) the

inferior to that of detonation, thus Le Chatelier gives 2990 metres per second for the 'detonation wave' and 2330 metres per second for the 'retonation wave' in an equimolecular mixture of acetylene and oxygen When, however, the 'retonation wave 'is developed just at the closed end of a tube (e.g. when the explosive mixture is fired at such a distance from the closed end that 'detonation' is set up just as the flame arrives at the end) it may be reinforced by a reflected wave, in which case its velocity cannot be distinguished from that of a true 'detonation

The explanation of the intense luminosity of the retonation wave, and its higher velocity than sound through the exploded gases, is to be found in the fact that the combustion during the initial stages of an explosion is very much




Fig. 7

combustion is much more prolonged than in detonation, and at the moment when detonation is set up combustion is still proceeding in the layers of gas for some distance behind the flame-front The 'retonation' wave, in passing backward through these layers, quickens this residual combustion, and is itself thereby rendered highly luminous This interthereby rendered highly luminous. This inter-pretation is supported by the repeated observations of Dixon, that the collision of two flames, in neither of which 'detonation' has been determined, will frequently give rise to reflected waves more rapid and more luminous than the There can be little doubt as incident waves to the important part played by reflected waves in determining the violent shattering effects associated with gaseous explosions on a large scale.

The Explosion Wave -Berthelot and Vieille, in announcing their discovery of the develop-

ment of detonation '(l'onde explosive') in gaseous explosions, described it as "une certaine surface régulaire, où se développe la transformation, et qui réalise un même état de combinaison, de température, de pressure, etc Cette surface, une fois produite, se propage ensuite, de couche en couche, dans la masse tout entière, par suite de la transmission des chocs successifs des molécules gazeuses amenées à un état vibratoire plus intense en raison de la chaleur dégagée dans leur combinaison, et transformées sur place, ou, plus exactement, avec un faible déplacement relati. ' Such conditions are comparable with those of a sound wave passing through the gaseous mixture, with, however, the important difference that, whereas a sound wave is propagated from layer to layer with a small compression and a velocity determined solely by the physical condition of the vibrating medium, it is an abrupt change in chemical condition which is propagated in the explosion wave, and which generates an enormous force as it passes through each successive layer of the medium.

In their experimental work Berthelot and Vieille proved that the velocity of the explosion wave is independent of the length of the column of gas traversed, and of the material or diameter of the tube employed (at least above a certain small limiting diameter) It is also immaterial whether the tube is laid out straight, coiled round a drum, or even zigzagged. They also concluded that the velocity is independent of the initial pressure; but this is not strictly correct, as H B Dixon has since shown. The rate increases slightly with the initial pressure, attaining a nearly constant value at a pressure of about two atmospheres

It must be understood that when detonation is once determined in a particular explosive mixture, the wave is propagated with constant velocity (i.e. for a given initial temperature and pressure), which velocity is indeed an important physical constant for the particular mixture. I'he following 'rates of explosion,' as determined by Dixon, may be considered as typical of the extremely high velocities attained .

RATES OF EXPLOSION FOR MIXTURES FIRED AT 10° AND 760 MM.

Mixture			M	Ictres	per second
$2H_{2}+O_{2}$					2820
$H_2 + Cl_2$		•			1729
$2CO + O_2$	(moist,	sat'd	at 28)		1703
$NH_3+3O$	٠ ، (		• ′		2390
$CH_4+2O$		•			2322

In detonation, each successive layer of the explosive mixture is fired by compression, and the chemical change in the wave occurs under strictly adiabatic conditions This means that the chemical action is more intense and of much shorter duration than during the initial phases of an explosion ('inflammation'); moreover, the pressures developed in the wave and the consequent shattering effects are much greater than in ordinary explosions. Thus, for example, in a mixture  $2H_2+O_2$ , fired at the ordinary atmospheric temperature and pressure, the initial velocity of the flame (i.e during the short period of 'inflammation') is only 20 metres per second, and chemical action endures in each uccessive layer of the mixture for as long as

3700°

13 750

7 641

3400°

13 102

7 608

as detonation is set up, the flame travels forward pression, and the abrupt variation in the with a constant velocity of 2820 metres per density and pressure of the medium is due to second, and the duration of chemical action in the chemical change. Chapman's formula for  $\frac{1}{5000}$  and possibly not much less than  $\frac{1}{10000}$  of a second. Needless to say 'detonation,' is not set up in a gas-engine cylinder; the terrific force developed in detonation would play havoc with the mechanism

Dixon's photographic researches, already alluded to, have demonstrated beyond doubt the important rôle of 'compression waves' in determining detonation in explosive mixtures, and it is probable that the disastious effect of colliery explosions are primarily due to the intensifying effects of reflected compression waves upon the burning mixture of gas and (oi) coal dust and air

In 1899 D L Chapman published a mathematical analysis of the explosion wave carding Berthelot's sound-wave theory, which indeed had led to values for rates of explosion which were difficult to reconcile with facts, he regards the explosion wave as a wave of compression not in a homogeneous medium, but in a medi m which is discontinuous in the vicinity of the wave-front It is assumed (1) that the · front ' of the wave (i e from the unexploded gas to the point of maximum pressure) does not alter in character, or, in other words, that every portion of the wave travels with the same velocity, (2) that the velocity is the minimum consistent with (1), and (3) that at the point of maximum pressure the chemical change concerned in the propagation of the wave is complete

Temperature .	$4300^{\circ}$	4000°
Cv steam .	14 750	$14\ 297$
Diatomic gases	7 707	7.674

With the aid of this series of numbers he proceeded to apply his formula to the calculations of the rates of explosions of some forty other mixtures investigated by Dixon, finding in all cases close agreement between the found and calculated values, of which the following may suffice as examples -

Mixture exploded	Products in the wave	Temp	Rate of sion I per — Calcu- lated	Metren
$2H_2 + 2N_2O$	$2H_{2}O + 2N_{2}$	3813°	2408	2305
$4H_{2} + 2N_{2}O$	$2H_{2}O+2N_{2}+2H_{2}$	3077	2604	2545
6H2+2N2O	$2H_{2}O + 2N_{2} + 4H_{2}$	2612	2720	2705
$2H_2 + 2N_2O + 2N_2$	$2H_{2}O + 4N_{2}$	3077	2097	1991
$C_2 II_4 + 2O_2$	2CO+2H2O	4365	2619	2581
$C_2H_4 + 3O_2$	$2CO + 2H_2O + O_2$	3882	<b>234</b> 8	2368
$C_2H_2+O_2$	2CO + H <sub>2</sub>	5029	3101	2961
$CH_4 + O_2$	$CO + H_2O + H_2$	2772	2502	2528
$2CH_4 + 3O_2$	$2CO + 4H_2O$	3764	2485	2470
$2CH_4 + 3O_2 + N_2$	$2CO + 4H_2O + N_2$	3513	2353	2349
1	ì	j		1

A characteristic feature of detonation is the extremely short duration of chemical action and subsequent rapid cooling of the products, as compared with ordinary combustion Summ

one-twentieth of a second. As soon, however, I front of the wave are, of course, fired by comeach successive layer certainly does not exceed, the velocity of the explosion wave in centimetres per second is-

when R=the gas constant (1985), J=the dynamical equivalent of heat (42+10° crgs), μ=the gram equivalents of the mixture exploded, n and m = the number of gaseous molecules before and after the chemical change in the wave, Cp and Cv=the mean specific heats of the products at constant pressure and volume respectively, h=the total heat generated in the wave, and  $t_0$  = the initial temperature (abs.) of the mixture exploded

From the fact that the dilution of electrolytic gas (2H<sub>2</sub>+O<sub>2</sub>) with oxygen lowers its rate of explosion a little more than a corresponding dilution with nitrogen, Chapman considers it improbable that there is any appreciable dissociation of steam in the wave He assumes that the molecular heat of steam rises more rapidly with temperature than that of a diatomic gas, and that the molecular heats of oxygen, hydrogen, nitrogen, and carbon monoxide may for all practical purposes be considered as equal at any given temperature Selecting some seventeen of Dixon's found rates of explosion, he has calculated by means of his formula the corresponding molecular heats and temperatures, arriving at the following results for Cv at inter-The unbuint gases immediately in mediate temperatures by interpolation .-

3100°

12 250

7 575

2800°

11 040

7 542

2500°

9 797

7 509

years ago, W. A Bone and B Lean found by a photographic method that the duration of luminosity in each successive layer of gas in the detonation of electrolytic gas is certainly less than boloo second, a much shorter interval of time than was required to shatter a tube of thin glass attached to the end of the explosion coll used This tube, although invariably smashed by the force of the explosion, always appeared perfectly intact in the photograph. Dixon's subsequent photographic researches have demonstrated the abrupt suddenness with which the gases attain the maximum temperature in detonation, the intensity and short duration of luminosity, and the subsequent rapid cooling, as compared with ordinary combustion over, high as is the temperature attained, there is no evidence of any considerable dissociation of steam in the wave, for, despite the instantaneous cooling of the products, there is less than 1 per cent of the gases left uncombined

gas  $(2H_2 + O_2)$ Influence of an Excess of an Inert Gas upon the Rate of Explosion of a particular Mixture -It has been found experimentally that whereas an excess of nitrogen or oxygen retards the rate of explosion of a particular mixture, an excess of hydrogen actually increases it. Thus in the

after the wave has passed through electrolytic

case of electrolytic gas, we have-

Mixture exploded			
			2817
			3268
			3527
			3532
			2328
			1927
			2426
			2055
	:	: : :	: .

The accelerating influence of an excess of hydrogen was for long an enigma, but it was perfectly explained when Chapman published his formula. It must, however, be understood that an excess of oxygen always has a somewhat greater influence than a corresponding excess of nitrogen by virtue of its relatively higher density

The Burning of Guseous Carbon—Of the many important facts brought to light during the course of Dixon's investigations none are of greater interest than those relating to the burning of gaseous carbon in the explosion wave, as illustrated by the case of cyanogen. In view of the fact that the molecular heat of combustion of cyanogen, when burnt completely to carbon dioxide, is 259 6 K C Us, whereas, if buint to the monoxide, it would only be 123 K C Us, it might be expected that the rate of explosion for a mixture  $C_2N_2+2O_2$  would be much higher than for  $C_2N_2+O_2$ , if gaseous carbon is primarily burnt to carbon dioxide in the wave. The exact opposite is the case, however, as the following results show.—

$$\begin{array}{ccc} \text{Mixture} & \text{Products} & \text{Rate} \\ C_2N_2+O_2 & 2CO+N_2 & 2728 \\ C_2N_2+2O_2 & 2CO_2+N_2 & 2321 \\ \end{array} \\ \text{metres per sec.}$$

Still more cogent is the evidence in favour of the initial formation of carbon monoxide in the wave, when the following figures are considered—

$$\begin{array}{cccc} & C_2N_2 + O_2 & & C_2N_2 + O_2 + N_2 & & C_2N_2 + O_2 + O_2 \\ \textbf{Rate} & 2728 & & 2398 & & 2321 \end{array}$$

The conclusion to be drawn from the preceding figures is that cyanogen is initially burnt to carbon monoxide and nitrogen in the wave itself, any excess of oxygen afterwards burning up the carbon monoxide as the gases cool down in the rear of the wave

This conclusion was driven home by Dixon, in conjunction with Strange and Graham, by photographing on a sensitive film rotated at a speed of about 1500 metres per minute, the explosion flame in the case of the three mixtures (a)  $C_2N_2+O_2$ , (b)  $C_2N_2+O_2+N_2$ , and (c)  $C_2N_2+2O_2$ . In each case the flame was photographed, after detonation had been set up, as it dashed past a glass window inserted into the lead explosion coil <sup>2</sup>. The image obtained in the case of (a) showed an intensely brilliant flame (the explosion wave), slightly drawn out in tapering form; in the case of (b), with nitrogen as diluent, the flame was less brilliant and somewhat more drawn out than in (a) but with (c) the flame, whilst no more luminous than in (b), was drawn out to great length, owing to the continued combustion of carbon monoxide in the rear of the wave. The following diagram, Fig. 8 (approximately to scale), will convey an idea of the relative durations of the flames in the three cases.

(a) 
$$MIXTURE - C_2N_2 + 0$$
.

(b)  $MIXTURE C N_2 + 0$ .

(c)  $MIXTURE C N_2 + 0$ .

(c)  $MIXTURE C N_2 + 0$ .

(d)  $MIXTURE C N_2 + 0$ .

Fig. 8.

The Burning of Hydrocarbons in the Wave—
It was during the course of measurements of
the rates of explosion of hydrocarbons with
varying proportions of oxygen in 1891-92 that
Dixon and his collaborators rediscovered facts
(originally set forth by Dalton in his 'New
System of Chemical Philosophy') which finally
disposed of the dogma that in a deficient supply
of oxygen a 'selective' burning of hydrogen
occurs Bone, in conjunction with Lean and
Cain, found that when either ethylene or acetylene
is detonated with its own volume of oxygen the
ultimate products consist almost entirely of
carbon monoxide and hydrogen <sup>1</sup> The rates of
explosion of hydrocarbon-oxygen mixtures,
whilst they do not reveal the real mechanism
of the combustion, show very clearly that,

1 Vide Lean and Bone, Trans. Chem. Soc 1892, 61, 873, and Bone and Cain, ibid 1897, 71, 26.

as in the case of cyanogen, the carbon of a hydrocarbon is burnt to the monoxide in the wave itself, the formation of the dioxide being an after-effect and taking place in the rear of the wave. In the cases of methane and acctylene the fastest rates are observed with equimolecular mixtures, whereas with ethylene the rate increases with the proportion of oxygen up to the limit  $C_2H_4+2O_2$ .

## C.—THE PRESSURES PRODUCED IN GASEOUS EXPLOSIONS.

Many investigators, from the time of Bunsen's well-known experiments in 1867 onwards, have measured the pressures produced in gaseous explosions, with doubtless considerable success so far as what may be termed the mean effective

- Trans. Chem. Soc. 1803, 69, 759

pressures are concerned The explosion vessel employed by Bunsen was a stout glass tube 8 15 cm long, 1.7 cm internal diameter, and of i8 cc capacity It was closed by a suitable valve, the load on which could be adjusted until it was just lifted when the explosive mixture was fired by means of a powerful spark passed along the axis of the tube Bunsen considered that the combustion would occur under adiabatic and 'constant volume' conditions, and he identified the rate of ignition of a particular mixture with that of the completion of chemical change. In calculating from his results the corresponding flame temperatures, he assumed the constancy of the specific heats of steam and carbon dioxide Finding that the pressures recorded in the cases of electrolytic gas and of a mixture of carbon monoxide and oxygen in their combining proportions (namely, 95 and 10.1 atmospheres respectively) were somewhat less than one-half of those theoretically re-

quired on the above assumption, he concluded that in each case combustion had proceeded per saltum, owing to the supposed theoretical flame temperatures exceeding the limits at which steam and carbon dioxide respectively are completely dissociated. The problem was again attacked independently by Mallard and Le Chatcher in 1883, and by Berthelot and Vieille in 1885 2 The last-named fired various gaseous mixtures at atmospheric pressure in a spherical non bomb, measuring the effective pressures by the movement of a light piston working against a spring in a tube attached to the bomb. In order to gain some information respecting the possible cooling influence of the walls upon the effective pressures recorded, three bombs of different capacities, namely, A of 300 ee, B of 1500 cc, and C of 4000 cc, were employed The results, expressed in each case as atmospheres in excess of the atmospheric pressure, were as follows .-

Bomb	Λ	В	C
Capa ity in cm 3 .	300	1500	4000
Surface in cm <sup>2</sup>	216	648	1200
Surface per unit volume	0 72	0 43	0 33
Diameter in cm	6.0	14 2	217
Length of firing piece in cm	6.3	5 3	6 3
Travel of flame before reaching the piston in cm	91	16 3	24 8
	Atms	Atms	Atme
$_{0.73}(2H_{2}+O_{2})$ .	7 41	9 69	9 80, 0
å <u>å</u> (200 + 0.	9 29	9 93	10 21 2
= The control of the	13 94	14 81	10 21 ding 16 31 g
$\exists \mathbb{R}   C_2 \mathbf{N}_2 + O_2 \qquad .$		_	25 11 2
$C_2N_2+2O_2 \qquad . \qquad .$			20 96)
Time required for flame to travel from the firing piece to piston of indicator.			
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	0 00104 sec		$0.00214~\mathrm{sec}$
$100000 \cdot 200 + 0$	0 01286 sec		$0.01551~\mathrm{sec}$

results is the smallness of the difference between the pressures observed with bombs A and C in any particular case. These differences might at first sight be attributed to the (supposed) much smaller cooling influence of the walls of the bomb C as compared with bomb A, but if this were the true explanation the difference should be greater with the slow-burning 200+0, mixture than with the fast-burning  $2H_2 + O_2$ , whereas the opposite was the case Morcover the fact that dilution of the mixture 2H2+O2 with twice its own volume of nitrogen almost obliterated the difference between the pressures observed in the cases of bombs A and C (the ratio  $\frac{pC}{c}$ ). for the diluted being 0.95, as against 0.76 for the undiluted mixture) is all against the cooling theory The more probable explanation is to be found in the fact that, owing to the much longer travel of the flame before it reached the piston of the indicator in bomb C as compared with A, the explosion would be in a more advanced phase of its development, and this would be most marked in the case of the fastest bu.ning mixtures. Berthelot and Vieille assumed

Perhaps the most notable feature about these alts is the smallness of the difference between pressures observed with bombs A and C m particular case. These differences might at the sight be attributed to the (supposed) much slight be attributed to the (supposed) much blief cooling influence of the walls of the bomb is compared with bomb A, but if this were true explanation the difference should be attributed the marked disparity between the observed pressures (bomb (') and those calculated from the heats of combustion (on the assumption of adiabatic conditions) to a rapid increase in the specific heats of steam and carbon dioxide at high temperatures with the fast-burning  $2\Omega + O_2$ , whereas opposite was the case. Moreover the fact

Mallard and Le Chatcher (l c) who used a Bourdon gauge for measuring pressures and applied a 'cooling correction' to their results, agrived at practically the same conclusion as Berthelot and Vieille (Ann Chim. Phys [vi] 4, 13) respecting both the increase of the specific heats of steam and carbon dioxide with temperature and the small if not negligible influence of dissociation. They calculated that the temperature of the flame when moist electrolytic gas is exploded in a closed vessel at atmospheric pressure is about 3350°, and that the mean molecular heat of steam at constant volume

Annales des Mines, sec viii vol 4 Ann. Chim Phys. (vi ) 4, 13

between  $0^{\circ}$  and  $3350^{\circ}$  is 16.6. In the case of REVOLVING DRUM the mixture 2CO+O, they concluded that the molecular heat (C2) of carbon dioxide rises to 136 at 2000°, above which temperature dissociation comes into play Dugald Clerk also made experiments with hydrogen-air mixtures in a large cast-iron vessel comparable in dimensions to the cylinder of a six-horse-power gas engine It was a cast-iron cylinder truly bound out and the end covers turned, internal dimensions 7 inches r manif diameter and 81 Minimin's 2 2 2 2 2 2 1 1 inches long, capacity INDICATOR 317 cubic inches Upon the cover was placed a Richards indicator, in which the reciprocating drum had been replaced by a re-FAN GOVERNOR volving one; the rate of revolution was adjusted by a small fan, a 11/1 weight and gear giving the power evlinder was filled with the explosive mixture to be tested, the drum set in motion, the pencil of the indicator pressed gently against it, and the EXPLOSION VESSEL electric spark passed within the cylin-

a line upon the drum which showed
the use and fall of pressure. The using line | Fig. 9 shows the arrangement of the apparatus traced the progress of the explosion, the falling | Experiments with hydrogen and air mixtures line the progress of loss of pressure by cooling gave the following results

The pressure of the explosion, acting upon the indicator piston, traced

Frg. 9.

## MIXTURES OF HYDROGEN AND AIR (CLERK).

		Tem	p. before explosion 16°.	Pressure 14 7 lbs (atm	ospheric)
Experi- ment	Mıv H	Air	Max pressure above atmosphere in lbs per sq_inch	Time of explosion	Max temp of explosion
	1 vol. 1 vol. 2 vols	6 vols. 4 vols. 5 vols	41 68 80	0 15 sec. 0 026 sec 0 01 sec	826° to 909° 1358° to 1539° 1615° to 1929°


The maximum temperatures were calculated | on the two extreme assumptions of complete contraction and no contraction

The curves of explosion from which these figures are deduced are given at Fig. 10

The rate of cooling was comparatively slow; for instance, in all these experiments, at the end | ing for loss of heat to the walls during time of

of 0 66 second from the beginning of explosion there still remains above 20 lbs per sq inch above atmospheric pressure. The explosion was followed by a very slow fall of pressure

Calculating from these curves the total heat accounted for by maximum pressure, and allow-



Vessel used 7 inches diam, and 84 inches long—Scale of indicator spring, 1 lb. =  $\frac{1}{2}s$  inch. Mixtures used, pure hydrogen and air—Experiment a, 1 vol hydrogen to 6 vols. air; b, 1 to 4; c, 1 to 5 Temperature of gases before ignition, 16°, pressure (atmospheric), 14 7 lbs.

explosion, as estimated from the falling curves, a great deficiency becomes evident.

Experiment Heat of combustion of hydrogen present 100 100 Heat accounted for by maximum temperature and loss previous to its attainment 55 70

Even with hydrogen and air mixtures the deficiency ranges from 30 to 46 p.c. It is remarkable that the deficiency at the temperature of about 900° is identical with that at about 1800° within the limits of error It should be noted, however, that this method of allowing

for heat loss will probably give a low value, as no account is taken of the difference in the action of the walls on the using and falling lines, nor of the possible difference in radiation

In 1915 W A Bone and his collaborators published the results of experiments in which mixtures corresponding to (1)  $2H_1+O_2+4N_2$ , (2)  $2CO+O_2+4N_2$ , and (3)  $CH_4+O_2+4N_2$ , were exploded at an initial pressure of 50 atmospheres m a spherical bomb (capacity=275 cc) to which was attached a Petavel recording manometer with its optical accessories. The results of two typical experiments with each of the three mixtures are recorded in the following table .-

W. A. BONE'S PRESSURE EXPERIMENTS.

Mixture fired	2112+0	) <sub>2</sub> +4N <sub>2</sub>	$\mathrm{CH_4} + \mathrm{C}$	) <sub>2</sub> +4N <sub>2</sub>	2CO+	()2+4N2
Per cent composition of mixture fired	(0,=14.5)	$H_2=297$ $O_2=143$ $N_2=569$		$(^{\circ}H_4 = 16.8)$ $(^{\circ}H_2 = 16.8)$ $(^{\circ}L_2 = 16.8)$ $(^{\circ}L_3 = 16.8)$	$O_2 = 110$	
Initial pressure $p_i$ in atrospheres .	53 0	50 0	44 9	50 0	510	498
Maximum explosion pressure in atmospheres $p_m$		400 0	240 0	280 0	423 0	420 0
Final pressure of cold products in atmospheres $p_f$	30 5	28 74	47 1	51 7	42 6	4146
Ratio $p_m/p_i$	80	80	5 35	56	83	84
Ratio $p_f/p_i$	0 575	0 575	1 049	1 034	0.832	0 837
Time required for attainment of maximum pressure in seconds	0 011	0 010	0 05	0 08 about	0 10	0 10

The pressure records which are reproduced | in the following diagrams (Figs 11, 12, and 13), in which pressures in atmospheres are plotted against time in seconds, prove conclusively the absence of any direct relation between the

explosive mixture is transferred on explosion as sensible heat to its products and the magnitude of the chemical affinity between its combining constituents This is hardly to be wondered at, seeing the extreme rapidity of chemical interactual rate at which the potential energy of an action at high temperature as compared with




Fig. 11.

the rate at which the liberated heat can be communicated to and uniformly distributed amongst the products by ordinary physical processes. Attention may be drawn to the extreme slowness of the cooling in each case after the attainment of maximum pressure; this has also been observed by previous workers in other similar cases of gaseous explosions This was particularly marked in the case of the methane-air mixture, in which there was hardly

0.22 seconds after the attainment of maximum pressure (in 0.08 sec), a circumstance which may be due in part to the combustion taking place in well-defined chemical stages, and in part also to the operation of the exothermic secondary interaction between carbon monoxide and steam during the cooling period. On the other hand, the curve for the hydrogen-air mixture, where the combustion to steam is a direct and comparatively simple transaction any appreciable cooling during an interval of suggests that the attainment of maximum

pressure is succeeded by a period of gradual cooling uninfluenced by chemical combination.

It may be taken as the common experience of all investigators since Bunsen's day that the maximum effective pressures recorded when gaseous mixtures are fired in closed vessels are always considerably less than those calculated on the old assumption (now discarded) that the

whole heat of combustion is communicated without loss to the products, and that the specific heats of the products do not vary with temperature Another feature of the phenomenon is the very short time required for the attainment of maximum pressure relative to the subsequent cooling period

The great disparity between the found and




Fig. 12.

'calculated' maximum pressures has been attributed by the various investigators ('concerned to one or other of the following causes,

namely
1 To the marked increase in the specific heats of steam and carbon dioxide with temperature.

2 To the fact that in ordinary gaseous explosions, where detonation has not been determined,

combustion is by no means instantaneous, and may not be completed within the period required for the attainment of maximum pressure. Sn Dugald Clerk put forward this suggestion as long ago as 1886 1 in criticising the conclusions of Mallard and Le Chatcher respecting the great increase in the specific heats of steam and of carbon dioxide with temperature. He considered it highly




Fig 13

probable that combustion extends far into the actual 'cooling period' in gaseous explosions (and hence the long drawn-out 'cooling curve'), so that the system loses a certain part of the heat of combustion before the chemical action is completed This idea of a continued combustion finds support in H. B. Dixon's photographic researches, and chemists generally will concede its reality in any gaseous combination in which But to what detonation is not determined extent it may be held to affect the pressures actually recorded by explosions is still a matter o. conjecture.

3. To loss of energy by direct radiation -Thus in the explosion of a mixture of hydrogen and oxygen it is conceivable that the initial action results in the formation of an intensely vibrating molecular complex from which steam issues as the first recognisable product. Some experiments made in 1890 by Robert von Helmholtz (Beiblatter, 14, 589), showed that non-luminous hydrocarbon flames radiate about

5 pc of the heat of combustion of the gas, and more recent experiments by Callendar and Nelson show that the heat radiated from an ordinary non-luminous Bunsen flame may amount to between 15 and 20 p c. of the total heat of combustion.

4 To dissociation of products (steam and carbon dioxide) In the case of two combining gases producing a dissociable product, it is clear that if the average temperature in the system reaches that at which dissociation begins the combustion must be delayed whilst heat escapes from the system by radiation and con-Qualitatively the partial dissociation of steam and carbon dioxide has been proved at temperatures which are certainly exceeded by those of explosion flames; but it may be urged that, masmuch as all experiments upon dissociation have up to the present involved contact with hot solid surfaces, there is no positive evidence that the phenomenon would <sup>1</sup> Proc. Inst. Civil Engineers, 85.

play any conspicuous part in an unconfineu On the other hand, there is gaseous system direct experimental evidence of the attainment of enormously high temperatures in the explosion wave, temperatures which would generally be considered as far beyond that of the initial, or serhaps even of the complete, dissociation of steam or of carbon dioxide. Moreover, the fact that the rate of explosion of electrolytic gas is retarded rather more by an excess of oxygen than by a corresponding excess of nitrogen is

inconsistent with the supposition of any appreciable dissociation of steam in the explosion wave, and photographic records give no evidence of continued combination in the rear of the wave except where there are two or more chemical stages in the combustion.

Explosion of mixtures of coal gas and air -Experiments were made many years ago by Clerk with mixtures of coal gas and air in the explosion apparatus already described, with the following results:-

## GLASGOW COAL GAS AND AIR MIXTURES (CLERK).

Experiment	Proportion of gas by volume	Maximum pressure in lbs per sq inch above atmosphere	Maximum tempera- ture centigrade	Time of explosion
$egin{array}{c} a \\ b \\ c \\ d \\ e \end{array}$	14 12 10 10 10 10	52 63 69 89 96	1047° 1265° 1384° 1780° 1918°	0 28 second 0 18 ", 0·13 ", 0 07 ", 0 05 ",

that the greatest possible pressure exerted by a mixture of Glasgow coal gas and air at atmospheric pressure and temperature was 96 lbs. per square inch above atmosphere, the maximum pressure falling as dilution with air increased, while the time of explosion also increased. The shortest time of explosion was  $\frac{1}{20}$  second, and the the following results .-

From these experiments, it will be observed longest over 4 second, at the end of 060 to the greatest possible pressure exerted by a second from 30 to 35 lbs per square inch pressure above atmosphere still remained in the vessel. It is very noticeable that, volume for volume, coal gas produced higher pressure than hydrogen.

Experiments with Oldham gas and air gave

OLDHAM COAL GAS AND AIR MIXTURES (CLERK).

Experiment	Proportion of gas	Maximum pressure in lbs per sq inch above atmosphere	Maximum temperature centigrade	Time of explosion
a b c d e f y h	15-14-15-15-15-15-15-15-15-15-15-15-15-15-15-	40 0 51 5 60 0 61 0 78 0 87 0 90 0 91 0 80 0	806° 1033° 1202° 1220° 1557° 1733° 1792° 1812° 1595°	0 45 second 0 31 " 0 24 " 0 17 " 0 08 " 0 06 " 0 04 " 0 055 " 0 16 "

The greatest pressure produced by a mixture | the same vessel with mixtures previously comof this gas and air was 91 lbs. per square inch above atmosphere Comparing these numbers with the Glasgow gas experiments, it appears that with 14 vol. of Oldham gas present, the pressure attained was 0 5 lb. per square inch less, and the time of explosion is 0 31 second against 0.28 second with the same proportion of Glasgow With 1 of the two gases, Oldham was 5 lbs. less maximum pressure and 0 055 second against 0.05 second Glasgow.

In all these experiments the suppression of heat at the maximum temperature is very considerable, ranging from 50 to 60 p.c.

Experiments were also made by Clera 'n

pressed, and it is found that the pressures produced with any given mixture are proportional to the pressure before ignition—that is, with a mixture of constant composition double the pressure before explosion, keeping the temperature constant, doubles the pressure of explosion.

In 1900 Clerk made further experiments with London coal gas and air, with very complete precautions to ensure mixing, and with more accurate timing devices. The curves obtained with this apparatus are shown at Fig. 14, and the maximum temperatures and pressures, and time of explosion for the different mixtures, are as follows :-

Explosion in a Closed Vessel (Clerk, 1900). Mixtures of air with London coal gas. Temperature before explosion 16°

Pressure before explosion . 1481bs persq in


М	ixture	Max pressure above atmosphere in 1bs per sq inch	Temp of ex- plosion from observed pressure	Time of ex- plosion
Gas	Air		°C	sec.
1 vol.	12 vols	4		
l "	11 ,,	58	1150	0 290
l "	10 ,,	60	1155	0 305
1 ,,	9 ,,	65	1270	0 155
1 ,,	8 ,,	77	1475	0 087
1 ,,	7 ,,	80	1565	0.067
1 ,,	6 ,,	85	1660	0.055
1 ,,	5 ,,	87	1710	0 042
1 ,,	4 ,,	93	1830	0 045
		_		

J. E. Petavel has also made experiments on the explosion of coal gas and air compressed before ignition to pressures of over 1000 lbs per square inch Fig 15 shows the diagrams obtained, with the particulars marked thereunder.

In all these experiments, it is clear that if the temperature be calculated from the maximum pressure of explosion, there is apparently a suppression of heat of a very considerable amount at maximum temperature, the amount

being about 60 p c

Important experiments were also made by Prof. B Hopkinson, of Cambridge University, who investigated the actual temperatures reached during explosions of coal gas and air in an explosion vessel of the form shown at Fig 16 and found very large temperature variations at different parts of the vessel, and that the temperature differences existed after complete inflammation; quite independently of the local cooling effect of the walls of the vessel.



Explosion in a closed vessel, London gas, 1900 (Clerk).

He found that after the explosion of coal gas and air in the vessel, the distribution of temperature at the moment of maximum pressure was roughly as follows :-

Mean temperature (inferred from pressure) 1600° Temperature at centre of vessel, thermometer B 1900° Temperature 10 cm (4 inch) within wall, 1700° thermometer C Temperature 1 cm (04 mch) from wall, 1100° to 1300° at end, thermometer D Temperature 1 cm. (04 inch) within wall at side 850°

Half a second after maximum pressure, the distribution was as follows :---

Mean temperature (inferred from pressure) 1100° Mean temperature, exclusive of layer l cm thick at walls, determined by long platinum wire from B to D 1166° Temperature vessel, at centre of thermometer B . 1100° to 1200°

Here the temperature differences are much smaller than at the moment of maximum pressure Hopkinson found that the explosion vessel is entirely filled with flame when the pressure reaches 70 lbs, although the maximum pressure attained later is 82 lbs per square inch above atmosphere; that is, maximum pressure is not attained until one-thirtieth of a second after the vessel is completely filled by flame. Hopkinson's experiments also distinctly prove that the highest temperature reached in the vessel is at the point of origin of the ignition, and is due to the compression of the gases which are first heated by explosion to about 1200°, and then compressed by the compression from the walls inwards, as the mixture near the walls inflames.


Explosion in gas-engine cylinders—In all internal-combustion engines in general use at the present time, the mixture of fuel and air is compressed before ignition, the maximum temperature ranging from 1200° to 2500°; while the temperature of the hot gases at the moment of opening the exhaust valve may be

about 1000° Fig. 17 is a reproduction of an indicator diagram of an internal-combustion engine working on the Otto cycle The effect of weakening the mixture is shown by the different lines on the diagram the line a shows a diagram obtained from a strong mixture; b, a diagram from a slightly weaker mixture, and c, a diagram obtained from a very weak mixture. The effect of apparent suppression of heat appears on the gas-engine diagram as well as in the diagrams obtained by explosion in closed vessels at constant volume, referred to above, and it is extremely important, for the purpose of a correct theory of the internalcombustion engine, to ascertain exactly what takes place on explosion, and immediately after explosion of the mixture used, and a complete knowledge of the thermal properties of the working fluid and

its chemical composition at any instant during


the cycle is necessary

In the normal gas-engine diagram, the expansion line is at first usually considerably above the adiabatic, showing that the whole of the heat of combustion is not added before maximum pressure, and that heat must be added to the gases during a portion of the expansion, to make up for the flow of heat through the cylinder walls If, however, the cooling on expansion is too great, the expansion line may fall below the adiabatic, although there is still heat being added. The adiabatic expansion line tor any given mixture of gases depends on the specific heat of the mixture, and therefore, in order to determine what are the conditions as



Hopkinson's Explosion Vessel.


regards addition of heat, it is necessary to know the specific heat or internal energy of the mixture at any given temperature



Clerk has investigated the properties of the


working fluid in the gas engine itself, by a method which allowed direct observation to be made upon an actual charge taken into the engine in the ordinary operations of its cycle. The method consisted in subjecting the whole of the heated products of combustion to alternate compression and expansion, within engine cylinder, while the cooling proceeded, and observing by the indicator the successive pressure falls from revolution to revolution, together with the temperature and pressure use and fall due to the alternate compression and expansion. The engine was set to run at a given speed, and after the charge of gas and air had been drawn in, compressed and ignited, the exhaust valve and inlet valve prevented from opening, so that when

the piston reached the



Spherical enclosure capacity,  $551.9\ c$  C  $\,$  Temperature of enclosure  $^{\circ}$  before firing,  $18^{\circ}\mathrm{C}$  , after firing,  $24^{\circ}\mathrm{C}$  . Initial pressure, 1136 lbs. per sq. inch. Ratio  $\frac{\text{Air}}{\text{Gas}} = 6.0$ Maximum pressure, 9508 lbs per sq inch

Ratio Maximum pressure =8 36. Initial pressure



Spherical enclosure capacity, 551 9 c c after fring, 27°C. Temperature of enclosure: before Initial pressure, 1094 lbs. per sq. inch Maximum pressure, 9618 lbs per sq inch Ratio  $\frac{A \text{ ir}}{Ga \sigma} = 5.71$ .

Ratio Maximum pressure =88 Initial pressure

Fig. 15

Pise and kall of Pressure during and after Explosion (Petavel).

end of its power stroke, the exhaust gases were retained within the cylinder, and compressed by the piston to the maximum volume, and again allowed to expand and again compressed, and so on. These operations gave a diagram as shown at Fig. 18. From this diagram, by determining the temperature fall due to work done on expansion, Clerk determined the apparent specific heat of the products of combustion, and found that for the gas-engine mixture there was a considerable rise of apparent specific heat with temperature, and it seemed probable that this was largely due to a real increase of the actual specific heat or internal energy of the gases at high temperatures. The experiments, however, could not differentiate between the possible effects produced by dissociation and real specific heat change. Further, the values obtained were subject to some uncertainty, due to the difficulty of apportioning heat loss between successive compression and expansion lines, and it is probable that the values are somewhat raised by some film phenomena, or by smouldering combustion, tending to keep the expansion line from falling as rapidly as it would otherwise do.

Work of the British Association Committee on Gaseous Explosions .- Owing to the great importance to the theory of the internal-combus-

Jon engine of a complete understanding of the phenomena of gaseous explosions, the British Association appointed a committee for the purpose of investigating the whole matter, and although the work of this committee is not yet completed, very important results have already been obtained. These are set out in the Reports of 1908, Dublin; 1909, Winnipeg; and 1910, Sheffield The committee regard it as proved that on the explosion of gaseous mixtures with air, the quantity of carbonic acid or steam dissociated at the temperature of the explosion is so small, that the suppression of heat cannot be due to this cause.

They also concluded that chemical equilibrium is complete within an extremely short period after inflammation at any point outside the influence of the walls, and that there is no time rate of combustion, such as will account for the


apparent suppression of heat

The committee have also made a very careful comparison of the specific heat found by Holborn and Henning by methods not depending on explosion, with the values found by explosion experiments, such as those of Clerk, Mallard, and Le Chatelier, and Langen. The results given in calories per gram-molecule for the internal energy calculated from 100° at temperatures of 800°, 1200°, 1600°, and 2000°, are as follows .--

_							8	00,	1	200°	1600°	2000°
							Clerk	Holborn and Henning	Clerk	Holborn and Henning	Langen	Langen
Aır				٠.				3570	**************************************	5840	8700	11500
$CO_2$						.		6460	*****	10880	17000	23300
$H_2O$								4670		7930	14400	19900
Gas-c	engino	mi	xture	•	•		4250	3840	6900	6340	9800	13200
Ideal	gas*						34	130	5	400	7350	9300

\*  $C_1 = 4.0$ .

In this table, in order to reduce to foot-lbs, the figures should be multiplied by the factor 3 96.



Clerk diagram from engine at a brake load of 50 H.P. at '60 revolutions per minute.

The agreement between the figures by the two methods shows that the suppression of heat is very largely due to merease in the real specific

It seems clear also that the cooling due to the walls of the explosion vessel will damp out a certain portion of the combustion, and thus give a certain time rate of combustion, so that all the heat is not actually evolved at the point of maximum pressure or maximum temperature. By arranging a moving piston so that the exhaust products can be collected, and immediately cooled at any instant during the expansion stroke, Clerk has shown from an analysis of the exhaust products that there is

still some uncombined gas after the point of maximum temperature has been reached, even with strong mixtures

The committee also consider that the apparent suppression of heat in explosion experiments, when temperature is calculated from pressure, is partly due to under-estimation of the heat loss due to radiation. Callendar predicted that this would be the case from his experiments on the radiation from a Bunsen flame; and Hopkinson has shown that the

END OF THE SECOND VOLUME

RINTED IN GREAT BRITAIN BY VITLIAM CLOWES AND SONS, LIMITED, BECOLES,